Building better genome annotations across the tree of life

Adam H. Freedman[1*], Timothy B. Sackton[1]

[1] Faculty of Arts and Sciences, Informatics Group, Harvard University, 52 Oxford Street,

Cambridge, MA 02138

* Corresponding author: adamfreedman@fas.harvard.edu

ABSTRACT

Recent technological advances in long read DNA sequencing accompanied by dramatic

reduction in costs have made the production of genome assemblies financially

achievable and computationally feasible, such that genome assembly no longer

represents the major hurdle to evolutionary analysis for most non-model organisms.

Now, the more difficult challenge is to properly annotate a draft genome assembly once

it has been constructed. The primary challenge to annotation is how to select from the

myriad gene prediction tools that are currently available, determine what kinds of data

are necessary to generate high quality annotations, and evaluate the quality of the

annotation. To determine which methods perform the best and determine whether the

inclusion of RNA-seq data is necessary to obtain a high-quality annotation, we

generated annotations with 10 different methods for 21 different species spanning

vertebrates, plants, and insects. We found that the RNA-seq assembler Stringtie and

the annotation transfer method TOGA were consistently top performers across a variety

of metrics including BUSCO recovery, CDS length, and false positive rate, with the

exception that TOGA performed less in plants with larger genomes. RNA-seq alignment

rate was best with RNA-seq assemblers. HMM-based methods such as BRAKER,

MAKER, and multi-genome AUGUSTUS mostly underperformed relative to Stringtie

and TOGA. In general, inclusion of RNA-seq data will lead to substantial improvements

to genome annotations, and there may be cases where complementarity among

methods may motivate combining annotations from multiple sources.

INTRODUCTION

The reporting in 2001 of the first draft of the human genome sequence (Lander et al.

2001; Venter et al. 2001) ushered in a new era of genome-scale analysis, with a

concomitant, rapid increase in the development of bioinformatics tools and resources to

interrogate genomes for evolutionary patterns and features of biomedical interest. But

even as genomes became available for other model organisms such as mouse (*Mus*

musculus) (Mouse Genome Sequencing Consortium et al. 2002) and rhesus macaque

(*Macaca mulatta*) (Rhesus Macaque Genome Sequencing and Analysis Consortium et

al. 2007)—and had been previously published for smaller genomes such as *Drosophila*

melanogaster (Adams et al. 2000)—the prohibitive cost of generating genome

assemblies meant that research groups working on non-model organisms continued to

operate in the genomic dark. Absent genome assemblies and annotations, such groups

were forced to embark on time-consuming efforts to sequence small sets of conserved

genes with Sanger sequencing using primers designed with other genomes, target

anonymous loci such as AFLPs or *de novo* assembled RAD-seq reads. These methods

imposed an analytical glass ceiling on the types of inferences that could be made and

prevented the framing of research findings in a genomic context. While the advent of

RNA-seq inched non-model organism research closer to understanding patterns at

functional loci, *de novo* assembled transcriptomes presented novel analytical

challenges and potential distortions of evolutionary patterns relative to what would be

obtained with access to a genome assembly (Freedman et al. 2021).

Recent technological advances in long read DNA sequencing such as Pacific

Biosciences HiFi (Wenger et al. 2019) and Oxford Nanopore (Jain et al. 2018),

accompanied by dramatic reduction in costs have made the production of genome

assemblies financially achievable and computationally feasible, such that genome

assembly no longer represents the major hurdle to evolutionary analysis for most non-

model organisms. Now, the more difficult challenge is to properly annotate a draft

genome assembly once it has been constructed. The challenge is not so much the

difficulty or computational resources required to run any one genome annotation tool,

but how to a) select from the myriad gene prediction tools that are currently available, b)

determine what kinds of data are necessary to generate high quality annotations, and c)

evaluate the quality of the predicted transcript and gene models.

Currently available genome annotation tools approach the genome annotation

problem in very different ways. Early computational tools for annotation used Hidden

Markov Models (HMMs) to scan genomes for sequences representing protein-coding

intervals, with AUGUSTUS (Stanke and Waack 2003) being the most widely used

example. Recent implementations of this approach, such as BRAKER1 (Hoff et al.

2016) and BRAKER2 (Brůna et al. 2021) wrap optimized implementations of

AUGUSTUS, using protein and RNA-seq evidence, respectively—and with the latest

release, both—to train HMMs. Transcript assemblers such as Stringtie (Pertea et al.

2015) implement a graph-based framework to directly assemble transcripts from splice-

aware alignments of RNA-seq reads to the genome. Tools such as Comparative

AUGUSTUS (CGP) (Nachtweide and Stanke 2019) and TOGA (Kirilenko et al. 2023)

use whole genome alignments to transfer annotation evidence between genomes, with

the former involving multi-way transfer of HMM-based gene predictions, and the latter

lifting over annotations from a high quality reference annotation in an exon-aware

fashion.

A large part of difficulty in determining what strategy will work for "my genome" is

annotation methods have, for the most part, been benchmarked and optimized with

genomes from a very small slice of the tree of life: small genomes such as *D.*

melanogaster, *Caenorhabditis elegans*, *Saccharomyces cerevisiae*, or with emphasis

on vertebrates (Pertea et al. 2015; Hoff et al. 2016; Cantarel et al. 2008; Shao and

Kingsford 2017) —no surprise given the implications for discoveries relevant to human

health and disease. A related problem is that genome annotation will in the not too far

future and, by necessity, need to be conducted by the group that has assembled a

genome, rather than relying on Ensembl or NCBI to implement their automated

pipelines. While the number of genomes annotated per year by NCBI has remained flat

over the last few years, the number of published genome assemblies continues to

increase at an unprecedented rate. In summary, publicly accessible generators of and

repositories for annotations cannot keep up, such that the wait times between genome

submission date and annotation completion will increase to the point of being

impractical.

Motivated by the practical challenges facing research groups seeking to annotate

new genome assemblies, here we evaluate the contents of genome annotations

produced by ten different methods across a broad swath of the tree of life. These

methods sample the current state of the art for different approaches to the annotation

problem. Our taxonomic sampling includes two vertebrate clades (birds and mammals),

two insect clades (*drosophila* spp., and heliconiine butterflies), and two plant clades

(rosids and monocots), totaling 21 species. While "genome annotation" is often treated

as an omnibus term that includes both the prediction of the genomic positions of genes

and constituent isoforms, and the assigning of gene symbols and functions, we focus on

the first of these two components. Our goals are to determine a) which methods are

consistently top performers with respect to various sensitivity and specificity metrics b)

the contents of individual annotations with respect to gene model fragmentation and

fusion, c) whether the inclusion of RNA-seq data is essential for producing a high-quality

annotation, d) whether species and taxonomic group affect annotation method

performance, and e) whether there is complementarity among methods, such that

integration of > 1 method might lead to detectable improvements in sensitivity.

RESULTS

We evaluated ten different methods for genome annotation. At the core of five of

these—MAKER (Cantarel et al. 2008) with both protein and RNA-seq evidence;

BRAKER1; BRAKER2; CGP using protein evidence; and CGP with RNA-seq

evidence—are HMM-based *ab initio* predictions by AUGUSTUS, with MAKER and

BRAKER including predictions from additional *ab initio* tools. CGP enables evidence-

based prediction within a species, as well as annotation transfer across species in the in

the whole-genome alignment. TOGA performs an exon-aware liftover of annotations

from a related genome that, ideally, has a complete, high quality genome annotation.

Four methods assembly transcripts directly from RNA-seq alignments to a genome with

one of two aligners: Stringtie with either HISAT2 (Kim et al. 2019) or STAR (Dobin et al.

2013) as aligner, and Scallop (Shao and Kingsford 2017) with either of these aligners.

Many of our performance metrics use NCBI annotations as benchmarks. Derived from

multiple lines of evidence, outputs of the NCBI annotation pipeline are good

approximations to complete annotations. Futhermore, the species we annotated are

either well-established model organisms or others with a long history of study and

substantial genomic resources, such that their NCBI annotations are of the highest

quality.

BUSCO recovery

The ability of a gene prediction method to recover genes known to be conserved across

a wide array of taxa is a good approximation for its ability to recover at least some sub-

sequence of any gene, including those showing less conservation. BUSCO (Simão et

al. 2015) scores –measuring the proportion of BUSCO targets in a search that have

matches to query transcripts—varied considerably across methods and among species

and broader taxonomic groups. Methods built on HMMs (BRAKER and CGP)

consistently produced BUSCO scores for dipterans, heliconiine butterflies, and rosid

plants that were higher than for other methods, or tied with TOGA (Fig. 1, Fig. S1). In

contrast, RNA-seq assemblers consistently outperform HMM-based methods in

mammals and monocots, and one of the three bird species (Fig. 1, Fig. S1).

$BRAKER_{RNA-seq}$ recovered more BUSCOs than $BRAKER_{protein}$ in 16 of 20 species (Fig.

1, Fig. S1), and in cases where $BRAKER_{protein}$ recovered more the difference in recovery

is typically small. BUSCO scores were very similar between $CGP_{protein}$ and $CGP_{RNA-seq}$,

with $CGP_{protein}$ recovering slightly more BUSCOs in 14 of 16 species (Fig. 1). CGP failed

to produce more than a handful of transcript predictions for heliconiines, and

$BRAKER_{protein}$ consistently failed with *B. oleracea*, hence why these results are not

included. For *D. pseudoobscura* and *B. oleracea*, CGP BUSCO recovery was poor, with

BUSCO scores of approximately 50% or less. TOGA consistently produced the highest

BUSCO scores in birds and mammals, and for most species in other groups had scores

comparable to those produced by the top-performing method (Fig. 1). TOGA BUSCO

scores relative to other methods was lower in plants, particularly in two of four monocots

(Fig. 1). Somewhat surprisingly, MAKER, a putative full annotation workflow that

leverages both protein and RNA-seq evidence, consistently lagged in performance

behind other standalone methods. These patterns are not influenced by variation in the

presence of BUSCOs in the underlying genome assemblies, which might produce

taxonomic effects (correlation between BUSCO score and annotation BUSCO

score/genome BUSCO score, Pearson's $\rho = 0.998$, $p = 2.2 \times 10^{-16}$).

While there is considerable overlap between the BUSCOs that are recovered

between classes of methods, there are species for which methods that use RNA-seq

will recover BUSCOs that methods that use protein evidence cannot (Fig. S1). This is

particularly the case for species with larger genomes, such as birds, mammals, and *Z.*

mays, the monocot in our sample with a genome > 2Gb, and that is six to 10-fold larger

than the genomes of other monocots in our study. In some cases, 10-15% of BUSCOs

are recovered by methods leveraging RNA-seq but not by methods relying on protein

evidence. However, there are very few BUSCOs recovered by methods leveraging

RNA-seq that TOGA does not also recover; the taxonomic exception is monocots, for

which TOGA can perform poorly, likely due to known issues with whole genome

alignment for plants.

Annotation composition: number and length of CDS

The constituent CDS predictions that underlie BUSCO recovery rates varied widely

among methods. Across diverse taxa, TOGA consistently produced CDS whose length

distributions closely approximated those generated by NCBI, and with few exceptions

Stringtie did as well (Fig. S2). CGP annotations contained larger proportions of short

predictions than other methods, with weaker trends towards shorter CDS observed in

BRAKER. Scallop length distributions were often shifted towards shorter CDS to a

similar degree as the HMM-based method with the greatest proportion of short CDS

transcripts.

Annotations with tendencies towards shorter and far larger numbers of CDS

relative to NCBI are likely indicative of fragmented transcript models. We observed this

tendency for several species, particularly those with larger genomes (Fig. 2, Fig. S3).

Notable extreme outliers were the 7-fold and 4-fold larger CDS counts produced by

CGP for macaque and human, respectively (Fig. 4A). The best approximations to

median CDS length and number of NCBI annotations were typically produced by TOGA

or a Stringtie assembly (Fig. 2, Fig. S3).

False positives: intergenic predictions

For all but *A. thaliana*, the false positive rate (FPR) at which predicted genes fell entirely

within intergenic regions relative to the respective NCBI annotations was lowest for

TOGA; for *A. thaliana*, Stringtie (with STAR alignments) had the lowest FPR, and that

for TOGA was nearly identical. In general, when TOGA did not have the lowest FPR, an

RNA-seq assembler did (Fig. 3). For all species, FPR for the best performing method

was ≤ 10% and for many species those predictions occurred < 5% of the time (Fig. 3).

Regardless of the species and evidence type, FPR for CGP was much larger, often

exceeding 50%, with predictions using RNA-seq evidence being less prone to FPR than

those relying on protein evidence (Fig. 3). While BRAKER FPR was typically less than

that of CGP, regardless of the evidence type used, FPR was higher than the best

performing RNA-seq assembler in 16 of 18 species. The disparity between the low rates

for RNA-seq assemblers compared to other methods was most evident in rosids,

monocots, and mammals (Fig. 3). Nevertheless, the observed FPR suggest that, even

for the best performing methods, hundreds or even thousands of gene predictions will

fall outside of the genomic intervals for known real coding sequence. This raises the

question of whether these sequences are false negatives in the NCBI annotation, or

whether they are, in fact, false positives. We take a conservative approach and assume

that most of these predictions are false positives, asking whether there are transcript

features that might be predictive of such putative false positives so they can be

removed. The same patterns held at the transcript level (gene vs. transcript intergenic

prediction rate, Pearson's $\sigma = 0.994$, $p < 2.2 \times 10^{-16}$)

To better understand whether features of predicted transcripts distinguish those

that at least partly overlap NCBI gene intervals from those that are entirely in NCBI

intergenic regions, we fit random forest models to predictors summarizing sequence

content, expression level, and whether the associated ORF had strong evidence to a

match in the reference proteins of a related species. We did this for the five reference

species for which NCBI annotations are thought to be the most complete. Overall, out-

of-bag error rates (OOB) were consistently low for RNA-seq assemblers and TOGA,

with OOB always being < 0.05 for TOGA, and for RNA-seq assemblers only exceeding

5% in *Z. mays* (Fig. S4). Except for CGP, the class (genic vs. intergenic) error rates that

comprise OOB are substantially higher for intergenic predictions (Fig. S5). This result is

undoubtedly due to the fact, that, for most species-method combinations, there are far

fewer intergenic than genic predictions (Fig. S6), making it harder for random forest to

optimally classify intergenic sequences. However, random forest is able to correctly

classify the majority of intergenic predictions as intergenic for methods that predict large

numbers of intergenic transcripts (Fig. S6).

Estimates of node purity by the Gini index—an estimate of variable importance

that quantifies the extent to which removing a predictor reduces the frequency of

predictions matching the true class on either side of a split—reveal that whether or not a

transcript ORF has a BLAST hit is frequently the most powerful predictor of whether or

not a transcript is genic or intergenic (Fig. 7A,B; Fig. S7), with expression level and

CDS length also frequently making large contributions to the models. These results

suggest that, in the absence of a truth-set of known CDS intervals, some CDS features

are potentially useful for setting filters to discriminating true from false positive intergenic

predictions (Fig. 7 C,D). The importance of individual expression metrics is likely

underestimated, due to the correlation between expression metrics, such that the effect

of removing one is mitigated by the presence of another in any one of the constituent

trees in the random forest. These results collectively suggest that short, lowly expressed

transcripts without hits to an external protein database are enriched for intergenic (and

likely spurious) predictions.

Gene fusions

We defined fusions as cases where for a predicted gene, the CDS of an associated

transcript overlapped with the CDS of $>$ 1 NCBI gene, or different CDS transcripts of a

predicted gene each overlapped with the CDS of a single NCBI gene, but different

predicted CDS transcripts overlapped with different NCBI genes; these definitions were

not mutually exclusive. With few exceptions, the rate of putatively false fusions fell

below 5% across species and methods, with notable exceptions for a handful of

MAKER and BRAKER$_{RNA}$ gene sets (Fig. 5). In general, fusion rates for HMM-based

methods were lower than for RNA-seq assemblers, likely due to the tendency for

shorter CDS lengths of the former; TOGA fusion rates were consistently among the

lowest (Fig. 5). Gene-level fusions were not merely due to individual predicted CDS

transcripts spanning the CDS of multiple NCBI genes, but were often dominated by

cases where different transcripts from the same predicted gene each overlapped the

CDS of a different NCBI gene (Fig. S8).

Protein sequence completeness

BRAKER and CGP consistently had the highest percentage of predicted proteins that

had proper start and stop codon without any internal stop codons (Fig. S9), with the

former usually outperforming the latter by a modest margin. TOGA predictions had up to

20% fewer complete proteins than these methods, with Stringtie performing either

slightly better or slightly worse, depending upon the species. Scallop consistently had

the smallest percentage of predictions that represented complete proteins. Despite

being a pipeline meant to process predictions from multiple *ab inito* tools (e.g.

AUGUSTUS), MAKER performed worse than BRAKER and CGP and was often, and for

plants had the smallest proportion of complete protein predictions. Nevertheless, most

predicted transcripts in an annotation have a structure consistent with complete

proteins, and with the exception of Scallop, rarely dipped below 80%.

The utility of "completeness" may be misleading as a stand-alone measure of

whether a prediction is correct (or representing a true protein-coding sequence), if the

wrong trinucleotide sequences are classified as start and stop codons due to incorrect

inference of splice sites. That this may happen is highlighted by contrasting rates of

completeness with the frequency with which predicted protein sequences match those

in high quality protein databases. For each species, BLASTP was performed against a

database comprised of proteins derived from the NCBI annotations of the species we

included in their taxonomic group, including the species in question. The overall

proportions of BLASTP hits were, with the exception of CGP, very high (Fig. 6A). TOGA

was consistently the top performer, but for dipterans and rosids, TOGA, RNA-seq

assemblers and BRAKER were barely distinguishable; for monocots, birds, and

mammals, BRAKER with protein evidence was the second highest rate of BLASTP hits,

with Stringtie following close behind (Fig. 6A). However, when we focused on our

reference species for which NCBI annotations are of highest quality, in looking at the

distributions of the coverage of NCBI proteins in BLASTP hits—defined as the number

of matching bases by the length of the best-hit target—with the exception of *D.*

melanogaster, there was broad overlap in coverage distributions between complete and

incomplete protein predictions for BRAKER, CGP, and MAKER (Fig. 6BC, Fig. S10).

This suggests these tools are frequently producing truncated protein predictions by

identifying the wrong start or stop codons. In contrast, there was far less overlap for

RNA-seq assemblers and TOGA. Stringtie and TOGA appear to do a better job than

other tools of reconstructing the amino acid sequences of high-quality reference

annotations that we defined as our truth set. The strong performance of TOGA cannot

be attributed solely to the fact that the annotations being transferred originate from one

of the species contained in our protein databases used for BLASTP searches, as

excluding the proteins originating from the species whose annotations are being

transferred led to negligible decreases in percentages of predicted proteins with hits: *H.*

sapiens, 99.9% vs. 98.9%; *G. gallus*, 100% vs. 99.7%; *D. melanogaster*, 100% vs.

99.7%; *Z. mays*, 99.4% vs. 99.3%; and *A. thaliana*, 99.9% vs. 99.8%.

Transcriptome representation: expression

Random forest models we constructed to distinguish intergenic predictions from ones

overlapping known protein-coding gene intervals indicate that the former may be

characterized by low expression. Predictions by BRAKER and CGP (regardless of the

evidence type) and to a lesser extent MAKER contain larger proportions of genes with

TPM < 1 (Fig. 7A, S11) relative to RNA-seq assemblers. HMM-base methods using

RNA-seq evidence often had larger or comparable fractions of lowly expressed genes

compared to their counterparts using protein evidence (Fig. 7A, S11). TOGA predictions

did not have substantially elevated proportions of low TPM genes (relative to RNA-seq

assemblers), and for monocots had the lowest fraction of lowly expressed genes (Fig.

7A, S11). This may be due to alignments of expressed sequences being part of the

evidence used by the NCBI annotation pipeline. In general, with their direct connection

to expression, transcript assemblies of RNA-seq data (particularly those based upon

Stringtie) consistently had the smallest fraction of lowly expressed transcripts.

The RNA-seq read alignment rate to an annotation provides an estimate of the

extent to which an annotation captures the underlying expressed transcriptome.

Because BRAKER2, CGP, MAKER, and TOGA do not predict UTRs, and because

BRAKER's UTR prediction option was an experimental features we chose not to

include, we removed UTR intervals from Stringtie and Scallop annotations to make

methods comparable. When UTRs were excluded from consideration, alignment rates

were relatively low, only ever exceeding 50% for rosids, *D. melanogaster,* and for two of

four monocots (Fig. 7B); this is partly, to the RNA-seq data containing reads originating

from UTR intervals. While RNA-seq assemblers had alignment rates that were

frequently the highest for a particular species, there were only modest differences

between these tools and the best performing implementations of BRAKER or CGP (Fig.

7B). Rates for TOGA were lower than these, and were at approximately 10% for two of

four monocots, with rates for MAKER also being low relative to most other tools (Fig.

7B). Regardless of method, alignment rates for human annotations always fell below

20%. Even so, alignment rates varied in a manner similar to NCBI annotations without

UTRs (Pearson's $\rho = 0.86$, p=2.2×10^{-16}), albeit in most species-by-method

combinations lower than those for NCBI (Fig. 7B).

Because these comparisons were based upon alignment of the same reads that

were assembled by Stringtie and Scallop, we considered the possibility that this would

provide an unfair advantage to the assemblers relative to tools that only used RNA-seq

data to generate splice hints (BRAKER$_{RNA}$ and CGP$_{RNA}$), or to filter HMM-based

predictions post hoc (MAKER), and even more so for methods that did not use RNA-seq

data (BRAKER$_{protein}$ and CGP$_{protein}$). Training and test data alignment rates were

strongly correlated (Pearson's $\rho = 0.86$, p=2.2×10^{-16}), although for all species except

human (and for the BRAKER$_{protein}$ annotation for chicken), test alignment rates were

lower than training rates (Fig. S12A). Nevertheless, the reduction in test data alignment

rates relative to training data were modest, only exceeding 10% for *D. melanogaster*

and *Z. mays* for several methods (Fig. S12B). Only for *D. melanogaster* was there a

clear drop in the test alignment rate for Stringtie and Scallop relative to other methods

(Fig. S12), suggesting that the recycling of training data for alignment does not generate

substantial bias in the broad patterns we observe.

RNA-seq assemblers are agnostic to the functional role of the sequence intervals

from which reads originate, while HMM-based approaches and TOGA do not predict

UTRs. The inclusion of UTR intervals predicted by the assemblers led to large

increases in their alignment rates, such that they outperformed other methods (Fig.

S13). The magnitude of this alignment rate difference raises questions regarding the

contents of predicted UTR intervals. We thus assessed whether the lengths of predicted

UTRs are consistent with those found in NCBI annotations, and whether the alignment

rate increase they provide is due, at least in part, to there being true CDS intervals

contained within UTRs predicted by Transdecoder—CDS that it failed to predict. We

address these questions immediately below.

UTRs in RNA-seq assemblers

Ratios of UTR to CDS length are greater for Scallop and Stringtie assemblies than for

NCBI annotations (Fig. S14). These disproportionately long (relative to NCBI) UTRs

constitute an excess of target sequence for alignment, such that their exclusion clearly

contributes to a large drop in alignment rates. We considered the possibility that the

greater proportional length of UTRs for RNA-seq assemblers relative to NCBI

annotations could be due to the NCBI pipeline computationally truncating UTRs when

mitigating for cases of putative transcriptional readthrough past stop codons. In this

case the gold standard genomes (*H. sapiens*, *M. musculus*, *D. melanogaster*, *Z. mays*,

and *A. thaliana*) for which there has been extensive curation would be expected to have

a lower ratio of UTR to CDS length. However, plotting the ratio of predicted UTR to CDS

ratios for the assemblers over that for NBCI predictions produces the opposite pattern,

where these ratios of ratios are lower for the gold standard genomes (Fig. S14).

Next, we evaluated whether Transdecoder pipeline fails to predict CDS at the

ends of transcripts, and be default assigns them to the UTR functional class. If this were

a pervasive problem, then we would expect a large fraction of UTRs to have BLASTX

hits to an NCBI protein database for the same species. As would be expected if

undetected CDS occur in the UTR intervals for RNA-seq assemblers, the percentage of

transcripts with a UTR BLASTX hit as the length of UTRs relative to CDS for

assemblers increases relative to that observed for NCBI annotations (Fig. 8).

Depending upon the species and particular assembler-aligner combination, up to 60%

of transcripts may potentially contain undetected CDS in regions annotated as UTRs.

This suggests the failure to detect CDS plays a substantial role in the drop in RNA-seq

alignment rates when UTRs are excluded. This then leads to an underestimate of the

realizable alignment rate for RNA-seq assemblers than if these CDS were properly

incorporated into the annotations. It seems likely that the frequency of such undetected

CDS exons may contribute to the tendency for assemblers to have lower percentages of

proteins classified as complete relative to HMM-based methods (Fig. S9).

Multi-method integration

Above, we have demonstrated that MAKER, a workflow for integrating and filtering

predictions from multiple *ab initio* prediction tools, underperforms compared to several

different stand-alone annotation methods and across a variety of metrics. Another

integration approach, TSEBRA (Gabriel et al. 2021a) selects transcripts and merges

transcript models from separate runs with protein and RNA-seq evidence, respectively.

TSEBRA integration produces an annotation with BUSCO scores that are no better than

the best of the two BRAKER runs (Fig. S15A), and in most cases have slightly worse

scores. Proportions of predicted transcripts with BLASTP hits to the NCBI proteins from

the species group we investigated are consistently worse than either BRAKER

implementation (Fig. S15B). Similarly, in four of five reference species, the percentage

of intergenic genes is higher for TSEBRA than either BRAKER implementation (Fig.

S15C), with RNA-seq alignment rates falling between the two BRAKER runs (Fig.

S15D). In short, MAKER and TSEBRA lead to a loss of meaningful transcriptome

information relative to the best of the alternative BRAKER approaches.

　　　　While not necessarily an optimized method for merging annotations, we

investigated whether adding genes (and their constituent child features) from a second

annotation to a base annotation—requiring that those added genes fell entirely outside

of the gene intervals of that base annotation—would leverage the complementary

strengths of different methods while minimizing information loss. We also explored

whether successive additions from different methods would continue to improve the

overall annotation. There was a high degree of variability in the benefits of such

integration. While stand-alone methods like TOGA had the highest BUSCO score,

adding TOGA annotations to those of Stringtie achieved the best balance of maximizing

both BUSCO score and RNA$_{seq}$ alignment rate, overcoming the tradeoff observed

between these two metrics often seen in individual methods (Fig. 9A). Which method

one chooses as the base annotation can impact BUSCO scores, the number of genes

that have BLASTP hits to NCBI proteins, the percentage of predicted genes with such

hits, and the RNA-seq alignment rate. For example, using Stringtie$_{STAR}$ as a base

annotation leads to lower BUSCO scores and higher alignment rates than for

integrations that start with TOGA (Figs. 9A,B, S16, S17), and while integration

increased, as expected by definition, an increase in predicted genes, the percentage of

genes with at least one transcript having a BLASTP hit to an NCBI protein decreased,

suggesting that both real and spurious predictions get added (Figs. S16-S19). Method

integrations that did not include TOGA produced high RNA$_{seq}$ alignment rates but lower

BUSCO scores than TOGA and integrations that used it as the base annotation (upon

which to add others), with reasonably high fractions of genes with BLASTP hits (Figs.

9C, S18). Finally, in a scenario where RNA-seq data is not available, adding

BRAKER$_{protein}$ annotations to those of TOGA did little else other than to minimally

increase the RNA$_{seq}$ alignment rate (Figs. 9D, S19). In these four scenarios, adding

BRAKER$_{protein}$ had negligible effect.

DISCUSSION

With genome assembly and annotation increasingly becoming part of the workflow for

researchers studying non-model organisms, the choice of an annotation method

depends upon knowing whether a particular method will perform well in the species in

question, as well as what data will need to be generated to generate the best quality

annotation possible. Previous efforts to evaluate and compare methods have sampled a

small slice of the tree of life, and often focused on small, tractable genomes with

extensive genomic resources, or other model organisms such as *H. sapiens* or *M.*

musculus. This can make choosing a method more difficult, because the species used

to benchmark annotation tools may be evolutionary distant from a newly assembled

genome of interest. Our investigation overcomes this problem by evaluating a large set

of methods across the broadest taxonomic swath investigated to date, revealing both

cross-species and taxon-restricted patterns. By identifying methods that performed well

across diverse species, we believe that researchers using those methods will likely be

able to generate reasonably high-quality annotations for their newly assembled genome

of interest. Our findings have implications for study design, data collection, and

annotation method choice, and highlight ongoing challenges that require further

methods development.

First and foremost, the inclusion of RNA-seq data will invariably improve

annotation quality, particularly when used to assemble transcripts directly from

sequence alignment with Stringtie. While HMM-based methods such as CGP and

BRAKER using either protein or RNA-seq evidence can produce high BUSCO scores,

they consistently lag behind RNA-seq assemblers with respect to their representation of

the underlying transcriptome—as characterized by read alignment rates. Furthermore,

HMM-based approaches typically produce thousands of false-positive predictions which

need to be filtered but may be difficult to identify. That HMM-based predictions may

consistently be "complete", with proper start and stop codons, yet also shorter than

NCBI transcripts and those produced by RNA-seq assemblers, highlights a shortcoming

of HMM-based methods—splicing patterns that are consistent with an inferred model of

protein-coding sequence structure may not be the real pattern, instead representing

truncated open reading frames or spurious predictions. While we did not assess the

utility of cDNA long reads for annotation, we expect our findings to be robust to their

adoption, and that the performance disparities between RNA-seq assemblers and

HMM-based methods will almost certainly widen. The direct evidence of splicing

patterns across full length reads will enable reconstruction of full-length transcripts,

while in the HMM context, longer reads will simply lead to more accurate detection of

splice sites, and more accurate model parameterization—to the extent that any model

can capture the diversity of sequence composition and splicing patterns observed in

higher organisms. We suspect that increasingly accurate model parameterization will

lead to diminishing returns relative to direct assembly of transcripts from reads. RNA-

seq assemblers are particularly valuable for larger, more complex genomes and

consistently rank among the top methods across diverse performance metrics.

Furthermore, and while not the focus of this study, assemblers permit the inclusion of

non-coding RNAs in an annotation, with the caveat that it is more difficult to distinguish

real non-coding transcripts from spurious assembly of low-coverage transcriptional

noise. Overall, while the advantages of using RNA-seq assembly over other methods

may be diminished for smaller, less complex genomes, it clearly produces better

annotations than HMM-based approaches for more complex genomes, and reliably

produces relatively complete annotations regardless of taxonomic group or genome

organization.

The demonstrated superiority of RNA-seq assemblers to HMM-based

approaches may in fact be an underestimate of their relative performance. Our

discovery that predicted UTRs have high fractions of sequence that with BLASTX hits to

NCBI proteins suggests that we failed to recover many CDS exons. Our pipeline for

producing CDS annotations uses Transdecoder, and our results suggest that it may

have a harder time correctly classifying CDS exons at the termini of a transcript than

those within the transcript body. While long-read technology might help overcome this

deficiency, we suggest that there is room for methods development to improve ORF

detection from predicted CDS transcripts. Improved ORF detection and CDS exon

boundary delineation would lead to improved performance with respect to several of the

metrics we used to compare methods in this study.

Our findings also highlight the power and limitations of annotation transfer from

another species with a high-quality annotation, as is done by TOGA. While TOGA often

had high sensitivity (BUSCO scores) accompanied by low rates of intergenic predictions

and gene fusions, we found that sensitivity could be much lower in plants, especially in

those with larger genomes such as *Z. mays*. This undoubtedly stems from known

difficulties performing whole-genome alignment with plant genomes (Song et al. 2024).

Given the strong performance of RNA-seq assemblers across all the species we

surveyed, researchers should consider RNA-seq assembly when there are known

issues performing whole genome alignment for the species in question. For some taxa,

there may be few if any closely related species with high-quality annotations, or genome

alignments may be fragmented or contain many missing intervals in the target species.

Our finding that, for some species, there may complementarity among methods

for which BUSCOs are recovered—and that RNA-seq based methods can recover

genes that protein-based methods fail to predict—suggests that integration of

annotations across multiple methods may potentially improve sensitivity. That MAKER

performed poorly and TSEBRA appears to filter out real annotations when integrating

protein and RNA-seq iterations of BRAKER, suggest that additional methods

development on annotation integration is needed. Our naïve approach of consecutively

adding annotations from one method that did not overlap with a base annotation also

increased sensitivity, leading to an increase in the number of protein-coding genes with

BLASTP hits to NCBI proteins. Nevertheless, we did not attempt to apply any filters to

remove the spurious annotations that were invariably added in that workflow.

While we did not explore in depth how applying various filters impacted

performance metrics, the frequent observation of high rates of intergenic predictions, as

well as the occurrence of gene fusions, strongly suggest that more work is needed to

identify filters that strike the balance between removing as many low-quality annotations

as possible, while minimizing the filtering out of real sequences. It is often the case that

filters applied in the literature appear *ad hoc*, even if guided by intuition and experience.

We suggest a more quantitative approach is needed. For example, our application of

random forest to classify genic and intergenic predictions suggests that machine

learning approaches, while not perfect, offer great promise in identifying which variables

should be used to set filtering thresholds. We found that whether a sequence had a

BLAST hit to a set of known proteins, and expression level were useful discriminatory

variables. Of course, a new genome does not benefit from the advantage of a "truth set"

of real transcripts. However, a random forest (or other) model could, in principle, be

trained with annotations from a related species, and that model could be applied to the

predicted transcripts for a new genome assembly. Future work is needed to explore the

utility of such an approach.

Even as much work remains to be done, our findings suggest some general

guidelines for a researcher deciding how to annotate their newly assembled genome.

1. Generate RNA-seq data for at least the tissues related to the most pressing

project needs, but ideally, across as many tissues as necessary to capture the

species' transcriptional complexity. Use Stringtie to assemble transcripts, and,

until a better option is available, the Transdecoder workflow for adding CDS

features to the annotation.

2. Consider using TOGA if RNA-seq data are not available and a well-annotated

high-quality genome is available for a closely related species. If there is

complementarity in the recovery of seemingly real protein coding sequences

(determined with BUSCOs or gene symbols extracted from BLAST hits to

established protein databases), consider an approach that integrates predictions

of TOGA with BRAKER$_{protein}$, giving more weight to TOGA that BRAKER.

3. If it is not possible to use TOGA and RNA-seq data are not available, use

BRAKER$_{protein}$.

4. If RNA-seq data are available and if there appears to be complementarity in the

recovery of real protein-coding sequences between the methods, consider using

an approach to integrate predictions from Stringtie and TOGA, giving more

weight to Stringtie.

5. Either through a statistical approach such as random forest, or through

heuristically-thresholded metrics (e.g. expression level, BLAST hits to an

established protein database), remove predictions with a high likelihood of being

intergenic.

6. Given the non-trivial frequency of fusions detected in the methods we analyzed

(with the exception of TOGA), consider flagging genes that are likely fusions, e.g.

if BLAST hits of different of different transcripts are to functionally distinct genes

produced by genes with clearly different symbols, or similarly, if subsequences of

individual transcripts have such divergent BLAST targets. Exclude these fusion

genes from downstream expression analyses.

7. Consider using CGP in edge cases where, for example, there are a handful of

incomplete annotations for some related species (perhaps generated by already

available RNA-seq data), and the genome of interest is of small or modest size.

Expect to do extensive filtering to exclude many spurious predictions.

In conclusion, the longer-term challenge for building genome annotations across the

tree of life is to make methodological advances suggested above, and to integrate them

into reproducible, automated workflows that can be deployed with minimal headaches

for biologists. When this happens, population and comparative genomics studies will be

easy to scale to hundreds, and even thousands of species, unleashing unprecedented

power to tackle long-standing questions regarding the genetic architecture of phenotypic

variation and the evolutionary mechanisms that generate and maintain biodiversity.

METHODS

Target taxa

Genome annotation tools are typically developed and optimized using high quality

genome assemblies from a small suite of model organisms, e. g. *Homo* sapiens,

Caenorhabditis elegans, and *Drosophila melanogaster*. As a result, it is difficult to

generalize their performance in this narrow context to taxonomic groups that are highly

divergent from those focal taxa, and for which the genome assemblies may not be of

comparable quality. To facilitate more accurate generalizations regarding the

performance of annotation methods, and, conversely, to explore whether there are

effects of taxonomy and genome structure on annotation quality, we generated genome

annotations for 21 species spanning six taxonomic groups: three species of heliconiine

butterflies, three *Drosophila* species (dipterans), three birds, four mammals, four rosids,

and four monocots (Supplemental Table 1). With the exception of the butterflies, we

included as a "reference" a species for which both a high-quality genome assembly and

annotation were available, and downloaded assemblies and annotations from NCBI.

Each group contains at least one species that is relatively closely related to this

reference. Because the NCBI genome versions and annotations for heliconiines are

older than those widely used by the heliconiine research community, we used lepbase

assemblies that were filtered to remove all scaffolds less than 1kb in length (Edelman et

al. 2019). High quality annotations for these species were either unavailable, or

generated by tools we evaluated and thus inappropriate to serve as a truth set. For

example, the annotation for *H. melpomene* was generated with BRAKER. Because of

the unavailability of gold standard annotations for heliconines, for these species we only

generate a subset of performance metrics that don't rely of an annotation truth set.

Furthermore, *H. melpomene* is used as a reference assembly in whole-genome

alignments (see below) but we did not generate annotations for this species.

Genome assembly and reference annotation quality

To understand how features of genome assembly structure and quality might impact the

quality of annotations, we generated several summary statistics describing genome

contiguity and completeness. We generated standard statistics such as the total

genome size, the contig and scaffold N50, and the fraction of genomic nucleotides that

were soft-masked. We also quantified the number of single copy orthologs (BUSCOs)

contained in a genome (Simão et al. 2015), and calculated a BUSCO score as 1 –

(number missing BUSCOs/total number of BUSCOs searched).

To assess the quality of the reference genome annotations generated by NCBI,

for each genome we extracted protein-coding transcripts and generated transcriptome

BUSCO scores, and statistics on the minimum, maximum and median CDS length and

fraction of CDS bases that were soft-masked. We also calculated the median RNA-seq

alignment rate (see below) for each species' annotation, as well as the fraction of

transcripts for which estimated transcripts per million (TPM) was < 1.

RNA-seq data acquisition and processing

For use with annotation methods that either build transcripts from RNA-seq reads or use

read alignments to generate splice hints, for each species we downloaded 15-20 fastq

file accessions from NCBI's short read archive (SRA). We used the following criteria to

choose accessions. We only considered 1) paired-end reads with a release data of

2011 or later, 2) at most one run per biosample, 3) Illumina reads sequence on HiSeq

2000, NextSeq or newer instruments (i.e. no Genome Analyzer II), 4) we excluded

experimental treatments such as knockdowns, infections, and CRISPR modifications. If

these criteria resulted in > 20 possible biosamples, we further required a minimum read

length of 100bp, and for Metazoans, preferentially selected brain or head samples. If <

20 samples were available, we relaxed read length, release date, and instrument criteria

with the goal of retaining 15 biosamples; with the exception of the heliconiine butterfly

D. plexippus (for which 8 of 19 libraries had 36bp reads), we strictly excluded paired-

end libraries where the read length was < 50bp. In a small number of cases where

libraries contained hundreds of millions of reads, we down-sampled libraries to

approximately 20 million read pairs with seqtk (https://github.com/lh3/seqtk).

To process the reads prior to sequence alignment, we stripped adapters with

TrimGalore (https://www.bioinformatics.babraham.ac.uk/projects/trim_galore/

). We did not trim low-quality bases at the ends of reads because the short read aligners

used in this study soft-clip such bases such that they do not impact sequence

alignment, and because such trimming can bias expression estimates (Williams et al.

2016). So as to avoid having to make inferences from SRA metadata regarding the

strandedness of RNA-seq libraries, and to avoid the likely variable effectiveness of

stranding protocols, we treated all libraries as unstranded, under the assumption that

such information, if used would result in modest improvements in performance for

annotation methods that leverage RNA-seq alignments.

Annotation tools

RNA-seq assembly

We used RNA-seq reads to directly assembly transcripts with StringTie v. 2.1.2 and

Scallop v. 0.10.5. These assemblers take as input spliced alignments of reads to the

genome. We evaluated the impact of aligner by generating assemblies with two different

aligners: HISAT2 v. 2.2.1 and STAR v. 2.7.9a. Following best practice, and to leverage

evidence for splice sites across multiple samples, we used a 2-pass approach to

generate STAR alignments. In the first pass, we generated an initial set of alignments

for each sample. In the second pass, we concatenated the splice site tables generated

for each sample's first-pass alignment, and supplied the concatenated table as splice-

site evidence for the second pass alignment of each sample. To generate a merged

transcript assembly combining information across all individual-level assemblies, for

StringTie and Scallop we use the stringtie-merge function and TACO (Niknafs et al.

2017) v. 0.7.3, respectively. With our heliconiine species, we initially evaluated two

additional assemblers: PsiCLASS (Song et al. 2019) and Scallop2 (Zhang et al. 2022,

2). Poor performance relative to StringTie and Scallop, as well as excessive run times

for PsiCLASS, led us to not consider these two tools further.

StringTie and Scallop annotations contain transcript and exon features: they do

not predict CDS. Therefore, to incorporate CDS predictions into the merged

annotations, we use a TransDecoder v. 5.5.0

(https://github.com/TransDecoder/TransDecoder) and an associated workflow

(https://github.com/TransDecoder/TransDecoder/wiki) that predicts orfs, and then

leverages orf predictions to predict CDS and UTR intervals associated with the gtf

format input annotation file. After initial prediction of likely candidate orfs, we run blastp

(Altschul et al. 1990) v. 2.12.0 searches against a protein database consisting of Uniprot

and Trembl entries from all the species that we are attempting to annotate in the

species group, e.g. for dipterans, the database consists of entries for *D. melanogaster*,

D. pseudoobscura, and *D. yakuba*. We provide the search results as an input to

transdecoder-predict, such that given two similarly scoring orfs, we preferentially keep

the one with a blastp hit. In the interest of minimizing the filtering out of real orfs, we set

the maximum e-value threshold for these blastp searches to 1×10^{-4}. It should be noted

that the workflow as described filters out of the final annotation any transcript without a

retained orf prediction. The filtered orfs contain an unknown fraction of real orfs that

TransDecoder failed to discover, as well as ncRNAs. While the focus of our research is

on prediction of protein-coding genes, we provide as part of our code repository an

accessory script for adding back into the final annotation these putative false negatives

and ncRNA annotations.

Single-species ab initio methods

In contrast to transcript assembly-from-reads approaches, a long-established approach

for predicting genes (and coding sequences in particular) is to parameterize hidden

Markov models (HMMs) that are designed to traverse scaffolds, identify exon

boundaries, and connect exons into transcript and gene-level features. The most

sophisticated single-species versions of this approach use external evidence to

parameterize HMMs and identify specific genomic locations where exon splice junctions

are located. We evaluate BRAKER1 and BRAKER2 (both v. 2.1.6) , which conduct

iterative training and gene prediction using RNA-seq read and protein alignment

evidence, respectively. Both BRAKER flavors wrap *ab initio* prediction with AUGUSTUS

(Stanke et al. 2006) and GeneMark, with BRAKER1 using GeneMark-ET (Lomsadze et

al. 2014) and BRAKER2 using GeneMark-EP+ (Brůna et al. 2020). Following developer

recommendations, we provide protein evidence to BRAKER2 in the form of a protein

fasta from OrthoDB v. 10 (Kriventseva et al. 2019) for the relevant taxonomic group,

generated from pre-partitioned raw files as provided by the BRAKER developers

(https://bioinf.uni-greifswald.de/bioinf), downloaded on 14 Septermber, 2018. For

BRAKER1, we provide a bam file of RNA-seq STAR alignments merged across all

libraries from the species being annotated.

While we consider multi-method annotation integration below, we also evaluate

TSEBRA (Gabriel et al. 2021b), a python script-based tool to combine BRAKER1 and

BRAKER2 runs, and select a well-supported subset of transcripts. We run TSEBRA

following guidelines available at the TSEBRA github repository

(https://github.com/Gaius-Augustus/TSEBRA), running it on the braker.gtf files (that

include AUGUSTUS and GeneMark predictions) rather than on the AUGUSTUS-only

annotations.

Single-species exon-aware liftover: TOGA

Using whole-genome alignments to transfer annotations across species from well-

annotated to poorly or un-annotated species has a long history, e.g. with the UCSC

Genome Browser LiftOver tool first becoming available in 2006 (Hinrichs et al. 2006). To

perform such "liftovers", we use TOGA (Kirilenko et al. 2023), which transfers CDS

annotations across genomes in an exon-aware fashion that minimizes disruptions of

ORFs. TOGA takes as input a whole genome alignment, and involves several steps, the

details of which we provide at https://github.com/harvardinformatics/GenomeAnnotation-

TOGA, and are an adaptation of the workflow described at

https://github.com/hillerlab/TOGA. To remove potential spurious or bad annotation

transfers, we filter out any transcripts in the primary annotation output

(query_annotation.bed) for which there was not a corresponding entry in

orthology_classification.tsv, i.e. transcripts for which TOGA could not determine an

orthology class. Within each taxonomic group, we transfer annotations from the high-

quality reference to all other species within the group, and from the species most closely

related to the reference back to the reference species. For example, for dipterans we

carry out three TOGA analyses, transferring *D. melanogaster* to both *D. pseudoobscura*

and *D. yakuba*, and from *D. yakuba* to *D. melanogaster*.

Multi-species ab initio annotation

In studies seeking to perform phylogenetic comparative analyses, annotate multiple

genome assemblies from related organisms, or where annotations or evidence (protein

or RNA-seq) already exist for a subset of species of interest, methods that transfer

evidence between lineages offer, in principle, a promising approach for performing

genome annotation. We evaluate the most well-established approach for doing this,

AUGUSTUS run in comparative mode (König et al. 2016), referred to hereafter as CGP.

CGP relies on whole-genome alignment (WGA). Thus, as a first step, for each

taxonomic group of genomes, we use Progressive Cactus (Armstrong et al. 2020) to

produce a WGA. We then use an AUGUSTUS accessory script, *hal2maf_split.pl*, to split

the hal-format cactus output file into multiple sub-files in multiple alignment (MAF)

format; in doing so, we set as the "reference" genome (with which to provide coordinate

anchors) a species with both a highly contiguous assembly and a high-quality

annotation, and split in such a way so as to avoid splits that bisect the genomic

coordinates of annotated genes in the reference. For each taxonomic group of species,

we run CGP twice, once with splice site evidence from protein alignments, and once

from RNA-seq alignments. For analysis with protein evidence, similar to analysis with

BRAKER2, we use OrthoDB v.100 data representing the taxonomic group. For analysis

with RNA-seq we used the merged STAR alignments across samples. In both

instances, following guidelines from the developers (Hoff and Stanke 2019), we

generate splice hints files for each species using scripts and code provided as part of

the AUGSTUS package. In both modes, we do not predict UTRs, nor do we predict

alternative isoform, i.e. one transcript prediction is made per putative gene. Detailed

instructions regarding how we generated hints and run CGP are found at

https://github.com/harvardinformatics/GenomeAnnotation-ComparativeAugustus.

MAKER

MAKER (Cantarel et al. 2008) is a genome annotation pipeline that has the ability to

integrate multiple *ab initio* gene prediction packages, and to use protein and RNA-seq

derived external evidence to perform post hoc curation of predictions. Because results

with MAKER usually involve > 1 runs in order to retrain gene-prediction models, it is not

a fully automated pipeline. Nevertheless, it has been used extensively due to its

purported ease of use. MAKER also has the option to perform quality filtering and

integration of annotations with EVidenceModeler (EVM) (Haas et al. 2008). For initial

testing with three heliconiine species, we ran MAKER v. 3.01.03 four different ways that

integrate predictions from AUGSTUS (Stanke et al. 2006), SNAP (Korf 2004), and

Genemark-ES (Lomsadze et al. 2005): 1) protein evidence only, without EVM; 2) protein

and RNA-seq evidence, without EVM; 3) protein evidence only, with EVM; and 4)

protein and RNA-seq evidence, with EVM. For protein evidence, we used the protein

accessions associated with the lepbase (lepbase.org) Hmel2 genome assembly, which

are proteins derived from BRAKER predictions. RNA-seq evidence was included as a

gff3 file generated from the Stringtie assembly using STAR alignments of the species'

RNA-seq samples. We used default settings for the EVM configuration scoring file. To

produce annotations, we ran MAKER twice closely following Daren Card's detailed

workflow (https://gist.github.com/darencard/bb1001ac1532dd4225b030cf0cd61ce2);

see also https://github.com/harvardinformatics/GenomeAnnotation-Maker . Because

with these test runs the use of EVM frequently produced lower quality annotations, and

because we wished to evaluate the potential of MAKER as a full-service annotation tool,

runs for other taxa were only performed with both protein and RNA-seq evidence and

without EVM. Furthermore, because MAKER is computationally intensive and can take

a considerable amount of time to run, for the other taxonomic groups we only generate

MAKER annotations for the "reference" species of each group, with protein evidence

being represented by the NCBI protein accession associated with the genome for the

next closely related species in the set of species we annotated within each taxonomic

group.

Annotation quality metrics

Accurate annotation of UTRs is challenging, and even more so for non-model

organisms for which RNA-seq data are typically sparse. Similarly, long non-coding

RNAs are also difficult to annotate (Uszczynska-Ratajczak et al. 2018). In most

contexts, neither feature is crucial to genome-enabled evolutionary studies in non-model

organisms. Thus, we focus our evaluation of genome annotation methods on protein

coding sequences, filtering out UTRs and non-coding loci. For annotation methods that

include the UTR portions of mRNAs, we strip UTR exons (and any UTR-labelled

features) from annotation gtf or gff3 files. We also do this for NCBI gff3 files prior to

comparisons with the annotation methods we evaluate.

782   We use the NCBI genome annotations as putative sets of "true" annotations.

Although the quality of these annotations certainly vary due to many factors – the quality

of the genome assembly, the amount and kind of experimental evidence available at the

time the annotation was generated, challenges in annotating larger, more complex

genomes—we believe they are a reasonable approximation to annotations one would

hope to achieve with a new genome assembly, using stand-alone annotation tools

deployed on local HPC clusters. Nevertheless, while we make comparisons to NCBI

annotations for all species-tool combinations, we pay particularly close attention to

those species for which we know the genomes and annotations are of the highest

quality: *H. sapiens*, *D. melanogaster*, *Z. mays*, *Arabidopsis thaliana*, and *Gallus gallus*.

Details on bioinformatics package command lines and custom python scripts are

available in the relevant GitHub repositories detailed in the DATA ACCESS section at

the end of this paper.

Annotation completeness: transcriptome BUSCOs and expression

To assess transcriptome completeness, we calculate transcriptome BUSCO scores and

compare them to scores for the NCBI transcriptomes. For comparisons across all

species-method combinations, and in order to normalize for varying degrees of genome

assembly completeness and quality, we calculate ratios of transcriptome BUSCO score

to that for the respective genome. To evaluate the extent to which the predicted

transcriptome represents the expressed transcriptome, we then use RSEM (Li and

Dewey 2011) v. 1.3.3 to wrap bowtie2 (Langmead and Salzberg 2012) alignment of

each RNA-seq library to the predicted transcriptome, and estimate both gene and

isoform-level expression. From those alignments, for each annotation we calculate the

median alignment rate (across the set of samples), and the proportion of genes and

transcripts for which TPM < 1. Because using the same RNA-seq libraries to generate

transcriptome assemblies with Stringtie and Scallop may bias alignment rates upwards

relative to tools that don't leverage evidence from those RNA-seq libraries, we also

perform alignments on an additional test set of six RNA-seq paired-end SRA accessions

for each of our five reference species.

Protein-level statistics

For each genome annotation we report the number of protein-coding genes, and the

number of CDS transcripts. We use GetProteinFastaStats.py, a custom python script

that leverages biopython (Cock et al. 2009) modules to calculate mean and median

protein lengths, the fraction of predicted proteins that have internal stop codons, and the

fraction that are complete, where complete is defined as having proper start and stop

codons, and no internal stop codons. For TOGA annotations, we generate these

statistics from the protein sequences the software outputs (after filtering out those

without ortholog classifications). For all other annotation tools we used the version of

gffread distributed with cufflinks (Trapnell et al. 2010) v. 2.2.1, to extract the protein

sequences.

To estimate the fraction of protein predictions that were real, we used blastp

(Camacho et al. 2009) v. 2.12.0 to search for matches, with a maximum e-value of 1 x

10^{-5}, against a database consisting of the NCBI protein accessions for all of the species

that were used in our study for the taxonomic group of interest.

False positives: intergenic predictions

Motivated, in part, by some tools predicting far more CDS transcripts that are recorded

in NCBI annotations, we assessed whether this could be due to (presumably incorrect)

intergenic predictions, where intergenic is defined as falling entirely outside of the CDS

intervals for all protein-coding genes annotated by NCBI. To do this, for each new

annotation, we generate transcript interval and gene interval bed files, where each entry

represented the genomic boundaries of the transcript or gene (excluding UTRs),

respectively. We then use bedtools v. 2.26.0 (Quinlan and Hall 2010) to intersect these

files with a bed file consisting of UTR-stripped NCBI gene boundaries, recording the

number of bases of overlap such that only same-strand overlaps were counted as

overlaps, e.g. *intersectBed -s -wao -a newannotation_intervals.bed -b*

NCBI_gene_intervals.bed. We then count the number of predicted transcripts and

genes lacking any overlap with NCBI gene coordinates.

Gene fusions

Real fusion events in which a transcript contains CDS from different annotated genes

should be extremely rare. Thus, the presence of non-trivial frequencies of predicted

transcripts for which exons span multiple NCBI genes most likely represent

bioinformatics pipeline-induced errors. We evaluate fusions at the gene level for which

we define three types: 1) an individual predicted CDS overlapping with CDS from

multiple NCBI genes, 2) different CDS originating from the same predicted gene

overlapping with the CDS of different NCBI genes, and 3) cases where both of these

types of fusions occur. To quantify the frequency of these fusion events, we first

converted the CDS features of a species' NCBI annotation and the CDS annotations of

a method being evaluated to bed format. Next, we used bedtools to perform a "left outer

join" of NCBI CDS features to those of the new annotation, e.g. *bedtools -loj -a*

newannotation_cds.bed -b ncbi_cds.bed > loj_overlaps.bed. We then evaluate each

gene in the new annotation relative to the first two classes of fusions using a custom

python script, BuildProteinCodingGeneCdsFusionSummaryTable.py and including the -

filter-nested switch that excludes from calculations known fusions that are present in the

NCBI annotation. We calculate the frequency of all three classes of fusions using basic

awk commaxnds. Our bedtools operation does not enforce same-strand matching, but

our python script does, such that we do not consider overlaps between new predictions

and NCBI predictions on opposite strands.

Undetected CDS in UTRs

Our RNA-seq assembly pipelines integrate ORF finding with Transdecoder such that

exon features are decomposed further into CDS and UTR features. While we exclude

UTRs to make assembler performance metrics comparable to tools that do not predict

UTRs, we observed a substantial drop-off in RNA-seq read alignment rates to the

Stringtie and scallop annotations when UTRs were filtered out: unfiltered annotations

had much higher alignment rates than all other methods, but were on par with those

methods after excluding UTR intervals. To examine the cause of this phenomenon, we

considered three possible causes. First, because for relatively complete high-quality

annotations the NCBI annotation pipeline will computationally truncate UTRs to prevent

stop-codon readthrough, we contrasted the length distributions RNA-seq assembler

UTRs and those from NCBI, expecting that the disparity would be greater for the

reference genomes for each of our taxonomic groups than for other, more recent

genome assemblies for which truncation would not be as severe. Under this scenario,

the reduction in alignment rate after UTR removal would be due to an excess of reads

originating from transcriptional readthrough (or because the NCBI UTR truncation was

overly conservative). Next, we considered the possibility that Transdecoder consistently

fails to predict CDS orfs at the terminal ends of transcripts, such that a large proportion

of real CDS sequences is being incorrectly filtered out when we strip out UTRs. To test

this hypothesis, we extracted the UTR sequences from the Stringtie and scallop

annotations and used blastx (Camacho et al. 2009) v. 2.12.0 to search for matches

against the NCBI protein sequences from the same species' NCBI accession, with a

maximum e-value of 1×10^{-5}. We calculated the fraction of transcripts for which at least

one of the UTRs had a hit to the protein database.

Mulit-method integration

To determine whether integration of multiple annotation methods could harness the

complementary strengths of individual methods, we tested an approach where we 1) set

one annotation as the "base annotation", 2) add features from a second annotation

methods that fall entirely outside of the gene intervals for the base annotation, and 3)

iterate this process for additional methods, e.g. adding features from a third method that

do not overlap genes from the integration of the first two annotations. We demonstrated

this approach for *Homo sapiens*, investigating the effects of different choices for base

annotation, exclusion of RNA-seq assembler or TOGA, and whether there are

decreasing returns with increasing number of integrated methods. For these

comparisons we focused on four metrics: BUSCO score, RNA-seq alignment rate, the

total number of genes with BLASTP hits to NCBI reference proteins, and the percentage

of all integrated genes that have such hits. While we do not compare this approach

directly to the other integration methods evaluated here (MAKER and TSEBRA,) our

results demonstrate that both of these annotation tools under-perform relative to other

standalone methods.

REFERENCES

Adams MD, Celniker SE, Holt RA, Evans CA, Gocayne JD, Amanatides PG, Scherer
908   SE, Li PW, Hoskins RA, Galle RF, et al. 2000. The genome sequence of
909   Drosophila melanogaster. *Science* **287**: 2185–2195.

910   Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. 1990. Basic local alignment
911   search tool. *J Mol Biol* **215**: 403–410.

912   Armstrong J, Hickey G, Diekhans M, Fiddes IT, Novak AM, Deran A, Fang Q, Xie D,
913   Feng S, Stiller J, et al. 2020. Progressive Cactus is a multiple-genome aligner for
914   the thousand-genome era. *Nature* **587**: 246–251.

915   Brůna T, Hoff KJ, Lomsadze A, Stanke M, Borodovsky M. 2021. BRAKER2: automatic
916   eukaryotic genome annotation with GeneMark-EP+ and AUGUSTUS supported
917   by a protein database. *NAR Genomics Bioinforma* **3**: lqaa108.

918   Brůna T, Lomsadze A, Borodovsky M. 2020. GeneMark-EP+: eukaryotic gene
919   prediction with self-training in the space of genes and proteins. *NAR Genomics*
920   *Bioinforma* **2**: lqaa026.

921   Camacho C, Coulouris G, Avagyan V, Ma N, Papadopoulos J, Bealer K, Madden TL.
922   2009. BLAST+: architecture and applications. *BMC Bioinformatics* **10**: 421.

923   Cantarel BL, Korf I, Robb SMC, Parra G, Ross E, Moore B, Holt C, Alvarado AS,
924   Yandell M. 2008. MAKER: An easy-to-use annotation pipeline designed for
925   emerging model organism genomes. *Genome Res* **18**: 188–196.

926   Cock PJA, Antao T, Chang JT, Chapman BA, Cox CJ, Dalke A, Friedberg I, Hamelryck
927   T, Kauff F, Wilczynski B, et al. 2009. Biopython: freely available Python tools for
928   computational molecular biology and bioinformatics. *Bioinforma Oxf Engl* **25**:
929   1422–1423.

930   Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, Batut P, Chaisson M,
931   Gingeras TR. 2013. STAR: ultrafast universal RNA-seq aligner. *Bioinformatics*
932   **29**: 15–21.

933   Edelman NB, Frandsen PB, Miyagi M, Clavijo B, Davey J, Dikow RB, García-Accinelli
934   G, Van Belleghem SM, Patterson N, Neafsey DE, et al. 2019. Genomic
935   architecture and introgression shape a butterfly radiation. *Science* **366**: 594–599.

Freedman AH, Clamp M, Sackton TB. 2021. Error, noise and bias in de novo
transcriptome assemblies. *Mol Ecol Resour* **21**: 18–29.

Gabriel L, Hoff KJ, Brůna T, Borodovsky M, Stanke M. 2021a. TSEBRA: transcript
selector for BRAKER. *BMC Bioinformatics* **22**: 566.

Gabriel L, Hoff KJ, Brůna T, Borodovsky M, Stanke M. 2021b. TSEBRA: transcript
selector for BRAKER. *BMC Bioinformatics* **22**: 566.

Haas BJ, Salzberg SL, Zhu W, Pertea M, Allen JE, Orvis J, White O, Buell CR,
Wortman JR. 2008. Automated eukaryotic gene structure annotation using
EVidenceModeler and the Program to Assemble Spliced Alignments. *Genome*
Biol **9**: R7.

Hinrichs AS, Karolchik D, Baertsch R, Barber GP, Bejerano G, Clawson H, Diekhans M,
Furey TS, Harte RA, Hsu F, et al. 2006. The UCSC Genome Browser Database:
update 2006. *Nucleic Acids Res* **34**: D590-598.

Hoff KJ, Lange S, Lomsadze A, Borodovsky M, Stanke M. 2016. BRAKER1:
Unsupervised RNA-Seq-Based Genome Annotation with GeneMark-ET and
AUGUSTUS. *Bioinforma Oxf Engl* **32**: 767–769.

Hoff KJ, Stanke M. 2019. Predicting Genes in Single Genomes with AUGUSTUS. *Curr*
Protoc Bioinforma **65**: e57.

Jain M, Koren S, Miga KH, Quick J, Rand AC, Sasani TA, Tyson JR, Beggs AD, Dilthey
AT, Fiddes IT, et al. 2018. Nanopore sequencing and assembly of a human
genome with ultra-long reads. *Nat Biotechnol* **36**: 338–345.

Kim D, Paggi JM, Park C, Bennett C, Salzberg SL. 2019. Graph-based genome
alignment and genotyping with HISAT2 and HISAT-genotype. *Nat Biotechnol* **37**:
907–915.

Kirilenko BM, Munegowda C, Osipova E, Jebb D, Sharma V, Blumer M, Morales AE,
Ahmed A-W, Kontopoulos D-G, Hilgers L, et al. 2023. Integrating gene
annotation with orthology inference at scale. *Science* **380**: eabn3107.

König S, Romoth LW, Gerischer L, Stanke M. 2016. Simultaneous gene finding in
multiple genomes. *Bioinformatics* **32**: 3388–3395.

Korf I. 2004. Gene finding in novel genomes. *BMC Bioinformatics* **5**: 59.

Kriventseva EV, Kuznetsov D, Tegenfeldt F, Manni M, Dias R, Simão FA, Zdobnov EM.
2019. OrthoDB v10: sampling the diversity of animal, plant, fungal, protist,
bacterial and viral genomes for evolutionary and functional annotations of
orthologs. *Nucleic Acids Res* **47**: D807–D811.

Lander ES, Linton LM, Birren B, Nusbaum C, Zody MC, Baldwin J, Devon K, Dewar K,
Doyle M, FitzHugh W, et al. 2001. Initial sequencing and analysis of the human
genome. *Nature* **409**: 860–921.

Langmead B, Salzberg SL. 2012. Fast gapped-read alignment with Bowtie 2. *Nat
Methods* **9**: 357–359.

Li B, Dewey CN. 2011. RSEM: accurate transcript quantification from RNA-Seq data
with or without a reference genome. *BMC Bioinformatics* **12**: 323.

Lomsadze A, Burns PD, Borodovsky M. 2014. Integration of mapped RNA-Seq reads
into automatic training of eukaryotic gene finding algorithm. *Nucleic Acids Res*
42: e119.

Lomsadze A, Ter-Hovhannisyan V, Chernoff YO, Borodovsky M. 2005. Gene
identification in novel eukaryotic genomes by self-training algorithm. *Nucleic
Acids Res* **33**: 6494–6506.

Mouse Genome Sequencing Consortium, Waterston RH, Lindblad-Toh K, Birney E,
Rogers J, Abril JF, Agarwal P, Agarwala R, Ainscough R, Alexandersson M, et
al. 2002. Initial sequencing and comparative analysis of the mouse genome.
Nature **420**: 520–562.

Nachtweide S, Stanke M. 2019. Multi-Genome Annotation with AUGUSTUS. *Methods
Mol Biol Clifton NJ* **1962**: 139–160.

Niknafs YS, Pandian B, Iyer HK, Chinnaiyan AM, Iyer MK. 2017. TACO produces robust
multi-sample transcriptome assemblies from RNA-seq. *Nat Methods* **14**: 68–70.

Pertea M, Pertea GM, Antonescu CM, Chang T-C, Mendell JT, Salzberg SL. 2015.
StringTie enables improved reconstruction of a transcriptome from RNA-seq
reads. *Nat Biotechnol* **33**: 290–295.

Quinlan AR, Hall IM. 2010. BEDTools: a flexible suite of utilities for comparing genomic
features. *Bioinformatics* **26**: 841–842.

Rhesus Macaque Genome Sequencing and Analysis Consortium, Gibbs RA, Rogers J,
Katze MG, Bumgarner R, Weinstock GM, Mardis ER, Remington KA, Strausberg
RL, Venter JC, et al. 2007. Evolutionary and biomedical insights from the rhesus
macaque genome. *Science* **316**: 222–234.

Shao M, Kingsford C. 2017. Accurate assembly of transcripts through phase-preserving
graph decomposition. *Nat Biotechnol* **35**: 1167–1169.

Simão FA, Waterhouse RM, Ioannidis P, Kriventseva EV, Zdobnov EM. 2015. BUSCO:
assessing genome assembly and annotation completeness with single-copy
orthologs. *Bioinforma Oxf Engl* **31**: 3210–3212.

Song B, Buckler ES, Stitzer MC. 2024. New whole-genome alignment tools are needed
for tapping into plant diversity. *Trends Plant Sci* **29**: 355–369.

Song L, Sabunciyan S, Yang G, Florea L. 2019. A multi-sample approach increases the
accuracy of transcript assembly. *Nat Commun* **10**: 5000.

Stanke M, Keller O, Gunduz I, Hayes A, Waack S, Morgenstern B. 2006. AUGUSTUS:
ab initio prediction of alternative transcripts. *Nucleic Acids Res* **34**: W435-439.

Stanke M, Waack S. 2003. Gene prediction with a hidden Markov model and a new
intron submodel. *Bioinforma Oxf Engl* **19 Suppl 2**: ii215-225.

Trapnell C, Williams BA, Pertea G, Mortazavi A, Kwan G, van Baren MJ, Salzberg SL,
Wold BJ, Pachter L. 2010. Transcript assembly and quantification by RNA-Seq
reveals unannotated transcripts and isoform switching during cell differentiation.
Nat Biotechnol **28**: 511–515.

Uszczynska-Ratajczak B, Lagarde J, Frankish A, Guigó R, Johnson R. 2018. Towards a
complete map of the human long non-coding RNA transcriptome. *Nat Rev Genet*
19: 535–548.

Venter JC, Adams MD, Myers EW, Li PW, Mural RJ, Sutton GG, Smith HO, Yandell M,
Evans CA, Holt RA, et al. 2001. The sequence of the human genome. *Science*
291: 1304–1351.

Wenger AM, Peluso P, Rowell WJ, Chang P-C, Hall RJ, Concepcion GT, Ebler J,
Fungtammasan A, Kolesnikov A, Olson ND, et al. 2019. Accurate circular
consensus long-read sequencing improves variant detection and assembly of a
human genome. *Nat Biotechnol* **37**: 1155–1162.

Williams CR, Baccarella A, Parrish JZ, Kim CC. 2016. Trimming of sequence reads
alters RNA-Seq gene expression estimates. *BMC Bioinformatics* **17**: 103.

Zhang Q, Shi Q, Shao M. 2022. Accurate assembly of multi-end RNA-seq data with
Scallop2. *Nat Comput Sci* **2**: 148–152.

1031

DATA ACCESS

Detailed explanation of steps in annotation pipelines, along with associated python

scripts for data processing are provided in several repositories listed below

• RNA-seq transcript assembly:

https://github.com/harvardinformatics/GenomeAnnotation-RNAseqAssembly/

• BRAKER: https://github.com/harvardinformatics/GenomeAnnotation-Braker

• TOGA: https://github.com/harvardinformatics/GenomeAnnotation-TOGA

• CGP: https://github.com/harvardinformatics/GenomeAnnotation-

ComparativeAugustus

• MAKER: https://github.com/harvardinformatics/GenomeAnnotation-MAKER

In addition, HPC slurm job scripts and specific command lines used to run annotation

tools in this paper, as well as python scripts to generate annotation quality metrics are

available at https://github.com/harvardinformatics/GenomeAnnotation.

Figure 1. BUSCO scores by annotation method for (A) vertebrates, (B) plants,, and (C) insects.

Figure 2. Joint distributions of number of predicted CDS (normalized by number of NCBI predictions) over median predicted CDS length (normalized by median NCBI CDS length) for (A) mammals, (B) monocots. Dotted lines indicated equivalence to NCBI annotation, such that methods that are closest to the intersection of those lines best approximate CDS length and number of NCBI annotations. For dipterans, birds and rosids, see Figure S4.

Figure 3. Proportion of predicted protein-coding genes that fall entirely outside the intervals for NCBI protein-coding genes, organized by species and method.

Figure 4. Decrease in Gini index, an estimator of variable importance for random forest predictions of NCBI intergenic versus NCBI genic region location of predicted transcripts for (A) *H. sapiens* and (B) *Z. mays*. For additional reference taxa, see Figure S7.

Figure 5. Frequency of gene fusions by species and method. Fusions are defined as when individual transcripts overlap the CDS of multiple NCBI genes, different transcripts from the same predicted gene each have overlaps to different genes, or a combination of both of these. Frequencies are calculated after filtering out NCBI-annotated fusion events.

Figure 6. (A) By species and method, proportion of predicted proteins with BLASTP hit to NCBI proteins of species in taxonomic group. For predicted proteins with BLASTP hits, proportion of NCBI best hit target covered by amino acid matches with the predicted protein for (B) *H. sapiens* and (C) *Z. mays,* broken into proteins that are complete (start and stop codons present, no internal stop codons), and those that are not. Benjamini-Hochberg adjusted p-values for Wilcoxon rank-sum tests p≤0.05 indicated by *.

Figure 7. By species and method, (A) proportions of predicted genes for which TPM < 1 , and (B) RNA-seq read alignment rates to predicted transcripts. For Stringtie and Scallop, UTR intervals have been removed, such that all annotations are for CDS only. Xs denote rates for the NCBI annotations (also excluding UTR intervals).

Figure 8. Evidence for undetected CDS in predicted UTR intervals for Stringtie and Scallop. Increasing percentage of RNA-seq assembler transcripts with UTRs that have a BLASTX hit to the NCBI protein database of the same species, as a function of an increase in the assembler UTR-to-CDS ratio relative to that for NCBI annotations.

Figure 9. Bi-plots of median sample RNA-seq alignment rate versus BUSCO score for individual base annotation methods and subsequent integrations for (A) Stringtie$_{STAR}$ as the base, with subsequent integration of TOGA and BRAKER$_{protein}$ (B) same as B but with TOGA as the base annotation, (C) the absence of a closely related high quality genome annotation, precluding the use of TOGA, with Stringtie$_{STAR}$ set as the base annotation, and (D) no RNA-seq data, with TOGA as the base annotation, and an integration of BRAKER$_{protein}$. Integration involves successively adding genes from one annotation that fall entirely outside of the annotation to which those genes are being added.

Figure S1. Percent of recovered BUSCOs that overlap between methods for species in addition to Figure 2. (A-D) mammals, (E-G) dipterans, (H-J) birds, (K-M) rosids, and (N-Q) monocots.

G

H

I

CDS length

J

Figure S2. CDS length distributions for annotation methods and NCBI benchmark for (A) *H.* sapiens, (B) C*. familiaris*, (C) *G.* gallus, (D) *A. platyrhynchos,* (E) *Z. mays* (F) *O. sativa*, (G) *A. thaliana*, (H) *C. rubella*, (I) *D. melanogaster*, and (J) *D. pseudoobscura*.

A

B

C

Figure S3. Joint distributions of number of predicted CDS (normalized by number of NCBI predictions) over median predicted CDS length (normalized by median NCBI CDS length) for (A) dipterans, (B) rosids, and (C) birds. Dotted lines indicated equivalence to NCBI annotation, such that methods that are closest to the intersection of those lines best approximate CDS length and number of NCBI annotations.

Figure S4. Out-of-bag error rates for random forest models classifying CDS transcripts as either overlapping NCBI gene intervals, or falling outside of those intervals, i.e. intergenic. Models are generated for the reference species for each taxonomic group for which annotations are thought to be complete or nearly so.

Figure S5. For reference species, out-of-bag error rates broken down by predicted class, for random forest predictions of CDS overlapping NCBI protein-coding genes, vs. those entirely outside of NCBI protein coding gene intervals.

Figure S6. Random forest confusion matrices for reference species, for prediction accuracy of models predicting the genic versus intergenic status of CDS transcripts based upon transcript features.

Figure S7. Decrease in Gini index, an estimator of variable importance for random forest predictions of intergenic versus genic region location of predicted transcripts for additional reference species displayed in Figure 4.

Figure S8. Frequency of gene fusions, broken down by fusion type for five reference species Fusions are defined as when individual transcripts overlap the CDS of multiple NCBI genes (fusion transcripts), different transcripts from the same predicted gene each have overlaps to different genes (1 tscript-per-gene, multi-gene), or a combination of both of these (mixed). Frequencies are calculated after filtering out NCBI-annotated fusion events.

Figure S9. Grouped by species and method, percentage of predicted proteins that are complete, having both initiating start and terminating stop codons, and no internal stop codons.

Figure S10. For predicted proteins with BLASTP hits, proportion of NCBI best hit target covered by amino acid matches with the predicted protein for (A) *G. gallus*, (B) *D. melanogaster*, and (C) *A. thaliana,* broken into proteins that are complete (start and stop codons present, no internal stop codons), and those that are not. Benjamini-Hochberg adjusted p-values for Wilcoxon rank-sum tests, p≤0.05 indicated with *.

E

Figure S11. Empirical cumulative distributions of TPM for (A) *H. sapiens*, (B) *G. gallus*, (C) *A. thaliana*, (D) *Z. mays*, and (E) *D. melanogaster.* 0.0001 is added to all TPM values before log-transformation. Dashed line indicates TPM of 1.

Figure S12. Comparison of RNA-seq alignment rates between training data used in annotations, and test data for the five reference species. (A) Pairwise plots of test again training rates, and (B) Rate differences by method a species.

Figure S13. By method and species, RNA-seq read alignment rates to predicted transcripts, analogous to Figure 11, but for Stringtie and Scallop UTR intervals are included in the transcript predictions.

Figure S14. Ratios of the ratio of the median predicted UTR length/median predicted CDS length for RNA-seq assemblers over that for NCBI protein coding transcripts. No clear increase is observed for the most complete and curated annotations (*H. sapiens*, *D. melanogaster*, *G. gallus*, *Z. mays*, *A. thaliana*) relative to other genomes, indicating that the filtering out of cases of transcriptional readthrough (of the sort that NCBI will filter out for high quality annotations) does not explain the reduction in alignment rates for RNA-seq assemblers when UTRs are not included.

Figure S15. Comparison of TSEBRA integration to BRAKER_{protein} and BRAKER_{RNA} with respect to (A) BUSCO scores, (B) percentage of predicted proteins that have BLASTP hits to NCBI proteins for the species group, (C) the percentage of predicted genes that are intergenic relative to NCBI annotations, and (D) the RNA-seq alignment rates.

Figure S16. For *Homo sapiens*, Comparisons of (A) BUSCO scores, (B) the number of protein-coding genes with BLASTP hits to NCBI proteins from the mammal species group (C) the proportion of protein-coding genes with BLASTP hits, and (D) the median RNA-seq alignment rate across samples for individual methods and integrations of methods that start with Stringtie$_{STAR}$, then add TOGA, and finally BRAKER$_{protein}$. For each annotation that is added to the base annotation, integration involves adding genes that fall outside of the base annotation. For example, Stringtie+TOGA is built by adding TOGA genes that fall outside of Stringtie gene intervals.

Figure S17. For *Homo sapiens*, Comparisons of (A) BUSCO scores, (B) the number of protein-coding genes with BLASTP hits to NCBI proteins from the mammal species group (C) the proportion of protein-coding genes with BLASTP hits, and (D) the median RNA-seq alignment rate across samples for individual methods and integrations of methods that start with TOGA, then add Stringtie$_{STAR}$, and finally BRAKER$_{protein}$. For each annotation that is added to the base annotation, integration involves adding genes that fall outside of the base annotation. For example, TOGA+Stringtie is built by adding Stringtie genes that fall outside of TOGA gene intervals.

Figure S18. For *Homo sapiens*, Comparisons of (A) BUSCO scores, (B) the number of protein-coding genes with BLASTP hits to NCBI proteins from the mammal species group (C) the proportion of protein-coding genes with BLASTP hits, and (D) the median RNA-seq alignment rate across samples for individual methods and integrations of methods that use both RNA-seq and protein evidence but that don't include annotation transfer with TOGA. For each annotation that is added to the base annotation, integration involves adding genes that fall outside of the base annotation. For example, TOGA+Stringtie is built by adding Stringtie genes that fall outside of TOGA gene intervals.

Figure S19. For *Homo sapiens*, Comparisons of (A) BUSCO scores, (B) the number of protein-coding genes with BLASTP hits to NCBI proteins from the mammal species group (C) the proportion of protein-coding genes with BLASTP hits, and (D) the median RNA-seq alignment rate across samples for individual methods and integrations of methods that do not include RNA-seq Integration of BRAKER$_{protein}$ with TOGA involves adding BRAKER genes that fall outside of TOGA gene intervals.