CCDC113 stabilizes sperm axoneme and head-tail coupling apparatus to ensure male fertility

Bingbing Wu[1,2,3#], Chenghong Long[1#], Yuzhuo Yang[4,5], Zhe Zhang[4,5], Shuang Ma[1,2,3], Yanjie Ma[1,2,3], Huafang Wei[1], Jinghe Li[1], Hui Jiang[4,5], Wei Li[1,2,3*], Chao Liu[1,2,3*]

[1] Guangzhou Women and Children's Medical Center, Guangzhou Medical University , Guangdong Provincial Clinical Research Center for Child Health, Guangzhou, 510623, China

[2] State Key Laboratory of Stem Cell and Reproductive Biology, Institute of Zoology, Stem Cell and Regenerative Medicine Innovation Institute, Chinese Academy of Sciences, Beijing 100101, China.

[3] University of Chinese Academy of Sciences, Beijing 100049, China.

[4] Department of Urology, Department of Reproductive Medicine Center, Peking University Third Hospital, Beijing 100191, China.

[5] Department of Urology, Peking University First Hospital Institute of Urology, Peking University, Beijing 100034, China

[#] These authors contributed equally to this work.

[*]Corresponding Authors:

Dr. Wei Li

    Guangzhou Women and Children's Medical Center,

    No. 9 Jinsui Road, Tianhe District,

    Guangzhou 510623, China.

    Email: leways@gwcmc.org

Dr. Chao Liu

    Guangzhou Women and Children's Medical Center,

    No. 9 Jinsui Road, Tianhe District,

    Guangzhou 510623, China.

    Email: liuchao@gwcmc.org

Abstract

The structural integrity of the sperm is crucial for male fertility, defects in sperm head-tail linkage and flagellar axoneme are associated with acephalic spermatozoa syndrome (ASS) and the multiple morphological abnormalities of the sperm flagella (MMAF). Notably, impaired head-tail coupling apparatus (HTCA) often accompanies defects in the flagellum structure, however, the molecular mechanisms underlying this phenomenon remain elusive. Here, we identified an evolutionarily conserved coiled-coil domain-containing (CCDC) protein, CCDC113, and found the disruption of CCDC113 produced spermatozoa with disorganized sperm flagella and HTCA, which caused male infertility. Further analysis revealed that CCDC113 could bind to CFAP57 and CFAP91, and function as an adaptor protein for the connection of radial spokes, nexin-dynein regulatory complex (N-DRC) and doublet microtubules (DMTs) in the sperm axoneme. Moreover, CCDC113 was identified as a structural component of HTCA, collaborating with SUN5 and CENTLEIN to connect sperm head to tail during spermiogenesis. Together, our studies reveal that CCDC113 serve as a critical hub for sperm axoneme and HTCA stabilization, providing insights into the potential pathogenesis of infertility associated with human *CCDC113* mutations.

Keywords: CCDC113, male infertility, HTCA formation, axoneme stabilization, radial spoke

Introduction

Male fertility relies on the continuous production of spermatozoa through a complex developmental process known as spermatogenesis. Spermatogenesis involves three primary stages: spermatogonia mitosis, spermatocyte meiosis, and spermiogenesis. During spermiogenesis, spermatids undergo complex differentiation processes to develop into spermatozoa, which includes nuclear elongation, chromatin remodeling, acrosome formation, cytoplasm elimination, and flagellum development (Hermo et al., 2010). The integrity of spermatozoa is essential for their migration through the female reproductive tract and subsequent successful fertilization. An intact spermatozoon contains properly formed sperm head, head-tail coupling apparatus (HTCA), and flagellum (Parker, 2020; Roldan, 2019). Numerous components have been identified in the sperm HTCA and flagellum that are essential for the sperm integrity (Lehti & Sironen, 2017; Wu et al., 2020). Defects in sperm flagellum and HTCA can lead to reduced sperm motility or abnormal sperm morphology, termed as multiple morphological abnormalities of the sperm flagella (MMAF) or acephalic spermatozoa syndrome (ASS), which, in turn, causes male infertility (Sudhakar et al., 2021; Tu et al., 2020).

The sperm flagellum possesses an evolutionarily conserved axonemal structure composed of "9+2" microtubules, specifically, nine peripheral doublet microtubules (DMTs) surrounding two central microtubules known as the central pair (CP) (Inaba & Mizuno, 2016). Axonemal dyneins, radial spokes (RS), and the nexin-dynein regulatory complex (N-DRC) are arranged on DMTs with a 96-nm repeating unit structures (Kumar & Singh, 2021). Within the axoneme, the N-DRC and RS are crucial for maintaining axonemal integrity, forming crossbridges between adjacent DMTs and linking the DMTs to the central apparatus, respectively (Canty et al., 2021; Ishikawa, 2017; Kumar & Singh, 2021). Recent advancements in artificial intelligence, biochemical techniques, and cryo-electron microscopy (cryo-EM) facilitated the analysis of the axonemal structures, revealing numerous components among RS, N-DRC and DMTs that may serve as hubs for axoneme stabilization (Bazan et al., 2021; Leung et al., 2023; Walton et al., 2023; Zhou et al., 2023). For example, CFAP91 has been identified as a

protein that extends from the base of RS2 through the N-DRC base plate to RS3, thus

stabilizing RS2 and RS3 on the DMTs (Bicka et al., 2022; Dymek et al., 2011; Gui et al.,

2021). Similarly, CFAP57 extends through the N-DRC and interacts with RS3 via its C-

terminal region (Ghanaeian et al., 2023). Recent analyses suggest that CCDC96 and

CCDC113 may form a complex that extends parallel to the N-DRC, connecting the base of

RS3 to the tail of dynein g (IDA g) and the N-DRC (Bazan et al., 2021; Ghanaeian et al.,

2023). However, the functions of these proteins in stabilizing the sperm flagellum remain

unknown.

The sperm flagellum is tightly anchored to the sperm head through the HTCA, a complex

structure based on the centrosome (Wu et al., 2020). This structure consists of two cylindrical

microtubule-based centrioles and associated components, including well-organized

segmented columns, capitulum plate and basal plate. The segmented columns and capitulum

plate, located below the basal plate, are thought to originate from the dense material

emanating from the proximal centriole (Fawcett & Phillips, 1969; Zamboni & Stefanini,

1971). Many proteins have been identified in the sperm HTCA, with mouse models exhibiting

phenotypes characteristic of acephalic spermatozoa syndrome (Wu et al., 2020). SPATA6 is

the first protein identified as a component of the HTCA using the knockout mouse model, and

is crucial for its formation (Yuan et al., 2015). Deficiencies in SUN5 (Elkhatib et al., 2017;

Fang et al., 2018; Liu et al., 2020; Sha et al., 2018; Shang et al., 2018; Shang et al., 2017;

Xiang et al., 2022; Zhang et al., 2021; Zhu et al., 2016) and PMFBP1 (Deng et al., 2022; Liu

et al., 2020; Liu et al., 2021; Lu et al., 2021; Nie et al., 2022; Sha et al., 2019; Zhu et al., 2018)

have been associated with ASS in both humans and mice. The centriole-related protein

CENTLEIN acts as a bona fide linker between SUN5 and PMFBP1, participating in the

HTCA assembly (Zhang et al., 2021). Notably, impaired HTCA often coincides with defects

in the sperm flagellum (Hall et al., 2013; Shang et al., 2017; Yuan et al., 2015; Zhang et al.,

2021; Zhu et al., 2018), suggesting that the stabilization of the HTCA may be closely

associated with the integrity of sperm flagellum. However, the mechanism that maintains the

stabilization of both the sperm flagellum and the HTCA remains to be clarified.

Here, we identified an evolutionarily conserved coiled-coil domain-containing (CCDC)

protein, CCDC113, and found that it forms a complex with CFAP57 and CFAP91, thereby

facilitating the connection of RS, N-DRC and DMTs in the axoneme. Knockout of *Ccdc113*

resulted in spermatozoa with flagellar defects and head-tail linkage detachment, leading to

male infertility. Ultrastructural analysis showed that the loss of CCDC113 disrupted both the

sperm axoneme and HTCA. CCDC113 localizes on the manchette, HTCA and flagellum in

elongating and elongated spermatids. Further analysis revealed that CCDC113 is

indispensable for the connection of CFAP91 and DRC2 with DMTs in the sperm axoneme,

and it interacts with SUN5 and CENTLEIN to stabilize the sperm HTCA. These results

suggest that CCDC113 serves as a critical hub in maintaining the structural integrity of both

the sperm flagellum and HTCA.

Results

CCDC113 complexes with CFAP57 and CFAP91

CCDC113 is an evolutionarily conserved coiled-coil domain-containing (CCDC) protein

identified in the ciliated species. Comparative analysis of CCDC113 structures from

Tetrahymena thermophila to *Homo sapiens* showed structural similarity among CCDC113

orthologs (Figure 1A). Recent cryo-EM analysis in the structure of the 96-nm modular repeats

of axonemes from the *Tetrahymena thermophila* cilia and human respiratory cilia revealed

that CCDC113 localizes to the linker region among RS, N-DRC and DMTs (Figure 1B),

suggesting it may serve as a structural component connecting RS, N-DRC and DMTs (Bazan

et al., 2021; Ghanaeian et al., 2023). To further investigate, we examined the interactions

between CCDC113 and its neighboring axoneme-associated proteins, CFAP57 and CFAP91

(Figure 1B). We transfected HEK293T cells with a GFP-tagged CCDC113 and FLAG-tagged

CFAP57 or CFAP91, then performed anti-FLAG-immunoprecipitations. CCDC113 was

present in both FLAG-CFAP57 and FLAG-CFAP91 immunoprecipitates (Figure 1C and D),

indicating CCDC113 interacts with both CFAP57 and CFAP91. Given that CFAP91 has been

reported to stabilize RS on the DMTs (Bicka et al., 2022; Dymek et al., 2011; Gui et al., 2021)

and cryo-EM analysis shows that CCDC113 is closed to DMTs, we speculated that CCDC113

may connect RS to DMTs by binding to CFAP91 and microtubules. To test this, we detected

the interaction between CCDC113 and β-tubulin (TUBB5) and found that CCDC113 was

present in MYC-TUBB5 immunoprecipitate (Figure 1E). Additionally, since CFAP57

extends through the N-DRC and CCDC113 is closed to the N-DRC (Ghanaeian et al., 2023),

we further examined the interaction between CCDC113 and N-DRC components adjacent to

DMTs. Co-immunoprecipitation (co-IP) analysis showed that CCDC113 could bind to DRC1,

DRC2, and DRC3 (Figure 1F-H). We also included two sperm outer dense fiber proteins,

ODF1 and ODF2 (Zhu et al., 2022), which are located far from the sperm axoneme, as

negative controls in the co-IP experiments. As shown in Figure 1-figure supplement 1A and

1B, neither ODF1 nor ODF2 bound to CCDC113. Therefore, CCDC113 may function as an

adaptor protein connecting RS, N-DRC, and DMTs, serving as a critical hub for axoneme

stabilization.

CCDC113 is required for male fertility

To investigate the physiological functions of CCDC113, we generated a *Ccdc113* knockout

mouse strain using the CRISPR/Cas9 system (Figure 2A). The *Ccdc113$^{-/-}$* mice were

genotyped by genomic DNA sequencing and further confirmed by polymerase chain reaction

(PCR). Genotypes were distinguished by a 539 bp band for *Ccdc113$^{+/+}$* mice, a 461 bp band

for *Ccdc113$^{-/-}$* mice, and two bands of 539 bp and 461 bp for *Ccdc113$^{+/-}$* mice (Figure 2B).

Further immunoblotting analysis confirmed the elimination of CCDC113 in total protein

extracts from *Ccdc113$^{-/-}$* testes (Figure 2C; Figure 2-figure supplement 1A). Mice lacking

Ccdc113 showed no gross abnormalities in their appearance or behavior, and no obvious

differences in body weight (Figure 2H). Additionally, no hydrocephalus or left-right

asymmetry defects were observed (Figure 2-figure supplement 1B). Additionally, the

deficiency of CCDC113 did not affect ciliogenesis in the lung and trachea (Figure 2-figure

supplement 1C-E). We then assessed the fertility of 2-month-old male and female *Ccdc113$^{-/-}$*

mice. *Ccdc113$^{-/-}$* female mice were able to produce offspring after mating with wild-type

(WT) adult males, similar to *Ccdc113$^{+/+}$* female mice (Figure 2D). However, *Ccdc113$^{-/-}$* male

mice exhibited normal mating behavior, as indicated by the presence of copulatory plugs, they

failed to produce offspring when mating with WT adult female mice (Figure 2E). Thus, the

knockout of *Ccdc113* results in male infertility.

***Ccdc113* knockout mice produce spermatozoa with flagellar defects and head-tail**

linkage detachment

To further investigate the cause of male infertility, we initially examined *Ccdc113$^{-/-}$* testis at

both macroscopic and histological levels. *Ccdc113* knockout did not affect testis size (Figure

2F and G) or the ratio of testis weight to body weight (Figure 2H and I). Histological sections

stained with hematoxylin-eosin (H&E) revealed that seminiferous tubules of *Ccdc113$^{+/+}$* mice

exhibited a tubular lumen with flagella emerging from the developing spermatids. In contrast,

flagellar staining appeared reduced in *Ccdc113$^{-/-}$* seminiferous tubules (Figure 2J, red

asterisks). Immunofluorescence staining for acetylated tubulin (ac-tubulin), a marker for

sperm flagellum (Martinez et al., 2020), further confirmed the flagellar defects in *Ccdc113$^{-/-}$*

mice (Figure 2K and L).

Subsequently, we examined spermatids at different stages in *Ccdc113$^{-/-}$* testes using periodic

acid Schiff (PAS) staining. Pioneering work in the mid-1950s used the PAS stain in histologic

sections of mouse testis to visualize glycoproteins of the acrosome and Golgi in seminiferous

tubules (Oakberg, 1956). The pioneers discovered in cross-sectioned seminiferous tubules the

association of differentiating germ cells with successive layers to define different stages that

in mice are twelve, indicated as Roman numerals (XII). For each stage, different associations

of maturing germ cells were always the same with early cells in differentiation at the

periphery and more mature cells near the lumen. In this way, progressive differentiation from

stem cells to mitotic, meiotic, acrosome-forming, and post-acrosome maturing spermatocytes

was mapped to define spermatogenesis with the XII stages in mice representing the

seminiferous cycle. The maturation process from acrosome-forming cells to mature

spermatocytes is defined as spermiogenesis with 16 different steps that are morphologically

distinct spermatids (O'Donnell L, 2015). While acrosome biogenesis and nuclear morphology

in $Ccdc113^{-/-}$ spermatids from steps 1 to 10 were comparable to those in $Ccdc113^{+/+}$

spermatids, abnormal club-shaped heads were observed in spermatids from steps 11 to 16 in

$Ccdc113^{-/-}$ mice (Figure 3-figure supplement 1A, black asterisk). In addition, the manchette

of $Ccdc113^{-/-}$ spermatids was more elongated compared to that of $Ccdc113^{+/+}$ spermatids

(Figure 3-figure supplement 1B). Therefore, the disruption of CCDC113 impaired

spermiogenesis.

Next, we examined the spermatozoa in the cauda epididymis and found that the sperm count

in the $Ccdc113^{-/-}$ cauda epididymis was significantly decreased compared to the control group

(Figure 3A and B). The motility of the released spermatozoa from $Ccdc113^{+/+}$ and $Ccdc113^{-/-}$

cauda epididymis showed that $Ccdc113^{-/-}$ spermatozoa were completely immotile (Figure 3C).

H&E staining of the cauda epididymis showed fewer hematoxylin-stained sperm heads in the

$Ccdc113^{-/-}$ cauda epididymis compared to the $Ccdc113^{+/+}$ cauda epididymis. Notably, unlike

the control group, which exhibited linear eosin staining in the epididymal lumen, the

$Ccdc113^{-/-}$ mice showed numerous coiled eosin-stained structures without sperm heads in the

epididymal lumen (Figure 3A, red circles). To determine the morphological characteristics of

the spermatozoa, we conducted single-sperm immunofluorescence using an anti-α/β-tubulin

antibody to label the sperm flagellum and lectin peanut agglutinin (PNA) to visualize the

sperm acrosome (Nakata et al., 2015). We noticed that $Ccdc113^{-/-}$ spermatozoa showed severe

morphological malformations, including sperm head-tail detachment (type 1), abnormal

sperm head with curly tail (type 2), normal sperm head with curly tail (type 3) (Figure 3D and

E). To determine the role of CCDC113 in flagellum assembly, we analyzed flagellum

formation during spermiogenesis. We found that in $Ccdc113^{-/-}$ mice, diffuse and curly

axonemal signals were observed in testicular germ cells starting from the round spermatid

stage (Figure 3F). Overall, these findings suggest that the deletion of CCDC113 leads to

sperm flagellum deformities and detachment of the sperm head-to-tail linkage, resulting in a

unique type of acephalic spermatozoa, which may be responsible for the $Ccdc113^{-/-}$ male

infertility.

CCDC113 localizes on the sperm neck and flagellum regions

To gain further insights into the functional role of CCDC113 during spermiogenesis, we examined its expression in different tissues and found that CCDC113 was predominantly expressed in mouse testis (Figure 4-figure supplement 1A and B). CCDC113 was first detected in testis at postnatal day 7 (P7), with expression levels increasing continuously from P21 onward, peaking in adult testes (Figure 4A), suggesting that CCDC113 is highly expressed throughout spermiogenesis. We then conducted immunofluorescence analysis of CCDC113 in *Ccdc113$^{+/+}$* and *Ccdc113$^{-/-}$* germ cells to determine its precise localization during spermatogenesis. CCDC113 appeared as punctate signals near the nuclei of spermatocyte and round spermatids, and localized to the manchette, sperm neck and flagellum regions in elongating and elongated spermatids (Figure 4-figure supplement 1C).

To further validate these findings, we co-stained CCDC113 with α/β-tubulin, which marks the manchette and flagellum in spermatids (Lehti & Sironen, 2016) (Figure 4B). The immunofluorescence analysis showed that CCDC113 localized to the manchette surrounding the spermatid head from step 9 to step 14, as well as to the sperm neck and flagellum in the testes (Figure 4B). Given that CCDC113 was initially identified as a component of centriolar satellites (Firat-Karalar et al., 2014), the punctate signals of CCDC113 in spermatocyte and spermatids may be localized around the centrosome. To test this, we performed the immunofluorescent staining of CCDC113 and centrosomal protein CENTRIN1/2 in spermatocytes and spermatids, and found that the signal of CCDC113 partially colocalized with CENTRIN1/2 (Figure 4C). Thus, CCDC113 is localized to the centrosome, manchette, sperm neck and flagellum regions in the developing germ cells.

Next, we examined the localization of CCDC113 in mature spermatozoa, and found that CCDC113 was localized in the sperm neck and flagellum regions (Figure 4D). Similar localization of CCDC113 was also observed in human mature spermatozoa (Figure 4E). The consistent localization of CCDC113 at the sperm neck and flagellum in mature spermatozoa suggests its importance for maintaining the integrity of the sperm flagellum and head-to-tail connection.

Ccdc113 knockout results in the disorganization of the sperm flagellum structures

To delineate the sperm flagellar defects in $Ccdc113^{-/-}$ mice, we conducted transmission

electron microscopy (TEM) examination of longitudinal sections of $Ccdc113^{-/-}$ spermatozoa.

TEM analysis revealed a significant presence of unremoved cytoplasm, including disrupted

mitochondria, damaged axonemes, and large vacuoles in $Ccdc113^{-/-}$ spermatozoa (Figure 5A,

red asterisks). Cross sections of the principal piece of $Ccdc113^{-/-}$ spermatozoa further

revealed partial loss or unidentifiable "9+2" axonemal structures, along with the disruption of

the fibrous sheath and outer dense fibers (Figure 5A). We further examined the axonemal

structure in $Ccdc113^{-/-}$ testicular spermatids using TEM and found disorganized axonemal

microtubules were detected in $Ccdc113^{-/-}$ testicular spermatids (Figure 5B). In contrast to the

regular positioning of the CP and nine peripheral DMTs in the $Ccdc113^{+/+}$ spermatid axoneme,

the $Ccdc113^{-/-}$ spermatids exhibited a scattered arrangement of DMTs, and no distinct radial

spokes were observed (Figure 5B, red arrowheads). These results indicate CCDC113 is

essential for the integrity of sperm flagellum.

CCDC113 has been shown to localize at the base of the RS3 and interact with adjacent

axoneme-associated proteins (Figure 1B-H). Given the disorganized "9+2" axonemal

structure was detected in cross-sectioned $Ccdc113^{-/-}$ flagellar specimen (Figure5A and B), we

speculated that CCDC113 likely served as an adaptor to connect the neighboring axoneme-

associated proteins to DMTs. To test this, we examined the flagellar localization of CFAP91

in $Ccdc113^{-/-}$ spermatozoa, which is positioned in close proximity to CCDC113 at the root

region of RS3 and is critical for the localizations of calmodulin-associated and spoke-

associated complex (CSC) proteins CFAP61 and CFAP251 (Bicka et al., 2022; Meng et al.,

2024). Immunofluorescence results indicated that the absence of *Ccdc113* leads to the

abnormal distribution of CFAP91 on the axoneme, where CFAP91 could not colocalize with

DMTs. (Figure 5C, white asterisks, E and F). Given that DRC2 serves as the core component

of the axonemal N-DRC (Jreijiri et al., 2023) and CCDC113 could bind to DRC2 (Figure 1G),

we further examined DRC2 localization in the $Ccdc113^{+/+}$ and $Ccdc113^{-/-}$ spermatozoa.

Immunofluorescence analysis showed that DRC2 exhibited distinct signals that did not

colocalize with DMTs of $Ccdc113^{-/-}$ spermatozoa (Figure 5D, white asterisks, G and H).

These findings collectively indicate that CCDC113 is indispensable for the connection of

CFAP91 and DRC2 to the DMTs, which is required for structural integrity of the sperm

axoneme.

$Ccdc113$ knockout impairs head-to-tail anchorage of the spermatids

To explore the mechanism behind acephalic spermatozoa in $Ccdc113$ knockout mice, we first

examined where the flagellum detached from sperm head in $Ccdc113^{-/-}$ mice. The proportion

of decapitated tails in the caput, corpus and cauda of $Ccdc113^{-/-}$ epididymis was similar

(Figure 6A), suggesting the separation of the sperm head and tail in $Ccdc113^{-/-}$ mice may

occur either within the seminiferous tubules or upon entering the caput of the epididymis. To

confirm this, we performed PAS staining to examine spermiogenesis stages in $Ccdc113^{+/+}$ and

$Ccdc113^{-/-}$ testes (Figure 6B). We found that in $Ccdc113^{+/+}$ testes, sperm heads at stages VII-

VIII were oriented towards the basal membrane, whereas in $Ccdc113^{-/-}$ testes, sperm heads

were oriented towards the tubule lumen during these stages (Figure 6C, arrows and D), which

may be due to the separation of sperm heads from the flagellum during spermiogenesis.

Additionally, mature sperm heads were still present at stages IX–X in $Ccdc113^{-/-}$ testes,

whereas mature spermatozoa were released into the lumen of the seminiferous tubule at stage

VIII in $Ccdc113^{+/+}$ testes (Figure 6B, red asterisk). These results suggest that the sperm head

and flagellum separation may occur during spermiation in the $Ccdc113^{-/-}$ testes.

Next, we examined the development of the HTCA in $Ccdc113^{+/+}$ and $Ccdc113^{-/-}$ spermatids

using TEM. In $Ccdc113^{+/+}$ step 9–11 spermatids, the well-defined coupling apparatus,

comprising the basal plate, capitulum plate, segmented columns, proximal centriole, distal

centriole, was tightly attached to the sperm head. However, in $Ccdc113^{-/-}$ step 9–11

spermatids, the abnormal HTCA was detached from the sperm head (Figure 6E, red asterisk).

Further observation of the HTCA structure revealed the absence of segmented columns and

capitulum plate; only dense material surrounding the proximal centriole and basal plate could

be detected (Figure 6E, white arrow). The basal plates were abnormally distant from their

native implantation site on the nucleus of *Ccdc113⁻/⁻* elongating and elongated spermatids

(Figure 6E). Taken together, our results indicate that the disruption of *Ccdc113* causes the

destroyed coupling apparatus detachment from the sperm head during spermiogenesis, and

CCDC113 is required for the integrity of the sperm HTCA.

CCDC113 cooperates with SUN5 and CENTLEIN to stabilize sperm HTCA

To elucidate the molecular function of CCDC113 in sperm head-tail linkage, we examined its

interaction with known HTCA-associated proteins, including SUN5, CENTLEIN, PMFBP1,

and SPATA6 (Shang et al., 2017; Yuan et al., 2015; Zhang et al., 2021; Zhu et al., 2018).

GFP-tagged CCDC113 and FLAG-tagged HTCA-associated proteins were co-transfected into

HEK293T cells, followed by the immunoprecipitation with anti-GFP antibody. We found that

CCDC113 interacted with SUN5 and CENTLEIN, but not with PMFBP1 and SPATA6

(Figure 7A-C and F). We further conducted co-IP assays in the reverse direction and verified

that both SUN5 and CENTLEIN could bind to CCDC113 (Figure 7D and E). To investigate

the localization of CCDC113 within the HTCA, we co-stained mature spermatozoa with

antibodies against CCDC113 and SUN5, which has been reported to be localized at the root

connecting the HTCA to the nuclear envelope (Shang et al., 2017; Zhang et al., 2021).

CCDC113 was found positioned below SUN5, showed partial overlap with it (Figure 7G).

The centriolar protein CENTLEIN was localized to the HTCA and served as the critical linker

protein between SUN5 and PMFBP1 in the elongating and elongated spermatid (Zhang et al.,

2021)). Given that CENTLEIN disappears in mature spermatozoa (Zhang et al., 2021), we

performed the immunofluorescent staining of CCDC113 and CENTLEIN in elongated

spermatids and found that CCDC113 partially colocalized with CENTLEIN at the HTCA

(Figure 7H). Thus, CCDC113 interacts with both SUN5 and CENTLEIN and localizes at the

sperm HTCA.

The HTCA localization of CCDC113 may be responsible for maintaining HTCA integrity. To

test this hypothesis, we examined the localization of classical HTCA component SPATA6 in

Ccdc113⁻/⁻ and *Ccdc113⁺/⁺* spermatozoa using immunofluorescent staining. We found that

SPATA6 was not attached to the implantation fossa of the sperm nucleus in *Ccdc113⁻/⁻*

spermatozoa (Figure 7I and J), indicating that CCDC113 is essential for the integrity of the

sperm HTCA. To further elucidate the functional relationships among CCDC113, SUN5,

CENTLEIN, and PMFBP1 at the sperm HTCA, we examined the localization of CCDC113 in

Sun5⁻/⁻, *Centlein⁻/⁻*, and *Pmfbp1⁻/⁻* spermatozoa. Compared to the control group, CCDC113

was predominantly localized on the decapitated flagellum in *Sun5⁻/⁻*, *Centlein⁻/⁻*, and *Pmfnp1⁻*

⁻ spermatozoa (Figure 7K and L), indicating SUN5, CENTLEIN, and PMFBP1 are crucial for

the proper docking of CCDC113 to the implantation site on the sperm head. Taken together,

these data demonstrate that CCDC113 cooperates with SUN5 and CENTLEIN to stabilize the

sperm HTCA and anchor the sperm head to the tail.

Discussion

CCDC113 is a highly evolutionarily conserved component of motile cilia/flagella. Studies in

the model organism, *Tetrahymena thermophila*, have revealed that CCDC113 connects RS3 to

dynein g and the N-DRC, which plays essential role in cilia motility (Bazan et al., 2021;

Ghanaeian et al., 2023). Recent studies have also identified the localization of CCDC113

within the 96-nm repeat structure of the human respiratory epithelial axoneme, and localizes

to the linker region among RS, N-DRC and DMTs (Walton et al., 2023). In this study, we

reveal that CCDC113 is indispensable for male fertility, as *Ccdc113* knockout mice produce

spermatozoa with flagellar defects and head-tail linkage detachment (Figure 3D). CCDC113

is localized to the sperm neck and flagellum regions in the elongating and elongated

spermatids. In the sperm flagellum, CCDC113 interacts with both CFAP57 and CFAP91,

serving as an adaptor protein to connect RS, N-DRC, and DMTs, thereby stabilizing sperm

flagellum. At the sperm head to tail connecting piece, CCDC113 binds to SUN5 and

CENTLEIN to stabilize sperm HTCA and anchor the sperm head to the tail. Thus, CCDC113

is essential for maintaining the integrity of both the sperm axoneme and the sperm HTCA.

Moreover, TEM analysis detected excess residual cytoplasm in spermatozoa, including

disrupted mitochondria, damaged axonemes, and large vacuoles, indicating defects in

cytoplasmic removal in *Ccdc113*$^{-/-}$ mice (Figure 5A).

Recent cryo-EM analysis of the axonemes from *Tetrahymena thermophila* cilia and human

respiratory cilia revealed that CCDC113 is localized at the linker region between RS3, N-

DRC and DMTs (Bazan et al., 2021; Ghanaeian et al., 2023). We also found that CCDC113

interacts with adjacent axoneme-associated proteins, CFAP57 and CFAP91 (Figure 1C and

D). CFAP57 has been identified as the adaptor protein responsible for assembling dynein g

and d, and it can interact with both N-DRC and RS3 (Ghanaeian et al., 2023; Ma et al., 2023).

Previous studies have demonstrated that CFAP91 extends from the base of RS2 through the

N-DRC base plate to RS3, playing a crucial role in stabilizing and localizing RS2 and RS3 on

the DMT (Bicka et al., 2022; Dymek et al., 2011; Gui et al., 2021). The CFAP91 ortholog,

FAP91, interacts with three N-DRC subunits (DRC1, DRC2, and DRC4), facilitating the

docking of the N-DRC in *Chlamydomonas* (Gui et al., 2021). In humans, pathogenic

mutations in CFAP91 and DRC2 disrupt sperm flagellum structure and result in MMAF

(Jreijiri et al., 2023; Martinez et al., 2020). TEM and immunostaining experiments in

spermatozoa showed severe CP and radial spokes defects in *CFAP91* mutant patients

(Martinez et al., 2020). In this study, we found that the absence of CCDC113 results in severe

axonemal disorganization characterized by defective RSs, scattered DMTs, and misplaced CP.

Further analysis demonstrated that the CCDC113 deficiency disrupts the localization of

CFAP91 and DRC2 on DMTs. Thus, CCDC113 may function as an adaptor protein to

stabilize CFAP91 and DRC2 on DMTs, facilitating the docking of RS and N-DRC to DMTs

and thereby maintaining the integrity of sperm axoneme.

Recent analyses have revealed that certain centrosomal proteins play crucial roles in the

assembly and maintenance of sperm HTCA (Avasthi et al., 2013; Liska et al., 2009; Zhang et

al., 2021). In early round spermatids, the centriole pair initially localizes to the caudal nuclear

pole and expands the electron-dense material, some of which exhibits striation (Wu et al.,

2020). As spermatids develop, the dense material around the centrioles gradually transforms

into a well-organized structure, clearly identified as the basal plate, capitulum plate and

segmented columns (Dooher & Bennett, 1973; Wu et al., 2020). CCDC113 was initially

identified through the isolation of centriolar proteins from bovine sperm (Firat-Karalar et al.,

2014). During spermiogenesis, CCDC113 colocalizes with CENTRIN1/2 at the centrosome in

developing spermatids and continues to localize to the sperm neck region in elongating and

elongated spermatids (Figure 4B and C). Additionally, CCDC113 can bind to HTCA

associated centrosomal protein, CENTLEIN, and the disruption of *Centlein* impairs the

attachment of CCDC113 to the sperm head. In *Ccdc113*$^{-/-}$ spermatids, the capitulum plate and

segmented columns are absent, and the basal plate is detached from the implantation site on

the nucleus of *Ccdc113*$^{-/-}$ elongating and elongated spermatids during spermiogenesis (Figure

6E). These observations raise the possibility that CCDC113 is an HTCA associated

centrosomal protein crucial for maintaining the structural integrity of the HTCA.

SUN5 is a transmembrane protein located in the nuclear envelope, and acts as the root

connecting the HTCA to the sperm nuclear envelope (Shang et al., 2017). CENTLEIN can

directly bind to both SUN5 and PMFBP1, serving as a linker between SUN5 and PMFBP1 to

maintain the integrity of HTCA (Zhang et al., 2021). CCDC113 can interact with SUN5 and

CENTLEIN, but not PMFBP1 (Figure 7A-C), and left on the tip of the decapitated tail in

Sun5$^{-/-}$ and *Centlein*$^{-/-}$ spermatozoa (Figure 7K and L). Furthermore, CCDC113 colocalizes

with SUN5 in the HTCA region, and immunofluorescence staining in spermatozoa shows that

SUN5 is positioned closer to the sperm nucleus than CCDC113 (Figure 7G and H). Therefore,

SUN5 and CENTLEIN may be closer to the sperm nucleus than CCDC113. PAS staining

revealed that *Ccdc113*$^{-/-}$ sperm heads are abnormally oriented in stages V–VIII seminiferous

epithelia (Figure6 C and D), and TEM analysis further demonstrated that the disruption of

CCDC113 causes the detachment of the destroyed coupling apparatus from the sperm head in

step 9–11 spermatids (Figure 6E). All these results suggest that the detachment of sperm head

and tail in *Ccdc113*$^{-/-}$ mice may not be a secondary effect of sperm flagellum defects.

Overall, we identified CCDC113 as a structural component of both the flagellar axoneme and

the HTCA, where it performs dual roles in stabilizing the sperm axonemal structure and

maintaining the structural integrity of HTCA. Given that the cryo-EM of sperm axoneme and

HTCA could powerfully strengthen the role of CCDC113 in stabilizing sperm axoneme and

head-tail coupling apparatus, it a valuable direction for future research. $Ccdc113^{-/-}$ mice did

not exhibit other ciliopathies, such as situs inversus, hydrocephalus, or abnormal ciliogenesis

of tracheal cilia (Figure 2-figure supplement 1B-E), which suggests that CCDC113 may

specifically function in spermiogenesis.

Materials and Methods

Phylogenetic analysis and structural similarity analysis

The amino acid sequence of CCDC113 of seven species were downloaded from UniProt. The

phylogenetic trees were constructed using MEGA 10.0 (Kumar et al., 2018) with the

Neighbor-Joining (NJ) method (Saitou & Nei, 1987). 3D structures of CCDC113 orthologs of

seven species were obtained from AlphaFold Protein Structure Database (Varadi et al., 2022).

Structural similarity Z scores were derived through the DALI webserver for all against all

structure comparison (Holm & Laakso, 2016).

Animals

The Sun5$^{-/-}$, $Pmfbp1^{-/-}$ and $Centlein^{-/-}$ mice have been reported previously (Shang et al., 2017;

Zhang et al., 2021; Zhu et al., 2018). The mouse $Ccdc113$ gene is 44.81 kb and contains 9

exons. Exon 2 to exon 6 of $Ccdc113$ were chosen as the target sites. The knockout mice were

generated using CRISPR-Cas9 system from Cyagen Biosciences. The gRNA and Cas9

mRNA were co-injected into fertilized eggs of C57BL/6J mice to generate $Ccdc113^{+/-}$ mice

with a 10215 bp base deletion and 64 bp insertion. The resulting heterozygotes were interbred,

and their offspring were genotyped by genomic DNA sequencing to identify wide-type and

homozygote mice. The genotyping primers for knockout were: F1: 5'-

TCAAATCATCACACCCTGCCTCT-3', R: 5'-GCTTGCACTCGGGTGATACATAA-3',

and for WT mice, the specific primers were: F2: 5'-CAGGTTCTCAACACCTACAAAGTA-

3', R: 5'-GCTTGCACTCGGGTGATACATAA-3'.

All the animal experiments were performed according to approved institutional animal care

and use committee (IACUC) protocols (# 08-133) of the Institute of Zoology, Chinese

Academy of Sciences.

Assessment of fertility

Fertility was assessed in 2-month-old male mice of various genotypes. Each male was paired

with two WT females (6–8 weeks old), and vaginal plugs were checked each morning.

Females with a vaginal plug were separated and housed individually, and their pregnancy

outcomes were recorded. If a female did not give birth by day 22 postcoitus, it was considered

not pregnant and euthanized for confirmation. Each male participated in six cycles of the

breeding assay with different females.

Human adult sperm sample preparation

The sperm donation candidates in this study were healthy young Chinese men. Each

participant underwent a thorough medical examination and completed an extensive medical

and social questionnaire to ensure the exclusion of individuals with genetic or significant

medical issues, as outlined in the Basic Standard and Technical Norms of Human Sperm Bank

published by the Chinese Ministry of Health. Individuals who smoked, abused drugs, or were

heavy drinkers were also excluded from the study. Those who remained eligible signed

informed consent forms for voluntary sperm donation and agreed to reside in Beijing for a

minimum of six months. The sperm bank documented each participant's age, date of birth,

and date of semen collection. Ethical approval for this study was granted by the Reproductive

Study Ethics Committee of Peking University Third Hospital (2017SZ-035). Semen samples

were processed using a 40% density gradient of PureSperm (Nidacon International, Molndal,

Sweden) through centrifugation at room temperature ($500 \times g$, 30 min) and washed three

times with phosphate-buffered saline (PBS). The obtained spermatozoa were utilized for

immunofluorescence staining.

Antibodies

Rabbit anti-CCDC113 generated from Dia-an Biotech (Wuhan, China) was diluted at 1:500

for western blotting and a 1: 25 dilution for immunofluorescence. Mouse anti-α-TUBULIN

antibody (AC012, Abclonal) was used at a 1:5000 dilution for western blotting. Mouse anti-

GFP antibody (M20004, Abmart) was used at a 1:2000 dilution for western blotting. Rabbit

anti-MYC antibody (BE2011, EASYBIO) was used at a 1:2000 dilution for western blotting.

Rabbit anti-FLAG antibody (20543-1 AP, Proteintech) was used at a 1:2000 dilution for

western blotting. Mouse anti-ac-TUBULIN antibody (T7451, Sigma-Aldrich) was used at a

1:200 dilution for immunofluorescence. Mouse anti-α/β-TUBULIN antibody (ab44928,

Abcam) was used at a 1: 100 dilution for immunofluorescence. Rabbit anti-DRC2 antibody

(NBP2-84617, Novus) was used at a 1:100 dilution for immunofluorescence. Rabbit anti-

CFAP91 antibody (bs-9823R, Bioss) was used at a 1:100 dilution for immunofluorescence.

In-house-generated mouse anti-SUN5 antibody targeting the SUN5 SUN domain (aa193–373)

was used at a 1:100 dilution for immunofluorescence analysis. Rat anti-CENTLEIN antibody

was generated by Absea Biotechnology Ltd (Beijing, China) was diluted at a 1:20 dilution for

immunofluorescence. Rabbit anti-SPATA6 antibody (11849 -1 AP, Proteintech) was used at a

1:100 dilution for immunofluorescence. The Alexa Fluor 488 conjugate of lectin PNA (1:400,

L21409, Thermo Fisher) was used for immunofluorescence. The secondary antibodies were

goat anti-rabbit FITC (1:200, ZF-0311, Zhong Shan Jin Qiao), goat anti-rabbit TRITC (1:200,

ZF-0316, Zhong Shan Jin Qiao), goat anti-mouse FITC (1:200, ZF-0312, Zhong Shan Jin

Qiao) and goat anti-mouse TRITC (1:200, ZF0313, Zhong Shan Jin Qiao).

Sperm motility and sperm count assays

The cauda epididymis was isolated from the male mice of the different genotypes. Sperm

were released in phosphate-buffered saline (PBS, Gibco, C14190500BT) from the incisions of

the cauda epididymis for 10 min at 37°C. And then the swim-up suspension was used for the

analysis of sperm motility with a microscope through a 20x phase objective. Viewing areas in

each chamber were imaged using a CCD camera. The samples were analyzed via computer-

assisted semen analysis (CASA) using the Minitube Sperm Vision Digital Semen Evaluation

System (12500/1300, Minitube Group, Tiefenbach, Germany) and were also analyzed by

CASA. The incubated sperm number was counted with a hemocytometer.

Histology staining

As previously reported (Wang et al., 2018), the testes and cauda epididymis were dissected

after euthanasia, and fixed with Bouin's fixative for 24 h at 4°C, then the testes were

dehydrated with graded ethanol and embedded in paraffin. For histological analysis, the 5 μm

sections were cut and covered on glass slides. Sections were stained with H&E and PAS for

histological analysis after deparaffinization.

Electron microscopy analysis

The cauda epididymis and testis were dissected and fixed in 2.5% (vol/vol) glutaraldehyde in

0.1 M cacodylate buffer at 4°C overnight. After washing in 0.1 M cacodylate buffer, samples

were cut into small pieces of approximately 1 mm^3, then immersed in 1% OsO4 for 1 h at 4°C.

Samples were dehydrated through a graded acetone series (50%, 60%, 70%, 80%, 90%, 95%,

100%) and embedded in resin (DDSA, NMA, enhancer, 812) for staining. Ultrathin sections

were cut and stained with uranyl acetate and lead citrate. Images were acquired and analyzed

using a JEM-1400 transmission electron microscope.

Scanning electron microscopy

The trachea was fixed in 2.5% glutaraldehyde solution overnight, and dehydrated in graded

ethanol, subjected to drying and coated with gold. The images were acquired and analyzed

using SU8010 scanning electron microscope.

Immunofluorescence staining

The testis albuginea was peeled and incubated with collagenase IV and hyaluronidase in PBS

for 15 min at 37°C, then washed twice with PBS. Next, fixed with 4% PFA for 5 min, and

then coated on slide glass to dry out. The slides were washed with PBS three times and then

treated with 0.5% Triton X-100 for 5 min, and blocked with 5% BSA for 30 min. Added the

primary antibodies and incubated at 4°C overnight, followed by incubating with a second

antibody and DAPI. The images were taken using LSM880 and SP8 microscopes.

The mouse testis was immediately dissected and fixed with 2% paraformaldehyde in 0.05%

PBST (PBS with 0.05% Triton X-100) at room temperature for 5 min. The fixed sample was

placed on a slide glass and squashed by placing a cover slip on top and pressing down. The

sample was immediately flash frozen in liquid nitrogen, and the slides were stored at −80°C

for further immunofluorescence experiments. After removing the coverslips, the slides were

washed with PBS three times and then treated with 0.1% Triton X-100 for 10 min, rinsed

three times in PBS, and blocked with 5% bovine serum albumin (Amresco, AP0027). The

primary antibody was added to the sections and incubated at 4°C overnight, followed by

incubation with the secondary antibody. The nuclei were stained with DAPI. The

immunofluorescence images were taken immediately using an LSM 780 microscope (Zeiss)

or SP8 microscope (Leica).

Spermatozoa were released from the cauda epididymis in PBS at 37°C for 15 min, then were

spread on glass slides for morphological observation or immunostaining. After air drying,

spermatozoa were fixed in 4% PFA for 5 min at room temperature, and slides were washed

with PBS three times and blocked with 5% BSA for 30 min at room temperature. The primary

antibodies were added to the sections and incubated at 4°C overnight, followed by incubation

with the secondary antibody. The nuclei were stained with DAPI and images were taken using

an LSM 880 microscope (Zeiss) or SP8 microscope (Leica).

Immunoprecipitation

Transfected cells were lysed in a lysis buffer (50mM HEPES, PH 7.4, 250mM NaCl, 0.1%

NP-40 containing PIC and PMSF) on ice for 30 min and centrifuged at 12000 rpm at 4°C

for 15 min, cell lysates were incubated with primary antibody about 12 hours at 4°C and then

incubated with protein A-Sepharose (GE, 17-1279-03) for 3 hours at 4°C, then washed three

times with lysed buffer and subjected to immunoblotting analysis.

Statistical analysis

All the experiments were repeated at least three times, and the results are presented as the

mean ± SD. The statistical significance of the differences between the mean values for the

different genotypes was measured by the Student's t-test with a paired, 2-tailed distribution.

The data were considered significant when the *P*-value was less than 0.05(*), 0.01(**),

0.001(***) or 0.0001(****).

Acknowledgements

This work was supported by the National Science Fund for Distinguished Young Scholars

(Grant No. 81925015), National Natural Science Foundation of China (Grant No. 32230029,

82371615), National Key Research and Development Program of China (Grant No.

2022YFC2702600), and Science and Technology Project of Guangzhou (Grant No.

2023A03J0886, 2023A03J0871).

Author Contributions

B.W. and C.L. performed most of the experiments and wrote the manuscript. Y.Y., Z.Z., and H.J. provided human adult sperm sample and performed some of the immunofluorescence experiments. S.M., Y.M., H.W., and J.L. contributed to the cell experiments and animal breeding. W.L. and C.L. supervised the whole project and revised the manuscript.

Declaration of interests

The authors declare that they have no conflict of interest.

References

Avasthi, P., Scheel, J. F., Ying, G., Frederick, J. M., Baehr, W., & Wolfrum, U. (2013). Germline deletion of Cetn1 causes infertility in male mice. *J Cell Sci*, *126*(Pt 14), 3204-3213. https://doi.org/10.1242/jcs.128587

Bazan, R., Schröfel, A., Joachimiak, E., Poprzeczko, M., Pigino, G., & Wloga, D. (2021). Ccdc113/Ccdc96 complex, a novel regulator of ciliary beating that connects radial spoke 3 to dynein g and the nexin link. *PLoS Genet*, *17*(3), e1009388. https://doi.org/10.1371/journal.pgen.1009388

Bicka, M., Joachimiak, E., Urbanska, P., Osinka, A., Konopka, A., Bulska, E., & Wloga, D. (2022). Cfap91-Dependent Stability of the RS2 and RS3 Base Proteins and Adjacent Inner Dynein Arms in Tetrahymena Cilia. *Cells*, *11*(24), 4048. https://doi.org/10.3390/cells11244048

Canty, J. T., Tan, R., Kusakci, E., Fernandes, J., & Yildiz, A. (2021). Structure and Mechanics of Dynein Motors. *Annu Rev Biophys*, *50*, 549-574. https://doi.org/10.1146/annurev-biophys-111020-101511

Deng, T. Q., Xie, Y. L., Pu, J. B., Xuan, J., & Li, X. M. (2022). Compound heterozygous mutations in PMFBP1 cause acephalic spermatozoa syndrome: A case report. *World J Clin Cases*, *10*(34), 12761-12767. https://doi.org/10.12998/wjcc.v10.i34.12761

Dooher, G. B., & Bennett, D. (1973). Fine structural observations on the development of the sperm head in the mouse. *Am J Anat*, *136*(3), 339-361. https://doi.org/10.1002/aja.1001360307

Dymek, E. E., Heuser, T., Nicastro, D., & Smith, E. F. (2011). The CSC is required for

complete radial spoke assembly and wild-type ciliary motility. *Mol Biol Cell*, *22*(14),

2520-2531. https://doi.org/10.1091/mbc.E11-03-0271

Elkhatib, R. A., Paci, M., Longepied, G., Saias-Magnan, J., Courbiere, B., Guichaoua, M. R.,

Levy, N., Metzler-Guillemain, C., & Mitchell, M. J. (2017). Homozygous deletion of

SUN5 in three men with decapitated spermatozoa. *Human Molecular Genetics*,

26(16), 3167-3171. https://doi.org/10.1093/hmg/ddx200

Fang, J., Zhang, J., Zhu, F., Yang, X., Cui, Y., & Liu, J. (2018). Patients with acephalic

spermatozoa syndrome linked to SUN5 mutations have a favorable pregnancy

outcome   from   ICSI.   *Hum   Reprod*,   *33*(3),   372-377.

https://doi.org/10.1093/humrep/dex382

Fawcett, D. W., & Phillips, D. M. (1969). The fine structure and development of the neck

region   of   the   mammalian   spermatozoon.   *Anat   Rec*,   *165*(2),   153-164.

https://doi.org/10.1002/ar.1091650204

Firat-Karalar, E. N., Sante, J., Elliott, S., & Stearns, T. (2014). Proteomic analysis of

mammalian sperm cells identifies new components of the centrosome. *J Cell Sci*,

127(Pt 19), 4128-4133. https://doi.org/10.1242/jcs.157008

Ghanaeian, A., Majhi, S., McCafferty, C. L., Nami, B., Black, C. S., Yang, S. K., Legal, T.,

Papoulas, O., Janowska, M., Valente-Paterno, M., Marcotte, E. M., Wloga, D., & Bui,

K. H. (2023). Integrated modeling of the Nexin-dynein regulatory complex reveals its

regulatory mechanism. *Nat Commun*, *14*(1), 5741. https://doi.org/10.1038/s41467-

023-41480-7

Gui, M., Ma, M., Sze-Tu, E., Wang, X., Koh, F., Zhong, E. D., Berger, B., Davis, J. H.,

Dutcher, S. K., Zhang, R., & Brown, A. (2021). Structures of radial spokes and

associated complexes important for ciliary motility. *Nat Struct Mol Biol*, *28*(1), 29-37.

https://doi.org/10.1038/s41594-020-00530-0

Hall, E. A., Keighren, M., Ford, M. J., Davey, T., Jarman, A. P., Smith, L. B., Jackson, I. J., &

Mill, P. (2013). Acute versus chronic loss of mammalian Azi1/Cep131 results in

distinct   ciliary   phenotypes.   *PLoS   Genet*,   *9*(12),   e1003928.

https://doi.org/10.1371/journal.pgen.1003928

Hermo, L., Pelletier, R. M., Cyr, D. G., & Smith, C. E. (2010). Surfing the wave, cycle, life

history, and genes/proteins expressed by testicular germ cells. Part 1: background to

spermatogenesis, spermatogonia, and spermatocytes. *Microscopy research and*

technique, *73*(4), 241–278. https://doi.org/10.1002/jemt.20783

Holm, L., & Laakso, L. M. (2016). Dali server update. *Nucleic Acids Res*, *44*(W1), W351-355.

https://doi.org/10.1093/nar/gkw357

Inaba, K., & Mizuno, K. (2016). Sperm dysfunction and ciliopathy. *Reprod Med Biol*, *15*(2),

77-94. https://doi.org/10.1007/s12522-015-0225-5

Ishikawa, T. (2017). Axoneme Structure from Motile Cilia. *Cold Spring Harbor Perspectives*

in Biology, *9*(1), a028076. https://doi.org/10.1101/cshperspect.a028076

Jreijiri, F., Cavarocchi, E., Amiri-Yekta, A., Cazin, C., Hosseini, S. H., El Khouri, E., Patrat,

C., Thierry-Mieg, N., Ray, P. F., Dulioust, E., Whitfield, M., & Touré, A. (2023).

CCDC65, encoding a component of the axonemal Nexin-Dynein regulatory complex,

is required for sperm flagellum structure in humans. *Clin Genet*, *105*(3), 317-322.

https://doi.org/10.1111/cge.14459

Kumar, N., & Singh, A. K. (2021). The anatomy, movement, and functions of human sperm

tail: an evolving mystery. *Biology of Reproduction*, *104*(3), 508-520.

https://doi.org/10.1093/biolre/ioaa213

Kumar, S., Stecher, G., Li, M., Knyaz, C., & Tamura, K. (2018). MEGA X: Molecular

Evolutionary Genetics Analysis across Computing Platforms. *Mol Biol Evol*, *35*(6),

1547-1549. https://doi.org/10.1093/molbev/msy096

Lehti, M. S., & Sironen, A. J. B. o. r. (2017). Formation and function of sperm tail structures

in association with sperm motility defects. *97*(4), 522-536.

https://doi.org/10.1093/biolre/iox096

Lehti, M. S., & Sironen, A. J. R. (2016). Formation and function of the manchette and

flagellum during spermatogenesis. *151*(4), R43-R54. https://doi.org/10.1530/REP-15-

0310

Leung, M. R., Zeng, J., Wang, X., Roelofs, M. C., Huang, W., Zenezini Chiozzi, R., Hevler, J.

F., Heck, A. J. R., Dutcher, S. K., Brown, A., Zhang, R., & Zeev-Ben-Mordehai, T. (2023). Structural specializations of the sperm tail. *Cell*, *186*(13), 2880-2896.e2817. https://doi.org/10.1016/j.cell.2023.05.026

Liska, F., Gosele, C., Rivkin, E., Tres, L., Cardoso, M. C., Domaing, P., Krejcí, E., Snajdr, P., Lee-Kirsch, M. A., de Rooij, D. G., Rooij, D. G., Kren, V., Krenová, D., Kierszenbaum, A. L., & Hubner, N. (2009). Rat hd mutation reveals an essential role of centrobin in spermatid head shaping and assembly of the head-tail coupling apparatus. *Biol. Reprod.*, *81*(6), 1196-1205. https://doi.org/10.1095/biolreprod.109.078980

Liu, G., Wang, N., Zhang, H., Yin, S., Dai, H., Lin, G., & Li, W. (2020). Novel mutations in PMFBP1, TSGA10 and SUN5: Expanding the spectrum of mutations that may cause acephalic spermatozoa. *Clin Genet*, *97*(6), 938-939. https://doi.org/10.1111/cge.13747

Liu, G., Xing, X., Zhang, H., Zhu, W., Lin, G., Lu, G., & Li, W. (2021). Patients with acephalic spermatozoa syndrome linked to novel TSGA10/PMFBP1 variants have favorable pregnancy outcomes from intracytoplasmic sperm injection. *Clin Genet*, *100*(3), 334-339. https://doi.org/10.1111/cge.14007

Lu, M., Kong, S., Xiang, M., Wang, Y., Zhang, J., Duan, Z., Zha, X., Wang, F., Cao, Y., & Zhu, F. (2021). A novel homozygous missense mutation of PMFBP1 causes acephalic spermatozoa syndrome. *J Assist Reprod Genet*, *38*(4), 949-955. https://doi.org/10.1007/s10815-021-02075-7

Ma, A., Zhou, J., Ali, H., Abbas, T., Ali, I., Muhammad, Z., Dil, S., Chen, J., Huang, X., Ma, H., Zhao, D., Zhang, B., Zhang, Y., Shah, W., Shah, B., Murtaza, G., Iqbal, F., Khan, M. A., Khan, A., Li, Q., Xu, B., Wu, L., Zhang, H., & Shi, Q. (2023). Loss-of-function mutations in CFAP57 cause multiple morphological abnormalities of the flagella in humans and mice. *JCI Insight*, *8*(3), e166869. https://doi.org/10.1172/jci.insight.166869

Martinez, G., Beurois, J., Dacheux, D., Cazin, C., Bidart, M., Kherraf, Z. E., Robinson, D. R., Satre, V., Le Gac, G., Ka, C., Gourlaouen, I., Fichou, Y., Petre, G., Dulioust, E., Zouari, R., Thierry-Mieg, N., Touré, A., Arnoult, C., Bonhivers, M., Ray, P., &

Coutton, C. (2020). Biallelic variants in MAATS1 encoding CFAP91, a calmodulin-associated and spoke-associated complex protein, cause severe astheno-teratozoospermia and male infertility. *J Med Genet*, *57*(10), 708-716. https://doi.org/10.1136/jmedgenet-2019-106775

Meng, X., Xu, C., Li, J., Qiu, B., Luo, J., Hong, Q., Tong, Y., Fang, C., Feng, Y., Ma, R., Shi, X., Lin, C., Pan, C., Zhu, X., Yan, X., & Cong, Y. (2024). Multi-scale structures of the mammalian radial spoke and divergence of axonemal complexes in ependymal cilia. *Nat Commun*, *15*(1), 362. https://doi.org/10.1038/s41467-023-44577-1

Nakata, H., Wakayama, T., Takai, Y., & Iseki, S. (2015). Quantitative analysis of the cellular composition in seminiferous tubules in normal and genetically modified infertile mice. *J Histochem Cytochem*, *63*(2), 99-113. https://doi.org/10.1369/0022155414562045

Nie, H., Tang, Y., Zhang, X., Tan, Y., & Qin, W. (2022). Novel mutations of PMFBP1 in a man with acephalic spermatozoa defects. *Mol Genet Genomic Med*, *10*(9), e2020. https://doi.org/10.1002/mgg3.2020

Oakberg, E. F. (1956). A description of spermiogenesis in the mouse and its use in analysis of the cycle of the seminiferous epithelium and germ cell renewal. *The American journal of anatomy*, *99*(3), 391-413. https://doi.org/10.1002/aja.1000990303

O'Donnell L. (2015). Mechanisms of spermiogenesis and spermiation and how they are disturbed. *Spermatogenesis*, *4*(2), e979623. https://doi.org/10.4161/21565562.2014.979623

Parker, G. A. (2020). Conceptual developments in sperm competition: a very brief synopsis. *Philosophical Transactions of the Royal Society B-Biological Sciences*, *375*(1813), 20200061. https://doi.org/10.1098/rstb.2020.0061

Roldan, E. R. S. (2019). Sperm competition and the evolution of sperm form and function in mammals. *Reproduction in Domestic Animals*, *54*, 14-21. https://doi.org/10.1111/rda.13552

Saitou, N., & Nei, M. (1987). The neighbor-joining method: a new method for reconstructing phylogenetic trees. *Mol Biol Evol*, *4*(4), 406-425. https://doi.org/10.1093/oxfordjournals.molbev.a040454

Sha, Y. W., Wang, X., Xu, X., Ding, L., Liu, W. S., Li, P., Su, Z. Y., Chen, J., Mei, L. B.,

Zheng, L. K., Wang, H. L., Kong, S. B., You, M., & Wu, J. F. (2019). Biallelic

mutations in PMFBP1 cause acephalic spermatozoa. *Clin Genet*, *95*(2), 277-286.

https://doi.org/10.1111/cge.13461

Sha, Y. W., Xu, X., Ji, Z. Y., Lin, S. B., Wang, X., Qiu, P. P., Zhou, Y., Mei, L. B., Su, Z. Y., Li,

L., & Li, P. (2018). Genetic contribution of SUN5 mutations to acephalic

spermatozoa in Fujian China. *Gene*, *647*, 221-225.

https://doi.org/10.1016/j.gene.2018.01.035

Shang, Y., Yan, J., Tang, W., Liu, C., Xiao, S., Guo, Y., Yuan, L., Chen, L., Jiang, H., Guo, X.,

Qiao, J., & Li, W. (2018). Mechanistic insights into acephalic spermatozoa syndrome-

associated mutations in the human SUN5 gene. *J Biol Chem*, *293*(7), 2395-2407.

https://doi.org/10.1074/jbc.RA117.000861

Shang, Y., Zhu, F., Wang, L., Ouyang, Y. C., Dong, M. Z., Liu, C., Zhao, H., Cui, X., Ma, D.,

Zhang, Z., Yang, X., Guo, Y., Liu, F., Yuan, L., Gao, F., Guo, X., Sun, Q. Y., Cao, Y.,

& Li, W. (2017). Essential role for SUN5 in anchoring sperm head to the tail. *Elife*, *6*,

e28199. https://doi.org/10.7554/eLife.28199

Sudhakar, D. V. S., Shah, R., & Gajbhiye, R. K. (2021). Genetics of Male Infertility - Present

and Future: A Narrative Review. *J Hum Reprod Sci*, *14*(3), 217-227.

https://doi.org/10.4103/jhrs.jhrs_115_21

Tu, C. F., Wang, W. L., Hu, T. Y., Lu, G. X., Lin, G., & Tan, Y. Q. (2020). Genetic

underpinnings of asthenozoospermia. *Best Practice & Research Clinical

Endocrinology & Metabolism*, *34*(6), 101472.

https://doi.org/10.1016/j.beem.2020.101472

Varadi, M., Anyango, S., Deshpande, M., Nair, S., Natassia, C., Yordanova, G., Yuan, D.,

Stroe, O., Wood, G., Laydon, A., Žídek, A., Green, T., Tunyasuvunakool, K., Petersen,

S., Jumper, J., Clancy, E., Green, R., Vora, A., Lutfi, M., Figurnov, M., Cowie, A.,

Hobbs, N., Kohli, P., Kleywegt, G., Birney, E., Hassabis, D., & Velankar, S. (2022).

AlphaFold Protein Structure Database: massively expanding the structural coverage

of protein-sequence space with high-accuracy models. *Nucleic Acids Res*, *50*(D1),

D439-d444. https://doi.org/10.1093/nar/gkab1061

Walton, T., Gui, M., Velkova, S., Fassad, M. R., Hirst, R. A., Haarman, E., O'Callaghan, C.,

Bottier, M., Burgoyne, T., Mitchison, H. M., & Brown, A. (2023). Axonemal

structures reveal mechanoregulatory and disease mechanisms. *Nature*, *618*(7965),

625-633. https://doi.org/10.1038/s41586-023-06140-2

Wang, L., Tu, Z., Liu, C., Liu, H., Kaldis, P., Chen, Z., & Li, W. (2018). Dual roles of TRF1

in tethering telomeres to the nuclear envelope and protecting them from fusion during

meiosis. *Cell Death Differ*, *25*(6), 1174-1188. https://doi.org/10.1038/s41418-017-

0037-8

Wu, B., Gao, H., Liu, C., & Li, W. (2020). The coupling apparatus of the sperm head and tail†.

Biol Reprod, *102*(5), 988-998. https://doi.org/10.1093/biolre/ioaa016

Xiang, M., Wang, Y., Wang, K., Kong, S., Lu, M., Zhang, J., Duan, Z., Zha, X., Shi, X., Wang,

F., Cao, Y., & Zhu, F. (2022). Novel Mutation and Deletion in SUN5 Cause Male

Infertility with Acephalic Spermatozoa Syndrome. *Reprod Sci*, *29*(2), 646-651.

https://doi.org/10.1007/s43032-021-00665-5

Yuan, S., Stratton, C. J., Bao, J., Zheng, H., Bhetwal, B. P., Yanagimachi, R., & Yan, W.

(2015). Spata6 is required for normal assembly of the sperm connecting piece and

tight head-tail conjunction. *Proc Natl Acad Sci U S A*, *112*(5), E430-439.

https://doi.org/10.1073/pnas.1424648112

Zamboni, L., & Stefanini, M. (1971). The fine structure of the neck of mammalian

spermatozoa. *Anat Rec*, *169*(2), 155-172. https://doi.org/10.1002/ar.1091690203

Zhang, D., Huang, W. J., Chen, G. Y., Dong, L. H., Tang, Y., Zhang, H., Li, Q. Q., Mei, X. Y.,

Wang, Z. H., & Lan, F. H. (2021). Pathogenesis of acephalic spermatozoa syndrome

caused by SUN5 variant. *Mol Hum Reprod*, *27*(5), gaab028.

https://doi.org/10.1093/molehr/gaab028

Zhang, Y., Liu, C., Wu, B., Li, L., Li, W., & Yuan, L. (2021). The missing linker between

SUN5 and PMFBP1 in sperm head-tail coupling apparatus. *Nat Commun*, *12*(1), 4926.

https://doi.org/10.1038/s41467-021-25227-w

Zhang, Y., Yang, L., Huang, L., Liu, G., Nie, X., Zhang, X., & Xing, X. (2021). SUN5

Interacting With Nesprin3 Plays an Essential Role in Sperm Head-to-Tail Linkage:

Research on Sun5 Gene Knockout Mice. *Front Cell Dev Biol*, *9*, 684826.

https://doi.org/10.3389/fcell.2021.684826

Zhou, L., Liu, H., Liu, S., Yang, X., Dong, Y., Pan, Y., Xiao, Z., Zheng, B., Sun, Y., Huang, P.,

Zhang, X., Hu, J., Sun, R., Feng, S., Zhu, Y., Liu, M., Gui, M., & Wu, J. (2023).

Structures of sperm flagellar doublet microtubules expand the genetic spectrum of

male infertility. *Cell*, *186*(13), 2897-2910.e2819.

https://doi.org/10.1016/j.cell.2023.05.009

Zhu, F., Liu, C., Wang, F., Yang, X., Zhang, J., Wu, H., Zhang, Z., He, X., Zhang, Z., Zhou, P.,

Wei, Z., Shang, Y., Wang, L., Zhang, R., Ouyang, Y. C., Sun, Q. Y., Cao, Y., & Li, W.

(2018). Mutations in PMFBP1 Cause Acephalic Spermatozoa Syndrome. *Am J Hum*

Genet, *103*(2), 188-199. https://doi.org/10.1016/j.ajhg.2018.06.010

Zhu, F., Wang, F., Yang, X., Zhang, J., Wu, H., Zhang, Z., Zhang, Z., He, X., Zhou, P., Wei, Z.,

Gecz, J., & Cao, Y. (2016). Biallelic SUN5 Mutations Cause Autosomal-Recessive

Acephalic Spermatozoa Syndrome. *Am J Hum Genet*, *99*(6), 1405.

https://doi.org/10.1016/j.ajhg.2016.11.002

Zhu, Z., Wang, Y., Wang, X., Yao, C., Zhao, L., Zhang, Z., Wu, Y., Chen, W., & Li, Z. (2022).

Novel mutation in ODF2 causes multiple morphological abnormalities of the sperm

flagella in an infertile male. *Asian journal of andrology*, *24*(5), 463 – 472.

https://doi.org/10.4103/aja202183

    Figures and Figure legends

Figure 1 CCDC113 is an evolutionarily conserved axoneme-associated protein. (A)

Multiple species phylogenetic tree of CCDC113. Structural similarity scores (Z scores) of

CCDC113 orthologs in *Homo sapiens*, *Mus musculus*, *Phascolarctos cinereus*, *Danio rerio*,

Chlamydomonas reinhardtii and *Tetrahymena thermophila* were derived through the DALI

webserver for pairwise structure comparisons (Holm & Laakso, 2016). (B) Positioning of

CCDC113 within the 96-nm repeat of human axoneme (Walton et al., 2023). CCDC113

forms a complex with CCDC96, is located at the base of RS3, and is adjacent to CFAP91 and

CFAP57. CFAP91 originates at the base of RS2 and links the RS3 subunits (CFAP251 and

CFAP61). (C-H) Neighboring axoneme-associated proteins were expressed alone or co-

expressed with CCDC113 in HEK293T cells, and the interactions between CCDC113 and

CFAP57, CFAP91, TUBB5, DRC1, DRC2, or DRC3 were examined by co-

immunoprecipitation. IB: immunoblotting; IP: immunoprecipitation. The % Input is displayed

below the corresponding figures for quantification. $n = 3$ independent experiments. Data are

presented as mean $\pm$ SD; $*P < 0.05$, $**P < 0.01$, $***P < 0.001$.

Figure 2 *Ccdc113* knockout leads to male infertility. (A) The CRISPR-Cas9 strategy for

generating the *Ccdc113* knockout mice. (B) Genotyping to identify *Ccdc113* knockout mice.

(C) Immunoblotting of CCDC113 in *Ccdc113*$^{+/+}$ and *Ccdc113*$^{-/-}$ testes. TUBULIN served as

the loading control. (D) The average litter size of *Ccdc113*$^{+/+}$ and *Ccdc113*$^{-/-}$ female mice in 2

months ($n = 5$ independent experiments). Data are presented as mean $\pm$ SD; ns indicates no

significant difference. (E) The average litter size of *Ccdc113*$^{+/+}$ and *Ccdc113*$^{-/-}$ male mice in 2

months ($n = 5$ independent experiments). Data are presented as mean $\pm$ SD; $****P < 0.0001$.

(F) The size of testes was similar in *Ccdc113*$^{+/+}$ and *Ccdc113*$^{-/-}$ mice. (G) The testis weights

of *Ccdc113*$^{+/+}$ and *Ccdc113*$^{-/-}$ male mice ($n = 5$ independent experiments). Data are presented

as mean $\pm$ SD; ns indicates no significant difference. (H) The body weights of *Ccdc113*$^{+/+}$ and

Ccdc113$^{-/-}$ male mice ($n = 5$ independent experiments). Data are presented as mean $\pm$ SD; ns

indicates no significant difference. (I) The ratio of testis weight/body weight in *Ccdc113*$^{+/+}$

and *Ccdc113*$^{-/-}$ male mice ($n = 5$ independent experiments). Data are presented as mean $\pm$ SD;

ns indicates no significant difference. (J) H&E staining of testis sections from $Ccdc113^{+/+}$ and

$Ccdc113^{-/-}$ male mice. Red asterisks indicate the abnormal sperm flagellum in the $Ccdc113^{-/-}$

testis seminiferous tubule. (K) Immunofluorescence of acetylated-tubulin (red) in testis

sections from $Ccdc113^{-/-}$ male mice showed flagellar defects. (L) Acetylated-tubulin

fluorescence intensity was measured per testis section in 155 sections from 3 $Ccdc113^{+/+}$ mice

and 153 sections from 3 $Ccdc113^{-/-}$ male mice. Data are presented as mean ± SD; ****$P <$

0.0001.

Figure 3 *Ccdc113* knockout results in sperm flagellar defects and sperm head-tail

detachment.

(A) H&E staining of the cauda epididymis from 2-month-old $Ccdc113^{+/+}$ and $Ccdc113^{-/-}$ male

mice. Red circles indicate coiled eosin-stained structures without sperm heads in the

epididymal lumen. (B) Analysis of sperm counts in $Ccdc113^{+/+}$ and $Ccdc113^{-/-}$ male mice ($n =$

5 independent experiments). Mature spermatozoa were extracted from the unilateral cauda

epididymis and dispersed in PBS. Sperm counts were measured using hemocytometers. Data

are presented as mean ± SD; ****P < 0.0001. (C) Motile sperm in $Ccdc113^{+/+}$ and $Ccdc113^{-/-}$

mice ($n = 5$ independent experiments). Data are presented as mean ± SD; ****P < 0.0001. (D)

$Ccdc113^{+/+}$ and $Ccdc113^{-/-}$ spermatozoa were co-stained with a flagellar marker α/β-tubulin

(red) and an acrosomal marker PNA. Nuclei were stained with DAPI (blue). (E)

Quantification of different categories of $Ccdc113^{+/+}$, $Ccdc113^{-/-}$ spermatozoa ($n = 3$

independent experiments). Data are presented as mean ± SD. (F) Immunofluorescence

analysis of acetylated-tubulin (green) and PNA (red) from $Ccdc113^{+/+}$ and $Ccdc113^{-/-}$

spermatids.

Figure 4. CCDC113 localizes to the HTCA, manchette and sperm flagellum. (A)

CCDC113 was expressed starting in P7 testes. TUBULIN served as the loading control. An

asterisk indicates nonspecific bands. (B) Immunofluorescence of CCDC113 (red) and

CENTRIN1/2 (green) in developing germ cells. CCDC113 partially colocalize with centriolar

protein CENTRIN1/2. (C) Immunofluorescence of CCDC113 (red) and α/β-tubulin (green) in

developing germ cells. The manchette was stained with the anti-α/β-tubulin antibody. (D-E) CCDC113 localizes to the HTCA and flagellum in mature mouse spermatozoa and human spermatozoa. Nuclei were stained with DAPI (blue).

Figure 5 CCDC113 is indispensable for the docking of CFAP91 and DRC2 to the DMTs to maintain the structural integrity of the axoneme. (A) TEM analysis of spermatozoa from the cauda epididymidis of $Ccdc113^{+/+}$ and $Ccdc113^{-/-}$ male mice. The flagellar longitudinal sections of $Ccdc113^{-/-}$ spermatozoa revealed unremoved cytoplasm, including disrupted mitochondria, damaged axonemes. and large vacuoles. Asterisks indicate large vacuoles. Cross sections of the principal piece of $Ccdc113^{-/-}$ spermatozoa further revealed partial loss or unidentifiable "9+2" structures, along with the disruption of the fibrous sheath and outer dense fibers. (B) TEM analysis of the axoneme in testicular spermatids from $Ccdc113^{+/+}$ and $Ccdc113^{-/-}$ male mice. The red arrowheads indicate the absence of significant radial spokes (RSs). MS: mitochondrial sheath, Mi: mitochondrial, AX: axoneme. FS: fibrous sheath. DMT: doublet microtubule, MT: microtubule, CP: central pair, ODF: outer dense fiber, RS: radial spokes. (C) The immunofluorescence analysis for CFAP91 (green) and α/β-tubulin (red) was performed in $Ccdc113^{+/+}$ and $Ccdc113^{-/-}$ spermatozoa. Nuclei were stained with DAPI (blue). White asterisks indicate regions not co-localized with tubulin. (D) The immunofluorescence analysis for DRC2 (green) and α/β-tubulin (red) was performed in $Ccdc113^{+/+}$ and $Ccdc113^{-/-}$ spermatozoa. Nuclei were stained with DAPI (blue). White asterisks indicate regions not co-localized with tubulin. (E, G) Line-scan analysis (white line) was performed using Image J software. (F, H) Percentage of abnormally distributed CFAP91 and DRC2 on the axoneme of $Ccdc113^{+/+}$ and $Ccdc113^{-/-}$ spermatozoa ($n = 3$ independent experiments). At least 200 spermatozoa were analyzed from each mouse. Data are presented as mean ± SD; ****$P < 0.0001$.

Figure 6 *Ccdc113* knockout spermatids display impaired HTCA. (A) The proportion of decapitated tails in $Ccdc113^{+/+}$ and $Ccdc113^{-/-}$ corpus, caput and cauda epididymis ($n = 3$ independent experiments). At least 200 spermatozoa were analyzed from each mouse. Data

are presented as mean ± SD; ****P < 0.0001, ***P < 0.001. (B) PAS staining of testis sections from $Ccdc113^{+/+}$ and $Ccdc113^{-/-}$ mice. The green arrows indicate the orientation of the sperm heads in stages V–VIII seminiferous epithelia. $Ccdc113^{-/-}$ sperm head could still be detected in stages IX–X seminiferous epithelia. P: pachytene spermatocyte, spz: spermatozoa, rSt: round spermatid, eSt: elongating spermatid, Z: zygotene spermatocyte, M: meiotic spermatocyte. (C) $Ccdc113^{-/-}$ spermatids lost their head orientation toward the basement membrane during spermiation in stages VII–VIII of the seminiferous epithelium. L: lumen, B: basement membrane. (D) Percentage of sperm heads with abnormal orientation in stages VII–VIII of the seminiferous epithelium in $Ccdc113^{+/+}$ and $Ccdc113^{-/-}$ mice (n = 3 independent experiments). At least 200 spermatozoa were analyzed from each mouse. Data are presented as mean ± SD; ****P < 0.0001. (E) Defective HTCA formation in $Ccdc113^{-/-}$ spermatids. TEM analysis of the stepwise development of the HTCA were performed in $Ccdc113^{+/+}$ and $Ccdc113^{-/-}$ testes. In $Ccdc113^{+/+}$ spermatids, the well-defined coupling apparatus was tightly attached to the sperm head. In $Ccdc113^{-/-}$ spermatids, segmented columns (Scs), the capitulum (Cp) were absent. The red asterisks indicate the distance between the sperm head and HTCA. The white arrows indicate the dense material surrounding the proximal centriole. Nu: nuclear, Bp: basal plate, Cp: capitulum, Sc: segmented column, Pc: proximal centriole, Dc: distal centriole, An: annulus, Ax: axoneme, Rn: redundant nuclear envelope.

Figure 7 CCDC113 interact with SUN5 and CENTLEIN, participating in sperm head-tail linkage. (A-C, F) HTCA-associated proteins (SUN5, CENTLEIN, PMFBP1, SPATA6) were expressed alone or co-expressed with CCDC113 in HEK293T cells, and the interactions between CCDC113 and these HTCA-associated proteins were examined by co-immunoprecipitation. CCDC113 interacted with SUN5 and CENTLEIN, but did not interact with PMFBP1 and SPATA6. IB: immunoblotting; IP: immunoprecipitation. (D) SUN5 interacted with CCDC113. pECMV-FLAG-$Ccdc113$ and pEGFP-GFP-$Sun5$ were transfected into HEK293T cells. At 48 h after transfection, the cells were collected for immunoprecipitation (IP) with anti-GFP antibody and analyzed with anti-FLAG and anti-GFP antibodies. (E) CENTLEIN interacted with CCDC113. pCDNA -FLAG-$Centlein$ and pEGFP-

GFP-*Ccdc113* were transfected into HEK293T cells. At 48 h after transfection, the cells were

collected for immunoprecipitation (IP) with anti-FLAG antibody and analyzed with anti-

FLAG and anti-GFP antibodies. The % Input is displayed below the corresponding figures for

quantification. $n = 3$ independent experiments. Data are presented as mean ± SD; *$P < 0.05$,

ns indicates no significant difference. (G) Immunofluorescence of CCDC113 (red) and SUN5

(green) in mature spermatozoa. Nuclei were stained with DAPI (blue). (H)

Immunofluorescence of CCDC113 (red) and CENTLEIN (green) in testicular step13-step14

spermatid. Nuclei were stained with DAPI (blue). (I) Immunofluorescence analysis for

SPATA6 (green) and α/β-tubulin (red) was performed in *Ccdc113*$^{+/+}$ and *Ccdc113*$^{−/−}$

spermatozoa. Nuclei were stained with DAPI (blue). (J) Quantification ratio of SPATA6 on

the detached sperm tail ($n = 3$ independent experiments). At least 200 spermatozoa were

analyzed for each mouse. (K) Quantification ratio of CCDC113 on the detached sperm tail (n

$= 3$ independent experiments). At least 200 spermatozoa were analyzed from each mouse.

Data are presented as mean ± SD; ***$P < 0.001$, ****$P < 0.0001$. (L) Immunofluorescence

analysis for CCDC113 (red) was performed in wild type (WT), *Sun5*$^{−/−}$, *Centlein*$^{−/−}$, and

Pmfbp1$^{−/−}$ spermatozoa. Nuclei were stained with DAPI (blue).