**Chalkophore mediated respiratory oxidase flexibility controls *M. tuberculosis* virulence**

John A. Buglino[1,*], Yaprak Ozakman[1,*], Chad Hatch[2], Anna Benjamin[1], Derek Tan[2,3], Michael S. Glickman[1,#]

1-Immunology Program, Sloan Kettering Institute
2-Chemical Biology Program, Sloan Kettering Institute
3-Tri-Institutional Research Program
Memorial Sloan Kettering Cancer Center, 1275 York Avenue, New York, New York 10065 USA

*Equal contribution

#Correspondence:
glickmam@mskcc.org
1275 York Ave
New York, NY 10065
646-888-2368

**Abstract**

Oxidative phosphorylation has emerged as a critical therapeutic vulnerability of *M. tuberculosis* (Mtb). However, it is unknown how intracellular bacterial pathogens such as Mtb maintain respiration during infection despite the chemical effectors of host immunity. *Mtb* synthesizes diisonitrile lipopeptides that tightly chelate copper, but the role of these chalkophores in host-pathogen interactions is also unknown. We demonstrate that *M. tuberculosis* chalkophores maintain the function of the heme-copper *bcc:aa$_3$* respiratory oxidase under copper limitation. Chalkophore deficiency impairs Mtb survival, respiration to oxygen, and ATP production under copper deprivation in culture, effects that are exacerbated by loss of the heme dependent Cytochrome BD respiratory oxidase. Our genetic analyses indicate that maintenance of respiration is the only cellular target of chalkophore mediated copper acquisition. *M. tuberculosis* lacking chalkophore biosynthesis is attenuated in mice, a phenotype that is also severely exacerbated by loss of the CytBD respiratory oxidase. We find that the host immune pressure that attenuates chalkophore deficient *Mtb* is independent of adaptive immunity and neutrophils. These data demonstrate that chalkophores counter host inflicted copper deprivation and highlight a multilayered system by which *M. tuberculosis* maintains respiration during infection.

**Introduction**

Bacterial pathogens are subjected to diverse stresses imposed by host immunity and deploy countermeasures to neutralize immune effectors. The ubiquity of metalloenzymes in all bacteria necessitates acquisition of trace metals such as iron, zinc and manganese, a vulnerability that is exploited by host nutritional immunity, which limits these metals (1). To counter this metal limitation by the host, pathogens deploy diverse high affinity metal acquisition systems including siderophores for iron and zinc (2-4), and transporters and metallophores for zinc and manganese (1, 3, 5-7). In addition to metal limitation, the host also deploys metals as antimicrobial effectors to kill bacteria (8). Copper and zinc are deposited into the phagosome of infected macrophages as antimicrobials and pathogens employ metal resistance systems such as metal efflux transporters (9-11) and metal binding proteins controlled by metal dependent repressors (9-11). Although both metal limitation and metal intoxication limit pathogen growth, in most cases the essential bacterial metalloenzymes rendered dysfunctional by nutritional immunity are incompletely defined.

*M. tuberculosis* is a successful global pathogen that can survive in both macrophages and neutrophils. *M. tuberculosis* experiences both high copper (10-15) and high zinc (12) in the macrophage phagosome and resists metal toxicity through several mechanisms including efflux and metal chelation. Recent data also indicates that *M. tuberculosis* experiences zinc starvation during infection, possibly imposed by calprotectin in caseum (16). However, it is unknown whether copper acquisition, and resistance to copper deprivation, are part of the virulence program of *M. tuberculosis. M. tuberculosis* synthesizes diisonitrile lipopeptide natural products directed by the 5 gene *nrp* operon, present in *M. tuberculosis* and *M. marinum*, but not nonpathogenic mycobacteria (17-20). The *nrp* operon is induced by copper deprivation and growth of *M. tuberculosis* lacking diisonitrile lipopeptide biosynthesis is inhibited in copper limiting conditions and rescued by a synthetic diisonitrile (21), establishing diisonitrile lipopeptides as mycobacterial chalkophores. However, the role of diisonitrile chalkophores in *M. tuberculosis* pathogenesis is not understood, including whether copper deprivation is imposed by the host during infection, and the specific bacterial pathways that require copper supplied by the chalkophores. In this study, we demonstrate that diisonitrile chalkophores supply copper to the heme-copper respiratory oxidase to maintain respiration and ATP production. The heme-copper oxidase of chalkophore deficient *M. tuberculosis* is compromised by the host during infection, but *M. tuberculosis* compensates with the heme-dependent cytochrome BD (CytBD).

76    *M. tuberculosis* lacking diisonitrile chalkophore biosynthesis and CytBD is severely attenuated,

77    demonstrating that this multilayered system for protecting respiration is a critical virulence

78    function of *M. tuberculosis*.

79 **Results**
80
81 **Chalkophore deficient *M. tuberculosis* upregulates respiratory chain components in**
82 **response to copper deprivation**
83
To understand the function of diisonitrile chalkophores in *M. tuberculosis*, we examined the

transcriptional profile of wild type (WT) and Δ*nrp M. tuberculosis* treated with 10 μM

tetrathiomolybdate (TTM), a copper chelator previously shown to inhibit the growth of *M*

*tuberculosis* lacking chalkophore biosynthesis (21). Copper chelation had relatively few effects

on gene expression in wild type *M. tuberculosis* (Figure 1A, Table S3). In contrast, in Δ*nrp M.*

*tuberculosis* we observed upregulation of the genes of the chalkophore cluster (other than *nrp*

itself) at baseline, consistent with prior data demonstrating autoregulation (21). In addition, a

cluster of genes was induced by copper deprivation in Δ*nrp* cells, but not wild type cells,

encoding components of the respiratory chain, including *cydABDC* (encoding components of the

heme dependent oxidase CytBD), *qcrABC* (encoding subunits of the *bcc:aa₃* heme-copper

oxidase), and subunits of ATP synthase (Fig 1A). Prior work demonstrated that genetic or

pharmacologic disruption of the *bcc:aa₃* respiratory oxidase, including by Q203, an inhibitor of

the QcrB subunit of the *bcc:aa₃* oxidase, transcriptionally upregulates genes encoding

components of the respiratory chain, including *cydAB* and ATP synthase (22, 23). Treatment of

wild type and Δ*nrp* with Q203 reproduced the published pattern in wild type cells and indicated

that diisonitrile chalkophore deficient cells still respond to Q203 (Figure 1). To confirm the RNA

sequencing results, we quantitated the transcript encoding the *cydA* encoded subunit of the

CytBD oxidase with different concentrations of TTM and observed no induction at any TTM

concentration in wild type cells, but progressive induction of CydA with escalating TTM

concentrations in the Δ*nrp* strain (Figure 1B). Longer durations of copper deprivation induced

*cydA* in both wild type and Δ*nrp*, with higher induction in Δ*nrp* (Figure 1C). Escalating

concentrations of TTM did not affect wild type *M. tuberculosis* growth but caused a graded

inhibition of growth of the Δ*nrp* strain in liquid media (Fig 1D). These data indicate that copper

deprivation in *M. tuberculosis* lacking diisonitrile chalkophore biosynthesis stimulates gene

expression that mimics inhibition of the *bcc:aa₃* respiratory oxidase.

109
110 **Chalkophores protect the *bcc:aa₃* oxidase from copper deprivation**
111
Inhibition of *bcc:aa₃* oxidase by Q203 (22, 24) or other QcrB inhibitors (25-28) is bacteriostatic

and incompletely inhibits respiration, but becomes bactericidal and abolishes oxidative

phosphorylation in *M. tuberculosis* lacking the alternative cytochrome BD (Δ*cydAB*, (22, 28)) or

treated with a CytBD inhibitor (29). Although the *bcc:aa₃* oxidase and CytBD are functionally

redundant, they differ in cofactor usage for electron transfer: *bcc:aa₃* is a heme-copper oxidase

with three copper ions in the electron transport path (30-33), whereas CytBD uses heme

prosthetic groups but not copper (34)(Figure 2A).

To test whether diisonitrile chalkophores maintain the function of the copper dependent

*bcc:aa₃* oxidase under copper limitation (Figure 2A), we generated *M. tuberculosis ΔnrpΔcydAB*,

along with control strains lacking *cydAB* alone and genetically complemented strains. We tested

the effect of TTM on growth in liquid media and observed mild inhibition of growth of *M.*

*tuberculosis Δnrp* and no effect on *ΔcydAB* (Figure 2B). However, growth of *ΔnrpΔcydAB* was

severely inhibited by TTM, an effect that was reversed by genetic complementation with either

*nrp* or *cydABDC* (Figure 2B). Assays on copper chelated agar media revealed a dramatic

sensitization of chalkophore deficient *M. tuberculosis* by loss of the secondary oxidase with 6

logs of killing, a phenotype that was also complemented by *nrp* or *cydABDC* (Figure 2C, S1). To

determine whether the phenotypes of *Δnrp* are due to diisonitrile chalkophore deficiency rather

than some function of the genetic element independent of the diisonitrile chalkophore itself, we

first repeated the same *cydAB* synergy test with *M. tuberculosis ΔfadD10*, a second biosynthetic

gene in the *nrp* operon that is required for chalkophore mediated resistance to copper chelation

(21). *ΔfadD10ΔcydAB M. tuberculosis* was also dramatically sensitized to

bathocuproinedisulfonic acid, disodium salt (BCS, another copper chelator) or TTM to a similar

degree as *ΔnrpΔcydAB* (Figure S1B, C). To demonstrate that the copper deprivation sensitivity

of the *ΔnrpΔcydAB* strain is due to absence of the diisonitrile lipopeptide, we synthesized an

analogue of a reported *M. tuberculosis* diisonitrile chalkophore (20) having 4-carbon side chains

(see Methods and SI Appendix). This C4 L-ornithyl-L-phenylalaninol acetate diisonitrile (C4-Orn-

Phin-OAc) was synthesized via a route analogous to our previously reported syntheses of a

*Streptomyces* diisonitrile (35). This synthetic diisonitrile efficiently rescued *ΔnrpΔcydAB* from the

growth inhibition imposed by TTM in liquid media (Figure 2D), consistent with direct mediation of

this effect by the diisonitrile lipopeptide product of the *nrp* locus.

The data above indicates that diisonitrile biosynthesis is necessary to resist the effects of

copper starvation and that one of the effects of this copper starvation is dysfunction of the

*bcc:aa₃* oxidase. To determine whether maintenance of respiratory oxidase function under

copper starvation is the only function of diisonitrile chalkophores, we executed genetic and

biochemical epistasis testing. Deleting *ctaD,* which encodes one subunit of the *bcc:aa₃* oxidase,

alone or in diisonitrile chalkophore deficient cells (*ΔnrpΔctaD*), impaired growth of *M.*

*tuberculosis* in copper replete media (Figure 2E), but copper deprivation had no additional

effect, indicating that the heme-copper oxidase complex is the only target impacted by copper

starvation (Figure 2E). Similarly, the growth inhibitory effect of copper starvation on Δ*nrp* cells

was similar in degree to treatment of WT or Δ*nrp* cells with Q203, but combined treatment was

not synergistic (Figure 2F), again indicating that the heme-copper oxidase is inactivated by

copper deprivation when the diisonitrile chalkophore is missing and that no additional targets

are relevant to copper deprivation-induced growth arrest.

155
156    **Chalkophores maintain oxidative phosphorylation and ATP production**
157
To determine whether diisonitrile chalkophores directly maintain respiration during copper

deprivation, as suggested by the growth and survival data above, we measured oxygen

consumption by *M. tuberculosis* using a qualitative methylene blue assay (22, 29). In this assay,

decolorization of methylene blue in a sealed tube indicates oxygen depletion, as observed in

WT or Δ*nrp*Δ*cydAB M. tuberculosis* treated with DMSO (Figure 3A), indicating intact respiration

in chalkophore deficient cells in basal conditions. Treatment with Q203 did not impair oxygen

consumption in WT cells but did in Δ*nrp*Δ*cydAB*, consistent with its inhibition of the *bcc*:*aa*$_3$

oxidase (Figure 3A). Treatment of wild-type cells with Q203 in combination with ND-011992, a

small-molecule inhibitor of CytBD, also blocked oxygen consumption, consistent with prior data

(Figure S4A) (22, 29). Copper chelation with 50 μM TTM had no effect on wild type cells but

abolished respiration in Δ*nrp*Δ*cydAB*, indicating that copper deprivation inhibits respiration

through the *bcc*:*aa*$_3$ oxidase (Figure 3A). Similarly, Δ*nrp* cells grown on agar media with TTM

and ND-011992 lost viability (Figure S4C). Taken together, these data are consistent with a

model in which *M. tuberculosis* can respire via either the *bcc*:*aa*$_3$ oxidase or CytBD, with the

former requiring diisonitrile chalkophore-mediated copper acquisition under copper starvation.

The full effect of loss of chalkophore biosynthesis is masked by the redundancy of the oxidases,

but in cells that rely only on *bcc*:*aa*$_3$ oxidase, respiration is abolished by copper deprivation.

To measure oxygen consumption more quantitatively, we adopted a spot sensor assay (36),

that can noninvasively measure oxygen in sealed vessels, allowing serial measurements. WT

*M. tuberculosis* consumed oxygen down to the lower limit of detection, and treatment with the

CytBD inhibitor ND-011992 (29) had no effect (Figure 3B). Treatment with the QcrB inhibitor

Q203 delayed but did not prevent oxygen consumption, consistent with the methylene blue

results and prior data (22, 24, 29). However, treatment with the combination of Q203 and ND-

011992 completely inhibited oxygen consumption (Figure 3B). Applying this assay to diisonitrile

chalkophore deficient strains, we observed that copper deprivation delayed but did not prevent

respiration in Δ*nrp* cells (Figure 3C), consistent with compensation by CytBD, but abolished

respiration in Δ*nrpΔcydAB* cells, a phenotype that was rescued by genetic complementation with

the *nrp* gene (Figure 3D). To confirm that the diisonitrile chalkophore-mediated protection of

respiration is accompanied by ATP depletion, we measured ATP levels in WT and chalkophore

deficient cells treated with copper deprivation or Q203. Copper deprivation with TTM or

treatment with Q203 had minimal effect on wild-type cells (Figure 3E), but inhibited ATP

production in the Δ*nrpΔcydAB* strain (Figure 3E).  Taken together, these data demonstrate that

diisonitrile chalkophores maintain oxidative phosphorylation by the heme-copper respiratory

oxidase under copper starvation. Because the electron transport path of the *bcc:aa*$_3$ oxidase

contains three copper sites, these findings are consistent with a model in which these copper

sites become dysfunctional when copper is limiting, leading to impaired supercomplex

biosynthesis or function.

In eukaryotic copper trafficking disorders in which copper becomes limiting in mitochondria,

cytochrome C oxidase becomes unstable, preventing assembly (37). To determine if similar

mechanism is operative in mycobacterial cells, we inserted an ALFA epitope tag (38) at the C

terminus of CtaD in both WT and Δ*nrp* cells and, after confirming functionality (Figure 2E),

examined CtaD protein levels with copper deprivation. CtaD levels remained unchanged with

either TTM or BCS (Figure 3F, S4B), indicating that copper deprivation is not affecting oxidase

biogenesis and is more likely acting on preexisting respiratory oxidase complexes.

204
205    **Chalkophores defend oxidative phosphorylation from nutritional immunity**
206
Respiration has emerged as an attractive target for antimycobacterials, with the ATP synthase

inhibitor bedaquiline now a cornerstone of MDR TB treatment (39, 40) and Q203 in clinical trials

(41). Although it is also clear that ATP generation via oxidative phosphorylation is required for

*M. tuberculosis* growth in mice (22, 29, 42, 43), it is less clear whether the reactive centers of

the electron transport chain are compromised by the host and whether *M. tuberculosis* must

defend or restore the integrity of the electron transport chain during infection. To examine this

question, we infected C57BL/6 mice with *M. tuberculosis* strains lacking diisonitrile chalkophore

biosynthesis, alone and in combination with deletion of the genes encoding CytBD (Δ*nrp* and

Δ*nrpΔcydAB*). The mouse attenuation phenotype of *M. tuberculosis* lacking *nrp* has been

reported by several groups (12, 20, 44) and indicates a mild early attenuation phenotype in the

lungs. Loss of *cydAB* in the Δ*nrp* background dramatically exacerbated the mild attenuation of

diisonitrile chalkophore deficient strain in the lung (Figure 4A) and caused severe attenuation in

the spleen (Figure 4B). Complementation with the *nrp* gene restored virulence to wild-type

levels, demonstrating that chalkophore mediated protection of the respiratory chain is a critical

virulence function of *M. tuberculosis*.

To investigate the host pressure that targets the respiratory chain, we hypothesized that

neutrophils, which express several metal binding proteins such as calprotectin, might be

relevant. However, efficient depletion of neutrophils by administration of a Ly6G antibody

(Figure S5A) did not reverse the severe attenuation of the Δ*nrp*Δ*cydAB* strain in the lungs or

spleen (Figure 4C, D). Similarly, infection of SCID mice, which lack adaptive immunity, also had

no effect on diisonitrile chalkophore deficient *M. tuberculosis* titers in lung or spleen (Figure 4E,

F) and the rapid mortality of SCID mice infected wild type *M. tuberculosis* was not evident in

SCID mice infected with Δ*nrp*Δ*cydAB* (Figure S5B), indicating that non adaptive immunity is fully

capable of controlling respiratory chain compromised *M. tuberculosis*.

233
## Discussion
235
We have identified diisonitrile chalkophore biosynthesis as an *M. tuberculosis* virulence

mechanism that defends against copper starvation during infection. Although it is well

established that phagosomal pathogens experience both metal excess and metal deprivation in

the host, the general paradigm of copper and zinc nutritional immunity is that high phagosomal

levels of these metals are deployed as antimicrobial effectors to limit pathogen growth (10, 11,

45-47). Our data indicate that *M. tuberculosis* must also cope with copper deprivation as a host

inflicted stress during infection and that *M. tuberculosis* deploys diisonitrile chalkophores to

acquire copper in the host. Diisonitrile chalkophores display an extremely high affinity for copper

ions (18, 35, 48), and their ability to overcome copper deprivation by strong copper chelators in

vitro and host immunity in vivo indicates that they perform an analogous function to bacterial

siderophores, which scavenge iron from host iron limitation. Given the abundant data indicating

a high copper environment in the macrophage phagosome, the sites and circumstances of the

copper deprived niche of *M. tuberculosis* remain to be determined but could include distinct

subcellular compartments (i.e., phagosomal vs cytosolic), phases of infection, or bacterial

localization within different lung compartments such as cavities.

Although deprivation of metals from pathogens, including iron and zinc, is a well-recognized

mechanism of innate immunity, in most cases the specific bacterial targets that ultimately

mediate the antimicrobial function of metal deprivation are not clearly defined and are assumed

to be pleiotropic due to the numerous essential metal dependent enzymes in the bacterial cell.

However, our data indicates that the copper deprivation countered by the diisonitrile

chalkophore system targets a single membrane enzyme complex, the *bcc:aa$_3$* supercomplex, a

heme-copper respiratory oxidase. *M. tuberculosis* deploys two respiratory oxidases, *bcc:aa$_3$* and

CytBD, which are redundant for bacterial viability. This dual oxidase arrangement provides

respiratory chain flexibility during chemical inhibition of each oxidase, and some evidence

suggests that respiratory chain flexibility promotes host adaptation (22, 42, 43). The copper

centers of respiratory oxidases that participate in electron flow to oxygen are conserved from

bacteria to mitochondria and represent a membrane exposed electroreactive center susceptible

to damage. Our data reveals that *M. tuberculosis* deploys multilayered strategies to maintain

oxidative phosphorylation in the face of host immune pressure. In addition, *M. tuberculosis*

deploys a backup oxidase, CytBD, which can support *M. tuberculosis* virulence when the

*bcc:aa$_3$* oxidase is dysfunctional, such as when QcrB is inhibited by Q203. However, the role of

this backup oxidase in pathogenesis was unclear. Although CytBD is induced in mouse lung,

peaking at 21 days post infection (49), is required for optimal fitness of *M. tuberculosis* in the

lung in competition experiments (43), *M. tuberculosis* lacking CytBD is fully virulent (22). Our

results reveal the essential role of this compensatory oxidase in virulence, which is only evident

when the $bcc:aa_3$ oxidase in inactivated in vivo in *M. tuberculosis* lacking chalkophore

biosynthesis, thereby revealing a complex system for maintaining respiration. These studies

further strengthen the rationale for targeting CytBD for antibiotic development (34, 50).

The respiratory chain of *M. tuberculosis* has emerged as a promising drug target. Our data

indicates that diisonitrile chalkophore biosynthesis, as a mechanism of protection for the

respiratory chain during infection, may provide an alternative approach to target this critical

energy generating system. Beyond its importance as a drug target, this study identifies a new

mechanism of resistance to copper deprivation that, in concert with CytBD, is part of a central

virulence strategy of *M. tuberculosis* to protect oxidative phosphorylation during infection.

283
**Materials and Methods**

285
**Reagents**

Middlebrook 7H10 Agar, 7H9 Broth, dextrose, Tween-80, bovine serum albumin (BSA), and

UltraPure DNase/RNase -free distilled water were purchased from Fisher Scientific. Ammonium

tetrathiomolybdate (TTM), bathocuproinedisulfonic acid, disodium salt (BCS), dimethyl sulfoxide

(DMSO), copper sulfate, zinc sulfate, magnesium sulfate, calcium chloride, and ATP were

purchased from Millipore Sigma. Biotechnology (BT) grade Chelex 100 resin, sodium form, was

purchased from BioRad.  Q203 (Telacebec) was purchased from AbMole.  ND-011992 was

synthesized as previously described (29).  The diisonitrile lipopeptide analogue was synthesized

and characterized as described below.

**General growth conditions, strains, and DNA manipulations**

*M. tuberculosis* Erdman WT and mutant stains were grown and maintained in 7H9 media

(broth), or on 7H10 (agar) supplemented with 10% OADC (oleic acid, albumin, dextrose, saline),

0.05% glycerol (7H9-OADC/7H10-OADC). Broth cultures were additionally supplemented with

0.02% Tween-80. Chromosomal deletion mutations were generated by specialized transduction

utilizing the temperature sensitive phage phAE87. Mutant strains were confirmed by PCR

followed by sequencing. For complete strain list with relevant features see **Table S1**. Plasmids

utilized in this study were generated using standard molecular techniques and are listed with

their features in **Table S2**.

**Growth assays**

For liquid growth assays, the indicated strains were pre grown in non-chelated media until

reaching an $OD_{600}$ value of ~1.0.  Cells were then collected by centrifugation (3,700 x *g*, 10 min)

and washed twice with chelexed phosphate buffered saline with 0.02% Tween-80 (PBS Tween-

80). Growth assays were initiated at a calculated $OD_{600}$ of 0.005 by the addition of 1 mL of

washed culture at an $OD_{600}$ of 0.05 to 9 mL of replete 7H9-ADS (albumin, dextrose, saline)

generated as described previously (21). TTM, BCS, Q203, ND-011992, and/or synthetic

diisonitrile chalkophore were then added at the indicated concentration.  Growth at 37°C was

assayed via daily $OD_{600}$ measurements.

For agar growth assays, cells were pre-grown and washed as indicated above. Washed cells

were then normalized to an $OD_{600}$ of 0.1, serially diluted from $10^0$-$10^{-6}$ in PBS Tween-80, and

spotted on 7H10-OADC plates containing the indicated concentrations of TTM, BCS, ND-

011992 or DMSO vehicle control in triplicate. Plates incubated at 37°C, 5% $CO_2$ were imaged

after 14 days of incubation to show relative growth, and CFU counts were enumerated after 21

days of incubation.

**RNA preparation**

Triplicate 25 mL cultures of strains indicated above were grown in 7H9-ADS until they reached

an $OD_{600}$ of 0.3 to 0.5. Cultures were then treated with the indicated concentration of TTM, 100

nM Q203, or a DMSO vehicle control, for 5 hrs. Following a duplicate wash with Chelex treated

PBS Tween-80, cells were collected by centrifugation (3,700 × $g$, 10 min), suspended, and

resuspended in 1 mL of TRIzol reagent. Cells in 1 mL of TRIzol were mechanically disrupted

with zirconia beads via 2x45 sec pulses in a BioSpec Mini24 BeadBeater with 5 min intervening

rest periods on ice. Following lysis, beads were removed by centrifugation (20,000 × $g$, 5 min)

and total RNA was isolated using the Direct-zol RNA miniprep kit (Zymo Research) as directed

by the manufacturer.

**RT-qPCR**

DNAse treatment of RNA purified as noted above was performed using the Turbo DNA-free kit

(Invitrogen). A total of 500 ng of resulting total RNA was used to synthesize cDNA via random

priming utilizing the iScript™ cDNA Synthesis Kit (Biorad). Real-time qPCR was performed on

QuantStudio 6 Pro (Applied Biosystems) using iTaq Universal SYBR Green Supermix (Biorad).

For *cydA* expression, normalized cycle threshold (CT) was determined relative to the

housekeeping gene *sigA*. Reactions with reverse transcriptase were included for each sample

and excluded DNA contamination as a source of amplification signal. Relative expression level

was calculated using the formula $2^{-(CT\mathit{cydA}-CT\mathit{sigA})}$. Sequences of gene specific primers are as

follows: For *cydA,* 5'-GTCATCGAAGTGCCCTATGT-3' and 5'- CTGGTATTCCTGCTGCAGAT-

3', and for *sigA,* 5'- CGTCTTCATCCCAGACGAAAT-3' and 5'-CGACGAAGACCACGAAGAC-3'.

**RNA Sequencing and Data Analysis**

After RiboGreen quantification and quality control by Agilent BioAnalyzer, 500 ng of total RNA

underwent ribosomal depletion with the NEBNext rRNA Depletion Kit (Bacteria) (NEB catalog #

E7850) and library preparation with the TruSeq Stranded Total RNA LT Kit (Illumina catalog #

RS-122-1202) according to instructions provided by the manufacturer with 8 cycles of PCR.

Samples were barcoded and run on a NovaSeq 6000 in a PE100 run, using the NovaSeq 6000

S4 Reagent Kit (200 Cycles) (Illumina). On average, 27 million paired reads were generated

per sample. Post-run demultiplexing and adapter removal were performed and fastq files were

inspected using fastqc (51). Trimmed fastq files were then aligned to the reference genome (*M.*

*tuberculosis* H37Rv; NC_000962.3) using bwa mem (52). Bam files were sorted and merged

using samtools (53) and gene counts were obtained using featureCounts from the Bioconductor

Rsubread package (54). Differentially expressed genes were identified using the DESeq2 R

package (55) and subsequent analysis of gene expression was performed in (56, 57).

364
**Immunoblotting**

Duplicate 20 mL cultures of chelated 7H9-ADS replete for all ions were inoculated at an $OD_{600}$

of 0.02. Upon reaching an $OD_{600}$ of 0.6-0.7, cultures were treated with the indicated

concentration of TTM, or DMSO vehicle control, for 24 hrs. Cells were collected by

centrifugation (3700 x *g*, 10 min) washed once with 1mL of lysis buffer (350 mM sodium

chloride, 20 mM Tris pH 8.0, 1 mM 2-mercaptoethanol) prior to suspension in 0.8 mL of lysis

buffer plus ~100µL of zirconia beads. Lysis was performed by 3x45 sec pulses in a BioSpec

Mini24 beadbeater with 5 min intervening rest periods on ice. Beads and debris were removed

by centrifugation at 20,000 x *g* for 15 min at 4°C, the resulting supernatant was mixed 1:1 with

2x Laemmli sample buffer supplemented with 0.1 M dithiothreitol (DTT). 20 µL of each sample,

heated for 10 min at 100ºC, was then separated on 4-12% NuPAGE Bis-Tris poly acrylamide

gels. Separated proteins were transferred to nitrocellulose and probed with the indicated

antibodies. Antibodies used in this study are monoclonal anti-*E. coli* RNA -polymerase β

(BioLegend), sdAb anti-ALFA tag-HRP (NanoTag Biotechnologies). Chemiluminescence was

visualized using SuperSignal™West Pico Plus chemiluminescent substrate (Thermo Scientific).

Blots were imaged on an iBright FL1000 imager (Themo Fisher Scientific). CtaD ALFA blots

were quantitated using ImageJ software. Relative ALFA signal was normalized to corresponding

Rpoβ loading control levels.

**Methylene blue assay**

Cultures were grown in 7H9-ADS to an $OD_{600}$ of 0.5-0.7, washed twice in Chelex treated PBS

Tween-80, and adjusted to an $OD_{600}$ of 0.15 with fresh 7H9-ADS and incubated with indicated

concentrations of TTM, Q203, ND-011992, or DMSO for 4 hrs. Following incubation, 4-ml

screw☐cap glass vials were filled with the cultures and methylene blue dye was added at a final

concentration of 0.001%. All vials were sealed tightly using a PTFE/rubber seal (Thermo

Scientific) and incubated at 37°C in an anaerobic container (Benton Dickinson GasPak EZ

container systems) for 3 days.

**Oxygen Consumption Measurements**

Oxygen consumption was assessed using PSt6 sensor spots and a Fibox 4 trace instrument

(PreSens Precision Sensing GmbH Am BioPark, Germany)(36). Screw cap glass vials (Thermo

Scientific) containing the sensor spot PSt6 were filled with cultures grown in 7H9-ADS at a

calculated $OD_{600}$ of 0.005 and the indicated concentrations of TTM, Q203, ND-011992, or

DMSO. The vials were tightly sealed using a PTFE/rubber seal (Thermo Scientific) and

incubated at 37°C in an anaerobic pouch (Benton Dickinson GasPak EZ pouch systems).

Percent oxygen was measured daily via the Fibox 4 trace oxygen meter without unsealing the

anaerobic environment.

**ATP quantitation**

ATP levels were quantified with the BacTiter-Glo$^{TM}$ (Promega). The indicated strains were first

pre-grown to an $OD_{600}$ of 0.5-1.0 in 7H9-OADC. Cells were then washed twice with PBS

Tween-80 and suspended in 7H9-OADC at an $OD_{600}$ of 0.05. 96-Well plates were inoculated

with 100 $\mu$L of washed culture in replicate wells. The indicated concentrations of TTM, Q203, or

DMSO vehicle control were then added, and plates were incubated at 37°C, 0.05% $CO_2$ for 24

hrs. 100 $\mu$L of BacTiter-Glo was then added to each well and plates were incubated 15

additional mins at 37°C, 5% $CO_2$ prior to reading on a SpectraMax M3 plate reader. [ATP] in

test samples was quantified against standard curves of ATP.

**Synthesis of diisonitrile chalkophores**

Details of synthesis and characterization are provided in SI Appendix 1.

**Aerosol infection of mice**

8–10-week-old C57BL/6J (JAX stock number 00064) and 8-10 week B6.Cg-$Prkdc^{scid}$/SzJ (stock number 001913, RRID: IMSR_JAX:001913) were purchased from The Jackson Laboratory. All purchased mice were rested within our animal facility to normalize microbiota for 2 weeks. Care, housing, and experimentation on laboratory mice were performed in accordance with the National Institute of Heath guidelines, and the approval of the Memorial Sloan Kettering Institutional Animal Care and Use Committee (IACUC). Strains for infection were grown to an $OD_{600}$ of 0.5-0.7 in 7H9-OADC, washed twice with PBS Tween-80 followed by brief sonication to disrupt aggregates. Final inoculums were prepared by suspending $8x10^7$ CFU in 10 mL of sterile water.  Mice were exposed to $4x10^7$ CFU in a Glas-Col aerosol exposure unit, a dose calibrated to deliver 10-50 CFU per mouse. At the indicated time points post infection, 5 individual mice were humanely euthanized and both lungs and spleens were harvested for CFU determination. Organ homogenates were cultured on 7H10-OADC plates, and CFU was enumerated after 28 days of incubation at 37ºC, 5% $CO_2$. 20 additional SCID mice (10: WT *M. tuberculosis*, 10: Δ*nrp*Δ*cydAB*) infected as described above, were used to assess survival.  Infected mice were monitored daily and humanely euthanized upon frank signs of morbidity.

**Neutrophil depletion**

Mice were treated intraperitoneally with anti-Ly6G (BioXcell, clone 1A8, RRID AB_1107721) or isotype control (BioXcell, clone 2A3, RRID AB_1107769) every 48 hrs. Lungs were harvested aseptically and single cell suspensions prepared by homogenization in a Bullet Blender with 2.0 mM zirconium oxide beads for 3 min followed by 100-micron filtration. After lysis of red blood cells with Gibco ACK lysing buffer and blocking of Fc receptors (Invitrogen CD16/CD32 clone 93, RRID AB_467133), cells were stained with live/dead aqua (Thermo) and antibodies to CD45 (FITC, clone 30-F11, BD Biosciences, RRID AB_394609), CD11b (BUV661, clone M1/70, BD Biosciences), CD11c (APC-R700, clone N418, RRID AB_2744277), Ly6C (BV605, clone AL-21, BD Biosciences, AB_2737949), CD80 (PerCP Cy5.5, clone 16-10A1 AB_1727514), CD86 (Pacific Blue, clone GL-1, BioLegend, RRID AB_493466), MHC II (BUV395, clone 2G9, BD Biosciences RRID AB_2741827), GR1 (PE, clone RB6-8C5, BD Biosciences, RRID AB_398532), and Ly6G (APC, clone 1A8, BD Biosciences, RRID AB_394208) and analyzed by flow cytometry on a BD Fortessa cytometer after paraformaldehyde fixation.

**Acknowledgements**

This work was supported by R01AI138446 and P30 CA008748. We acknowledge the use of the Integrated Genomics Operation Core, funded by the NCI Cancer Center Support Grant (CCSG, P30 CA08748), Cycle for Survival, and the Marie-Josée and Henry R. Kravis Center for Molecular Oncology. We thank Christina Stallings for advice about neutrophil depletion and George Sukenick and Rui Wang (MSK) for expert NMR and mass spectral support.

**Author Contributions**

J.A.B., Y.O., and M.S.G. conceived and planned the experiments. C.H. synthesized the diisonitrile analogue under the supervision of D.T.  J.A.B., Y.O., and M.S.G. carried out the experiments with support and technical expertise from A.B. for neutrophil depletion.  J.A.B., Y.O., C.H., D.T., and M.S.G. contributed to the interpretation of the results. M.S.G. took the lead in writing the manuscript. All authors provided critical feedback and helped shape the research, analysis, and manuscript.

**Conflicts of Interest**

MSG declares equity and consulting fees from Vedanta biosciences and consulting fees from Fimbrion therapeutics. DST declares no conflicts of interest pertaining to this work.

**Figure 1 Copper deprivation in chalkophore-deficient _M. tuberculosis_ mimics _bcc:aa₃_ oxidase inhibition**

**A.** Heat map of transcripts encoding selected respiratory chain components determined by RNA sequencing of _M. tuberculosis_ WT or Δ*nrp* treated with TTM or Q203. WT_GSE is the published dataset GSE159080 of _M. tuberculosis_ H37Rv treated with Q203.

492      **B.** RT-qPCR of the transcript encoding CydA in *M. tuberculosis* WT, Δ*nrp*, and

493           complemented strain treated with varying TTM concentrations for 4 hours. Error bars

494           represent standard error of the mean (SEM).  Statistical significance determined via two-

495           way ANOVA with Tukey correction for multiple comparisons.  $*=p<0.05$, $**=p<0.01$, $****=$

496           $p<0.0001$.

497      **C.** RT-qPCR of the transcript encoding CydA in *M. tuberculosis* WT, Δ*nrp*, and

498           complemented strain treated with 20 μM TTM for 24 hours.  Error bars are SEM.

499           Statistical significance determined via two-way ANOVA with Tukey correction for multiple

500           comparisons.  $**=p<0.01$, $***=p<0.001$, $****=p<0.0001$.

501      **D.** Dose dependent effect of TTM on growth of the indicated *M. tuberculosis* strains at 7

502           days post inoculation. The dotted line indicates the starting inoculum.  Error bars are

503           SEM.  Statistical significance determined via two-way ANOVA with Tukey correction for

504           multiple comparisons. $*=p<0.05$.

**Figure 2 Chalkophores maintain *M. tuberculosis* viability through the heme-copper**

*bcc:aa₃* **oxidase during copper starvation**

      **A. Schematic of the terminal respiratory oxidases of *M. tuberculosis*.** The *bcc:aa₃*

oxidase is a heme-copper oxidase and CytBD is a copper-independent heme oxidase.

Both transfer electrons to oxygen. Q203 is an inhibitor of *bcc:aa₃* by targeting the QcrB

subunit, whereas ND-011992 targets CytBD. The two oxidases are individually

dispensable due to compensation by the other oxidase, but *M. tuberculosis* lacking both

is nonviable. The model to be tested is that copper chelation deprives the *bcc:aa₃*

oxidase of copper and that diisonitrile chalkophores counter this copper deprivation
stress.
**B.** Liquid growth assays of the indicated strains with or without 20 µM TTM treatment.
$OD_{600}$ at day 10 post inoculation displayed. Dotted line indicates starting inoculum. Error
bars are SEM. Statistical significance determined via two-way ANOVA with Tukey
correction for multiple comparisons. ****= $p<0.0001$.
**C.** Bacterial survival of the indicated strains on agar media containing DMSO, 1 mM
BCS or 40 µM TTM. Dotted line indicates lower limit of detection (LLOD). Error bars are
SEM. Statistical significance determined via two-way ANOVA with Tukey correction for
multiple comparisons. **= $p<0.01$, *=$p<0.05$
**D. The copper deprivation sensitivity of *M. tuberculosis* Δ*nrp*Δ*cydAB* strain can be**
**rescued with a synthetic diisonitrile chalkophore.** Liquid growth assays of
Δ*nrp*Δ*cydAB* with DMSO, 20 µM TTM, or 20 µM TTM with 10 µM of the diisonitrile
chalkophore pictured in panel A. Error bars are SEM.
**E. The *bcc:aa₃* oxidase is the only target of copper starvation countered by**
**diisonitrile chalkophores.** Liquid growth assays of the indicated strains treated with 10
or 20 µM TTM, or DMSO vehicle control. $OD_{600}$ at day 10 post inoculation displayed.
Dotted line indicates starting inoculum. Error bars are SEM. Statistical significance
determined via two-way ANOVA with Tukey correction for multiple comparisons. ns= not
significant, ****= $p<0.0001$.
**F. The effect of copper deprivation is masked by inhibition of QcrB subunit of**
***bcc*:*aa₃*.** Liquid growth assays of the indicated strains treated with TTM or Q203 alone or
co-treated with both at the indicated concentrations. $OD_{600}$ at day 7 post inoculation
displayed. Dotted line indicates starting inoculum. Error bars are SEM. Statistical
significance determined via two-way ANOVA with Tukey correction for multiple
comparisons. ns= not significant.
540

**Figure 3 Chalkophore biosynthesis maintains oxidative phosphorylation through the heme-copper *bcc:aa₃* oxidase**

    **A.** Methylene blue decolorization assay of oxygen consumption under copper deprivation (TTM) or treatment with Q203 in WT or Δ*nrp*Δ*cydAB M. tuberculosis*. Clear vials indicate oxygen consumption by respiration.

**B.** Quantitative measurement of oxygen consumption using oxygen sensitive optical

sensors. WT *M. tuberculosis* treated with DMSO, ND-011992, Q203, or both ND-011992

and Q203. Oxygen measurements were taken daily. Each point represents three

measurements of two biologic replicates. Error bars are SEM. Statistical significance

between Q203 and ND-011992 + Q203 determined via two-way ANOVA with Tukey

correction for multiple comparisons. ****= $p<0.0001$.

**C.** Same assay as in panel B with WT and Δ*nrp M. tuberculosis* treated with DMSO or 25

μM TTM. Error bars are SEM.

**D.** Same assay as in panel B with WT, Δ*nrp*Δ*cydAB,* or Δ*nrp*Δ*cydAB+ nrp* treated with 25

μM TTM.  Error bars are SEM. Statistical significance between WT and Δ*nrp*Δ*cydAB*

treated with 25 μM TTM determined via two-way ANOVA with Tukey correction for

multiple comparisons. ****= $p<0.0001$.

**E.** Cellular ATP levels determined by BacTiter-Glo in the indicated strains treated with

DMSO, 20 or 40 μM TTM, or 100 nM Q203. [ATP] determined by standard curve

determined in growth media containing the same quantities of DMSO, TTM or Q203.

Error bars are SEM.  Statistical significance determined via two-way ANOVA with Tukey

correction for multiple comparisons. *= $p<0.05$, ***=$p<0.001$, ****=$p<0.0001$.

**F.** Relative abundance of a CtaD-ALFA protein in *M. tuberculosis* of the indicated genotype

treated with BCS or TTM. See Figure S4 for primary immunoblot data. Error bars are

SEM. Statistical significance determined via two-way ANOVA with Tukey correction for

multiple comparisons. ns= not significant.

**Figure 4 Respiratory chain flexibility is critical for *M. tuberculosis* virulence**

A, B.        Bacterial titers in the lung (A) or spleen (B) in mice infected with *M. tuberculosis*

WT, Δ*nrp*, Δ*nrp*Δ*cydAB,* or Δ*nrp*Δ*cydAB*+ *nrp.*  Error bars are SEM. Statistical significance

determined via two-way ANOVA with Tukey correction for multiple comparisons.  Not

significant (ns), **=p<0.01, and ***=p<0.001.

C, D.        **Copper deprivation by the host is independent of neutrophils**. Bacterial titers

in the lung (C) or spleen (D) in mice infected with *M. tuberculosis* WT or Δ*nrp*Δ*cydAB,* or

*ΔnrpΔcydAB* + *nrp* treated with isotype control antibodies or anti-Ly6G antibodies to deplete

neutrophils. Flow cytometric quantitation of neutrophil depletion is provided in Figure S5.

Error bars are SEM.

E, F.        Copper deprivation by the host is independent of adaptive immunity. Bacterial

titers in the lung (E) or spleen (F) in C57BL/6J or C57BL/6 SCID mice infected with *M.*

*tuberculosis* WT or *ΔnrpΔcydAB.* Error bars are SEM.

588
589
590

**References**

1.  Murdoch CC, Skaar EP. 2022. Nutritional immunity: the battle for nutrient metals at the host–pathogen interface. Nature Reviews Microbiology doi:10.1038/s41579-022-00745-6.

2.  Sheldon JR, Laakso HA, Heinrichs DE. 2016. Iron Acquisition Strategies of Bacterial Pathogens. Microbiology Spectrum 4:10.1128/microbiolspec.vmbf-0010-2015.

3.  Behnsen J, Zhi H, Aron AT, Subramanian V, Santus W, Lee MH, Gerner RR, Petras D, Liu JZ, Green KD, Price SL, Camacho J, Hillman H, Tjokrosurjo J, Montaldo NP, Hoover EM, Treacy-Abarca S, Gilston BA, Skaar EP, Chazin WJ, Garneau-Tsodikova S, Lawrenz MB, Perry RD, Nuccio SP, Dorrestein PC, Raffatellu M. 2021. Siderophore-mediated zinc acquisition enhances enterobacterial colonization of the inflamed gut. Nat Commun 12:7016.

4.  Bobrov AG, Kirillina O, Fetherston JD, Miller MC, Burlison JA, Perry RD. 2014. The Y ersinia pestis siderophore, yersiniabactin, and the ZnuABC system both contribute to zinc acquisition and the development of lethal septicaemic plague in mice. Molecular microbiology 93:759-775.

5.  Diaz-Ochoa VE, Lam D, Lee CS, Klaus S, Behnsen J, Liu JZ, Chim N, Nuccio SP, Rathi SG, Mastroianni JR, Edwards RA, Jacobo CM, Cerasi M, Battistoni A, Ouellette AJ, Goulding CW, Chazin WJ, Skaar EP, Raffatellu M. 2016. Salmonella Mitigates Oxidative Stress and Thrives in the Inflamed Gut by Evading Calprotectin-Mediated Manganese Sequestration. Cell Host Microbe 19:814-25.

6.  Liu JZ, Jellbauer S, Poe AJ, Ton V, Pesciaroli M, Kehl-Fie TE, Restrepo NA, Hosking MP, Edwards RA, Battistoni A, Pasquali P, Lane TE, Chazin WJ, Vogl T, Roth J, Skaar EP, Raffatellu M. 2012. Zinc sequestration by the neutrophil protein calprotectin enhances Salmonella growth in the inflamed gut. Cell Host Microbe 11:227-39.

7.  Nairn BL, Lonergan ZR, Wang J, Braymer JJ, Zhang Y, Calcutt MW, Lisher JP, Gilston BA, Chazin WJ, de Crécy-Lagard V, Giedroc DP, Skaar EP. 2016. The Response of Acinetobacter baumannii to Zinc Starvation. Cell Host Microbe 19:826-36.

8.  Sheldon JR, Skaar EP. 2019. Metals as phagocyte antimicrobial effectors. Curr Opin Immunol 60:1-9.

9.  Shi X, Darwin KH. 2015. Copper homeostasis in Mycobacterium tuberculosis. Metallomics 7:929-34.

624     10.     Shi X, Festa RA, Ioerger TR, Butler-Wu S, Sacchettini JC, Darwin KH, Samanovic MI.
625            2014. The copper-responsive RicR regulon contributes to Mycobacterium tuberculosis
626            virulence. mBio 5.

627     11.     Festa RA, Jones MB, Butler-Wu S, Sinsimer D, Gerads R, Bishai WR, Peterson SN,
628            Darwin KH. 2011. A novel copper-responsive regulon in Mycobacterium tuberculosis.
629            Mol Microbiol 79:133-48.

630     12.     Buglino JA, Sankhe GD, Lazar N, Bean JM, Glickman MS. 2021. Integrated sensing of
631            host stresses by inhibition of a cytoplasmic two-component system controls M.
632            tuberculosis acute lung infection. Elife 10.

633     13.     Rowland JL, Niederweis M. 2012. Resistance mechanisms of Mycobacterium
634            tuberculosis against phagosomal copper overload. Tuberculosis (Edinb) 92:202-10.

635     14.     Wolschendorf F, Ackart D, Shrestha TB, Hascall-Dove L, Nolan S, Lamichhane G, Wang
636            Y, Bossmann SH, Basaraba RJ, Niederweis M. 2011. Copper resistance is essential for
637            virulence of *Mycobacterium tuberculosis*. Proceedings of the National Academy of
638            Sciences 108:1621-1626.

639     15.     Wagner D, Maser J, Lai B, Cai Z, Barry CE, Höner zu Bentrup K, Russell DG, Bermudez
640            LE. 2005. Elemental Analysis of <em>Mycobacterium avium</em>-,
641            <em>Mycobacterium tuberculosis</em>-, and <em>Mycobacterium smegmatis</em>-
642            Containing Phagosomes Indicates Pathogen-Induced Microenvironments within the Host
643            Cell's Endosomal System. The Journal of Immunology 174:1491-1500.

644     16.     Dow A, Sule P, O'Donnell TJ, Burger A, Mattila JT, Antonio B, Vergara K, Marcantonio
645            E, Adams LG, James N, Williams PG, Cirillo JD, Prisic S. 2021. Zinc limitation triggers
646            anticipatory adaptations in Mycobacterium tuberculosis. PLoS Pathog 17:e1009570.

647     17.     Harris NC, Born DA, Cai W, Huang Y, Martin J, Khalaf R, Drennan CL, Zhang W. 2018.
648            Isonitrile Formation by a Non-Heme Iron(II)-Dependent Oxidase/Decarboxylase.
649            Angewandte Chemie International Edition 57:9707-9710.

650     18.     Wang L, Zhu M, Zhang Q, Zhang X, Yang P, Liu Z, Deng Y, Zhu Y, Huang X, Han L, Li
651            S, He J. 2017. Diisonitrile Natural Product SF2768 Functions As a Chalkophore That
652            Mediates Copper Acquisition in Streptomyces thioluteus. ACS Chem Biol 12:3067-3075.

653     19.     Harris NC, Sato M, Herman NA, Twigg F, Cai W, Liu J, Zhu X, Downey J, Khalaf R,
654            Martin J, Koshino H, Zhang W. 2017. Biosynthesis of isonitrile lipopeptides by conserved
655            nonribosomal peptide synthetase gene clusters in Actinobacteria. Proc Natl Acad Sci U
656            S A 114:7025-7030.

657    20.    Mehdiratta K, Singh S, Sharma S, Bhosale RS, Choudhury R, Masal DP, Manocha A,
658             Dhamale BD, Khan N, Asokachandran V, Sharma P, Ikeh M, Brown AC, Parish T, Ojha
659             AK, Michael JS, Faruq M, Medigeshi GR, Mohanty D, Reddy DS, Natarajan VT, Kamat
660             SS, Gokhale RS. 2022. Kupyaphores are zinc homeostatic metallophores required for
661             colonization of Mycobacterium tuberculosis. Proc Natl Acad Sci U S A 119.

662    21.    Buglino JA, Ozakman Y, Xu Y, Chowdhury F, Tan DS, Glickman MS. 2022. Diisonitrile
663             Lipopeptides Mediate Resistance to Copper Starvation in Pathogenic Mycobacteria.
664             mBio 13:e0251322.

665    22.    Kalia NP, Hasenoehrl EJ, Ab Rahman NB, Koh VH, Ang MLT, Sajorda DR, Hards K,
666             Grüber G, Alonso S, Cook GM, Berney M, Pethe K. 2017. Exploiting the synthetic
667             lethality between terminal respiratory oxidases to kill <em>Mycobacterium
668             tuberculosis</em> and clear host infection. Proceedings of the National Academy of
669             Sciences 114:7426-7431.

670    23.    Matsoso LG, Kana BD, Crellin PK, Lea-Smith DJ, Pelosi A, Powell D, Dawes SS, Rubin
671             H, Coppel RL, Mizrahi V. 2005. Function of the cytochrome bc1-aa3 branch of the
672             respiratory network in mycobacteria and network adaptation occurring in response to its
673             disruption. J Bacteriol 187:6300-8.

674    24.    Lamprecht DA, Finin PM, Rahman MA, Cumming BM, Russell SL, Jonnala SR,
675             Adamson JH, Steyn AJ. 2016. Turning the respiratory flexibility of Mycobacterium
676             tuberculosis against itself. Nat Commun 7:12393.

677    25.    Harrison GA, Mayer Bridwell AE, Singh M, Jayaraman K, Weiss LA, Kinsella RL, Aneke
678             JS, Flentie K, Schene ME, Gaggioli M, Solomon SD, Wildman SA, Meyers MJ, Stallings
679             CL. 2019. Identification of 4-Amino-Thieno[2,3-d]Pyrimidines as QcrB Inhibitors in
680             Mycobacterium tuberculosis. mSphere 4.

681    26.    O'Malley T, Alling T, Early JV, Wescott HA, Kumar A, Moraski GC, Miller MJ, Masquelin
682             T, Hipskind PA, Parish T. 2018. Imidazopyridine Compounds Inhibit Mycobacterial
683             Growth by Depleting ATP Levels. Antimicrob Agents Chemother 62.

684    27.    Lupien A, Foo CS, Savina S, Vocat A, Piton J, Monakhova N, Benjak A, Lamprecht DA,
685             Steyn AJC, Pethe K, Makarov VA, Cole ST. 2020. New 2-Ethylthio-4-
686             methylaminoquinazoline derivatives inhibiting two subunits of cytochrome bc1 in
687             Mycobacterium tuberculosis. PLoS Pathog 16:e1008270.

688    28.    Foo CS, Lupien A, Kienle M, Vocat A, Benjak A, Sommer R, Lamprecht DA, Steyn AJC,
689             Pethe K, Piton J, Altmann KH, Cole ST. 2018. Arylvinylpiperazine Amides, a New Class
690             of Potent Inhibitors Targeting QcrB of Mycobacterium tuberculosis. mBio 9.

29.   Lee BS, Hards K, Engelhart CA, Hasenoehrl EJ, Kalia NP, Mackenzie JS, Sviriaeva E,
Chong SMS, Manimekalai MSS, Koh VH, Chan J, Xu J, Alonso S, Miller MJ, Steyn AJC,
Gruber G, Schnappinger D, Berney M, Cook GM, Moraski GC, Pethe K. 2021. Dual
inhibition of the terminal oxidases eradicates antibiotic-tolerant Mycobacterium
tuberculosis. EMBO Mol Med 13:e13207.

30.   Gong H, Li J, Xu A, Tang Y, Ji W, Gao R, Wang S, Yu L, Tian C, Li J, Yen HY, Man Lam
S, Shui G, Yang X, Sun Y, Li X, Jia M, Yang C, Jiang B, Lou Z, Robinson CV, Wong LL,
Guddat LW, Sun F, Wang Q, Rao Z. 2018. An electron transfer path connects subunits
of a mycobacterial respiratory supercomplex. Science 362.

31.   Kim MS, Jang J, Ab Rahman NB, Pethe K, Berry EA, Huang LS. 2015. Isolation and
Characterization of a Hybrid Respiratory Supercomplex Consisting of Mycobacterium
tuberculosis Cytochrome bcc and Mycobacterium smegmatis Cytochrome aa3. J Biol
Chem 290:14350-60.

32.   Mathiyazakan V, Wong CF, Harikishore A, Pethe K, Grüber G. 2023. Cryo-Electron
Microscopy Structure of the Mycobacterium tuberculosis Cytochrome bcc:aa(3)
Supercomplex and a Novel Inhibitor Targeting Subunit Cytochrome cI. Antimicrob
Agents Chemother 67:e0153122.

33.   Wiseman B, Nitharwal RG, Fedotovskaya O, Schäfer J, Guo H, Kuang Q, Benlekbir S,
Sjöstrand D, Ädelroth P, Rubinstein JL, Brzezinski P, Högbom M. 2018. Structure of a
functional obligate complex III(2)IV(2) respiratory supercomplex from Mycobacterium
smegmatis. Nat Struct Mol Biol 25:1128-1136.

34.   Safarian S, Opel-Reading HK, Wu D, Mehdipour AR, Hards K, Harold LK, Radloff M,
Stewart I, Welsch S, Hummer G. 2021. The cryo-EM structure of the bd oxidase from M.
tuberculosis reveals a unique structural framework and enables rational drug design to
combat TB. Nature Communications 12:5236.

35.   Xu Y, Tan DS. 2019. Total Synthesis of the Bacterial Diisonitrile Chalkophore SF2768.
Org Lett 21:8731-8735.

36.   Kalia NP, Singh S, Hards K, Cheung CY, Sviriaeva E, Banaei-Esfahani A, Aebersold R,
Berney M, Cook GM, Pethe K. 2023. M. tuberculosis relies on trace oxygen to maintain
energy homeostasis and survive in hypoxic environments. Cell Rep 42:112444.

37.   Soma S, Latimer AJ, Chun H, Vicary AC, Timbalia SA, Boulet A, Rahn JJ, Chan SSL,
Leary SC, Kim BE, Gitlin JD, Gohil VM. 2018. Elesclomol restores mitochondrial function
in genetic models of copper deficiency. Proc Natl Acad Sci U S A 115:8161-8166.

38.   Gotzke H, Kilisch M, Martinez-Carranza M, Sograte-Idrissi S, Rajavel A, Schlichthaerle

T, Engels N, Jungmann R, Stenmark P, Opazo F, Frey S. 2019. The ALFA-tag is a

highly versatile tool for nanobody-based bioscience applications. Nat Commun 10:4403.

39.   Conradie F, Bagdasaryan TR, Borisov S, Howell P, Mikiashvili L, Ngubane N, Samoilova

A, Skornykova S, Tudor E, Variava E. 2022. Bedaquiline–pretomanid–linezolid regimens

for drug-resistant tuberculosis. New England Journal of Medicine 387:810-823.

40.   Conradie F, Diacon AH, Ngubane N, Howell P, Everitt D, Crook AM, Mendel CM, Egizi

E, Moreira J, Timm J. 2020. Bedaquiline, pretomanid and linezolid for treatment of

extensively drug resistant, intolerant or non-responsive multidrug resistant pulmonary

tuberculosis. The New England Journal of Medicine 382:893.

41.   de Jager VR, Dawson R, van Niekerk C, Hutchings J, Kim J, Vanker N, van der Merwe

L, Choi J, Nam K, Diacon AH. 2020. Telacebec (Q203), a new antituberculosis agent.

New England Journal of Medicine 382:1280-1281.

42.   Beites T, O'Brien K, Tiwari D, Engelhart CA, Walters S, Andrews J, Yang HJ, Sutphen

ML, Weiner DM, Dayao EK, Zimmerman M, Prideaux B, Desai PV, Masquelin T, Via LE,

Dartois V, Boshoff HI, Barry CE, 3rd, Ehrt S, Schnappinger D. 2019. Plasticity of the

Mycobacterium tuberculosis respiratory chain and its impact on tuberculosis drug

development. Nat Commun 10:4970.

43.   Cai Y, Jaecklein E, Mackenzie JS, Papavinasasundaram K, Olive AJ, Chen X, Steyn

AJC, Sassetti CM. 2021. Host immunity increases Mycobacterium tuberculosis reliance

on cytochrome bd oxidase. PLoS Pathog 17:e1008911.

44.   Bhatt K, Machado H, Osorio NS, Sousa J, Cardoso F, Magalhaes C, Chen B, Chen M,

Kim J, Singh A, Ferreira CM, Castro AG, Torrado E, Jacobs WR, Jr., Bhatt A, Saraiva M.

2018. A Nonribosomal Peptide Synthase Gene Driving Virulence in Mycobacterium

tuberculosis. mSphere 3.

45.   Botella H, Peyron P, Levillain F, Poincloux R, Poquet Y, Brandli I, Wang C, Tailleux L,

Tilleul S, Charriere GM, Waddell SJ, Foti M, Lugo-Villarino G, Gao Q, Maridonneau-

Parini I, Butcher PD, Castagnoli PR, Gicquel B, de Chastellier C, Neyrolles O. 2011.

Mycobacterial p(1)-type ATPases mediate resistance to zinc poisoning in human

macrophages. Cell Host Microbe 10:248-59.

46.   Boudehen YM, Faucher M, Marechal X, Miras R, Rech J, Rombouts Y, Seneque O,

Wallat M, Demange P, Bouet JY, Saurel O, Catty P, Gutierrez C, Neyrolles O. 2022.

Mycobacterial resistance to zinc poisoning requires assembly of P-ATPase-containing

membrane metal efflux platforms. Nat Commun 13:4731.

47.  Darwin KH. 2015. Mycobacterium tuberculosis and Copper: A Newly Appreciated
Defense against an Old Foe? J Biol Chem 290:18962-6.
48.  Chen T-Y, Chen J, Ruszczycky MW, Hilovsky D, Hostetler T, Liu X, Zhou J, Chang W-c.
2024. Variation in Biosynthesis and Metal-Binding Properties of Isonitrile-Containing
Peptides Produced by Mycobacteria versus Streptomyces. ACS Catalysis 14:4975-
4983.
49.  Shi L, Sohaskey C, Kana B, Dawes S, North R, Mizrahi V, Gennaro M. 2005. Changes
in energy metabolism of Mycobacterium tuberculosis in mouse lung and under in vitro
conditions affecting aerobic respiration. Proceedings of the National Academy of
Sciences of the United States of America 102:15629-15634.
50.  Harikishore A, Mathiyazakan V, Pethe K, Grüber G. 2023. Novel targets and inhibitors of
the Mycobacterium tuberculosis cytochrome bd oxidase to foster anti-tuberculosis drug
discovery. Expert Opinion on Drug Discovery 18:917-927.
51.  Andrews S. 2010. FastQC: a quality control tool for high throughput sequence data.
http://www.bioinformatics.babraham.ac.uk/projects/fastqc. Accessed
52.  Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, Marth G, Abecasis G,
Durbin R. 2009. The Sequence Alignment/Map format and SAMtools. Bioinformatics
25:2078-9.
53.  Li H, Durbin R. 2009. Fast and accurate short read alignment with Burrows-Wheeler
transform. Bioinformatics 25:1754-60.
54.  Liao Y, Smyth GK, Shi W. 2014. featureCounts: an efficient general purpose program for
assigning sequence reads to genomic features. Bioinformatics 30:923-30.
55.  Love MI, Huber W, Anders S. 2014. Moderated estimation of fold change and dispersion
for RNA-seq data with DESeq2. Genome Biol 15:550.
56.  Kolde R. pheatmap, https://rdocumentation.org/packages/pheatmap/versions/1.0.12.
57.  R Core Team R. 2013. R: A language and environment for statistical computing.
784
785

**Supplementary Materials**

Figures S1-S5

Supplementary Tables:

Table S1: strains used

Table S2: plasmids used

Table S3: RNA sequencing dataset

SI appendix 1: chemical synthesis methods