Order of amino acid recruitment into the genetic code resolved by
Last Universal Common Ancestor's protein domains

Sawsan Wehbi[1], Andrew Wheeler[1], Benoit Morel[2], Nandini Manepalli[3], Bui Quang Minh[4], Dante
S. Lauretta[5], Joanna Masel[6]

[1]Genetics Graduate Interdisciplinary Program, University of Arizona, Tucson, Arizona, 85721,
USA

[2]Computational Molecular Evolution Group, Heidelberg Institute for Theoretical Studies,
Heidelberg, Germany

[3]Department of Molecular and Cellular Biology, University of Arizona, Tucson, AZ, 85721, USA

[4]School of Computing, Australian National University, Canberra, ACT, Australia

[5]Lunar and Planetary Laboratory, University of Arizona, Tucson, AZ 85721, USA

[6]Department of Ecology and Evolutionary Biology, University of Arizona, Tucson, AZ, 85721,
USA

Corresponding author: Joanna Masel

Email: masel@arizona.edu

Author Contributions: S.W. and J.M. conceived the study, with input from D.S.L. N.M. produced
Figure 3, and S.W. conducted all other analyses, with advice from J.M., A.W., B.Q.M., and B.M.
B.M. implemented a new feature in GeneRax to assist the analyses. S.W. wrote the first draft of
the manuscript, with subsequent editing from S.W., J.M., D.S.L., A.W., and B.Q.M.

Competing Interest Statement: The authors declare no competing interest.

Classification: Biological Sciences, Evolution.

Keywords: Origins of life, early life, astrobiology, phylostratigraphy, antioxidants, metalloproteins

Abstract

The current "consensus" order in which amino acids were added to the genetic code is based on potentially biased criteria, such as absence of sulfur-containing amino acids from the Urey-Miller experiment which lacked sulfur. More broadly, abiotic abundance might not reflect biotic abundance in the organisms in which the genetic code evolved. Here, we instead identify which protein domains date to the last universal common ancestor (LUCA), then infer the order of recruitment from deviations of their ancestrally reconstructed amino acid frequencies from the still-ancient post-LUCA controls. We find that smaller amino acids were added to the code earlier, with no additional predictive power in the previous "consensus" order. Metal-binding (cysteine and histidine) and sulfur-containing (cysteine and methionine) amino acids were added to the genetic code much earlier than previously thought. Methionine and histidine were added to the code earlier than expected from their molecular weights, and glutamine later. Early methionine availability is compatible with inferred early use of S-adenosylmethionine, and early histidine with its purine-like structure and the demand for metal-binding. Even more ancient protein sequences — those that had already diversified into multiple distinct copies prior to LUCA — have significantly higher frequencies of aromatic amino acids (tryptophan, tyrosine, phenylalanine and histidine), and lower frequencies of valine and glutamic acid than single copy LUCA sequences. If at least some of these sequences predate the current code, then their distinct enrichment patterns provide hints about earlier, alternative genetic codes.

Significance Statement

The order in which the amino acids were added to the genetic code was previously inferred from consensus among forty metrics. Many of these reflect abiotic abundance on ancient Earth. However, the abundances that matter are those within primitive cells that already had sophisticated RNA and perhaps peptide metabolism. Here, we directly infer the order of recruitment from the relative ancestral amino acid frequencies of ancient protein sequences. Small size predicts ancient amino acid enrichment better than the previous consensus metric does. We place metal-binding and sulfur-containing amino acids earlier than previously thought, highlighting the importance of metal-dependent catalysis and sulfur metabolism to ancient life. Understanding early life has implications for our search for life elsewhere in the universe.

Main Text

Introduction

The modern genetic code was likely assembled in stages, hypothesized to begin with "early" amino acids present on Earth before the emergence of life (possibly delivered by extraterrestrial sources such as asteroids or comets), and ending with "late" amino acids requiring biotic synthesis (1, 2). For example, the Urey-Miller experiment (3) has been used to identify which amino acids were available abiotically and are thus likely to have come earlier than those requiring biotic synthesis. The order of amino acid recruitment, from early to late, was inferred by taking statistical consensus among 40 different rankings (4), none of which constitute strong evidence on their own. On the basis of this ordering, Moosmann (5) hypothesized that the first amino acids recruited into the genetic code were those that were useful for membrane anchoring, then those useful for halophilic folding, then for mesophilic folding, then for metal binding, and finally for their antioxidant properties. However, a late role for metal-binding amino acids is puzzling; many metalloproteins date back to the Last Universal Common Ancestor's (LUCA)'s proteome, where they are presumed to be key to the emergence of biological catalysis (6).

Indeed, the late status of some amino acids is disputed (7). For example, the Urey-Miller experiment (3) did not include sulfur, and so should not have been used to infer that the sulfur-containing amino acids cysteine and methionine were late additions. Methionine and homocysteine (a product of cysteine degradation) were detected in hydrogen sulfide (H_2S)-rich

spark discharge experiments, suggesting that methionine and cysteine could be abiotically
produced (8). A nitrile-activated dehydroalanine pathway can produce cysteine from abiotic serine
that is produced from a Strecker reaction (9), further demonstrating the possibility of its early
chemical availability.

Histidine's classification as abiotically unavailable also contributed to its annotation as late (4).
While histidine can be abiotically synthesized from erythrose reacting with formamidine followed
by a Strecker synthesis reaction (10), the reactant concentrations might have been insufficient in
a primitive earth environment (11). More importantly, because histidine resembles a purine, even
if histidine were abiotically unavailable, it might have had cellular availability at the time of genetic
code construction (12), in an organism that biotically synthesized ribosomes, and that might also
have already utilized amino acids and peptides. Indeed, histidine is the most commonly
conserved residue in the active site of enzymes (13).

To directly infer the order of recruitment from protein sequence data, without reference to abiotic
availability arguments, we consider that some of LUCA's proteins were born prior to the
completion of the genetic code (14). We predict that ancestrally reconstructed sequences from
this era will be enriched in early amino acids and depleted in late amino acids. Previous analyses
relied on conserved residues within a small number of LUCA proteins (15, 16). Here, we classify
a larger set of protein-coding domains that date back to LUCA, rather than being more recently
born, e.g., *de novo* from non-coding sequences or alternative reading frames (17, 18). We
compare reconstructed ancient amino acid frequencies of the most ancient vs. moderately
ancient protein cohorts, to deduce the order in which amino acids were incorporated into the
genetic code.

We take advantage of gene-tree species-tree reconciliation methods (19) to infer LUCA's protein
sequences. Previous analyses focused on the age of orthologous gene families (20-22); ours is
the first to infer which protein domains date back to LUCA. Protein domains are the basic units of
proteins, that can fold, function, and evolve independently (23). Proteins often contain multiple
protein domains, each of which might have a different age (Figure 1). For the purpose of inferring
ancient amino acid usage, what matters is the age of the protein domain, not that of the whole
protein that it is part of. We use protein domain annotations from the Pfam database (24). We
recognize Pfams present in LUCA by trimming horizontal gene transfer (HGT) events, and by
exploiting long archaeal-bacterial branches (Figure 2; see Methods for details).

Figure 1. The evolutionary history of a protein domain may date back further in time than
that of the whole-gene ortholog that it is part of. Multi-domain genes 3 and 4 originated

around the same time. However, they are made up of two protein domains (blue & orange boxes)
that emerged and diverged at different points in time – domain 1 is older than domain 2.

110
Figure 2. Criteria for (a) LUCA Pfam annotation, (b) identifying HGT to be filtered, and (c)
pre-LUCA Pfam annotation. Details are in Methods, with a brief summary here. a) Pruning HGT
between archaea and bacteria reveals a LUCA node as dividing bacteria and archaea at the root.
Colored circles are indicated just upstream of the most recent common ancestor (MRCA) of all
copies of that Pfam found within the same taxonomic supergroup. We recognize a total of five
bacterial supergroups (FCB, PVC, CPR, Terrabacteria and Proteobacteria (75, 76)) and four
archaeal supergroups (TACK, DPANN, Asgard and Euryarchaeota (77, 78)); only 4 out of 5
bacterial supergroups and 3 out of 4 archaeal supergroups are shown. The yellow diamond
indicates LUCA as a speciation event between archaea and bacteria. We do not assume that the
LUCA coalescence timing was the same for every Pfam (94). Prior to HGT pruning, PVC
sequences can be found on either side of the two lineages divided by the root. After pruning
intradomain HGT, four MRCAs are found one node away from the root, and three more MRCAs
are found two nodes away from the root, fulfilling our other LUCA criterion described in the
Methods, namely presence of at least three bacterial and at least two archaeal supergroup
MRCAs one to two nodes away from the root. b) Criteria for pruning likely HGT between archaea
and bacteria (see Methods for details). We partition into monophyletic groups of sequences in the
same supergroup; in this example, there are four such groups, representing two bacterial
supergroups and one archaeal supergroup. There is one 'mixed' node, separating an archaeal
group (HG1) from a bacterial group (HG2). It is also annotated by GeneRax (19) as a transfer 'T'.
The bacterial nature of groups 3 and 4 indicates a putative HGT direction from group 2 to group
1. Group 2 does not contain any Euryarchaeota sequences, meeting the third and final
requirement for pruning of group 1. If neither Proteobacteria or Euryarchaeota sequences were
present among the other descendants of the parent node, both groups 1 and 2 would be
considered acceptors of a transferred Pfam and would both be pruned from the tree. c) Pre-LUCA
Pfams have at least two nodes annotated as LUCA.

136
137　**Results**

Ancient protein domain classifications agree with whole-gene classifications

We classify 969 Pfams and 445 clans (sets of one or more Pfams that are evolutionary related) as present in LUCA (Figures 3a and 3b; detailed lists in Supplementary Tables 1 and 2). We compare these to the 3055 Pfams and 1232 clans that we classify as ancient but post-LUCA (including Last Bacterial Common Ancestor (LBCA) and Last Archaeal Common Ancestor (LACA) candidates). Encouragingly, 88.6% of Pfams that we annotate as pre-LUCA or LUCA are contained within genes annotated by Moody et al. (21) as present in LUCA with more than 50% confidence, when present in their dataset (Figure 3c). This level of agreement far exceeds earlier works (22).

Figure 3. Pfams (a) and clans (b) classified as ancient are well validated by the whole gene annotations of Moody et al. (21) (c). a) Ancient post-LUCA Pfam classifications include 285 LACA candidates and 2770 LBCA candidates (more analysis would be required to rule out extensive HGT within archaea or bacteria). Modern Pfams are distributed among the prokaryotic supergroups as follows: 9 CPR, 210 FCB, 942 Proteobacteria, 51 PVC, 1111 Terrabacteria, 2 Asgard, 49 TACK, and 177 Euryarchaeota. In addition to supergroup-specific modern Pfams, we classified another 1097 Pfams, present in exactly two bacterial supergroups, as modern post-LBCA. We deemed 15 Pfams unclassifiable due to high inferred HGT rates, 397 due to uncertainty in rooting, and 198 due to ancient rooting combined with absence from too many supergroups (see Methods). b) Pre-LUCA clans contain at least two LUCA-classified Pfams or one pre-LUCA Pfam, whereas LUCA clans contain exactly one LUCA Pfam. Ancient post-LUCA clans contain no LUCA, pre-LUCA, or unclassified Pfams; they include an ancient post-LUCA Pfam or at least two modern Pfams covering at least two supergroups from only one of either bacteria or archaea. Modern clans include Pfams whose root is assigned at the origin of one

supergroup. Finally, unclassifiable clans did not meet any of our clan classification criteria, e.g.,
because they included both post-LUCA and unclassifiable Pfams. c) 98% of our pre-LUCA Pfams
and 87% of our LUCA Pfams are present in genes annotated by as present in LUCA with more
than 50% confidence, when present in their dataset. We mapped all Clusters of Orthologous
Genes (COGs) (95) in the Moody et al. (21) supplementary dataset (STable_1.csv) to UniProt IDs
(96) using the EggNOG 5.0 database (97). We then identified their associated Pfams using the
'Pfam-A.regions.uniprot.tsv' file downloaded from the Pfam FTP site (https://pfam-
docs.readthedocs.io/en/latest/ftp-site.html#current-release) (24) on May 28[th], 2024. Our protein to
Pfam ID mappings are available in 'Protein2Domain_mappings' at
https://doi.org/10.6084/m9.figshare.27191274.v1.

In agreement with the Moody et al. (21) classification of LUCA metabolism, almost all Pfams
associated with enzymes in hydrogen metabolism, assimilatory nitrate and sulfate reduction
pathways, and the Wood-Ljungdahl pathway date back to LUCA (Supplementary Table 3). Our
results also support a, post-LUCA, bacterial origin of nitrogen fixation (21, 25) (Supplementary
Table 3). We assign to LUCA the complete set of amino acid-tRNA synthetase-associated anti-
codon binding domains found in modern prokaryotes. Here, focusing on complete genes would
have been problematic, because accessory amino acid-tRNA synthetase-associated domains
(e.g. PF04073 and PF13603, which deacylate misacylated tRNA) were sometimes added later.

We also checked the antiquity of the cofactor/cosubstrate S-adenosylmethionine (SAM) (26), both
with respect to SAM biosynthesis and SAM usage. In agreement with past work attributing the
SAM biosynthesis enzyme methionine adenosyltransferase to LUCA (27, 28), we assign its single
Pfam (PF01941) to LUCA (the corresponding COG1812 is not analyzed by Moody et al. (21)). In
agreement with past work attributing SAM-dependent methyltransferases to LUCA (29), Moody et
al. (21) assign the RsmB/RsmF family (COG0144), which methylates 16S rRNA, more than 75%
confidence of being present in LUCA, and we also classify its SAM-binding Rossman fold Pfam
(PF01189) as LUCA. In agreement with (30, 31), Moody et al. (21) assign the SAM-binding tRNA
methylthiolase (COG0621) to LUCA with more than 75% confidence, and we confirm the pre-
LUCA status of its associated Radical SAM, TIM-barrel-related Pfam (PF04055). In agreement
with attribution of polyamines to LUCA (32) we assign to LUCA the one Pfam (PF02675) of S-
adenosylmethionine decarboxylase, which acts on SAM in the first step of polyamine synthesis;
the antiquity of corresponding COG1586 is not further confirmed by Moody et al. (21).

Hydrophobic amino acids are more interspersed within ancient proteins

Interspersion of hydrophobic amino acids away from one another along the primary sequence is
believed to mitigate risks from protein misfolding, while still enabling correct folding (33-35). Older
sequences have previously been found to have greater interspersion among their hydrophobic
residues, indicating more sophisticated protein folding (14, 36), likely due to survivorship bias
(37). Our Pfam age classifications confirm the antiquity of this trend, previously observed only for
animal sequences. LUCA Pfams show even more hydrophobic interspersion than the still-ancient
'post-LUCA' Pfams that include LACA candidates and LBCA candidates (Supplementary Figure
1; Wilcoxon rank sum test; $p = 0.02$). Post-LUCA Pfams in turn have more hydrophobic
interspersion than 'modern' Pfams that are specific to particular prokaryotic supergroups
(Wilcoxon rank sum test; $p = 0.02$).

LUCA's protein sequences were depleted in larger amino acids

Clans present in LUCA were born before the divergence of Archaea and Bacteria, some
potentially prior to the completion of the genetic code. If newly recruited amino acids were added
slowly, the contemporary descendants of LUCA clans will show signs of ancestral depletion in
amino acids that were added late to the genetic code. We first focus on clans present in one copy

in LUCA (denoted "LUCA clans"), excluding those that had already duplicated and diverged into
multiple surviving lineages (denoted "pre-LUCA clans"). We score ancestral amino acid
enrichment and depletion as relative to still-ancient post-LUCA clans, which represent amino acid
usage from the standard genetic code of all 20 amino acids, plus any ascertainment biases. This
ratio, reflecting ancient amino acid usage, is not confounded with the effects of temperature, pH,
oxygen tolerance, salinity, GC content, or transmembrane status on amino acid frequencies
(Supplementary Figures 2a-f). Indeed, LUCA usage is similar in the very different biophysical
context of a transmembrane site (Supplementary Figure 3).

Smaller amino acids are enriched in LUCA (Figure 4a; weighted R^2 = 0.48, p = 0.0005). Results
are similar using a restricted set of Pfams validated by Moody et al. (21) (weighted R^2 = 0.44, p =
0.001). As a negative control for methodological artifacts, the ancestral amino acid usage of post-
LUCA clans relative to modern clans is not correlated with molecular weight (p = 0.9).

Figure 4. LUCA is enriched for smaller amino acids, with subtle differences between single
copy LUCA vs. multi-copy pre-LUCA sequences. Ancestrally reconstructed amino acid
frequencies in LUCA and pre-LUCA clans are shown relative to those in ancient post-LUCA
clans. a) LUCA clans and b) pre-LUCA clans are enriched for amino acids of smaller molecular
weight. Weighted model 1 regression lines are shown in black with 95% confidence interval grey

shading. Error bars indicate standard errors. c) Character colors show the assignments of
Moosmann (5); colored circles indicate our re-assignments. We reclassify phenylalanine (F)
because it is enriched in proteins in mesophiles compared to their orthologs in thermophiles and
hyperthermophiles (98). We reclassify aspartic acid (D) because the surfaces of proteins within
halophilic bacteria are highly enriched in aspartic acid compared to in the surfaces of non-
halophilic mesophilic and thermophilic bacteria, in a manner that cannot be accounted for by the
dinucleotide composition of the halophilic genomes (99). The brown circle around methionine
highlights that while it might not be utilized against reactive oxygen species, it might once have
been against ancient reactive sulfur species. d) Model 2 Deming regression (accounting for
standard errors in both variables, implemented in deming() version 1.4-1 (100)) in blue shows that
pre-LUCA enrichments are not more extreme versions of LUCA enrichments, lying on the wrong
side of the y=x red line. We include the imidazole-ring-containing H as aromatic. Asterisks (*)
indicate statistically different amino acid frequencies between pre-LUCA and LUCA (Welch two
sample t-test, $p<0.05$ and $p<0.01$).

Revised Order of Amino Acid Recruitment

Figure 4c visualizes how LUCA's amino acid enrichments compare to Trifonov's consensus order
(4). While they are correlated (weighted R^2 = 0.37, p = 0.003), this association disappears in a
weighted multiple regression with both molecular weight (p = 0.03) and Trifonov's (4) order (p =
0.9) as predictors (weighted R^2 = 0.48). This is also true using Trifonov's revised 2004 order
based on 60 metrics (38) (weighted R^2 = 0.34, p = 0.006 on its own; p = 0.9 when molecular
weight is also a predictor of LUCA usage). This suggests that some of Trifonov's 40-60 metrics
made his estimates of the order of recruitment worse rather than better. We use enrichment in
LUCA to re-classify VGIMTAHEPC as 'early' and depletion to classify KSDLNRFYQW as 'late'.
More precise estimation of the order of recruitment, with standard errors, is given in Table 1.

We place glutamine (Q or Gln) as the second last amino acid, much later than Trifonov (4)
inferred. Consistent with its late addition, Gln-tRNA synthetase (GlnRS) is either absent in
prokaryotes, or acquired via horizontal gene transfer from eukaryotes (39). Prokaryotes that lack
GlnRS perform tRNA-dependent amidation of Glu mischarged to Gln-tRNA by GluRS, forming
Gln-acylated Gln-tRNA via amidotransferase. The core catalytic domain (PF00587), shared
between the GlnRS and GluRS paralogs, is present in LUCA and can indiscriminately acylate
both Gln-tRNA and Glu-tRNAs with Glu (40).

Metal-binding and sulfur-containing amino acids were added early to the genetic code

Methionine (M), cysteine (C), and histidine (H) are all enriched in LUCA, despite previous
annotation as late additions to the genetic code (Figure 4c). C and H are the most frequently used
amino acids for binding iron, zinc, copper, and molybdenum, and H, aspartic acid (D) and
glutamic acid (E or Glu) for binding manganese and cobalt (Figure 2D of (41)). Binding can either
be to a metal ion, or to iron-sulfur (FeS) clusters, usually via C but sometimes via H or D (42).
Binding these transition metals is key to catalysis (43). Figure 4a is incompatible with C, H, D, or
E being late additions, and indeed H is more enriched than one would expect from its molecular
weight.

C and M are the only sulfur-containing amino acids in the contemporary genetic code.
Contemporary prokaryotes living in H_2S-rich environments use more C and M than matched
species (Supplementary Figure 4); LUCA's C and M enrichment might thus reflect an
environment rich in H_2S.

Moosmann (5) classified M, tryptophan (W), and tyrosine (Y) as antioxidants, because he
believed them to protect the overall protein structure from oxidative stress via sacrificial
oxidization. For instance, surface M residues can be reversibly oxidized to form methionine

sulfoxide (44). This might have driven isoleucine recoding to methionine in mitochondria (45, 46).
However, proteins in aerobes are enriched in W and Y but not in M (47). Our results also
separate early M from late Y and W (Figure 4). We speculate that methionine, abundant due to
early life's use of SAM, might have protected against reactive sulfur species such as sulfide (S^{2-}),
which were present in early, H_2S-rich environments (48). Our results are then partially compatible
with Granold et al.'s (49) view that Y and W (but not M) were added to complete the modern
genetic code after reactive oxygen species became the main oxidizing threat.

Amino acid	LUCA usage	LUCA usage standard error	Pre-LUCA usage	Pre-LUCA usage standard error
V	1.12	0.0241	1.04	0.0205
G	1.11	0.0283	1.09	0.0241
I	1.1	0.0325	1.07	0.0351
M	1.08	0.0386	1.1	0.0383
A	1.07	0.0317	1.03	0.0297
T	1.07	0.0369	1.05	0.0362
H	1.04	0.0416	1.17	0.0486
E	1.03	0.0357	0.911	0.0357
C	1.01	0.0722	1.03	0.0844
P	1.01	0.0282	1.04	0.0255
K	0.974	0.038	0.901	0.0334
S	0.972	0.0265	1.02	0.0239
D	0.968	0.027	0.988	0.0363
L	0.942	0.0256	0.962	0.032
N	0.934	0.0374	0.996	0.0432
R	0.916	0.0265	0.915	0.0271
F	0.895	0.032	1.02	0.0394
Y	0.858	0.0341	0.982	0.0309
Q	0.827	0.031	0.847	0.0304
W	0.649	0.0476	0.865	0.0526

Table 1. LUCA and pre-LUCA clans' ancestral amino acid frequencies are divided by post-LUCA
clan's ancestral amino acid frequencies to produce measures of relative usage. The standard
errors of the amino acid usages were calculated using an approximation derived from a Taylor
expansion of the ratio (90). For each of the 20 ancestral amino acid frequencies, the standard
errors of the weighted means across all the clans within the LUCA and pre-LUCA phylostrata
(weighted by the maximum number of ancestral sites across all Pfams in a given clan) were
calculated using the weighted_se() function in the diagis R package (89)(See Methods for more
detail).

Pre-LUCA clans hint at more ancient genetic codes

We expected pre-LUCA enrichments and depletions to be more extreme than for LUCA, but only
H fits this prediction (Figure 4d), with significantly higher frequencies in pre-LUCA than in LUCA.

There is nevertheless a strong overall correlation between LUCA and pre-LUCA usages (R^2 =
0.51, p = 0.0003). Pre-LUCA, like LUCA, is strongly depleted in Q, supporting the inference that
Q, not Y, was the 19th amino acid recruited into the standard genetic code. Pre-LUCA usage does
not correlate with Trifonov's consensus order (4) (p = 0.2), and correlates more weakly with
molecular weight (Figure 4b) (weighted R^2 = 0.33, p = 0.007).

H is one of six amino acids with significantly different frequencies in pre-LUCA vs. LUCA. All
three of the canonical, benzene-ring bearing, aromatic amino acids (W, Y, and phenylalanine (F)),
as well as the imidazole-ring containing H, are more common in pre-LUCA than in LUCA (Figure
4d, Welch 2-sample t-test; p = 0.03, 0.001, 0.03 and 0.01, respectively; 2.4% vs 2.1% H, 1.2% vs
0.9% W, 3.1% vs. 2.8% Y, and 4.1% vs. 3.7% F). Glutamic acid (E) and Valine (V) are less
common in pre-LUCA than in LUCA (Welch 2-sample t-test; p = 0.01 and 0.004, respectively;
7.3% vs. 8.2% E, 7.5% vs. 8.1% V).

More W in pre-LUCA than LUCA is particularly surprising, because there is scientific consensus
that W was the last of the 20 canonical amino acids to be added to the genetic code. Therefore,
we manually inspected the pre-LUCA Pfam with the highest tryptophan frequency (3.1%):
PF00133, the core catalytic domain of the tRNA synthetases of leucine (L), isoleucine (I), and
valine (V). Each of these three synthetases has well-separated archaeal and bacterial branches,
confirming its pre-LUCA dating (Supplementary Figure 5). Highly conserved tryptophan sites
regulate the size of the amino acid binding pocket, allowing the synthetases to discriminate
among I, L, and V (50). There are also conserved I and V sites in the common ancestor of the I
and V tRNA synthetases, indicating that discrimination between the two happened prior to the
evolution of the synthetases currently responsible for the discrimination (51). This suggests that
an alternative, more ancient system predated the modern genetic code, and in particular predated
the evolution of super-specific, cognate aaRSs (51).

319
Discussion

The evolution of the current genetic code proceeded via stepwise incorporation of amino acids,
driven in part by changes in early life's environment and requirements. Contemporary proteins
retain information about which amino acids were part of the code at the time of their birth,
allowing us to infer the order of recruitment on the basis of enrichment or depletion in LUCA's
protein domains. Smaller amino acids were added to the code first, and when this is accounted
for, there is no further information in Trifonov's (4) widely used 'consensus' order based on 40
metrics, some of dubious relevance. The sulfur-containing amino acids C and M were
incorporated earlier than previously thought, likely because those metrics included experiments
conducted in the absence of sulfur. Q was added later than previously thought, in agreement with
evidence from glutamyl-tRNA synthetases. M and H were added to the code earlier than
expected from their molecular weights, and Q later. Even more ancient amino acid usage, in
sequences that had already duplicated and diverged pre-LUCA, shows significantly higher
frequencies of the aromatic amino acids W, Y, F, and H, and significantly lower frequencies of E
and V.

If LUCA lived in a H_2S-rich environment (48, 52), M residues could have protected proteins
against sulfur-mediated oxidative stress. M would furthermore have had high biotic availability as
the precursor (53) and product (54) of SAM, given our finding that LUCA made and used SAM.
The potentially sulfur-rich nature of early terrestrial life is context for astrobiology investigations of
sulfur-rich environments on Mars and Europa, with associated biosignatures key to life detection
(55).

An early role for H is compatible with a key role for metal binding in early life. It also resolves the
previous puzzle that the ancestral, conserved region of all Class I aaRSs contains a histidine-rich
HIGH motif (56, 57). The lack of abiotic availability was key to H's previous annotation as late, but
biotic availability of H in an RNA-dominant biotic context would have been sufficient. The

importance of abiotic availability (58, 59) to the origins of the genetic code remains unclear. We
note that ongoing research on plausible prebiotic syntheses in cyanosulfidic environments (60)
and alkaline hydrothermal vents (61) is reshaping our understanding of which amino acids were
accessible to early life. Amino acid abundances obtained from asteroid sample returns will also
soon contribute (62).

Our results offer an improved approximation of the order of recruitment of the twenty amino acids
into the genetic code under which contemporary protein-coding sequences were born. This order
need not match the importance or abundance with which amino acids were used by still earlier
life forms, nor during the prebiotic to biotic transition. Instead of using Trifonov's assignments (4)
to capture the order in which amino acids were recruited into our genetic code, we recommend
using the LUCA amino acid enrichment values plotted on the y-axis of Figure 4a, which can be
found together with their standard errors in Table 1.

More broadly, coding for different amino acids might have emerged at similar times but in
different biogeochemical environments. The temporal order of recruitment that we infer based on
LUCA sequences is not the temporal order for coding as a whole, but for the ancestor of the
modern translation machinery. Indeed, horizontal gene transfer of the tRNAs coupled with their
cognate aminoacyl tRNA synthetases might have brought the diverse components of the modern
translation machinery together (63). This further emphasizes that the time of origin of the
translation machinery's components need not match the time of their incorporation into the
surviving ancestral lineage.

The construction of the genetic code was tethered to the evolution of the ribosome (64). If the
ribosome's exit tunnel, whose formation and subsequent extension was key to ribosome evolution
(65), limited the size of the amino acids passing through, its progressive dilation could explain the
strong relationship between amino acid size and order of recruitment evidenced in LUCA clans. If
older, alternative codes were not similarly limited, this would explain why amino acid size is a
weaker predictor of pre-LUCA's amino acid usage compared to LUCA's amino acid usage.

To explain the different enrichments of pre-LUCA versus LUCA sequences, as well as the
surprising conservation of some sites prior to the emergence of the aaRSs that distinguish the
relevant amino acids, we propose that some pre-LUCA sequences are older than the current
genetic code, perhaps even tracing back to a peptide world at the dawn of precellular life (7).
Stepwise construction of the current code and competition among ancient codes could have
occurred simultaneously (66, 67). Ancient codes might also have used non-canonical amino
acids, such as norvaline and norleucine (68) which can be recognized by LeuRS (69, 70). Along
with having different genetic codes, we speculate that pre-LUCA and LUCA might have existed in
different geochemical settings. For instance, if pre-LUCA ancestors inhabited alkaline
hydrothermal vents, where abiotically produced aromatic amino acids have been found (61), this
would explain their enrichment in pre-LUCA relative to LUCA. We note that abiotic synthesis of
aromatic amino acids might be possible in the water-rock interface of Enceladus's subsurface
ocean, which is speculated to be analogous to terrestrial alkaline hydrothermal vents (71).

Perhaps the biggest mystery is how sequences such as the common ancestor of L/I/V-tRNA
synthetase, which were translated via alternative or incomplete genetic codes, ended up being re-
coded for translation by the direct ancestor of the canonical genetic code. Harmonization of
genetic codes facilitated innovation sharing via HGT, making it advantageous to use the most
common code, driving code convergence (72, 73). Only once a common code was established
did HGT drop to levels such that a species tree became apparent, i.e. the LUCA coalescence
point corresponds to convergence on a code (72). Our identification of pre-LUCA sequences
provides a rare source of data about early, alternative codes.

Materials and Method

Pfam sequences

We downloaded genomes of 3562 prokaryotic species from NCBI that were present in the Web of
Life (WoL): Reference phylogeny of microbes (74) in August 2022. We classified them into five
bacterial supergroups (FCB, PVC, CPR, Terrabacteria and Proteobacteria (75, 76)) and four
archaeal supergroups (TACK, DPANN, Asgard and Euryarchaeota (77, 78)). We included
incomplete genomes, to enhance coverage of underrepresented supergroups.

We assign ages not to whole proteins but to each of their protein domain constituents. We used
InterProScan (79) to identify instances of each Pfam domain (24). We excluded Pfams with fewer
than 50 instances across all downloaded genomes. We also excluded 9 Pfams marked "obsolete"
starting July 2023. Among the remaining 8282 Pfams, 2496 Pfams had more than 1000
instances. We downsampled these to balance representation across the two taxonomic domains
(archaea and bacteria). For instance, a Pfam with 2000 bacterial and 500 archaeal instances was
downsampled by retaining all 500 archaeal sequences plus a subset (randomly sampled without
replacement) of 500 bacterial sequences.

The Pfam database includes annotations of "clans" of Pfams that share a common ancestor
despite limited sequence similarity; for many analyses, we used clans rather than Pfams to
ensure independent datapoints. We treated Pfams that were not annotated as part of a clan as
single-entry clans, with clan ID equal to their Pfam ID.

Pfam trees

We aligned downsampled sequences for each Pfam using MAFFT v.7 (80), to infer a preliminary
tree with IQ-Tree (81), using a time non-reversible amino acid substitution matrix trained on the
Pfam database (NQ.PFAM) (82), and no rate heterogeneity among sites. Because most Pfams
are too short for reliable tree inference, we next reconciled preliminary Pfam trees with the WoL
prokaryotic species tree (74) using GeneRax (19). While there is no perfect species tree for
prokaryotes, reconciliation even with a roughly approximate tree can still provide benefits. We ran
GeneRax twice. The first run used the LG amino acid substitution model, a gamma distribution
with four discrete rate categories, and a Subtree Prune and Regraft (SPR) radius of 3. The
second run used the output of reconciled trees from the first run as input, and switched to an SPR
radius of 5, and the Q.PFAM amino acid substitution model (83), which was trained on the Pfam
dataset. We did not use NQ.PFAM, because time non-reversible models are only implemented in
IQ-Tree (82), and not in GeneRax. In both runs, we used the UndatedDTL probabilistic model to
compute the reconciliation likelihood. The second run of GeneRax reduced estimated transfer
rates by an additional 7% (Welch two sample t-test, $p = 10^{-12}$), indicating continued improvements
to the phylogenies.

We re-estimated the branch lengths of the reconciled Pfam trees in IQ-Tree using the NQ.PFAM
substitution model with no rate heterogeneity, then performed midpoint rooting using the phytools
R package (84) on these re-estimated branch lengths. As alternative rooting methods, we also
explored and rejected minimum variance (85), minimal ancestral deviation (86), and rootstraps
based on time non-reversible substitution models (87). The first two methods work best when
deviations from the molecular clock average out on longer timescales, which is not true for
phylogenies in which evolution e.g. at different temperatures causes sustained differences in
evolutionary rate. Indeed, minimum variance failed to resolve the prokaryotic supergroups as
separate clades, in visual inspection of PF00001, due to presumed genuine rate variation among
taxa. The latter produced very low confidence roots. In contrast, midpoint rooting largely
conformed to expectations for aaRSs once we implemented the procedure for outlier removal
described under "Classifying Pfam domains into ancient phylostrata" below.

We then implemented a new --enforce-gene-tree-root option in GeneRax, and ran GeneRax in
evaluation mode, with Q.PFAM+G as the substitution and rate heterogeneity models,
respectively. Evaluation mode re-estimates the reconciliation likelihood and the duplication,
transfer and loss (DTL) rates on a fixed tree, without initiating a tree search. Fifteen reconciled

Pfam trees had inferred transfer rates higher than 0.6, three times the seed transfer rate
implemented by GeneRax. We took this as a sign of poor tree quality, and annotated these 15
Pfams as of unclassifiable age.

Filtering out HGT between archaea and bacteria

Exclusion of horizontal gene transfer (HGT) between bacteria and archaea facilitates the
classification of a Pfam into LUCA (Figure 2a). To achieve this, we divided sequences into
"homogeneous groups", meaning the largest monophyletic group in the Pfam tree for which the
corresponding species all belong to the same prokaryotic supergroup. Each homogeneous group
was considered as a candidate for exclusion, via its "focal node" separating it from its sister
group. To avoid over-pruning, we do not consider deep focal nodes that are 2 or fewer nodes
away from the root.

To be excluded, we first require the focal node to be 'mixed', meaning its descendants are found
within both Bacteria and Archaea. We next require the focal node to be labelled by GeneRax as
most likely a transfer (T), rather than a duplication (D) or speciation (S). Finally, to identify
homogeneous groups likely to be receivers rather than the donors of transferred sequences, we
require the sister lineage to contain no sequences present in the same supergroup as that
defining the homogeneous group in question. An example of filtering is shown in Figure 2b.

We ran the filtering process twice to address rare occasions of an intradomain HGT nested within
another intradomain HGT group. In the second filter, we apply the third criterion after pruning the
homogenous groups identified as HGT during the first filter.

Classifying Pfam domains into ancient phylostrata

We re-rooted the HGT-pruned Pfam trees using the midpoint.root function in the 'phytools' R
package (84), before classifying them into phylostrata (i.e. cohort of sequences of similar age).
Classification was based on the locations of the most recent common ancestors (MRCAs) of each
supergroup. For a LUCA Pfam, we require the root to separate the MRCAs of all bacterial
supergroups from the MRCAs of all archaeal supergroups (Figure 2a).

If there were no horizontal transfer, and the tree of a Pfam present in one copy in LUCA were
error-free, then the MRCAs for the nine supergroups would be two to four branches away from
the root. This is true even if our Pfam tree and/or species tree do not correctly capture the true
phylogenetic relationships among supergroups. However, we cannot ignore HGT; we did not filter
out the products of HGT between supergroups within Archaea or within Bacteria, only that of HGT
between Archaea and Bacteria. HGT from a more derived supergroup to a more basal
supergroup will move the inferred MRCA of the former further back in time. Given rampant HGT,
whether real or erroneously implied by Pfam tree error, we required Pfams to have their
supergroups' MRCA two branches away from the root (Figure 2a).

Phylogenies with three or more basal bacterial supergroups and two or more basal archaeal
supergroups were classified as LUCA. In other words, we allow the absence of up to two
supergroups per taxonomic domain, as compatible with ancestral presence followed by
subsequent loss. Trees with three or more basal bacterial supergroups but fewer than two basal
archaeal supergroups, as well as trees with two or more basal archaeal supergroups but fewer
than three basal bacterial supergroups, were classified as ancient but post-LUCA. These are
candidate Pfams for the Last Bacterial Common Ancestor (LBCA) and the Last Archaeal
Common Ancestor (LACA) phylostrata, respectively, but the necessary HGT filtering for sufficient
confidence in this classification is beyond the scope of the current work. If only one basal
supergroup is present, then the Pfam is classified into the corresponding supergroup-specific
phylostratum, meaning it emerged relatively recently (modern post-LUCA). If two basal bacterial
supergroups (and no archaeal supergroups) were present, the Pfam was classified as post-LBCA

which was also considered modern post-LUCA (younger than LBCA but older than the
supergroup-specific phylostrata). The remaining Pfams were considered unclassifiable.

We also classify into a pre-LUCA phylostratum the subset of LUCA-classified Pfams for which
there is evidence that LUCA contained at least two copies that left distinct descendants. This is
motivated by the assumption that LUCA domains that were born earlier are more likely to have
duplicated and diverged prior to the archaeal-bacterial split (88). We require that both the nodes
that are only one branch from the root be classified as LUCA nodes. This means that each of
these nodes should, after HGT filtering: i) split a pure-bacterial lineage from a pure-archaeal
lineage, and ii) include as descendants at least three bacterial and two archaeal basal MRCAs no
more than two nodes downstream of the potential LUCA nodes (Figure 2c).

Assignment of a Pfam to a phylostratum is sensitive to the root's position. Midpoint rooting is
based on the longest distance between two extant sequences. A single inaccurately placed
sequence can yield an abnormally long terminal branch, upon which the root is then based. This
phenomenon was readily apparent upon manual inspection of rooted Pfam trees. To ensure the
robustness of age classifications to the occasional misplaced sequence, we removed the Pfam
instance with the longest root-to-tip branch length in each HGT-filtered tree as potentially faulty,
re-calculated the midpoint root, and then re-classified each Pfam. We repeated this for ten
iterations, then retained only those Pfams that were classified into the same phylostratum at least
7 out of 10 times. Our HGT filtering algorithm does not act on nodes near the root, making it
robust to small differences in root position; we therefore did not repeat the HGT-filtering during
these iterations.

We classified clans that contained at least two LUCA Pfams as pre-LUCA clans. Clans that
contained both ancient archaeal and ancient bacterial post-LUCA Pfams (i.e. candidate LACA
and LBCA Pfams) were classified as LUCA. Clans that contained at least two different archaeal
but no bacterial supergroup-specific Pfams, or three different bacterial supergroup-specific Pfams
but no archaeal supergroup-specific Pfams, were classified as ancient post-LUCA clans. Clans
that meet neither of these criteria, and that contain at least one unclassified Pfam, were
considered unclassifiable due to the possibility that the unclassified Pfam might be older than the
classified Pfams present in the clan. All other clans were assigned the age of their oldest Pfam.

For a more stringent analysis of amino acid usage, we restrict our Pfam dataset to those present
in proteins annotated by Moody et al. (21) as >75% likely to be in LUCA. We then re-classified
clan ages. Data on the likelihood of Pfams being present in LUCA, as annotated by Moody et al.
(21), can be found in 'MoodyPfams_probabilities.csv' on GitHub.

Ancestral amino acid usages

Ancestral sequence reconstruction (ASR) can introduce a variety of biases. ASR methods do not
resolve alignment gaps well, to infer indel evolution, instead inferring ancestral sequences far
longer than any contemporary descendant. To avoid bias among amino acids regarding which
contemporary sequences appear in the ancestral sequence more often than they should, we
retain only sites where more than 50% of the sequences contain an amino acid (i.e. no indel).
This ensures that no amino acid can be double counted.

For Pfams classified as pre-LUCA or LUCA, we require that a given site contain an amino acid
and not a gap in at least five bacterial sequences and five archaeal sequences. This additional
filter helps ensure that the ancestrally reconstructed sites were not inserted post-LUCA (even
when the Pfam itself dates back to LUCA). It also reduces the impact of any Pfams misclassified
as ancient on the inferred ancient amino acid usage.

Following these filters, we ran the remaining sites in each Pfam alignment (prior to HGT filtering)
through IQ-Tree with the -asr option, the NQ.PFAM substitution model, and R10 rate
heterogeneity. We then excluded low confidence sites from subsequent analyses, based on the

most likely amino acid having an ancestral probability estimate <0.4. Combined with the other two
filters described above, the concatenated sequence length for all four phylostrata (pre-LUCA,
LUCA, post-LUCA, and modern) fell by ~11%, presumably preferentially excluding rapidly
evolving sites to a similar degree in all four cases, such that amino acid exclusion biases cancel
out when ratios are taken.

We then summed over the amino acid probability distributions at each site at the deepest node,
and divided by the number of sites, to obtain per-Pfam estimated ancestral amino acid
frequencies. For each clan, we took the ancestral amino acid frequencies across Pfams,
weighted by the number of ancestral sites in the Pfams. For each phylostratum, we averaged
across clans, weighted by the maximum number of ancestral sites across all Pfams in a given
clan. We calculated a standard error associated with each phylostratum mean using the
weighted_se() function in the diagis R package (89).

We divided ancestral amino acid frequencies for the LUCA and pre-LUCA phylostrata by post-
LUCA ancestral amino acid frequencies to produce measures of relative usage. Standard errors
of each of these ratios L/P were calculated using an approximation derived from a Taylor

expansion of the ratio: $\sqrt{\frac{\sigma_L{}^2}{P^2} + \frac{L^2 \sigma_P{}^2}{P^4}}$ (90). These were used in weighted linear model 1
regressions, using the lm() function with the 'weights' argument in the 'stats' package in base R
(91). Uncertainty in the ancestral states arising over 4 billion years of evolution is expected to
bring values of L/P closer to one, without entirely erasing the signal. As a negative control for
bias, we calculate the relative amino acid usage of post-LUCA clans by dividing the ancestral
amino acid frequencies for post-LUCA clans by the ancestral amino acid frequencies for modern
clans.

Standard errors in Trifonov's (4) average rank reflect but underestimate uncertainty; we therefore
treat Trifonov's (4) rankings as the dependent variable and use its weights rather than errors on
L/P to weight the regression model in Figure 4c. Standard errors are not available for alternative
results based on Trifonov's 2004 order (38).

Hydrophobic interspersion

The degree to which hydrophobic are clustered vs. interspersed along the primary sequence was
calculated as a normalized index of dispersion for each Pfam instance (35). This metric uses the
ratio of the variance to the mean in the number of the most hydrophobic amino acids (leucine,
isoleucine, valine, phenylalanine, methionine, and tryptophan) within consecutive blocks of six
amino acids. The values of this index of dispersion were then normalized, to make them
comparable across Pfams with different lengths and hydrophobicities. In cases where the Pfam
length was not a multiple of 6, the average across all possible 6-amino acid frames was
computed, trimming the ends as needed. For additional details, see Foy et al. (36) or James et al.
(14). For each Pfam, we then took the average across all its instances (prior to downsampling
species).

Transmembrane annotation

We identified transmembrane sites within each Pfam using DeepTMHMM (92) on a consensus
sequence generated from the original multiple sequence alignments (prior to HGT filtering) using
the majority-rule seq_consensus() function in the R package 'bioseq' (93).

Data and Code Availability

Data files and R scripts used to generate the results and figures are available at
[sawsanwehbi/Pfam-age-classification GitHub repository](). Pfam sequences, alignments, trees and
mappings to protein IDs are available at https://figshare.com/projects/Pfam-age-classification-
data/201630.

Acknowledgments

We thank NASA [80NSSC24K0384] and the John Templeton Foundation [62220] for funding SW,
NM, and JM, Chan-Zuckerberg Initiative [EOSS4-0000000312] for funding BQM, the DFG [STA
860/6-2] for funding BM, the NSF [DEB-2333243] for funding AW, NM, and JM, the Arizona
NASA Space Grant Consortium [80NSSC20M0041] for funding NM, and the NIH [T32GM132008]
for funding AW. We thank Mike Barker, Alan Moses, and Elisa Tomat for helpful discussions, and
Cat Wolner for comments on the manuscript.

References

1. F. H. C. Crick, The origin of the genetic code. *Journal of Molecular Biology* **38**, 367-379 (1968).
2. J. T.-F. Wong, A Co-Evolution Theory of the Genetic Code. *Proceedings of the National Academy of Sciences* **72**, 1909-1912 (1975).
3. S. L. Miller, A Production of Amino Acids Under Possible Primitive Earth Conditions. *Science* **117**, 528-529 (1953).
4. E. N. Trifonov, Consensus temporal order of amino acids and evolution of the triplet code. *Gene* **261**, 139-151 (2000).
5. B. Moosmann, Redox Biochemistry of the Genetic Code. *Trends in Biochemical Sciences* **46**, 83-86 (2021).
6. W. Nitschke, S. E. McGlynn, E. J. Milner-White, M. J. Russell, On the antiquity of metalloenzymes and their substrates in bioenergetics. *Biochimica et Biophysica Acta (BBA) - Bioenergetics* **1827**, 871-881 (2013).
7. S. D. Fried, K. Fujishima, M. Makarov, I. Cherepashuk, K. Hlouchova, Peptides before and during the nucleotide world: an origins story emphasizing cooperation between proteins and nucleic acids. *Journal of The Royal Society Interface* **19** (2022).
8. E. T. Parker *et al.*, Primordial synthesis of amines and amino acids in a 1958 Miller H2S-rich spark discharge experiment. *Proceedings of the National Academy of Sciences* **108**, 5526-5531 (2011).
9. C. S. Foden *et al.*, Prebiotic synthesis of cysteine peptides that catalyze peptide ligation in neutral water. *Science* **370**, 865-869 (2020).
10. C. Shen, L. Yang, S. L. Miller, J. Oró, Prebiotic synthesis of histidine. *Journal of Molecular Evolution* **31**, 167-174 (1990).
11. A. Vázquez-Salazar, A. Becerra, A. Lazcano, Evolutionary convergence in the biosyntheses of the imidazole moieties of histidine and purines. *PLOS ONE* **13**, e0196349 (2018).
12. A. D. Goldman, B. Kacar, Cofactors are Remnants of Life's Origin and Early Evolution. *Journal of Molecular Evolution* **89**, 127-133 (2021).
13. A. J. M. Ribeiro, J. D. Tyzack, N. Borkakoti, G. L. Holliday, J. M. Thornton, A global analysis of function and conservation of catalytic residues in enzymes. *Journal of Biological Chemistry* **295**, 314-324 (2020).
14. J. E. James *et al.*, Universal and taxon-specific trends in protein sequences as a function of age. *Elife* **10** (2021).
15. D. J. Brooks, J. R. Fresco, A. M. Lesk, M. Singh, Evolution of Amino Acid Frequencies in Proteins Over Deep Time: Inferred Order of Introduction of Amino Acids into the Genetic Code. *Molecular Biology and Evolution* **19**, 1645-1655 (2002).
16. G. P. Fournier, J. P. Gogarten, Signature of a Primitive Genetic Code in Ancient Protein Lineages. *Journal of Molecular Evolution* **65**, 425-436 (2007).
17. P. K. Keese, A. Gibbs, Origins of genes: "big bang" or continuous creation? *Proceedings of the National Academy of Sciences* **89**, 9489-9493 (1992).
18. S. B. Van Oss, A.-R. Carvunis, De novo gene birth. *PLoS Genet.* **15**, e1008160 (2019).
19. B. Morel, A. M. Kozlov, A. Stamatakis, G. J. Szöllősi, GeneRax: A tool for species tree-aware maximum likelihood based gene family tree inference under gene duplication, transfer, and loss. (2019).
20. M. C. Weiss *et al.*, The physiology and habitat of the last universal common ancestor. *Nat Microbiol* **1**, 16116 (2016).
21. E. R. R. Moody *et al.*, The nature of the last universal common ancestor and its impact on the early Earth system. *Nature Ecology & Evolution* 10.1038/s41559-024-02461-1 (2024).
22. A. J. Crapitto, A. Campbell, A. Harris, A. D. Goldman, A consensus view of the proteome of the last universal common ancestor. *Ecology and Evolution* **12** (2022).
23. Y. Wang, H. Zhang, H. Zhong, Z. Xue, Protein domain identification methods and online resources. *Comput Struct Biotechnol J* **19**, 1145-1153 (2021).

24. J. Mistry *et al.*, Pfam: The protein families database in 2021. *Nucleic Acids Research* **49**,
D412-D419 (2021).
25. H. W. Pi *et al.*, Origin and Evolution of Nitrogen Fixation in Prokaryotes. *Mol Biol Evol* **39**
(2022).
26. P. Laurino, D. S. Tawfik, Spontaneous Emergence of S-Adenosylmethionine and the
Evolution of Methylation. *Angewandte Chemie International Edition* **56**, 343-345 (2017).
27. B. P. S. Chouhan, M. Gade, D. Martinez, S. Toledo-Patino, P. Laurino, Implications of
divergence of methionine adenosyltransferase in archaea. *FEBS Open Bio* **12**, 130-145
(2022).
28. C. Minici *et al.*, Structures of catalytic cycle intermediates of the Pyrococcus furiosus
methionine adenosyltransferase demonstrate negative cooperativity in the archaeal
orthologues. *Journal of Structural Biology* **210**, 107462 (2020).
29. P. Laurino *et al.*, An Ancient Fingerprint Indicates the Common Ancestry of Rossmann-
Fold Enzymes Utilizing Different Ribose-Based Cofactors. *PLOS Biology* **14**, e1002396
(2016).
30. A. D. Goldman, J. T. Beatty, L. F. Landweber, The TIM Barrel Architecture Facilitated the
Early Evolution of Protein-Mediated Metabolism. *J Mol Evol* **82**, 17-26 (2016).
31. P. Z. Kozbial, A. R. Mushegian, Natural history of S-adenosylmethionine-binding proteins.
BMC Struct Biol **5**, 19 (2005).
32. A. J. Michael, Polyamine function in archaea and bacteria. *Journal of Biological*
Chemistry **293**, 18693-18701 (2018).
33. R. Schwartz, S. Istrail, J. King, Frequencies of amino acid strings in globular protein
sequences indicate suppression of blocks of consecutive hydrophobic residues. *Protein*
Science **10**, 1023-1031 (2001).
34. E. Monsellier *et al.*, The Distribution of Residues in a Polypeptide Sequence Is a
Determinant of Aggregation Optimized by Evolution. *Biophysical Journal* **93**, 4382-4391
(2007).
35. A. Irbäck, C. Peterson, F. Potthast, Evidence for nonrandom hydrophobicity structures in
protein chains. *Proceedings of the National Academy of Sciences* **93**, 9533-9538 (1996).
36. S. G. Foy, B. A. Wilson, J. Bertram, M. H. J. Cordes, J. Masel, A Shift in Aggregation
Avoidance Strategy Marks a Long-Term Direction to Protein Evolution. *Genetics* **211**,
1345-1355 (2019).
37. J. E. James, P. G. Nelson, J. Masel, Differential Retention of Pfam Domains Contributes
to Long-term Evolutionary Trends. *Mol Biol Evol* **40** (2023).
38. E. N. Trifonov, The Triplet Code From First Principles. *Journal of Biomolecular Structure*
and Dynamics **22**, 1-11 (2004).
39. V. Lamour *et al.*, Evolution of the Glx-tRNA synthetase family: the glutaminyl enzyme as
a case of horizontal gene transfer. *Proceedings of the National Academy of Sciences* **91**,
8670-8674 (1994).
40. J. Lapointe, L. Duplain, M. Proulx, A single glutamyl-tRNA synthetase aminoacylates
tRNAGlu and tRNAGln in Bacillus subtilis and efficiently misacylates Escherichia coli
tRNAGln1 in vitro. *Journal of Bacteriology* **165**, 88-93 (1986).
41. J. Li *et al.*, The Metal-binding Protein Atlas (MbPA): An Integrated Database for Curating
Metalloproteins in All Aspects. *J Mol Biol* **435**, 168117 (2023).
42. C. Vallières *et al.*, Iron-sulfur protein odyssey: exploring their cluster functional versatility
and challenging identification. *Metallomics* **16** (2024).
43. C. Andreini, I. Bertini, G. Cavallaro, G. L. Holliday, J. M. Thornton, Metal ions in biological
catalysis: from enzyme databases to general principles. *JBIC Journal of Biological*
Inorganic Chemistry **13**, 1205-1218 (2008).
44. R. L. Levine, L. Mosoni, B. S. Berlett, E. R. Stadtman, Methionine residues as
endogenous antioxidants in□proteins. *Proceedings of the National Academy of Sciences*
93, 15036-15040 (1996).

45. A. Bender, P. Hajieva, B. Moosmann, Adaptive antioxidant methionine accumulation in
respiratory chain complexes explains the use of a deviant genetic code in mitochondria.
Proceedings of the National Academy of Sciences **105**, 16496-16501 (2008).
46. M. Schindeldecker, B. Moosmann, Protein-borne methionine residues as structural
antioxidants in mitochondria. *Amino Acids* **47**, 1421-1432 (2015).
47. S. Vieira-Silva, E. P. C. Rocha, An Assessment of the Impacts of Molecular Oxygen on
the Evolution of Proteomes. *Molecular Biology and Evolution* **25**, 1931-1942 (2008).
48. A. Neubeck, F. Freund, Sulfur Chemistry May Have Paved the Way for Evolution of
Antioxidants. *Astrobiology* **20**, 670-675 (2020).
49. M. Granold, P. Hajieva, M. I. Toşa, F.-D. Irimie, B. Moosmann, Modern diversification of
the amino acid repertoire driven by oxygen. *Proceedings of the National Academy of*
Sciences **115**, 41-46 (2018).
50. S. Fukai *et al.*, Structural Basis for Double-Sieve Discrimination of L-Valine from L-
Isoleucine and L-Threonine by the Complex of tRNAVal and Valyl-tRNA Synthetase. *Cell*
103, 793-803 (2000).
51. G. P. Fournier, C. P. Andam, E. J. Alm, J. P. Gogarten, Molecular Evolution of Aminoacyl
tRNA Synthetase Proteins in the Early History of Life. *Origins of Life and Evolution of*
Biospheres **41**, 621-632 (2011).
52. K. R. Olson, Hydrogen sulfide, reactive sulfur species and coping with reactive oxygen
species. *Free Radical Biology and Medicine* **140**, 74-83 (2019).
53. J. Komoto, T. Yamada, Y. Takata, G. D. Markham, F. Takusagawa, Crystal Structure of
the S-Adenosylmethionine Synthetase Ternary Complex: A Novel Catalytic Mechanism
of S-Adenosylmethionine Synthesis from ATP and Met. *Biochemistry* **43**, 1821-1831
(2004).
54. M. A. Grillo, S. Colombatto, S-adenosylmethionine and its products. *Amino Acids* **34**,
187-193 (2008).
55. A. Moreras-Marti, M. Fox-Powell, C. R. Cousins, M. C. Macey, A. L. Zerkle, Sulfur
isotopes as biosignatures for Mars and Europa exploration. *Journal of the Geological*
Society **179**, jgs2021-2134 (2022).
56. G. Eriani, M. Delarue, O. Poch, J. Gangloff, D. Moras, Partition of tRNA synthetases into
two classes based on mutually exclusive sets of sequence motifs. *Nature* **347**, 203-206
(1990).
57. J. Douglas, R. Bouckaert, C. W. Carter, R. Wills, Peter, Enzymic recognition of amino
acids drove the evolution of primordial genetic codes. *Nucleic Acids Research* **52**, 558-
571 (2024).
58. P. G. Higgs, R. E. Pudritz, A Thermodynamic Basis for Prebiotic Amino Acid Synthesis
and the Nature of the First Genetic Code. *Astrobiology* **9**, 483-490 (2009).
59. N. Kitadai, S. Maruyama, Origins of building blocks of life: A review. *Geoscience*
Frontiers **9**, 1117-1153 (2018).
60. J. Fairchild, S. Islam, J. Singh, D.-K. Bučar, M. W. Powner, Prebiotically plausible
chemoselective pantetheine synthesis in water. *Science* **383**, 911-918 (2024).
61. B. Menez *et al.*, Abiotic synthesis of amino acids in the recesses of the oceanic
lithosphere. *Nature* **564**, 59-63 (2018).
62. D. S. C. Lauretta, Harold C. Jr; Grossman, Jeffrey N. ; Polit, Anjani T. ; the OSIRIS-REx
Sample Analysis Team, OSIRIS-REx Sample Analysis Plan -- Revision 3.0. (2023).
63. T. Froese, J. I. Campos, K. Fujishima, D. Kiga, N. Virgo, Horizontal transfer of code
fragments between protocells can explain the origins of the genetic code without vertical
descent. *Scientific Reports* **8** (2018).
64. J. C. Bowman, A. S. Petrov, M. Frenkel-Pinter, P. I. Penev, L. D. Williams, Root of the
Tree: The Significance, Evolution, and Origins of the Ribosome. *Chemical Reviews* **120**,
4848-4878 (2020).
65. A. S. Petrov *et al.*, Evolution of the ribosome at atomic resolution. *Proceedings of the*
National Academy of Sciences **111**, 10251-10256 (2014).

66. D. W. Morgens, A. R. O. Cavalcanti, An Alternative Look at Code Evolution: Using Non-
canonical Codes to Evaluate Adaptive and Historic Models for the Origin of the Genetic
Code. *Journal of Molecular Evolution* **76**, 71-80 (2013).
67. E. V. Koonin, A. S. Novozhilov, Origin and evolution of the genetic code: The universal
enigma. *IUBMB Life* **61**, 99-111 (2009).
68. C. Alvarez-Carreño, A. Becerra, A. Lazcano, Norvaline and Norleucine May Have Been
More Abundant Protein Components during Early Stages of Cell Evolution. *Origins of Life*
and Evolution of Biospheres **43**, 363-375 (2013).
69. Y. Tang, D. A. Tirrell, Attenuation of the editing activity of the Escherichia coli leucyl-tRNA
synthetase allows incorporation of novel amino acids into proteins in vivo. *Biochemistry*
41, 10635-10645 (2002).
70. I. Apostol *et al.*, Incorporation of Norvaline at Leucine Positions in Recombinant Human
Hemoglobin Expressed in Escherichia coli. *Journal of Biological Chemistry* **272**, 28980-
28988 (1997).
71. C. R. Glein, J. A. Baross, J. H. Waite, The pH of Enceladus' ocean. *Geochimica et*
Cosmochimica Acta **162**, 202-219 (2015).
72. K. Vetsigian, C. Woese, N. Goldenfeld, Collective evolution and the genetic code.
Proceedings of the National Academy of Sciences **103**, 10696-10701 (2006).
73. G. Sella, D. H. Ardell, The Coevolution of Genes and Genetic Codes: Crick's Frozen
Accident Revisited. *Journal of Molecular Evolution* **63**, 297-313 (2006).
74. Q. Zhu *et al.*, Phylogenomics of 10,575 genomes reveals evolutionary proximity between
domains Bacteria and Archaea. *Nature Communications* **10** (2019).
75. C. Rinke *et al.*, Insights into the phylogeny and coding potential of microbial dark matter.
Nature **499**, 431-437 (2013).
76. C. T. Brown *et al.*, Unusual biology across a group comprising more than 15% of domain
Bacteria. *Nature* **523**, 208-211 (2015).
77. B. J. Baker *et al.*, Diversity, ecology and evolution of Archaea. *Nat Microbiol* **5**, 887-900
(2020).
78. W.-S. Shu, L.-N. Huang, Microbial diversity in extreme environments. *Nature Reviews*
Microbiology **20**, 219-235 (2021).
79. P. Jones *et al.*, InterProScan 5: genome-scale protein function classification.
Bioinformatics **30**, 1236-1240 (2014).
80. K. Katoh, D. M. Standley, MAFFT multiple sequence alignment software version 7:
improvements in performance and usability. *Mol Biol Evol* **30**, 772-780 (2013).
81. B. Q. Minh *et al.*, IQ-TREE 2: New Models and Efficient Methods for Phylogenetic
Inference in the Genomic Era. *Mol Biol Evol* **37**, 1530-1534 (2020).
82. C. C. Dang *et al.*, nQMaker: Estimating Time Nonreversible Amino Acid Substitution
Models. *Syst Biol* **71**, 1110-1123 (2022).
83. B. Q. Minh, C. C. Dang, L. S. Vinh, R. Lanfear, QMaker: Fast and Accurate Method to
Estimate Empirical Models of Protein Evolution. *Syst Biol* **70**, 1046-1060 (2021).
84. L. J. Revell, phytools: an R package for phylogenetic comparative biology (and other
things). *Methods in Ecology and Evolution* **3**, 217-223 (2012).
85. U. Mai, E. Sayyari, S. Mirarab, Minimum variance rooting of phylogenetic trees and
implications for species tree reconstruction. *PLOS ONE* **12**, e0182238 (2017).
86. F. D. K. Tria, G. Landan, T. Dagan, Phylogenetic rooting using minimal ancestor
deviation. *Nature Ecology & Evolution* **1** (2017).
87. S. Naser-Khdour, B. Quang Minh, R. Lanfear, Assessing Confidence in Root Placement
on Phylogenies: An Empirical Study Using Nonreversible Models for Mammals.
Systematic Biology **71**, 959-972 (2022).
88. G. P. Fournier, E. J. Alm, Ancestral Reconstruction of a Pre-LUCA Aminoacyl-tRNA
Synthetase Ancestor Supports the Late Addition of Trp to the Genetic Code. *J Mol Evol*
80, 171-185 (2015).

89.   J. Helske, diagis: Diagnostic Plot and Multivariate Summary Statistics of Weighted
Samples from Importance Sampling. https://doi.org/10.32614/CRAN.package.diagis
(2023).
90.   H. Seltman, Approximations for mean and variance of a ratio.
https://www.stat.cmu.edu/~hseltman/files/ratio.pdf (2012).
91.   R Core Team, R: A Language and Environment for Statistical Computing. https://www.R-
project.org/ (2024).
92.   J. Hallgren *et al.*, DeepTMHMM predicts alpha and beta transmembrane proteins using
deep neural networks. *bioRxiv* 10.1101/2022.04.08.487609, 2022.2004.2008.487609
(2022).
93.   F. Keck, Handling biological sequences in R with the bioseq package. *Methods in*
Ecology and Evolution **11**, 1728-1732 (2020).
94.   W. F. Doolittle, The nature of the universal ancestor and the evolution of the proteome.
Current Opinion in Structural Biology **10**, 355-358 (2000).
95.   R. L. Tatusov, M. Y. Galperin, D. A. Natale, E. V. Koonin, The COG database: a tool for
genome-scale analysis of protein functions and evolution. *Nucleic Acids Research* **28**,
33-36 (2000).
96.   The UniProt Consortium, UniProt: the universal protein knowledgebase. *Nucleic Acids*
Research **45**, D158-D169 (2016).
97.   J. Huerta-Cepas *et al.*, eggNOG 5.0: a hierarchical, functionally and phylogenetically
annotated orthology resource based on 5090 organisms and 2502 viruses. *Nucleic Acids*
Research **47**, D309-D314 (2018).
98.   A. Szilágyi, P. Závodszky, Structural differences between mesophilic, moderately
thermophilic and extremely thermophilic protein subunits: results of a comprehensive
survey. *Structure* **8**, 493-504 (2000).
99.   S. Fukuchi, K. Yoshimune, M. Wakayama, M. Moriguchi, K. Nishikawa, Unique amino
acid composition of proteins in halophilic bacteria. *J Mol Biol* **327**, 347-357 (2003).
100. T. Therneau, deming: Deming, Theil-Sen, Passing-Bablock and Total Least Squares
Regression. https://doi.org/10.32614/CRAN.package.deming (2024).
840