Optimizing Phylogenetic Eigenvector Regression: Union Eigenvectors,

Robust Estimation, and Flexible Application to Comparative Analyses

Zheng-Lin Chen and Deng-Ke Niu[*]

MOE Key Laboratory for Biodiversity Science and Ecological Engineering, College of Life

Sciences, Beijing Normal University, Beijing, 100875, China.

E-mail addresses: chenzljx@gmail.com (Z.-L. Chen), dkniu@bnu.edu.cn,

dengkeniu@hotmail.com (D.-K. Niu)

*Correspondence author: Deng-Ke Niu, College of Life Sciences, Beijing Normal University,

Beijing, 100875, China.

Abstract

Phylogenetic eigenvector regression (PVR) is widely used in ecology and evolution by representing phylogenetic structure through separable eigenvectors. Despite this flexibility, its implementation faces three key challenges: (1) the selection of eigenvectors, (2) the reduced robustness of ordinary least-squares (OLS) regression under shift-like evolutionary heterogeneity, and (3) the applicability of conventional model complexity rules such as the "samples-per-variable (SPV) $\geq$ 10" guideline. Here, we propose an optimized PVR framework that addresses these limitations. First, we show that trait-specific selections of eigenvectors often diverge, sometimes producing inconsistent results, and that using their union offers stronger control of phylogenetic non-independence. Second, we evaluate robust regression estimators within PVR, demonstrating that PVR-MM – and in most cases PVR-L2, the standard OLS estimator – maintains high accuracy under non-stationary evolutionary shifts where other non-robust methods fail. Third, through simulation, we reassess the SPV $\geq$ 10 rule, showing that PVR tolerates eigenvector counts well beyond this threshold, offering greater flexibility while requiring attention to potential overfitting. Extensive simulations across diverse trees and evolutionary scenarios confirm that the optimized framework improves accuracy and robustness. By addressing key aspects of eigenvector selection, regression, and model complexity, our findings strengthen the reliability and applicability of PVR.

Keywords: phylogenetic comparative methods; robust regression; evolutionary shift; phylogenetic autocorrelation; phylogenetic signal; trait correlation.

Introduction

Comparative studies of traits across species must account for phylogenetic non-independence (Cornwallis & Griffin, 2024; Dewar et al., 2025; Felsenstein, 1985; Garamszegi, 2014). Among the available approaches, phylogenetic eigenvector regression (PVR) has become widely used in ecology and evolutionary biology because it expresses phylogenetic information as separable eigenvectors (EVs) obtained by decomposing the phylogenetic distance or covariance structure encoded in the tree (Diniz-Filho et al., 2012a; Diniz-Filho et al., 2014; Diniz-Filho et al., 2011; Diniz-Filho et al., 1998; Diniz-Filho & Tôrres, 2002). This representation provides flexibility because it captures phylogenetic structure directly from the tree without requiring the specification of an explicit evolutionary model, while also accommodating extensions to other forms of autocorrelation such as spatial or temporal structure (Kühn et al., 2009; Maynard et al., 2022; Peres-Neto et al., 2012; Safi & Pettorelli, 2010; Wang et al., 2019) and with machine learning methods (Debastiani et al., 2021; Penone et al., 2014). Such versatility has helped promote the broad adoption of PVR in ecological applications and highlights the importance of refining its implementation (Divíšek et al., 2018; Gao et al., 2023; Lanuza et al., 2023; LeRoy et al., 2020).

Despite its utility, two issues limit PVR performance. First, EVs selected from each trait separately may yield inconsistent results, raising concerns when analyzing relationships between traits. Second, conventional PVR relies on ordinary least squares regression, which can be sensitive to non-stationary evolutionary shifts or outliers. These challenges motivate the development of optimized strategies that improve reliability while retaining the flexibility of PVR.

Here, we propose an optimized PVR framework that combines (i) a union-based EV set (*EV_U*) to capture shared phylogenetic structure, (ii) robust regression estimators to reduce sensitivity to non-stationarity (Adams et al., 2024; Huber, 1964; Slater & Pennell, 2014; Yu & Yao, 2017), and (iii) a reassessment of the conventional "samples-per-variable (SPV, i.e., events per variable) $\geq$ 10" rule (Babyak, 2004), showing that PVR remains reliable with far fewer samples per predictor and emphasizing the need to check EV counts before model fitting. Using extensive simulations under a range of evolutionary scenarios and tree structures, we assess the accuracy of these strategies relative to established comparative methods and highlight the conditions under which optimized PVR offers particular advantages.

Materials and Methods

Simulation of phylogenetic trees

We evaluated PVR on a set of fully bifurcating trees chosen to vary in branch-length scaling,

taxon sampling, and topological balance, generated with *stree* (**ape** 5.8) and, where necessary,

combined with *paste.tree* (**phytools** 2.3-0) (Paradis & Schliep, 2019; Revell, 2024):

1. **Baseline**: 128-tip balanced tree, Grafen (1989) scaling $\rho = 1$. (ρ is the exponent in

Grafen's branch-length transformation; $\rho = 1$ yields equal branch-length scaling across

hierarchical levels.)

2. **Compressed deep branches**: 128-tip balanced tree, $\rho = 0.1$ (short ancestral, long

terminal branches).

3. **Elongated deep branches**: 128-tip balanced tree, $\rho = 2$ (long ancestral, short terminal

branches).

4. **Extreme imbalance**: 128-tip ladder (comb-shaped) tree, $\rho = 1$.

5. **Small-sample setting**: 16-tip tree generated under the same four branch-length

conditions (balanced with $\rho = 1$, 0.1, 2, and ladder).

Simulated trait generation

We generated paired traits X_1 and X_2 on each of the eight phylogenetic trees under six

evolutionary settings adapted from Revell (2010), Adams et al. (2024), and Chen et al.

(2025). In all scenarios, the generative model for X_2 is $X_2 = \beta X_1 + \varepsilon$, where β denotes the

regression slope specifying the intended linear dependence of X_2 on X_1, and ε contributes

additional noise (Table 1).

These six settings fall into two broad categories: (i) four gradual-evolution scenarios, in

which traits evolve without abrupt changes, and (ii) two shift scenarios, in which an abrupt

change is introduced on a single deep branch. The four gradual-evolution scenarios (Table 1)

involve traits with noise that either evolved under Brownian motion (BM) or with

independent and identically distributed (i.i.d.) normal (non-phylogenetic) noise (Norm). In all

scenarios, X_1 is generated first, and X_2 is then constructed as a linear function of X_1 plus a

noise term ε. The four gradual-evolution scenarios differ in whether X_1 and ε evolve under

Brownian motion (BM) or independent normal noise (Norm), yielding the combinations BM–

BM, BM–Norm, Norm–BM, and Norm–Norm. For example, in the BM–Norm scenario, X_1

follows Brownian motion, whereas X_2 inherits the phylogenetic structure of X_1 through the

βX_1 term, with additional independent and identically distributed (i.i.d.) normal noise ε. The

variance of ε is determined by a numerical gradient (10^{-4}, 0.01, 1, 4, 16, 64, 256, 1024),

modifying noise intensity and phylogenetic dilution, here referring to the progressive

weakening of phylogenetic signal in trait correlations as non-phylogenetic noise increases. In contrast, in the two shift scenarios (X_1-shift and Dual-shift, Table 1), the same numerical gradient specifies the shift magnitude s applied to one of the two sister branches descending directly from the root, producing non-stationary, abrupt evolutionary changes in X_1 alone or in both X_1 and ε, where non-stationary refers to localized deviations from homogeneous evolutionary dynamics along the phylogeny. Together, these six scenarios span a wide range of phylogenetic signals, noise intensities, and stationary versus non-stationary dynamics, providing a robust and diverse framework for benchmarking PVR variants and comparison methods. For clarity, the full combination of evolutionary scenarios, variance or shift magnitudes, and replicate counts is summarized in Table 1.

Defining process-level ground truth

A commonly used approach in comparative-method simulations is to define true and false positives directly by the generating effect size β, with $\beta = 0$ representing an absence of true association and $\beta \neq 0$ representing a true association (Adams & Collyer, 2022). This strategy is intuitive and effective for simulation designs in which the structural signal specified by β is expressed consistently across datasets, and where stochastic variation and finite sample effects do not substantially alter the resulting data-level association patterns.

However, our simulation framework differs in a key respect: because trait evolution is stochastic on a fixed phylogeny, each replicate expresses its own realized statistical association, determined jointly by the coupling parameter β and the stochastic variation encoded by the noise component ε. As a result, the association observed in any given dataset may diverge from that implied by the generative value of β.

Most notably, when $\beta = 1$, the noise term ε can be sufficiently large – particularly under higher-variance settings such as 256 or 1024 – to mask or overwhelm the β-induced linear trend. In such cases, the realized population-level association may be weak, often leading to nonsignificant statistical results in finite samples despite a nonzero generative β. By the same logic, when $\beta = 0$, any apparent association reflects stochastic evolutionary variation and may be observed in individual realizations despite the absence of structural coupling. Consequently, a method's "true" performance cannot be assessed by whether it recovers statistical relationship implied by the coupling parameter β; instead, evaluating performance requires a dataset-specific benchmark that reflects the realized association in each replicate.

135 Table 1. Six evolutionary scenarios for simulating paired traits X_1 and X_2 (generated as $X_2 = \beta X_1 + \varepsilon$).

Scenario	X_1	ε	Shift?	β	Explanation/purpose
Gradual-evolution scenarios (stationary)					
BM–BM	BM	BM	No	1	X_1 and ε both follow BM, so X_2 is a linear combination of two BM components. As a result, realized associations between X_1 and X_2 can vary substantially across replicates, particularly as stochastic variance increases, despite a fixed structural relationship in the data-generating process.
BM–Norm	BM	Normal	No	1	X_1 follows BM, while ε is drawn from an i.i.d. normal (non-phylogenetic) distribution. The non-phylogenetic ε reduces the phylogenetic signal in X_2, causing realized associations to vary across replicates and, under high noise variance, potentially weaken the correlation relative to that implied by the generative β.
Norm–BM	Normal	BM	No	1	X_1 is i.i.d. normal; ε adds phylogenetic structure to X_2. Realized correlations between X_1 and X_2 vary across replicates and may be substantially attenuated by noise relative to that implied by the generative β.
Norm–Norm	Normal	Normal	No	1	Neither trait carries phylogenetic signal. X_1 and X_2 vary independently, and realized associations may be weak or highly variable depending on noise variance.
Shift scenarios (non-stationary, single-branch abrupt shifts)					
X_1-shift	BM + shift	Normal	X_1 only	1	Extends the BM–Norm scenario by adding a single-branch shift to X_1, while fixing $\varepsilon \sim \mathcal{N}(0, 0.01)$. Because ε is very small, the background evolutionary association between X_1 and X_2 remains tightly coupled across most of the phylogeny, whereas the shifted branch introduces a localized deviation from this otherwise strong relationship.
Dual-shift	BM + shift	BM + shift	Both	0	X_1 and ε both undergo a shift on the same branch, and no causal association is intended between X_1 and X_2. Both X_1 and ε are simulated independently under BM, with the same root value (1) and variance ($\sigma^2 = 1$), ensuring that shifts occur on the same scale. Such synchronized shifts may induce spurious overall correlation signals in methods that lack robustness to localized shifts.

The generative model for X_2 consists of a linear combination of X_1 and random noise ε, with β controlling the intended association. In the two shift scenarios, an additional
shift parameter s displaces trait means on a single deep branch. A series of values (10^{-4}, 0.01, 1, 4, 16, 64, 256, 1024) was used to model both the shift magnitude and the
gradient of ε's variance, influencing either the variance of ε in stationary models or the shift size in non-stationary scenarios, and thereby modulating the strength of
population-level correlations. Brownian-motion (BM) traits were simulated using *rTraitCont* (**ape** 5.8) with rate = 1, and normal values were generated using *rnorm* (**stats**
4.1.3). Each magnitude was replicated 1,000 times, yielding 8,000 datasets per scenario and 48,000 datasets per tree. For implementation details of the shift simulations, see
Chen et al. (2025).

To accomplish this, we used branchwise standardized changes ($\Delta X/L$, where L denotes

branch length) as a replicate-specific, phylogenetically independent benchmark. For every

simulation, trait values at all nodes were recorded and standardized changes were computed

on each branch ($\Delta X_1/L$ and $\Delta X_2/L$, for each trait respectively). Because these branchwise

changes are independent across the tree, they provide a tree-wide ground-truth sample for the

statistical association realized in that dataset. This branchwise differencing approach is

conceptually related to Felsenstein's (1985) phylogenetically independent contrasts (PIC),

which also leverage the independence of evolutionary changes along branches. But unlike

PIC, our procedure does not require ancestral-state reconstruction or any Brownian-motion

assumptions, as all node states are known in our simulations.

We quantified the realized association using Spearman's rank correlation across the

branchwise changes, which provides a robust, rank-based summary of the association realized

in each dataset. We then evaluated each comparative method against this benchmark: a

method was considered correct only if it recovered both (i) the direction of association and (ii)

the significance decision ($p < 0.05$) for that replicate. Across 1,000 replicates for each

experimental condition, method accuracy was defined as the proportion of correct inferences.

Phylogenetic eigenvectors selection methods

EVs were obtained by principal-coordinate analysis of the double-centered cophenetic

distance matrix, yielding $n - 1$ axes. For each trait (X_1 or X_2), we selected a subset of EVs

with three published criteria (Diniz-Filho et al., 2012a; Diniz-Filho & Nabout, 2009):

1. **ESRRV**: keep EVs whose Pearson or Spearman correlation with the focal trait is

significant, choosing the test according to the trait's normality.

2. **mAIC**: forward stepwise regression adding EVs that lower Akaike Information

Criterion (AIC), implemented with *stepAIC()* (**MASS** 7.3-55) (Venables & Ripley,

2002).

3. **mMorI**: forward selection until the residual Moran's I is minimized. Because

phylogenetic autocorrelation is almost always positive, we modified this rule to stop

as soon as residual Moran's I becomes negative; if it never does, selection continues to

the usual non-significant threshold.

Custom R wrappers *ESRRV()* and *MoransI()* implement rules 1 and 3, while all other

calculations use the original **PVR** package (Diniz-Filho et al., 2012a; Diniz-Filho et al., 1998;

Diniz-Filho et al., 2012b).

Correcting phylogenetic autocorrelation in trait regressions

After EV selection we compared four ways to purge phylogenetic signal:

1. **EV_X_1 model**: $X_2 \sim X_1 + EV\_X_1$ — covariates are the EVs chosen for X_1.

2. **EV_X_2 model**: $X_2 \sim X_1 + EV\_X_2$ — uses the EVs chosen for X_2. Writing it in reverse form, $X_1 \sim X_2 + EV\_X_2$, controls the same EVs, so the correlation (and its p-value) is identical, although the regression slope becomes the reciprocal of the original.

3. **EV_U model**: $X_2 \sim X_1 + EV\_U$ — EV_U is the union of EVs selected from both traits.

4. **Residual correlation**: Regress each trait on its own EVs, then take the Pearson correlation of the residuals. To curb type-I error from over-fitting, p-values were based on $df = n - k - 1$, where n is the number of species and k the number of EVs in EV_U. While this conservative adjustment may reduce power, it controls for inflated false positives due to shared structure among EVs (Diniz-Filho & Tôrres, 2002).

Robust regressions in PVR

To test the sensitivity of PVR to outliers and non-normal errors we compared it with five different regression estimators:

1. **L2**: as termed in robust regression (Huber, 1964), corresponds to the standard ordinary least-squares (OLS) estimator that minimizes squared residuals under homoscedastic normal errors. It was implemented using *lm()* (package **stats**, R 4.1.3).

2. **L1**: least-absolute-deviation (*lad()*, **L1pack** 0.41-245) (Phillips, 2002).

3. **M**: Huber-type M-estimator (*rlm()*, **MASS** 7.3-55) (Venables & Ripley, 2002).

4. **S**: high-breakdown S-estimator (*lmRob()*, **robust** 0.7-4) (Gervini & Yohai, 2002).

5. **MM**: S start followed by an M step (*lmRob()*, **robust**).

Among these estimators, L2 serves as the baseline; L1, M, S, and MM offer progressively stronger resistance to leverage points while retaining efficiency. For detailed properties and phylogenetic applications of these estimators, see Adams et al. (2024).

Results

Unless otherwise noted, in the subsequent analyses, PVR regressions control phylogeny by including as covariates the EVs selected for each trait separately (EV_X_1, EV_X_2) or their union (EV_U).

Eigenvector selection and budgeting

We applied three EV-selection rules (ESRRV, mAIC, mMorI) separately to X_1 and X_2; the

resulting EV sets, together with their union (EV_U), were carried forward as candidate

covariates for examination. As detailed later through mathematical derivations and PVR

accuracy assessments in the section "*Why use the union of eigenvectors?*", EV_U plays an

important role in controlling phylogenetic autocorrelation, so here we give special attention to

|EV_U| (the number of eigenvectors in EV_U) when comparing outcomes across rules,

without disregarding the other EV sets.

Across all four tree scenarios with 128 species, the size of the EV_U set differed

markedly among selection rules (Fig. 1). mAIC consistently returned the largest sets, with

medians ranging from 123 to 125 EVs across topologies. By comparison, ESRRV produced

EV_U medians ranging from 6 to 31, and mMorI from 1 to 37; under the balanced 128-

species tree with ρ =1, both methods returned a median of 9.

The four scenarios with 16 species (Fig. S1) showed the same relative ordering across

selection rules and tree types, even though the absolute EV_U counts scaled downward

proportionally with the reduced tree size. This confirms that EV counts are influenced

primarily by tree topology and the selection rule, rather than sample size.

Fig. 1. Number of eigenvectors (EVs) retained by three selection criteria (ESRRV, mAIC and mMorI) for a 128-
species phylogenetic tree. EVs were picked separately for X_1 and X_2; EV_U is their union. Violin outlines depict
the full distribution of selected EV counts; white boxes show the inter-quartile range, black whiskers the 5th–
95th percentiles, and red dots mark the median of each distribution. The red dashed line marks the upper bound
implied by the conventional samples-per-variable ≥ 10 guideline, calculated here assuming one non-EV
predictor (128 spp./10 = 12.8, meaning 11.8 EVs plus 1 predictor). Selection criteria are described in Materials
and Methods.

In our SPV calculations, the sample size is the number of species at the tips of the tree,

and the variable count includes both the retained EVs and the single predictor trait (X_1),

giving |EV| + 1 variables in total. The widely recommended "SPV ≥ 10" rule (Austin et al.,

2017; Babyak, 2004; Peduzzi et al., 1996) originally proposed to prevent over-

parameterization and unstable inference in conventional regression – states that SPV should

be at least 10. So for our 128 species trees, this would imply an upper bound of 12.8 variables

(i.e., ~12 EVs + 1 predictor). However, because EVs are orthogonal components derived from

the phylogenetic distance matrix rather than biological or environmental predictors, this rule

does not carry the same meaning in PVR as it does in ecological regressions.

To test this idea, we analyzed the subset of simulated datasets corresponding to balanced

trees with ρ = 1 and n = 128 species, using six evolutionary scenarios. For each scenario, we

examined five |EV_U| settings, corresponding to SPV ratios of approximately 1.5, 2, 3, 6, and

the EV count selected by mMorI. In this setting, mMorI typically returned fewer EVs than the

conventional SPV ≥ 10 threshold yet was already sufficient to control phylogenetic

autocorrelation and avoid false positives due to residual structure. We then augmented the

mMorI-selected EV_U core set (median |EV_U| = 9) by adding additional EVs in descending

eigenvalue order to reach the 6, 3, 2, and 1.5 SPV ratio targets (|EV_U| = 20, 42, 63, and 84,

respectively). When independence was effectively addressed by the EV selected using

mMorI, differences in PVR accuracy across these EV_U-count gradients are attributable

mainly to potential overfitting from excessive predictors.

In the six simulation scenarios, the impact of EV count on PVR accuracy varied. Here

'PVR accuracy' denotes the accuracy of the PVR-controlled trait–trait regression $X_2 \sim X_1 +$

EV_U, as defined in the Materials and Methods, where the benchmark for determining correct

inference is described in detail.

Across the four gradual-evolution scenarios (Figs. 2a–d), accuracy showed distinct

patterns as EV count increased. In the three gradual-evolution cases (Figs. 2a, 2b, and 2d),

accuracy began to decline when |EV_U| = 63 and declined further at |EV_U| = 84. In the

Norm–BM scenario (Fig. 2c), accuracy exhibited only modest variation across EV counts,

and the mMorI-selected set (median = 9 EVs) was not among the best-performing choices.

We next turn to the two non-stationary scenarios (Figs. 2e–f), which introduce an abrupt

evolutionary shift on a deep branch, either in X_1 alone or in both X_1 and ε. Under the single-

shift condition (Fig. 2e), accuracy was nearly identical across all EV counts, indicating no

sensitivity to EV number. In the dual-trait shift scenario (Fig. 2f), accuracy differed only

slightly across EV counts. Unlike Figs. 2a, 2b and 2d – where the *EV_U* obtained directly by mMorI was among the strongest choices and modest EV increases performed similarly – the larger EV sets (63 and 84) showed a marginal but consistent accuracy advantage over the *EV_U* set obtained directly by mMorI.

Fig. 2. Accuracy of PVR under six simulation scenarios with varying numbers of eigenvectors (*EV_U*). Panels (a–d) show the four gradual-evolution (stationary) scenarios, whereas panels (e–f) show the two shift scenarios involving a single-branch abrupt evolutionary change. All simulated on a balanced phylogenetic tree with 128 species and uniformly scaled branch lengths ($\rho = 1$). Lines represent PVR accuracy for different *EV_U* sizes. The mMorI-selected *EV_U* (median $|EV\_U| = 9$) serves as the core set, and larger *EV_U* sizes ($|EV\_U| = 20, 42, 63,$ and 84) were obtained by augmenting this core set with additional eigenvectors in descending eigenvalue order. Accuracy is defined with respect to each dataset's realized association, based on replicate-specific branchwise standardized changes ($\Delta X/L$). For details on EV selection, scenario and accuracy definitions, see Materials and Methods; the data used to plot this figure appear in Table S1.

Taken together, these results in Fig. 2 show that modest increases in EV count – such as expanding from the mMorI-selected set (median = 9 EVs) to intermediate sizes around 40 EVs, corresponding to a SPV ratio of roughly 3 – do not dramatically reduce accuracy under gradual-evolution scenarios, and under shift scenarios the PVR results are more insensitive to EV number. These patterns reflect the behavior of EVs themselves rather than the

performance of any particular EV-selection strategy evaluated in Fig. 1 and Fig. S1, and also

illustrate why the conventional SPV-based heuristic does not directly translate to PVR, given

the orthogonality of phylogenetic eigenvectors.

These findings also clarify the implications for EV-selection rules. Across all tree

topologies (Fig. 1 and Fig. S1), mAIC consistently returned very large EV sets, often far

exceeding the range in which accuracy stabilizes in Fig. 2, and is therefore unsuitable for

PVR. ESRRV generally produced moderate EV counts and can be applied across topologies,

but mMorI frequently returned very few EVs – including zero in some unbalanced trees –

which limits its ability to control phylogenetic autocorrelation. However, from a

methodological perspective, mMorI has the conceptual advantage of explicitly minimizing

residual Moran's I – the most direct criterion for removing phylogenetic dependence.

Accordingly, in balanced trees with $\rho = 1$, where ESRRV and mMorI return similar EV

counts, the comparative results presented in subsequent sections indicate that mMorI may be

preferable because it directly optimizes phylogenetic residual structure.

Trait-specific selection of EVs and its downstream effects on PVR estimation

Although retaining somewhat larger EV sets does not markedly reduce PVR accuracy overall,

most empirical applications of PVR select eigenvectors only for the predictor variable,

implicitly assuming that the EV sets selected for different traits do not differ enough to

influence inference (e.g., Álamo et al., 2024; Carotenuto et al., 2015; Gao et al., 2023; Liu et

al., 2024; Seger et al., 2025; Vilela et al., 2014). However, whether the EV sets selected for

different traits coincide or diverge – and, if they diverge, whether such differences materially

affect PVR estimation – remains unclear. This motivates a systematic evaluation of trait-

specific selection of EVs and its consequences.

To explore this, we conducted simulations using balanced phylogenetic trees with 128

species and uniformly scaled branches ($\rho = 1$). We analyzed paired predictors under various

evolutionary scenarios to quantify how divergence in EV sets impacts regression outcomes.

Across 8,000 simulations per scenario (Table 2), the EV sets selected for two traits (X_1

and X_2) disagreed in 76.4% of cases on average. The highest rate of disagreement occurred

under the dual-trait shift scenario, with notable inconsistencies observed across all scenarios.

These results indicate that EV selection is highly sensitive to which trait is treated as the

dependent variable.

Given the high disagreement rates between the EV sets selected for X_1 and X_2 (Table 2),

we next examined whether these differences translate into divergent PVR outcomes. To do so,

we fit parallel PVR models using either EV_X_1 or EV_X_2 as covariates and compared the

resulting inferences on the X_1–X_2 relationship. Table 3 reports the proportion of replicates in

which the two regressions yielded different conclusions (i.e., differing in direction or

significance). These analyses quantify the practical consequences of EV-set discrepancies:

when X_1 and X_2 select different eigenvectors, the downstream PVR conclusions may also

diverge. The magnitude of this effect varies across evolutionary scenarios, with the largest

discrepancies occurring under the BM–BM and dual-shift conditions when EVs were selected

by ESRRV.

Table 2. Frequency with which the EV sets chosen for X_1 and X_2 do not match.

Simulation scenarios	ESRRV	mMorI
BM–BM	83.3%	84.1%
BM–Norm	80.4%	81.4%
Norm–BM	86.3%	75.8%
Norm–Norm	86.0%	4.5%
X_1-shift	91.0%	49.2%
Dual-shift	94.5%	100%

333 A disagreement is recorded whenever the EV sets selected for X_1 and X_2 differ in at least one eigenvector. Values
334 represent the percentage of inconsistent selections across 8,000 simulations per scenario. Methods (ESRRV and
335 mMorI) and scenarios are detailed in Materials and Methods. The raw data underlying this summary appear in
336 Table S2.

337
338 Table 3. Proportion of simulations in which PVR models using the EV set from X_1 versus that from X_2 yield
339 discordant results.

Simulation scenarios	ESRRV	mMorI
BM–BM	19.8%	11.4%
BM–Norm	9.18%	18.2%
Norm–BM	11.5%	10.8%
Norm–Norm	6.44%	0.13%
X_1-shift	14.1%	6.02%
Dual-shift	30.7%	14.9%

This table evaluates the downstream effects of the EV-set discrepancies reported in Table 2. The results of a
sample are counted as discordant when the two regressions differ in statistical significance ($p < 0.05$ vs ≥ 0.05)
or slope sign (≥ 0 vs < 0). Percentages are based on 8 000 datasets per scenario. See Materials and Methods for
details on EV selection methods (ESRRV and mMorI) and simulation scenarios. The raw data underlying this
summary appear in Table S3.

To further quantify the extent of divergence, we computed the absolute difference in

partial correlation coefficients for the discordant cases:

$$|\Delta\rho| = |pcor(X_1, X_2 + EV\_X_1) - pcor(X_2, X_1 + EV\_X_2)|.$$

Over 73% of discordant comparisons showed $|\Delta\rho| > 0.1$, and 15% exceeded 0.5,

demonstrating substantial variation in effect size and even sign reversals in biological

interpretation (Fig. 3). The X_1-only shift scenario was especially extreme, with median $|\Delta\rho|$

values of 1.04 (ESRRV) and 0.97 (mMorI). Even among the 73,318 cases where models

agreed in significance and slope sign, 31% showed $|\Delta\rho| > 0.1$, and 4% exceeded 0.5 (Fig. S2).

Thus, consistent significance does not necessarily imply consistent effect estimation.

Fig. 3. Distributions of absolute differences in partial correlation coefficients between comparing PVR models
using EV_X_1 versus EV_X_2 as covariates under six evolutionary scenarios. The violin plot illustrates the absolute
differences in partial correlation coefficients $|pcor(X_1, X_2 + EV\_X_1) - pcor(X_2, X_1 + EV\_X_2)|$ for cases with
conflicting PVR results; numbers beneath each violin give the sample size (n). Partial correlations were
computed with ppcor::pcor() (Pearson). Red dots indicate the median values, black vertical bars span the inter-
quartile range, and black whiskers extend to the 5th–95th percentiles. Paired t-tests revealed significant
differences in partial-correlation estimates between EV_X_1 and EV_X_2 in every scenario (all $p < 0.001$) except
when mMorI was applied to the Norm−Norm scenario data, where the test was non-significant ($n = 10$, $p =$
0.373). See Materials and Methods for details of the EV-selection procedures (ESRRV, mMorI) and the BM,
Norm and Shift simulation variants; the raw values plotted here are provided in Table S4.
367

In summary, trait-specific selections of EVs frequently diverge (Table 2), and in a

measurable subset of cases these divergences lead to different PVR conclusions, manifested

as differences in significance or moderate-to-large coefficient shifts. The magnitude of these

effects depends on the evolutionary context, with certain scenarios (e.g., dual-shift or trait-

specific shifts) showing particularly pronounced instability. Moreover, preliminary results

(Fig. 1b) indicate that under alternative branch-length scaling ($\rho = 2$), the EV sets selected for

X_1 versus X_2 – when under the mMorI rule – differ far more than in the baseline $\rho = 1$ setting

(Fig. 1a), suggesting an even greater potential for distorted inference.

Why use the union of eigenvectors?

The EV sets selected for X_1 and X_2 frequently differ (Table 2), and these differences lead to

distinct PVR outcomes in a subset of replicates (Table 3). This inconsistency prompts a more

fundamental theoretical question: can such per-trait EV sets adequately control for phylogeny

when two traits share only part of their phylogenetic signal?

When either trait is filtered through only its own eigenvector set, phylogenetic structure

unique to the partner trait remains unmodeled, potentially biasing the estimated association.

Our derivation (Appendix A) shows that using the union of both EV sets (EV_U) more fully

captures the phylogenetic component (P) in Cheverud's (1985) decomposition of phenotypic

variation (T = P + S), resulting in more stable and unbiased regression estimates. This aligns

with Westoby et al.'s (2023) critique that phylogenetic generalised least squares (PGLS)

corrects for phylogeny in the response but not in the predictor.

Motivated by the same concern, we additionally considered a two-step alternative: each

trait is first regressed on its own EVs to remove trait-specific phylogenetic signal, and then an

OLS regression is applied to the resulting residuals to quantify their association. This

residual-based approach offers a different path to adjusting for phylogeny in both variables

and provides a comparative baseline for evaluating the EV_U method.

Building on the earlier results (Fig. 1a), where both ESRRV and mMorI produced

admissible EV counts under a balanced tree with 128 species and $\rho = 1$, we next used

simulations to assess whether the choice of selection method materially influences PVR

outcomes. Additionally, we tested whether regressions based on EV_U outperform those

relying on EVs selected separately for each trait or on residual approaches.

Figure 4 compares eight PVR configurations: two EV-selection methods (ESRRV,

mMorI), each combined with four regression strategies (EV_X_1, EV_X_2, EV_U, and residual).

Because accuracy is averaged over 1,000 replicates – and EV-set differences affect outcomes

only in a subset of cases (Table 3) – the four regression strategies appear broadly similar.

When EV_U equals the larger of EV_X_1 or EV_X_2, the corresponding curves overlap, further

reducing visual differences among methods. Despite this overall similarity, systematic

differences still emerge. Across scenarios, EV_U-based regressions (red lines) – regardless of

whether paired with ESRRV (dashed) or mMorI (solid) – generally perform better than the other regression strategies (blue, green, and black lines). Among these, mMorI + EV_U (red solid line) consistently ranked first or jointly first across all six evolutionary scenarios and shift magnitudes, making it the most reliable configuration overall.

Fig. 4. Performance across different phylogenetic eigenvector (EV) selection methods and regression strategies. This figure compares two EV selection methods (ESRRV and mMorI), each combined with four regression approaches: (1) using EV_X_1 as covariates; (2) using EV_X_2; (3) using their union, EV_U; and (4) regressing the residuals of X_1 and X_2, each pre-adjusted by their respective EVs. Simulations were conducted on balanced trees with 128 species ($\rho = 1$) across six scenarios: (a–d) the four gradual-evolution (stationary) scenarios – BM–BM, BM–Norm, Norm–BM, and Norm–Norm – and (e–f) the two single-branch shift scenarios – X_1-only shift and dual-trait shift. In some cases, when multiple method combinations produced identical results, their lines overlap and may appear as a single curve in the plot. Accuracy values are evaluated against each dataset's realized association, estimated from replicate-specific branchwise standardized changes. For details on EV selection, scenario and accuracy definitions, see Materials and Methods. The source data for this figure are provided in Table S5.

While residual-based regressions (black lines) also account for phylogenetic signals in both traits, their performance was generally inferior to that of EV_U-based strategies (Fig. 4). This discrepancy likely arises because removing phylogenetic signal independently for each trait can also eliminate shared structure that underlies true biological associations. In contrast, EV_U integrates information from both traits within a single model, thereby preserving shared

structure while controlling for phylogenetic effects, and consequently achieved higher

accuracy across scenarios.

A notable special case occurs in Fig. 4e, where the two traits are highly correlated but X_1

undergoes a large, localized shift on a deep branch. Under this condition, all configurations –

except those relying on EV_X_2 – yield nearly identical results. The overlapping lines illustrate

this convergence among EV_U, EV_X_1, and residual approaches, likely reflecting the fact that

they each capture a critical subset of eigenvectors that effectively model the key phylogenetic

signal introduced by the shift in X_1.

Across scenarios where both ESRRV and mMorI produced admissible EV counts, we

retained mMorI as the default selection method, owing to its consistent superiority. When

considered together with results from Fig. 1 and Fig. 2, our findings suggest that mMorI is

preferable for balanced trees with $\rho = 1$ or $\rho = 2$, while ESRRV should be used for $\rho = 0.1$

balanced trees and $\rho = 1$ ladder-shaped trees, where mMorI often fails to return sufficient

EVs.

EV_U-based PVR as an intrinsically robust framework

We evaluated five regression estimators – L2 (ordinary least squares), L1 (least absolute

deviations), S (high-breakdown S-estimator), M (Huber-type M-estimator), and MM (S-start

followed by an M-step) – within the EV_U-based PVR framework. Throughout this section,

L2-based PVR is used as the baseline for comparison. To assess robustness across

evolutionary contexts, we focused on the combinations of tree topology and EV-selection

method identified in the preceding analyses. Specifically, this analysis included 128-species

trees with four phylogenetic topologies: balanced trees with $\rho = 1$ and $\rho = 2$ analyzed under

mMorI, and balanced trees with $\rho = 0.1$ and ladder-shaped trees with $\rho = 1$ analyzed under

ESRRV (Figs. S3–S6). In addition, 16-species balanced trees with $\rho = 1$ were analyzed under

mMorI (Fig. S7). Across these settings, L2, M, and MM yielded nearly identical accuracies in

most scenarios, with their curves largely overlapping in the plots. Only in ladder-shaped trees

under strong shift conditions did MM show a clear robustness advantage, while L2 and M

remained indistinguishable. Compared with these three estimators, the performance of L1

varied depending on the scenario; in contrast, S generally underperformed across multiple

shift settings and was thus excluded from subsequent analyses. Estimator M was also omitted

from later figures, as its outputs were nearly indistinguishable from L2 and were almost

entirely overlapping in all scenarios, making its inclusion redundant.

Comparative evaluation of optimized PVR, PGLS, and PIC

To compare optimized PVR with established phylogenetic regression methods, we evaluated multiple PVR configurations alongside PGLS and PIC-based approaches. Here, L2 serves as the standard least-squares baseline, whereas L1 and MM represent increasingly robust alternatives. We used EV_U constructed from mMorI for 128-species balanced trees ($\rho = 1, 2$) and 16-species balanced trees ($\rho = 1$), and from ESRRV for 128-species balanced trees ($\rho = 0.1$) and ladder-shaped trees ($\rho = 1$). These approaches were compared against three established methods, PGLS, PIC-L2 (i.e., PIC-OLS), and PIC-MM (a robust variant recommended by Adams et al. (2024). For PGLS, the best-fitting evolutionary model was selected from five candidates – BM, Pagel's λ, Ornstein–Uhlenbeck (OU) with fixed α, OU with random α, and the Early-Burst (EB) model – using AIC and is hereafter referred to as PGLS-optimal.

Across all phylogenetic structures examined, the four gradual-evolution scenarios (BM–BM, BM–Norm, Norm–BM, and Norm–Norm) produced broadly similar accuracy patterns among most methods (Figs. 5a–d; Figs. S8–S11a–d). In these settings, the various PVR variants generally performed slightly below – or occasionally comparable to – PGLS-optimal and PIC-L2, and often showed accuracy similar to PIC-MM. Under the BM–Norm condition, however, all methods except PGLS-optimal performed similarly, with most showing somewhat higher accuracy than PGLS-optimal. These similarities across figures reflect that, under gradual or near-gradual evolutionary dynamics, the compared phylogenetic regression approaches capture the major components of phylogenetic structure to a comparable degree.

In contrast, the shift scenarios revealed clearer performance differences (Figs. 5e–f; Figs. S8–S11e–f). Across all tree shapes – whether 128-species or 16-species, and across balanced trees with varying values of ρ – PGLS-optimal and PIC-L2 consistently showed poor robustness when shift magnitudes increased. By comparison, PIC-MM and the PVR variants remained markedly more robust to strong evolutionary shifts, with even the standard PVR-L2 regression showing substantial robustness under most tree shapes (Figs. S8–S11e–f). On ladder-shaped trees (Figs. 5e-f), however, the robustness of PVR-L1 and PVR-L2 deteriorated, yielding performance closer to that of PGLS-optimal and PIC-L2. In this setting, only the MM-based approaches – PIC-MM and PVR-MM – sustained high accuracy and effectively accommodated large shifts.

Fig. 5. Performance of PVR, PGLS, and PIC on 128-species ladder-shaped phylogenetic trees ($\rho = 1$).
Comparisons were conducted under six evolutionary scenarios: (a–d) the four gradual-evolution (stationary)
scenarios – BM–BM, BM–Norm, Norm–BM, and Norm–Norm – and (e–f) the two non-stationary single-branch
shift scenarios – X_1-only shift and dual-trait shift. All eigenvectors (EVs) were selected using the ESRRV
criterion. Accuracy is quantified using a replicate-level benchmark derived from branchwise standardized
changes. For details on ESRRV, scenario definitions, and accuracy measures, see Materials and Methods. The
data underlying this figure are provided in Table S6.

Discussion

PVR, first proposed to model trait variation and trait–trait correlations using phylogenetic

eigenvectors (Diniz-Filho et al., 1998), is widely used across ecological studies. Its strength

lies in capturing separable, orthogonal eigenvectors, without requiring an explicit

evolutionary model. This flexibility allows easy incorporation into machine learning methods

such as random forests and neural networks (Debastiani et al., 2021; Penone et al., 2014),

supporting tasks like trait imputation and classification beyond regression. Moreover, because

the eigenvectors are orthogonal, they can be jointly modeled with other sources of

autocorrelation (e.g., spatial or temporal), a flexibility particularly valuable in ecological

settings where multiple forms of structure coexist (Kühn et al., 2009; Maynard et al., 2022;

Safi and Pettorelli, 2010; Wang et al., 2019).

Methodological improvements to PVR

In this study, we revisit and optimize the PVR framework. We show that eigenvectors derived from different traits often diverge, and using their union (EV_U) more effectively controls phylogenetic non-independence. While our focus is on trait pairs, the logic extends to multi-trait settings, where incorporating all relevant eigenvectors selected for each trait helps retain phylogenetic signal. However, care must be taken to avoid overfitting by limiting the number of retained eigenvectors.

The "SPV ≥ 10" rule is widely recommended in conventional regression to guard against overfitting and unstable inference (Austin et al., 2017; Babyak, 2004; Peduzzi et al., 1996). Because eigenvectors are orthogonal rather than correlated predictors, uncritically importing such heuristics into phylogenetic contexts risks misrepresenting model reliability. Our analyses indicate that PVR remains reliable with much lower SPV, under gradual evolution, with accuracy still acceptable when SPV falls to ~3, whereas under shift conditions PVR performance shows virtually no such lower bound. This tolerance affords greater flexibility, crucial for small datasets: with ≤19 species, SPV ≥ 10 precludes eigenvectors entirely, whereas a relaxed threshold of SPV ≥ 3 allows inclusion of up to four eigenvectors. Our analyses further suggest a practical guideline: regardless of the selection method, researchers should inspect the number of retained eigenvectors before incorporating them into regression models. For example, mAIC typically returns excessively large sets that risk overfitting, whereas mMorI may frequently yield very few or even zero eigenvectors.

PVR under non-stationary and shift-like evolutionary dynamics

Unlike likelihood-based methods such as PGLS, whose performance is inherently tied to a single evolutionary model (Freckleton et al., 2011), PVR extracts phylogenetic structure through eigenvector decomposition and thus remains flexible when traits deviate from standard processes (e.g., BM, OU, EB) or when model uncertainty is high. Because phylogenetic eigenvectors are derived solely from the topology and branch-length structure of the tree, they capture variation expressed at both broad and fine phylogenetic scales. This property makes them well suited for describing complex empirical patterns – such as clade-specific differences in trait lability, heterogeneous evolutionary rates, or multiple adaptive regimes – that cannot be adequately summarized by a single global parameter (Diniz-Filho et al., 2015; Diniz-Filho et al., 2014; Diniz-Filho et al., 2019; Diniz-Filho et al., 2012b; Diniz-Filho et al., 2010; Gouveia et al., 2013). Within this perspective, the abrupt evolutionary shifts examined in our simulations represent an extreme form of non-stationarity: substantial localized departures from Brownian covariance that arise when evolutionary rates or selective

regimes change along particular branches of the tree. With appropriately selected EV sets

(e.g., union-based selection guided by mMorI or ESRRV), EVs can capture this structure,

yielding near-i.i.d. residuals. Because M-estimator reweighting is primarily triggered by

influential outliers (Huber, 1964), it is rarely activated under such conditions; consequently,

default PVR-L2 (OLS) typically matches PVR-M in robustness, outperforming PGLS or

typical PIC. Under extreme heterogeneity (e.g., strongly ladderized trees with multiple sharp

shifts), PVR-MM affords added stability. Rather than overturning prior assessments

characterizing PVR as inferior to likelihood-based approaches (Freckleton et al., 2011;

Laurin, 2010; Pennell & Harmon, 2013), our results build upon earlier demonstrations that

PVR's performance depends critically on how eigenvectors are selected, and show that

systematic selection strategies can substantially improve stability and accuracy under non-

stationary evolutionary conditions (Diniz-Filho et al., 2015; Diniz-Filho et al., 2012a; Diniz-

Filho et al., 2010).

PGLS under signal imbalance and unmodeled phylogenetic structure

Our analyses reveal that the performance of PGLS is strongly shaped by how phylogenetic

signal is distributed between the two traits. In Fig. 5 and Figs. S8–S11, PGLS-optimal

performed worst in the BM–Norm condition, where the dependent variable (X_2) carried

markedly weaker phylogenetic signal than the predictor (X_1). In contrast, PGLS achieved the

best or near-best performance in the other stationary conditions, in which X_2 carried

substantially stronger or at least comparable signal. This pattern is consistent with the fact that

PGLS inherits from OLS the assumption that predictors are fixed and measured without error

(Casson & Farmer, 2014; Jarantow et al., 2023; McArdle, 1988; Osborne & Waters, 2002),

creating a mismatch when the predictor carries the stronger phylogenetic signal. Preliminary

analyses by Chen et al. (2023, preprint) similarly show that such signal imbalance can distort

regression estimates, underscoring the importance of accounting for the distribution of

phylogenetic signal when selecting the response variable.

Adding phylogenetic eigenvectors to PGLS – a framework recently explored by

Schraiber et al. (2024) could, in principle, also mitigate problems arising when the predictor

carries substantially stronger phylogenetic signal than the response, as in the BM–Norm

scenario. Although Schraiber et al. did not study this mechanism, their empirical results

highlight a related phenomenon. In fungal gene-expression data, including eigenvectors in

PGLS markedly reduced spurious correlations, revealing unmodeled phylogenetic structure

that is unlikely to reflect differences in gradual signal strength between paired genes. Instead,

their results point to abrupt, localized deviations from Brownian expectations – a shift-like

form of non-stationarity closely paralleling the abrupt dual-shifts examined in our

simulations. These findings illustrate a broader function of eigenvectors: they can capture

residual phylogenetic structure not encoded in the assumed covariance matrix, whether arising

from gradual imbalance or from localized evolutionary shifts. Within this broader context,

determining which eigenvectors should be added to PGLS remains an open question. A

natural possibility – yet to be formally evaluated – is that including only those eigenvectors

present in the predictor but absent in the response may provide a better complement to PGLS

than using EV_U, or perhaps even using the full set of predictor-selected eigenvectors,

including those that overlap with the response. Because PGLS already accounts for the

phylogenetic structure of the response through its covariance model, re-introducing

eigenvectors that primarily reflect that same structure could risk unnecessary complexity or

overfitting. Exploring whether predictor-specific eigenvectors yield more stable inference

therefore represents a promising direction for future work.

Extending asymmetry considerations to spatial autocorrelation models

Although our focus is on phylogenetic comparative methods, similar issues arise in spatial

analyses, where spatial autocorrelation must also be accounted for (Cottenie, 2005; Dale &

Fortin, 2009; Diniz-Filho & Bini, 2005; He et al., 2003; Keitt et al., 2002; Legendre, 1993;

Peres-Neto & Legendre, 2010). Preliminary tests show that the asymmetry introduced by

variable designation – discussed for PVR – also affects spatial frameworks such as spatial

eigenvector mapping, CAR, SAR, and trend-surface GAMs, with performance differing when

dependent and independent variables are swapped (Table S7; see Spatial_methods in

Supplementary material for implementation details). By analogy, the union-based strategy we

adopted for PVR may also apply to spatial analyses, potentially helping to preserve shared

spatial signal while controlling autocorrelation. In addition, our findings regarding effective

sample sizes – particularly the observation that reliable performance can be achieved with

substantially lower SPV than conventionally assumed – may also be informative for spatial

models, where limited sample sizes and large sets of spatial eigenvectors pose similar

challenges.

Concluding remarks: Practical guidelines for selecting eigenvectors in PVR

Based on our theoretical derivations and extensive simulations across evolutionary scenarios and tree structures, we recommend the following practical guidelines for selecting eigenvectors in PVR:

1. **Use the union of eigenvectors selected separately for each trait as the default.** This strategy consistently outperforms per-trait selections by controlling phylogenetic structure in both variables and preventing asymmetric corrections.

2. **Prefer mMorI when it does not return too few eigenvectors; otherwise use ESRRV.** mMorI typically provides a compact set of eigenvectors by minimizing residual Moran's I, ensuring strong phylogenetic correction. However, when mMorI selects very few – or even zero – eigenvectors, ESRRV offers a more reliable alternative by providing a more adequate number of components.

3. **Avoid mAIC for regression-based PVR**, because the resulting EV sets are typically excessively large, risking overfitting without improving phylogenetic control.

4. **Inspect EV counts before model fitting.** For datasets with $n \leq 19$, SPV ≥ 10 would prohibit the use of eigenvectors entirely. However, our simulations show that PVR remains reliable with SPV ≈ 3, while shift-like non-stationarity allows an even more relaxed SPV threshold.

5. **Prefer PVR under non-stationary or shift-like evolutionary conditions.** Under these scenarios, PVR maintains strong performance even without robust estimation. However, under more extreme heterogeneity – such as strongly ladderized trees with sharp shifts – using a robust MM-estimator (PVR-MM) can provide additional stability.

Supplementary material

Supplementary material is submitted with this manuscript as online Supporting Information.

Data availability

The scripts used to generate the results in the main text—including phylogenetic tree construction, trait data simulation, regression model fitting, and statistical analysis—are provided in Main_code.zip. The code used to evaluate spatial statistical methods is provided in Spatial_methods.zip. In addition, we developed an R package, ROBPVRU, which implements phylogenetic eigenvector regression with union operations on eigenvectors combined with robust estimators. All code and data used to generate Fig. 1 and the

supplementary figures have been uploaded in a review-only format to preserve author
anonymity during peer review and will be made freely accessible via the first author's GitHub
page upon acceptance.

References

Adams, D. C., & Collyer, M. L. (2022). Consilience of methods for phylogenetic analysis of variance. *Evolution, 76*(7), 1406-1419. https://doi.org/10.1111/evo.14512

Adams, R., Cain, Z., Assis, R., & DeGiorgio, M. (2024). Robust phylogenetic regression. *Systematic Biology, 73*(1), 140-157. https://doi.org/10.1093/sysbio/syad070

Álamo, M., García-Barros, E., & Romo, H. (2024). Adult thermoregulatory behaviour does not provide, by itself, an adaptive explanation for the reflectance–climate relationship (Bogert's pattern) in Iberian butterflies. *Ecological Entomology, 49*(1), 77-90. https://doi.org/https://doi.org/10.1111/een.13281

Austin, P. C., Allignol, A., & Fine, J. P. (2017). The number of primary events per variable affects estimation of the subdistribution hazard competing risks model. *Journal of Clinical Epidemiology, 83*, 75-84. https://doi.org/https://doi.org/10.1016/j.jclinepi.2016.11.017

Babyak, M. A. (2004). What you see may not be what you get: a brief, nontechnical introduction to overfitting in regression-type models. *Psychosomatic Medicine, 66*(3), 411-421. https://doi.org/10.1097/01.psy.0000127692.23278.a9

Carotenuto, F., Diniz-Filho, J. A. F., & Raia, P. (2015). Space and time: The two dimensions of Artiodactyla body mass evolution. *Palaeogeography, Palaeoclimatology, Palaeoecology, 437*, 18-25. https://doi.org/https://doi.org/10.1016/j.palaeo.2015.07.013

Casson, R. J., & Farmer, L. D. (2014). Understanding and checking the assumptions of linear regression: a primer for medical researchers. *Clinical & Experimental Ophthalmology, 42*(6), 590-596. https://doi.org/https://doi.org/10.1111/ceo.12358

Chen, Z.-L., Guo, H.-J., & Niu, D.-K. (2023). Dependent variable selection in phylogenetic generalized least squares regression analysis under Pagel's lambda model. *bioRxiv*, 2023.2005.2021.541623. https://doi.org/10.1101/2023.05.21.541623

Chen, Z.-L., Huang, R., Guo, H.-J., & Niu, D.-K. (2025). Improving the robustness of

phylogenetic independent contrasts: addressing abrupt evolutionary shifts with outlier-

and distribution-guided correlation. *bioRxiv*, 2024.2006.2016.599156.

https://doi.org/10.1101/2024.06.16.599156

Cheverud, J. M., Dow, M. M., & Leutenegger, W. (1985). The quantitative assessment of

phylogenetic constraints in comparative analyses: sexual dimorphism in body weight

among primates. *Evolution, 39*(6), 1335-1351. https://doi.org/10.1111/j.1558-

5646.1985.tb05699.x

Cornwallis, C. K., & Griffin, A. S. (2024). A guided tour of phylogenetic comparative

methods for studying trait evolution. *Annual Review of Ecology, Evolution, and*

Systematics, 55, 181-204. https://doi.org/10.1146/annurev-ecolsys-102221-050754

Cottenie, K. (2005). Integrating environmental and spatial processes in ecological community

dynamics. *Ecology Letters, 8*(11), 1175-1182. https://doi.org/10.1111/j.1461-

0248.2005.00820.x

Dale, M. R. T., & Fortin, M.-J. (2009). Spatial autocorrelation and statistical tests: Some

solutions. *Journal of Agricultural, Biological, and Environmental Statistics, 14*(2), 188-

206. https://doi.org/10.1198/jabes.2009.0012

Debastiani, V. J., Bastazini, V. A. G., & Pillar, V. D. (2021). Using phylogenetic information

to impute missing functional trait values in ecological databases. *Ecological*

Informatics, 63, 101315. https://doi.org/10.1016/j.ecoinf.2021.101315

Dewar, A. E., Belcher, L. J., & West, S. A. (2025). A phylogenetic approach to comparative

genomics. *Nature Reviews Genetics, 26*, 395-405. https://doi.org/10.1038/s41576-024-

00803-0

Diniz-Filho, J. A. F., Alves, D. M. C. C., Villalobos, F., Sakamoto, M., Brusatte, S. L., &

Bini, L. M. (2015). Phylogenetic eigenvectors and nonstationarity in the evolution of

theropod dinosaur skulls. *Journal of Evolutionary Biology, 28*(7), 1410-1416.

https://doi.org/10.1111/jeb.12660

Diniz-Filho, J. A. F., & Bini, L. M. (2005). Modelling geographical patterns in species

richness using eigenvector-based spatial filters. *Global Ecology and Biogeography,*

14(2), 177-185. https://doi.org/10.1111/j.1466-822X.2005.00147.x

Diniz-Filho, J. A. F., Bini, L. M., Rangel, T. F., Morales-Castilla, I., Olalla-Tárraga, M. Á.,

Rodríguez, M. Á., & Hawkins, B. A. (2012a). On the selection of phylogenetic

eigenvectors for ecological analyses. *Ecography 35*(3), 239-249.

https://doi.org/10.1111/j.1600-0587.2011.06949.x

Diniz-Filho, J. A. F., Bini, M., Sakamoto, M., & Brusatte, S. (2014). Phylogenetic

eigenvector regression in paleobiology. *Revista Brasileira de Paleontologia, 17*(2), 107-

122. https://doi.org/10.4072/rbp.2014.2.01

Diniz-Filho, J. A. F., Cianciaruso, M. V., Rangel, T. F., & Bini, L. M. (2011). Eigenvector

estimation of phylogenetic and functional diversity. *Functional Ecology, 25*(4), 735-

744. https://doi.org/10.1111/j.1365-2435.2011.01836.x

Diniz-Filho, J. A. F., de Sant'Ana, C. E. R., & Bini, L. M. (1998). An eigenvector method for

estimating phylogenetic inertia. *Evolution, 52*(5), 1247-1262.

https://doi.org/10.1111/j.1558-5646.1998.tb02006.x

Diniz-Filho, J. A. F., Jardim, L., Mondanaro, A., & Raia, P. (2019). Multiple components of

phylogenetic non-stationarity in the evolution of brain size in fossil Hominins.

Evolutionary Biology, 46(1), 47-59. https://doi.org/10.1007/s11692-019-09471-z

Diniz-Filho, J. A. F., & Nabout, J. C. (2009). Modeling body size evolution in Felidae under

alternative phylogenetic hypotheses. *Genet Mol Biol, 32*(1), 170-176.

https://doi.org/10.1590/s1415-47572009005000004

Diniz-Filho, J. A. F., Rangel, T. F., Santos, T., & Mauricio Bini, L. (2012b). Exploring

patterns of interspecific variation in quantitative traits using sequential phylogenetic

eigenvector regressions. *Evolution, 66*(4), 1079-1090. https://doi.org/10.1111/j.1558-

5646.2011.01499.x

Diniz-Filho, J. A. F., Terribile, L. C., da Cruz, M. J. R., & Vieira, L. C. G. (2010). Hidden

patterns of phylogenetic non-stationarity overwhelm comparative analyses of niche

conservatism and divergence. *Global Ecology and Biogeography, 19*(6), 916-926.

https://doi.org/https://doi.org/10.1111/j.1466-8238.2010.00562.x

Diniz-Filho, J. A. F., & Tôrres, N. M. (2002). Phylogenetic comparative methods and the

geographic range size – body size relationship in new world terrestrial carnivora.

Evolutionary Ecology, 16(4), 351-367. https://doi.org/10.1023/A:1020210321776

Divíšek, J., Chytrý, M., Beckage, B., Gotelli, N. J., Lososová, Z., Pyšek, P., Richardson, D.

M., & Molofsky, J. (2018). Similarity of introduced plant species to native ones

facilitates naturalization, but differences enhance invasion success. *Nature*

Communications, 9(1), 4631. https://doi.org/10.1038/s41467-018-06995-4

Felsenstein, J. (1985). Phylogenies and the comparative method. *The American Naturalist,*

125(1), 1-15. https://doi.org/10.1086/284325

Freckleton, R. P., Cooper, N., & Jetz, W. (2011). Comparative methods as a statistical fix: the

dangers of ignoring an evolutionary model. *American Naturalist, 178*(1), E10-E17.

https://doi.org/10.1086/660272

Gao, X., Dai, J., Tao, Z., Shahzad, K., & Wang, H. (2023). Autumn phenology of tree species

in China is associated more with climate than with spring phenology and phylogeny.

Frontiers in Plant Science, 14, 1040758. https://doi.org/10.3389/fpls.2023.1040758

Garamszegi, L. Z. (Ed.) (2014). *Modern Phylogenetic Comparative Methods and Their*

Application in Evolutionary Biology: Concepts and Practice. Berlin: Springer.

Gervini, D., & Yohai, V. J. (2002). A class of robust and fully efficient regression estimators.

The Annals of Statistics, 30(2), 583-616. https://doi.org/10.1214/aos/1021379866

Gouveia, S. F., Hortal, J., Cassemiro, F. A. S., Rangel, T. F., & Diniz-Filho, J. A. F. (2013).

Nonstationary effects of productivity, seasonality, and historical climate changes on

global amphibian diversity. *Ecography, 36*(1), 104-113. https://doi.org/10.1111/j.1600-

0587.2012.07553.x

Grafen, A. (1989). The phylogenetic regression. *Philosophical Transactions of the Royal*

Society of London. Series B, Biological Sciences, 326(1233), 119-157.

https://doi.org/10.1098/rstb.1989.0106

He, F., Zhou, J., & Zhu, H. (2003). Autologistic regression model for the distribution of

vegetation. *Journal of Agricultural, Biological, and Environmental Statistics, 8*(2), 205-

222. https://doi.org/10.1198/1085711031508

Huber, P. J. (1964). Robust estimation of a location parameter. *The Annals of Mathematical*

Statistics, 35(1), 73-101.

Jarantow, S. W., Pisors, E. D., & Chiu, M. L. (2023). Introduction to the use of linear and

nonlinear regression analysis in quantitative biological assays. *Current Protocols, 3*(6),

e801. https://doi.org/10.1002/cpz1.801

Keitt, T. H., Bjørnstad, O. N., Dixon, P. M., & Citron-Pousty, S. (2002). Accounting for

spatial pattern when modeling organism-environment interactions. *Ecography, 25*(5),

616-625. https://doi.org/10.1034/j.1600-0587.2002.250509.x

Kühn, I., Nobis, M. P., & Durka, W. (2009). Combining spatial and phylogenetic eigenvector

filtering in trait analysis. *Global Ecology and Biogeography, 18*(6), 745-758.

https://doi.org/https://doi.org/10.1111/j.1466-8238.2009.00481.x

Lanuza, J. B., Rader, R., Stavert, J., Kendall, L. K., Saunders, M. E., & Bartomeus, I. (2023).

Covariation among reproductive traits in flowering plants shapes their interactions with

pollinators. *Functional Ecology, 37*(7), 2072-2084. https://doi.org/10.1111/1365-

2435.14340

Laurin, M. (2010). Assessment of the relative merits of a few methods to detect evolutionary

trends. *Systematic Biology, 59*(6), 689-704. https://doi.org/10.1093/sysbio/syq059

Legendre, P. (1993). Spatial autocorrelation: trouble or new paradigm? *Ecology, 74*(6), 1659-

1673. https://doi.org/10.2307/1939924

LeRoy, C. J., Hipp, A. L., Lueders, K., Follstad Shah, J. J., Kominoski, J. S., Ardón, M.,

Dodds, W. K., Gessner, M. O., Griffiths, N. A., Lecerf, A., Manning, D. W. P.,

Sinsabaugh, R. L., & Webster, J. R. (2020). Plant phylogenetic history explains in-

stream decomposition at a global scale. *Journal of Ecology, 108*(1), 17-35.

https://doi.org/10.1111/1365-2745.13262

Liu, Y., Dunker, S., Durka, W., Dominik, C., Heuschele, J. M., Honchar, H., Hoffmann, P.,

Musche, M., Paxton, R. J., Settele, J., & Schweiger, O. (2024). Eco-evolutionary

processes shaping floral nectar sugar composition. *Scientific Reports, 14*(1), 13856.

https://doi.org/10.1038/s41598-024-64755-5

Maynard, D. S., Bialic-Murphy, L., Zohner, C. M., Averill, C., van den Hoogen, J., Ma, H.,

Mo, L., Smith, G. R., Acosta, A. T. R., Aubin, I., Berenguer, E., Boonman, C. C. F.,

Catford, J. A., Cerabolini, B. E. L., Dias, A. S., González-Melo, A., Hietz, P., Lusk, C.

H., Mori, A. S., . . . Crowther, T. W. (2022). Global relationships in tree functional

traits. *Nature Communications, 13*(1), 3185. https://doi.org/10.1038/s41467-022-30888-

2

McArdle, B. H. (1988). The structural relationship: regression in biology. *Canadian Journal*

of Zoology, 66(11), 2329-2339. https://doi.org/10.1139/z88-348

Osborne, J. W., & Waters, E. (2002). Four assumptions of multiple regression that researchers

should always test. *Practical Assessment, Research, and Evaluation, 8*(1), 2.

https://doi.org/10.7275/r222-hv23

Paradis, E., & Schliep, K. (2019). ape 5.0: an environment for modern phylogenetics and

evolutionary analyses in R. *Bioinformatics, 35*(3), 526-528.

https://doi.org/10.1093/bioinformatics/bty633

Peduzzi, P., Concato, J., Kemper, E., Holford, T. R., & Feinstein, A. R. (1996). A simulation

study of the number of events per variable in logistic regression analysis. *Journal of*

Clinical Epidemiology, 49(12), 1373-1379.

https://doi.org/https://doi.org/10.1016/S0895-4356(96)00236-3

Pennell, M. W., & Harmon, L. J. (2013). An integrative view of phylogenetic comparative

methods: connections to population genetics, community ecology, and paleobiology.

Annals of the New York Academy of Sciences, 1289(1), 90-105.

https://doi.org/https://doi.org/10.1111/nyas.12157

Penone, C., Davidson, A. D., Shoemaker, K. T., Di Marco, M., Rondinini, C., Brooks, T. M.,

Young, B. E., Graham, C. H., & Costa, G. C. (2014). Imputation of missing data in life-

history trait datasets: which approach performs the best? *Methods in Ecology and*

Evolution, 5(9), 961-970. https://doi.org/10.1111/2041-210X.12232

Peres-Neto, P. R., & Legendre, P. (2010). Estimating and controlling for spatial structure in

the study of ecological communities. *Global Ecology and Biogeography, 19*(2), 174-

184. https://doi.org/10.1111/j.1466-8238.2009.00506.x

Peres-Neto, P. R., Leibold, M. A., & Dray, S. (2012). Assessing the effects of spatial

contingency and environmental filtering on metacommunity phylogenetics. *Ecology,*

93(sp8), S14-S30. https://doi.org/10.1890/11-0494.1

Phillips, R. F. (2002). Least absolute deviations estimation via the EM algorithm. *Statistics*

and Computing, 12(3), 281-285. https://doi.org/10.1023/A:1020759012226

Revell, L. J. (2010). Phylogenetic signal and linear regression on species data. *Methods in*

Ecology and Evolution, 1(4), 319-329. https://doi.org/10.1111/j.2041-

210X.2010.00044.x

Revell, L. J. (2024). phytools 2.0: an updated R ecosystem for phylogenetic comparative

methods (and other things). *PeerJ, 12*, e16505. https://doi.org/10.7717/peerj.16505

Safi, K., & Pettorelli, N. (2010). Phylogenetic, spatial and environmental components of

extinction risk in carnivores. *Global Ecology and Biogeography, 19*(3), 352-362.

https://doi.org/10.1111/j.1466-8238.2010.00523.x

Schraiber, J. G., Edge, M. D., & Pennell, M. (2024). Unifying approaches from statistical

genetics and phylogenetics for mapping phenotypes in structured populations. *Plos*

Biology, 22(10), e3002847. https://doi.org/10.1371/journal.pbio.3002847

Seger, G. D. d. S., Hartz, S. M., & Duarte, L. d. S. (2025). Contrasting Phylogenetic Signals

in Flowering and Fruiting Phenology at Species and Assemblage Scales in a Climbing

Plant Community. *Journal of Vegetation Science, 36*(6), e70087.

https://doi.org/https://doi.org/10.1111/jvs.70087

Slater, G. J., & Pennell, M. W. (2014). Robust regression and posterior predictive simulation

increase power to detect early bursts of trait evolution. *Systematic Biology, 63*(3), 293-

308. https://doi.org/10.1093/sysbio/syt066

Venables, W. N., & Ripley, B. D. (2002). *Modern Applied Statistics with S* (4th ed.):

Springer.

Vilela, B., Villalobos, F., Rodríguez, M. Á., & Terribile, L. C. (2014). Body Size, Extinction

Risk and Knowledge Bias in New World Snakes. *PLoS ONE, 9*(11), e113429.

https://doi.org/10.1371/journal.pone.0113429

Wang, Z. H., Jiang, Y., Deane, D. C., He, F. L., Shu, W. S., & Liu, Y. (2019). Effects of host

phylogeny, habitat and spatial proximity on host specificity and diversity of pathogenic

and mycorrhizal fungi in a subtropical forest. *New Phytologist, 223*(1), 462-474.

https://doi.org/10.1111/nph.15786

Westoby, M., Yates, L., Holland, B., & Halliwell, B. (2023). Phylogenetically conservative

trait correlation: Quantification and interpretation. *Journal of Ecology, 111*(10), 2105-

2117. https://doi.org/10.1111/1365-2745.14150

Yu, C., & Yao, W. (2017). Robust linear regression: A review and comparison.

Communications in Statistics - Simulation and Computation, 46(8), 6261-6282.

https://doi.org/10.1080/03610918.2016.1202271