Optimized Phylogenetic Eigenvector Regression Provides Robust and Flexible Trait Correlation Analysis under Diverse Evolutionary Conditions

Zheng-Lin Chen and Deng-Ke Niu[*]

MOE Key Laboratory for Biodiversity Science and Ecological Engineering and Beijing Key Laboratory of Gene Resource and Molecular Development, College of Life Sciences, Beijing Normal University, Beijing, 100875, China

*Correspondence author:

Deng-Ke Niu, College of Life Sciences, Beijing Normal University, Beijing, 100875, China.

dkniu@bnu.edu.cn, dengkeniu@hotmail.com

ABSTRACT

Phylogenetic comparative methods are essential tools for analyzing trait correlations among

species while controlling for evolutionary dependencies. Phylogenetic eigenvector

regression (PVR) uses eigenvectors (EVs) derived from the phylogenetic tree as covariates

to remove shared evolutionary history. Although conceptually simple and independent of

predefined evolutionary models, PVR has traditionally been viewed as inferior to

phylogenetic generalized least squares (PGLS), which explicitly incorporates evolutionary

models. Here, we systematically optimized the PVR framework by integrating robust

statistical estimators (L1, L2, M, and MM) and developing a theoretically justified approach

to selecting EVs, particularly by using the union of EVs from both traits rather than a single

trait alone. Using extensive simulations, we compared optimized PVR methods to

established approaches, including PGLS and phylogenetic independent contrasts (PIC),

across a range of evolutionary scenarios, spanning strong, weak, and asymmetric

phylogenetic signals as well as non-stationary shifts; and across multiple tree structures

(balanced and ladder-shaped), branch-length scalings ($\rho = 0.1, 1, 2$), and sample sizes (128

and 16 species). Our results challenge the prevailing assumption that PGLS universally

outperforms PVR. While PGLS and standard PIC generally excelled when dependent traits

exhibited strong phylogenetic signals, optimized PVR methods—particularly PVR-MM—

demonstrated superior robustness under scenarios involving mixed or weak phylogenetic

signals, as well as abrupt, non-stationary evolutionary shifts. Even the simplest optimized

PVR variant (PVR-L2) exhibited inherent robustness, highlighting a fundamental advantage

of phylogenetic eigenvectors in controlling evolutionary structure. Furthermore, PVR

provides greater methodological flexibility compared to model-based methods like PGLS, as

it is not constrained by predefined evolutionary models and can readily extend to advanced

machine-learning approaches. Lastly, unlike PGLS, PVR effectively addresses the

directional asymmetry inherited from ordinary least squares regression, allowing researchers

to select predictors based on biological reasoning or measurement accuracy without

compromising model validity. Overall, this study highlights that optimized PVR is not

merely a competitive alternative to traditional methods, but also a flexible and robust

framework that adapts well to a wide range of evolutionary scenarios and analytical needs.

KEYWORDS: phylogenetic comparative methods; phylogenetic eigenvector regression

(PVR); robust regression; evolutionary shift; phylogenetic autocorrelation; phylogenetic

signal; dependent and independent variable asymmetry; trait correlation.

INTRODUCTION

Phylogenetic comparative methods correct for non-independence among species due to

shared ancestry (Felsenstein 1985, Garamszegi 2014, Cornwallis and Griffin 2024, Dewar et

al. 2025). Among these, phylogenetic generalized least squares (PGLS) is widely regarded as

the standard method, modeling trait covariation using predefined evolutionary models such

as Brownian motion (BM) or Ornstein-Uhlenbeck (OU) (Grafen 1989, Pagel 1999,

Garamszegi 2014). By contrast, phylogenetic eigenvector regression (PVR) uses

phylogenetic eigenvectors (EVs)—orthogonal axes extracted from the phylogenetic tree—to

statistically isolate trait correlations from phylogenetic structure (Diniz-Filho et al. 1998,

Diniz-Filho and Tôrres 2002, Diniz-Filho et al. 2014). These EVs represent species'

positions on the phylogenetic tree and can be incorporated as covariates into models to

account for phylogenetic relationships, allowing for partial regression between traits while

controlling for shared phylogenetic history (Desdevises et al. 2003, Safi and Pettorelli 2010,

Diniz-Filho et al. 2011, Peres-Neto et al. 2012).

Despite its intuitive appeal and flexibility, PVR has been considered less reliable than

PGLS in controlling phylogenetic non-independence (Pennell and Harmon 2013). This

conventional view—articulated in foundational comparative studies (Laurin 2010,

Freckleton et al. 2011)—suggests that PVR lacks the explicit evolutionary modeling

necessary for producing robust estimates, and may suffer from inflated Type I error rates

particularly when broken-stick criterion was used to select EVs. Moreover, PVR has been

viewed as a purely descriptive tool rather than one suitable for detecting adaptation or

making evolutionary inferences (Hansen 2014). As a result, PVR has been underutilized,

despite its potential for broader applications.

A key challenge to PVR lies in the selection of EVs. Including all EVs as covariates

may lead to overfitting, where the model excessively attributes variation in the dependent

variable to phylogenetic effects, thus obscuring meaningful trait correlations (Rohlf 2001).

On the other hand, excluding relevant EVs can result in the loss of crucial phylogenetic

information. Fortunately, researches have demonstrated that not all EVs are necessary for

robust analysis, as many contribute minimally to the overall model. By carefully selecting a

subset of EVs, it is often possible to capture sufficient phylogenetic information and avoid

overfitting (Peres-Neto et al. 2005, Griffith and Peres-Neto 2006, Diniz-Filho et al. 2009,

Diniz-Filho et al. 2012a). These selected EVs help to isolate the effects of phylogeny from

other sources of variation, ultimately providing a clearer understanding of trait relationships

that are independent of phylogenetic influence (Diniz-Filho and Nabout 2009, Diniz-Filho et

al. 2009, Burns et al. 2015, Divíšek et al. 2018, Diniz-Filho et al. 2019, LeRoy et al. 2020,

Gao et al. 2023, Lanuza et al. 2023).

Although EVs are derived solely from the phylogenetic tree, the selection of which EVs

to include in a regression model is inherently trait-specific. Different biological traits

emphasize different components of the tree's structure, leading to divergent EV sets

depending on which trait is analyzed. This divergence is not trivial: when EVs are selected

separately based on different traits, subsequent regression analyses may yield conflicting

correlation results. Rather than focusing on distinctions between dependent and independent

variables, which are often ambiguous in biological systems, we addressed this issue by

treating the two traits symmetrically. Inspired by Cheverud's (1985) decomposition of trait

covariation into phylogenetic and specific components ($T = P + S$), we hypothesized that

using the union of EVs selected from both traits (EV_U) would provide a more stable and

comprehensive representation of the phylogenetic structure relevant to either trait.

Subsequent simulation analyses confirmed this hypothesis, showing that EV_U led to more

accurate estimates of trait correlations across a wide range of evolutionary conditions.

In addition to improving EV selection, we explored a second optimization strategy for

PVR: enhancing its robustness to model violations and outliers through the integration of

robust regression estimators. Traditional ordinary least squares (OLS) regression, as used in

standard PVR, assumes normally distributed errors and is sensitive to extreme values.

Robust regression techniques—such as L1, M-, S-, and MM-estimators—have been

developed to mitigate these limitations by down-weighting the influence of outliers (Huber

1964, Yu and Yao 2017). Robust regression estimators have increasingly been adapted for

use in phylogenetic comparative methods (Slater and Pennell 2014, Adams et al. 2024). In

particular, Adams et al. (2024) demonstrated that the MM-estimator applied to phylogenetic

independent contrasts (PIC), PIC-MM, not only outperformed other robust variants of PIC

and PGLS (e.g., PIC-L1, PIC-M, PGLS-L1), but also consistently surpassed standard

implementations such as PIC-L2 and PGLS-L2—especially under non-stationary

evolutionary conditions.

This study revisits and systematically improves the PVR framework through two

methodological enhancements: (1) using EV_U, the full union of eigenvectors selected

separately from X_1 and X_2, including their shared components, to more comprehensively

account for overlapping and trait-specific phylogenetic signals; (2) integrating robust

regression estimators to improve model resilience to outliers and deviations from normality.

These optimized variants are then benchmarked against established methods such as PGLS

and PIC across a wide range of evolutionary conditions. Contrary to the long-standing view

that PGLS consistently outperforms alternative approaches (Laurin 2010, Freckleton et al.

2011, Pennell and Harmon 2013, Hansen 2014), our findings demonstrate that optimized

PVR methods—particularly those incorporating robust estimators—can match or exceed

PGLS in accuracy, primarily under non-stationary or weak signal scenarios. These results

highlight the context-dependent nature of methodological performance and suggest that

comparative researchers should move beyond one-size-fits-all approaches when selecting

analytical tools.

124   MATERIALS AND METHODS

125   *Simulation of Phylogenetic Trees*

We simulated five fully bifurcating phylogenetic trees to evaluate the performance of PVR,

PGLS, and PIC under diverse evolutionary conditions. (1) A 128-species balanced tree with

uniformly scaled branch lengths ($\rho = 1$) was used as the baseline structure. (2) To assess the

effects of branch-length scaling (Grafen 1989), we constructed two additional 128-species

balanced trees: one with compressed ancestral and elongated terminal branches ($\rho = 0.1$),

and one with elongated ancestral and shortened terminal branches ($\rho = 2$). (3) To evaluate

the impact of taxonomic sampling, we included a 16-species balanced tree with uniformly

scaled branch lengths ($\rho = 1$). (4) To examine performance under topological asymmetry, we

simulated a 128-species ladder-shaped (comb-shaped) tree with $\rho = 1$. This tree represents

extreme imbalance, with each bifurcation producing one descendant lineage that does not

split further—modeling prolonged unbranched evolutionary trajectories.

All balanced and ladder-shaped phylogenetic trees were constructed using the *stree*

function from the R package ape (Version 5.8) (Paradis and Schliep 2019) and the *paste.tree*

function from the R package phytools (Version 2.3-0) (Revell 2024).

141   *Simulated Trait Generation*

142 We simulated the evolution of two traits (X_1 and X_2) along each of the five phylogenetic

143 trees, following the simulation logic proposed in Revell (2010) and Adams et al. (2024). In

144 all scenarios, X_2 was modeled as:

145                                    $$X_2 = \beta X_1 + \varepsilon$$

where β is a scalar coefficient controlling the trait association, and ε is an independently

simulated noise term. Unless otherwise noted, β was set to 1 to introduce a correlation

between the traits. The magnitude and structure of ε were varied to simulate different

evolutionary dynamics. We tested eight levels of ε magnitude: 10^{-4}, 0.01, 1, 4, 16, 64, 256,

and 1024. Each setting was repeated 1,000 times per tree, totaling 8,000 simulations per

scenario and 40,000 simulations per tree type.

Five simulation scenarios were defined. (1) **BM1 & BM1+BM2**, where X_1 evolves

under Brownian motion (BM) with unit variance. ε also evolves under a separate BM

process with the variable magnitude. Both X_1 and X_2 thus exhibit strong phylogenetic

autocorrelation. (2) **BM & BM+Norm**, where X_1 evolves under BM (variance = 1). ε is

drawn from an i.i.d. normal distribution with varying magnitude. This setup retains strong

phylogenetic signal in X_1, but the phylogenetic signal in X_2 is diluted by non-phylogenetic

noise. (3) **Norm & Norm+BM**, where X_1 follows a normal distribution (mean = 0, variance

= 1) without phylogenetic structure. ε evolves under BM. X_2 thus acquires some

phylogenetic structure from ε, while X_1 remains unstructured. (4) **Norm1 &**

Norm1+Norm2, where Both X_1 and ε follow i.i.d. normal distributions. Neither trait is

expected to carry phylogenetic signal. This scenario serves as a null benchmark, though

some simulations may exhibit weak incidental phylogenetic structure due to noise. (5) **Shift**,

where β was set to 0 to eliminate inherent correlation between X_1 and X_2. Both X_1 and ε

evolved independently under a BM model with unit variance. To simulate non-stationary

evolutionary change, a synchronous, instantaneous shift was introduced along a single

internal branch near the root. In the ladder-shaped tree, this shift was applied to the branch

leading to the lineage that continues branching. The shift simultaneously displaced the

values of X_1 and ε by a specified magnitude s, which was varied across eight values: 10^{-4},

0.01, 1, 4, 16, 64, 256, and 1024. The case $s = 1$ served as the baseline (no shift), while other

values captured greater degrees of evolutionary displacement without introducing causal

association between traits.

We used the *rTraitCont* function from the *ape* package to generate trait data under the

BM evolutionary model and the *rnorm* function from the *stats* package to generate data

under a normal distribution.

177   ***Phylogenetic Eigenvectors Selection Methods***

Building on Cheverud's (1985) autoregressive framework (T = P + S), where T denotes trait

values, P represents the phylogenetic component, and S accounts for other influences, PVR

applies principal coordinate analysis to a double-centered phylogenetic distance matrix to

derive $n - 1$ EVs, from which a relevant subset is selected as covariates in regression models.

EVs were selected separately for each trait (X_1 or X_2) by fitting the models: $X_1 = EV\_X_1*$

$\beta + \varepsilon$, and $X_2 = EV\_X_2* \beta + \varepsilon$, where β is the regression coefficient vector and ε is the

residual (S component). We adopted three non-sequential EV selection methods, adapted

from references (Diniz-Filho and Nabout 2009, Diniz-Filho et al. 2012a). (1) In the ESRRV

approach, EVs significantly correlated with the dependent variable were identified via

Pearson or Spearman correlations, depending on normality. (2) The mAIC method applied

forward stepwise regression to sequentially add EVs that minimized the Akaike Information

Criterion (AIC), with the selection process terminating when no further improvement could

be achieved. (3) The mMorI approach used forward selection to include EVs until the

residuals minimized Moran's I, a statistic ranging from –1 to +1 that quantifies the strength

of phylogenetic autocorrelation, with 0 indicating no detectable autocorrelation.

To simplify interpretation and improve computational performance, we made a practical

adjustment to the mMorI algorithm. Originally developed to detect spatial autocorrelation,

Moran's I is capable of capturing both positive and negative structure by quantifying the

clustering of similar values. In conventional applications, variable selection continues until

the residuals reach non-significance (e.g., $p > 0.05$), indicating that spatial autocorrelation

has been adequately removed. However, in phylogenetic contexts, trait values among closely

related species tend to be positively correlated—negative phylogenetic autocorrelation is

theoretically possible but biologically rare. Therefore, rather than relying solely on p-values,

we modified the mMorI procedure to terminate the eigenvector selection process once a

negative value of Moran's I is observed in the residuals. If no negative value was reached,

selection proceeded until a non-significant p-value threshold was met.

To accomplish this, we rewrote the relevant code in the PVR package for selecting EVs

by minimizing Moran's I, resulting in a standalone function named *MoransI*. This custom

script is now used for Moran's I method of EV selection in PVR, while all other

computations, such as EV decomposition, still utilize the original *PVR* package (Diniz-Filho

et al. 1998, Diniz-Filho et al. 2012a, Diniz-Filho et al. 2012b). For the ESRRV method, we

created the *ESRRV* function, and for the mAIC method, we employed the *stepAIC* function

from the *MASS* package (Version 7.3-55) (Venables and Ripley 2002).

213       *Accounting for Phylogenetic Autocorrelation in Trait Regression Models*

214   Following EV selection, we implemented four approaches to account for phylogenetic

215   autocorrelation. (1) In the ***EV_X₁-corrected regression***, the EVs selected for trait X_1 were

216   included as covariates in the model $X_1 \sim X_2 + EV\_X_1$. (2) In the ***EV_X₂-corrected***

217   **regression**, the EVs from trait X_2 were used as covariates in the model $X_2 \sim X_1 + EV\_X_2$. It is

218   important to note that the relationship between X_1 and X_2 derived from $X_2 \sim X_1 + EV\_X_2$ is

219   equivalent to that from $X_1 \sim X_2 + EV\_X_2$, since both models adjust for phylogenetic

220   autocorrelation in trait X_2, and the direction of the relationship does not influence the

221   correlation or slope between the two traits. (3) In the ***EV_U-corrected regression***, the union

222   of EVs selected from both traits (denoted EV_U) was incorporated as covariates in the model

223   $X_2 \sim X_1 + EV\_U$. (4) Finally, in the **residual correlation analysis**, each trait was regressed

224   on its own EVs (X_1 on EV_X_1 and X_2 on EV_X_2), and the Pearson correlation between the

225   resulting residuals was calculated. To avoid inflated type I error due to overfitting, degrees

226   of freedom were adjusted to $n - k - 1$, where n is the number of species and k is the number

227   of eigenvectors in EV_U (Diniz-Filho and Tôrres 2002).

229       *Robust Estimators Used in PVR Analysis*

230   To address potential violations of model assumptions—such as outliers and non-normal

231   residuals—we incorporated several robust regression estimators into our PVR analysis. The

232   **L2** estimator, equivalent to ordinary least squares (OLS), represents the conventional

233   regression method in robust regression contexts. It minimizes the sum of squared residuals

234   but is highly sensitive to outliers due to its quadratic loss function, which disproportionately

235   amplifies large errors. The **L1** estimator, or least absolute deviation, minimizes the sum of

236   absolute residuals. Compared to L2, it is less affected by extreme values and performs better

237   when outliers are present. **M**-estimators generalize the likelihood framework and down-

238   weight the influence of large residuals. They behave like L2 for small errors and like L1 for

239   large errors, striking a balance between efficiency and robustness. **S**-estimators minimize a

240   robust measure of residual spread—typically based on median absolute deviation—offering

241   a high breakdown point and strong resistance to outliers. **MM**-estimators combine S- and M-

242   estimators, using the former for initial robustness and the latter for efficient final estimation.

243   This hybrid approach ensures both high robustness and statistical efficiency, making it well-

244   suited for noisy or heterogeneous datasets. For comparative purposes, we applied these

robust estimators alongside the traditional L2 estimator. All estimators were implemented

using R packages, with M-estimators via the *rlm* function from the *MASS* package (Version

7.3-55) (Venables and Ripley 2002), L1-estimators via *lad* function from *L1PACK* package

(Version 0.41-245) (Phillips 2002), and S- and MM-estimators via the *lmRob* function from

the *robust* package (Version 0.7-4) (Gervini and Yohai 2002).

Gold Standard for Trait Correlations in Each Simulation

Trait evolution for X_1 and X_2 was simulated along phylogenetic trees, with values recorded at

both tips and internal nodes. This design enabled the calculation of standardized

evolutionary changes per branch, expressed as $\Delta X_1/L$ and $\Delta X_2/L$, where L denotes is branch

length. These changes are independent from one branch to another, which allows us to use

conventional correlation tests to evaluate the relationship between trait changes across the

tree.

Our approach, while conceptually similar to PIC, avoids the need for model-based

ancestral state reconstruction by directly recording internal node values during simulations.

In contrast, PIC relies on reconstructing these states under a BM assumption, which may not

hold in many empirical datasets. Notably, PIC is mathematically equivalent to PGLS only

when the BM assumption is satisfied (Rohlf 2001, Blomberg et al. 2012). Because our

simulations are not constrained by such assumptions, they provide a more accurate and

flexible benchmark for evaluating comparative methods, including PIC, PGLS, and

optimized PVR.

To ensure methodological consistency across datasets and facilitate fair comparisons, we

uniformly applied Spearman's rank correlation to assess the association between $\Delta X_1/L$ and

$\Delta X_2/L$. This nonparametric method is robust to non-normality and outliers, and avoids

assumptions such as linearity or homoscedasticity.

271   **RESULTS**

To evaluate regression outcomes under different EV sets, we applied two models: OLS($X_1 \sim$

$X_2 + EV\_X_1$), where X_1 is the dependent variable; and OLS($X_2 \sim X_1 + EV\_X_2$), where X_2 is the

dependent variable. In each model, the corresponding eigenvectors (EV_X_1 or EV_X_2) were

included as covariates to account for phylogenetic structure. Unless otherwise specified, all

results in this section are based on 128-species balanced phylogenetic trees with uniformly

scaled branch lengths.

Eigenvector-Set Size and Exclusion of mAIC

To assess model complexity and mitigate the risk of overfitting, we first examined the

number of EVs selected by each method, ESRRV, mAIC, and mMorI. These methods were

applied separately to traits X_1 and X_2, and the resulting EVs were used as covariates in

regression models. As shown in Figure 1, the mAIC method produced substantially larger

EV sets, with median counts of 57 for EV_X_1, 56 for EV_X_2, and a striking 123 for their

union (EV_U), which is close to the total number of species (N = 128) in the dataset.

Such dimensionality severely compromises degrees of freedom and increases the risk of

overfitting. In contrast, ESRRV and mMorI consistently produced much smaller EV sets,

preserving model simplicity and statistical power. As a result, mAIC was excluded from

subsequent comparative analyses, and only ESRRV and mMorI were retained for

downstream evaluations.

Figure 1. Number of phylogenetic eigenvectors (EVs) selected by each method across trait

sets. For each method, EVs were selected separately for trait X_1 and trait X_2. The mAIC

method produced notably larger sets, with median counts reaching 57, 56, and 123 for

EV_X_1, EV_X_2, and their union EV_U, respectively. By contrast, ESRRV and mMorI

consistently selected fewer EVs, with union set medians remaining below 10. The total

number of species in the dataset was 128. See MATERIALS AND METHODS for details

regarding the three EV selection methods, ESRRV, mAIC, and mMorI. The data used to plot

this figure are available in Supplementary Table S1.

Impact of Variable Designation on Eigenvector Selection

The designation of dependent and independent variables significantly influences the results

of EV selection. As shown in Table 1, the discrepancy rate between EV selections was

notably high, with an average of 77% across all methods and simulation scenarios.

This demonstrates that relying on EVs selected from only one trait in regression models—

without accounting for discrepancies between the two sets—can introduce considerable risk

of bias, potentially compromising the robustness and reliability of the analysis.

Table 1. Inconsistency rates between phylogenetic eigenvector (EV) sets selected from traits

X_1 and X_2.

Simulation scenarios	ESRRV	mMorI
BM1 & BM1+BM2	83%	84%
BM & BM+Norm	80%	81%
Norm & Norm+BM	86%	76%
Norm1 & Norm1+Norm2	86%	4%
Shift	94%	100%

311 Values represent the percentage of inconsistent selections across 8,000 simulations per

312 scenario. Please refer to the MATERIALS AND METHODS for details regarding the EV

313 selection methods (ESRRV and mMorI) and the five simulation scenarios.

Divergent PVR Results Due to Different Eigenvector Sets

To assess how discrepancies between EV_X_1 and EV_X_2 affect PVR outcomes, we compared

two regression models: OLS($X_1 \sim X_2 + EV\_X_1$) and OLS($X_2 \sim X_1 + EV\_X_2$), applied across

various evolutionary scenarios and EV selection methods. Divergence was defined by either

a change in statistical significance ($p < 0.05$ vs. $p \geq 0.05$) or a reversal in slope direction (≥ 0

vs. < 0). As summarized in Table 2, divergence rates varied from 0.13% to 30.7%,

depending on the scenario and EV selection method. While mMorI generally produced more

consistent results, ESRRV showed greater divergence in several scenarios, most notably in

the Shift scenario (30.7%). To quantify these discrepancies, we computed Cohen's Kappa

values for each combination (Landis and Koch 1977). Asterisks in the table indicate the

degree of disagreement based on Kappa statistics: more stars denote greater inconsistency.

These results highlight that the inconsistencies in EV selection can substantially affect

regression outcomes—especially under non-stationary evolutionary conditions—

underscoring the need for trait-agnostic strategies such as *EV_U*.

Table 2. PVR model divergence due to differences in EV sets derived from X_1 and X_2.

Simulation scenarios	ESRRV	mMorI
BM1 & BM1+BM2	19.8% **	11.4% **
BM & BM+Norm	9.18% *	18.2% **
Norm & Norm+BM	11.5% **	10.8% **
Norm1 & Norm1+Norm2	6.44% *	0.13% *
Shift	30.7% ***	14.9% ***

Values represent the percentage of divergent regression results across 8,000 simulations per

scenario. Divergence was defined as differences in statistical significance ($p < 0.05$ vs. $p \geq$

0.05) or slope direction (≥ 0 vs. < 0) between the two models. Asterisks indicate the degree

of inconsistency between regression results based on Cohen's Kappa values (reversed scale):

*, $\kappa \geq 0.8$; **, $0.6 \leq \kappa < 0.8$; ***, $0.2 \leq \kappa < 0.4$. See Materials and Methods for details on

phylogenetic eigenvector (EV) selection methods (ESRRV and mMorI) and simulation

scenarios.

339   ***Divergence in Correlation Coefficients Due to EV Set Differences***

To quantify the sensitivity of PVR outcomes to differences in EV selection, we examined the

absolute differences in partial correlation coefficients between OLS($X_1 \sim X_2 + EV\_X_1$) and

OLS($X_2 \sim X_1 + EV\_X_2$) across five simulation scenarios. To this end, we computed the

absolute difference in partial correlation coefficients between the two regression models:

$|\mathrm{pcor}(X_1, X_2 + EV\_X_1) - \mathrm{pcor}(X_2, X_1 + EV\_X_2)|$.

As shown in Figure 2, coefficient discrepancies were consistently observed across all

five evolutionary scenarios, with many differences exceeding 10% in absolute value. This

suggests that EV selection based on variable designation can substantially affect not just the

significance of results, but also the estimated strength of trait associations. Notably, in the

Norm1 & Norm1+Norm2 scenario—characterized by minimal phylogenetic signal—

coefficient differences were markedly smaller, particularly when EVs were selected using

the mMorI method. These results reinforce the methodological risks associated with trait-

specific EV selection and motivate the use of more stable, trait-agnostic alternatives.

Figure 2. Distributions of differences in partial correlation coefficients between X_1 and X_2,

comparing PVR models using EV_X_1 versus EV_X_2 as covariates. The violin plot illustrates

the absolute differences in partial correlation coefficients ($|\mathrm{pcor}(X_1, X_2 + EV\_X_1) - \mathrm{pcor}(X_2,$

$X_1 + EV\_X_2|$) for cases with conflicting PVR results. Red dots indicate the median values,

and the vertical lines within the violins show the range of error bars. Paired t-tests and

Wilcoxon signed-rank tests both showed that the differences in partial correlation

coefficients between using EV_X_1 and EV_X_2 as covariates were highly significant across all

scenarios (all p-values $< 10^{-23}$). See MATERIALS AND METHODS for details regarding the EV

selection methods (ESRRV and mMorI) and the simulation scenarios (BM1 & BM1+BM2,

BM & BM+Norm, Norm & Norm+BM, and Norm1 & Norm1+Norm2, and Shift). The data

used to plot this figure are available in Supplementary Table S2.

Theoretical Rationale for Using the Union of Trait Eigenvectors

Our empirical findings demonstrated that regression outcomes based on EV sets selected

from different traits (EV_X_1 vs. EV_X_2) often diverged—both in terms of statistical

significance and estimated coefficients. This variability underscores a key methodological

issue: the conventional approach of selecting EVs based on a single trait may not adequately

capture the phylogenetic structure of the traits.

To clarify this point, we revisited the trait decomposition framework T = P + S,

originally proposed by Cheverud (1985), where where P represents the phylogenetic

component and S the specific or residual variation. Our derivation (see Appendix 1) shows

that when two traits share overlapping but non-identical phylogenetic structure, relying

solely on trait-specific EVs may lead to incomplete or biased control of phylogenetic effects.

In contrast, using the union of EVs selected from both traits (denoted EV_U) can offer more

stable and comprehensive adjustment. This insight also aligns with critiques of PGLS by

Westoby et al. (2023), who emphasized that PGLS typically accounts for phylogenetic

structure in the response variable but not in the predictor—potentially undermining model

reliability when both traits exhibit phylogenetic signal.

This theoretical insight also inspired an alternative strategy for jointly controlling the

phylogenetic non-independence of both traits: a two-step residual regression approach. In

this framework, each trait is first regressed against its own EVs to remove trait-specific

phylogenetic signals, and then the resulting residuals are used in a standard regression to

assess the trait–trait relationship. In the following sections, we compare this residual-based

method with models using EV_U and EVs derived independently from each trait across five

evolutionary scenarios.

Performance of Eigenvector Selection Methods and Regression Approaches

In addition to evaluating different regression strategies, we compared the performance of

two eigenvector selection methods—ESRRV and mMorI—to examine how the choice of EV

selection influences regression outcomes. As shown in Figure 3, mMorI (solid lines)

generally outperformed ESRRV (dashed lines), particularly in terms of regression accuracy,

except in one case—when using EV_X_2 under the BM & BM+Norm scenario.

When comparing regression approaches under a fixed EV selection method, we found

that models using EV_U consistently achieved the highest accuracy across five evolutionary

scenarios when mMorI was used (red solid lines). With ESRRV, EV_U was still preferred in

scenarios such as BM1 & BM1+BM2, BM & BM+Norm, and Shift, where it was nearly the

top performer (red dashed lines), though not always the best. Notably, under conditions of

minimal phylogenetic autocorrelation (Norm1 & Norm1+Norm2), the union method

performed the worst among all tested approaches.

We also assessed a two-step residual regression strategy, where each trait was regressed

on its own EVs before analyzing the correlation between residuals. However, this approach

yielded lower accuracy overall and was therefore not selected for further analyses.

Based on these findings, we chose the mMorI method combined with *EV_U* as the

representative PVR configuration in subsequent comparisons, which included integration

with robust estimators and benchmarking against PIC and PGLS methods.

Figure 3. Performance across different phylogenetic eigenvector (EV) selection methods and

regression approaches. This figure shows the performance of different EV selection methods

(ESRRV and mMorI) combined with four regression strategies: (1) using EV_X_1 as

covariates; (2) using EV_X_2 as covariates; (3) using the union set of EV_X_1 and EV_X_2,

EV_U, as covariates; and (4) regressing the residuals of X_1 and X_2, each obtained from

separate regressions against their respective EVs. Simulations were conducted on 128-

species balanced trees ($\rho = 1$). The comparisons were performed across five simulation

scenarios: a) BM1 & BM1+BM2; b) BM & BM+Norm; c) Norm & Norm+BM; d) Norm1 &

Norm1+Norm2; and e) Shift. See MATERIALS AND METHODS for details regarding the EV

selection methods (ESRRV and mMorI) and the definitions of each simulation scenario. The

data used to plot this figure are available in Supplementary Table S3.

422   ***Performance of Robust Estimators in PVR Models***

We evaluated the performance of five regression estimators—L1, L2, M, MM, and S—

within the optimized PVR framework (using mMorI and EV_U), focusing on the accuracy of

regression outcomes across five evolutionary scenarios (Fig. 4). The S estimator generally

exhibited the lowest accuracy, ranking last in three scenarios and performing suboptimally in

the others, particularly under the Shift condition. In contrast, L1, L2, M, and MM produced

broadly similar results, with each showing modest advantages under certain conditions. No

single estimator consistently outperformed the others across all scenarios. Based on these

findings, we excluded S from further analysis and retained the remaining four estimators for

integration into the PVR framework in subsequent comparisons with PIC and PGLS.

Figure 4. Comparison of robust estimator performance in phylogenetic eigenvector

regression (PVR). This figure compares the performance of five robust estimators (L1, L2,

M, MM, and S) integrated into the optimized PVR framework (selected based on mMorI and

EV_U). The five evolutionary scenarios—a) BM1 & BM1+BM2, b) BM & BM+Norm, c)

Norm & Norm+BM, d) Norm1 & Norm1+Norm2, and e) Shift—represent different

combinations of evolutionary models and relationships between X_1 and X_2. See MATERIALS

AND METHODS for details on the phylogenetic EV selection method (mMorI), the definitions

of robust estimators, the construction of *EV_U*, and the definitions of each simulation

scenario. The data used to plot this figure are available in Supplementary Table S4.

Comparative Evaluation of Optimized PVR, PGLS, and PIC

Overall research setup.—Having integrated four estimators (L1, L2, M, and MM) into the

PVR framework using mMorI and *EV_U*, we proceeded to compare these optimized PVR

approaches with PGLS and PIC. For PGLS, the most appropriate model for each dataset was

selected from five candidate evolutionary models—BM, Pagel's λ, OU fixed, OU random,

and EB—via AIC minimization (hereafter referred to as PGLS-best). Following Adams et al.

(2024), who demonstrated that PIC-MM outperformed other PIC variants (e.g., PIC-L1,

PIC-M) and various robust-estimator PGLS approaches under non-stationary evolutionary

scenarios, we included PIC-MM and PIC-L2 (representing standard PIC) in our analyses to

compare their performance with optimized PVR across different evolutionary contexts.

To comprehensively assess the performance of the optimized PVR framework relative to

PIC and PGLS, we conducted evaluations under five distinct evolutionary scenarios, applied

consistently across all tree types. These scenarios were tested on multiple phylogenetic

structures, beginning with 128-species balanced trees under three different branch-length

scaling conditions: (1) uniformly scaled branches (ρ = 1), (2) compressed ancestral and

elongated terminal branches (ρ = 0.1), and (3) elongated ancestral branches (ρ = 2). We then

extended the analyses to two additional tree types representing key structural contrasts: a

reduced sampling scenario using 16-species balanced trees (ρ = 1) and a topologically

imbalanced scenario using 128-species ladder-shaped trees (ρ = 1).

128-species balanced trees with uniformly scaled branches (ρ = 1).—Across five

evolutionary scenarios differing in the strength and structure of phylogenetic autocorrelation,

method accuracy was plotted against increasing variance (shift) gradient values (Fig. 5). At

low variance, all methods—including optimized PVR variants (solid lines) and benchmarks

PIC-L2, PIC-MM, and PGLS-best (dashed lines)—performed similarly. However,

performance diverged significantly as variance (shift) increased.

Under BM1 & BM1+BM2, both X_1 and X_2 evolved under strong phylogenetic signal,

with PGLS-best and PIC-L2 consistently outperforming all PVR variants (Fig. 5a). Under

BM & BM+Norm, which features substantial differences in phylogenetic signal between

traits, PGLS-best exhibited significant accuracy loss, while PVR-MM and PVR-L1

performed slightly better than the other methods (Fig. 5b). The opposite pattern emerged

under Norm & Norm+BM, where PGLS-best and PIC-L2 again dominated, while PVR-MM

and PVR-L1 performed most poorly (Fig. 5c). In Norm1 & Norm1+Norm2, all methods

performed similarly, with the exception of PVR-L1, which exhibited a noticeable drop in

accuracy at one variance level (Fig. 5d). Among the remaining methods, the three other PVR

variants and PGLS-best demonstrated marginally higher accuracy than the PIC-based

methods.

The most pronounced contrasts emerged in the Shift scenario, modeling non-stationary

evolutionary change (Fig. 5e). Whereas PGLS-best and PIC-L2 suffered dramatic accuracy

reductions as shift magnitude increased, PVR methods and PIC-MM maintained consistently

high performance, demonstrating robustness against abrupt, branch-specific evolutionary

changes. Surprisingly, PVR-L2, without explicit robust estimators, performed comparably to

PIC-MM, suggesting inherent robustness conferred by phylogenetic eigenvectors

themselves. Among PVR variants, while L2 and M estimators showed slightly lower

accuracy, L1 and MM—together with PIC-MM—consistently maintained higher accuracy

across different shift magnitudes.

Figure 5. Comparative accuracy of PVR, PGLS, and PIC methods on 128-species balanced

phylogenetic trees ($\rho = 1$). The comparisons are performed under five evolutionary

scenarios: a) BM1 & BM1+BM2, both traits exhibit strong phylogenetic signal; b) BM &

BM+Norm and c) Norm & Norm+BM, large disparity in phylogenetic signal strength

between traits; d) Norm1 & Norm1+Norm2, both traits lack phylogenetic structure; e) Shift,

a synchronous trait shift was introduced on a single internal branch, while the two traits

evolved independently. See MATERIALS AND METHODS for details on the definitions of

each simulation scenario. The data used to plot this figure are available in Supplementary

Table S5.

128-species balanced trees with compressed ancestral branches ($\rho = 0.1$).—With

compressed ancestral branches, methods exhibited closely aligned performance across

stationary scenarios, except for PVR-L1 (Fig. 6a-d). Under BM & BM+Norm, PVR-L1 was

distinctly superior; however, it underperformed in other stationary scenarios (BM1 &

BM1+BM2, Norm & Norm+BM, and Norm1 & Norm1+Norm2). PVR-L2, PVR-M, and

PVR-MM consistently matched PIC-L2, PIC-MM, and PGLS-best. In the Shift scenario,

PGLS-best and PIC-L2 again showed accuracy declines, while PIC-MM and all PVR

methods remained robust (Fig. 6e).

Figure 6. Performance of PVR, PGLS, and PIC on 128-species balanced phylogenetic trees

($\rho = 0.1$). The comparisons are performed under five evolutionary scenarios: a) BM1 &

BM1+BM2; b) BM & BM+Norm; c) Norm & Norm+BM; d) Norm1 & Norm1+Norm2; e)

Shift. See MATERIALS AND METHODS for details on the definitions of each simulation

scenario. The data used to plot this figure are available in Supplementary Table S6.

128-species balanced trees with elongated ancestral branches ($\rho = 2$).—When deep

ancestral branches were elongated, accuracy patterns diverged notably from the other

branch-length scenarios (Fig. 7). In BM1 & BM1+BM2, PGLS-best, PIC-L2, and PIC-MM

substantially outperformed PVR methods. Under BM & BM+Norm, PGLS-best accuracy

declined dramatically at intermediate variance, whereas other methods remained stable.

Interestingly, in Norm & Norm+BM with elongated ancestral branches ($\rho = 2$), all

methods achieved perfect (100%) accuracy (Fig. 7c). Analysis of raw simulation data

indicated strongly positive correlations between traits (Supplementary Table S7). With

elongated ancestral branches and extremely short terminal branches, BM—which scales

proportionally with branch length—contributed minimally to trait differences along the short

terminal branches. In contrast, the independent normal (Norm) component, being

independent of branch length, dominated these short branches. Consequently, the minimal

terminal BM contributions made X_2 nearly collinear with X_1, effectively eliminating

methodological performance differences.

In the Shift scenario, PGLS-best and PIC-L2 experienced marked accuracy reductions at

high shift magnitudes (Fig. 7c). PIC-MM and PVR methods remained relatively robust. PIC-

MM generally maintained a slight advantage at this larger ρ values (Fig.7), whereas PVR-L1

performed slightly better at smaller ρ (see Figure 6 and Supplementary Table S6).

Figure 7. Performance of PVR, PGLS, and PIC on 128-species balanced phylogenetic trees

($\rho = 2$). The comparisons are performed under five evolutionary scenarios: a) BM1 &

BM1+BM2; b) BM & BM+Norm; c) Norm & Norm+BM; d) Norm1 & Norm1+Norm2; e)

Shift. See MATERIALS AND METHODS for details on the definitions of each simulation

scenario. The data used to plot this figure are available in Supplementary Table S8.

16-species balanced trees with uniformly scaled branches.— The overall performance

patterns under reduced taxonomic sampling were broadly consistent with those observed

under 128-species balanced trees (Fig. 8). Although minor differences existed in the exact

performance ranks of certain methods, the scenario-dependent relative advantages among

methods remained unchanged.

Figure 8. Performance of PVR, PGLS, and PIC on 16-species balanced trees with uniformly

scaled branches ($\rho = 1$). The comparisons are performed under five evolutionary scenarios:

a) BM1 & BM1+BM2; b) BM & BM+Norm; c) Norm & Norm+BM; d) Norm1 &

Norm1+Norm2; e) Shift. See MATERIALS AND METHODS for details on the definitions of

each simulation scenario. The data used to plot this figure are available in Supplementary

Table S9.

128-species Ladder-Shaped Trees (ρ = 1)—During simulation analyses using ladder-shaped

phylogenies, we observed that the mMorI method frequently failed to select a sufficient

number of eigenvectors, often returning none at all. Quantitatively, the median number of

eigenvectors in the union set (*EV_U*) was only 1. In contrast, the mAIC method produced

excessively large EV sets, with a median *EV_U* count of 80—raising concerns about model

overfitting. The ESRRV method yielded a more moderate and balanced result (median

EV_U = 31), effectively supporting phylogenetic correction. Accordingly, ESRRV was

adopted for all subsequent analyses involving ladder-shaped trees.

Given these patterns, we further examined the underlying reasons why mMorI

underperformed in ladder-shaped trees. Cumulative eigenvalue distributions differed notably

between balanced and ladder-shaped trees (Fig. 9). The extreme topological imbalance of

ladder-shaped trees distorts the distribution of phylogenetic signal, resulting in smaller

disparities among eigenvalues. This in turn hinders the identification of informative

eigenvectors—likely explaining why mMorI failed to select sufficient EVs under these

conditions. These findings highlight the importance of tailoring eigenvector selection

strategies to tree topology and signal structure.

Figure 9. Contribution of phylogenetic eigenvectors across different tree structures. Both the

balanced and ladder-shaped trees shown here are 128-species phylogenies with uniformly

scaled branches (ρ = 1). The data used to plot this figure are available in Supplementary

Table S10.

We next evaluated the comparative performance of PVR, PGLS, and PIC methods under

ladder-shaped tree conditions (Fig. 10). While overall trends resembled those observed under

balanced trees, important distinctions emerged in the Shift scenario. PVR-L1, PVR-L2, and

PVR-M showed notable accuracy declines at higher shift magnitudes—similar to PGLS-best

and PIC-L2—whereas PIC-MM and PVR-MM maintained robustness, reaffirming their

superior performance under non-stationary conditions.

Collectively, these results illustrate that method performance varies markedly depending

on tree structure and evolutionary scenario. No single method consistently outperformed the

others across all conditions, highlighting the need for context-specific model selection in

phylogenetic comparative analysis.

Figure 10. Performance of PVR, PGLS, and PIC on on 128-species ladder-shaped

phylogenetic trees ($\rho = 1$). The comparisons are performed under five evolutionary

scenarios: a) BM1 & BM1+BM2; b) BM & BM+Norm; c) Norm & Norm+BM; d) Norm1 &

Norm1+Norm2; e) Shift. This figure differs from Figures 5-8 in that all PVR methods use

ESRRV for eigenvector selection, rather than mMorI. See MATERIALS AND METHODS for

details on the definitions of each simulation scenario and the two EV selection methods. The

data used to plot this figure are available in Supplementary Table S11.

DISCUSSION

PVR was originally proposed as a flexible alternative to model trait–trait correlations while

accounting for phylogenetic structure through eigenvectors derived from phylogenetic

distance matrices (Diniz-Filho et al. 1998). However, this conventional view (Laurin 2010,

Freckleton et al. 2011, Pennell and Harmon 2013, Hansen 2014) holds that PVR is less

reliable in controlling for phylogenetic non-independence and produces less robust estimates

than likelihood-based approaches, particularly PGLS.

In this study, we revisited the PVR framework and introduced several key optimizations.

First, we evaluated how the selection of phylogenetic eigenvectors varies depending on

which trait is treated as the dependent variable. Our analysis revealed that the sets of

eigenvectors selected based on X_1 and X_2 often differ substantially (Table 1), and when these

trait-specific eigenvectors are used as covariates in regression models, they can lead to

markedly different correlation estimates (Table 2 and Fig. 2). Both theoretical derivation and

simulation-based validation demonstrated that using the union of eigenvectors selected from

both traits (*EV_U*) better control for phylogenetic independence than using those selected

from a single trait (Appendix 1 and Fig.3).

Building on this foundation, we further integrated robust estimators into the PVR

framework to improve its resilience to noise and data irregularities. The resulting variants—

PVR-L1, PVR-L2, PVR-M, and PVR-MM—were then systematically compared against

established methods including PGLS-best, PIC-L2 (standard PIC), and PIC-MM across a

wide range of evolutionary scenarios.

Our findings challenge the conventional view that PGLS universally outperforms PVR.

While PGLS-best and PIC-L2 excelled when dependent traits exhibited strong phylogenetic

signal, optimized PVR methods—particularly PVR-MM—demonstrated superior robustness

in scenarios characterized by weak or asymmetric phylogenetic signal and non-stationary

evolutionary shifts. These results underscore the context-dependent nature of method

performance and suggest that researchers should move away from universally applying

PGLS, instead tailoring model choice to the evolutionary properties of their data. The

optimized PVR framework, especially when coupled with robust estimators, offers a

valuable and versatile analytical alternative for phylogenetic comparative studies.

PVR Robustness Under Non-Stationary Evolutionary Shift

One of the most challenging conditions for phylogenetic models is non-stationary

evolutionary change. Under such shift scenarios, our results showed that the optimized PVR

framework—especially PVR-MM and even PVR-L2—maintained high and stable accuracy,

closely matching PIC-MM across most tree structures and shift magnitudes. In contrast,

PGLS-best and PIC-L2 suffered pronounced accuracy declines under increasing shift

intensity.

Remarkably, PVR-L2, which does not incorporate any explicit robust estimator,

performed comparably to PIC-MM—one of the most robust models available (Adams et al.

2024)— under balanced tree conditions. This suggests that the use of phylogenetic

eigenvectors may confer inherent robustness, likely due to their capacity to absorb structured

variation across the tree. Among PVR variants, L1 and MM estimators provided additional

but modest gains, especially under large shifts and in most tree conditions. However, under

ladder-shaped phylogenies, only PVR-MM and PIC-MM retained robustness, highlighting

their stability in particularly asymmetric topologies.

Theoretically, this robustness stems from PVR's model-agnostic design: it does not rely

on global stationarity assumptions but instead decomposes phylogenetic structure via

orthogonal eigenvectors, with some exhibiting greater sensitivity to variation in specific

regions of the tree. This multiscale sensitivity allows PVR to flexibly accommodate

localized shifts in evolutionary rates (Diniz-Filho et al. 2012b, Gouveia et al. 2013, Diniz-

Filho et al. 2014, Diniz-Filho et al. 2015, Diniz-Filho et al. 2019).

Together, these empirical and theoretical insights reinforce the view that PVR is not

only a competitive alternative to conventional phylogenetic comparative methods under

stationary models, but also a particularly robust tool for modeling evolutionary

nonstationarity—especially when paired with robust estimators such as MM.

651   ***PVR Advantage Under Weak Phylogenetic Signal***

In addition to its robustness under non-stationary evolutionary shifts, the PVR framework

also exhibited a notable advantage in scenarios characterized by weak phylogenetic signal.

In scenarios with weak phylogenetic signal, such as Norm1 & Norm1+Norm2, PVR

methods, particularly L2 and M, showed a clear advantage.

We found that the phylogenetic signal (λ) in X_1 and X_2 under this scenario was

significantly greater than zero in a proportion ranging from 0.5% to 6% (depending on the

structures of phylogenetic trees), with the maximum value reaching 1.13. While such signals

are generally weak, their presence indicates that assessing the performance of phylogenetic

methods in these contexts is not without value. Note that although λ is typically bounded

between 0 and 1, values slightly above 1 can occur due to maximum likelihood estimation

and are not uncommon in empirical datasets.

When the phylogenetic signal is weak, the coefficients of eigenvectors as covariates tend

to be small. Since these coefficients are estimated freely based on the data, without

constraint to a specific range, they provide modeling flexibility even when phylogenetic

effects are minimal. Therefore, PVR methods may be more effective than PIC and PGLS in

handling cases with weak phylogenetic signals.

669   ***Comparison with Two-Step Residual Regression Approaches***

In addition to comparing EV sets derived from individual traits and their union (*EV_U*), we

also evaluated a two-step residual regression approach as an alternative means of controlling

phylogenetic non-independence. In this method, each trait is first regressed against its own

set of EVs to remove trait-specific phylogenetic signals. The resulting residuals are then used

in a standard regression to assess the relationship between traits. This residual-based strategy

offers a symmetric structure. However, our results showed that this method consistently

underperformed relative to the *EV_U*-based regression approach across all evolutionary

scenarios (Fig. 4). While the residual method did not exhibit clear advantages over trait-

specific EV regressions, it showed notably lower accuracy in certain scenarios—particularly

under shift conditions.

Theoretically, this outcome is not unexpected. When two traits share overlapping

phylogenetic structure—as often occurs due to shared ancestry—independently removing

phylogenetic signal from each trait can result in the unintended removal of shared

components that may underlie genuine biological associations. As a result, residual-based

approaches may reduce correlation accuracy or introduce bias under certain conditions.

In contrast, the *EV_U* approach directly incorporates the phylogenetic information from

both traits within a single model, without presuming a causal direction or requiring

sequential correction. This unified strategy helps preserve meaningful shared signal while

effectively controlling for phylogenetic effects. Our simulation results support this

reasoning: *EV_U* consistently yielded higher accuracy across diverse evolutionary scenarios,

reinforcing its utility as the preferred method in our optimized PVR framework.

Flexibility of Eigenvector-Based Approaches in Phylogenetic Modeling

Unlike PGLS and other model-based approaches that rely on predefined evolutionary models

(Freckleton et al. 2011, O'Meara and Beaulieu 2016), PVR extracts phylogenetic structure

through eigenvector decomposition, offering greater modeling flexibility. This feature is

particularly valuable when the true evolutionary process deviates from canonical models

(e.g., BM, OU or EB), or when model uncertainty is high (Diniz-Filho et al. 2014). Given

the limited repertoire of evolutionary models currently available, it is unlikely that all real-

world evolutionary patterns can be adequately captured by likelihood-based methods. Even

if model libraries were expanded in the future, distinguishing among them would require

repeated model fitting and comparison, rendering the process both cumbersome and prone to

error.

PVR's modular use of phylogenetic eigenvectors offers a key advantage: unlike PGLS,

where phylogenetic information is embedded in model-specific assumptions, PVR expresses

it through separable components, enabling seamless integration with machine learning

algorithms such as random forests and neural networks (Penone et al. 2014, Debastiani et al.

2021). This portability enables applications beyond traditional regression, including trait

imputation and classification in a phylogenetic context.

Third, PVR offers a structural advantage in handling directional asymmetry in trait

relationships. Unlike correlation analysis, which treats both variables symmetrically, OLS

requires a clear distinction between the independent variable (X) and the dependent variable

(Y). This distinction is built into the model's structure: X is assumed to be fixed and

measured without error, while all residual variation is attributed to Y (McArdle 1988,

Osborne and Waters 2002, Casson and Farmer 2014, Jarantow et al. 2023). Moreover, OLS

implicitly encodes a causal direction—predicting changes in Y as a function of X. As a

result, OLS is inherently asymmetrical: reversing X and Y leads to a different model,

different estimates, and potentially different conclusions. PGLS inherited the asymmetry

from OLS regression, requiring a clear assignment of dependent and independent variables.

However, when the two traits differ in the strength of their phylogenetic signal, assigning the

trait with weaker signal as the dependent variable can distort the estimated relationship

between them (Chen et al. 2023). In situations where the trait that is either logically

considered causal or measured with less error happens to exhibit a stronger phylogenetic

signal, researchers face a dilemma: treating it as the independent variable aligns with

biological reasoning or data reliability, but doing so may compromise phylogenetic model

performance in PGLS.

PVR, by contrast, avoids this conflict. By including the union of eigenvectors selected

from both traits as covariates, PVR allows either trait to be designated as the independent

variable without affecting the accuracy of the model. This design enables researchers to

select predictors based on causal logic or measurement quality, without being constrained by

the phylogenetic properties of the traits—removing a key asymmetry-related limitation

present in OLS-derived methods like PGLS.

Dependence of PGLS-best Performance on Dependent Variable Selection

From Figures 5-8 and Figure 10, we observed a consistent pattern in the relative

performance of PGLS-best compared to other methods across the first four evolutionary

scenarios. In BM1 & BM1+BM2, PGLS-best generally performed the best or was close to

the best; in BM & BM+Norm, it was the worst or close to the worst; in Norm & Norm+BM,

it reversed again to be the best or close to the best; and in Norm1 & Norm1+Norm2, its

performance was average. In BM1 & BM1+BM2 and Norm & Norm+BM, X_2 exhibited

stronger phylogenetic signal than X_1, while in BM & BM+Norm, X_2 had a weaker

phylogenetic signal compared to X_1. In Norm1 & Norm1+Norm2, there was little to no

difference in the phylogenetic signals of the two traits. In this study, X_2 was treated as the

dependent variable and X_1 as the independent variable. It seems that the performance of

PGLS-best depends on whether the trait with the stronger phylogenetic signal is chosen as

the dependent variable, which is the focus of our unpublished manuscript (Chen et al. 2023).

As the study has faced multiple rejections in various journals and still requires further

refinement, it should not be considered a mature benchmark method for evaluating PVR

performance. However, it can be speculated that, if such optimization were performed, the

relative advantage of PGLS-best might become even more pronounced.

Broader Implications: Variable Asymmetry Beyond Phylogenetic Models

While this study focuses primarily on phylogenetic comparative methods, similar challenges

may arise in other contexts involving structured data—particularly those with spatial

autocorrelation. Spatial autocorrelation refers to the tendency for geographically proximate

observations to exhibit greater similarity than those farther apart. It is common in ecological

and evolutionary datasets and requires statistical models that account for spatial structure to

avoid biased estimates (Doreian 1980, Dow et al. 1982, Legendre 1993, Keitt et al. 2002, He

et al. 2003, Cottenie 2005, Diniz-Filho and Bini 2005, Dormann et al. 2007, Dale and Fortin

2009, Peres-Neto and Legendre 2010).

Preliminary analyses suggest that the asymmetry introduced by variable designation—

previously discussed in the context of PGLS and PVR—also occurs in spatial models. In

methods such as spatial eigenvector mapping, conditional autoregressive (CAR) models,

simultaneous autoregressive (SAR) models, and trend-surface generalized additive models

(GAMs), we observed outcome discrepancies when the dependent and independent variables

were swapped. These findings, summarized in Supplementary Table S12, indicate that

variable swapping may affect model performance across a wide range of autocorrelation-

adjusted frameworks. A full analysis of these effects is beyond the scope of this study, but

future work will explore their implications more systematically, with potential relevance to

both spatial and ecological inference.

In many real-world applications, multiple forms of autocorrelation—such as spatial and

phylogenetic structure—must be addressed simultaneously. This introduces additional

complexity in both variable designation and eigenvector selection. As discussed by Safi and

Pettorelli (2010), properly balancing spatial, phylogenetic, and environmental components

depends on the specific analytical goals. Although our current study emphasizes

phylogenetic relationships, their work underscores the importance of aligning

methodological choices with the biological structure of the data. Future extensions of PVR

and related models may benefit from more flexible integration of multiple autocorrelation

sources to improve model reliability across complex ecological scenarios.

DATA AVAILABILITY

Supplementary Tables S1-S12 are included in Supplementary_tables.xlsx. The raw data for

the five simulation scenarios are provided in the file Raw_data.zip. The scripts used to

generate the results presented in the main text, including phylogenetic tree construction, trait

data simulation, regression model fitting, and statistical analysis, are available in

Main_code.zip. The relevant codes for evaluating spatial statistical methods are included in

Spatial_methods.zip. Additionally, standalone functions for eigenvector selection

(MoransI.R and ESRRV.R) are available separately. All these files and .R scripts can be

found in the Dryad data repository: http://datadryad.org/stash/share/dLL0-

zHGm_Wyrli08AJ9P4pR-yFycPeYOqktVZhEUVA.

ACKNOWLEDGMENTS

We are deeply grateful to the reviewers for their valuable comments and suggestions, which significantly improved the quality of our work. In particular, we would like to thank Prof. J. A. F. Diniz-Filho for his insightful feedback. We also appreciate the contributions from the anonymous reviewer, whose detailed suggestions greatly helped us refine our analysis and presentation. This study was supported in part by the National Natural Science Foundation of China (grant number 31671321), with leftover resources used for occasional research expenses. To enhance language and readability, this article has undergone refinement by ChatGPT 4o. Subsequently, the authors conducted a thorough review and made necessary edits, taking full responsibility for the publication's content.

AUTHOR CONTRIBUTIONS

DKN and ZLC conceived the research hypothesis; ZLC performed the simulations and analyzed the data. ZLC drafted the manuscript, and DKN revised the manuscript. Both authors agree with the final version and the submission.

REFERENCES

Adams R., Cain Z., Assis R., DeGiorgio M. 2024. Robust phylogenetic regression. Syst. Biol., 73:140-157.

Blomberg S.P., Lefevre J.G., Wells J.A., Waterhouse M. 2012. Independent contrasts and PGLS regression estimators are equivalent. Syst. Biol., 61:382-391.

Burns J.H., Anacker B.L., Strauss S.Y., Burke D.J. 2015. Soil microbial community variation correlates most strongly with plant species identity, followed by soil chemistry, spatial location and plant genus. AoB PLANTS, 7:plv030.

Casson R.J., Farmer L.D. 2014. Understanding and checking the assumptions of linear regression: a primer for medical researchers. Clin. Exp. Ophthalmol., 42:590-596.

Chen Z.-L., Guo H.-J., Niu D.-K. 2023. Dependent variable selection in phylogenetic generalized least squares regression analysis under Pagel's lambda model. bioRxiv:2023.2005.2021.541623.

Cheverud J.M., Dow M.M., Leutenegger W. 1985. The quantitative assessment of

phylogenetic constraints in comparative analyses: sexual dimorphism in body weight among

primates. Evolution, 39:1335-1351.

Cornwallis C.K., Griffin A.S. 2024. A guided tour of phylogenetic comparative methods for

studying trait evolution. Annu. Rev. Ecol. Evol. Syst., 55:181-204.

Cottenie K. 2005. Integrating environmental and spatial processes in ecological community

dynamics. Ecol. Lett., 8:1175-1182.

Dale M.R.T., Fortin M.-J. 2009. Spatial autocorrelation and statistical tests: Some solutions.

JABES, 14:188-206.

Debastiani V.J., Bastazini V.A.G., Pillar V.D. 2021. Using phylogenetic information to

impute missing functional trait values in ecological databases. Ecol. Inform., 63:101315.

Desdevises Y., Legendre P., Azouzi L., Morand S. 2003. Quantifying phylogenetically

structured environmental variation. Evolution, 57:2647-2652.

Dewar A.E., Belcher L.J., West S.A. 2025. A phylogenetic approach to comparative

genomics. Nat. Rev. Genet., 10.1038/s41576-024-00803-0.

Diniz-Filho J.A.F., Alves D.M.C.C., Villalobos F., Sakamoto M., Brusatte S.L., Bini L.M.

2015. Phylogenetic eigenvectors and nonstationarity in the evolution of theropod dinosaur

skulls. J. Evol. Biol., 28:1410-1416.

Diniz-Filho J.A.F., Bini L.M. 2005. Modelling geographical patterns in species richness

using eigenvector-based spatial filters. Glob. Ecol. Biogeogr., 14:177-185.

Diniz-Filho J.A.F., Bini L.M., Rangel T.F., Morales-Castilla I., Olalla-Tárraga M.Á.,

Rodríguez M.Á., Hawkins B.A. 2012a. On the selection of phylogenetic eigenvectors for

ecological analyses. Ecography 35:239-249.

Diniz-Filho J.A.F., Bini M., Sakamoto M., Brusatte S. 2014. Phylogenetic eigenvector

regression in paleobiology. Rev. Bras. Paleontol., 17:107-122.

Diniz-Filho J.A.F., Cianciaruso M.V., Rangel T.F., Bini L.M. 2011. Eigenvector estimation

of phylogenetic and functional diversity. Funct. Ecol., 25:735-744.

Diniz-Filho J.A.F., de Sant'Ana C.E.R., Bini L.M. 1998. An eigenvector method for

estimating phylogenetic inertia. Evolution, 52:1247-1262.

Diniz-Filho J.A.F., Jardim L., Mondanaro A., Raia P. 2019. Multiple components of

phylogenetic non-stationarity in the evolution of brain size in fossil Hominins. Evol. Biol.,

46:47-59.

Diniz-Filho J.A.F., Nabout J.C. 2009. Modeling body size evolution in Felidae under

alternative phylogenetic hypotheses. Genet Mol Biol, 32:170-176.

Diniz-Filho J.A.F., Rangel T.F., Santos T., Mauricio Bini L. 2012b. Exploring patterns of

interspecific variation in quantitative traits using sequential phylogenetic eigenvector

regressions. Evolution, 66:1079-1090.

Diniz-Filho J.A.F., Rodríguez M.Á., Bini L.M., Olalla-Tarraga M.Á., Cardillo M., Nabout

J.C., Hortal J., Hawkins B.A. 2009. Climate history, human impacts and global body size of

Carnivora (Mammalia: Eutheria) at multiple evolutionary scales. J. Biogeogr., 36:2222-

2236.

Diniz-Filho J.A.F., Tôrres N.M. 2002. Phylogenetic comparative methods and the

geographic range size – body size relationship in new world terrestrial carnivora. Evol.

Ecol., 16:351-367.

Divíšek J., Chytrý M., Beckage B., Gotelli N.J., Lososová Z., Pyšek P., Richardson D.M.,

Molofsky J. 2018. Similarity of introduced plant species to native ones facilitates

naturalization, but differences enhance invasion success. Nat. Commun., 9:4631.

Doreian P. 1980. Linear models with spatially distributed data: spatial disturbances or spatial

effects? Soc. Methods Res., 9:29-60.

Dormann C.F., McPherson J.M., Araújo M.B., Bivand R., Bolliger J., Carl G., Davies R.G.,

Hirzel A., Jetz W., Kissling W.D., Kühn I., Ohlemüller R., Peres-Neto P.R., Reineking B.,

Schröder B., Schurr F.M., Wilson R. 2007. Methods to account for spatial autocorrelation in

the analysis of species distributional data: a review. Ecography, 30:609-628.

Dow M.M., Burton M.L., White D.R. 1982. Network autocorrelation: A simulation study of

a foundational problem in regression and survey research. Social Networks, 4:169-200.

Felsenstein J. 1985. Phylogenies and the comparative method. Amer. Natur., 125:1-15.

Freckleton R.P., Cooper N., Jetz W. 2011. Comparative methods as a statistical fix: the

dangers of ignoring an evolutionary model. Amer. Natur., 178:E10-E17.

Gao X., Dai J., Tao Z., Shahzad K., Wang H. 2023. Autumn phenology of tree species in

China is associated more with climate than with spring phenology and phylogeny. Front.

Plant Sci., 14:1040758.

Garamszegi L.Z. 2014. Modern Phylogenetic Comparative Methods and Their Application

in Evolutionary Biology: Concepts and Practice. Berlin, Springer.

Gervini D., Yohai V.J. 2002. A class of robust and fully efficient regression estimators. Ann.

Statist., 30:583-616.

Gouveia S.F., Hortal J., Cassemiro F.A.S., Rangel T.F., Diniz-Filho J.A.F. 2013.

Nonstationary effects of productivity, seasonality, and historical climate changes on global

amphibian diversity. Ecography, 36:104-113.

Grafen A. 1989. The phylogenetic regression. Philos. Trans. R. Soc. Lond. B Biol. Sci.,

326:119-157.

Griffith D.A., Peres-Neto P.R. 2006. Spatial modeling in ecology: the flexibility of

eigenfunction spatial analyses. Ecology, 87:2603-2613.

Hansen T.F. 2014. Use and misuse of comparative methods in the study of adaptation. In:

Garamszegi LZ editor. Modern Phylogenetic Comparative Methods and Their Application in

Evolutionary Biology: Concepts and Practice. Berlin, Heidelberg, Springer Berlin

Heidelberg, p. 351-379.

He F., Zhou J., Zhu H. 2003. Autologistic regression model for the distribution of

vegetation. JABES, 8:205-222.

Huber P.J. 1964. Robust estimation of a location parameter. Ann. Math. Statist., 35:73-101.

Jarantow S.W., Pisors E.D., Chiu M.L. 2023. Introduction to the use of linear and nonlinear

regression analysis in quantitative biological assays. Curr. Protoc., 3:e801.

Keitt T.H., Bjørnstad O.N., Dixon P.M., Citron-Pousty S. 2002. Accounting for spatial

pattern when modeling organism-environment interactions. Ecography, 25:616-625.

Landis J.R., Koch G.G. 1977. The measurement of observer agreement for categorical data.

Biometrics, 33:159-174.

Lanuza J.B., Rader R., Stavert J., Kendall L.K., Saunders M.E., Bartomeus I. 2023.

Covariation among reproductive traits in flowering plants shapes their interactions with

pollinators. Funct. Ecol., 37:2072-2084.

Laurin M. 2010. Assessment of the relative merits of a few methods to detect evolutionary

trends. Syst. Biol., 59:689-704.

Legendre P. 1993. Spatial autocorrelation: trouble or new paradigm? Ecology, 74:1659-

1673.

LeRoy C.J., Hipp A.L., Lueders K., Follstad Shah J.J., Kominoski J.S., Ardón M., Dodds

W.K., Gessner M.O., Griffiths N.A., Lecerf A., Manning D.W.P., Sinsabaugh R.L., Webster

J.R. 2020. Plant phylogenetic history explains in-stream decomposition at a global scale. J

Ecol, 108:17-35.

McArdle B.H. 1988. The structural relationship: regression in biology. Can. J. Zool.,

66:2329-2339.

O'Meara B.C., Beaulieu J.M. 2016. Past, future, and present of state-dependent models of

diversification. Am. J. Bot., 103:792-795.

Osborne J.W., Waters E. 2002. Four assumptions of multiple regression that researchers

should always test. Pract. Assess. Res. Eval., 8:2.

Pagel M. 1999. Inferring the historical patterns of biological evolution. Nature, 401:877-884.

Paradis E., Schliep K. 2019. ape 5.0: an environment for modern phylogenetics and

evolutionary analyses in R. Bioinformatics, 35:526-528.

Pennell M.W., Harmon L.J. 2013. An integrative view of phylogenetic comparative

methods: connections to population genetics, community ecology, and paleobiology. Ann

NY Acad Sci, 1289:90-105.

Penone C., Davidson A.D., Shoemaker K.T., Di Marco M., Rondinini C., Brooks T.M.,

Young B.E., Graham C.H., Costa G.C. 2014. Imputation of missing data in life-history trait

datasets: which approach performs the best? Methods Ecol. Evol., 5:961-970.

Peres-Neto P.R., Jackson D.A., Somers K.M. 2005. How many principal components?

stopping rules for determining the number of non-trivial axes revisited. Comput. Statist. Data

Anal., 49:974-997.

Peres-Neto P.R., Legendre P. 2010. Estimating and controlling for spatial structure in the

study of ecological communities. Glob. Ecol. Biogeogr., 19:174-184.

Peres-Neto P.R., Leibold M.A., Dray S. 2012. Assessing the effects of spatial contingency

and environmental filtering on metacommunity phylogenetics. Ecology, 93:S14-S30.

Phillips R.F. 2002. Least absolute deviations estimation via the EM algorithm. Statistics and

Computing, 12:281-285.

Revell L.J. 2010. Phylogenetic signal and linear regression on species data. Methods Ecol.

Evol., 1:319-329.

Revell L.J. 2024. phytools 2.0: an updated R ecosystem for phylogenetic comparative

methods (and other things). PeerJ, 12:e16505.

Rohlf F.J. 2001. Comparative methods for the analysis of continuous variables: geometric

interpretations. Evolution, 55:2143-2160.

Safi K., Pettorelli N. 2010. Phylogenetic, spatial and environmental components of

extinction risk in carnivores. Glob. Ecol. Biogeogr., 19:352-362.

Slater G.J., Pennell M.W. 2014. Robust regression and posterior predictive simulation

increase power to detect early bursts of trait evolution. Syst. Biol., 63:293-308.

Venables W.N., Ripley B.D. 2002. Modern Applied Statistics with S. 4th ed., Springer.

Westoby M., Yates L., Holland B., Halliwell B. 2023. Phylogenetically conservative trait

correlation: Quantification and interpretation. J Ecol, 111:2105-2117.

Yu C., Yao W. 2017. Robust linear regression: A review and comparison. Commun. Stat.

Simul. Comput., 46:6261-6282.