Chromosome-level assembly of *Cucumis sativus cv.* 'Tokiwa' as a reference genome

of Japanese cucumber.

Takashi Seiko[1], Chiaki Muto[1], Koichiro Shimomura[2], Ryoichi Yano[3], Yoichi Kawazu[2], Mitsuhiro

Sugiyama[2], Kenji Kato[4], Norihiko Tomooka[1], Ken Naito[1*]

1. Research Center of Genetic Resources, National Agriculture and Food Research Organization, 2-

1-2 Kannondai, Tsukuba, Ibaraki 305-8602, Japan

2. Institute of Vegetable and Floriculture Science, National Agriculture and Food Research

Organization, 360 Kusawa, Ano, Tsu, Mie 514-2392, Japan

3. Research Center of Advanced Analysis, National Agriculture and Food Research Organization, 2-

1-2 Kannondai, Tsukuba, 305-8602, Japan

4. Graduate school of Environment, Life, Natural Science and Technology, Okayama University, 1-

1-1 Tsushimanaka, Kita, Okayama, Okayama 700-8530, Japan

*Corresponding author: Ken Naito

Research Center of Genetic Resources, National Agriculture and Food Research

Organization, 2-1-2 Kannondai, Tsukuba, Ibaraki 305-8602, Japan

knaito@affrc.go.jp

Keywords: Cucumis sativus, cucumber, whole genome sequencing, tandem repeat

Abstract

Cucumber is one of the most important vegetables in the Japanese market. To facilitate genomics-

based breeding, there is a demand for reference genome of Japanese cucumber. However, although

cucumber genome is relatively small, its assembly is a challenging issue because of tandem repeats

comprising ~30% (~100 Mbp) of the genome. To overcome, we deployed the Oxford nanopore

sequencing that produces long reads with N50 length of >30 kbp. With this technology we achieved a

chromosome-level assembly of cv 'Tokiwa', a founder line of Japanese cucumber represented with

the elongated fruit shape and high-crisp texture. Compared to the existing cucumber genomes, the

Tokiwa genome is 20% longer and annotated with 10% more genes. The assembly with nanopore long

reads also resolved tandem repeats spanning >100 kbp, demonstrating its strength in overcoming

repetitive sequences.

Introduction

Cucumber (*Cucumis sativus* L.) shares the 3rd place in the value production in the Japanese vegetable market, after tomato, and onion (https://www.usdajapan.org/wpusda/wp-content/uploads/2019/03/Japanese-Fresh-Vegetable-Market-Overview-2018_Osaka-ATO_Japan_12-21-2018-1.pdf). The Japanese cultivars of cucumber are represented with a long and straight fruit shape (Shimomura et al. 2016) and a crispy texture (Yoshioka et al., 2009), due to specific needs of the customers. Thus, breeding efforts of cucumber mainly target resistance to diseases including melon yellow spot virus and powdery mildews without spoiling the important traits described above. Heat tolerance has also become an important target because of the global warming. To facilitate breeding of such cultivars, we need to reconstruct a reference genome of Japanese cucumber and implement marker-assisted selection, allele mining via GWAS and prediction models for genomic selection.

However, efforts to sequence cucumber genomes have faced difficulties despite its relatively small size (~350 Mbp) (Huang et al., 2009, Yang et al., 2012, Li et al., 2019). The current reference genome is "Chinese long" inbred line 9930, which was sequenced with single molecule real-time (SMRT) sequencing technology (Li et al., 2019). The 9930 genome consists of 7 pseudomolecules covering 211 Mbp + 15 Mbp unanchored scaffolds, leaving ~120 Mbp sequences missing (Huang et al., 2009). More recently 11 more accessions have also been sequenced with the SMRT sequencing, but the assemblies were only slightly better than the 9930 genome (Li et al., 2022). The difficulty in assembling cucumber genome derives from long arrays of tandemly repeated sequences around centromeric regions (Ganal et al., 1986, Ganal and Hemleben, 1988).

To overcome the difficulties, increasing read length is the simplest solution. We adopted a recently-developed technology by Oxford Nanopore, which can generate ultra long reads spanning more than 100 kbp (Wang et al, 2021). Many plant genomes have already been sequenced with this technology, some achieving the NG50 length of more than 10 Mbp (Dumschott et al., 2020). Thus, we

expect it may improve *de novo* assembly of a cucumber genome.

Here, we sequenced and assembled the whole genome of a Japanese cucumber cultivar, cv.

'Tokiwa' (Fig. 1). 'Tokiwa' was developed from a cross between North China type 'Shimo-shirazu'

and South China type 'Shindome' around 1960. Given its Japanese favored fruit morphology and

texture and adaptability to both summer and fall cropping system, it has been a founder line of various

modern cultivars, which is now called as "Tokiwa group" (Fujieda, 2006). The Nanopore-only

assembly generated greatly improved contiguity and coverage compared to the 9930 genome,

providing a qualified knowledge-base for Japanese breeders.

Figure 1. Photographs of *Cucumis sativus* L. cv 'Tokiwa'. a. Plants growing in a greenhouse. b. A green fresh fruit. c. Mature fruites. d. Seeds.

Materials and methods

Plant material and DNA extraction

The cucumber cultivar 'Tokiwa' was provided by National Genebank Program in Research Center of Genetic Resources, NARO. We extracted genomic DNA from 3 g of young leaves using Nucleobond HMW DNA kit (MACHEREY-NAGEL GmbH & Co. KG, Düren, Germany). We basically followed the DNA extraction protocol provided by the manufacturer, but we extended the lysis step up to 3 h instead of 30 min. In the precipitation + washing step with isopropanol and 70% (v/v) ethanol, we did not centrifuge but directly transfer the precipitated DNA by pipetting.

We also extracted RNA from the leaves for gene annotation used by NucleoSpin RNA Plant kit (MACHEREY-NAGEL GmbH & Co. KG, Düren, Germany).

Sequence of DNA and RNA

For long-read sequencing, we followed the protocol as described by Naito (2023). First, we size-selected the genomic DNA using Short Read Eliminatior XL kit (Pacific Biosciences of California, Inc., California,USA) to remove the DNA fragments <10 kbp. Second, we put 3 μg of the size-selected DNA for end-repair using NEBNext FFPE DNA Repair Mix (NEW INGLAND BioLabs inc., Massachusetts, USA) and NEBNext Ultra II End Repair/dA-Tailing Module. Third, we ligated the end-repaired DNA with the sequencing adaptor in SQK-LSK109 (Oxford Nanopore Technologies KK, Tokyo, Japan) using NEBNext Quick Ligation Module (NEW INGLAND BioLabs inc., Massachusetts, USA). In each step, we purified DNA using AMPure XP (BECKMAN COULTER inc., California, USA). The prepared library was then loaded to R9.4.1 Flowcell (Oxford Nanopore Technologies KK, Tokyo, Japan) and was run in PromethION24 (Oxford Nanopore Technologies PLC, Oxford, UK) for 10 h. We also obtained short-reads for polishing the draft assembly with Illumina Hiseq 4000, which was provided as a customer service by GeneBay, Inc. (Yokohama, Japan). For

scaffolding, we obtained optical mapping data with Bionano Sahyr, provided as a customer service by

AS ONE corp. (Osaka, Japan). We also sequenced the RNA with Hiseq4000 (Illumina Inc., San Diego,

USA) for gene annotation, provided as a customer service by GeneBay Inc. (Yokohama, Japan). All

the data is uploaded to the Sequence Read Archive in NCBI website, with BioSample ID

SAMN40100976 in BioProject ID PRJNA1080095.

Genome assembly and scaffolding

We also followed the protocol described by Naito (2023) for genome assembly. Fist, we used necat-

0.0.1 (Chen et al., 2021) with default parameters except "PREP_OUTPUT_COVERAGE=60" and

"CNS_OUTPUT_COVERAGE=40". The draft contigs were then polished twice with racon-1.4.3

(Vaser et al., 2017) with options of "-m 8 -x -6 -g -8 -w 500" and medaka-1.0.3

(https://github.com/nanoporetech/medaka) with options of "-g -m r941_prom_high_g360". To check

misassemblies, we first aligned the polished contigs to the 9930 genome by D-genies (Cabanettes and

Klopp 2018) to visualize conflicted sites. We then mapped the long-reads with minimap2-2.14 (Li,

2018) and manually checked the conflicted sites. If the mapped reads were discontinuous as presented

by Sakai et al. (2015), we broke the contig at the site. The corrected contigs were further polished with

Illumina short-reads twice with Hypo-1.0.3 (https://github.com/kensung-lab/hypo) with default

parameters. The Hypo-polished contigs were then run with purge_haplotigs-1.1.1 (Roach et al., 2018)

to remove low-coverage contigs and those derived from heterozygous regions.

After assembling, polishing and filtering the contigs, we obtained optical mapping data

provided as a customer service by AS ONE, corp. (Osaka, Japan). High-molecule genomic DNA was

extracted from young leaves with the Plant DNA Isolation Kit (Bionano Genomics, San Diego, CA,

USA) according to Bionano Prep Plant Tissue DNA Isolation Base Protocol. The DNA was treated

with DLE-1 nickase, labelled with DLS DNA Labeling Kit (Bionano Genomics), and run on the

Saphyr Genome Mapping Instrument (Bionano Genomics). The obtained Saphyr reads were

assembled and then merged with the draft assembly to generate hybrid scaffolds sequences with

Bionano Solve (Bionano Genomics) with the default parameters. Tha Bionano data is also available

from NCBI website with BioProject ID: PRJNA1098469.

Finally, pseudomolecules were reconstructed by anchoring the hybrid scaffolds to the

genetic map, which was previously constructed (Sugiyama et al., 2016).

To compare the assembly with 9930 genome, we made a whole genome alignment by

minimap2 and ran the output with D-genies to generate a dotplot.

Assembling organelle genomes

Given there are hundreds of chloroplasts and tens of mitochondria in a single cell, the coverage depth

is usually too high for assembling the organellar genomes. Thus, we had to extract and downsample

the long reads derived from the organellar genomes.

To do so, we first identified the organellar contigs by BLASTing the reference genomes of

chloroplast (Pląder et al., 2007) and mitochondrion (Alverson et al., 2011) as query to the draft

assembly. Then we mapped the long reads to the draft assembly and recovered the reads that were

mapped to the organellar contigs. We then downsampled the recovered reads to ~50x coverage and

ran Flye-2.9 (Kolmogorov et al., 2019). The assembled sequences were then polished with long reads

with medaka.

Gene predictions

Briefly, we predicted gene structures using Braker-2.0 (Brůna et al., 2021) using protein sequences

and GeMoMa-1.9 (Keilwagen et al., 2018) using our own RNA-seq data and the coding sequences of

9930 genome.

142   First, we softmasked the Tokiwa genome by running RepeatModeler-2.0.5

143 (https://www.repeatmasker.org/RepeatModeler/) and RepeatMasker-4.0.9

144 (https://www.repeatmasker.org/RepeatMasker/) with default parameters. Second, we downloaded the

145 sequences of plant proteins from OrthoDB (Kuznetsov et al., 2022) as well as the license for

146 GeneMark-EP+ (Brůna et al., 2020), which was required for running Braker-2.0. Third, we ran braker-

147 2.0 on the softmasked genome with an option of "--softmasking".

148   Before running GeMoMa, we removed noisy reads from the RNA-seq data. To do so, we *de*

149 *novo* assembled the RNA-seq data with Trinity-2.13.0 (Grabherr et al., 2011) with default parameters.

150 Then we mapped the RNA-seq reads to the assembled transcript sequences using HISAT-2.2.1 (Kim

151 et al., 2019) and recovered only concordantly mapped reads. The recovered RNA-seq reads were then

152 mapped to Tokiwa genome by HISAT-2.2.1 with an option of "--max-intronlen 15000", which was

153 used for intron prediction by GeMoMa. Then we ran GeMoMa using 9930 genome and gene

154 annotations as reference, with parameters for unstranded RNA-seq data. After running the annotation

155 tools, we merged the outputs of GeMoMa and Braker, by filtering the overlapping annotations. Then

156 we ran EnTAP-0.10.8 (Hart et al., 2020) with default parameters to functionally annotate the predicted

157 genes. The final annotations retained those identified as "Viridiplantae" and "Eukaryotes" for

158 Taxonomy scope.

159   We also reannotated the 9930 genome and those of the 11 accessions assembled by Li et al.

160 (2022). The 11 genome sequences were downloaded from the website of NCBI datasets. The

161 downloaded genomes were annotated by GeMoMa-1.9 with the predicted gene models of 'Tokiwa' as

162 reference. JCVI MCscan (https://github.com/tanghaibao/jcvi/wiki/MCscan-(Python-version)) was

163 used to predict synteny block between each genome.

164   The predicted genes were evaluated with eudicots_odb10 (Kriventseva et al., 2018) by

165 BUSCO-5.5 (Manni et al., 2021).

Phylogenetic analysis

To place the Tokiwa genome in a phylogenetic position, we first ran orthofinder-2.5.5 (Emms et al.,

2019) on the protein sequences of the 12 accessions, with "-M msa" option. Then we reconstructed a

maximum likelihood tree from the ouput mafft file of single copy orthologs by running iqtree-2.2.6

with options of "-m MFP -bb 1000 -alrt 1000 -nt AUTO".

Repeat annotation

To identify sequences derived from transposable elements, we ran EDTA-2.2.2 (Ou et al., 2019) with

the predicted coding sequences of 'Tokiwa' in a sensitive mode. To identify tandem repeats, we ran

trf-4.0.9 with parameters suggested by the official protocol.

Results

De novo *Assembly of Tokiwa genome*

We have successfully reconstructed the pseudomolecules of the 7 chromosomes in Tokiwa genome,

covering 50 Mbp more than the 9930 genome (Table 1, Fig. 2).

Using the Oxford Nanopore sequencer, we obtained 34.2 Gbp of long reads with the N50

length of 33.2 kbp (Supplemental Table 1). The draft assembly with NECAT was ~300 Mbp sequences

in 645 contigs, with the N50 length of 12.1 Mbp and the maximum length of 20.1 Mbp (Table 1).

Purge_haplotigs identified 260 contigs as repetitive or low-coverage junk and thus retained 385

contigs covering 278.7 Mbp. The following optical mapping scaffolded only 71 contigs but improved

the N50 length to 23.4 Mbp and the maximum length to 45.1 Mbp. Finally, we anchored the 14

scaffolds to the genetic linkage map (Sugiyama et al., 2015) and reconstructed the pseudomolecules

of the 7 chromosomes (Fig. 2). After removing the scaffolds/contigs derived from organelle genome

and possible contaminations, 23.6 Mbp in 130 scaffolds/contigs remained unplaced. All the seven

pseudomolecules contained telomeric repeats in both ends.

We also separately assembled the genome sequences of the chloroplast and mitochondrion.

The chloroplast genome was successfully assembled into a single circular genome. However, due to

the highly recombining feature of mitochondrion, we could not circularize the mitochondrial genome

and ended up with 6 contigs (Supplemental Table 2).

Table 1. The statistics of genome assemblies.

	NECAT	Purge_haplotigs	BioNano	Tokiwa v1.0	9930 v3
Num	645	385	314	7	7
Sum	300,877,148	278,732,282	299,675,636	257,281,677	210,974,704
Min	208	4,095	4,095	27,903,525	22,466,726
Ave	466,476	723,980	954,381	36,754,525	30,139,243
Max	20,967,667	20,967,667	45,117,507	45,117,507	40,877,379
N50	12,103,241	12,256,096	23,376,839	36,771,724	31,913,682
GAP	-	-	20,923,098	15,419,875	36,170
Total ATGC	-	-	278,752,538	241,861,802	210,938,534
Non-placed scaffolds	-	-	-	130	81
Non-placed bps	-	-	-	23,587,426	15,666,262
Protein coding genes				27,905	24,317

High duplication rate in the Tokiwa genome than others

The pseudomolecules of Tokiwa genome reached 257 Mbp, which was >40 Mbp longer than those of

the 9930 genome (Fig. 2B). Even though the extended sequences were mostly in centromeric, highly

repetitive regions, gene coding sequences were also present (Fig. 2A). Although we used the Tokiwa's

gene models to reannotate the 9930 and other genomes for fair comparison, the Tokiwa genome

contained ~3,000 more genes than others (Table 2). The BUSCO scores were not greatly different

among them, but the predicted gene models in the Tokiwa genome contained ~10 times more

duplicated BUSCO genes (Table 2).

Figure 2. Overview of Taokiwa genome. A. Circos plot indicating densities of, from the outer to inner tracks, protein coding sequences, LTR retrotransposons, DNA transposable elements, and tandem repeats. B. Coliniarity between the physical positions and the genetic linkage map of the Tokiwa genome. White and blue boxes indicate the physical and genetic map, respectively. Dashed lines indicate the corresponding positions. C. Dotplot of a whole genome alignment of 9930 and Tokiwa. *x*- and *y*-axes indicate Chromosome numbers. Cultivar names are indicated above the *x*-axis and right to the *y*-axis.

The genome annotation revealed highly conserved synteny between the cucumber genomes

including wild accessions, except several inversions in Chromosomes 4, 5, and 7 of Cuc64 and W8,

which were also revealed by genome alignment-based analysis (Li et al., 2022). (Fig. 3). However, it

also revealed that the genes present in the newly assembled regions of the Tokiwa genomes were

mostly the duplicated copies of those in other regions (Fig. 3). The high rate of duplicated genes was

also noted when we extracted single-copy orthologs for phylogenetic analysis. Without the Tokiwa

genome, there were 17,619 single-copy orthologs among the 11 genomes. However, with the Tokiwa

genome, the number decreased to 13,542. This was because the ~4,000 genes, present as single copy

in all the Pacbio-sequenced genomes, had duplicated copies in the Tokiwa genome.

Table 2. BUSCO scores of the Tokiwa other genomes

	Complete		Single		Duplicate		Fragmented		Missing	
	Number	(%)	Number	(%)	Number	(%)	Number	(%)	Number	(%)
Tokiwa	2,281	(98.1)	1,797	(77.3)	484	(20.8)	6	(0.3)	39	(1.6)
9930	2,268	(97.5)	2,217	(95.3)	51	(2.2)	12	(0.5)	46	(2.0)
9110gt	2,250	(96.7)	2,213	(95.1)	37	(1.6)	11	(0.5)	65	(2.8)
Cu2	2,266	(97.4)	2,212	(95.1)	54	(2.3)	10	(0.4)	50	(2.2)
Cuc37	2,262	(97.3)	2,221	(95.5)	41	(1.8)	11	(0.5)	53	(2.2)
Cuc64	2,273	(97.7)	2,231	(95.9)	42	(1.8)	8	(0.9)	45	(2.0)
Cuc80	2,272	(97.7)	2,225	(95.7)	47	(2.0)	8	(0.3)	46	(2.0)
Gy14	2,256	(97.0)	2,212	(95.1)	44	(1.9)	10	(0.4)	60	(2.6)
Hx14	2,251	(96.7)	2,213	(95.1)	38	(1.6)	11	(0.5)	64	(2.8)
Hx117	2,233	(96.0)	2,189	(94.1)	44	(1.9)	15	(0.6)	78	(3.4)
W4	2,267	(97.4)	2,224	(95.6)	43	(1.8)	9	(0.4)	50	(2.2)
W8	2,251	(96.7)	2,206	(94.8)	45	(1.9)	13	(0.6)	62	(2.7)
XMTC	2,259	(97.1)	2,217	(95.3)	42	(1.8)	12	(0.5)	55	(2.4)

238   With the 13,542 single-copy orthologs, we reconstructed the phylogenetic tree (Fig. 3). Our

239   phylogenetic tree grouped the cultivars (9110gt, 9930, XMTC, Tokiwa, Cu2, Cuc37, Gy14, Cuc80,

240   Hx117 and HX14) and the wild cucumbers (*C. sativus* var. *hardwickii*) (W4, Cuc64 and W8) into

241   monophyletic clades, respectively. 'Tokiwa' formed an East-Asian group with the Chinese cultivars

242   9110gt, 9930, XMTC, and Cu2 (Fig. 3).

Figure 3. Phylogenetic tree and synteny plot of the 12 chromosome-level genome assemblies. Gray lines indicate synteny blocks between the genomes. Light red lines indicate pseudo synteny blocks between the newly assembled regions of the Tokiwa genome and XMTC or Cu2. Bootstrap values were 100 in all the nodes.

Distribution of transposable elements in Tokiwa genome

Because of the relatively small size of the genome, transposable elements shared only 23.9% of the

Tokiwa genome (Supplemental Table 2). Overall, DNA transposons were more abundant than LTR

retrotransposons were. The most abundant was Mutator family, comprising nearly 10% (22.8 Mbp) of

the Tokiwa genome, followed by Gypsy, Copia, and CACTA.

In the Tokiwa genome, the distribution of LTR retrotransposons and DNA transposons were

basically exclusive to each other (Fig. 2A). While the DNA transposons were highly condensed near

centromeric and the telomeric regions, the LTR retrotransposons were more broadly distributed across

the pericentromeric regions.

Figure 4. Distribution of unit size, repeat count and total length of the tandem repeats in the Tokiwa genome. A. Scatter plot of unit size (*x*-axis) and repeat count (*y*-axis) of each tandem repeat. B. Scatter plot of unit size (*x*-axis) and the total length of each tandem repeat (*y*-axis). C. Schematic of a Mutator-like element containing tandem repeats. Triangles indicate the units of each tandem repeat.

Tandem repeat in Tokiwa genome

As cucumber genome is peculiar for harboring long stretches of tandem repeats, we also surveyed the

Tokiwa genome with tandem repeat finder (Benson 1999). The survey revealed that tandem repeats

comprise 35.2 Mbp (12.6%) of the Tokiwa genome.

Although the size of the repeat unit varied, those of ~180 bp, ~360 bp, ~540 bp and ~720 bp

had extremely high copy numbers as well as those <10 bp (Fig. 4). The longest stretch of the tandem

repeats was that of 712.6 kbp on Chromosome 4, where a unit of 531 bp sequence was tandemly

repeated by 1,331 times (Table 3). Other chromosomes also harbored tandem repeats spanning >100

kbp (Table. 3), which could be even longer, as many of them ended with gaps.

Interestingly, the long stretches of tandem repeats (>10 kbp) were often overlapped with

annotations of Mutator and CACTA transposons (Fig 2). In addition, the Mutator and CACTA

elements identified by EDTA often contained the tandem repeats as their internal sequences (Fig. 4C).

Namely, the tandem repeats did not cover the full length of the transposable elements. Thus, increase

in tandem repeats did not necessarily occur via TE amplifications.

Table 3. The top 10 longest tandem repeats in the Tokiwa Genome.

Chromosome	Start	End	Unit size	Repeat Count	Length
Chromosome4	12,049,307	12,761,914	531	1331.1	712,608
Chromosome3	25,304,414	25,792,696	531	915.6	488,283
Chromosome5	109,973	578,952	360	1301	468,980
Chromosome1	14,073,064	14,531,408	501	903.2	458,345
Chromosome7	7,938,677	8,366,883	354	1203.4	428,207
Chromosome7	8,006,713	8,377,207	531	693.4	370,495
Chromosome7	27,538,573	27,895,514	360	989.3	356,942
Chromosome7	10,064,253	10,414,004	531	654.1	349,752
Chromosome5	37,052,513	37,392,146	720	463.2	334,450

Discussion

Here we present the most contiguous assembly of the cucumber genome, owing to the long reads by

Oxford Nanopore. The assembled Tokiwa genome is ~20% longer and contains 10% more gene coding

sequences compared to the 9930 genome, or other existing assemblies (Table 2, Figs. 2, 3). The

annotation predicted more duplicated genes, less fragmented genes, and much more repeat sequences

including tandem repeats. As 'Tokiwa' is a founder line of modern Japanese cultivars, the highly

contiguous and accurate genome could facilitate breeding for urgent and long-term demands in

Japanese market.

Our results have demonstrated the importance of read length to overcome the repetitive

sequences in *de novo* assembly. The nanopore sequencer produced long reads with N50 length of 33.2

kbp, which was >3 time longer than those used for assembling the 9930 genome (Supplementary Table

1). Such long reads have enabled us to resolve and assemble long stretches of tandem repeats,

including the one over 700 kbp (Table 3).

However, we should also note that we could not assemble centromeric regions without the

optical mapping by Bionano. Although the draft assembly already generated contigs containing long

stretches of tandem repeats, such contigs could not be directly anchored to the linkage map because

there were no genetic markers in such highly repetitive regions. Thus, optical mapping was the only

way to place the repeat-rich contigs in centromeric or pericentromeric regions.

The newly resolved repeat sequences also give insight into the relationship between the

DNA transposons and the tandem repeats. The distribution of the unit size in the tandem repeats

indicated that those with high copy numbers are overrepresented with specific unit sizes. This led us

to suspect that such tandem repeats could arise from cis-transposition of transposable elements.

However, many of the tandem repeats are actually "internal" sequences of transposons. Namely, one

single copy of a transposon could be very long, containing a long stretch of tandem repeats within its

sequence (Fig. 4). Thus, the tandem repeats could have increased its copy number via other

mechanisms than transposition, such as illegitimate recombination or double strand break repair.

Although the long reads by Oxford Nanopore greatly facilitated assembling highly repetitive

regions, the Tokiwa genome is still far from gapless assembly. Given the estimated size of cucumber

genome is ~350 Mbp, there remains ~70 Mbp sequences unassembled. As the Oxford Nanopore have

greatly improved the accuracy of raw reads (Zhao et al., 2023), new assemblers optimized for lower

error rate would further facilitate resolving the tandem repeats.

To conclude, this study presented a *de novo* assembly of Tokiwa genome using long reads

of Oxford Nanopore, which could serve as a reference of Japanese cucumber. We hope the Tokiwa

genome would be a fundamental knowledge-base for future breeding of Japanese cultivars.

Funding

This study was supported by Public/Private R&D Investment Strategic Expansion PrograM

(PRISM) of Japan.

Author's contributions

KN, NT and **KK** conceived the study.

TS and **CM** performed sequencing and assembly.

CM, **KS**, **YK**, **MS**, **RY** and **KN** fixed misassemblies and performed scaffolding.

TS and **KN** performed genome annotation and data analysis.

TS, KS and **KN** wrote the paper.

All the authors read and approved the final manuscript.

Data availability

All the sequence data were deposited at Biosample IDs SAMN40100976, under Bioproject

PRJNA1080095.

Literature Cited

Alverson, A.J., D.W. Rice, S. Dickinson, K. Barry and J.D. Palmer (2011) Origins and

Recombination of the Bacterial-Sized Multichromosomal Mitochondrial Genome of

Cucumber. The Plant Cell 23: 2499-2513.

Benson, G. (1999) Tandem repeats finder: a program to analyze DNA sequences. Nucleic Acids

Research 27: 573-580.

Brůna, T., A. Lomsadze and M. Borodovsky (2020) GeneMark-EP+: eukaryotic gene prediction

with self-training in the space of genes and proteins. NAR Genomics and

Bioinformatics 2.

Brůna, T., K.J. Hoff, A. Lomsadze, M. Stanke and M. Borodovsky (2021) BRAKER2:

automatic eukaryotic genome annotation with GeneMark-EP+ and AUGUSTUS

supported by a protein database. NAR Genomics and Bioinformatics 3.

Cabanettes, F. and C. Klopp (2018) D-GENIES: dot plot large genomes in an interactive,

efficient and simple way. PeerJ 6: e4958.

Chen, Y., F. Nie, S.-Q. Xie, Y.-F. Zheng, Q. Dai, T. Bray, Y.-X. Wang, J.-F. Xing, Z.-J. Huang,

D.-P. Wang, *et al.* (2021) Efficient assembly of nanopore reads via highly accurate and

intact error correction. Nature Communications 12: 60.

Dumschott, K., M.H.-W. Schmidt, H.S. Chawla, R. Snowdon and B. Usadel (2020) Oxford

Nanopore sequencing: new opportunities for plant genomics? Journal of Experimental

Botany 71: 5313-5322.

Emms, D.M. and S. Kelly (2019) OrthoFinder: phylogenetic orthology inference for

comparative genomics. Genome Biology 20: 238.

Fujieda, K. 2006. Cucumber. p. 136-142. In: Organizing Committee (ed.). Horticulture in Japan

2006. Nakanishi Printing, Kyoto.

Ganal, M., I. Riede and V. Hemleben (1986) Organization and sequence analysis of two related

satellite DNAs in cucumber (Cucumis sativus L.). Journal of Molecular Evolution 23:

23-30.

Ganal, M. and V. Hemleben (1988) Insertion and amplification of a DNA sequence in satellite

DNA of Cucumis sativus L. (cucumber). Theoretical and Applied Genetics 75: 357-361.

Grabherr, M.G., B.J. Haas, M. Yassour, J.Z. Levin, D.A. Thompson, I. Amit, X. Adiconis, L.

Fan, R. Raychowdhury, Q. Zeng, *et al.* (2011) Full-length transcriptome assembly from

RNA-Seq data without a reference genome. Nature Biotechnology 29: 644-652.

Hart, A.J., S. Ginzburg, M. Xu, C.R. Fisher, N. Rahmatpour, J.B. Mitton, R. Paul and J.L.

Wegrzyn (2020) EnTAP: Bringing faster and smarter functional annotation to non-

model eukaryotic transcriptomes. Molecular Ecology Resources 20: 591-604.

Huang, S., R. Li, Z. Zhang, L. Li, X. Gu, W. Fan, W.J. Lucas, X. Wang, B. Xie, P. Ni, *et al.*

(2009) The genome of the cucumber, Cucumis sativus L. Nature Genetics 41: 1275-

1281.

Keilwagen, J., F. Hartung, M. Paulini, S.O. Twardziok and J. Grau (2018) Combining RNA-seq

data and homology-based gene prediction for plants, animals and fungi. BMC

Bioinformatics 19: 189.

Kim, D., J.M. Paggi, C. Park, C. Bennett and S.L. Salzberg (2019) Graph-based genome

alignment and genotyping with HISAT2 and HISAT-genotype. Nature Biotechnology

37: 907-915.

Kolmogorov, M., J. Yuan, Y. Lin and P.A. Pevzner (2019) Assembly of long, error-prone reads

using repeat graphs. Nature Biotechnology 37: 540-546.

Kriventseva, E.V., D. Kuznetsov, F. Tegenfeldt, M. Manni, R. Dias, F.A. Simão and E.M.

Zdobnov (2018) OrthoDB v10: sampling the diversity of animal, plant, fungal, protist,

bacterial and viral genomes for evolutionary and functional annotations of orthologs.

Nucleic Acids Research 47: D807-D811.

Kuznetsov, D., F. Tegenfeldt, M. Manni, M. Seppey, M. Berkeley, Evgenia V. Kriventseva and

E.M. Zdobnov (2022) OrthoDB v11: annotation of orthologs in the widest sampling of

organismal diversity. Nucleic Acids Research 51: D445-D451.

Li, H. (2018) Minimap2: pairwise alignment for nucleotide sequences. Bioinformatics 34: 3094-

3100.

Li, Q., H. Li, W. Huang, Y. Xu, Q. Zhou, S. Wang, J. Ruan, S. Huang and Z. Zhang (2019) A

chromosome-scale genome assembly of cucumber (Cucumis sativus L.). GigaScience 8.

Li, H., S. Wang, S. Chai, Z. Yang, Q. Zhang, H. Xin, Y. Xu, S. Lin, X. Chen, Z. Yao, *et al.*

(2022) Graph-based pan-genome reveals structural and sequence variations related to

agronomic traits and domestication in cucumber. Nature Communications 13: 682.

Manni, M., M.R. Berkeley, M. Seppey, F.A. Simão and E.M. Zdobnov (2021) BUSCO Update:

Novel and Streamlined Workflows along with Broader and Deeper Phylogenetic

Coverage for Scoring of Eukaryotic, Prokaryotic, and Viral Genomes. Molecular

Biology and Evolution 38: 4647-4654.

Nakamura, T., K.D. Yamada, K. Tomii and K. Katoh (2018) Parallelization of MAFFT for

large-scale multiple sequence alignments. Bioinformatics 34: 2490-2492.

Ou, S., W. Su, Y. Liao, K. Chougule, J.R.A. Agda, A.J. Hellinga, C.S.B. Lugo, T.A. Elliott, D.

Ware, T. Peterson, *et al.* (2019) Benchmarking transposable element annotation

methods for creation of a streamlined, comprehensive pipeline. Genome Biology 20:

275.

Pląder, W., Y. Yukawa, M. Sugiura and S. Malepszy (2007) The complete structure of the

cucumber (Cucumis sativus L.) chloroplast genome: Its composition and comparative

analysis. Cellular and Molecular Biology Letters 12: 584-594.

Ritu, K., C. Joshua and S. Wing-Kin (2019) HyPo: Super Fast & Accurate Polisher

for Long Read Genome Assemblies. bioRxiv: 2019.12.19.882506.

Roach, M.J., S.A. Schmidt and A.R. Borneman (2018) Purge Haplotigs: allelic contig

reassignment for third-gen diploid genome assemblies. BMC Bioinformatics 19: 460.

Sakai, H., K. Naito, E. Ogiso-Tanaka, Y. Takahashi, K. Iseki, C. Muto, K. Satou, K. Teruya, A.

429    Shiroma, M. Shimoji, *et al.* (2015) The power of single molecule real-time sequencing

430    technology in the de novo assembly of a eukaryotic genome. Scientific Reports 5:

431    16780.

432 Shimomura, K., H. Horie, M. Sugiyama, Y. Kawazu and Y. Yoshioka (2016) Quantitative

433    evaluation of cucumber fruit texture and shape traits reveals extensive diversity and

434    differentiation. Scientia Horticulturae 199: 133-141.

435 Sugiyama, M., Y. Kawazu, N. Fukino, Y. Yoshioka, K. Shimomura, Y. Sakata and M. Okuda

436    (2015) Mapping of quantitative trait loci for Melon yellow spot virus resistance in

437    cucumber (Cucumis sativus L.). Euphytica 205: 615-625.

438 Vaser, R., I. Sovic, N. Nagarajan and M. Sikic (2017) Fast and accurate de novo genome

439    assembly from long uncorrected reads. Genome Research.

440 Wang, Y., Y. Zhao, A. Bollas, Y. Wang and K.F. Au (2021) Nanopore sequencing technology,

441    bioinformatics and applications. Nature Biotechnology 39: 1348-1365.

442 Yang, L., D.-H. Koo, Y. Li, X. Zhang, F. Luan, M.J. Havey, J. Jiang and Y. Weng (2012)

443    Chromosome rearrangements during domestication of cucumber as revealed by high-

444    density genetic mapping and draft genome assembly. The Plant Journal 71: 895-906.

445 Yoshioka, Y., H. Horie, M. Sugiyama and Y. Sakata (2009) Quantifying cucumber fruit

446    crispness by mechanical measurement. Breeding Science 59: 139-147.

447 Zhao, W., W. Zeng, B. Pang, M. Luo, Y. Peng, J. Xu, B. Kan, Z. Li and X. Lu (2023) Oxford

448    nanopore long-read sequencing enables the generation of complete bacterial and plasmid

449    genomes without short-read sequencing. Frontiers in Microbiology 14.

 Figure legends

 Figure 1. Photographs of *Cucumis sativus* L. cv ʹTokiwaʹ. a. Plants growing in a greenhouse. b. A

 green fresh fruit. c. Mature fruites. d. Seeds.

 Figure 2. Overview of Taokiwa genome. A. Circos plot indicating densities of, from the outer to inner

 tracks, protein coding sequences, LTR retrotransposons, DNA transposable elements, and tandem

 repeats. B. Coliniarity between the physical positions and the genetic linkage map of the Tokiwa

 genome. White and blue boxes indicate the physical and genetic map, respectively. Dashed lines

 indicate the corresponding positions.   C. Dotplot of a whole genome alignment of 9930 and Tokiwa.

 x- and *y*-axes indicate Chromosome numbers. Cultivar names are indicated above the *x*-axis and right

 to the *y*-axis.

 Figure 3. Phylogenetic tree and synteny plot of the 12 chromosome-level genome assemblies. Gray

 lines indicate synteny blocks between the genomes. Light red lines indicate pseudo synteny blocks

 between the newly assembled regions of the Tokiwa genome and XMTC or Cu2. Bootstrap values

 were 100 in all the nodes.

 Figure 4. Distribution of unit size, repeat count and total length of the tandem repeats in the Tokiwa

 genome. A. Scatter plot of unit size (*x*-axis) and repeat count (*y*-axis) of each tandem repeat. B. Scatter

 plot of unit size (*x*-axis) and the total length of each tandem repeat (*y*-axis). C. Schematic of a Mutator-

 like element containing tandem repeats. Triangles indicate the units of each tandem repeat.