Structural basis of iron piracy by human gut *Bacteroides*

Short title: *Bacteroides* iron piracy

Augustinas Silale[1]*, Yung Li Soo[1], Hannah Mark[1], Rachel N. Motz[2], Arnaud Baslé[1], Elizabeth M. Nolan[2], Bert van den Berg[1]*

[1]Biosciences Institute, Newcastle University, Framlington Place, Newcastle upon Tyne, NE2 4HH, United Kingdom.
[2]Department of Chemistry, Massachusetts Institute of Technology, Cambridge, Massachusetts 02139, United States.

*Correspondence to: augustinas.silale@newcastle.ac.uk
                                      bert.van-den-berg@newcastle.ac.uk

Abstract

Iron is an essential element that can be growth-limiting in microbial communities, particularly those present within host organisms. To acquire iron, many bacteria secrete siderophores, secondary metabolites that chelate ferric iron. These iron chelates can be transported back into the cell via TonB-dependent transporters in the outer membrane, followed by intracellular liberation of the iron. Pathogenic *Escherichia coli* and *Salmonella* produce siderophores during gut infection. In response to iron starvation, the human gut symbiont *Bacteroides thetaiotaomicron* upregulates an iron piracy system, XusABC, which steals iron-bound siderophores from the invading pathogens. Here, we investigated the molecular details of xenosiderophore uptake across the outer membrane by the XusAB complex. Our crystal and cryogenic electron microscopy structures explain how the XusB lipoprotein recognises iron-bound xenosiderophores and passes them on to the XusA TonB-dependent transporter. Moreover, we show that Xus homologues can transport a variety of siderophores with different iron-chelating functional groups.

Introduction

Iron is essential for most organisms, including almost all bacteria. It is a protein cofactor required for enzymatic reactions and electron transport during various cellular processes. While iron is one of the most abundant transition metals on Earth, its bioavailability is limited due to the extremely low solubility ($\sim 10^{-18}$ M) of its predominant ferric (Fe^{3+}) form at physiological pH (*1*). Many Gram-negative bacteria release siderophores, iron-chelating secondary metabolites, into the environment which, once bound to ferric iron, can be transported back into the cell via TonB-dependent transporters in the outer membrane (OM) (*2*). A classic example is enterobactin (Ent), which is produced by Enterobacteriaceae and consists of three iron-chelating catecholate groups connected via amide linkers to a cyclic triserine lactone (*3, 4*). Ferric enterobactin (FeEnt) is taken up by the FepA and IroN transporters in *E. coli* and *Salmonella* (*5–7*), and iron is liberated from the FeEnt complex inside the cell (*8, 9*). Siderophore-mediated iron scavenging is important for pathogens during infection, when the host restricts iron availability to starve invading bacteria via nutritional immunity (*10–12*). Additionally, secreted siderophores are not necessarily taken up by the same bacterium that produced them, resulting in complex iron availability-dependent interactions in microbial communities (*13*).

The gastrointestinal tract is home to a diverse community of microorganisms that cooperate and compete for available nutrients both with the host and between themselves (*14, 15*).

Depriving the gut microbiota of iron eventually results in irreversible structural changes in gut
microbial communities (*16*, *17*). Bacteroidota, the dominant phylum of diderm bacteria in the
distal gut, are not known to produce siderophores. *Bacteroides thetaiotaomicron* (*B. theta*) has
been shown to prefer heme as an iron source over soluble ferrous iron (*18*). *B. theta* likely
acquires heme from the diet (*19*) as well as from dead intestinal cells and dead bacteria. Iron-
bound siderophores produced by commensal bacteria and fungi as well as various pathogens
are another source of bioavailable iron for gut bacteria that have transporters to take them up
(*20*). It was recently shown that during *Salmonella* infection in the mouse gut, when iron
availability becomes limiting, *B. theta* upregulates a novel iron uptake system that steals iron-
bound siderophores produced by the pathogenic *Salmonella* (*21*). This xenosiderophore (i.e.,
foreign siderophore) utilization system (Xus) consists of the TonB-dependent transporter
XusA, the surface-exposed lipoprotein XusB and the PepSY domain-containing inner
membrane protein XusC (*21*, *22*). The XusABC system provides resilience to *B. theta* during
infection, but XusB secreted by *B. theta* and bound to xenosiderophores also acts as an iron
reservoir for the pathogen (*21*, *22*). It is therefore important to understand the molecular details
of this iron piracy system to obtain a deeper understanding of pathogen-symbiont interactions
inside the gut.

Here, we present crystal structures of the apo- and xenosiderophore-bound XusB lipoproteins
from several Bacteroidota species, which reveal the mechanism of xenosiderophore capture.
We also use single particle cryo-EM to investigate how XusB interacts with the XusA TonB-
dependent transporter. In vitro binding studies and structural bioinformatics analyses reveal
distinct subclasses of xenosiderophore utilization systems in Bacteroidota. Together, our results
provide mechanistic insights into xenosiderophore uptake across the OM of the gut commensal
B. theta and the pathobiont *Bacteroides fragilis*.

Results

Crystal structures of apo and xenosiderophore-bound *B. theta* XusB
We expressed recombinant XusB of *B. theta* (BtXusB; UniProt accession Q8A622) lacking the
signal sequence and the lipid anchor cysteine in *E. coli* BL21(DE3) and determined its crystal
structure using data to 1.56 Å resolution (Fig. 1A and B, Table S1). BtXusB has a seven-bladed
β-propeller fold with a distinct 15-residue loop, which we termed the hook, inserted into the
β2 blade and protruding outwards from the β-propeller (Fig. 1A). The β-propeller pocket (Fig.
1B) has been suggested to bind FeEnt based on recent computational docking (*22*).

We co-crystallized recombinant BtXusB with FeEnt and determined the crystal structure of the
complex to 1.50 Å. BtXusB binds FeEnt via four loops (Fig. 1C), including the hook.
Interestingly, FeEnt binds off the β-propeller pocket rather than in it, and the distance to the
previously modelled binding site is ~10 Å (Fig. 1D) (*22*), underscoring the importance and
value of experimental protein-ligand structure determination. The FeEnt electron density in the
co-crystal structure is of very high quality (Fig. 1E), with the cyclic triserine lactone, amide
linkers and Fe^{3+}-ligating catecholate groups fully resolved. The catecholate arms of FeEnt are
not perpendicular to the triserine lactone ring plane but slanted to one side (Fig. 1E, Movie S1),
in agreement with the structure of FeEnt in isolation (*4*). FeEnt does not undergo
conformational changes upon binding to BtXusB. Instead, BtXusB loops close in on FeEnt,
forming a 419 $Å^2$ interaction interface (Fig. S1 and S2, Movie S2). All three catecholate arms
of FeEnt interact with BtXusB loops L1-4 (Fig. 1C and F). Sidechains of N120, R167 and
Q413 slot between the slanted catecholate arms, while S121, S168 and Y411 sidechains form
hydrogen bonds with the *meta* oxygens of the catecholate groups (Fig. 1F, Movie S1). One

amide linker of FeEnt hydrogen-bonds to a single water molecule, another to a water molecule
and the sidechain of Q413, and the third to the sidechain of R255 (Fig. 1F). The sidechain of
R167 forms a hydrogen bond with the triserine lactone ring—the only interaction between
BtXusB and this part of FeEnt.

BtXusB binds FeEnt with a dissociation constant value of ~55 nM as determined by isothermal
titration calorimetry (ITC) (Fig. 1G). We constructed a BtXusB variant, BtXusB$^{R167A,\Delta hook}$, with
the R167A substitution and deletion of residues 119-122 which form the tip of the hook.
Titration of FeEnt into BtXusB$^{R167A,\Delta hook}$ resulted in reduced injection heats and no saturation
(Fig. 1G), which we interpret as lack of binding. The ITC results strongly suggest that the
binding site observed in the co-crystal structure is the only FeEnt binding site on BtXusB.

100
Fig. 1. BtXusB binding to FeEnt. (**A**) Crystal structure of apo BtXusB at 1.56 Å. The cartoon is
coloured in rainbow; the N-terminus is blue, the C-terminus is red. Blades of the β-propeller (β1-7) are
labelled starting from the N-terminus. (**B**) Surface representation of apo BtXusB. (**C**) Co-crystal

structure of BtXusB bound to FeEnt at 1.5 Å. FeEnt is depicted as an orange stick model. The four
FeEnt binding loops (L1-L4) are shown in cyan. (**D**) Surface representation of BtXusB bound to FeEnt.
(**E**) FeEnt model fit into the $2mF_o$-DF_c electron density map at 2σ. (**F**) BtXusB residues interacting with
FeEnt. The yellow dashed lines show likely hydrogen bonds. (**G**) Representative ITC experiments
where 250 μM FeEnt was titrated into 25 μM BtXusB (n=4 experiments) or 25 μM BtXusB$^{R167A,\Delta hook}$
variant (n=2 experiments). Integrated heats were fitted to a single binding site model, giving the
apparent K_d value for the XusB-FeEnt titration. No K_d value could be determined for the XusB$^{R167A,\Delta hook}$
titration.

Ent secreted by pathogens and commensals can be taken up by any other Gram-negative
bacterium that expresses suitable TonB-dependent transporters, such as FepA and XusA (*5*,
21). Furthermore, as part of the innate immune response host cells secrete lipocalin-2, which
sequesters Ent and deprives pathogens of iron (*23*, *24*). Therefore, production of Ent by the
pathogen can be less effective during infection. *Salmonella* and some *E. coli*, e.g. many
uropathogenic strains (*25*), have in turn evolved a strategy to prevent waste of resources and
secure iron by making chemically modified versions of Ent that require the TonB-dependent
transporter IroN for import and do not bind to lipocalin-2, such as di-*C*-glucosylenterobactin
(DGE), also known as salmochelin S4 (*26*, *27*). DGE has the same core structure as Ent, with
two of the catechol groups *C*-glucosylated at the C5 position (Fig. 2A). Notably, *E. coli* FepA
does not import DGE (*7*). We soaked apo BtXusB crystals with iron-bound DGE (FeDGE) and
determined the crystal structure of FeDGE-bound BtXusB to 1.80 Å resolution (Fig. 2B). One
of two protein chains in the asymmetric unit had FeDGE bound, which we could confidently
build into the ligand density (Fig. 2B,C). BtXusB binds FeDGE via the same loops as FeEnt
with additional interactions with the two glucosyl modifications, Glc1 and Glc2 (Fig. 2D). Glc1
hydroxyl groups hydrogen-bond to two water molecules and the hook loop via the backbone
carbonyl oxygen of N120 and the side chain of S123. Glc2 contacts the hook via the sidechain
of S122, L2 via the backbone of R167, and L3 via the backbone of S254. Additionally, Glc2
interacts with a single water molecule and the hydroxyl group of Y189, which is not part of the
four loops that bind FeEnt. The electron density for Glc1 was weaker than for Glc2 (Fig. 2C),
which suggests that Glc1 is bound less tightly than Glc2. The two glucosyl modifications slot
between the ligand-binding loops (Fig. 2E). One notable difference between the FeEnt- and
FeDGE-bound structures is that the amide linker of one of the catechol arms of FeDGE does
not hydrogen-bond with the sidechain of L3 R255 as observed for FeEnt (Fig. 2F). This
rearrangement is likely the result of the C6 atom of Glc2 nudging the sidechain of S254 by
~1.7 Å and pushing the entire L3 further away from the siderophore. Consequently, the
sidechain of R255 swings away by ~5 Å and can no longer contact the amide linker. ITC data
suggest that the amide-R255 interaction is important for tight xenosiderophore binding, as the
FeDGE-BtXusB interaction has a ~2-3-fold lower apparent dissociation constant value
compared to FeEnt (Fig. 2G and Table S2), even though the glucosyl modifications of FeDGE
form additional contacts with BtXusB. In addition, the decreased mobility of the constrained
Glc modifications likely results in an entropic penalty for bound FeDGE. ITC data fitting
results support this prediction, as the BtXusB-FeDGE interaction has an estimated -TΔS value
~2.3 kcal/mol smaller than the BtXusB-FeEnt interaction, while the ΔG terms are similar for
both with a difference of ~0.6 kcal/mol (Table S2).

147
Fig. 2. BtXusB binding to FeDGE. (**A**) Chemical structure of DGE. The glucosyl groups are in red.
(**B**) Crystal structure of BtXusB (green) bound to FeDGE (yellow). The siderophore-binding loops are
shown in cyan. (**C**) FeDGE model fit into the $2mF_o$-DF_c electron density map at 1σ. The arrowheads
point to the C-C bonds between the catechol groups and the C1 atoms of the glucosyl moieties. (**D**)
Hydrogen bonding network formed between the glucosyl groups of FeDGE and BtXusB. (**E**) The Glc1
and Glc2 groups of FeDGE occupy slots between the xenosiderophore-binding loops of BtXusB (green
surface). (**F**) Superposition of BtXusB-FeEnt (grey and orange, respectively) and BtXusB-FeDGE
(green and yellow, respectively) crystal structures. Hook loop residues N120-S123 and L2 R167 are in
almost identical conformations in the two structures, but L3 is pushed away in the FeDGE structure due
a clash between Glc2 of FeDGE and S254 of L3. The black dashed line indicates the hydrogen bond
between R255 and the amide carbonyl of FeEnt. (**G**) Representative ITC experiment where 154 μM
FeDGE was titrated into 17.3 μM BtXusB (n=2 experiments). Integrated heats were fitted to a single
binding site model, giving the apparent K_d value.

Cryo-EM structure of native XusAB complex from *B. theta*

FeEnt is sufficient to rescue growth of *B. theta* under iron limiting conditions (Fig. S3). We
reasoned that the proteins encoded by the Xus operon would be expressed to high enough copy
number for purification and structural characterisation if the cells were cultured under iron-
limiting conditions. We purified the native XusAB complex from the *B. theta bt2064-his* strain
grown in minimal medium supplemented with the iron chelator bathophenanthroline
disulfonate (BPS) (Fig. S3, Methods). We determined the structure of XusAB by single particle
cryo-EM to a global resolution of 2.7 Å, with local resolution extending to 2.2 Å (Fig. 3A, Fig.
S4, Table S3). The resolved portion of XusA has a classic TonB-dependent transporter fold: a
micelle-embedded 22-strand β-barrel occluded by an N-terminal plug domain (*2*). The XusA
extreme N-terminal carboxypeptidase-like domain of unknown function was not resolved in
our structure likely because it is connected to the barrel via a flexible linker. The conformation
of XusB in the XusAB complex is almost identical to that observed in the apo XusB crystal
structure, with Cα-Cα RMSD = 0.9 Å. We did not observe any alternative conformations, sub-
complexes or movement of XusA and XusB in the cryo-EM data.
175

XusB sits on top of the extracellular side of XusA like a lid, reminiscent of *B. theta* TonB-
dependent transporter and surface-exposed lipoprotein complexes that take up glycans and
vitamin B$_{12}$ (*28, 29*). The interaction interface of XusA and XusB is extensive: PISA analysis
indicates 52 hydrogen bonds and 6 salt bridges, with a total interaction surface area of 2,795
Å^2 (Fig. S5). XusB interacts with every extracellular loop of XusA, except for the loop between
barrel strands β9 and β10, which is disordered (Fig. 3B). Although we did not observe FeEnt
in the cryo-EM structure, the XusB hook facing towards a cavity enclosed by the XusAB
complex (Fig. 3A,B) indicates that FeEnt binds inside this cavity. We speculate that XusB
moves in a hinge-like motion opening and closing the XusAB cavity like a lid, thus transiently
allowing extracellular xenosiderophores access to their binding site on the inward-facing side
of XusB.

XusB binds FeEnt with high affinity (Fig. 1G), but FeEnt must somehow be transferred from
XusB to XusA to achieve transport across the OM. Structural alignment of the cryo-EM XusB
structure and the XusB-FeEnt co-crystal structure reveals that Q413, part of L4, must undergo
a shift of 7.7 Å to interact with FeEnt (Fig. 3C). However, the position of L4 in the XusB-FeEnt
structure would result in clashes with the XusA β7-8 extracellular loop within the apo complex
(Fig. 3D). Furthermore, L4 in the apo complex is stabilised via a salt bridge between E416 and
K354 of XusA β5-6 extracellular loop. This suggests that the FeEnt-bound state of XusB when
in complex with XusA might be short-lived despite the high affinity of XusB for FeEnt. We
speculate that the XusA β7-8 extracellular loop displaces L4 of XusB, aided by formation of
the E416-K354 salt bridge, thus disrupting the FeEnt binding pocket and releasing the
xenosiderophore to diffuse towards XusA, which would transport it across the OM. We
envisage that the transfer of FeDGE from XusB to XusA proceeds via an identical mechanism
to FeEnt.

We observed unexplained density extending from the sidechain of T401, which is part of the
XusA β7-8 extracellular loop (Fig. S6). Together with flanking residues this threonine forms a
DTA sequence, which matches the Bacteroidota O-glycosylation motif (*30*). We therefore
conclude that the cryo-EM density extending from T401 corresponds to a glycan chain. Four
sugar units can be discerned, including the branching deoxyhexose previously identified in the
B. fragilis O-glycan (*31*). The T401 glycosylation site is located near residues 421-423 which
are implicated in FeEnt release from XusB (Fig. 3D and Fig. S6). The sidechain of T401 and
the O-glycan face the solvent rather than XusB, but we cannot rule out that the glycan

modification affects the conformation of neighbouring XusA extracellular loops and that it
could be functionally important.

210
Fig. 3. Cryo-EM structure of the native XusAB complex from *B. theta*. (**A**) Single particle cryo-EM
reconstruction of the XusAB complex at 2.7 Å global resolution. (**B**) Model built into the cryo-EM
density in cartoon representation. The red dashed line depicts the disordered extracellular loop between
β strands 9 and 10 of the XusA β-barrel, corresponding to residues 464-473. (**C**) Structural alignment
of the apo-XusB structure observed in the cryo-EM reconstruction (hot pink, hook in green) and the
XusB-FeEnt co-crystal structure (grey, FeEnt binding loops in cyan). Cα-Cα RMSD = 1.6 Å. The
dashed yellow line corresponds to a distance of 7.7 Å between the Cα atoms of Q413, which is part of
the L4 FeEnt binding loop, observed in the two structures. (**D**) Close-up view of XusB L4 in the apo
cryo-EM structure and the XusB-FeEnt co-crystal structure, viewed from a different orientation
compared to (**C**). Colouring as in (**C**), XusA is in light sea green. The yellow dashes correspond to a
distance of 11.3 Å between the Cα atoms of E416 in the two structures. The black dashes show a salt
bridge between K354 of XusA and E416 of XusB in the cryo-EM structure. Residues E421, Q422 and
S423 are part of the β7-8 extracellular loop of XusA.

Structural variety of XusB homologues

BtXusB is annotated as a DUF4374-containing protein in UniProt (*32*), which has a β-propeller
fold as demonstrated by our structural data. We investigated the predicted structures of
DUF4374-containing homologues. The top hits from protein BLAST searches of the BtXusB
sequence against Bacteroidota genomes have a sequence identity of around 60%. The reason

for the relatively low sequence identity for the top BLAST hits is that the BtXusB hook position
inside the β2 propeller blade is unique (Fig. 4A). Other Bacteroidota XusB homologues, such
as *Parabacteroides distasonis* BDI_3402 (59.5% identity; UniProt accession A6LHE0) and
Barnesiella viscericola BARVI_05925 (43.4% identity; UniProt accession W0ET74), are
predicted to have their hook inserted in the β4 blade or, in the case of *B. fragilis* BF9343_4228
(20.2% identity; UniProt accession Q5L7E3), do not appear to have a hook at all (Fig. 4B-D
and Fig. S7). These DUF4374-containing proteins are likely part of genuine xenosiderophore
utilization systems, rather than performing some other function. They all contain a lipoprotein
export signal that directs proteins to the outer leaflet of the OM (*33*). Additionally, they are
encoded next to TonB-dependent transporters and PepSY domain-containing inner membrane
proteins that likely reduce siderophore-bound ferric iron to facilitate dissociation of the iron-
siderophore complex (*34*) (Fig. 4E). *P. distasonis* also encodes a periplasmic esterase
(BDI_3403) in the same operon as the XusB homologue which might liberate siderophore-
bound iron via siderophore hydrolysis, as shown for FeEnt in *E. coli* (*8*, *9*).

Conservation of the hook and L2 amino acid sequences between the *B. theta, P. distasonis* and
B. viscericola homologues (Fig. S7) suggests that, despite differences in position of the hook
within the β-propeller, *P. distasonis* and *B. viscericola* homologues might still bind FeEnt,
FeDGE and perhaps other catecholate siderophores. However, the *B. fragilis* homologue is
unlikely to bind FeEnt due to lack of all FeEnt-binding regions observed in the BtXusB-FeEnt
co-crystal structure, which is consistent with a previous report that *B. fragilis* cannot utilize
iron bound to catecholate xenosiderophores (*35*).

The amino acid sequence length of BtXusB BLAST hits follows a bimodal distribution with
peaks at 410 and 465 amino acids (Fig. S9). *B. theta, P. distasonis* and *B. viscericola*
homologues (464, 471 and 491 residues, respectively) belong to the longer group, while the *B.*
fragilis homologue (406 residues) belongs to the shorter group. Together with the differences
in the xenosiderophore-binding loop regions, the bimodal distribution suggests there are at least
two different subtypes of DUF4373-containing proteins.

256
Fig. 4. Structural variety of DUF4374-containing proteins. (**A**) BtXusB-FeEnt co-crystal structure.
FeEnt atoms are displayed as spheres. (**B**) *P. distasonis* DSM 20701 BDI_3402 (59.52% identity to *B.*
theta XusB), (**C**) *B. viscericola* DSM 18177 BARVI_05925 (43.36% identity to BtXusB), and (**D**) *B.*
fragilis NCTC 9343 BF9343_4228 (20.21% identity to BtXusB) AlphaFold2 (*36*) models. Views were
generated from a superposition, β-propeller blades are labelled β1-7 starting from the N-terminus. All
models are coloured in rainbow, from the N-terminus in blue to the C-terminus in red; the hook is in
black. AlphaFold2 model prediction confidence is shown in Fig. S8. (**E**) Genetic context of the selected
XusB homologues. DUF, domain of unknown function; LES, lipoprotein export signal (*33*).

B. viscericola XusB binds ferric enterobactin

265
We produced recombinant XusB from *B. viscericola* (BvXusB) to investigate its
xenosiderophore-binding properties. We observed binding of FeEnt to BvXusB in ITC with
similar affinity to BtXusB (Fig. 5A). We attempted to investigate the molecular details of the
interaction between BvXusB and FeEnt and obtained the crystal structure of apo BvXusB (Fig.
5B and Table S1), which confirmed the computational prediction. However, we could not co-
crystallise BvXusB with FeEnt, and BvXusB crystals soaked with FeEnt did not diffract.
Superposition of the BtXusB-FeEnt and apo BvXusB structures suggests that most BtXusB
residues involved in FeEnt binding are present in BvXusB, even though the hook loop in
BvXusB is inserted in the β4 blade rather than in the β2 blade as in BtXusB (Fig. 5C). Based
on these structural similarities and AlphaFold 3 predictions of BvXusB in complex with FeEnt
and FeDGE (Fig. S10), we expect that BvXusB and other XusB homologues with the hook
inserted in the β4 blade interact with FeEnt and FeDGE in a very similar manner to BtXusB.

278
Fig. 5. *B. viscericola* XusB binds FeEnt. (**A**) Representative ITC experiment where 250 μM FeEnt
was titrated into 25 μM BvXusB (n=2). Integrated heats were fitted to a single binding site model,
giving the apparent K_d value. (**B**) Crystal structure of apo BvXusB to 3.2 Å. (**C**) Comparison of apo
BvXusB (cyan) and BtXusB-FeEnt (grey and orange) crystal structures. The models were superposed
using the Matchmaker tool in ChimeraX (Cα-Cα RMSD between 252 pruned atom pairs was 1.0 Å;
across all 301 pairs—3.6 Å). BtXusB residues involved in FeEnt binding and the equivalent residues in
BvXusB are shown as stick models. The dashed line indicates a disordered loop in the BvXusB crystal
structure.

B. fragilis XusB binds ferrichrome

287
We produced recombinant *B. fragilis* XusB (BfXusB) and determined its crystal structure using
data to 1.77 Å, which was consistent with the AlphaFold 2 prediction (Fig. 4D and 6A).
BfXusB lacks the hook loop observed in BtXusB crystal structures (Fig. 6B). A previous study
reported that *B. fragilis* can grow on ferrichrome as the sole iron source, but the ferrichrome
transporter could not be conclusively identified (*35*). Ferrichrome binds iron via hydroxamate
groups, rather than catecholate groups as in Ent and DGE (Fig. 6C). FeEnt titrations into
BfXusB did not result in substantial injection heats indicating lack of binding, but titrations of
ferrichrome into BfXusB showed binding (Fig. 6D,E). Titration of ferrichrome into BtXusB
and BvXusB did not show heat changes that would suggest binding (Fig. 6E and Fig. S11). We
then determined the structure of BfXusB from crystals soaked with ferrichrome using data to

3.32 Å resolution (Fig. 6F,G and Table S1). We saw additional electron density consistent with
the structure of ferrichrome inside the β-propeller pocket without any changes in protein
conformation compared to the apo structure (Fig. 6G,H). The moderate resolution makes it
difficult to reliably discern which residues of BfXusB interact with ferrichrome, but it is clear
the binding residues are completely different to those in BtXusB and BvXusB. The side chains
of L93 and W235 slot between two grooves formed by the hydroxamate arms, while the third
groove remains unoccupied and exposed to solvent. The side chains of Y52, Y77 and F398 and
backbones of W323 and D352-G354 interact with the cyclic peptide backbone (Fig. 6H), in
contrast to the BtXusB-FeEnt interaction where there are no contacts with the triserine lactone
ring (Fig. 1F). Despite these differences, the xenosiderophore binding sites in BfXusB and
BtXusB occupy the same location in the context of the DUF4374 fold (Fig. 6I). This points to
a common evolutionary ancestry despite low sequence homology and suggests that other
DUF4374-containing proteins also bind their ligands at the same site.

Xenosiderophore utilization bioassay
The BfXusB-ferrichrome structure and binding data strongly suggested that *B. fragilis* takes
up ferrichrome across its outer membrane via a XusAB-like complex formed by BfXusB and
a XusA homologue encoded in the same operon (BF9343_4229; Fig. 4E). We wanted to
experimentally confirm this and used a modified assay by Rocha & Krykunivsky (*35*) to test
xenosiderophore utilization in *B. theta* and *B. fragilis xusAB* deletion strains. We made *xusAB*
knockouts in *B. theta* and *B. fragilis* thymidine kinase deletion (*tdk*⁻) strains using the
pExchange-tdk allelic exchange system (*37*). As shown previously (*21*), *B. theta* could grow
under iron-limiting conditions on both FeEnt and FeDGE as the sole iron source in a *xusAB*-
dependent manner (Fig. 7A). We found that growth of *B. fragilis* on ferrichrome is also *xusAB*-
dependent (Fig. 7B). This observation confirms the in vivo relevance of our in vitro findings.
Surprisingly, we found that *B. fragilis* with and without the *xusAB* locus could utilize FeEnt
(Fig. 7A). This result implies that there are more xenosiderophore utilization systems in *B.*
fragilis, though BLAST searches did not reveal any paralogous loci to *xusABC* (BF9343_4229-
4227).

326
327 **Fig. 6. *B. fragilis* XusB binds ferrichrome.** (**A**) Crystal structure of apo BfXusB at 1.77 Å shown in
328 cartoon (left) and surface (right) representations. (**B**) Superposition of apo BfXusB and apo BtXusB
329 crystal structures generated using Matchmaker in ChimeraX (Cα-Cα RMSD = 1.2 Å between 140

pruned atom pairs; 3.8 Å across all 249 pairs). (**C**) Chemical structure of desferrichrome. The ferric iron-ligating hydroxamate groups are coloured in blue. (**D**) Representative ITC experiment where 250 μM ferrichrome was titrated into 25 μM BfXusB (n=4 repeats). Integrated heats were fitted to a single binding site model, giving the apparent K_d value. (**E**) Representative ITC experiments where 250 μM FeEnt was titrated into 25 μM BfXusB (n=2 experiments) and 250 μM ferrichrome was titrated into 25 μM BtXusB (n=2 experiments). (**F**) Crystal structure of ferrichrome-bound BfXusB to 3.32 Å. (**G**) Surface representation of BfXusB with ferrichrome shown as a stick model inside the shallow β-propeller pocket. (**H**) Ferrichrome model fit into the $2mF_o$-DF_c electron density map at 1σ. Residues that likely interact with ferrichrome are shown as yellow stick models. The same residues in the apo BfXusB crystal structure are shown as salmon stick models. (**i**) Superposition of BfXusB-ferrichrome (yellow and purple) and BtXusB-FeEnt (grey and orange) crystal structures (Cα-Cα RMSD between 140 pruned atom pairs was 1.2 Å; across all 249 pairs—3.8 Å). The distance between the two Fe^{3+} ions in the aligned structures is 2.3 Å.

Fig. 7. Xenosiderophore utilization bioassay. *B. theta* and *B. fragilis tdk⁻* and *ΔxusAB* strains were spotted on minimal medium agar plates under iron-replete (200 μM FeCl₃) or iron-limiting conditions (100 μM BPS). Iron-siderophores or the solvent controls were supplied via sterile filter discs (white). The *tdk⁻* strains are thymidine kinase knockouts that are sensitive to 5-fluoro-2′-deoxyuridine that is used for counter-selection during allelic exchange but otherwise have wild-type genotypes. (**A**) shows FeEnt and FeDGE (both supplied in 50% DMSO) utilization. (**B**) shows ferrichrome utilization.

Discussion

There are many TonB-dependent transporters in Bacteroidota that transport unknown molecules (*38*). It is important to elucidate what these transporters are transporting and how, so that we can obtain a more complete understanding of microbial interactions inside the gut. Our study provides insights into xenosiderophore uptake via Bacteroidota TonB-dependent transporter-lipoprotein complexes and contributes to the ever-widening substrate range described for lipoprotein-assisted TonB-dependent transporters (*2, 28, 29, 38–41*).

We observed an excess of XusB in our native XusAB purifications (Fig. S3). The excess XusB is likely destined for secretion in outer membrane vesicles (OMVs) (*22*). It has been shown that OMVs containing FeEnt-loaded XusB can be used as an iron source by *B. theta* (*22*). This observation implies that the OMVs can somehow deliver FeEnt bound to XusB to XusA in the OM of recipient bacteria. However, our cryo-EM structure shows that XusAB form a tight, closed complex even in the absence of FeEnt. It is unclear how the FeEnt bound to XusB could

access the apo-XusB bound to XusA. We speculate that XusB transiently opens like a lid and gives access to the XusAB cavity to incoming xenosiderophores, analogous to the mechanism proposed for vitamin B_{12} import by *Bacteroides* (*28*). Alternatively, the disordered β9-10 extracellular loop of XusA might act as a lateral gate for ingress of xenosiderophores. The latter hypothesis is favoured by lack of any XusAB particles in an open state in the cryo-EM data and the many interactions between the closed XusB lid and XusA (Fig. S5). On the other hand, closure of a mobile XusB lid appears more effective in dislodging the XusB-bound siderophore via clashes with XusA loop β7-8, and we therefore favour the hypothesis of a mobile XusB lid delivering the xenosiderophore to the XusA transporter. It is noteworthy that, while open lipoprotein lids have been observed experimentally for all characterised dimeric SusCD transporters (*29, 39–41*), for unknown reasons this has not been the case for BtuBG (*28*) and XusAB systems, which are all monomeric. It is also unclear why we could not load the purified XusAB complex with FeEnt or FeDGE. The in vitro conditions may somehow preferentially stabilise the detergent-purified complex in a closed state, while in vivo the presence of negatively charged lipo-oligosaccharide in the *B. theta* OM could destabilise the XusAB interaction. It is also conceivable that excess free, but OM-anchored XusB plays a role in complex opening.

B. theta XusAB can transport at least two structurally similar xenosiderophores, FeEnt and FeDGE (*21, 35*). BtXusB interacts extensively with the siderophore triscatecholate structure, and the binding site can also accommodate the two glucosyl groups of FeDGE. Furthermore, the triserine lactone of FeEnt and FeDGE is exposed to solvent, suggesting that the binding site could accommodate siderophores with different backbones connecting the catecholate arms, such as vibriobactin and bacillibactin, or synthetic Ent mimics (*20, 42*). This interaction mode raises the question of how many different xenosiderophores a single XusAB complex can import, which could have a substantial impact on our understanding of competition for iron at the host-pathogen-commensal interface.

Together, our structures, functional data and bioinformatics analyses suggest that the presence or absence of the hook in DUF4374-containing proteins indicates preference for catecholate or hydroxamate xenosiderophores, respectively. Whether the shorter, BfXusB-like homologues can bind multiple related xenosiderophores as well remains unknown. Cyclic hydroxamate siderophores structurally related to ferrichrome, such as ferricrocin and ferrichrysin, have hydroxymethyl modifications on the peptide backbone (*43–45*). Because BfXusB interacts extensively with the backbone, these substitutions may either form additional contacts with BfXusB or clash with its sidechains. Further functional and structural analyses are required to determine the substrate range of hydroxamate-binding XusB homologues.

While catecholate siderophores are produced by most Enterobacteriaceae, hydroxamate siderophores are mainly produced by fungi (*45*). Our results suggest that many Bacteroidota species, such as *B. fragilis*, can steal fungal siderophores. Interactions between bacteria and fungi in microbial communities are complex. *Candida albicans* has been shown to promote growth of *B. fragilis* in vitro (*46*). On the other hand, bacteria that have type VI secretion systems, including *B. fragilis,* can directly inject antifungal effectors into fungal cells (*47–49*). The role of iron piracy by *B. fragilis* in the context of its interactions with fungi remains unknown and requires further study.

Methods

Bacterial strains and culture conditions

All strains used in this study are listed in Table S4. *E. coli* BL21(DE3) (*50*) and TOP10 strains
were genetically manipulated using standard laboratory procedures. The kanamycin
concentration used in liquid and solid lysogeny broth (LB) was 50 µg/ml. *B. theta* VPI-5482
tdk⁻ was cultured either in brain-heart infusion (BHI, Oxoid) supplemented with 1 µg/ml hemin
or defined minimal medium (50 mM potassium phosphate buffer pH 7.2, 15 mM sodium
chloride, 7.5 mM ammonium sulfate, 9.4 mM sodium carbonate, 4.1 mM L-cysteine, 0.37 nM
vitamin B_{12}, 5.8 µM vitamin K, 180 µM calcium chloride, 100 µM magnesium chloride, 50
µM manganese(II) chloride and 42 µM cobalt(II) chloride)(*51*) supplemented with 1 µg/ml
hemin and 0.5% fructose. *B. theta* was grown under anaerobic conditions at 37 °C in a Don
Whitley A35 workstation. When required, 200 µg/ml gentamycin and 25 µg/ml erythromycin
was used for selection of *B. theta*.

Construction of expression plasmids
All plasmids used in this study are listed in Table S5. The nucleotide sequence coding for amino
acid residues 35-464 of *B. theta* VPI-5482 XusB (*bt_2064*) and 30-406 of *B. fragilis* NCTC
9343 XusB (*bf9349_4228*) which excludes the signal sequence, the lipid anchor cysteine and
a disordered linker, was amplified by PCR using genomic DNA as the template and primers
that introduce overhangs containing NcoI and XhoI restriction sites. The PCR product was
digested with NcoI and XhoI FastDigest restriction enzymes (ThermoFisher Scientific) and
ligated into pET28b, resulting in a C-terminal His$_6$-tag fusion. The XusB$^{R167A,\Delta hook}$ binding site
variant nucleotide sequence with flanking NcoI and XhoI restriction sites was synthesised and
ligated into the pTwist Amp High Copy plasmid by Twist Bioscience. The synthetic DNA
sequence was sub-cloned into pET28b using the flanking restriction sites. Similarly, the
nucleotide sequence encoding amino acid residues 29-491 of *B. viscericola* DSM 18177 XusB
(*BARVI_05925*) was synthesised by Twist Bioscience and sub-cloned into pET28b using NcoI
and XhoI restriction sites. Cloning was carried out in *E. coli* TOP10 cells. Clones were
screened for successful insert ligation by colony PCR using EmeraldAmp GT PCR master mix
(Takara Bio) with T7 promoter and T7 terminator primers. All constructs were verified by
Sanger sequencing (Eurofins).

Construction of chromosomal deletions
B. theta and *B. fragilis xusAB* chromosomal deletions were generated using the pExchange-tdk
plasmid (*37*). The pExchange-tdk plasmids containing ~700 bp flanking the *xusAB* coding
sequences in *B. theta* VPI-5482 (*bt2064-65*) or *B. fragilis* NCTC 9343 (BF9343_4228-29) were
used to transform *E. coli* S-17 λ pir cells (*52*). The *E. coli* cells were used to introduce the
pExchange-tdk plasmids into recipient *Bacteroides* strains lacking thymidine kinase (*tdk⁻*).
Conjugants, which underwent a single recombination event, were selected on BHI-hemin plates
containing gentamicin (200 µg/ml) and erythromycin (25 µg/ml). Single colonies were cultured
in enriched BHI medium overnight, pooled, and plated on BHI-hemin agar plates containing
5-fluoro-2′-deoxyuridine (FUdR; 200 µg/ml) to select for cells that have eliminated the vector
backbone from their genome in a second recombination event. After 48 h of growth, FUdR-
resistant colonies were restreaked on fresh BHI-hemin-FUdR plates. Clones were screened for
successful *xusAB* deletion using PCR with primers that bind outside the homologous region
used to make the pExchange-tdk plasmids and confirmed by Sanger sequencing (Eurofins).

Protein expression and purification in *E. coli*
E. coli BL21(DE3) cells were transformed with plasmids carrying the coding sequences for
soluble variants of XusB, plated on LB kanamycin agar plates and incubated at 37 °C overnight.
The following day, approximately one third of transformants were scraped off the agar plate
and used to start an LB kanamycin preculture, which was incubated at 37 °C with shaking for

1.5-3 hours. 12-15 ml of preculture was used to inoculate 1 l flasks of pre-warmed LB
kanamycin medium. The cultures were incubated at 37 °C with shaking until OD_{600} reached
0.4-0.8. Protein expression was then induced by adding isopropyl β-d-1-thiogalactopyranoside
to a final concentration of 0.1 mM, and the cultures were grown for another 16-20 h at 18 °C
with 150 rpm shaking. Cells were harvested by centrifugation at 8,000 × g at 4 °C for 20 min.
Pellets were resuspended in cold Tris-buffered saline (TBS, 20 mM Tris-HCl, 300 mM NaCl)
and stored at -20 °C until protein purification. Cell pellets were thawed and homogenized in
TBS using a Dounce tissue grinder and supplemented with DNase I (Roche). Cells were lysed
by passing the suspension once through a cell disruptor (Constant Systems) at 20 kpsi. The
lysates were supplemented with 1 mM phenylmethylsulfonyl fluoride (PMSF) and clarified by
centrifugation at 30,000 g for 30 min at 4 °C. The supernatants were loaded on 5 ml chelating
Sepharose columns charged with Ni^{2+} ions using gravity flow. The columns were washed with
30 column volumes of TBS with 30 mM imidazole. Bound proteins were eluted with 4 column
volumes of TBS with 250 mM imidazole. The eluate was concentrated with an Amicon Ultra
filtration device (30 kDa cut-off membrane), loaded on a HiLoad Superdex 200 16/600 pg
column and eluted in 10 mM HEPES-NaOH pH 7.5, 100 mM NaCl. Fractions were analysed
by SDS-PAGE. Fractions containing the soluble XusB variants were pooled, concentrated,
flash-frozen and stored at -80 °C.

Synthesis of DGE
DGE was prepared chemoenzymatically based on previously reported procedures (*53*, *54*). A
reaction solution containing 500 μM Ent and 3 mM UDP-glucose (UDP-Glc) in buffer (75 mM
Tris-HCl pH 8.0, 5 mM $MgCl_2$, 2.5 mM TCEP) was prepared and aliquoted into 10 × 750 μl
portions. The *C*-glucosyltransferase IroB was added to each sample to a final concentration of
5 μM and the reactions were incubated at room temperature. After 1 h, each aliquot was
quenched by addition of 75 μl 6% trifluoroacetic acid (TFA)/H_2O and diluted with ~300 μl
acetonitrile (MeCN). Quenched solutions were centrifuged (13,000 rpm, 10 min, 4 °C) then
purified by preparative reversed phase HPLC (0−100% B in 30 min, H_2O/MeCN + 0.1% TFA,
10 mL/min). The peak corresponding to DGE was collected and lyophilized to dryness.

Preparation of ferric siderophores
Ent and DGE were dissolved in dimethyl sulfoxide (DMSO) to make 10 mM solutions. The 10
mM siderophore solution was mixed in a 1:1 volume ratio with 10 mM aqueous $FeCl_3$ solution
to yield a 5 mM ferric siderophore stock solution in 50% (v/v) DMSO. The concentrations of
FeEnt and FeDGE stock solutions were determined spectrophotometrically using the extinction
coefficient $\varepsilon_{495} = 5{,}600\ M^{-1}\ cm^{-1}$ (in 20 mM Tris-HCl pH 7.0, 50% methanol)(*55*). The stock
solutions were stored in small aliquots at -80 ºC to minimise siderophore hydrolysis. Similarly,
desferrichrome (Merck) was dissolved in sterile Milli-Q water to make a 10 mM solution,
aliquoted and stored at -20 ºC. After thawing, 10 mM desferrichrome aliquots were mixed with
10 mM $FeCl_3$ in a 1:1 ratio, resulting in a 5 mM aqueous ferrichrome solution. Ferrichrome
concentration was confirmed spectrophotometrically using the extinction coefficient
$\varepsilon_{425} = 2{,}900\ M^{-1}\ cm^{-1}$ (in 20 mM Tris-HCl pH 7.0)(*56*).

Crystal structure determination
The purified soluble BtXusB variant was concentrated to 36 mg/ml. Sitting drop vapour
diffusion crystallisation screens were set up using a Mosquito robot (SPT Labtech) either with
apo BtXusB or with a 1:1.1 molar ratio of BtXusB and FeEnt. The crystallization plates were
incubated at 20 °C. Apo XusB crystals appeared after less than a week in the PACT premier
screen (Molecular Dimensions) containing 0.1 M MES pH 6.0, 0.2 M calcium chloride and
20% PEG 6000. Crystals were cryo-protected in mother liquor supplemented with ~20% PEG

400 and flash-cooled in liquid nitrogen. BtXusB-FeEnt co-crystals appeared in the Index screen (Hampton Research) condition containing 0.1 M citric acid pH 3.5 and 2 M ammonium sulphate after 2 weeks. Crystals were cryo-protected by passing through a drop of 3.5 M ammonium sulphate and flash-cooled in liquid nitrogen. Apo BtXusB crystals grown in hanging vapour diffusion drops (0.1 M HEPES pH 6.7, 0.2 M calcium chloride, 20% PEG 6000, and 5% DMSO) were soaked with 1 mM FeDGE in mother liquor for 24 h, cryoprotected with mother liquor, 1 mM FeDGE and 20% PEG 400, and flash-cooled in liquid nitrogen.

BvXusB was concentrated to 35 mg/ml and sitting drop crystallization trials using commercial screens were set up as above. An initial hit was observed in the Structure 1+2 screen (Molecular Dimensions) and further optimized using hanging drop vapour diffusion and streak seeding. The final condition for diffracting apo BvXusB crystals was 0.1 M sodium acetate pH 4.8, 0.2 M ammonium sulfate and 35% PEG 2000 monomethyl ether. The crystals were cryoprotected in mother liquor with 20% PEG 400 and flash-cooled in liquid nitrogen.

BfXusB was concentrated to 30 mg/ml and sitting drop crystallization trials using commercial screens were set up as above. Apo crystals were harvested directly from the Index screen (Hampton Research) condition containing 0.1 M HEPES pH 7.5, 0.02 M magnesium chloride hexahydrate, 22% w/v poly(acrylic acid sodium salt) 5100. Mother liquor with 20% PEG 400 was used to cryoprotect the crystals before flash-cooling in liquid nitrogen. Crystals from an identical condition in the JCSG+ screen (Molecular Dimensions) were soaked with 1 mM ferrichrome in mother liquor for 3 weeks, cryoprotected in mother liquor, 1 mM ferrichrome and 20% PEG 400 and flash-cooled in liquid nitrogen.

X-ray diffraction data were collected at the Diamond Light Source synchrotron (UK) at a temperature of -173 °C (Table S1). Datasets were processed with XIA2-dials (*57*), scaled with Aimless (*58*), and the space group was confirmed with Pointless (*59*). The apo structures were solved by molecular replacement with computational models generated by AlphaFold2 (*36*). The siderophore-bound structures were solved by molecular replacement using the apo experimental structures. All models underwent cycles of manual building in Coot (*60*) and refinement in Phenix (*61*) until no further improvement in R factors could be achieved. The models were validated using MolProbity (*62*). Refinement statistics are summarised in Table S1.

Isothermal titration calorimetry

ITC was carried out in 10 mM HEPES–NaOH pH 7.5 and 100 mM NaCl supplemented with 5% DMSO to improve FeEnt and FeDGE solubility. A 250 µM FeEnt solution was made in ITC buffer, and any precipitate was removed by centrifugation. The clarified FeEnt solution was titrated into 25 µM protein at 25 °C using a Microcal PEAQ-ITC instrument (Malvern Panalytical). After an initial delay of 60 s, a single injection of 0.4 µl was carried out, which was discarded from data analysis, followed by 18 injections of 2 µl spaced in 150 s intervals. For FeDGE titrations only, the initial 0.4 µl injection was followed by 12 injections of 3 µl spaced in 150 s intervals. The sample cell was stirred at 750 rpm during titration. Ligand to buffer control titrations were subtracted from all experiments. The experiments were repeated at least twice (Table S2). Data were fitted to a single-binding-site model using the Microcal PEAQ-ITC Analysis software v1.40. For FeDGE titrations only, the stoichiometry (n) was fixed to one and the ligand concentration was allowed to float as the fits would not converge otherwise. We speculate that the FeDGE concentration may have been inaccurate due to hydrolysis of the siderophore during repeated freeze-thawing. Data fitting results for successful binding experiments are shown in Table S2.

Growth curve experiments

B. theta tdk⁻ strain was cultured anaerobically in BHI supplemented with hemin at 37 ºC overnight. 0.2 ml of the overnight culture was used to inoculate 5 ml of fresh supplemented BHI the next morning, followed by a 4 h incubation. The cells were collected by centrifugation for 5 min at 2,800 × *g*, 20 ºC, and resuspended in 1 ml of fresh, pre-warmed minimal medium. Minimal medium supplemented with fructose and hemin was aliquoted into tubes, to which BPS, FeEnt and FeCl$_3$ was added as required. Washed cells were diluted in the appropriate tubes to OD$_{600}$=0.04 and dispensed in 200 µl aliquots into the wells of a sterile 96-well plate in triplicate. Growth at 37 ºC was monitored for 48 h using a Biotek Epoch microplate reader housed inside an anaerobic workstation. The experiment was repeated three times with similar results.

Purification of the XusAB complex from *B. theta*

A C-terminal His$_6$-tag was fused to XusB by introducing the tag coding sequence into the *B. theta tdk⁻* chromosome before the stop codon of *bt_2064* via allelic exchange using the pExchange plasmid (*37*). Presence of the tag on the chromosome was confirmed by PCR and Sanger sequencing. Conditions under which the tagged XusB is expressed were identified by Western blotting. The *B. theta bt_2064-his* strain was cultured anaerobically in tubes containing 2 ml minimal medium supplemented with 6.25-100 µM BPS for 18 h. Equivalents of 1 ml culture at OD$_{600}$=2 were pelleted by centrifugation. The pellets were resuspended in 80 µl BugBuster (Sigma), supplemented with 1 mM PMSF, and incubated for 15 min at room temperature. Cell debris was pelleted by centrifugation in a benchtop microcentrifuge. Samples from cells grown in the presence of different amounts of BPS were separated by SDS-PAGE and transferred onto a PVDF membrane via wet transfer. The PVDF membrane was stained with Ponceau S stain to confirm successful transfer and blocked with 1% milk solution in PBS supplemented with 0.1% (v/v) Tween 20 for 20 min at room temperature. The membrane was probed with anti-His-horseradish peroxidase conjugate antibody (Roche; 1:500 dilution in 1% milk solution) for 1 h at room temperature and washed three times with PBS-Tween. The blots were developed using SuperSignal West Pico Plus chemiluminescent substrate (Thermo Fisher Scientific) and imaged using a Gel Doc XR+ system (Bio-Rad).

Frozen *B. theta bt_2064-his* glycerol stocks were used to inoculate BHI, followed by overnight incubation at 37 °C under anaerobic conditions. After autoclaving, minimal medium was stored in an A35 Don Whitley anaerobic workstation overnight. The following day, the minimal medium was supplemented 50 µM BPS, fructose (0.5%) and hemin (1 µg/ml). Overnight *B. theta* BHI cultures were pelleted, resuspended in minimal medium to original volume and used to inoculate 0.5 l bottles of minimal medium at a ratio of 1:250. Bacteria were cultured at 37 °C under anaerobic conditions for 18-20 h. Cultures were pelleted by centrifugation at 6,000g for 30 min at 4 °C, resuspended in TBS, and stored at -20 °C.

Pellets were thawed, supplemented with DNase I and homogenised. Cells were lysed by passing the cell suspension once through a cell disruptor at 22 kpsi. The lysate was clarified by centrifugation at 30,000 × g, 4 °C for 30 min. Membranes were isolated from the clarified lysate by ultracentrifugation at 140,000 × g, 4 °C for 50 min, followed by solubilization in 1.5% lauryldimethylamine N-oxide (LDAO) in TBS for 1 h at 4 °C. Insoluble material was pelleted by centrifugation at 44,000 × g, 4 °C for 30 min. The solubilised material was passed through ~3 ml of chelating Sepharose resin charged with Ni^{2+} ions using gravity flow. The column was washed with 25 column volumes of TBS with 30 mM imidazole and 0.1% dodecyl-β-D-maltopyranoside (DDM, Anatrace), and bound protein was eluted with TBS supplemented with 200 mM imidazole and 0.03% DDM. The eluate was concentrated using an Amicon Ultra

595 filtration device (100 kDa cut-off membrane), loaded on a Superdex 200 10/300 Increase
596 column and eluted in 10 mM HEPES-NaOH pH 7.5, 100 mM NaCl, 0.03% DDM. Fractions
597 corresponding to the XusAB peak were pooled, concentrated, flash-frozen in liquid nitrogen,
598 and stored at −80 °C.

Cryo-EM structure determination

Pure XusAB complex at 7 mg/ml was incubated with a four-fold molar excess of FeEnt for 45
min at room temperature. 3.5 μl of the complex was then applied to glow-discharged Quantifoil
R1.2/1.3 copper 200 mesh holey carbon grids. The grids were immediately blotted for 1-5 s
and plunge-frozen in liquid ethane using a Vitrobot Mark IV (ThermoFisher Scientific) device
operating at 4 °C and 100% humidity. The grids were initially screened on a 200kV FEI Glacios
microscope at the University of York (UK). Data were collected at the Astbury Centre (Leeds,
UK) on a FEI Titan Krios microscope operating at 300 kV using a Falcon 4i direct electron
detector (ThermoFisher Scientific) operating in counting mode (Table S3). A total of 6,645
movies were recorded in electron event representation (EER) format at 165,000×
magnification, corresponding to a pixel size of 0.74 Å.

The cryo-EM workflow is shown in Fig. S4. All data processing was done in cryoSPARC
v4.4.1. Movies were motion-corrected using patch motion correction. CTF parameters were
fitted using patch CTF correction. Initially, ~2,000 particles were picked manually and used to
make 2D classes for template picking. 1,110,843 picked particles were extracted in 480-pixel
boxes (0.74 Å/pixel), Fourier cropped to a box size of 240 pixels (1.48 Å/pixel) and subjected
to two rounds of 2D classification. Ab initio models were generated from classes exhibiting
protein density and decoy models were generated from classes containing predominately noise.
Particles selected after 2D classification were subjected to heterogeneous refinement against
four ab initio models, two of which were decoys. A single class from heterogeneous refinement
refined to high resolution in non-uniform refinement (*63*). Most particles clustered into a single
class in 3D classification (8 classes in total). Particles from this class were re-extracted in 480-
pixel boxes (0.74 Å/pixel) and refined using non-uniform refinement (per-particle defocus, tilt
and trefoil refinement enabled), followed by local refinement. Per-particle motion correction
was performed using reference-based motion correction, followed by local refinement and re-
estimation of per-particle defocus, tilt and trefoil parameters. The particle stack was subjected
to a final round of 2D classification with the noise model (sigma) annealing turned off to
remove any remaining poor-quality particles. Another round of reference-based motion
correction was performed, followed by a final round of local refinement that gave a map with
a global resolution of 2.7 Å based on the FSC=0.143 criterion. The final stack had 76,979
particles.

The XusAB protein sequences were supplied to ModelAngelo (*64*) for automated model
building. The resulting model was iteratively adjusted in Coot (*60*) and ISOLDE (*65*) and
refined using Phenix real space refinement (*61*). The model was validated using MolProbity
(*62*). Refinement statistics are shown in Table S3.

Structure analysis and visualisation

635 Atomic models, electron density maps and cryo-EM maps were analysed in Coot (*60*) and
636 UCSF ChimeraX (*66*). All figures depicting structural data were generated using UCSF
637 ChimeraX. The ISOLDE (*65*) plugin was used to visualise electron density maps.

Xenosiderophore utilisation bioassay

639 *B. theta tdk⁻*, *B. theta ΔxusAB*, *B. fragilis tdk⁻* and *B. fragilis ΔxusAB* strains were grown
640 overnight in enriched BHI (EBHI; 37 g/l Oxoid BHI powder, 5 g/l yeast extract) supplemented

with 1 µg/ml hemin. 0.3 ml of the overnight cultures were used to inoculate 5 ml of fresh EBHI-hemin medium. The strains were cultured anaerobically at 37 ºC for 3.5 h. Cells were pelleted, resuspended in 5 ml fresh minimal medium with hemin and 0.5% fructose, and diluted to OD~0.05. 10 µl aliquots were spotted on minimal medium agar plates supplemented with 0.5% fructose, 10 µg/ml protoporphyrin IX and either 200 µM $FeCl_3$ or 100 µM BPS. The spots were placed 1.5 cm from the centre of sterile filter discs embedded in agar. Four 5 µl drops of 0.5 mM FeEnt in 50% DMSO, 0.5 mM FeDGE in 50% DMSO, 0.5 mM ferrichrome in sterile water, 50% DMSO only or sterile water only were added to the filter discs. The plates were incubated at 37 ºC anaerobically for 18 h before imaging.

References

1. S. C. Andrews, A. K. Robinson, F. Rodríguez-Quiñones, Bacterial iron homeostasis. *FEMS Microbiol. Rev.* **27**, 215–237 (2003).
2. A. Silale, B. Van Den Berg, TonB-Dependent Transport Across the Bacterial Outer Membrane. *Annu. Rev. Microbiol.* **77**, 67–88 (2023).
3. K. N. Raymond, E. A. Dertz, S. S. Kim, Enterobactin: An archetype for microbial iron transport. *Proc. Natl. Acad. Sci. U S A* **100**, 3584–3588 (2003).
4. T. C. Johnstone, E. M. Nolan, Determination of the Molecular Structures of Ferric Enterobactin and Ferric Enantioenterobactin Using Racemic Crystallography. *J. Am. Chem. Soc.* **139**, 15245–15250 (2017).
5. R. Annamalai, B. Jin, Z. Cao, S. M. C. Newton, P. E. Klebba, Recognition of ferric catecholates by FepA. *J. Bacteriol.* **186**, 3578–3589 (2004).
6. S. K. Buchanan, B. S. Smith, L. Venkatramani, D. Xia, L. Esser, M. Palnitkar, R. Chakraborty, D. Van Der Helm, J. Deisenhofer, Crystal structure of the outer membrane active transporter FepA from *Escherichia coli*. *Nat. Struct. Biol.* **6**, 56–63 (1999).
7. P. Chairatana, T. Zheng, E. M. Nolan, Targeting virulence: salmochelin modification tunes the antibacterial activity spectrum of β-lactams for pathogen-selective killing of *Escherichia coli*. *Chem. Sci.* **6**, 4458–4471 (2015).
8. T. J. Brickman, M. A. McIntosh, Overexpression and purification of ferric enterobactin esterase from *Escherichia coli*. Demonstration of enzymatic hydrolysis of enterobactin and its iron complex. *J. Biol. Chem.* **267**, 12350–12355 (1992).
9. I. G. O'Brien, G. B. Cox, F. Gibson, Enterochelin hydrolysis and iron metabolism in *Escherichia coli*. *BBA General Subjects* **237**, 537–549 (1971).
10. C. C. Murdoch, E. P. Skaar, Nutritional immunity: the battle for nutrient metals at the host–pathogen interface. *Nat. Rev. Microbiol.* **20**, 657–670 (2022).
11. M. Miethke, M. A. Marahiel, Siderophore-Based Iron Acquisition and Pathogen Control. *Microbiol. Mol. Biol. Rev.* **71**, 413–451 (2007).
12. E. D. Weinberg, Nutritional Immunity: Host's Attempt to Withhold Iron From Microbial Invaders. *JAMA* **231**, 39–41 (1975).
13. J. Kramer, Ö. Özkaya, R. Kümmerli, Bacterial siderophores in community and host interactions. *Nat. Rev. Microbiol.* **18**, 152–163 (2019).
14. K. Z. Coyte, S. Rakoff-Nahoum, Understanding Competition and Cooperation within the Mammalian Gut Microbiome. *Current Biol.* **29**, R538–R544 (2019).
15. C. Huttenhower, D. Gevers, R. Knight, S. Abubucker, J. H. Badger, A. T. Chinwalla, H. H. Creasy, A. M. Earl, M. G. Fitzgerald, R. S. Fulton, M. G. Giglio, K. Hallsworth-Pepin, E. A. Lobos, R. Madupu, V. Magrini, J. C. Martin, M. Mitreva, D. M. Muzny, E. J. Sodergren, J. Versalovic, A. M. Wollam, K. C. Worley, J. R. Wortman, S. K. Young, Q. Zeng, K. M. Aagaard, O. O. Abolude, E.

Allen-Vercoe, E. J. Alm, L. Alvarado, G. L. Andersen, S. Anderson, E. Appelbaum, H. M. Arachchi, G. Armitage, C. A. Arze, T. Ayvaz, C. C. Baker, L. Begg, T. Belachew, V. Bhonagiri, M. Bihan, M. J. Blaser, T. Bloom, V. Bonazzi, J. Paul Brooks, G. A. Buck, C. J. Buhay, D. A. Busam, J. L. Campbell, S. R. Canon, B. L. Cantarel, P. S. G. Chain, I. M. A. Chen, L. Chen, S. Chhibba, K. Chu, D. M. Ciulla, J. C. Clemente, S. W. Clifton, S. Conlan, J. Crabtree, M. A. Cutting, N. J. Davidovics, C. C. Davis, T. Z. Desantis, C. Deal, K. D. Delehaunty, F. E. Dewhirst, E. Deych, Y. Ding, D. J. Dooling, S. P. Dugan, W. Michael Dunne, A. Scott Durkin, R. C. Edgar, R. L. Erlich, C. N. Farmer, R. M. Farrell, K. Faust, M. Feldgarden, V. M. Felix, S. Fisher, A. A. Fodor, L. J. Forney, L. Foster, V. Di Francesco, J. Friedman, D. C. Friedrich, C. C. Fronick, L. L. Fulton, H. Gao, N. Garcia, G. Giannoukos, C. Giblin, M. Y. Giovanni, J. M. Goldberg, J. Goll, A. Gonzalez, A. Griggs, S. Gujja, S. Kinder Haake, B. J. Haas, H. A. Hamilton, E. L. Harris, T. A. Hepburn, B. Herter, D. E. Hoffmann, M. E. Holder, C. Howarth, K. H. Huang, S. M. Huse, J. Izard, J. K. Jansson, H. Jiang, C. Jordan, V. Joshi, J. A. Katancik, W. A. Keitel, S. T. Kelley, C. Kells, N. B. King, D. Knights, H. H. Kong, O. Koren, S. Koren, K. C. Kota, C. L. Kovar, N. C. Kyrpides, P. S. La Rosa, S. L. Lee, K. P. Lemon, N. Lennon, C. M. Lewis, L. Lewis, R. E. Ley, K. Li, K. Liolios, B. Liu, Y. Liu, C. C. Lo, C. A. Lozupone, R. Dwayne Lunsford, T. Madden, A. A. Mahurkar, P. J. Mannon, E. R. Mardis, V. M. Markowitz, K. Mavromatis, J. M. McCorrison, D. McDonald, J. McEwen, A. L. McGuire, P. McInnes, T. Mehta, K. A. Mihindukulasuriya, J. R. Miller, P. J. Minx, I. Newsham, C. Nusbaum, M. Oglaughlin, J. Orvis, I. Pagani, K. Palaniappan, S. M. Patel, M. Pearson, J. Peterson, M. Podar, C. Pohl, K. S. Pollard, M. Pop, M. E. Priest, L. M. Proctor, X. Qin, J. Raes, J. Ravel, J. G. Reid, M. Rho, R. Rhodes, K. P. Riehle, M. C. Rivera, B. Rodriguez-Mueller, Y. H. Rogers, M. C. Ross, C. Russ, R. K. Sanka, P. Sankar, J. Fah Sathirapongsasuti, J. A. Schloss, P. D. Schloss, T. M. Schmidt, M. Scholz, L. Schriml, A. M. Schubert, N. Segata, J. A. Segre, W. D. Shannon, R. R. Sharp, T. J. Sharpton, N. Shenoy, N. U. Sheth, G. A. Simone, I. Singh, C. S. Smillie, J. D. Sobel, D. D. Sommer, P. Spicer, G. G. Sutton, S. M. Sykes, D. G. Tabbaa, M. Thiagarajan, C. M. Tomlinson, M. Torralba, T. J. Treangen, R. M. Truty, T. A. Vishnivetskaya, J. Walker, L. Wang, Z. Wang, D. V. Ward, W. Warren, M. A. Watson, C. Wellington, K. A. Wetterstrand, J. R. White, K. Wilczek-Boney, Y. Wu, K. M. Wylie, T. Wylie, C. Yandava, L. Ye, Y. Ye, S. Yooseph, B. P. Youmans, L. Zhang, Y. Zhou, Y. Zhu, L. Zoloth, J. D. Zucker, B. W. Birren, R. A. Gibbs, S. K. Highlander, B. A. Methé, K. E. Nelson, J. F. Petrosino, G. M. Weinstock, R. K. Wilson, O. White, Structure, function and diversity of the healthy human microbiome. *Nature* **486**, 207–214 (2012).

16. G. L. Coe, N. V. Pinkham, A. I. Celis, C. Johnson, J. L. DuBois, S. T. Walk, Dynamic Gut Microbiome Changes in Response to Low-Iron Challenge. *Appl. Environ. Microbiol.* **87**, 1–14 (2021).

17. A. I. Celis, D. A. Relman, K. C. Huang, The impact of iron and heme availability on the healthy human gut microbiome in vivo and in vitro. *Cell Chem. Biol.* **30**, 110-126.e3 (2023).

18. M. M. Meslé, C. R. Gray, M. Dlakić, J. L. DuBois, *Bacteroides thetaiotaomicron*, a Model Gastrointestinal Tract Species, Prefers Heme as an Iron Source, Yields Protoporphyrin IX as a Product, and Acts as a Heme Reservoir. *Microbiol. Spectr.* **11** (2023).

739   19.   N. Martínez-Navarrete, M. M. Camacho, J. Martínez-Lahuerta, J. Martínez-
740   Monzó, P. Fito, Iron deficiency and iron fortified foods—a review. *Food Res. Int.*
741   **35**, 225–231 (2002).
742   20.   I. J. Schalk, Bacterial siderophores: diversity, uptake pathways and applications.
743   *Nat. Rev. Microbiol.* **23**, 24–40 (2024).
744   21.   W. Zhu, M. G. Winter, L. Spiga, E. R. Hughes, R. Chanin, A. Mulgaonkar, J.
745   Pennington, M. Maas, C. L. Behrendt, J. Kim, X. Sun, D. P. Beiting, L. V. Hooper,
746   S. E. Winter, Xenosiderophore Utilization Promotes *Bacteroides*
747   *thetaiotaomicron* Resilience during Colitis. *Cell Host Microbe* **27**, 376-388.e8
748   (2020).
749   22.   L. Spiga, R. T. Fansler, Y. R. Perera, N. G. Shealy, M. J. Munneke, H. E. David,
750   T. P. Torres, A. Lemoff, X. Ran, K. L. Richardson, N. Pudlo, E. C. Martens, E.
751   Folta-Stogniew, Z. J. Yang, E. P. Skaar, M. X. Byndloss, W. J. Chazin, W. Zhu,
752   Iron acquisition by a commensal bacterium modifies host nutritional immunity
753   during *Salmonella* infection. *Cell Host Microbe* **31**, 1639-1654.e10 (2023).
754   23.   D. H. Goetz, M. A. Holmes, N. Borregaard, M. E. Bluhm, K. N. Raymond, R. K.
755   Strong, The Neutrophil Lipocalin NGAL Is a Bacteriostatic Agent that Interferes
756   with Siderophore-Mediated Iron Acquisition. *Mol. Cell.* **10**, 1033–1043 (2002).
757   24.   T. H. Flo, K. D. Smith, S. Sato, D. J. Rodriguez, M. A. Holmes, R. K. Strong, S.
758   Akira, A. Aderem, Lipocalin 2 mediates an innate immune response to bacterial
759   infection by sequestrating iron. *Nature* **432**, 917–921 (2004).
760   25.   J. P. Henderson, J. R. Crowley, J. S. Pinkner, J. N. Walker, P. Tsukayama, W. E.
761   Stamm, T. M. Hooton, S. J. Hultgren, Quantitative Metabolomics Reveals an
762   Epigenetic Blueprint for Iron Acquisition in Uropathogenic *Escherichia coli.*
763   *PLoS Pathog* **5**, e1000305 (2009).
764   26.   M. A. Fischbach, H. Lin, L. Zhou, Y. Yu, R. J. Abergel, D. R. Liu, K. N. Raymond,
765   B. L. Wanner, R. K. Strong, C. T. Walsh, A. Aderem, K. D. Smith, The pathogen-
766   associated *iroA* gene cluster mediates bacterial evasion of lipocalin 2. *Proc. Natl.*
767   *Acad. Sci. U S A* **103**, 16502–16507 (2006).
768   27.   K. Hantke, G. Nicholson, W. Rabsch, G. Winkelmann, Bioinorganic Chemistry
769   Special Feature: Salmochelins, siderophores of *Salmonella enterica* and
770   uropathogenic *Escherichia coli* strains, are recognized by the outer membrane
771   receptor IroN. *Pro.c Natl. Acad. Sci. U S A* **100**, 3677 (2003).
772   28.   J. Abellon-Ruiz, K. Jana, A. Silale, A. M. Frey, A. Baslé, M. Trost, U.
773   Kleinekathöfer, B. van den Berg, BtuB TonB-dependent transporters and BtuG
774   surface lipoproteins form stable complexes for vitamin B12 uptake in gut
775   Bacteroides. *Nat. Commun.* **14**, 1–18 (2023).
776   29.   J. B. R. White, A. Silale, M. Feasey, T. Heunis, Y. Zhu, H. Zheng, A. Gajbhiye, S.
777   Firbank, A. Baslé, M. Trost, D. N. Bolam, B. van den Berg, N. A. Ranson, Outer
778   membrane utilisomes mediate glycan uptake in gut Bacteroidetes. *Nature* **618**,
779   583–589 (2023).
780   30.   M. J. Coyne, C. M. Fletcher, M. Chatzidaki-Livanis, G. Posch, C. Schaffer, L. E.
781   Comstock, Phylum-wide general protein O-glycosylation system of the
782   Bacteroidetes. *Mol. Microbiol.* **88**, 772–783 (2013).
783   31.   G. Posch, M. Pabst, L. Neumann, M. J. Coyne, F. Altmann, P. Messner, L. E.
784   Comstock, C. Schäffer, "Cross-glycosylation" of proteins in Bacteroidales
785   species. *Glycobiology* **23**, 568 (2013).
786   32.   A. Bateman, M. J. Martin, S. Orchard, M. Magrane, S. Ahmad, E. Alpi, E. H.
787   Bowler-Barnett, R. Britto, H. Bye-A-Jee, A. Cukura, P. Denny, T. Dogan, T. G.
788   Ebenezer, J. Fan, P. Garmiri, L. J. da Costa Gonzales, E. Hatton-Ellis, A. Hussein,

A. Ignatchenko, G. Insana, R. Ishtiaq, V. Joshi, D. Jyothi, S. Kandasaamy, A. Lock, A. Luciani, M. Lugaric, J. Luo, Y. Lussi, A. MacDougall, F. Madeira, M. Mahmoudy, A. Mishra, K. Moulang, A. Nightingale, S. Pundir, G. Qi, S. Raj, P. Raposo, D. L. Rice, R. Saidi, R. Santos, E. Speretta, J. Stephenson, P. Totoo, E. Turner, N. Tyagi, P. Vasudev, K. Warner, X. Watkins, R. Zaru, H. Zellner, A. J. Bridge, L. Aimo, G. Argoud-Puy, A. H. Auchincloss, K. B. Axelsen, P. Bansal, D. Baratin, T. M. Batista Neto, M. C. Blatter, J. T. Bolleman, E. Boutet, L. Breuza, B. C. Gil, C. Casals-Casas, K. C. Echioukh, E. Coudert, B. Cuche, E. de Castro, A. Estreicher, M. L. Famiglietti, M. Feuermann, E. Gasteiger, P. Gaudet, S. Gehant, V. Gerritsen, A. Gos, N. Gruaz, C. Hulo, N. Hyka-Nouspikel, F. Jungo, A. Kerhornou, P. Le Mercier, D. Lieberherr, P. Masson, A. Morgat, V. Muthukrishnan, S. Paesano, I. Pedruzzi, S. Pilbout, L. Pourcel, S. Poux, M. Pozzato, M. Pruess, N. Redaschi, C. Rivoire, C. J. A. Sigrist, K. Sonesson, S. Sundaram, C. H. Wu, C. N. Arighi, L. Arminski, C. Chen, Y. Chen, H. Huang, K. Laiho, P. McGarvey, D. A. Natale, K. Ross, C. R. Vinayaka, Q. Wang, Y. Wang, J. Zhang, UniProt: the Universal Protein Knowledgebase in 2023. *Nucleic Acids Res.* **51**, D523–D531 (2023).

33. F. Lauber, G. R. Cornelis, F. Renzi, Identification of a new lipoprotein export signal in gram-negative bacteria. *mBio* **7**, 1232–1248 (2016).

34. I. Josts, K. Veith, V. Normant, I. J. Schalk, H. Tidow, Structural insights into a novel family of integral membrane siderophore reductases. *Proc. Natl. Acad. Sci. U S A* **118**, e2101952118 (2021).

35. E. R. Rocha, A. S. Krykunivsky, Anaerobic utilization of Fe(III)-xenosiderophores among *Bacteroides* species and the distinct assimilation of Fe(III)-ferrichrome by *Bacteroides fragilis* within the genus. *Microbiologyopen* **6**, e00479 (2017).

36. J. Jumper, R. Evans, A. Pritzel, T. Green, M. Figurnov, O. Ronneberger, K. Tunyasuvunakool, R. Bates, A. Žídek, A. Potapenko, A. Bridgland, C. Meyer, S. A. A. Kohl, A. J. Ballard, A. Cowie, B. Romera-Paredes, S. Nikolov, R. Jain, J. Adler, T. Back, S. Petersen, D. Reiman, E. Clancy, M. Zielinski, M. Steinegger, M. Pacholska, T. Berghammer, S. Bodenstein, D. Silver, O. Vinyals, A. W. Senior, K. Kavukcuoglu, P. Kohli, D. Hassabis, Highly accurate protein structure prediction with AlphaFold. *Nature* **596**, 583–589 (2021).

37. N. M. Koropatkin, E. C. Martens, J. I. Gordon, T. J. Smith, Starch Catabolism by a Prominent Human Gut Symbiont Is Directed by the Recognition of Amylose Helices. *Structure* **16**, 1105–1115 (2008).

38. R. M. Pollet, L. M. Martin, N. M. Koropatkin, TonB-dependent transporters in the Bacteroidetes: Unique domain structures and potential functions. *Mol. Microbiol.* **115**, 490-501 (2021).

39. M. Madej, J. B. R. White, Z. Nowakowska, S. Rawson, C. Scavenius, J. J. Enghild, G. P. Bereta, K. Pothula, U. Kleinekathoefer, A. Baslé, N. A. Ranson, J. Potempa, B. van den Berg, Structural and functional insights into oligopeptide acquisition by the RagAB transporter from *Porphyromonas gingivalis*. *Nat. Microbiol.* **5**, 1016–1025 (2020).

40. A. J. Glenwright, K. R. Pothula, S. P. Bhamidimarri, D. S. Chorev, A. Baslé, S. J. Firbank, H. Zheng, C. V. Robinson, M. Winterhalter, U. Kleinekathöfer, D. N. Bolam, B. Van Den Berg, Structural basis for nutrient acquisition by dominant members of the human gut microbiota. *Nature* **541**, 407–411 (2017).

41. M. Tong, J. Xu, W. Li, K. Jiang, Y. Yang, Z. Chen, X. Jiao, X. Meng, M. Wang, J. Hong, H. Long, S.-J. Liu, B. Lim, X. Gao, A highly conserved SusCD

transporter determines the import and species-specific antagonism of *Bacteroides*
ubiquitin homologues. *Nat. Commun.* **15**, 1–17 (2024).

42. R. J. Abergel, E. G. Moore, R. K. Strong, K. N. Raymond, Microbial Evasion of
the Immune System: Structural Modifications of Enterobactin Impair Siderocalin
Recognition. *J. Am. Chem. Soc.* **128**, 10998 (2006).

43. M. Llinás, J. B. Neilands, Structure of ferricrocin. *Bioinorg. Chem.* **2**, 159–165
(1973).

44. R. Norrestam, B. Stensland, C. I. Brändén, On the conformation of cyclic iron-
containing hexapeptides: The crystal and molecular structure of ferrichrysin. *J
Mol Biol* **99**, 501–506 (1975).

45. D. Al Shaer, O. Al Musaimi, B. G. de la Torre, F. Albericio, Hydroxamate
siderophores: Natural occurrence, chemical synthesis, iron binding affinity and
use as Trojan horses against pathogens. *Eur. J. Med. Chem.* **208**, 112791 (2020).

46. M. Valentine, E. Benadé, M. Mouton, W. Khan, A. Botha, Binary interactions
between the yeast *Candida albicans* and two gut-associated *Bacteroides* species.
Microb. Pathog. **135**, 103619 (2019).

47. M. J. Coyne, K. G. Roelofs, L. E. Comstock, Type VI secretion systems of human
gut Bacteroidales segregate into three genetic architectures, two of which are
contained on mobile genetic elements. *BMC Genomics* **17**, 1–21 (2016).

48. K. Trunk, J. Peltier, Y. C. Liu, B. D. Dill, L. Walker, N. A. R. Gow, M. J. R. Stark,
J. Quinn, H. Strahl, M. Trost, S. J. Coulthurst, The type VI secretion system
deploys antifungal effectors against microbial competitors. *Nat. Microbiol.* **3**,
920–931 (2018).

49. K. Trunk, S. J. Coulthurst, J. Quinn, A New Front in Microbial Warfare—Delivery
of Antifungal Effectors by the Type VI Secretion System. *J. Fungi* **5**, 50 (2019).

50. F. W. Studier, B. A. Moffatt, Use of bacteriophage T7 RNA polymerase to direct
selective high-level expression of cloned genes. *J. Mol. Biol.* **189**, 113–130
(1986).

51. D. A. Gray, J. B. R. White, A. O. Oluwole, P. Rath, A. J. Glenwright, A. Mazur,
M. Zahn, A. Baslé, C. Morland, S. L. Evans, A. Cartmell, C. V. Robinson, S.
Hiller, N. A. Ranson, D. N. Bolam, B. van den Berg, Insights into SusCD-
mediated glycan import by a prominent gut symbiont. *Nat. Commun.* **12**, 1–14
(2021).

52. V. de Lorenzo, L. Eltis, B. Kessler, K. N. Timmis, Analysis of *Pseudomonas* gene
products using *lacIq/Ptrp-lac* plasmids and transposons that confer conditional
phenotypes. *Gene* **123**, 17–24 (1993).

53. M. A. Fischbach, H. Lin, D. R. Liu, C. T. Walsh, In vitro characterization of IroB,
a pathogen-associated C-glycosyltransferase. *Proc. Natl. Acad. Sci. U S A* **102**,
571–576 (2005).

54. R. N. Motz, J. K. Anderson, E. M. Nolan, Re-evaluation of the C-
Glucosyltransferase IroB Illuminates Its Ability to C-Glucosylate Non-native
Triscatecholate Enterobactin Mimics. *Biochemistry* **64**, 224–237 (2025).

55. C. K. Murphy, V. I. Kalve, P. E. Klebba, Surface topology of the Escherichia coli
K-12 ferric enterobactin receptor. *J. Bacteriol.* **172**, 2736–2746 (1990).

56. T. Emery, J. B. Neilands, Contribution to the Structure of the Ferrichrome
Compounds: Characterization of the Acyl Moieties of the Hydroxamate
Functions. *J. Am. Chem. Soc.* **82**, 3658–3662 (1960).

57. G. Winter, Xia2: An expert system for macromolecular crystallography data
reduction. *J. Appl. Crystallogr.* **43**, 186–190 (2010).

58. P. R. Evans, G. N. Murshudov, How good are my data and what is the resolution? *Acta Crystallogr. D Biol. Crystallogr.* **69**, 1204–1214 (2013).

59. P. Evans, Scaling and assessment of data quality. *Acta Crystallogr. D Biol. Crystallogr.* **62**, 72–82 (2006).

60. P. Emsley, B. Lohkamp, W. G. Scott, K. Cowtan, Features and development of Coot. *Acta Crystallogr. D Biol. Crystallogr.* **66**, 486–501 (2010).

61. D. Liebschner, P. V. Afonine, M. L. Baker, G. Bunkoczi, V. B. Chen, T. I. Croll, B. Hintze, L. W. Hung, S. Jain, A. J. McCoy, N. W. Moriarty, R. D. Oeffner, B. K. Poon, M. G. Prisant, R. J. Read, J. S. Richardson, D. C. Richardson, M. D. Sammito, O. V. Sobolev, D. H. Stockwell, T. C. Terwilliger, A. G. Urzhumtsev, L. L. Videau, C. J. Williams, P. D. Adams, Macromolecular structure determination using X-rays, neutrons and electrons: recent developments in Phenix. *Acta Crystallogr. D Struct. Biol.* **75**, 861–877 (2019).

62. V. B. Chen, W. B. Arendall, J. J. Headd, D. A. Keedy, R. M. Immormino, G. J. Kapral, L. W. Murray, J. S. Richardson, D. C. Richardson, MolProbity: all-atom structure validation for macromolecular crystallography. *Acta Crystallogr. D Biol. Crystallogr.* **66**, 12–21 (2010).

63. A. Punjani, H. Zhang, D. J. Fleet, Non-uniform refinement: adaptive regularization improves single-particle cryo-EM reconstruction. *Nat. Methods* **17**, 1214–1221 (2020).

64. K. Jamali, L. Käll, R. Zhang, A. Brown, D. Kimanius, S. H. W. Scheres, Automated model building and protein identification in cryo-EM maps. *Nature* **628**, 450–457 (2024).

65. T. I. Croll, ISOLDE: A physically realistic environment for model building into low-resolution electron-density maps. *Acta Crystallogr. D Struct. Biol.* **74**, 519–530 (2018).

66. E. F. Pettersen, T. D. Goddard, C. C. Huang, E. C. Meng, G. S. Couch, T. I. Croll, J. H. Morris, T. E. Ferrin, UCSF ChimeraX: Structure visualization for researchers, educators, and developers. *Protein Sci.* **30**, 70–82 (2021).

Acknowledgements

The *B. fragilis tdk⁻* strain was a gift from Janet Quinn (Newcastle). We thank Matthew Nodwell (Simon Fraser University) for providing the Ent used in this study. We also thank Diamond Light Source for access to macromolecular crystallography beamlines (proposals mx-24948 and mx-32736). We acknowledge use of the crystallization facilities and the GPU cluster for cryo-EM data processing at the Newcastle University Structural Biology Facility. We acknowledge use of the Glacios microscope for screening cryo-EM grids at the York Structural Biology Laboratory (University of York). We acknowledge use of the Astbury Centre (University of Leeds) Titan Krios microscope.

Funding:

Wellcome Trust Investigator award 214222/Z/18/Z (B.v.d.B.)
National Institutes of Health grant R01AI176390 (E.M.N)
National Science Foundation Graduate Research Fellowship (R.N.M.)

Author contributions:

A.S. conceived the project, designed experiments, made constructs, purified and crystallized proteins, made cryo-EM grids and collected cryo-EM data, determined crystal and cryo-EM structures, and wrote the manuscript with input from R.N.M,

E.M.N. and B.v.d.B. Y.L.S. and H.M. made expression constructs, purified and
crystallized proteins, and carried out ITC titrations, supervised by A.S. and B.v.d.B.
R.N.M. synthesized DGE, supervised by E.M.N. A.B. collected X-ray diffraction data
and provided computational support. B.v.d.B. acquired funding, designed experiments,
and collected cryo-EM data.

Competing interests:
The authors declare that they have no competing interests.

Data availability:
Electron microscopy volumes of the XusAB complex have been deposited in the
Electron Microscopy Data Bank with the accession code EMD-51210, and the atomic
coordinates have been deposited in the Protein Data Bank under the accession code
9GBC. Atomic coordinates and the associated crystallographic structure factors have
been deposited in the Protein Data Bank under the following accession codes: 9GCY
(apo BtXusB), 9GCZ (BtXusB-FeEnt), 9HQ1 (BtXusB-FeDGE), 9GAR (apo BvXusB),
9HQE (apo BfXusB), and 9HQK (BfXusB-ferrichrome). Other data and materials
related to the manuscript are available from the corresponding authors upon request.

Supplementary Materials for

Structural basis of iron piracy by human gut *Bacteroides*

Augustinas Silale *et al.*

*Corresponding authors: augustinas.silale@newcastle.ac.uk
bert.van-den-berg@newcastle.ac.uk

This file includes:

962 **Other Supplementary Materials for this manuscript include the following:**

A

FeEnt			BtXusB			Interface area (Å^2)	Δ^{i}G, kcal/mol	N_{HB}	N_{SB}	CSS
$^iN_{at}$	$^iN_{res}$	Surface (Å^2)	$^iN_{at}$	$^iN_{res}$	Surface (Å^2)					
42	N/A	718	59	16	17,327	418.8	3.1	10	0	0.005

B

FeEnt	Dist. (Å)	BtXusB
EB4 802 (N2)	3.13	GLN 413 (OE1)
EB4 802 (O2)	3.57	GLN 413 (OE1)
EB4 802 (O4)	2.44	SER 121 (OG)
EB4 802 (O5)	2.57	SER 168 (OG)
EB4 802 (O6)	2.59	TYR 411 (OH)
EB4 802 (O15)	3.40	ASN 120 (ND2)
EB4 802 (O4)	3.33	SER 121 (N)
EB4 802 (O4)	3.23	SER 168 (N)
EB4 802 (O5)	3.02	SER 168 (N)
EB4 802 (O7)	2.92	ARG 255 (NE)

964
965 **Fig. S1. PISA (*1*) analysis of BtXusB-FeEnt interactions.**
966 (**A**) Complex surface area calculations. (**B**) Hydrogen bonds formed between FeEnt and BtXusB.

968
969 **Fig. S2. Comparison of apo and FeEnt-bound BtXusB structures.**
970 Structural alignment of apo (hot pink) and FeEnt-bound (grey) BtXusB crystal structures. Cα-Cα
971 RMSD = 1.07 Å. FeEnt is in orange; BtXusB FeEnt-binding loops are highlighted in cyan in the co-
972 crystal structure.

973
Fig. S3. Expression and purification of the native XusAB complex.
(**A**) *B. theta tdk⁻* strain grown anaerobically in minimal medium with 0.5% fructose and 1 µg/ml hemin.
The iron chelator bathophenanthroline disulfonate (BPS) was added at indicated concentrations. (**B**) *B.*
theta tdk⁻ strain grown anaerobically in the presence of BPS and indicated iron sources. All growth
curves shown are averages from 3 wells of a single 96-well growth experiment. The experiment was
repeated three times under identical conditions with similar results. (**C**) Western blot of whole cell
lysates from *B. theta bt2064-his* cells grown overnight in minimal medium and the indicated
concentrations of BPS. (**D**) Size exclusion chromatography trace of XusAB complex after immobilized
metal affinity chromatography on a Superdex 200 10/300 Increase column. (**E**) SDS-PAGE analysis of
0.5 ml elution fractions from (**D**) indicated by the dashed arrow (approximately 10-12 ml elution
volume). The indicated bands were identified as XusA and XusB by peptide mass fingerprinting. Both
bands around 100 kDa correspond to XusA.

A

6,645 movies
Patch motion correction
Patch CTF estimation

2,000 particles
manually picked

Template picker

2D classification
Generate 2 templates
for particle picking

1,110,843 particles
240 pix box, 1.42 Å/pix

2D classification (2x)
174,239 particles

Ab initio reconstruction
with good particles   with bad particles

Heterogeneous refinement
174,239 particles
2 noise decoys, 2 good ab initio models

NU refinement
102,772 particles

3.03 Å

3D classification
8 classes

Re-extract particles
480 pix box, 0.74 Å/pix
94,987 particles

NU refinement + local refinement

2.83 Å

Reference-based motion correction
Local refinement
CTF refinement
Per-particle defocus
Beam tilt, trefoil

2D classification
Disabled noise model annealing

Local refinement
Reference-based motion correction

Local refinement
76,976 particles
2.7 Å

Final map

B

60 nm

C

150 Å

D

Gold-standard FSC resolution: 2.70 Å

- No Mask (3.6Å)
- Spherical (3.3Å)
- Loose (2.8Å)
- Tight (2.7Å)

FSC

FSC=0.143

DC   7.1Å   3.6Å   2.4Å   1.8Å
Resolution

E

Elevation

Azimuth

# of images

F

180°

Local resolution, Å
2.2 2.6 3.0 3.4 3.8

 Fig. S4. XusAB cryo-EM data processing.
(A) Data processing workflow carried out in cryoSPARC v4.4.1 (*2*). **(B)** Representative motion-corrected micrograph (n=6,645). **(C)** Representative 2D class averages. **(D)** Gold-standard Fourier shell correlation curve and **(E)** viewing direction distribution plot for the final map. **(F)** The final cryo-EM map coloured by estimated local resolution.

A

XusB			XusA			Interface area (Å²)	ΔᴵG, kcal/mol	N_HB	N_SB	CSS
ᴵN_at	ᴵN_res	Surface (Å²)	ᴵN_at	ᴵN_res	Surface (Å²)					
327	98	18,536	292	81	32,277	2,794.6	-12.4	52	6	1.000

B

XusB	Dist. (Å)	XusA
THR 438 (OG1)	3.10	GLU 289 (OE2)
MET 54 (N)	3.21	SER 297 (O)
GLU 436 (N)	3.02	ILE 299 (O)
LYS 432 (NZ)	3.83	THR 301 (OG1)
VAL 59 (N)	3.14	LYS 416 (O)
ASN 412 (ND2)	3.70	GLU 421 (O)
GLN 413 (NE2)	3.85	SER 423 (OG)
ARG 255 (NH2)	3.47	ASP 568 (O)
ARG 167 (NH2)	3.04	GLU 621 (OE2)
GLY 258 (N)	3.57	THR 623 (O)
ALA 259 (N)	3.60	THR 623 (O)
GLY 258 (N)	3.31	SER 624 (O)
THR 260 (OG1)	2.88	ASN 625 (OD1)
THR 249 (OG1)	3.23	ASN 625 (OD1)
THR 310 (N)	2.95	THR 674 (O)
THR 310 (N)	3.81	THR 674 (OG1)
ASN 387 (ND2)	3.20	GLU 728 (O)
ASN 387 (N)	3.16	ALA 729 (O)
ASN 386 (N)	3.43	ALA 729 (O)
ASN 387 (ND2)	2.82	GLY 731 (O)
GLU 391 (N)	3.20	THR 735 (OG1)
MET 54 (O)	3.11	SER 297 (N)
GLU 436 (O)	3.13	ILE 299 (N)
SER 434 (O)	3.33	THR 301 (N)
GLU 436 (OE1)	2.84	THR 301 (OG1)
GLU 436 (OE1)	3.15	LYS 354 (NZ)

XusB	Dist. (Å)	XusA
ASN 56 (O)	3.05	LYS 416 (NZ)
ARG 57 (O)	2.80	ARG 418 (NH1)
ARG 57 (O)	2.90	ARG 418 (NH2)
ILE 44 (O)	3.00	ARG 418 (NH2)
ASP 459 (O)	3.86	LYS 419 (N)
ASP 194 (OD1)	3.15	SER 516 (OG)
ASP 194 (OD1)	3.22	THR 517 (OG1)
ILE 192 (O)	3.11	THR 517 (OG1)
ASP 194 (OD1)	3.54	SER 518 (N)
ASP 194 (OD1)	2.84	SER 518 (OG)
ASP 194 (OD2)	3.08	THR 520 (OG1)
ASP 194 (OD2)	3.50	SER 521 (N)
SER 289 (O)	3.00	THR 623 (N)
ALA 259 (O)	3.69	THR 623 (OG1)
SER 289 (O)	3.79	THR 623 (OG1)
SER 289 (O)	3.47	SER 624 (N)
PRO 308 (O)	2.55	THR 674 (OG1)
PRO 308 (O)	3.30	GLN 676 (NE2)
ASN 316 (O)	2.81	GLN 676 (NE2)
THR 310 (OG1)	3.17	GLN 676 (NE2)
PRO 384 (O)	3.29	GLY 731 (N)
GLY 389 (O)	2.87	ALA 732 (N)
GLY 389 (O)	3.66	THR 735 (OG1)
GLU 391 (OE2)	2.99	LYS 736 (NZ)
GLU 391 (OE1)	3.57	ASP 737 (N)

C

XusB	Dist. (Å)	XusA
ARG 167 (NH2)	3.04	GLU 621 (OE2)
GLU 436 (OE2)	3.39	LYS 354 (NZ)
GLU 436 (OE1)	3.15	LYS 354 (NZ)
GLU 416 (OE2)	3.12	LYS 354 (NZ)
GLU 391 (OE1)	3.95	LYS 736 (NZ)
GLU 391 (OE2)	2.99	LYS 736 (NZ)

993
994 **Fig. S5. PISA** (*1*) **analysis of XusA-XusB interactions.**
995 (**A**) Complex surface area calculations. (**B**) Hydrogen bonds formed between XusA and XusB. (**C**) Salt
996 bridges formed between XusA and XusB.

997
Fig. S6. XusA O-glycosylation.
(**A**) The glycan density extending from the sidechain hydroxy oxygen of T401 is shown in grey. Sugar
units that likely constitute the glycan chain are indicated: Hex, hexose; HexA, hexuronic acid;
(me)dHex, (methyl)deoxyhexose. Sugars were assigned based on the structure of the *B. fragilis* O-
glycan (*3*). (**B**) Residues 421-423, which are in the C-terminal part of the β7-8 extracellular loop of
XusA and are implicated in FeEnt release from XusB, are somewhat close to T401. The sidechain of
T401 faces away from residues 421-423 towards the solvent.

Fig. S7. XusB homologue sequence alignment.

Alignment of BtXusB, *P. distasonis* DSM 20701 BDI_3402, *B. viscericola* DSM 18177 BARVI_05925, and *B. fragilis* NCTC 9343 BF9343_4228 amino acid sequences. The secondary structure features of BtXusB in the BtXusB-FeEnt structure are annotated above the alignment. Visualised in ESPript v3.0 (*4*) (https://espript.ibcp.fr).

Parabacteroides distasonis
AF-A6LHE0-F1

Barnesiella viscericola
AF-W0ET74-F1

Bacteroides fragilis
AF-Q5L7E3-F1

25   61.8   98.9

1011
1012 **Fig. S8. XusB homologue AlphaFold2 (*5*) model prediction confidence.**
1013 The indicated entries were downloaded from AlphaFold DB. The views were generated from
1014 superposition with the BtXusB-FeEnt crystal structure. Each model is coloured according to pLDDT
1015 value (colour key). The blue (low confidence) N-terminal regions correspond to the signal peptides.
1016

1017

Fig. S9. XusB BLAST hit amino acid sequence length distribution.
The BtXusB amino acid sequence was submitted to the EFI Enzyme Similarity Tool server (*6*). BLAST results were filtered to remove hits with an E-value higher than 10^{-5} and sequences shorter than 380 and longer than 500 amino acids, resulting in 681 hits. The amino acid sequence lengths of these hits are plotted in the histogram, clearly showing a bimodal distribution with peaks around 410 and 465 amino acids. The shorter group includes BfXusB, while BtXusB, BvXusB and *P. distasonis* XusB belong to the longer group.

1025
1026 **Fig. S10. Comparison of BvXusB apo structure and ligand-bound structure predictions.**
1027 BvXusB apo crystal structure (cyan) superposed with AF3-predicted BvXusB-FeEnt (**A**) and
1028 BvXusB-FeDGE (**B**) models.

1029

1030 **Fig. S11. BvXusB does not bind ferrichrome.**
1031 Representative ITC experiment where 250 µM ferrichrome was titrated into 25 µM BvXusB (n=2). No
1032 substantial injection heats were observed, consistent with lack of binding.

33   **Table S1**.

34   X-ray data collection and refinement parameters. Values for highest resolution shell in parentheses.

	Apo BtXusB	BtXusB-FeEnt	BtXusB-FeDGE	Apo BvXusB	Apo BfXusB	BfXusB-ferrichrome
Data collection						
DLS beamline	I03	I24	I03	I03	I03	I03
Wavelength	0.9537	0.6199	0.9686	0.9686	0.9686	0.9686
Space Group	P2$_1$	P2$_1$	P2$_1$	I4$_1$22	P2$_1$2$_1$2$_1$	P2$_1$
Unit cell parameters						
a, b, c (Å)	45.7, 108.63, 93.89	59.99, 48.18, 135.47	45.44, 108.71, 93.83	184.87, 184.87, 82.31	50.99, 86.93, 154.39	57.20, 155.97, 89.62
α, β, γ (°)	90, 96.57, 90	90, 96.79, 90	90, 96.66, 90	90, 90, 90	90, 90, 90	90, 95.56, 90
Molecules in AU	2	2	2	1	2	4
Resolution range (Å)	54.32-1.56 (1.59-1.56)	39.17-1.50 (1.53-1.50)	54.35-1.80 (1.83-1.80)	58.46-3.2 (3.43-3.2)	48.42-1.77 (1.81-1.77)	58.71-3.32 (3.59-3.32)
I/σI	11.0 (1.2)	6.7 (1.4)	9.4 (1.2)	6.8 (1.8)	15.2 (2.0)	5.3 (2.4)
Completeness (%)	100 (99.5)	99.6 (94.4)	99.9 (99.4)	99.9 (99.3)	100 (99.9)	100 (100)
Multiplicity	6.8 (5.3)	6.7 (4.9)	6.8 (6.9)	27.3 (28.3)	13.4 (13.9)	7.1 (7.4)
R$_{pim}$	0.039 (0.652)	0.096 (0.955)	0.045 (0.468)	0.104 (0.444)	0.029 (0.355)	0.116 (0.318)
R$_{merge}$	0.091 (1.283)	0.162 (1.049)	0.110 (1.151)	0.538 (2.335)	0.102 (1.289)	0.290 (0.812)
CC$_{1/2}$	0.998 (0.621)	0.774 (0.545)	0.998 (0.609)	0.998 (0.751)	0.999 (0.799)	0.981 (0.862)
No. of unique reflections	129,093 (6,316)	122,900 (5,684)	83,651 (4,580)	12,010 (2,123)	67,890 (3,826)	23,146 (4,780)
Phasing						
Molecular replacement model	AlphaFold2 model (AF-Q8A622-F1)	apo XusB structure (PDB 9GCY)	apo XusB structure (PDB 9GCY)	AlphaFold2 model (AF-W0ET74-F1)	AlphaFold2 model (AF-Q5L7E3-F1)	Apo BfrXusB structure (PDB 9HQE)
Refinement						
Resolution (Å)	46.94-1.56	33.94-1.50	54.36-1.80	49.33-3.20	48.42-1.77	58.71-3.32
Rwork/Rfree	0.1757/0.2068	0.2038/0.2439	0.1916/0.2282	0.2338/0.2768	0.1987/0.2368	0.2484/0.2958
Reflections	128,930	121,657	83,497	11,963	67,737	22,931
Non-hydrogen atoms	7,736	7,583	7,471	3,594	6,081	11,812
Protein only	6,522	6,585	6,605	3,585	5,708	11,616
Mean B-factor (Å^2)	25.41	17.44	26.00	64.54	32.43	51.06
Rmsd						
Bond lengths (Å)	0.010	0.006	0.007	0.001	0.006	0.003
Bond angles (°)	0.99	0.80	0.81	0.39	0.79	0.63
Clashscore	1.94	2.14	2.00	4.29	3.01	8.45
Rotamer outliers (%)	0.14	0.71	0.85	0	0.16	0
Ramachandran plot						
Favoured (%)	96.90	97.26	96.79	95.15	94.38	94.00
Outliers (%)	0	0	0.12	0	0.14	0.27
PDB ID	9GCY	9GCZ	9HQ1	9GAR	9HQE	9HQK

1035 **Table S2.** Isothermal titration calorimetry data fitting results.

Titration	Replicate	[Protein] (M)	[Ligand] (M)	N	Kd (M)	ΔH (kcal/mol)	ΔG (kcal/mol)	-TΔS (kcal/mol)
BtXusB + FeEnt	1	2.50E-04	2.50E-05	1.3	9.82E-08	-24.5	-9.57	14.9
BtXusB + FeEnt	2	2.50E-04	2.50E-05	1.46	8.42E-08	-22.8	-9.67	13.1
BtXusB + FeEnt	3	2.50E-04	2.50E-05	1.47	4.94E-08	-21	-9.97	11
BtXusB + FeEnt	4	2.50E-04	2.50E-05	1.22	5.52E-08	-22.6	-9.9	12.7
BtXusB + FeDGE	1	1.54E-04	1.73E-05	1 (fixed)	1.69E-07	-20.1	-9.24	10.8
BtXusB + FeDGE	2	1.57E-04	1.73E-05	1 (fixed)	1.99E-07	-19.5	-9.14	10.4
BvXusB + FeEnt	1	2.50E-04	2.50E-05	1.35	1.01E-07	-15.5	-9.55	5.98
BvXusB + FeEnt	2	2.50E-04	2.50E-05	1.21	7.87E-08	-14.8	-9.69	5.15
BfXusB + ferrichrome	1	2.50E-04	2.50E-05	1.18	2.68E-07	-11.5	-8.97	2.52
BfXusB + ferrichrome	2	3.48E-04	2.54E-05	1.27	2.99E-07	-10.1	-8.9	1.24
BfXusB + ferrichrome	3	2.46E-04	2.54E-05	1.09	1.76E-07	-10.3	-9.22	1.11
BfXusB + ferrichrome	4	2.26E-04	2.27E-05	1.06	2.98E-07	-12.7	-8.9	3.78

1036

1037   **Table S3.** Cryo-EM data collection and structure refinement parameters.

	XusAB
Data collection	
Electron microscope	FEI Titan Krios
Voltage (kV)	300
Spherical aberration (μm)	2.7
Camera	Falcon 4i (counting)
Energy filter	Selectris (10 eV slit)
Magnification	165,000
Pixel size (Å)	0.74
Total dose (e$^-$/Å^2)	40.83
Defocus range (μm)	-2.0 to -0.8
Number of movies collected	6,645
Image Processing	
Symmetry	C1
Initial number of particles	1,110,843
Final number of particles	76,969
Global resolution (FSC = 0.143)	2.7 Å
Map sharpening B-factor (Å^2)	-57.2
Refinement	
Model composition	
Non-hydrogen atoms	8,742
Protein residues	1,093
R.m.s. deviations	
Bonds lengths (Å)	0.002
Bond angles (°)	0.543
Validation	
MolProbity score	1.12
Clash score	2.84
Rotamer outliers (%)	0
Ramachandran plot	
Favoured (%)	97.79
Outliers (%)	0
PDB	9GBC
EMDB	EMD-51210

1038

1039 **Table S4.** Strains used in this study.

Strain	Description	Source
E. coli TOP10	Used for constructing expression plasmids	Invitrogen
E. coli BL21(DE3)	Used for recombinant protein production	(*7*)
E. coli S17-1λpir	Used for constructing and conjugating pExchange plasmids into *Bacteroides* spp.	(*8*)
B. theta VPI-5482 *tdk⁻*	Thymidine kinase knockout used for allelic exchange with pExchange vector.	(*9*)
B. theta VPI-5482 *tdk⁻ bt2064-his*	Chromosomal His$_6$-tag for XusAB complex purification	This study
B. theta VPI-5482 *tdk⁻ bt2064⁻ bt2065⁻*	XusAB knockout	This study
B. fragilis NCTC 9343 *tdk⁻*	Thymidine kinase knockout used for allelic exchange with pExchange vector.	Janet Quinn (Newcastle), unpublished
B. fragilis NCTC 9343 *tdk⁻ bf9343_4228⁻ bf4393_4229⁻*	XusAB knockout	This study

1040

1041 **Table S5.** Plasmids used in this study.

Strain	Description	Source
pET28b	Used for recombinant protein production	EMD Biosciences
pET28b_BtXusB	Used for expression of BtXusB (residues 35-464)	This study
pET28b_BvXusB	Used for expression of BvXusB (residues 29-491)	This study
pET28b_BfXusB	Used for expression of BfXusB (residues 39-406)	This study
pExchange	Used for *Bacteroides* spp. allelic exchange.	(*9*)
pExchange_bt2064-his	Used to insert a chromosomal His$_6$-tag on BtXusB	This study
pExchange_bt2065-64_KO	Used to delete *xusAB* from *B. theta* chromosome	This study
pExchange_bf9343_4229-4228	Used to delete *xusAB* from *B. fragilis* chromosome	This study

1042

Movie S1. Morph of the apo BtXusB crystal structure to the FeEnt-bound BtXusB crystal structure.
The protein model is depicted as a cartoon and as a surface. The siderophore-binding loops are in blue.
The position of FeEnt, shown as an orange stick model, is fixed for reference.

Movie S2. Fit of FeEnt and the BtXusB residues and interacting water molecules to the $2mF_o\text{-}DF_c$
electron density map contoured at 1.5σ. FeEnt is in orange, BtXusB residues are in grey, and water
molecules are shown as red spheres.

Supplementary References

1. E. Krissinel, K. Henrick, Inference of macromolecular assemblies from crystalline state. *J. Mol. Biol.* **372**, 774–797 (2007).

2. A. Punjani, J. L. Rubinstein, D. J. Fleet, M. A. Brubaker, cryoSPARC: algorithms for rapid unsupervised cryo-EM structure determination. *Nat. Methods* **14**, 290–296 (2017).

3. G. Posch, M. Pabst, L. Neumann, M. J. Coyne, F. Altmann, P. Messner, L. E. Comstock, C. Schäffer, "Cross-glycosylation" of proteins in Bacteroidales species. *Glycobiology* **23**, 568 (2013).

4. X. Robert, P. Gouet, Deciphering key features in protein structures with the new ENDscript server. *Nucleic Acids Res.* **42** (2014).

5. J. Jumper, R. Evans, A. Pritzel, T. Green, M. Figurnov, O. Ronneberger, K. Tunyasuvunakool, R. Bates, A. Žídek, A. Potapenko, A. Bridgland, C. Meyer, S. A. A. Kohl, A. J. Ballard, A. Cowie, B. Romera-Paredes, S. Nikolov, R. Jain, J. Adler, T. Back, S. Petersen, D. Reiman, E. Clancy, M. Zielinski, M. Steinegger, M. Pacholska, T. Berghammer, S. Bodenstein, D. Silver, O. Vinyals, A. W. Senior, K. Kavukcuoglu, P. Kohli, D. Hassabis, Highly accurate protein structure prediction with AlphaFold. *Nature* **596**, 583–589 (2021).

6. N. Oberg, R. Zallot, J. A. Gerlt, EFI-EST, EFI-GNT, and EFI-CGFP: Enzyme Function Initiative (EFI) Web Resource for Genomic Enzymology Tools. *J. Mol. Biol.* **435**, 168018 (2023).

7. F. W. Studier, B. A. Moffatt, Use of bacteriophage T7 RNA polymerase to direct selective high-level expression of cloned genes. *J. Mol. Biol.* **189**, 113–130 (1986).

8. V. de Lorenzo, L. Eltis, B. Kessler, K. N. Timmis, Analysis of Pseudomonas gene products using *lacIq/Ptrp-lac* plasmids and transposons that confer conditional phenotypes. *Gene* **123**, 17–24 (1993).

9. N. M. Koropatkin, E. C. Martens, J. I. Gordon, T. J. Smith, Starch Catabolism by a Prominent Human Gut Symbiont Is Directed by the Recognition of Amylose Helices. *Structure* **16**, 1105–1115 (2008).