1   **Human adherent cortical organoids in a multiwell format**

3   Mark van der Kroeg[1], Sakshi Bansal[1], Maurits Unkel[1], Hilde Smeenk[1,] Steven A. Kushner[1,2,3#*],

4   Femke M.S. de Vrij[1,4#*]

6   [1]Department of Psychiatry, Erasmus MC, Rotterdam, 3015 CN, The Netherlands.

7   [2]Dept. of Psychiatry, Columbia University Irving Medical Center, New York, NY, USA
8
9   [3]Stavros Niarchos Foundation (SNF) Center for Precision Psychiatry & Mental Health,
10   Columbia University, New York, NY, USA
11
12   [4]ENCORE Expertise Center for Neurodevelopmental Disorders, Erasmus MC, Rotterdam, NL
13
14
15
16
[#]These authors contributed equally

Contact information: *Correspondence: sk2602@cumc.columbia.edu, f.devrij@erasmusmc.nl.

Summary

In the growing diversity of human iPSC-derived models of brain development, we present

here a novel method that exhibits 3D cortical layer formation in a highly reproducible

topography of minimal dimensions. The resulting adherent cortical organoids develop by

self-organization after seeding frontal cortex patterned iPSC-derived neural progenitor cells

in 384-well plates during eight weeks of differentiation. The organoids have stereotypical

dimensions of 3 x 3 x 0.2 mm, contain multiple neuronal subtypes, astrocytes and

oligodendrocyte lineage cells, and are amenable to extended culture for at least 10 months.

Longitudinal imaging revealed morphologically mature dendritic spines, axonal myelination,

and robust neuronal activity. Moreover, adherent cortical organoids compare favorably to

existing brain organoid models on the basis of robust reproducibility in obtaining

topographically-standardized singular radial cortical structures and circumvent the internal

necrosis that is common in free-floating cortical organoids. The adherent human cortical

organoid platform holds considerable potential for high-throughput drug discovery

applications, neurotoxicological screening, and mechanistic pathophysiological studies of

brain disorders.

 Introduction

 Using human embryonic or induced pluripotent stem cell (hiPSC) derived models to

 investigate the developing brain in health and disease has yielded considerable success

 (Marton and Paşca, 2020). The approaches have been varied, including single cell hiPSC-

 derived models grown in a monolayer (Sarkar et al., 2018; Zhang et al., 2013), multiple

 neural cell types in 2D neural networks (Astick and Vanderhaeghen, 2018; Bardy et al., 2015;

 Gunhanlar et al., 2018; Shi et al., 2012), 3D free-floating regionalized neural organoids

 (Paşca et al., 2015; Qian et al., 2016a; Xiang et al., 2019; Zhang et al., 2023), and hiPSC-

 derived free-floating unguided neural organoids (Gomes et al., 2020; Lancaster et al., 2013;

 Paşca et al., 2022; Pellegrini et al., 2020; Renner et al., 2017; Sawada et al., 2020). Among

 the consistent findings across models is that increasing cellular and topographical

 complexity has appeared to come at the cost of increased variability (Eichmüller and

 Knoblich, 2022; Kelava and Lancaster, 2016). Therefore, a major current technical challenge

 is to identify hiPSC-derived models that recapitulate higher-order neural complexity with

 reduced heterogeneity.

 Existing 3D models suffer from considerable variability due to the complex and

 heterogeneous nature of the free-floating structures (Cederquist et al., 2019; Lancaster et

 al., 2013, 2017; Paşca et al., 2015; Renner et al., 2017; Velasco et al., 2019; Yoon et al.,

 2019). A further challenge, in particular with free-floating organoids, is the necrotic core

 that emerges when tissue volumes exceed the limits of oxygen and nutrient diffusion

 beyond a radius of ~300-400μm (Lancaster et al., 2017; Qian et al., 2016b). Although recent

 progress has been made with slicing organoids prior to the emergence of necrosis followed

 by organotypic air-liquid interface culture (Giandomenico et al., 2019; Qian et al., 2020),

 even sliced organoids have to be repeatedly re-cut to prevent necrosis (Qian et al., 2020),

which is both laborious and risks introducing another potential source of variability. The

generation of vascularized organoids would be the ultimate solution to reduce the necrotic

core and the inherent stress that is observed in cortical organoids (Bhaduri et al., 2020; Fan

et al., 2022). It will be beneficial to have increased diffusion through microfluidics or

vascularization (current efforts summarized in Matsui et al., 2021) and further study the

mutually beneficial interaction of neural cells and vascular cells (Crouch et al., 2022;

Mansour et al., 2018; Wang et al., 2023).

Here we propose a simplified approach to generating long-term hiPSC-derived adherent

cortical organoids with reproducible dimensions and the potential for high-throughput

screening in a 384-well format. The resulting adherent cortical organoids can be maintained

in long-term culture and contain neurons with dendritic spines and robust activity, as well as

several classes of glial cells including oligodendrocyte precursor cells, myelinating

oligodendrocytes, and morphologically distinct sub-types of astrocytes.

78 Results

79 *Self-organized topography of iPSC-derived adherent cortical organoids*

Adherent cortical organoids reproducibly self-organized into layered radial structures in

384-wells plates within 8 weeks of seeding with hiPSC-derived forebrain-patterned neural

progenitor cells (NPCs). Three different hiPSC source cell lines were used in this study,

including commercially available NPCs, and NPCs generated using a modified version of our

previously described protocol (Gunhanlar et al., 2018) (**Figure 1A**). NPCs were capable of

neural rosette formation and expressed SOX2, Nestin, and the frontal cortical NPC-marker

FOXG1 (**Figure 1B, C, D, Figure S1A).** The initial 4 weeks after seeding the NPCs in Neural

Differentiation Medium (ND) were characterized by proliferative expansion of NPCs and the

emergence of early neural differentiation markers (**Figure 1E, Figure S2**). Between 4- and 8-

weeks post-seeding, neurons and glial cells emerged with a consistent spatial organization

(**Figure 1E, F, Figure S1B, Figure S2**), in which the central region was densely packed with

cell bodies while the periphery contained circumferentially and radially organized processes

originating from cells in the centre. Typically, cortical organoids defined as a single radial

structure per well were observed in ~80% of the wells seeded with NPCs after 60 days of

differentiation. The structural integrity of single structure organoids remained quite stable

and slowly diminished over time to about 50% of intact single structure organoids after 1

year in culture (**Figure S1D**). Organoid structure formation was highly dependent on the

proliferation rate of NPCs, that can differ substantially between differentiation batches and

hiPSC clones. Plotting the interaction between proliferation and the amount of NPCs

required to be seeded for the successful generation of adherent cortical organoids, showed

a significant correlation (r^2=0.67) that can be used as a guideline for testing a range of NPC

densities (**Figure S1C**). For each NPC line an optimal seeding density was estimated based on

the proliferation rate of that NPC line. Multiple densities were seeded around the estimated

optimal density and after 6 weeks it was visually determined which NPC density enabled

adherent cortical organoid generation. Typically, too sparse seeding density generated

neural networks lacking structure, while excessive densities resulted in overgrowth, leading

to reduced structural organisation and compromised long-term survival.

Cell type distribution and layer formation

The spatial organisation that evolved over the first 8 weeks after seeding was paralleled

by a shift in cell type distribution. Tau+/MAP2- axons exhibited long extensions in a circular

pattern, while MAP2+ dendrites exhibited orthogonally-oriented radial outgrowth (**Figure**

2A, Figure S3).

Overall, a reduction in progenitor markers (SOX2 day 14: 58.4%, day 56: 18.8% *P* =

<0.001, PAX6 day 14: 34.5%, day 56: 8.0% *P* = 0.20) and a significant increase in neuronal

cortical layer markers (CTIP2 day 14: 0.5%, day 56: 14.0% *P* = <0.001, CUX1 day 14: 1.3%,

day 56: 24.4% *P* = <0.001) were observed (**Figure S2**). Cortical layer markers exhibited an

inside-out pattern of development in which expression of the deep layer excitatory

neuronal marker CTIP2 emerged before the upper layer marker CUX1 (**Figure S2**, **Figure 2C**).

After 6-8 weeks following seeding, a self-organised rudimentary segregation of deep- and

upper layer neurons emerged as shown by a clear macroscopic separation of deep- and

upper layer neurons, although some neurons were spatially intermixed and some neurons

were double-positive for CTIP2 and CUX1. Segregation was also observed between CUX1

and CUX2 positive cells as CUX2 is typically expressed over a wider range of upper cortical

layers than CUX1 and also marks intermediate progenitors (Molyneaux et al., 2007) (**Figure**

2C, Figure S4). Analogous to the broad distribution of cortical cell subclasses, the majority of

the neurons were glutamatergic, while GAD67+ interneurons were also present (**Figure 2D**),

constituting ~10% of the NeuN-positive neuronal population consistently for all three

source hiPSC lines (**Figure S5**).

Adherent cortical organoids contain multiple glial cell types

Within 8 weeks of seeding, a population of GFAP+/S100β+ astrocytes emerged. Many

astrocytes had their soma located in the central region with process outgrowth radially

(**Figure 2E, Figure S6**), while other astrocytes exhibited subtype-specific morphologies

including fibrous astrocytes (**Figure 2F**), protoplasmic-like astrocytes (**Figure 2G**) and

interlaminar astrocytes (**Figure 2H**). GFAP/PAX6 double-positive radial glia were present at

the outskirts of the densely populated centre of the organoid with processes growing

radially outwards (**Figure 2I**). Similar to free-floating organoids, adherent cortical organoids

survived for longer periods compared to monolayer neural cultures grown on larger

surfaces. The longevity allows for the development of cell types not usually seen in a

monolayer culture that can typically be cultured up to a maximum of 2 to 3 months. By 6

weeks after seeding NPCs we observed the emergence of oligodendrocyte precursor cells

(OPC), as shown by the expression of NG2 (**Figure 3A**). NG2+ cells remained present until at

least 4 months **(Figure 3B/C)**. Staining for Myelin Basic Protein (MBP) revealed the

emergence of MBP+ oligodendrocytes around 4 months after NPC seeding when the

organoids were continuously grown in the presence of T3 (2ng/ml) **(Figure 3D)**. At 5 months,

the oligodendrocytes showed increasingly mature morphologies (**Figure 3E-I**) and exhibited

MBP co-localization along NF200+ axons (**Figure 3F/I**).

Adherent cortical organoids show synaptic connectivity and functional activity

Neurons within adherent cortical organoids exhibited clear evidence of synaptogenesis

(**Figure 4A-D**). Sparse labelling of excitatory neurons with AAV9.CamKII.eGFP revealed the

presence of Synapsin-positive (Syn+) mushroom-shaped dendritic spines (**Figure 4E**). To

assess the functional activity of the cortical organoids, we used the genetically-encoded

calcium indicator GCaMP6s under the control of the human Synapsin promoter (**Figure 4F**),

allowing cell-type specific quantification of neuronal activity. Calcium imaging revealed

robust synchronous network-level bursting (NB) (1.4 $\pm$ 0.07 NB/min) in which the vast

majority of recorded neurons participated. In addition, substantial desynchronized activity

was also observed (3.9 $\pm$ 0.5 events/min) during time periods outside of network-level

bursting (**Figure 4G-J, Figure 4 – Video 1**).

Discussion

The study of early human brain development and related diseases has long been hampered

by the inherent complexity of the human brain and the inaccessibility of living brain tissue at

cellular resolution. Technological advances in induced pluripotent stem cell technology have

now facilitated the opportunity to obtain living human neurons derived from specific

individuals.

We describe here a platform to model early human frontal cortical development with high

reproducibility and simplified organization. While 3D floating organoids, or sliced organoids,

nicely recapitulate layered cortex formation, they are subject to variation in the relative

contribution of cortical tissue within the organoid, forming multiple cortical patches along

the edges and complicating structured analysis (Eichmüller and Knoblich, 2022;

Giandomenico et al., 2019; Quadrato et al., 2017). Moreover, 3D floating organoids suffer

from necrosis in the core of the organoid due to lack of oxygen and nutrient diffusion.

Recently, other protocols have been published starting from rosette formation with a focus on very early development and leading to single structure free floating cortical organoids(Pagliaro et al., 2023; Tidball et al., 2023). Our platform predefines a rosette-forming iPSC-derived cortical NPC population that self-organizes into adherent singular radial structures in a standard 384-well format. Other examples have highlighted the benefits of a multi-well format or systematic individual structure formation (Knight et al., 2018; Medda et al., 2016). Our platform now integrates these features to yield individual adherent layered cortical structures with robust functional synaptic connectivity and neuronal activity and including advanced glial cell types such as myelinating oligodendrocytes and subclasses of astrocytes. The small reproducible format of the organoids in a 384-well format has the distinct advantage of being able to image entire organoids without slicing or clearing and to perform spatiotemporal functional analysis by fluorescence-based calcium imaging. We confirmed the self-organizing potential and reproducibility of adherent cortical organoids across multiple hiPSC lines and using different sources of NPCs, controlling the seeding density for the proliferation rate of the specific NPC batch. Seeding NPCs with frontal cortical identity in the defined geometry of a 384-well plate enabled the development of long-term functional neural networks in a complex radial structure resembling early human cortical development. Future studies aimed at single cell gene expression analysis and advanced image-based analysis solutions in this platform will be interesting in the context of the increasing knowledge on single cell topographical, typological and temporal hierarchies in the developing human cortex (Bhaduri et al., 2021; Nowakowski et al., 2017, 2016; Uzquiano et al., 2022).

These functional adherent cortical organoids in a multi-well format should be amenable to

high-throughput screening applications, mechanistic pathophysiological studies of

neurodevelopmental and neuropsychiatric disorders, and pharmacological and phenotypic

screening of disease phenotypes during early cortical development. Moreover, toxicological

studies for novel therapeutic compounds also show an increasing need for testing specific

effects in human neuronal models, whereas studies using rodent models have often shown

poor predictive power for drug safety and efficacy in human central nervous system

disorders (van Esbroeck et al., 2017).

Adherent cortical organoids also have some limitations, as the current iteration of adherent

cortical organoids exhibits limited cortical layering and regional specification, that seem to

be more advanced in floating whole brain organoids. However, these drawbacks are offset

by significantly enhanced ease and higher-throughput possibilities of downstream analysis

applications.

Taken together, we present a novel platform for cellular-level human brain modeling using

adherent cortical organoids that exhibit high reproducibility and robust neuronal activity.

The ability to reliably generate human cortical organoids in multi-well plates combined with

neural network functionality offers a unique potential for brain disease modeling and

therapeutic screening applications.

Materials and Methods

Generation of Neural Progenitor Cells

NPCs from 3 different source hiPSC lines were used. NPC-line 1: in house generated NPCs from human iPSC line WTC11 (Provided by Bruce R. Conklin, The Gladstone Institutes and UCSF, #GM25256, RRID:CVCL_Y803, Miyaoka et al.,2014). NPC-line 2: commercially available hNPCs from Axol Biosciences (ax0015). NPC-line 3: NPCs derived using the protocol of Shi et al., (Shi et al., 2012) from hiPSC line IPSC0028 (Sigma-Aldrich, RRID:CVCL_EE38). Line 1 NPCs were generated as previously described from hiPSCs grown on a mouse embryonic fibroblast (MEF) feeder layer (Gunhanlar et al., 2018). After passage 3, NPC cultures were purified using fluorescence-activated cell sorting (FACS). NPCs were detached from the culture plate using Accutase (Stem Cell Technologies) and $CD184^+/CD44^-/CD271^-/CD24^+$ cells (Yuan et al., 2011) were collected on a FACSAria III Cell Sorter (BD Bioscience) and expanded in NPC medium consisting of: DMEM/F12, 1% N2 supplement, 2% B27-RA supplement (Thermo Fisher Scientific), 1 µg/ml laminin (L2020, Sigma-Aldrich), 20 ng/ml basic fibroblast growth factor (Merck-Millipore, Darmstadt, Germany) and 1% penicillin/streptomycin (Thermo Fisher Scientific). NPCs were differentiated to adherent cortical organoids between passage 3 and 7 after sorting.

Neural differentiation

384-well plates (M1937-32EA. Life Technologies) were coated with 50 µg/ml laminin in dH_2O (Sigma, L2020) for 30 minutes at 37 °C. The NPCs in NPC medium were dissociated with Accutase (Stem Cell Technologies), live cells were counted with Trypan Blue (Stem Cell Technologies) in a Burker counting chamber. The NPCs were seeded in the wells of a 384-well plate at defined densities for each cell line. Specifically, the optimal seeding density was

determined by visual inspection of the organoids between 28 to 42 days after seeding a

range of cell densities in the 384-well plate wells. NPCs were seeded and differentiated in

Neural Differentiation Medium: Neurobasal medium (Thermo Fisher Scientific), 1% N2

supplement (Thermo Fisher Scientific), 2% B27-RA supplement (Thermo Fisher Scientific),

1% minimum essential medium/non-essential amino acid (Stem Cell Technologies),

20 ng/ml brain-derived neurotrophic factor (ProSpec Bio), 20 ng/ml glial cell-derived

neurotrophic factor (ProSpec Bio), 1 µM dibutyryl cyclic adenosine monophosphate (Sigma-

Aldrich), 200 µM ascorbic acid (Sigma-Aldrich), 2 µg/ml laminin (Sigma-Aldrich) and 1%

penicillin/streptomycin (Thermo Fisher Scientific). For oligodendrocyte maturation the cells

were grown in the presence of 2 ng/ml T3 (Sigma-Aldrich). Cells were refreshed every 2-3

days.

Immunocytochemistry

For live-dead staining, living cultures were incubated with LIVE/DEAD™ Viability/Cytotoxicity

Kit according to manufacturer's instructions (Thermo Fisher Scientific). For

immunocytochemistry adherent cortical organoids were fixed for 20-30 minutes using 4%

formaldehyde in phosphate-buffered saline (PBS), washed with PBS and blocked for 1 hour

by pre-incubation in staining buffer containing 0.05 M Tris, 0.9% NaCl, 0.25% gelatin and

0.5% Triton-X-100 (pH 7.4). Primary antibodies were incubated for 48-72h at 4 °C in staining

buffer, washed with PBS and incubated with the secondary antibodies in staining buffer for

2h at room temperature. The cultures were embedded in Mowiol 4-88 (Sigma-Aldrich), after

which confocal imaging was performed with a Zeiss LSM700 and Zeiss LSM800 confocal

microscope using ZEN software (Zeiss, Oberkochen, Germany). The following primary

antibodies were used: SOX2 (Merck-Millipore AB5603, 1:200); Nestin (Merck-Millipore

MAB5326, 1:200); MAP2 (Synaptic Systems 188004, 1:100); NeuN (Merck ABN78, 1:200);

GFAP (Merck-Millipore AB5804, 1:300); FOXG1 (Abcam AB18259, 1:200); CUX1 (Abcam

AB54583, 1:200); CTIP2 (Abcam AB18465, 1:100); Synapsin 1/2 (Synaptic Systems 106003,

1:200); and PSD95 (Thermo Fisher Scientific MA1-046, 1:100); Tau (Cell Signaling Technology

4019, 1:200); S100ß (Sigma-Aldrich S2532, 1:200); Pax6 (Santa Cruz sc-81649, 1:100); NG2

(Gift from W. Stallcup Lab, 1:100); NF200 (Sigma-Aldrich 083M4833 1:200); MBP (Abcam

AB7349, 1:100); GFP (Abcam ab13970, 1:100); GAD67 (Merck-Millipore MAB5406, 1:100)

DAPI (Thermo Fisher Scientific D1306). The following secondary antibodies were used 1:200:

Alexa-488, Alexa-555, Alexa-647 (Jackson ImmunoResearch, West Grove, PA, USA).

Sparse labelling of excitatory neurons

pENN.AAV9.CamKII.4.eGFP.WPRE.rBG (Addgene viral prep # 105541-AAV9) was added to

the cortical organoids at day 278 (1.68×10^8 GC/well). The transduced cortical organoids

were fixed and stained at day 310.

Calcium imaging

For calcium imaging the genetically encoded calcium indicator

AAV1.Syn.GCaMP6s.WPRE.SV40 (Penn Vector Core, 100843-AAV1) was added to the

organoids at day 42 of differentiation (1.5×10^8 GC/well). Recordings were performed at day

60 on a Zeiss LSM800 confocal microscope using ZEN software (Zeiss, Oberkochen,

Germany). The recordings were made with a 20x/0.8NA Ph2 Plan-Apochromat objective,

with a field of view of 150 × 100 μm and a pixel size of 0,3 μm. The acquisition rates of the

recordings were between 4-5 f.p.s. 24 hours before the recordings, the medium was

switched to BrainPhys Neural Differentiation Medium. BrainPhys Neuronal Media (Stem Cell

Technologies), 1% N2 supplement (Thermo Fisher Scientific), 2% B27-RA supplement

(Thermo Fisher Scientific), 1% minimum essential medium/non-essential amino acid (Stem

Cell Technologies), 1% penicillin/streptomycin (Thermo Fisher Scientific), 20 ng/ml brain-

derived neurotrophic factor (ProSpec Bio), 20 ng/ml glial cell-derived neurotrophic factor

(ProSpec Bio), 1 µM dibutyryl cyclic adenosine monophosphate (Sigma-Aldrich),

200 µM ascorbic acid (Sigma-Aldrich) and 2 µg/ml laminin (Sigma-Aldrich). The calcium

imaging recordings were processed using CNMF-E (Pnevmatikakis et al., 2016). Calcium

traces were then analysed using a custom script for event and network burst detection

using an algorithm written in Python (v3.8.2) (code accessibility can be requested via

Github).

Cyquant Proliferation Assay

CyQUANT™ Direct Cell Proliferation Assay, C35011 (Thermo Fisher Scientific) was used

according to manufacturer specifications. For each time point and each of the 3 NPC lines

10-12 wells of a 96-well plate were seeded with NPCs (2500 NPCs per well). In addition, NPC

lines generated from 2 different clones from IPS line MH0159020 (Rutgers University Cell

and DNA Repository) were added to increase the dynamic range of the proliferation curve.

At 24h and 96h NPCs were frozen at -80℃. All NPC lines and timepoints were thawed, lysed

and measured together. Doubling time was calculated between 24h and 96h. None of the

wells was confluent at 96h.

Statistical analysis

All data represent mean ± SEM. When comparing developmental markers in Figure S2 we

used one-way ANOVA followed by Tukey- Kramer's multiple correction test. n=3-6 images

taken over two wells, for each time point.

Acknowledgements

This work was supported by the Netherlands Organ-on-Chip Initiative, an NWO Gravitation

project funded by the Ministry of Education, Culture and Science of the government of the

Netherlands (024.003.001) to SAK, Hersenstichting Fellowship (F2012(1)-39) to FMSdV, an

Erasmus MC Human Disease Model Award to FMSdV and a Dutch ZonMw MKMD

Create2Solve grant (114025201) to SAK and FMSdV.

Author contributions

MvdK, SB and HS performed experiments, MU developed calcium imaging analysis

algorithms, FdV and SAK supervised the work, MvdK, FdV and SAK wrote the paper.

Declaration of interests

The authors declare no competing interests.

References

Astick M, Vanderhaeghen P. 2018. From Human Pluripotent Stem Cells to Cortical Circuits.

Curr Top Dev Biol **129**:67–98. doi:10.1016/bs.ctdb.2018.02.011

Bardy C, van den Hurk M, Eames T, Marchand C, Hernandez R V, Kellogg M, Gorris M, Galet

B, Palomares V, Brown J, Bang AG, Mertens J, Böhnke L, Boyer L, Simon S, Gage FH.

2015. Neuronal medium that supports basic synaptic functions and activity of human

neurons in vitro. *Proc Natl Acad Sci U S A* **112**. doi:10.1073/pnas.1504393112

Bhaduri A, Andrews MG, Mancia Leon W, Jung DD, Shin D, Allen D, Jung DD, Schmunk G,

Haeussler M, Salma J, Pollen AA, Nowakowski TJ, Kriegstein AR. 2020. Cell stress in

cortical organoids impairs molecular subtype specification. *Nature* 1–7.

doi:10.1038/s41586-020-1962-0

Bhaduri A, Sandoval-Espinosa C, Otero-Garcia M, Oh I, Yin R, Eze UC, Nowakowski TJ,

Kriegstein AR. 2021. An atlas of cortical arealization identifies dynamic molecular

signatures. *Nature* **598**:200–204. doi:10.1038/s41586-021-03910-8

Cederquist GY, Asciolla JJ, Tchieu J, Walsh RM, Cornacchia D, Resh MD, Studer L. 2019.

Specification of positional identity in forebrain organoids. *Nat Biotechnol* **37**:436–444.

doi:10.1038/s41587-019-0085-3

Crouch EE, Bhaduri A, Andrews MG, Cebrian-Silla A, Diafos LN, Birrueta JO, Wedderburn-

Pugh K, Valenzuela EJ, Bennett NK, Eze UC, Sandoval-Espinosa C, Chen J, Mora C, Ross

JM, Howard CE, Gonzalez-Granero S, Lozano JF, Vento M, Haeussler M, Paredes MF,

Nakamura K, Garcia-Verdugo JM, Alvarez-Buylla A, Kriegstein AR, Huang EJ. 2022.

Ensembles of endothelial and mural cells promote angiogenesis in prenatal human

brain. *Cell* **185**:3753-3769.e18. doi:10.1016/j.cell.2022.09.004

Eichmüller OL, Knoblich JA. 2022. Human cerebral organoids — a new tool for clinical

neurology research. *Nat Rev Neurol* **18**:661–680. doi:10.1038/s41582-022-00723-9

Fan W, Christian KM, Song H, Ming G li. 2022. Applications of Brain Organoids for Infectious

Diseases. *J Mol Biol* **434**:167243. doi:10.1016/j.jmb.2021.167243

Giandomenico SL, Mierau SB, Gibbons GM, Wenger LMD, Masullo L, Sit T, Sutcliffe M,

Boulanger J, Tripodi M, Derivery E, Paulsen O, Lakatos A, Lancaster MA. 2019. Cerebral

organoids at the air–liquid interface generate diverse nerve tracts with functional

output. *Nat Neurosci* **22**:669–679. doi:10.1038/s41593-019-0350-2

Gomes AR, Fernandes TG, Vaz SH, Silva TP, Bekman EP, Xapelli S, Duarte S, Ghazvini M,

Gribnau J, Muotri AR, Trujillo CA, Sebastião AM, Cabral JMS, Diogo MM. 2020.

Modeling Rett Syndrome With Human Patient-Specific Forebrain Organoids. *Front Cell*

Dev Biol **8**:1–18. doi:10.3389/fcell.2020.610427

Gunhanlar N, Shpak G, van der Kroeg M, Gouty-Colomer LA, Munshi ST, Lendemeijer B,

Ghazvini M, Dupont C, Hoogendijk WJG, Gribnau J, de Vrij FMS, Kushner SA. 2018. A

simplified protocol for differentiation of electrophysiologically mature neuronal

networks from human induced pluripotent stem cells. *Mol Psychiatry* **23**:1336–1344.

doi:10.1038/mp.2017.56

Kelava I, Lancaster MAA. 2016. Stem Cell Models of Human Brain Development. *Cell Stem*

Cell **18**:736–748. doi:10.1016/j.stem.2016.05.022

Knight GT, Lundin BF, Iyer N, Ashton LMT, Sethares WA, Willett RM, Ashton RS. 2018.

Engineering induction of singular neural rosette emergence within hPSC-derived

tissues. *Elife* **7**. doi:10.7554/eLife.37549

Lancaster M a, Renner M, Martin C-A, Wenzel D, Bicknell LS, Hurles ME, Homfray T,

Penninger JM, Jackson AP, Knoblich JA. 2013. Cerebral organoids model human brain

development and microcephaly. *Nature* **501**:373–379. doi:10.1038/nature12517

Lancaster MA, Corsini NS, Wolfinger S, Gustafson EH, Phillips AW, Burkard TR, Otani T,

Livesey FJ, Knoblich JA. 2017. Guided self-organization and cortical plate formation in

human brain organoids. *Nat Biotechnol* **35**:659–666. doi:10.1038/nbt.3906

Mansour AA, Gonçalves JT, Bloyd CW, Li H, Fernandes S, Quang D, Johnston S, Parylak SL, Jin

X, Gage FH. 2018. An in vivo model of functional and vascularized human brain

organoids. *Nat Biotechnol* **36**:432–441. doi:10.1038/nbt.4127

Marton RM, Pașca SP. 2020. Organoid and Assembloid Technologies for Investigating

Cellular Crosstalk in Human Brain Development and Disease. *Trends Cell Biol*.

doi:10.1016/j.tcb.2019.11.004

Matsui TK, Tsuru Y, Hasegawa K, Kuwako K ichiro. 2021. Vascularization of human brain

organoids. *Stem Cells* **39**:1017–1024. doi:10.1002/stem.3368

Medda X, Mertens L, Versweyveld S, Diels A, Barnham L, Bretteville A, Buist A, Verheyen A,

Royaux I, Ebneth A, Cabrera-Socorro A. 2016. Development of a Scalable, High-

Throughput-Compatible Assay to Detect Tau Aggregates Using iPSC-Derived Cortical

Neurons Maintained in a Three-Dimensional Culture Format. *J Biomol Screen*.

doi:10.1177/1087057116638029

Molyneaux BJ, Arlotta P, Menezes JRL, Macklis JD. 2007. Neuronal subtype specification in

the cerebral cortex. *Nat Rev Neurosci* **8**:427–37. doi:10.1038/nrn2151

Nowakowski TJ, Bhaduri A, Pollen AA, Alvarado B, Mostajo-Radji MA, Di Lullo E, Haeussler

M, Sandoval-Espinosa C, Liu SJ, Velmeshev D, Ounadjela JR, Shuga J, Wang X, Lim DA,

West JA, Leyrat AA, Kent WJ, Kriegstein AR. 2017. Spatiotemporal gene expression

trajectories reveal developmental hierarchies of the human cortex. *Science (80-)* **358**.

doi:10.1126/science.aap8809

Nowakowski TJ, Pollen AA, Sandoval-Espinosa C, Kriegstein AR. 2016. Transformation of the

Radial Glia Scaffold Demarcates Two Stages of Human Cerebral Cortex Development.

Neuron **91**. doi:10.1016/j.neuron.2016.09.005

Pagliaro A, Finger R, Zoutendijk I, Bunschuh S, Clevers H, Hendriks D, Artegiani B. 2023.

Temporal morphogen gradient-driven neural induction shapes single expanded

neuroepithelium brain organoids with enhanced cortical identity. *Nat Commun* **14**.

doi:10.1038/s41467-023-43141-1

Paşca AM, Sloan SA, Clarke LE, Tian Y, Makinson CD, Huber N, Kim CH, Park J-Y, O'Rourke

NA, Nguyen KD, Smith SJ, Huguenard JR, Geschwind DH, Barres BA, Paşca SP. 2015.

Functional cortical neurons and astrocytes from human pluripotent stem cells in 3D

culture. *Nat Methods*. doi:10.1038/nmeth.3415

Paşca SP, Arlotta P, Bateup HS, Camp JG, Cappello S, Gage FH, Knoblich JA, Kriegstein AR,

Lancaster MA, Ming G-L, Muotri AR, Park I-H, Reiner O, Song H, Studer L, Temple S,

Testa G, Treutlein B, Vaccarino FM. 2022. A nomenclature consensus for nervous

system organoids and assembloids. *Nat 2022 6097929* **609**:907–910.

doi:10.1038/s41586-022-05219-6

Pellegrini L, Bonfio C, Chadwick J, Begum F, Skehel M, Lancaster MA. 2020. Human CNS

barrier-forming organoids with cerebrospinal fluid production. *Science (80-)* **369**.

doi:10.1126/science.aaz5626

Qian X, Nguyen HN, Song MM, Hadiono C, Ogden SC, Hammack C, Yao B, Hamersky GR,

Jacob F, Zhong C, Yoon KJ, Jeang W, Lin L, Li Y, Thakor J, Berg DA, Zhang C, Kang E,

Chickering M, Nauen D, Ho CY, Wen Z, Christian KM, Shi PY, Maher BJ, Wu H, Jin P,

Tang H, Song H, Ming GL. 2016b. Brain-Region-Specific Organoids Using Mini-

bioreactors for Modeling ZIKV Exposure. *Cell* **165**:1238–1254.

doi:10.1016/j.cell.2016.04.032

Qian X, Su Y, Adam CD, Deutschmann AU, Pather SR, Goldberg EM, Su K, Li S, Lu L, Jacob F,

Nguyen PTT, Huh S, Hoke A, Swinford-Jackson SE, Wen Z, Gu X, Pierce RC, Wu H, Briand

LA, Chen HI, Wolf JA, Song H, Ming G. 2020. Sliced Human Cortical Organoids for

Modeling Distinct Cortical Layer Formation. *Cell Stem Cell* **26**:766-781.e9.

doi:10.1016/j.stem.2020.02.002

Qian Xuyu, Nguyen Ha Nam, Song MM, Hadiono C, Ogden Sarah C., Hammack Christy, Yao

Bing, Hamersky GR, Jacob F, Zhong C, Yoon K, Jeang W, Lin L, Li Yujing, Thakor J, Berg

DA, Zhang Ce, Kang E, Chickering M, Nauen David, Ho C-Y, Wen Zhexing, Christian KM,

Shi P-Y, Maher Brady J., Wu H, Jin P, Tang Hengli, Song H, Ming G, Bayer A, Lennemann

NJ, Ouyang Y, Bramley JC, Morosky S, Marques ETJ, Cherry S, Sadovsky Y, Coyne CB,

Ben-Ari Y, Spitzer NC, Blackshaw S, Scholpp S, Placzek M, Ingraham H, Simerly R,

Shimogori T, Calvet G, Aguiar RS, Melo AS, Sampaio SA, Filippis I de, Fabri A, Araujo ES,

Sequeira PC de, Mendonça MC de, Oliveira L de, al. et, Cauchemez S, Besnard M,

Bompard P, Dub T, Guillemette-Artur P, Eyrolle-Guignot D, Salje H, Kerkhove MD Van,

Abadie V, Garel C, al. et, Chung S, Hedlund E, Hwang M, Kim DW, Shin BS, Hwang DY,

Kang UJ, Isacson O, Kim KS, Danjo T, Eiraku M, Muguruma K, Watanabe K, Kawada M,

Yanagawa Y, Rubenstein JL, Sasai Y, Driggers RW, Ho CY, Korhonen EM, Kuivanen S,

Jääskeläinen AJ, Smura T, Rosenberg A, Hill DA, DeBiasi RL, Vezina G, al. et, Faria NR,

Azevedo RD, Kraemer MU, Souza R, Cunha MS, Hill SC, Thézé J, Bonsall MB, Bowden

TA, Rissanen I, al. et, Garcez PP, Loiola EC, Costa RM da, Higa LM, Trindade P,

Delvecchio R, Nascimento JM, Brindeiro R, Tanuri A, Rehen SK, Goodwin TJ, Schroeder

WF, Wolf DA, Moyer MP, Haddow AD, Schuh AJ, Yasuda CY, Kasper MR, Heang V, Huy

R, Guzman H, Tesh RB, Weaver SC, Hansen DV, Lui JH, Parker PR, Kriegstein AR,

Heymann DL, Hodgson A, Sall AA, Freedman DO, Staples JE, Althabe F, Baruah K,

Mahmud G, Kandun N, Vasconcelos PF, al. et, Jaffe AE, Shin J, Collado-Torres L, Leek JT,

Tao R, Li C, Gao Y, Jia Y, Maher B.J., Hyde TM, al. et, Jessup JM, Goodwin TJ, Spaulding

G, Kadoshima T, Sakaguchi H, Nakano T, Soen M, Ando S, Eiraku M, Sasai Y, Kim JY, Liu

CY, Zhang F, Duan X, Wen Z., Song J, Feighery E, Lu B, Rujescu D, Clair DS, al. et, Kriks S,

Shim JW, Piao J, Ganat YM, Wakeman DR, Xie Z, Carrillo-Reid L, Auyeung G, Antonacci

C, Buch A, al. et, Kundakovic M, Gudsnuk K, Franks B, Madrid J, Miller RL, Perera FP,

Champagne FA, Lancaster MA, Knoblich JA, Lancaster MA, Renner M, Martin CA,

Wenzel D, Bicknell LS, Hurles ME, Homfray T, Penninger JM, Jackson AP, Knoblich JA,

Lazear HM, Govero J, Smith AM, Platt DJ, Fernandez E, Miner JJ, Diamond MS, Lui JH,

Hansen DV, Kriegstein AR, Mariani J, Coppola G, Zhang P, Abyzov A, Provini L, Tomasini

L, Amenduni M, Szekely A, Palejev D, Wilson M, al. et, Mlakar J, Korva M, Tul N,

Popović M, Poljšak-Prijatelj M, Mraz J, Kolenc M, Rus KR, Vipotnik TV, Vodušek VF, al.

et, Muguruma K, Nishiyama A, Kawakami H, Hashimoto K, Sasai Y, Paşca AM, Sloan SA,

Clarke LE, Tian Y, Makinson CD, Huber N, Kim CH, Park JY, O'Rourke NA, Nguyen KD, al.

et, Petanjek Z, Berger B, Esclapez M, Pollen AA, Nowakowski TJ, Chen J, Retallack H,

Sandoval-Espinosa C, Nicholas CR, Shuga J, Liu SJ, Oldham MC, Diaz A, al. et, Roost MS,

Iperen L van, Ariyurek Y, Buermans HP, Arindrarto W, Devalla HD, Passier R, Mummery

CL, Carlotti F, Koning EJ de, al. et, Tang H., Hammack C., Ogden S.C., Wen Z., Qian X., Li

Y., Yao B., Shin J, Zhang F, Lee EM, al. et, Thomsen ER, Mich JK, Yao Z, Hodge RD, Doyle

AM, Jang S, Shehata SI, Nelson AM, Shapovalova NV, Levi BP, al. et, Wang W, Lufkin T,

Wen Z., Nguyen H.N., Guo Z, Lalli MA, Wang X, Su Y, Kim NS, Yoon KJ, Shin J, Zhang C.,

al. et, Yin X, Mead BE, Safaee H, Langer R, Karp JM, Levy O, Yoon KJ, Nguyen H.N.,

Ursini G, Zhang F, Kim NS, Wen Z., Makri G, Nauen D., Shin JH, Park Y, al. et, Yu X,

Zecevic N. 2016a. Brain-Region-Specific Organoids Using Mini-bioreactors for Modeling

ZIKV Exposure. *Cell* **165**:1238–1254. doi:10.1016/j.cell.2016.04.032

Quadrato G, Nguyen T, Macosko EZ, Sherwood JL, Min Yang S, Berger DR, Maria N, Scholvin

J, Goldman M, Kinney JP, Boyden ES, Lichtman JW, Williams ZM, McCarroll SA, Arlotta

P. 2017. Cell diversity and network dynamics in photosensitive human brain organoids.

Nature **545**:48–53. doi:10.1038/nature22047

Renner M, Lancaster MA, Bian S, Choi H, Ku T, Peer A, Chung K, Knoblich JA. 2017. Self-

organized developmental patterning and differentiation in cerebral organoids. *EMBO J*

36:1316–1329. doi:10.15252/embj.201694700

Sarkar A, Mei A, Paquola ACM, Stern S, Bardy C, Klug JR, Kim S, Neshat N, Kim HJ, Ku M,

Shokhirev MN, Adamowicz DH, Marchetto MC, Jappelli R, Erwin JA, Padmanabhan K,

Shtrahman M, Jin X, Gage FH. 2018. Efficient Generation of CA3 Neurons from Human

Pluripotent Stem Cells Enables Modeling of Hippocampal Connectivity In Vitro. *Cell*

Stem Cell **22**:684-697.e9. doi:10.1016/j.stem.2018.04.009

Sawada T, Chater TE, Sasagawa Y, Yoshimura M, Fujimori-Tonou N, Tanaka K, Benjamin KJM,

Paquola ACM, Erwin JA, Goda Y, Nikaido I, Kato T. 2020. Developmental excitation-

inhibition imbalance underlying psychoses revealed by single-cell analyses of

discordant twins-derived cerebral organoids. *Mol Psychiatry* **25**:2695–2711.

doi:10.1038/s41380-020-0844-z

Shi Y, Kirwan P, Smith J, Robinson HPC, Livesey FJ. 2012. Human cerebral cortex

development from pluripotent stem cells to functional excitatory synapses. *Nat*

Neurosci **15**:477–86, S1. doi:10.1038/nn.3041

Tidball AM, Niu W, Ma Q, Takla TN, Walker JC, Margolis JL, Mojica-Perez SP, Sudyk R, Deng

L, Moore SJ, Chopra R, Shakkottai VG, Murphy GG, Yuan Y, Isom LL, Li JZ, Parent JM.

2023. Deriving early single-rosette brain organoids from human pluripotent stem cells.

Stem Cell Reports **18**:2498–2514. doi:10.1016/j.stemcr.2023.10.020

Uzquiano A, Kedaigle AJ, Pigoni M, Paulsen B, Adiconis X, Kim K, Faits T, Nagaraja S, Antón-

Bolaños N, Gerhardinger C, Tucewicz A, Murray E, Jin X, Buenrostro J, Chen F, Velasco

S, Regev A, Levin JZ, Arlotta P. 2022. Proper acquisition of cell class identity in

organoids allows definition of fate specification programs of the human cerebral

cortex. *Cell* **185**:3770-3788.e27. doi:10.1016/j.cell.2022.09.010

van Esbroeck ACM, Janssen APA, Cognetta AB, Ogasawara D, Shpak G, van der Kroeg M,

Kantae V, Baggelaar MP, de Vrij FMS, Deng H, Allarà M, Fezza F, Lin Z, van der Wel T,

Soethoudt M, Mock ED, den Dulk H, Baak IL, Florea BI, Hendriks G, De Petrocellis L,

Overkleeft HS, Hankemeier T, De Zeeuw CI, Di Marzo V, Maccarrone M, Cravatt BF,

Kushner SA, van der Stelt M. 2017. Activity-based protein profiling reveals off-target

proteins of the FAAH inhibitor BIA 10-2474. *Science (80-)* **356**:1084–1087.

doi:10.1126/science.aaf7497

Velasco S, Kedaigle AJ, Simmons SK, Nash A, Rocha M, Quadrato G, Paulsen B, Nguyen L,

Adiconis X, Regev A, Levin JZ, Arlotta P. 2019. Individual brain organoids reproducibly

form cell diversity of the human cerebral cortex. *Nature* 1. doi:10.1038/s41586-019-

1289-x

Wang M, Gage FH, Schafer ST. 2023. Transplantation Strategies to Enhance Maturity and

Cellular Complexity in Brain Organoids. *Biol Psychiatry* **93**:616–621.

doi:10.1016/j.biopsych.2023.01.004

Xiang Y, Tanaka Y, Cakir B, Patterson B, Kim K-Y, Sun P, Kang Y-J, Zhong M, Liu X, Patra P, Lee

S-H, Weissman SM, Park I-H. 2019. hESC-Derived Thalamic Organoids Form Reciprocal

Projections When Fused with Cortical Organoids. *Cell Stem Cell* **24**:487-497.e7.

doi:10.1016/J.STEM.2018.12.015

Yoon S-J, Elahi LS, Paşca AM, Marton RM, Gordon A, Revah O, Miura Y, Walczak EM,

Holdgate GM, Fan HC, Huguenard JR, Geschwind DH, Paşca SP. 2019. Reliability of

human cortical organoid generation. *Nat Methods* **16**:75–78. doi:10.1038/s41592-018-

0255-0

Yuan SH, Martin J, Elia J, Flippin J, Paramban RI, Hefferan MP, Vidal JG, Mu Y, Killian RL,

Israel MA, Emre N, Marsala S, Marsala M, Gage FH, Goldstein LSB, Carson CT. 2011.

Cell-Surface Marker Signatures for the Isolation of Neural Stem Cells, Glia and Neurons

Derived from Human Pluripotent Stem Cells. *PLoS One* **6**:e17540.

doi:10.1371/journal.pone.0017540

Zhang Y, Pak C, Han Y, Ahlenius H, Zhang Z, Chanda S, Marro S, Patzke C, Acuna C, Covy J, Xu

W, Yang N, Danko T, Chen L, Wernig M, Südhof TCCC. 2013. Rapid Single-Step Induction

of Functional Neurons from Human Pluripotent Stem Cells. *Neuron* **78**:785–798.

doi:10.1016/j.neuron.2013.05.029

Zhang Z, Wang X, Park S, Song H, Ming GL. 2023. Development and Application of Brain

Region–Specific Organoids for Investigating Psychiatric Disorders. *Biol Psychiatry*

93:594–605. doi:10.1016/j.biopsych.2022.12.015

 Figure Titles and Legends

Figure 1 Adherent cortical organoid model. A Schematic representation of the

differentiation protocol. **B/C/D** Representative NPCs from three different iPSC lines with

markers SOX2, Nestin and FOXG1 (scale bar B, 50 µm; C/D, 20 µm). **E** Representative time

course showing self-organization during differentiation, starting with radial organization

between day 28 and day 42, seeding density 1500 NPCs per well (green is Live-stain from

Viability/Cytotoxicity kit; scale bars left to right 100 µm, 150 µm, 100 µm, 100 µm). **F** Full

well showing radial organization at day 42 in culture were only a few dead cells are visible in

red in the dense centre of the structure, seeding density 1500 NPCs per well (scale bar, 500

µm).

Figure 2 Adherent cortical organoids show an organized network of neuronal and

astrocyte subtypes. A MAP2+ somas and dendrites alongside Tau+/MAP2- axons show

segregation of dendritic and axonal compartments, with SOX2+ progenitors concentrated in

the center of the organoid (Day 75, 200µm). **B** MAP2+ and NeuN+ cells indicate mature

neurons (Day 72, 50 µm). **C** Deep-layer cortical marker CTIP2 and upper layer marker CUX1

show rudimentary segregation of cortical layers in expected inside-out pattern (Day 64, 50

µm). **D** A subset of the MAP2-positive neurons are GAD67+ (white arrows), indicating the

presence of an interneuron population in the cortical organoids (Day 60, 50 µm). **E** Astrocyte

markers GFAP and S100β show the general radial pattern of astrocyte outgrowth (Day 66,

500 µm). **F/G/H** GFAP staining reveals the morphologies of different astrocyte subtypes

(white arrows), including fibrous astrocytes (**F**, Day 65, 100 µm), protoplasmic astrocytes (**G**,

Day 65, 50 µm) and interlaminar astrocytes (**H**, Day 65, 100 µm). **I** Co-localization of

astrocyte marker GFAP and PAX6 marks radial glia (white arrows) (Day 65, 50 µm).

Figure 3 Adherent cortical organoids form oligodendrocyte lineage cells. A Adherent

cortical organoids show OPCs as early as 44 days, indicated by OPC marker NG2 (Day 44, 20

µm). **B/C** The NG2+ OPCs are still present in the cortical organoids after 4 months (Day 119,

B 50 µm, C 20 µm). **D** Young oligodendrocytes start to emerge after 4 months indicated by

rudimentary MBP staining (Day 119 100 µm and 20 µm). **E/F** After 5 months, MBP-positive

oligodendrocytes show more mature morphology and initial wrapping of NF200+ axons (Day

148; E, 50 µm and 20 µm, F 5 µm). **G/H/I** Oligodendrocyte distribution at day 161 where the

MBP+ oligodendrocytes sit between axons bundles and co-localize with NF200+ axons.

(G/H/I Day 161, G 500 µm, H 100µm, I 10 µm)

Figure 4 Adherent cortical organoids show synaptic connectivity and network bursts. A

Synapsin staining shows synapse formation along MAP2+ dendrites (Day 70, 20 µm). **B** Co-

localization of pre-synaptic marker Synapsin and post-synaptic marker PSD-95 (Day 205, 20

µm). **C** MAP2+ dendrites and soma are decorated with Synapsin+ synapses (Day 251, 100

µm) **D** Overview of entire well, showing alignment of Synapsin staining with MAP2 (Day 251,

500 µm)**. E** Sparse labeling of neurons with AAV9.CamKII.eGFP allows detailed imaging of

glutamatergic dendritic spines with mature mushroom morphology, contacting pre-synaptic

Synapsin puncta (Day 310, 5 µm). **F** Snapshot of radially organized neurons transduced with

AAV1.Syn.GCaMP6s.WPRE.SV40 (Day 60, 100 µm). **G** Calcium events per minute from 36

cells from 7 recordings (each showing a cluster of neurons) from 2 different cortical

organoids at day 61. **H** Network bursts per minute from 37 cells from 6 recordings from 2

different cortical organoids. **I** Percentage of events that are part of network bursts indicates

that the majority of cells are involved in network burst activity. **J** Representative calcium

traces from 2 different clusters in which cells are showing individual activity as well as

network bursts where multiple cells are active simultaneously.

Supplementary Figure 1 Reproducibility of Neural Progenitor Cells and Adherent Cortical

Organoids. A NPCs are positive for SOX2, Nestin, FOXG1, PAX6 and TBR2 (scale bars 20 μm)

B LIVE/DEAD staining of adherent organoids (N=4) showing reproducibility of the structure

formation. Seeding density line 1; 1250 NPCs, line 2; 750 NPCs, line 3; 1000 NPCs (scale bars

500 μm). **C** Function of the necessary NPC seeding density to form adherent cortical

organoids in relation to the doubling time as a measure of the proliferation rate of the NPCs.

The doubling time of the NPCs explains more than half the variation (r^2 = 0.67) of the

required seeding density showing that fewer cells need to be seeded for NPC lines with a

higher proliferation rate. **D** Proportion of successful, single structure adherent cortical

organoids. Each dot represents a different batch of adherent cortical organoids where the

size of the batch ranges from 10-40 organoids.

Supplementary Figure 2 Neuronal maturation in cortical organoids is shown by a temporal

shift in NPC and neuronal markers. **A** Representative images exhibiting change in

expression of NPC and neuronal markers over time. Number of cells positive for NPC

markers SOX2 and PAX6 decreases over time. Deep-layer CTIP2+ neurons start appearing at

day 28, while upper-layer CUX1+ positive neurons appear in higher numbers around day 56.

(Scale bars 20 μm). **B** The percentages of DAPI+ cells expressing each of the markers were

quantified at four time points. Significant differences between week 2 and week 8

expression were seen for SOX2+ (down by 39.6 %), CTIP2+ (up by 13.5 %) and CUX1+ cells

(up by 23.1 %). *** $p < 0.0001$, ANOVA one-way followed by Tukey- Kramer's multiple

correction test. Error bars indicate SEM, n=3-6 images taken over two wells, for each time

point.

Supplementary Figure S3 Distribution of axons and dendrites in adherent cortical

organoids Radially organized MAP2+/Tau+ dendrites along with both radially and

circumferentially organized NF200+/Tau+ axons in duplicate for each cell line (Day 63, scale

bars 500 μm)

Supplementary Figure S4 Cortical layering in the adherent cortical organoids A

Rudimentary separation of SOX2+ NPCs, CTIP2+ deep layer neurons and CUX1+ upper layer

neurons (Day 67, scale bar 100 μm) **B-E** Separation of CUX1 and CUX2 expression (B day 70,

scale bar 100 μm; C day 70, scale bar 500 μm; D day 67, scale bar 500 μm; E day 67, scale

bar 100 μm)

Supplementary Figure S5 GAD67+ interneurons make up a small proportion of NeuN+

neurons in the adherent cortical organoids A Example image of GAD67+ interneuron

proportion of NeuN+ neurons (Day 67, 50 μm). **B** Quantification of GAD67+ proportion of

NeuN+ neurons. Every point is the percentage of GAD67+ interneurons out of all NeuN+

neurons in a spatially randomized selected image with on average 130.6 (± 16.1) NeuN+

nuclei per frame. (Line 1: N= 26 images of 6 different cortical organoids at day 65-67; Line

2/3: 6 images of 2 different organoids at day 65).

Supplementary Figure S6 Astrocyte distribution within Adherent Cortical Organoids The

abundantly present GFAP+ and S100B+ astrocytes have a comparable distribution with the

MAP2+ neurons in the adherent cortical organoids. The astrocytes show radial patterns with

many somas located in the centre as well as a large presence outside the centre (Day 63,

scale bars 500 µm).

Figure 4 – Video 1

Video showing 10 minutes of calcium imaging of an adherent cortical organoid at Day 61

using the genetically encoded calcium indicator AAV1.Syn.GCaMP6s.WPRE.SV40 (rainbow

filter indicates increasing calcium signal ranging from blue to red, speed increase indicated

by time stamp in top left corner).

A

hIPSCs on MEFs → Embryoid bodies → Neural Progenitor mix → FACS

CD184+
CD44-/CD271-
CD24+

Purified NPCs → 384-well plate → Differentiation

B
Nestin
SOX2
DAPI

C
Nestin
SOX2
DAPI

D
FOXG1
Nestin
DAPI

E
Day 14   Day 28   Day 42   Day 56
Live/Dead

F
Day 42
Live/Dead