Genetic basis of reproductive isolation in Torrey pine (*Pinus torreyana* Parry): insights

from hybridization and adaptation

Lionel N. Di Santo[1,2,3*], Alayna Mead[4,*], Jessica W. Wright[5], and Jill A. Hamilton[1,4,6]

[1] North Dakota State University, Department of Biological Sciences, Fargo, ND.

[2] University of Geneva, Department of Genetics and Evolution, Geneva, Switzerland.

[3] University of Basel, Department of Environmental Sciences, Basel, Switzerland.

[4] Pennsylvania State University, Department of Ecosystem Science and Management, State

College, PA.

[5] Pacific Southwest Research Station, USDA-Forest Service, Placerville, CA.

[6] Corresponding author: jvh6349@psu.edu

[*] Contributed equally to the study

Keywords: conifers, forest trees, genetic rescue, reproductive isolation, Torrey Pine.

Abstract

Tree species are often locally adapted to their environments, but the extent to which

environmental adaptation contributes to incipient speciation is unclear. One of the rarest

pines in the world, Torrey Pine (*Pinus torreyana* Parry), persists naturally across one island

and one mainland population in southern California. The two populations are

morphologically and genetically differentiated, but experience some connectivity, making it

an ideal system for assessing the evolution of reproductive isolation. Previous work has found

evidence of heterosis in F1 mainland-island hybrids, suggesting genetic rescue could be

beneficial in the absence of reproductive barriers. Using ddRADseq and GWAS for a

common garden experiment of island, mainland, and F1 individuals, we identified candidate

loci for environmentally-driven reproductive isolation, their function, and their relationship to

fitness proxies. By simulating neutral evolution and admixture between the two populations,

we identified loci that exhibited reduced heterozygosity in the F1s, evidence of selection

against admixture. SNPs with reduced F1 heterozygosity were enriched for growth and

pollination functions, suggesting genetic variants that could be involved in the evolution of

reproductive barriers between populations. One locus with reduced F1 heterozygosity

exhibited strong associations with growth and reproductive fitness proxies in the common

garden, with the mainland allele conferring increased fitness. If this locus experiences

divergent selection in the two natural populations, it could promote their reproductive

isolation. Finally, although hybridization largely reduced allele fixation in the F1s initially,

indicating heterosis is likely due to the masking of deleterious alleles, the emergence of

reproductive isolation between populations may diminish the longer-term benefits of genetic

rescue in F2 or advanced-generation hybrids. As Torrey pine is a candidate for

interpopulation genetic rescue, caution is warranted where longer-term gene flow between

diverged populations may result in reduced fitness if barriers have evolved.

Background

41
Forest trees are well-known for maintaining signatures of local adaptation among populations

even when experiencing high levels of homogenizing gene flow (Petit and Hampe 2006;

Janes and Hamilton 2017; Cannon and Petit 2020). This apparent paradox is explained by

strong competition and selection at early life stages: deleterious non-adaptive alleles

introduced by migrant individuals or pollen are likely to be removed from the gene pool

(Petit and Hampe 2006). In some cases, population differentiation may lead to reproductive

isolation and eventually speciation (Schluter 2001; Hendry et al. 2007; Schluter 2009;

Hendry 2009; Nosil et al. 2009b; Andrew and Rieseberg 2013). However, the evolutionary

and genomic mechanisms underlying the shift from local adaptation with gene flow to the

evolution of reproductive isolation is poorly documented in trees, particularly conifers (Bolte

and Eckert 2020). The prevalence of local adaptation even under high levels of gene flow

makes trees a good system for better understanding how and when divergent selection may

drive the evolution of reproductive isolation, a prerequisite for speciation.

Speciation requires the evolution of genetic divergence, which can arise in response to

environmental differences, as a result of isolation, or a combination of both (Schluter 2001;

Nosil et al. 2009b,a). Allopatry may cause intrinsic reproductive incompatibilities to evolve

between populations, resulting in reproductive isolation if they come into secondary contact.

However, reproductive isolation can evolve even when gene flow is occurring if selection

against migrants and hybrids is strong enough (Schluter 2009). If non-local alleles are

deleterious, selection against them may favor the evolution of reproductive barriers. Extrinsic

postzygotic reproductive barriers may occur when environmental selection results in lower

hybrid or migrant fitness (as found by Lowry, Rockwood, et al., 2008; Melo et al., 2014;

Richards & Ortiz-Barrientos, 2016), while intrinsic postzygotic barriers result in lower hybrid

fitness regardless of the environment (Coughlan and Matute 2020). While both types of

barriers can be involved in speciation, with great variation in the strength of each type of

barrier across systems, extrinsic postzygotic barriers are generally stronger than intrinsic

postzygotic barriers (Christie et al. 2022). Additionally, extrinsic postzygotic barriers are

stronger in ecotypes than in species, suggesting they may be more important at earlier stages

of speciation (Christie et al. 2022) with intrinsic postzygotic isolation being important at later

stages, either as a result of divergence or by contributing to reinforcement (Coughlan and

Matute 2020).

As reproductive isolation can arise through many mechanisms, including both

prezygotic and postzygotic barriers, identifying the genes underlying these reproductive

barriers can help us better understand the process of speciation at the genomic level (Lowry

et al. 2008a; Rieseberg and Blackman 2010; Feder et al. 2012; Strasburg et al. 2012; Choi et

al. 2020; Schluter and Rieseberg 2022). One way to identify mechanisms underlying the

evolution of reproductive isolation is through signatures left in the genome (Nosil et al.

2009a; Schluter 2009; Feder et al. 2012; Strasburg et al. 2012). When reproductive isolation

is developing amidst ongoing gene flow, as in the early stages of ecological speciation where

species diverge as a consequence of natural selection among contrasting environments, loci

that are under selection will become more differentiated between two populations than other

genomic regions (Feder et al. 2012). As speciation proceeds, variants linked to those regions

under selection diverge, eventually leading to differentiation throughout the genome.

Analysis of the functions of these divergent regions of the genome can be complemented with

phenotypic data where both populations are grown in a common environment. Using

genotype-phenotype associations, differentiated loci can be linked to variance in reproductive

or fitness-associated traits. As reproductive-aged common gardens are uncommon for long-

lived species, our understanding of reproductive barriers and their genomic underpinnings in

trees is limited.

In this study, we took advantage of a 10-year-old common garden with Torrey Pine

(*Pinus torreyana* Parry), established in 2007 by the USDA-Forest Service Pacific Southwest

Research Station. Torrey pine is one of the rarest pines in the world with only two native

populations remaining (Di Santo et al. 2022). Separated by approximately 280 km (170

miles), the island population native to Santa Rosa Island, one of the Channel Islands,

experiences cooler temperatures and greater precipitation on average than the mainland

population located at the Torrey Pine State Natural Reserve in La Jolla, California (e.g., mean

temperature of the driest quarter is 16.7 °C at the island population and 18.9 °C at the

mainland population; precipitation of the warmest quarter is 17 mm at the island population

and 12 mm at the mainland population, Appendix S1 and S2). When grown in a common

environment, the two populations exhibit differences in growth, cone, and needle morphology

consistent with adaptation to their native environments (Hamilton et al. 2017), but also

exhibit relatively low genetic differentiation (F_{ST} = 0.0129) and some evidence of ongoing

gene flow (Di Santo et al. 2022). In an open-pollinated common garden including mainland

and island trees, the only F1s produced were from island maternal trees fertilized by

mainland pollen, with no F1s observed between mainland maternal trees fertilized by island

pollen (Hamilton et al. 2017). Asymmetric crossing barriers can be evidence of cytonuclear

or maternal incompatibility (Tiffin et al. 2001; Turelli and Moyle 2007; Case et al. 2016), and

may be a sign that reproductive barriers between the two populations have already begun to

evolve (Barnard-Kubow et al. 2016). Thus, Torrey pine is an ideal system for investigating

the early stages of reproductive isolation between two geographically separated populations

occurring in contrasting environments. To evaluate the genomic and fitness consequences of

inter-population gene flow, we leverage a common garden experiment of reproductive-aged

trees of both parental populations and their F1 hybrids.

Understanding the nature of the divergence between Torrey pine populations will be

critical to the species' management. Both populations exhibit exceedingly low genetic

diversity, particularly for a conifer (Ledig and Conkle 1983; Farjon 2013; Di Santo et al.

2022), and F1 hybrids between the two populations appear to have higher fitness proxies than

the parental populations, suggesting genetic rescue may be beneficial (Hamilton et al. 2017).

However, increased F1 fitness may be the result of heterosis, in which excess homozygosity

of deleterious recessive alleles is reduced (Williams and Savolainen 1996; Tallmon et al.

2004; van de Kerk et al. 2019). Given this, it is unclear whether increased hybrid fitness

would persist in future generations of hybrids backcrossed with either parental population. If

the parental populations have evolved partial reproductive isolation as a result of ecological

speciation, F2s and backcrossed individuals may ultimately have lower fitness despite

heterosis in F1s, limiting the value of inter-population genetic rescue (Walter et al. 2020;

Christie et al. 2022).

In this paper, we investigate whether there is evidence for incipient ecological

speciation between Torrey pine populations by testing whether loci with reduced admixture

in F1 hybrids are associated with fitness-related phenotypic differences. We address three

main questions: 1) Are there genomic signatures of reduced admixture between the two

Torrey Pine populations? 2) If so, are alleles with reduced admixture linked to phenotypes or

functions that indicate they may be involved in local adaptation and underlie extrinsic

postzygotic isolation? 3) What are the implications for conservation of this endangered

species – would genetic rescue be beneficial or should these two populations be managed as

separate taxonomic groups?

Methods

Multigenerational common garden

Prior to 1960, two plots of 20 mainland and 20 island Torrey pine trees were established

adjacent to each other at the USDA Horticultural Field Station (now the Scripps Institute) in

La Jolla, CA (Haller 1967; Hamilton et al. 2017). In 2004, open-pollinated cones from 10

mainland and 16 island trees were sampled in the plots, yielding a total of 643 seeds. In 2006,

seeds were planted and grown in a greenhouse at the USDA-Forest Service Pacific Southwest

Station, Institute of Forest Genetics, Placerville, CA. As seedlings emerged, one cotyledon

was clipped from each plant for genetic analyses and identification of seedlings' ancestry

(island, mainland, or F1 hybrid) using two population-specific allozyme markers (Ledig and

Conkle 1983). Of the 643 seeds planted, 128 were genetically confirmed as being of pure

mainland ancestry, 134 were identified as being of pure island ancestry, and 381 were

identified as mixed ancestry (F1 hybrids) (Ledig and Conkle 1983; Hamilton et al. 2017).

These F1 hybrids solely reflected crossings of island females with mainland pollen donors,

indicating hybrids were the product of unidirectional gene flow between parental populations.

In 2007, a subset of 360 seedlings representing each ancestry (n=120 each of island,

mainland, and F1 hybrids) were planted following a randomized complete block design with

six replicate blocks at the common garden site in Montecito, CA (34.42963, -119.56396).

Each of the 360 trees planted were measured for four quantitative traits once a year

(Appendix S3), including tree height in cm (measured between 2008 and 2018), the number

of conelets or first-year cones produced (measured between 2013 and 2018), the number of

immature cones or second-year cones produced (measured between 2015 and 2018), and the

number of mature or third-year cones produced (measured between 2015 and 2018). These

traits were selected as they provide multiple proxies of fitness, incorporating both growth and

reproductive output needed to examine the impact of interpopulation gene flow to Torrey

pine fitness (see below). Given that selection likely acts on each reproductive stage and there

was limited correlation across traits over time (see below), we analyzed each cone production

trait separately using all available years. For full details about the common garden experiment

see Hamilton et al. (2017).

To understand the effects of climate on fitness, we compared climatic variables from

the two native Torrey Pine populations and the common garden site, extracting data from

climate rasters from *BioClim* version 2.1 for the years 1970-2000 at 30 second resolution

(Fick and Hijmans 2017) using the R package *terra*, version 1.7.71 (Hijmans 2024) in R 4.3.2

(R Core Team 2023). The common garden site is a novel environment compared to either

native site, and is more similar to the island site along PC1 (Appendix S2). However, because

the island site generally experiences more moderate conditions (lower temperatures and more

precipitation during the warmest quarter, Appendix S1 and S2), and because the severity of

the summer drought is the primary stressor in Mediterranean climates and is likely to be a

more important factor in adaptation (Granda et al. 2014; Nardini et al. 2014; Riordan et al.

2016; DeSilva and Dodd 2020), we considered the common garden site to be more similar to

the native mainland site.

DNA extraction and reduced representation sequencing

During the summer of 2016, needle tissue was collected from the 262 individuals surviving

of the 360 seedlings originally planted within the common garden (Hamilton et al. 2017),

including 89 pure mainland, 88 pure island, and 85 F1 hybrid trees. Following collection,

needles were dried and maintained on silica gel until genomic DNA could be extracted using

approximately 20 to 35 mg of dried needle tissue based on a modified version of the CTAB

protocol (Doyle and Doyle 1987). In this modified version, rough mixing (e.g., vortex) was

replaced with slow manual shaking of samples to reduce DNA shearing. The concentration

and purity of extracted DNA was evaluated for each sample separately using a NanoDrop

1000 Spectrophotometer (Thermo Scientific). Overall, purity ratios averaged 1.86 and 2.21

for 260/280 and 260/230 respectively, with concentrations of DNA ranging from 153.78 to

4,463.31 ng/µL (average = 1,138.41 ng/µL). Following extraction, DNA concentration was

standardized to 85 ng/µL for all 262 samples and reduced-representation sequencing libraries

were prepared using the protocol described in Di Santo et al. (2022). Once constructed, all

libraries were pooled and sent to the Genomic Sequencing and Analysis Facility (GSAF;

Austin, TX) for single-end sequencing (1x100 bp) on 5 lanes of an Illumina HiSeq 2500.

Prior to sequencing, the pooled library was size-selected for fragments within the range of

450-500 bp.

Reference-guided assembly and calling of genetic variants

Raw sequenced libraries were demultiplexed using *ipyrad* version 0.9.12 (Eaton and Overcast

2020). To reduce the probability of assigning a read to the wrong individual, *ipyrad* was

parameterized to allow only a single mismatch in the barcode sequence. Raw reads were

subsequently filtered for quality using *dDocent* version 2.8.13 (Puritz et al. 2014a,b). Within

dDocent, using *fastp* (Chen et al. 2018), base pairs with PHRED score below 20 at the

beginning and end of reads and Illumina adapter sequences were removed, and reads were

trimmed once the average PHRED score dropped below 15. Lastly, reads where more than

50% of base pairs had a PHRED score below 15 were discarded. Cleaned reads were mapped

to the *Pinus taeda* (loblolly pine) draft genome version 2.0 (GeneBank accession:

GCA_000404065.3) using the *BWA-MEM* algorithm (Li 2013) with default values, except for

the mismatch penalty (-B, default: 4) and the gap open penalty (-O, default: 6), which were

relaxed to 3 and 5, respectively, to account for additional genomic differences that may have

evolved between Torrey pine and closely related loblolly pine. Genetic variants were called

from sequence alignments using the *SAMtools* and *BCFtools* pipeline (Li 2011). Specifically,

we used the multiallelic and rare-variant calling model (-m) with no prior expectation on the

substitution rate (--prior 0). In total, the pipeline identified 44,493,992 genetic variants,

including single-nucleotide (SNP) and insertion/deletion (INDEL) polymorphisms, that were

subjected to quality filtering. First, variants with a genotype quality (GQ) < 20 and a

genotype depth (DP) < 3 were marked as missing. Next, variants with PHRED scores $\leq$ 20

(QUAL), minor allele counts (MAC) < 3, minor allele frequencies (MAF) < 0.01, genotyping

rates across individuals < 0.8, average depth across individuals > 23 (based on the equation

from Li 2014), and inbreeding coefficients (F_{IS}) < -0.5 or > 0.5 were filtered out of the raw

data. Of the initial 262 genotyped individuals, 53 were removed from subsequent analyses as

they exhibited greater than 50% missing values. In total, following the removal of INDELs

and SNPs with 3 or more alleles, 11,379 biallelic SNPs across 209 individuals, including 68

pure island, 75 pure mainland, and 66 F1 hybrid individuals were identified and used for

analysis.

Parental population hybridization simulations

To evaluate whether barriers to gene flow may have evolved between Torrey pine

populations, we quantified genome-wide admixture in F1 hybrids, measured as locus-specific

observed heterozygosity (H_O), specifically searching for SNPs with reduced heterozygosity.

Using a customized R script (R version 4.1.3) relying on R packages *adegenet* version 2.1.5

(Jombart 2008; Jombart and Ahmed 2011) and *hierfstat* version 0.5.10 (Goudet and Jombart

2021), we simulated the distribution of H_O values for each of the 11,379 SNPs expected in

the F1 hybrids under the null hypothesis that H_O at any locus would be based on the

frequency of alleles in the island and mainland population. To do so, we used a 2-step

approach within the function *hybridize()* that simulates hybridization between two

populations. First, allele frequencies are derived from genotypes specific to each parental

population. Then, hybrid genotypes are generated by sampling gametes in each of these

populations using a multinomial probability distribution. SNP-specific null distributions were

produced by repeating this process 1000 times, estimating locus-specific H_O each time from

simulated hybrid genotypes using the function *basic.stats()*. To mirror the empirical SNP data

set, we simulated 11,379 genotypes for 66 F1 hybrids within each iteration, using allele

frequencies derived from 68 and 75 pure island and pure mainland individuals, respectively.

To identify SNPs that may exhibit reduced heterozygosity in F1s relative to

expectation, we compared simulated null H_O distributions (expected H_O) with H_O values

estimated from empirical F1 hybrid genotypes (observed H_O). For each locus, we computed a

one-sided p-value, defined as the probability that an expected H_O value is equal or lower than

the observed H_O value at this locus. All p-values were corrected for multiple testing using

Benjamini and Hochberg's (Benjamini and Hochberg 1995) False Discovery Rate procedure

implemented within the R function *p.adjust()*. SNPs that exhibited significantly reduced

heterozygosity relative to expectation based on parental allele frequencies (FDR < 0.1) were

classified as potential candidate regions of the genome associated with the evolution of

barriers to gene flow between island and mainland populations of Torrey pine.

Functional plant ontology (PO) and gene ontology (GO) annotation

SNPs exhibiting reduced heterozygosity relative to expectations were functionally annotated

to determine whether their functions, processes, or anatomical and temporal expressions may

be important to Torrey pine fitness using *blastx* 2.9.0+ (Altschul et al. 1990, 1997; Camacho

et al. 2009) and a combination of customized R scripts. First, a region of 5,000 bps (5 kbps)

from *Pinus taeda* draft genome version 2.0 (GenBank accession: GCA_000404065.3) was

extracted 2,500 bps before and after each target locus using *samtools* (Li et al. 2009). If a

SNP was within the first 2,500 bp of a scaffold, the first 5,000 bps were extracted. If a

scaffold harboring a SNP was shorter than 5 kbps, the whole scaffold length was used

instead. Extracted sequences were then blasted against the TAIR10 peptide blastset

(TAIR10_pep_20101214_updated). Homology between query and blasted sequences were

assessed using *blastx* default parameters, an expectation value threshold of 10^{-3} (-evalue

0.001), and a maximum number of database hits of 5 (-max_target_seqs 5). Lastly, mapping

of gene ontology (GO) and plant ontology (PO) terms onto annotated sequences was

performed in R using a customized script and GO (ATH_GO_GOSLIM.txt.gz) as well as PO

(grow.txt.gz) annotations available online. All databases mentioned throughout this section

are available from https://www.arabidopsis.org/download/. While neither conifer- nor pine-

specific, using the TAIR database provides a more exhaustive set of GO and PO annotations

to associate with reduced admixture than is available for conifers. This database remains the

most complete with respect to plant gene functions and shares orthologous and homologous

sequences with pine that will aid in determining functional categories critical to plant fitness

(e.g., Eckert et al. 2010). Functional gene annotation is limited for conifers, particularly for

genes involved in reproduction and development (De La Torre et al. 2020). Furthermore,

some genes involved in flowering in angiosperms pre-date their divergence with

gymnosperms (Moyroud et al. 2017; Liu et al. 2018; De La Torre et al. 2020). However,

because of the evolutionary distance between *Arabidopsis* and Torrey pine, the annotations

will be best understood within broad functional categorizations.

Following the same procedure as above, we also annotated 5 kb regions around 500

randomly selected SNPs of the total 11,379 called to investigate whether the set of candidate

loci for reproductive isolation was enriched or depleted for specific functions or processes

using Fisher's exact test for count data as implemented in the R function *fisher.test()*. For

each annotation (GO or PO), we constructed a two by two contingency table recording the

number of successes (the number of times an annotation was observed) and failures (the sum

of successes across all annotations minus the number of successes for the annotation of

interest) in the candidate and the random set of loci, and tested the null hypothesis of

independence of rows (candidate, random) and columns (success, failure). When an

annotation was absent from one of the sets of loci, we considered the number of successes

associated with that annotation and set of loci to be zero. Resulting p-values were corrected

for multiple testing within each annotation data set (GO or PO) and category within

annotation data sets (for GO: biological process, molecular function, and cellular component)

using Benjamini and Hochberg's (Benjamini and Hochberg 1995) False Discovery Rate

procedure implemented within the R function *p.adjust()*. We assumed an annotation to be

either enriched or depleted within a particular set of loci when the false discovery rate

associated with the odd ratio for that annotation was inferior to 10% (FDR < 0.1).

Correlation analysis between measured phenotypes

To limit redundancy across traits and ensure each phenotype largely reflects unique inter-

individual variation, we conducted a repeated measures correlation analysis in R based on

temporal measurements taken on all 209 samples using the package *rmcorr* (Bakdash and

Marusich 2022). Significance of estimated correlation coefficients was evaluated considering

a threshold $\alpha = 0.05$. A correlogram with correlations between all pairs of phenotypes was

generated using the R package *corrplot* (Wei and Simko 2021) and is shown in Appendix S4.

While some correlation coefficients were significant, no phenotypes were highly correlated (r

≥ 0.6). Consequently, all four temporally assessed traits – tree height in cm, the number of

conelets (first-year cones), the number of immature cones (second-year cones), and the

number of mature cones (third-year cones) produced – were kept for subsequent analyses.

Genome-wide association analysis (GWAS)

To assess whether genotypic variation at select SNPs may be associated with variability in

phenotypes measured across all available years data (Appendix S3), we conducted a

repeated-measures genome-wide association analysis based on all 209 samples and 11,379

genetic variants using the R package *RepeatABEL* (Rönnegård et al. 2016). For each trait, the

function *preFitModel()* was used to fit a linear mixed model to the data without including

SNP effects to estimate variance components for the trait, including a random polygenic

(based on a genetic relationship matrix, Appendix S5) and permanent environmental effect.

Including these effects (computed and assessed internally within the function) accounted for

genetic relatedness among individuals and repeated annual measures within the model. In

addition, "block" (spatial position of a sample within the common garden) and "year" were

considered as random variables within the model. Following this, we used the function

rGLS(), fitting a generalized least square model to each genetic marker given a covariance

matrix (estimated using *preFitModel*), to test for associations between SNPs (considered as

fixed effects) and phenotypes while correcting for the effect of random variables. Lastly,

SNP-specific p-values were corrected for multiple testing using Benjamini and Hochberg's

(Benjamini and Hochberg 1995) False Discovery Rate procedure implemented within the R

function *p.adjust()*. Of the 11,379 SNPs tested for association with each of the four

phenotypic traits, only those with FDR < 0.1 were considered potential candidate markers

underlying genotype-phenotype association.

Evaluating fitness consequences following interpopulation genetic mixing

One locus was identified that exhibited both lower F1 heterozygosity than expected under a

neutral model and was statistically associated with one or more of the four fitness or fitness-

related phenotypes (hereafter referred to as the shared locus, see Results). To suggest whether

this locus could be a potential candidate associated with extrinsic reproductive isolation

between Torrey pine populations, we required that (1) genotype frequencies for the locus

would be biased by population, with each population favoring a different homozygous

genotype, and (2) individuals with non-local or admixed genotypes would exhibit reduced

fitness relative to individuals with the more local genotype. As the common garden was

planted in Montecito near Santa Barbara, CA, we considered mainland trees as local and

island trees as non-local. To determine whether the frequency of homozygous genotypes (i.e.,

0/0 and 1/1) statistically differed between parental ancestries, we counted genotype

occurrences across pure island and pure mainland individuals and conducted a Pearson's Chi-

squared test with Yates' continuity correction as implemented in the function *chisq.test()* in R

for the resulting contingency table.

A linear mixed model and expected marginal means approach was used to evaluate

whether average phenotypes statistically differed among genotypes by leveraging R packages

lme4 (Bates et al. 2015) and *emmeans* (Lenth 2023). For each trait, we built a model where

phenotypic variation was explained by the fixed effect of individuals' genotypes at the shared

locus, the fixed effect of among-individual genetic relationships, the random effect of blocks

assigned to individuals within the common garden, the random effect of year the phenotypes

were measured, and the random effect of individuals themselves (to account for the repeated

nature of response variables). Genetic relationships among common garden individuals were

estimated using a principal component analysis implemented within the *dudi.pca()* function

from the *adegenet* R package. This analysis was performed on allele frequencies of the full

genomic data set (11,379 SNPs across 209 individuals), with missing values replaced by the

average allele frequencies across individuals. The first two principal components were used

in all four linear mixed models to summarize genetic relationships among individuals, as

these components provide good proxies for among-population (PC1) and within-population

genetic structure (PC2) (Appendix S6).

The *emmeans()* function, computing expected marginal means, was used for the

assessment of phenotype average differences among genotypes. Briefly, the function

computes genotype-specific trait averages and standard errors corrected for all effects

included within linear mixed models. The function also computes adjusted p-values estimated

from t-ratios calculated between all possible pairwise genotype comparisons. Ultimately, we

used this statistic to infer statistical differences in average phenotypes for all traits measured

between genotypes. For this analysis, 175 of the total 209 Torrey pine trees present within the

full genomic data set were used, as genotypes at the shared locus were missing for 34

individuals.

Evaluating the distribution of genetic variation across Torrey pine ancestries and SNP sets

The distribution of genetic diversity estimated as observed heterozygosity (H_O) was compared

between pure island, pure mainland individuals, and F1 hybrids for the whole genomic

dataset (N=11,379 SNPs), SNPs significantly explaining variation in fitness-related traits

(N=12 SNPs), and SNPs exhibiting reduced F1 heterozygosity relative to expectations

(N=185 SNPs). In addition, for each ancestry separately, we quantified the number of loci

across the whole genomic dataset that were fixed. A locus was considered fixed within an

ancestry when all individuals of that ancestry at that locus were homozygous for either the

reference or alternate allele (0/0 or 1/1). H_O and numbers of fixed loci were both estimated in

R. While H_O was computed using the *basic.stats()* function implemented within the package

hierfstat, numbers of fixed loci were computed manually.

To determine whether differences in genetic diversity exist among ancestries within

SNP datasets, we performed three Dunn's tests using the package *dunn.test* in R. This test is a

nonparametric equivalent of analyses of variance and post-hoc tests, as normality for

estimated observed heterozygosities within ancestries and SNP sets, assessed either visually

or using Shapiro-Wilk normality test, could not be assumed. We accounted for multiple

testing by correcting p-values associated with each comparison using Benjamini and

Hochberg's (Benjamini and Hochberg 1995) False Discovery Rate procedure. Two H_O

averages were considered significantly different when the false discovery rate dropped below

10% (FDR < 0.1).

Finally, Fisher's exact test for count data as implemented within the function

fisher.test() in R was used to compare the number of loci that are homozygous for either

reference or alternate alleles between mainland, island, and F1 individuals. We first leveraged

a three by two contingency table recording the number of successes (the number of fixed

loci) and the number of failures (the total number of loci minus the number of fixed loci)

across all three ancestries, testing the null hypothesis of the independence of rows (island,

mainland, F1 hybrid) and columns (success, failure). In addition, we used three two by two

contingency tables of the number of successes and failures (defined as above) to test the null

hypothesis of the independence of rows (F1 hybrid, island; F1 hybrid, mainland; island,

mainland) and columns (success, failure). The p-value associated with each pairwise

comparison was corrected for multiple testing using Benjamini and Hochberg's (Benjamini

and Hochberg 1995) False Discovery Rate procedure implemented within the R function

p.adjust(). We considered a difference to be significant when the false discovery rate dropped

below 10% (FDR < 0.1).

Results

SNPs exhibiting lower-than-expected F1 heterozygosity and their functional relevance

Comparison of simulated F1 hybrids between island and mainland individuals for 11,379

SNPs using empirical allele frequencies in each parental population revealed some loci that

exhibited a higher degree of homozygosity in the F1s relative to the expectation. A

comparison between observed and simulated H_O estimates at each locus identified 185 SNPs

(FDR < 0.1) with reduced F1 heterozygosity. Of the 184 5kb-long sequences containing these

SNPs (two SNPs were located on the same segment), 75 (41%) could be annotated with the

TAIR 10 peptide blastset, with hits in 59 described *A. thaliana* genes. Plant ontology terms

could be retrieved for 37 (63%) of the 59 gene hits, while gene ontology terms were retrieved

for all 59 gene hits.

Regions of the genome that exhibited reduced F1 heterozygosity relative to

expectation were associated with a variety of plant developmental stages (Appendix S7). This

includes embryo development stages (e.g., C globular stage, D bilateral stage, E expanded

cotyledon stage, or F mature embryo stage), vegetative development stages (e.g., 2-, 4-, 6-, 8-

, 10-, 12-leaf stages, or seedling development stage), and reproductive development stages

(e.g., 4 anthesis stage, petal differentiation and expansion stage, M germinated pollen stage,

or L mature pollen stage). Functionally, the majority of these loci are located or active in the

nucleus with molecular functions including protein binding, catalytic activity, RNA binding,

as well as kinase and transferase activities (Appendix S8A, B). Biological processes

associated with these molecular functions covered responses to various stimuli and stresses,

and plant development (Appendix S8C). Enrichment analyses demonstrated that while no PO

terms were preferentially associated with SNPs exhibiting lower heterozygosity than

expected in the F1, some GO terms were significantly enriched or depleted in the latter SNP

set (Figure 1, Appendix S9). Interestingly, assessment of enriched GO terms indicated the

products of genes associated with these loci may be preferentially located in the plasma

membrane and involved in important developmental and reproductive processes, including

cell growth, plant growth, and pollination, with molecular functions associated with signaling

receptor activity.

SNPs exhibiting statistical association with fitness traits and their proxies

Genome-wide association analysis indicated alleles at several loci may play an important role

in contributing to fitness variation in Torrey pine. In total, 12 SNPs were significantly

associated (FDR < 0.1) with variation in phenotypes (Table 1). While 8 (67%) of the 12 SNPs

were associated with a single trait, the remaining 4 were associated with two or more traits.

Phenotypic values of these traits all increased with the number of derived alleles at associated

SNPs except for one, locus_8778 [APFE031129984.1:16200] (Table 1), where it decreased.

Interestingly, comparing SNPs associated with fitness and fitness-related traits with SNPs

exhibiting low-level F1 heterozygosity identified one common locus (locus_4218

[APFE030529380.1:392010]). This locus, in addition to exhibiting a reduced degree of F1

heterozygosity (Appendix S10), was also the only locus that significantly explained

phenotypic variation across all four traits measured (Table 1). Unfortunately, functional

annotations for the 5-kbps region surrounding this locus were limited. While this region may

be located in the mitochondrion (GO:0005739), nothing is known about its molecular

function or the biological process it might be involved in.

Fitness consequences of hybridization at a locus exhibiting reduced admixture and

significant genotype-phenotype association

The chi-squared test performed on homozygous genotype counts for locus_4218

[APFE030529380.1:392010] demonstrated unequal distribution of homozygous genotypes

between pure parental ancestries (χ^2_1 = 104.1, P < 0.001, Appendix S10). Homozygotes for

the reference allele (0/0) were dominant among pure island individuals (94% of all

genotypes) while homozygotes for the derived allele (1/1) formed the core of pure mainland

individuals (97% of all genotypes). Given this result, we considered the reference allele (0) as

the island allele, and the derived allele (1) as the mainland allele.

The evaluation of expected marginal means (phenotype averages corrected for within-

and among-ancestry genetic structure, blocks within the common garden, year of trait

measurements, and sample ID) revealed unequal fitness across reference, derived, and

heterozygous genotypes. For three out of the four phenotypic traits, individuals homozygous

for the island allele exhibited reduced fitness, while individuals homozygous for the mainland

allele exhibited greater fitness, with heterozygous genotypes intermediate between the two

(Figure 2; Appendix S11). These traits included tree height, the number of conelets produced,

and the number of mature cones produced. For the number of immature cones produced, the

distribution of fitness among genotypes was slightly different. Individuals homozygous for

the island allele still exhibited the lowest fitness, but fitness of individuals homozygous for

the mainland allele and heterozygous individuals were the highest, with no significant

differences between groups. We considered the mainland allele (1) as the local, and the island

allele (0) as non-local because both the common garden site and the mainland population

experience warmer and drier summers (Appendix S1). Our results thus indicate that carrying

at least one copy of the local allele increased tree fitness, with a direct additive relationship

based on the dosage of mainland allele to fitness for tree height (cm), the number of conelets

produced, and the number of mature cones produced (Figure 2).

Distribution of genetic variation across island, mainland and F1 hybrid individuals

For the whole genomic dataset (N=11,379 SNPs), estimates of observed heterozygosity were

similar across ancestries (H_O = 0.229, 0.230, and 0.230 for F1 hybrid, island, and mainland

individuals, respectively; Figure 3). However, despite almost-identical average observed

heterozygosities, the number of loci homozygous for either the reference or the derived allele

(1/1 or 0/0) varied significantly among ancestries (Fisher's exact test, $P < 0.001$), with island

and mainland individuals exhibiting four times greater number of fixed alleles than that

estimated for F1 hybrids (Appendix S12). While these results may seem contradictory, they

can be explained by a proportionally higher number of loci with extremely low

heterozygosity in the F1s relative to island or mainland individuals (Appendix S13).

Average observed heterozygosity estimated for loci exhibiting reduced F1

heterozygosity for hybrids (0.159) was significantly lower than that for island (0.230) and

mainland (0.232) individuals (Figure 3). If they were to contribute to extrinsic reproductive

isolation, we would expect heterozygosity at these loci to be much lower in parental

populations than in F1s as well as to be highly differentiated between island and mainland

populations due to divergent selection promoting fixation of distinct alleles. Because average

heterozygosity in F1 hybrids at these loci was lower than that found for both island and

mainland individuals and given that these loci did not only comprise those highly

differentiated between parental populations (Appendix S14), our results suggest intrinsic

genetic incompatibilities may potentially have evolved between populations, with some

allelic combinations following hybridization leading to reduced hybrid fitness.

Finally, for SNPs associated with fitness proxies, we observed exceedingly low

genetic variability (Figure 3), with average estimates of heterozygosity for F1, island, and

mainland individuals evaluated at 0.113, 0.066, and 0.071, respectively. These results

indicate that all three groups have reduced genetic variation available for selection to act

upon. However, F1 hybrids did exhibit significantly greater heterozygosity for genetic

variants associated with fitness when compared to island individuals.

Discussion

Local adaptation is common in tree species (Savolainen et al. 2007; Lind et al. 2017, 2018;

Gugger et al. 2021), but it is unclear when and how population differentiation may drive the

evolution of reproductive isolation. Understanding whether reproductive isolation has

developed between two populations is essential to species management, particularly where

intra-specific gene flow may be a beneficial source of genetic variation or may be a

hindrance, leading to outbreeding depression. In Torrey pine, previous work has shown that

the island and mainland populations are genetically and phenotypically different despite some

recurrent gene flow, consistent with adaptation to differing environments (Hamilton et al.

2017; Di Santo et al. 2022). Furthermore, asymmetric gene flow between populations when

grown together provides some evidence for the evolution of reproductive isolation (Hamilton

et al. 2017). Together, this suggests that the two Torrey pine populations could be at an early

stage of speciation. Here, we tested for further evidence for the evolution of reproductive

isolation by identifying loci exhibiting less heterozygosity in F1 hybrids than expected given

allele frequencies in parental populations, suggesting selection against heterozygotes. One

locus that was highly diverged among populations and showed signatures of selection against

F1 hybrids was associated with fitness differences in the common garden, and could be a

candidate locus for ecologically-driven reproductive isolation between island and mainland

Torrey pine populations. Overall, loci with low heterozygosity in F1 hybrids were enriched

for functions important in reproduction, such as growth and pollination, suggesting these

could be the mechanisms underlying the evolution of reproductive isolation between the two

populations.

While demographic modeling has provided evidence for historic speciation with gene

flow in pines (Menon et al. 2018; Bolte et al. 2022), the specific molecular mechanisms

underlying reproductive isolation in conifers are not well understood (Bolte and Eckert

2020). We identified loci that may be involved in the evolution of reproductive isolation

between two populations, including pollination- and growth-related genes, and one locus

associated with fitness differences expected when hypothesizing local adaptation. In

combination with asymmetric gene flow, this suggests that both intrinsic and extrinsic

barriers could be contributing to reproductive isolation in Torrey pine. These results thus

identify potential molecular mechanisms underlying the process of reproductive isolation in a

critically endangered pine and suggest that differential adaptation could be a driving force,

which is critical information required for optimal conservation management of the species.

One locus shows patterns suggestive of extrinsic postzygotic reproductive isolation

We identified a locus that is divergent between the two populations and is also associated

with fitness differences (locus_4218 [APFE030529380.1:392010]). Individuals with the

derived "mainland" variant of this locus had higher growth and reproduction in the mainland

common garden than individuals homozygous for the reference "island" allele, consistent

with local adaptation to the hotter, drier environments shared by the mainland population and

the common garden environment (Appendix S1, S2). Torrey Pine is likely a relictual endemic

species that once had a larger range and became restricted to coastal climates as climate

became warmer and drier (Axelrod 1967; Williams et al. 2008), and thus the island

population may have a climate more similar to its ancestral climate. Demographic modeling

suggests that the two modern-day Torrey Pine populations formed after an ancestral

population split, forming a slightly larger island population (N_e = 2305) and smaller mainland

population (N_e = 1715) approximately 1.2 million years ago, with gene flow occurring

following divergence (Di Santo et al. 2022). It is possible that the mainland population has

undergone additional evolutionary change following divergence to adapt to the combined

stresses of warm and drier summers associated with the mainland environment, and that this

locus is a candidate for local adaptation and divergence of the mainland population. The

differentiation between populations for this allele could result from differential selection if

heterozygous hybrids have lower fitness in both natural populations relative to individuals

that are homozygous for the local variant. If this locus experiences differential selection

between the two Torrey pine populations, it could contribute to reproductive isolation. While

our results suggest that this locus may be beneficial in the hotter, drier summers experienced

by mainland environments relative to the island population (Appendix S1), a reciprocal

transplant of the two populations and their F1 hybrids would be required to determine

whether the non-local variant decreases fitness in both natural populations.

Reduced average heterozygosity at this locus in approximately 10-year-old F1 hybrids

(0.028 compared to a neutral expectation of 0.936) suggests selection against heterozygotes

with one copy of the island and one copy of the mainland allele, or that this locus is linked to

such a gene under selection. Reduced heterozygosity could result from intrinsic prezygotic or

postzygotic barriers (e.g., incompatibilities affecting pollination, germination, or

development of seedlings), extrinsic environmental selection in the common garden in the

following ten years, or both. In the common garden, heterozygotes were intermediate in

fitness metrics between island homozygotes with lower fitness and mainland homozygotes

with higher fitness (although only the number of immature cones was significantly different

between island homozygotes and heterozygotes, Figure 2). Our evidence suggests that

selection against heterozygotes occurred at early developmental stages, reducing the expected

number of heterozygous individuals, and also that the surviving heterozygous individuals in

the common garden had intermediate fitness across multiple years of growth and

reproduction measurements. This may seem contradictory, but it could indicate that this allele

has different fitness effects at different developmental stages, or that pleiotropic interactions

with other alleles result in selection against heterozygotes only at certain developmental

stages or in combination with specific variants of other alleles.

While there is limited functional annotation for genomic regions proximal to this

locus, it does have the annotation "mitochondrion" (GO:0005739). As heterozygotes are

present, it is probably not encoded in the mitochondrion and may represent ancient paralogy

between nuclear and mitochondrial genomes. Alternatively, the locus could also be part of a

nuclear mitochondrial gene present in the nucleus following mito-nuclear gene transfer, with

its product interacting with mitochondria. Consequently, the locus could be involved in

cytoplasmic incompatibility between the maternally-inherited mitochondrion and nuclear

paternal genes (such as those involved in pollen tube growth) (Turelli and Moyle 2007;

Rieseberg and Blackman 2010), or with the chloroplast, which is paternally inherited in pines

(Neale and Sederoff 1989; Mogensen 1996).

Local adaptation is generally polygenic in forest trees (Yeaman et al. 2016;

MacLachlan et al. 2021), suggesting ecological speciation may also have a broad genetic

basis (Rose et al. 2018; Schluter and Rieseberg 2022). However, we only find one locus

showing evidence consistent with ecological speciation. Because we used ddRADseq data

and only sampled a fraction of the genome of Torrey pine, it is likely that other loci

underlying reproductive isolation and local adaptation exist, but were not captured in this

study. Nonetheless, the fact that we identified a locus linked to both reproductive isolation

and fitness despite this limitation suggests that the pattern may be more prevalent throughout

the genome and could be better assessed using whole-genome sequence data.

Functions of genes exhibiting reduced heterozygosity

If loci exhibiting reduced heterozygosity in F1s are "speciation genes" underlying hybrid

incompatibilities, differential adaptation, or both (or linked to involved genes), they should be

enriched for related functions (Rieseberg and Blackman 2010; Wright et al. 2013; Walter et

al. 2020; Schluter and Rieseberg 2022). Intrinsic barriers are likely to be related to pollination

or reproduction, and extrinsic barriers may be involved in traits under divergent selection,

such as abiotic or biotic conditions. We found that highly differentiated loci were located

near genes that were enriched for a wide range of functions (Figure 1, Appendix S9). Similar

to previous studies, enriched functions included those related to pollination and development

("growth", "pollination", "cell growth") (Rieseberg and Blackman 2010; Leroy et al. 2020).

These regions may be involved in intrinsic reproductive barriers, such as pollination

incompatibilities associated with female cones or failed embryo development post-

fertilization (McWilliam 1959; Fernando et al. 2005). Other enriched functions include

"plasma membrane", "signaling receptor activity", or "protein metabolic process". Given

these broad categorizations it is difficult to determine whether these genes could be involved

in adaptation to contrasting environments, particularly as only one differentiated gene was

associated with fitness differences in the common garden. Previously, island seedlings were

found to germinate later and have a reduced growth rate relative to mainland seedlings

(Hamilton et al. 2017). The enrichment of loci with the "growth" and "cell growth" GO terms

may be related to the evolution of differential growth rates between parental populations.

However, it is also possible that the enriched functions are associated with fitness in traits or

environmental conditions that were not measured in the common garden. Because the loci

with reduced heterozygosity in F1 hybrids were investigated after ten years of growth in the

garden, we cannot determine the stage at which selection against heterozygotes occurred in

order to distinguish between intrinsic and extrinsic postzygotic barriers. Crossing

experiments and monitoring of hybrids produced between the two populations would provide

a more direct approach to determine the developmental stage at which potential barriers act

(Christie et al. 2022).

Conservation implications

Similar to previous studies, genome-wide estimates of heterozygosity for the island and

mainland populations were low (Di Santo et al. 2022), suggesting a lack of evolutionary

potential. Additionally, we found in the present study exceedingly low diversity at loci

putatively underlying trait variation in the species, further supporting these results.

F1 hybrids between mainland and island populations exhibit heterosis, with a higher

growth rate and fecundity than either parental population (Hamilton et al. 2017). The four-

fold reduction in the number of fixed alleles when compared to pure island and pure

mainland Torrey pine trees, and the almost two-fold increase in average heterozygosity at loci

exhibiting significant genotype-phenotype associations when compared to pure island Torrey

pine trees indicate that heterosis in F1 hybrids may result from the masking of deleterious

mutations. Alone, these results would suggest that natural populations could benefit from

genetic rescue, where hybridization between the two populations could introduce genetic

variation to alleviate genetic load. However, our results here warrant caution before

introducing non-local or hybrid trees into the wild. If the natural populations have already

developed reproductive isolation or are locally adapted, hybridization may result in decreased

fitness that only presents in F2s or later generation hybrids (Lowry et al. 2008a; Walter et al.

2020; Christie et al. 2022), or only in field conditions where they are exposed to stresses not

present in the common garden (Melo et al. 2014). Our data suggests that the island and

mainland populations should be managed separately at present, but that future work should

use crossing experiments and reciprocal transplants to determine whether genetic rescue

could have long-term benefits to this endangered tree species.

Conclusions

Taken together, we find evidence of asymmetric barriers to gene flow, reduced

heterozygosity in genomic regions related to varying reproductive and developmental

functions, and one low-heterozygosity and highly differentiated locus that associates with

reduced fitness in individuals carrying one or two non-local (island) alleles for all four

measured fitness metrics in Torrey Pine. These results suggest that Torrey Pine populations

may be beginning to evolve reproductive isolation, and this may partly be driven by

adaptation to contrasting island and mainland environments, in which case introducing non-

local individuals or hybrids to the natural populations for genetic rescue may actually be

harmful in the long term. In the future, whole-genome sequencing combined with reciprocal

transplant experiments and the development of experimental crosses including F2s and

backcrosses will be poised to identify the genes that may be under divergent environmental

selection in the two populations, and/or contributing to the evolution of reproductive isolation

(Schluter and Rieseberg 2022). Furthermore, understanding the extent of reproductive

isolation among these two populations will inform management strategies for this rare species

that balance the benefits of genetic diversity and local adaptation.

Data accessibility

Genomic and phenotypic data generated for this study, as well as scripts used for analysis are

available from Dryad: doi:10.5061/dryad.pc866t1xg.

Author's contributions

L.N.D.S. designed the study, contributed to data collection, generated, processed, and analyzed genomic data, interpreted the data, and led the writing of the manuscript. *A.M.* contributed to data collection, analyzed the climate data, interpreted the results, and led the writing of the manuscript. *J.W.W.* contributed to data collection and edited the manuscript. *J.A.H.* designed the study, contributed to data collection, interpreted the results, and wrote the manuscript. *L.N.D.S.* and *A.M.* contributed equally to the study.

Funding

This work was funded by USDA Forest Service Health Protection Gene Conservation Program and Western Wildlands Environmental Threat Assessment Center to JAH and JWW, Morton Arboretum Center for Tree Science Fellowship to JAH, a new faculty award from the office of North Dakota Experimental Program to Stimulate Competitive Research (ND-EPSCoR NSFIIA-1355466), the Schatz Center for Tree Molecular Genetics, and the NDSU Environmental and Conservation Sciences Graduate Program to LNDS and an NSF-PRFB (Award #2209410) to AM.

Acknowledgements

Tom Ledig, USDA-Forest Service, PSW, had the foresight to establish the common garden study at the end of his career. We thank him for sharing this project with us. For their help collecting needle tissue and measuring phenotypes within the common garden, the authors would like to thank Alexis Pearson, Tyler Stadel, Courtney Canning, and Madison Snider. The authors also would like to thank John Gabbert for allowing the scientific crew to access the common garden study. Finally, the authors would like to acknowledge the Michumash

References

694   **References**

695   Altschul, S. F., W. Gish, W. Miller, E. W. Myers, and D. J. Lipman. 1990. Basic local

696   alignment search tool. J. Mol. Biol. 215:403–410. Netherlands.

697   Altschul, S. F., T. L. Madden, A. A. Schäffer, J. Zhang, Z. Zhang, W. Miller, and D. J.

698   Lipman. 1997. Gapped BLAST and PSI-BLAST: a new generation of protein

699   database search programs. Nucleic Acids Res. 25:3389–3402. England.

700   Andrew, R. L., and L. H. Rieseberg. 2013. Divergence is focused on few genomic regions

701   early in speciation: Incipient speciation of sunflower ecotypes. Evolution 67:2468–

702   2482.

703   Axelrod, D. I. 1967. Geologic history of the Californian insular flora. Pp. 267–314 *in* 1st

704   Symposium on the Biology of the California Islands. National Park Service.

705   Bakdash, J. Z., and L. R. Marusich. 2022. rmcorr: Repeated Measures Correlation.

706   Barnard-Kubow, K. B., N. So, and L. F. Galloway. 2016. Cytonuclear incompatibility

707   contributes to the early stages of speciation. Evolution 70:2752–2766.

708   Bates, D., M. Mächler, B. Bolker, and S. Walker. 2015. Fitting linear mixed-effects models

709   using lme4. J. Stat. Softw. 67:1–48.

710   Benjamini, Y., and Y. Hochberg. 1995. Controlling the false discovery rate: a practical and

711   powerful approach to multiple testing. J. R. Stat. Soc. Ser. B Methodol. 57:289–300.

712   Wiley Online Library.

713   Bolte, C. E., and A. J. Eckert. 2020. Determining the when, where and how of conifer

714   speciation: a challenge arising from the study 'Evolutionary history of a relict conifer

715   Pseudotaxus chienii.' Ann. Bot. 125:v–vii.

716   Bolte, C. E., T. M. Faske, C. J. Friedline, and A. J. Eckert. 2022. Divergence amid recurring

717   gene flow: complex demographic histories for two North American pine species

718   (Pinus pungens and P. rigida) fit growing expectations among forest trees. Tree

Genet. Genomes 18:35.

Camacho, C., G. Coulouris, V. Avagyan, N. Ma, J. Papadopoulos, K. Bealer, and T. L.

Madden. 2009. BLAST+: architecture and applications. BMC Bioinformatics 10:421.

England.

Cannon, C. H., and R. J. Petit. 2020. The oak syngameon: more than the sum of its parts.

New Phytol. 226:978–983.

Case, A. L., F. R. Finseth, C. M. Barr, and L. Fishman. 2016. Selfish evolution of cytonuclear

hybrid incompatibility in Mimulus. Proc. R. Soc. B Biol. Sci. 283:20161493. Royal

Society.

Chen, S., Y. Zhou, Y. Chen, and J. Gu. 2018. fastp: an ultra-fast all-in-one FASTQ

preprocessor. Bioinformatics 34:i884–i890.

Choi, J. Y., M. Purugganan, and E. A. Stacy. 2020. Divergent Selection and Primary Gene

Flow Shape Incipient Speciation of a Riparian Tree on Hawaii Island. Mol. Biol.

Evol. 37:695–710.

Christie, K., L. S. Fraser, and D. B. Lowry. 2022. The strength of reproductive isolating

barriers in seed plants: Insights from studies quantifying premating and postmating

reproductive barriers over the past 15 years. Evolution 76:2228–2243.

Coughlan, J. M., and D. R. Matute. 2020. The importance of intrinsic postzygotic barriers

throughout the speciation process. Philos. Trans. R. Soc. B Biol. Sci. 375:20190533.

Royal Society.

De La Torre, A. R., A. Piot, B. Liu, B. Wilhite, M. Weiss, and I. Porth. 2020. Functional and

morphological evolution in gymnosperms: A portrait of implicated gene families.

Evol. Appl. 13:210–227.

DeSilva, R., and R. S. Dodd. 2020. Association of genetic and climatic variability in giant

sequoia, Sequoiadendron giganteum, reveals signatures of local adaptation along

moisture-related gradients. Ecol. Evol. 10:10619–10632.

Di Santo, L. N., S. Hoban, T. L. Parchman, J. W. Wright, and J. A. Hamilton. 2022. Reduced

representation sequencing to understand the evolutionary history of Torrey pine

(*Pinus torreyana* parry) with implications for rare species conservation. Mol. Ecol.

31:4622–4639.

Doyle, J. J., and J. L. Doyle. 1987. A rapid DNA isolation procedure for small quantities of

fresh leaf tissue. Phytochem. Bull. 19:11–15.

Eaton, D. A. R., and I. Overcast. 2020. ipyrad: Interactive assembly and analysis of RADseq

datasets. Bioinformatics 36:2592–2594. Oxford University Press.

Eckert, A. J., J. Van Heerwaarden, J. L. Wegrzyn, C. D. Nelson, J. Ross-Ibarra, S. C.

González-Martínez, and D. B. Neale. 2010. Patterns of Population Structure and

Environmental Associations to Aridity Across the Range of Loblolly Pine (Pinus

taeda L., Pinaceae). Genetics 185:969–982.

Farjon, A. 2013. IUCN Red List of Threatened Species: *Pinus torreyana*. International Union

for Conservation of Nature and Natural Resources.

Feder, J. L., S. P. Egan, and P. Nosil. 2012. The genomics of speciation-with-gene-flow.

Trends Genet. 28:342–350.

Fernando, D. D., S. M. Long, and R. A. Sniezko. 2005. Sexual reproduction and crossing

barriers in white pines: the case between Pinus lambertiana (sugar pine) and P.

monticola (western white pine). Tree Genet. Genomes 1:143–150.

Fick, S. E., and R. J. Hijmans. 2017. WorldClim 2: new 1-km spatial resolution climate

surfaces for global land areas. Int. J. Climatol. 37:4302–4315.

Goudet, J., and T. Jombart. 2021. hierfstat: Estimation and Tests of Hierarchical F-Statistics.

Granda, E., C. Scoffoni, A. E. Rubio-Casal, L. Sack, and F. Valladares. 2014. Leaf and stem

physiological responses to summer and winter extremes of woody species across

temperate ecosystems. Oikos 123:1281–1290.

Gugger, P. F., S. T. Fitz-Gibbon, A. Albarrán-Lara, J. W. Wright, and V. L. Sork. 2021.

Landscape genomics of *Quercus lobata* reveals genes involved in local climate

adaptation at multiple spatial scales. Mol. Ecol. 30:406–423.

Haller, J. R. 1967. Proceedings of the Symposium on the Biology of the California Islands.

Santa Barbara Botanic Garden.

Hamilton, J. A., R. Royauté, J. W. Wright, P. Hodgskiss, and T. F. Ledig. 2017. Genetic

conservation and management of the California endemic, Torrey pine (Pinus

torreyana Parry): Implications of genetic rescue in a genetically depauperate species.

Ecol. Evol. 7:7370–7381. Wiley Online Library.

Hendry, A. P. 2009. Ecological speciation! Or the lack thereof? Can. J. Fish. Aquat. Sci.

66:1383–1398. NRC Research Press.

Hendry, A. P., P. Nosil, and L. H. Rieseberg. 2007. The Speed of Ecological Speciation.

Funct. Ecol. 21:455–464. [British Ecological Society, Wiley].

Hijmans, R. J. 2024. terra: Spatial Data Analysis.

Janes, J. K., and J. A. Hamilton. 2017. Mixing It Up: The Role of Hybridization in Forest

Management and Conservation under Climate Change. Forests 8:237.

Multidisciplinary Digital Publishing Institute.

Jombart, T. 2008. adegenet: a R package for the multivariate analysis of genetic markers.

Bioinformatics 24:1403–1405. Oxford University Press.

Jombart, T., and I. Ahmed. 2011. adegenet 1.3-1: new tools for the analysis of genome-wide

SNP data. Bioinformatics 27:3070–3071. Oxford University Press.

Ledig, T. F., and T. M. Conkle. 1983. Gene diversity and genetic structure in a narrow

endemic, Torrey Pine (Pinus torreyana Parry ex Carr.). Evolution 37:79–85.

Lenth, R. V. 2023. emmeans: Estimated Marginal Means, aka Least-Squares Means.

Leroy, T., Q. Rougemont, J.-L. Dupouey, C. Bodénès, C. Lalanne, C. Belser, K. Labadie, G.

Le Provost, J.-M. Aury, A. Kremer, and C. Plomion. 2020. Massive postglacial gene

flow between European white oaks uncovered genes underlying species barriers. New

Phytol. 226:1183–1197.

Li, H. 2011. A statistical framework for SNP calling, mutation discovery, association

mapping and population genetical parameter estimation from sequencing data.

Bioinformatics 27:2987–2993.

Li, H. 2013. Aligning sequence reads, clone sequences and assembly contigs with BWA-

MEM. arXiv.

Li, H. 2014. Toward better understanding of artifacts in variant calling from high-coverage

samples. Bioinformatics 30:2843–2851. Oxford University Press.

Li, H., B. Handsaker, A. Wysoker, T. Fennell, J. Ruan, N. Homer, G. Marth, G. Abecasis, R.

Durbin, and P. S. 1000 G. P. Data. 2009. The Sequence Alignment/Map format and

SAMtools. Bioinformatics 25:2078–2079.

Lind, B. M., C. J. Friedline, J. L. Wegrzyn, P. E. Maloney, D. R. Vogler, D. B. Neale, and A.

J. Eckert. 2017. Water availability drives signatures of local adaptation in whitebark

pine (Pinus albicaulis Engelm.) across fine spatial scales of the Lake Tahoe Basin,

USA. Mol. Ecol. 26:3168–3185.

Lind, B. M., M. Menon, C. E. Bolte, T. M. Faske, and A. J. Eckert. 2018. The genomics of

local adaptation in trees: are we out of the woods yet? Tree Genet. Genomes 14:29.

Liu, Z.-X., H.-Y. Xiong, L.-Y. Li, and Y.-J. Fei. 2018. Functional Conservation of an

AGAMOUS Orthologous Gene Controlling Reproductive Organ Development in the

Gymnosperm Species Taxus chinensis var. mairei. J. Plant Biol. 61:50–59.

Lowry, D. B., J. L. Modliszewski, K. M. Wright, C. A. Wu, and J. H. Willis. 2008a. The

strength and genetic basis of reproductive isolating barriers in flowering plants.

Philos. Trans. R. Soc. B Biol. Sci. 363:3009–3021. Royal Society.

Lowry, D. B., R. C. Rockwood, and J. H. Willis. 2008b. ECOLOGICAL REPRODUCTIVE

ISOLATION OF COAST AND INLAND RACES OF MIMULUS GUTTATUS.

Evolution 62:2196–2214.

MacLachlan, I. R., T. K. McDonald, B. M. Lind, L. H. Rieseberg, S. Yeaman, and S. N.

Aitken. 2021. Genome-wide shifts in climate-related variation underpin responses to

selective breeding in a widespread conifer. Proc. Natl. Acad. Sci. 118. National

Academy of Sciences.

McWilliam, J. R. 1959. Interspecific Incompatibility in Pinus. Am. J. Bot. 46:425–433.

Melo, M. C., A. Grealy, B. Brittain, G. M. Walter, and D. Ortiz-Barrientos. 2014. Strong

extrinsic reproductive isolation between parapatric populations of an Australian

groundsel. New Phytol. 203:323–334.

Menon, M., J. C. Bagley, C. J. Friedline, A. V. Whipple, A. W. Schoettle, A. Leal-Sàenz, C.

Wehenkel, F. Molina-Freaner, L. Flores-Rentería, M. S. Gonzalez-Elizondo, R. A.

Sniezko, S. A. Cushman, K. M. Waring, and A. J. Eckert. 2018. The role of

hybridization during ecological divergence of southwestern white pine (Pinus

strobiformis) and limber pine (P. flexilis). Mol. Ecol. 27:1245–1260.

Mogensen, H. L. 1996. INVITED SPECIAL PAPER: The hows and whys of cytoplasmic

inheritance in seed plants. Am. J. Bot. 83:383–404.

Moyroud, E., M. Monniaux, E. Thévenon, R. Dumas, C. P. Scutt, M. W. Frohlich, and F.

Parcy. 2017. A link between LEAFY and B-gene homologues in Welwitschia

mirabilis sheds light on ancestral mechanisms prefiguring floral development. New

Phytol. 216:469–481.

Nardini, A., M. A. Lo Gullo, P. Trifilò, and S. Salleo. 2014. The challenge of the

Mediterranean climate to plant hydraulics: Responses and adaptations. Environ. Exp.

Bot. 103:68–79.

Neale, D. B., and R. R. Sederoff. 1989. Paternal inheritance of chloroplast DNA and maternal

inheritance of mitochondrial DNA in loblolly pine. Theor. Appl. Genet. 77:212–216.

Nosil, P., D. J. Funk, and D. Ortiz-Barrientos. 2009a. Divergent selection and heterogeneous

genomic divergence. Mol. Ecol. 18:375–402.

Nosil, P., L. J. Harmon, and O. Seehausen. 2009b. Ecological explanations for (incomplete)

speciation. Trends Ecol. Evol. 24:145–156. Elsevier.

Petit, R. J., and A. Hampe. 2006. Some evolutionary consequences of being a tree. Annu.

Rev. Ecol. Evol. Syst. 37:187–214.

Puritz, J. B., C. M. Hollenbeck, and J. R. Gold. 2014a. dDocent: a RADseq, variant-calling

pipeline designed for population genomics of non-model organisms. PeerJ 2:e431–

e431. PeerJ Inc.

Puritz, J. B., M. V. Matz, R. J. Toonen, J. N. Weber, D. I. Bolnick, and C. E. Bird. 2014b.

Demystifying the RAD fad. Mol. Ecol. 23:5937–5942. Wiley Online Library.

R Core Team. 2023. R: A Language and Environment for Statistical Computing. R

Foundation for Statistical Computing, Vienna, Austria.

Richards, T. J., and D. Ortiz-Barrientos. 2016. Immigrant inviability produces a strong barrier

to gene flow between parapatric ecotypes of Senecio lautus. Evolution 70:1239–1248.

Rieseberg, L. H., and B. K. Blackman. 2010. Speciation genes in plants. Ann. Bot. 106:439–

455.

Riordan, E. C., P. F. Gugger, J. Ortego, C. Smith, K. Gaddis, P. Thompson, and V. L. Sork.

2016. Association of genetic and phenotypic variability with geography and climate in

three southern California oaks. Am. J. Bot. 103:73–85.

Rönnegård, L., S. E. McFarlane, A. Husby, T. Kawakami, H. Ellegren, and A. Qvarnström.

2016. Increasing the power of genome wide association studies in natural populations

using repeated measures – evaluation and implementation. Methods Ecol. Evol.

7:792–799.

Rose, N. H., R. A. Bay, M. K. Morikawa, and S. R. Palumbi. 2018. Polygenic evolution

drives species divergence and climate adaptation in corals. Evolution 72:82–94.

Savolainen, O., T. Pyhäjärvi, and T. Knürr. 2007. Gene Flow and Local Adaptation in Trees.

Annu. Rev. Ecol. Evol. Syst. 38:595–619.

Schluter, D. 2001. Ecology and the origin of species. Trends Ecol. Evol. 16:372–380.

Schluter, D. 2009. Evidence for Ecological Speciation and Its Alternative. Science 323:737–

741. American Association for the Advancement of Science.

Schluter, D., and L. H. Rieseberg. 2022. Three problems in the genetics of speciation by

selection. Proc. Natl. Acad. Sci. 119:e2122153119. Proceedings of the National

Academy of Sciences.

Strasburg, J. L., N. A. Sherman, K. M. Wright, L. C. Moyle, J. H. Willis, and L. H.

Rieseberg. 2012. What can patterns of differentiation across plant genomes tell us

about adaptation and speciation�? 364–373.

Tallmon, D. A., G. Luikart, and R. S. Waples. 2004. The alluring simplicity and complex

reality of genetic rescue. Trends Ecol. Evol. 19:489–496. Elsevier.

Tiffin, P., S. Olson, and L. C. Moyle. 2001. Asymmetrical crossing barriers in angiosperms.

Proc. R. Soc. Lond. B Biol. Sci. 268:861–867. Royal Society.

Turelli, M., and L. C. Moyle. 2007. Asymmetric Postmating Isolation: Darwin's Corollary to

Haldane's Rule. Genetics 176:1059–1088.

van de Kerk, M., D. P. Onorato, J. A. Hostetler, B. M. Bolker, and M. K. Oli. 2019.

Dynamics, Persistence, and Genetic Management of the Endangered Florida Panther

Population. Wildl. Monogr. 203:3–35.

Walter, G. M., T. J. Richards, M. J. Wilkinson, M. W. Blows, J. D. Aguirre, and D. Ortiz-

Barrientos. 2020. Loss of ecologically important genetic variation in late generation

hybrids reveals links between adaptation and speciation. Evol. Lett. 4:302–316.

Wei, T., and V. Simko. 2021. R package "corrplot": Visualization of Correlation Matrix.

Williams, A. P., C. J. Still, D. T. Fischer, and S. W. Leavitt. 2008. The influence of

summertime fog and overcast clouds on the growth of a coastal Californian pine: a

tree-ring study. Oecologia 156:601–611.

Williams, C. G., and O. Savolainen. 1996. Inbreeding Depression in Conifers: Implications

for Breeding Strategy. For. Sci. 42:102–117.

Wright, K. M., D. Lloyd, D. B. Lowry, M. R. Macnair, and J. H. Willis. 2013. Indirect

Evolution of Hybrid Lethality Due to Linkage with Selected Locus in Mimulus

guttatus. PLOS Biol. 11:e1001497. Public Library of Science.

Yeaman, S., K. A. Hodgins, K. E. Lotterhos, H. Suren, S. Nadeau, J. C. Degner, K. A.

Nurkowski, P. Smets, T. Wang, L. K. Gray, K. J. Liepe, A. Hamann, J. A. Holliday,

M. C. Whitlock, L. H. Rieseberg, and S. N. Aitken. 2016. Convergent local adaptation

to climate in distantly related conifers. Science 353:1431–1433.

Tables

Table 1. SNPs significantly associated with one or more phenotypes (FDR < 0.1), including
tree height (cm), number of conelets, number of immature cones, and number of mature
cones. Listed are generic SNP IDs given for this study, SNP IDs retrieved from *Pinus taeda*
draft genome (scaffold ID:position on the scaffold), the trait(s) each SNP is associated with,
the effect of increasing the number of derived alleles on the associated phenotype(s), and the
FDR of the genotype-phenotype association.
916

Generic SNP ID	SNP ID within *P. taeda* genome	Phenotypic trait association	Phenotypic effect	FDR
locus_4218	APFE030529380.1:392010	Tree height (cm)	+	0.047
		Nb conelets	+	<0.001
		Nb immature cones	+	0.085
		Nb mature cones	+	0.037
locus_12498	APFE031619559.1:67487	Tree height (cm)	+	0.047
locus_433	APFE030047412.1:104020	Nb conelets	+	0.078
		Nb mature cones	+	0.057
locus_3004	APFE030371338.1:302521	Nb conelets	+	0.078
locus_3676	APFE030458498.1:93076	Nb conelets	+	0.024
locus_9630	APFE031249435.1:61968	Nb conelets	+	0.011
		Nb immature cones	+	0.085
		Nb mature cones	+	0.027
locus_13093	APFE031688147.1:172170	Nb conelets	+	0.072
		Nb mature cones	+	0.027
locus_1164	APFE030138697.1:151998	Nb mature cones	+	0.032
locus_6624	APFE030828725.1:2769	Nb mature cones	+	0.027
locus_8019	APFE031020621.1:50406	Nb mature cones	+	0.053
locus_8778	APFE031129984.1:16200	Nb mature cones	-	0.057
locus_10668	APFE031380592.1:30882	Nb mature cones	+	0.053

918 **Figures**

919
Figure 1. Haldane-Anscombe-corrected log-transformed odd ratios $\pm$ standard error (x axis)
for significantly (FDR < 0.1) enriched (positive values) and depleted (negative values) GO
terms (y axis) for the set of alleles with reduced F1 heterozygosity, including cellular
component (CC, black), molecular function (MF, dark gray), and biological process (BP,
light gray) terms. Here, we present Haldane-Anscombe-corrected estimates of log-transform
odd ratios as the presence of zero success probabilities within the data would otherwise lead
to nonsensical estimates of odd ratios (see Appendix S9 for details).

927
Figure 2. Expected Marginal Means (EMM) ± standard error of all four phenotypes
investigated (y axis) given the genotype (x axis) of island, mainland, and hybrid individuals
at locus_4218. Phenotypes measured include tree height (cm), number of conelets, number of
immature cones, and number of mature cones. Sample sizes are 47, 5, and 123 individuals
with genotype 0/0, 0/1, and 1/1, respectively. Distinct bolded lowercase letters indicate
significant differences in phenotypes (adjusted $P < 0.05$) among genotypes (see Appendix
S11 for details). Based on genotype frequencies (see Results), 0 was defined as the island
allele, and 1 as the mainland allele.

937
Figure 3. Distribution of observed heterozygosity estimates across ancestries (F1 hybrid,
Island, and Mainland) for three SNP sets (All, GWAS, and Low). Distinct lowercase letters
indicate significant difference in observed heterozygosities (FDR < 0.1) among ancestries
(see Appendix S12 for details). All: data set containing all retained SNPs after filtering
(11,379 SNPs across 209 individuals), GWAS: data set containing SNPs exhibiting a
significant genotype-phenotype association (12 SNPs across 209 individuals), Low; data sets
containing SNPs exhibiting reduced heterozygosity relative to expectations (185 SNPs across
209 individuals).