Molecular fingerprints of cell size sensing and mating type differentiation in pennate diatoms

Darja Belišová*[1], Gust Bilcke*[1,2,3], Sien Audoor[1,a], Sofie D'hondt[1], Lieven De Veylder[2,3], Klaas Vandepoele[2,3,4], Wim Vyverman[1†]

[1]Protistology and Aquatic Ecology, Department of Biology, Ghent University, 9000 Ghent, Belgium

[2]Department of Plant Biotechnology and Bioinformatics, Ghent University, 9052 Ghent, Belgium

[3]VIB Center for Plant Systems Biology, 9052 Ghent, Belgium

[4]VIB Center for AI & Computational Biology, VIB, Ghent, Belgium

[a]Present address: Institut de Biologie Physico-Chimique, Laboratory of Chloroplast Biology and Light Sensing in Microalgae, UMR7141 Centre National de la Recherche Scientifique (CNRS), Sorbonne Université, 75005 Paris, France

[†]Corresponding author - wim.vyverman@ugent.be

*These authors contributed equally to this work

The authors declare that they have no conflict of interest.

Summary

- A unique cell size sensing mechanism is at the heart of the life cycle of diatoms. During population growth, cell size decreases until a Sexual Size Threshold (SST) is reached, below which cells become sexually competent. In most pennate diatoms, the two mating types undergo biochemical and behavioral differentiation below the SST, although the molecular pathways underlying their size-dependent maturation remain unknown.

- Here, we developed a method to shorten the generation time of *Cylindrotheca closterium* through single-cell microsurgery, enabling the transcriptomic comparison of genetically identical large and undifferentiated cells with small, sexually competent cells for six different genotypes.

- We identified 21 genes upregulated in small cells regardless of their mating type, revealing how cells undergo specific transcriptional reprogramming when passing the SST. Furthermore, we revealed a size-regulated gene cluster with three mating type-specific genes susceptible to sex-inducing pheromones. In addition, comparative transcriptomics confirmed the shared mating type specificity of Mating-type Related Minus 2 homologs in three pennate diatoms, suggesting them to be part of a conserved partner-recognition mechanism.

- This study sheds light on how diatoms acquire sexual competence in a strictly size-dependent manner, revealing a complex machinery underlying size-dependent maturation, mating behavior, and heterothally in pennate diatoms.

Introduction

Diatoms are an extremely successful group of unicellular primary producers in freshwater and marine environments (Field *et al.*, 1998). Despite their extraordinary species diversity underlying the colonization of widely different habitats spanning terrestrial to aquatic realms, all diatoms share a unique silica cell wall consisting of two halves (thecae) slightly differing in size. During cell division, the two daughter cells each inherit one of the rigid parental thecae that are dissimilar in size and fit as a "lid on a box" (**Figure 1**). As a result, an asexually growing population of cells will gradually decrease in mean cell size, a phenomenon known as the MacDonald-Pfitzer rule (Pfitzer, 1869; Macdonald, 1869). Different mechanisms of cell size restoration mechanisms have been described (Nagai *et al.*, 1995; Chepurnov *et al.*, 2004; Kaczmarska *et al.*, 2022); but most commonly this happens via the sexual phase during which diploid vegetative cells form haploid gametes. The ability to engage in sexual reproduction is strictly cell-size controlled and triggered once cells drop below a species-specific Sexual Size Threshold (SST). The cellular mechanisms and molecular changes associated with this unique ability of diatom cells to sexually mature by measuring their own cell size are still completely unknown.

In the generally homothallic centric diatoms, cells produce oogonia and sperm cells, while in raphid pennate diatoms vegetative cells become responsive to sexual cues after passing the SST and transform into morphologically identical but functionally differentiated gametangial cells (Moeys *et al.*, 2016; Basu *et al.*, 2017; Bilcke *et al.*, 2021; Klapper *et al.*, 2021). Recent advances revealed the complex nature of diatom sexual behavior, including the identification of multiple sex pheromones involved in the mating process (Gillard *et al.*, 2013; Moeys *et al.*, 2016; Klapper *et al.*, 2021, 2023).

Most raphid pennate diatoms have a heterothallic mating system with two mating types (MT+ and MT-) (reviewed in (Chepurnov *et al.*, 2004; Davidovich *et al.*, 2015; Poulíčková *et al.*,

2019; Bilcke *et al.*, 2022)). Although the elucidation of the diploid sex-determination system

in heterothallic pennate diatoms is still in its infancy, crossing experiments have revealed that

one of the two mating types is typically heterogametic, i.e. produces different gametes that

determine the mating type of the offspring (Davidovich 2002, Vanstechelman *et al.*, 2013).

Additionally, recent studies in *Seminavis robusta* and *Pseudo-nitzschia multistriata* have

demonstrated the existence of a single sex-determining locus that drives the expression of

downstream mating type related genes (Vanstechelman *et al.*, 2013; Russo *et al.*, 2018).

Specifically, the sex-determining A allele of the *P. multistriata* Mating-type Related Plus 3

(*MRP3*) gene segregates with MT+ strains (Russo *et al.*, 2018). A simple control mechanism,

where differential methylation of the sex locus leads to mono-allelic expression of *MRP3* only

in MT+, has been described by Ruggiero et al. (2023). MRP3 acts as a master regulator,

activating downstream MT+ genes *MRP1* and *MRP2*, while inactivating the MT- specific

genes *MRM1* and *MRM2*. Importantly, despite the widespread occurrence of heterothally in

pennate diatoms, no homologs of this sex determining system have yet been identified in

other pennate species outside the *Pseudo-nitzschia - Fragilariopsis* clade (Russo *et al.*, 2018).

Figure 1: **A typical life cycle of a raphid pennate diatom**, illustrated with the species *C. closterium*. (**1**) Cells divide vegetatively and the mean size in the population drops below the sexual size threshold (SST) (**2**). Small cells become sexually mature and seek a compatible mate (**3**). In *C. closterium*, mating type minus (MT-) produces pheromone (SIP-, blue dots) while mating type plus (MT+) evinces oriented motility towards the partner (**3,4**). (**5**) Each gametangial cell produces two gametes. (**6**) Gametes fuse and form zygotes (**7**). Zygotes expand and become auxospores (**8**), and finally, an initial cell (**9**) starts a new generation.

Here, we studied size sensing and mating type differentiation in the raphid pennate diatom *Cylindrotheca closterium,* a common and globally widespread member of marine, brackish water phytoplankton and microphytobenthos (Audoor *et al.*, 2024). We developed a protocol

to significantly shorten generation time by mechanically reducing cell length, allowing us to

compare gene expression of cells above and below the SST for the same genotypes. We

carried out differential gene expression analysis using robust transcriptome profiling of six

different genotypes, encompassing both mating types in both size variants. Phylogenetic

analysis and cross-species transcriptomic analysis were used to reveal conserved size- and

mating type-specific genes.

Materials & Methods

Culture conditions for *C. closterium* and *S. robusta* cells

Cultures of *C. closterium* and *S. robusta* were grown in artificial seawater (ASW) composed

of Tropic Marin BIO-ACTIF sea salt (34.5 g/l; Tropic Marin, Wartenberg, Germany) and

sodium bicarbonate (0.08 g/l) supplemented with Guillard's F/2 Marine Water enrichment

solution (Sigma-Aldrich, Belgium) and an antibiotic mixture (500 mg/l penicillin, 500 mg/l

ampicillin, 100 mg/l streptomycin and 50 mg/l gentamicin). Cells were sustained under

constant illumination (25 µmol/m2/s) at 18°C, if not stated otherwise. An initial set of strains

was provided by the Belgian Coordinated Collection of Microorganisms (BCCM/DCG)

(https://bccm.belspo.be/about-us/bccm-dcg) (**Supplementary Table 1**).

Experimental size manipulation and recovery of *C. closterium*

Four parental *C. closterium* strains of two different mating types were crossed to create

progeny strains. More information about the parental strains can be found in **Supplementary**

Methods. Single cells were isolated to start monoclonal progeny strains and subsequently

their subpopulations were reduced in size using a novel microsurgery approach. Particularly,

long apices (cell protrusions) of single cells were cut off using an extremely thin tip of a glass

Pasteur pipette, with an open diameter ranging from 50 to 200 µm (**Supplementary Figure**

1). Prior to the experiment, pipettes used for cutting and isolation were manually thinned

using a Bunsen Burner and clipped by a tweezer to create open and sharp tips

(**Supplementary Video**). In healthy and unstressed cultures, ~50-70% of cells recovered

successfully from the cutting procedure, forming new monoclonal, smaller-celled populations.

Such manipulated single cells were immediately isolated in a 96 well-plate (Greiner

CELLSTAR®, Sigma Aldrich Belgium) and were then maintained two weeks in low light

conditions (10 – 15 µmol photons/m2/s) for wounding recovery, giving rise to new

monoclonal populations that were genetically identical to their larger counterparts but with a

smaller cell size. After this immediate recovery period, cultures were maintained in the

growth phase for at least three months to ensure their complete recovery. An identical period

of vegetative growth was enabled for large non-manipulated paired cultures. During this

extended period of growth, cultures were grown in 24-well plates under 12/12 day/night

cycles. Once a week, each culture was scraped from the bottom of the well and 15 µl of a

dispersed culture was inoculated into 1 ml of fresh medium to ensure that cultures remained in

exponential growth phase. Finally, all cultures were transferred to small culture flasks (25

cm^2, VWR ®) to be grown under continuous light for at least four weeks to acclimate before

harvesting for the RNA-seq experiment.

Determination of mating types and selection of genotypes for RNA-sequencing

Before selecting *C. closterium* cultures for the RNA-seq experiment, the average length of

cells (the distance between the tip of each apex, **Supplementary Figure 2**) was determined

and the mating type was identified by backwards crossing of each culture with both parental

genotypes below the SST (Audoor *et al*., 2024). Specifically, synchronized cultures were dark

arrested for 36 hours followed by mixing various combinations of two strains in a 96 well-

plate with a cell density of 40 cells/µl per strain in a total volume of 200 µl. The potential

mating of strains was microscopically inspected after 12, 24 and 48 hours. The

experimentally determined mating types are indicated in **Supplementary Table 1**. Backwards

crosses were performed for both the small (after microsurgery) and large (original) variants to

verify whether small cells can form all sexual cell stages (mating pairs, auxospores, initial

cells), while large cells are incapable of sexual reproduction.

Altogether, six *C. closterium* genotypes were selected for RNA-seq of their matched large and

small size classes. We picked three MT+ genotypes: C2 (86 µm large cells vs 34 µm small

cells), MC4 (68 µm vs 29 µm) and, MC5 (80 µm vs 26 µm); and three MT- genotypes: – A5

(81 µm vs 47 µm), A6 (67 µm vs 37 µm) and, MC3 (74 µm vs 45 µm) (**Supplementary**

Table 1). For *S. robusta*, six strains below the SST, three of each mating type, were selected

for additional RNA-seq (**Supplementary Table 1**) to complement existing datasets (Osuna-

Cruz *et al.*, 2020).

RNA extraction and sequencing

Exponentially growing cultures of *C. closterium* and *S. robusta* were harvested on membrane

filters (3µm, 25mm, Versapor®, Pall Laboratory, VWRTM) and RNA was extracted using a

Qiagen RNeasy extraction kit with optimizations. Detailed protocols can be found in

Supplementary Methods.

Samples with the highest RNA integrity number and sufficient quantity were processed by

TruSeq Stranded mRNA Library Prep with polyA selection. The library was further subjected

to 2x150bp paired-end sequencing using the Illumina NovaSeq 6000 instrument at VIB

Nucleomics Core (Leuven, Belgium www.nucleomics.be). On average, 33.8 ± 4.4 million and

32.0 ± 3.7 million reads per sample were sequenced for *C. closterium* and *S. robusta*

respectively.

Differential expression analysis and functional annotation

Quality trimmed paired-end reads (as explained in **Supplementary Methods**) were mapped to transcripts belonging to the gene annotation model v1.2 of the *C. closterium* reference genome (Audoor *et al.*, 2024) using Salmon v1.3 (Patro *et al.*, 2017). Read counts were imported into R with tximport v1.22.0 (Soneson *et al.*, 2016), followed by filtering, normalization and generalized linear model fitting as detailed in the **Supplementary Methods**. For *C. closterium*, we defined four contrasts to execute a differential expression (DE) analysis with EdgeR v3.36.0 (Robinson *et al.*, 2009) (**Figure 2**): (1) cell sizes above and below the SST irrespective of the MT, (2) cells of MT+ versus MT- irrespective of the size, (3) cells of MT+ below the SST versus all other samples, and (4) cells of MT- below the SST versus all other samples. For the "Size" contrast (1), an additional paired analysis was performed, controlling for the variable "genotype" in the generalized linear model. Next, likelihood ratio tests were performed using EdgeR, and p-values were controlled at the 5 % false discovery rate (FDR) level using the Benjamini-Hochberg correction. The analysis of existing (Osuna-Cruz *et al.*, 2020) and newly generated RNA-seq data of *S. robusta* used for comparative purposes is explained in **Supplementary Methods**. In addition, differential expression calls during the different stages of sexual reproduction of *C. closterium* and *S. robusta* were retrieved from (Audoor *et al.*,2024).

Protein domain and gene family annotation of *C. closterium* genes was performed using TRAPID 2.0 (Bucchini *et al.*, 2021), with default parameters (E-value threshold 10^{-5}) and PLAZA Diatoms as a database for similarity searches. Meanwhile, InterPro and gene family functional annotation for *S. robusta* was retrieved from PLAZA Diatoms 1.0 (Osuna-Cruz *et al.*, 2020). Additionally, computational predictions of subcellular localization, phylostratigraphic gene age as well as enrichment analyses were performed, as detailed in the **Supplementary Methods**.

Reverse transcriptase quantitative PCR (RT-qPCR) analysis of selected candidate genes

194 An RT-qPCR experiment was performed on six genes (Ccl_4437, Ccl_5484, Ccl_13556, Ccl_14859, Ccl_14861 and Ccl_16260) to validate results of the size-vs-mating type RNA-seq experiment. An identical set-up as for the RNA-seq experiment was used to prepare the cDNA of four strains (A5, C2, MC3, MC5), two biological replicates for each mating type, in two size variants above and below the SST.

200 Details about cDNA preparation, primers (**Supplementary Table 2**), cycling conditions and statistical analysis of the results can be found in **Supplementary Methods**.

Transcriptomic response to sex-inducing pheromone minus (SIP-)

204 Further, we performed an RNA-seq experiment to specifically asses the response of three MT+ specific genes (Ccl_14859, Ccl_14861 and Ccl_14874 to sex-inducing pheromone of the opposing mating type. Prior to the experiment SIP containing medium from MT- strain CA1.15 (MT-, DCG 0923) was harvested as described previously (Klapper *et al.*, 2021). The activity of the filtered medium was tested by its ability to induce a partial cell cycle arrest after 6 h of treatment in two biological replicates of MT+ strain ZW2.20 using an Amnis ImageStreamX Mk II imaging Flow Cytometer (Luminex), following the protocol of Audoor *et al.*, 2024 (**Supplementary Figure 3**). Next, cultures of MT+ strain ZW2.20 (MT+, DCG 0922) growing in the mid-exponential phase were placed in the dark for 24 h. Two hours before re-illumination, the supernatant medium of ZW2.20 was replaced with either filtered medium from its mating partner CA1.15 (treatment) or fresh culture medium (control). Triplicate ZW2.20 cultures were harvested at two time points (3 h, 9 h) after re-illumination for both control and SIP treatment conditions. Culture harvesting and RNA

extraction was performed as explained above. Library preparation (including poly-A

enrichment) and paired-end Illumina 150 bp mRNA sequencing was performed at Novogene

(UK) Co., Ltd. Details about quality trimming, mapping and differential expression analysis

of the results can be found in **Supplementary Methods**.

Phylogenetic analysis of MRM2 homologs

A hidden Markov model (HMM) was constructed to define the MRM2 gene family

(HOM02SEM009752) and identify more distant homologs. A protein sequence alignment

created with MAFFT v7.453 (Katoh & Standley, 2013) was trimmed to retain only the

conserved LRR domain, after which a profile HMM was created using the hmmbuild

command of HMMER (hmmer.org). The profile HMM is included in **Supplementary Data**

as a reference for the MRM2 family. Next, a targeted search for MRM2 homologs in

eukaryotic genomes was performed with the hmmsearch command and the top-50 proteins

with the highest domain-specific E-value score were selected for phylogenetic analysis. A

detailed pipeline can be found in **Supplementary Methods**.

Data availability

New RNA-seq libraries generated for *C. closterium* and *S. robusta* are available at ENA with

project number PRJEB52586 (size and mating type) and PRJEB81513 (SIP response).

Differential expression calls for both *C. closterium* and *S. robusta* as well as gene family

predictions for *C. closterium* , curated gene models of MRM2 family proteins in selected

pennate diatoms, and the MRM2 family HMM model are available as **Supplementary Data**.

Results & Discussion

Transcriptome profiling reveals size- and mating type-dependent gene expression

We profiled gene expression of large- and small-sized cells for three independent *C. closterium* cultures of each mating type. To identify only genes that show a robust response across strains, three different genotypes of MT+ (C2, MC4, MC5) and MT- (A5, A6, MC3) each were used as biological replicates. Only genotypes that belong to *C. closterium* clade II were selected, to avoid sequence divergence affecting the mapping to the reference genome (CA1.15, clade II) (Stock *et al.*, 2019; Audoor *et al.*, 2024).Small cells were generated by microsurgery of the apices of large cells, shortening the time required for cells to become sexually inducible (**Supplementary Figure 2**). Indeed, we estimate that more than one year of optimal growth (with a daily size decrease of ~0.11 µm) would be necessary for cells with an initial cell size of around 100 µm to naturally drop below the SST, which we determined to occur at a cell size between 55 - 60 µm, unlike the previously reported 66µm (Vanormelingen *et al.*, 2013). (**Figure 2A**). Using the microsurgery technique, we obtained matched cultures for each genotype in two size variants, before and after size manipulation. This eventually resulted in paired cultures with sizes on average 76±7µm (>SST) and 36±8µm (<SST) that were harvested for RNA-seq (**Figure 2B**). All cultures were tested for their ability to engage in sexual reproduction prior to the experiment. In none of the large-celled cultures, any stage of sexual reproduction (paired cells, auxospores, initial cells) could be identified after mixing compatible mating types, while all sexual stages were observed in all cultures whose cell size was experimentally decreased.

To desynchronize the cell cycle and thus prevent the enrichment of a particular cycle phase cultures were maintained in constant light regime for at least four weeks. RNA-seq reads of all samples were mapped to the reference genome of *C. closterium* strain CA1.15 (MT-) (Audoor *et al.*, 2024). The average mapping rate was 62.5 %, with no significant differences

in mapping rate between large (63.1±8.0%) and small cells (61.9±4.9%) or between MT+

(62.2±7.0%) and MT- (62.9±6.4%) (p>0.05, two-sample t-test). Multidimensional scaling

showed clustering of samples by parental genotype (A5, A6, C2 offspring of CA1.15xZW2.20

versus MC3, MC4, MC5, offspring of NH3.13xWS3.7) and not by condition (size or mating

type), in agreement with recent observations of significant genotype-specific differences in

expression patterns of the centric diatom *Skeletonema marinoi* (Pinseel *et al.*, 2022)

(**Supplementary Figure 4**).

Differential expression (DE) analysis was performed over four contrasts to detect genes

responsive to cell size, mating type, or the interaction of cell size and mating type (**Figure**

2C). Overall, a set of just 112 out of 24.633 annotated genes was identified as differentially

expressed in one or more of these contrasts (**Supplementary Data**). Given that these DE

genes show a conserved expression pattern throughout three independent genotypes for each

mating type, we expect a significant role for these genes in the size- and mating type-related

response in *C. closterium*. The small number of DE genes suggests that transcriptomic

differentiation between cell size classes and mating types in diatoms is limited to specific

traits related to size sensing and behavioral sexual dimorphism, while the overall metabolism

of cells remained unchanged. RT-qPCR was performed on independent cultures belonging to

both mating types and size classes for validation, confirming the mating type- and size-

dependent expression of five genes (Ccl_14859, Ccl_14861, Ccl_16260, Ccl_4437 and

Ccl_13556) while failing to find a response for Ccl_5484 (**Supplementary Figure 5**).

Whereas 21 genes were upregulated in small cells regardless of their MT, no genes were

upregulated in large cells. The cellular changes that take place in both mating types when cells

pass the SST thus appears to be mainly driven by the activation of genes below rather than by

suppression of genes expressed above the SST. ▯We extended the search for size-related

genes by performing a paired analysis in which the effects of genotype were subtracted for

each corresponding large-vs-small pair (**Supplementary Data**). In this way, we identified a

larger set of 182 differential expressed genes, comprising 169 genes upregulated in small

cells, and 13 genes that were biased towards large cells.

A number of genes was differentially expressed between mating types regardless of their size:

8 and 54 genes were induced in MT- and MT+, respectively (**Figure 2D**). Hence, the two

mating types can be distinguished by their unique gene expression profiles throughout their

entire life cycle, i.e. across large and small stages. Once cells decrease below the SST, an

additional activation of mating type specific genes occurs. Indeed, 29 genes were upregulated

in cells of a specific mating type but only below the SST, with a marked predominance of

MT+ specific genes (26 versus only 3 MT- specific) (**Figure 2D**). Hence, size-dependent gene

expression changes also reflect the physiological and behavioral differentiation between

mating types, even in the absence of a compatible mating partner.

Figure 2: Illustration of the microsurgery-based set-up for size manipulation of *C. closterium*, experimental design of RNA sequencing and contrasts tested in the differential expression analyses. (**A**) Large single cells were isolated to form monoclonal vegetative cultures. These were manipulated in size to drop below the sexual size threshold (SST) and significantly reduce life cycle duration. In such a manner cells above and below SST of the same genotype were cultured simultaneously and harvested for RNA-seq. Cells are coloured to represent the different mating types and size classes, with the line between two shades representing the raphe. (**B**) Cell lengths at the time of harvesting for RNA-seq experiment. (**C**) Transcriptomic differences between mating types, size classes and their interaction were studied by differential

expression analysis over four different contrasts. (**D**) Bar plots showing the number of

differentially expressed *C. closterium* genes for each contrast, each time subdivided in genes

upregulated in group A vs group B, and genes upregulated in group B vs group A.

Regulatory genes and retrotransposon activity mark small and MT+ cells

respectively

For 72 of the differentially expressed genes, InterPro protein domains and/or PLAZA Diatoms

gene families could be assigned, whereas 40 genes in our dataset remained without functional

annotation (**Supplementary Data**). Enrichment tests were performed to identify InterPro

domains and gene families that were significantly more common among the DE genes

compared to the background of all expressed genes. We detected eight gene families and five

InterPro domains that were significantly enriched among the DE genes for specific contrasts

(**Figure 3**). Moreover, one additional gene family was enriched in the set of all DE genes.

Genes upregulated in small cells of both mating types appear to be dominated by regulatory

functions. These include a family encoding DNA-binding helicase-like proteins

(HOM02SEM000018) (**Figure 3**). Several other genes that interact with DNA or RNA were

upregulated in small cells: a nuclease (Ccl_23878), a 5'-Nucleotidase (Ccl_12722) and a

ribonuclease H (Ccl_21492). Furthermore, five genes upregulated in cells below the SST

contain a potential nuclear localization signal, among which two genes from the family of

unknown function HOM02SEM018909. The identification of putative regulatory genes that

are transcribed in both mating types below the SST is especially relevant in light of recent

reports that the sex-determining gene *MRP3* of *P. multistriata* and a methyltransferase

situated nearby are differentially methylated between mating types (Ruggiero *et al.*, 2023)*.* A

general size-responsive transcription factor is thought to preferentially activate the non-

methylated sex-determining allele, causing mating type differentiation below the SST

(Ruggiero *et al.*, 2023). Hence, the nuclear and DNA-binding genes identified here are prime

candidates as size-dependent molecular switches in diatoms.

Two families of proteins containing reverse transcriptase and DNA integrase domains,

indicating a retrotransposon origin, were significantly enriched among MT+ biased genes:

HOM02SEM000149 and HOM02SEM000048 (**Figure 3**). Indeed, reverse transcriptase and

integrase InterPro domains were significantly more abundant among MT+ biased genes than

in the rest of the expressed transcriptome (**Figure 3**). The role of retrotransposons in the

evolution of mating type identity may be similar to mechanisms observed in other eukaryotic

groups where the mating type locus is protected through suppression of recombination caused

by retrotransposon accumulation (Bachtrog, 2006; Kent *et al.*, 2017). Alternatively, an overall

higher activity of (former) transposable elements in MT+ may be the result of differences in

chromatin accessibility between mating types similar to the above-mentioned differential

methylation in *P. multistriata* and as is the case in sex chromosomes of higher animals and

plants (Slotkin & Martienssen, 2007; Hobza *et al.*, 2017), or may be the result of random

differences in transposon activity between selected genotypes for each mating type.

Figure 3: Enrichment of gene families and protein domains among differentially expressed (DE) genes from different contrasts (x-axis). Features that are significantly enriched in a contrast are shown by circles, whose size scales with the significance level (-log10). Pie charts show the taxonomic composition of gene families. MT: mating type, RT: reverse transcriptase, PDE: phosphodiesterase, GPCR: G-protein coupled receptor, LRR: Leucine-rich repeat, vWF-D: Von Willebrand factor type D, TM: transmembrane domain.

A gene cluster of MT+ and small size specific genes plays a role during sex pheromone signaling

Among the differentially expressed genes for size and mating type, we identified three *C. closterium* genes that are located nearby on contig81: Ccl_14859, Ccl_14861 and Ccl_14874

(**Figure 4A**). Gene expression of each of the three genes was extremely specific for small

cells and restricted to the MT+ (**Figure 4A**). Consulting publicly available RNA-seq datasets,

we found that the three aforementioned genes were on average 109-fold upregulated during

early sexual reproduction in available RNA-seq data on top of their baseline expression in

small cells (**Figure 4A**) (Audoor *et al.*, 2024). Inspecting the expression pattern following the

crossing of compatible strains ZW2.20 and CA1.15 (T1: 9 h, T2: 14 h, T3: 27 h), we observed

that the induction of Ccl_14859 and Ccl_14861 was the strongest at the first time point T1,

which is the stage when sex pheromone signaling, mate finding and mating pair formation is

most intense (**Figure 4A**). The expression of Ccl_14874 peaked somewhat later: between

stage T1 and T2, when gametogenesis is at its maximum. Since the rapid induction of these

three genes during the early stages of mating may indicate a role in the response to sex

inducing pheromone (SIP) of the opposite mating type (Klapper *et al.*, 2021), we set up a

time-series RNA-seq experiment (**Figure 4B**). We specifically focused on the interval during

which cells become conditioned by SIP before the formation of mating pairs, by treating MT+

cells with SIP- containing filtrate and assessing global gene expression after 3 h and 9 h of re-

illumination following a dark-arrest. Compared to the untreated control, the expression of

each of the three genes was significantly induced by a treatment of 9 h with SIP-containing

filtrate (**Figure 4B**). Compared to their initial expression in small MT+ cells, both Ccl_14861

and Ccl_14874 were stimulated about 30-fold after 9h.

In terms of functional annotation, the amino acid sequence of Ccl_14859 shares similarity

with the *P. tricornutum* phytotransferrin Phatr3_J54465 (BLASTp against PLAZA Diatoms

v1.0 database, E-value: 3e-88, **Supplementary Figure 6**) (McQuaid *et al.*, 2018; Turnšek *et*

al., 2021) and encodes an N-terminal signal peptide. The other two genes (Ccl_14861,

Ccl_14874) which differ only by two amino acids at positions 303 and 306 (**Supplementary**

Figure 7), had no functional annotation except for an N-terminal signal peptide, indicating a

secretory pathway. Furthermore, no homologs could be detected in any other diatom genome

in the PLAZA Diatoms v1.0 database (BLASTp, E < 1e-3), making these two genes highly

taxon-specific markers for MT+ identity.

Figure 4: Gene expression of adjacent genes on *C. closterium* contig81 in different size classes, mating types and during three stages of sexual reproduction. (**A**) Bar plots show the average expression over replicates. The expression of individual replicates is visualized in Supplementary Figure 8 and Supplementary Table 3. Heatmaps show the log2 fold change between sexual crosses and vegetative control cultures across three time points: T1 (9 h post-cross, dominated by paired gametangia), T2 (14 h post-cross, gametes and zygotes) and T3 (27 h post-cross, auxospores). Expression data of sexual reproduction was obtained from Audoor *et al.*, (2024). CPM: counts per million, MT: mating type, SST: sexual size threshold, log2FC: log2(fold changes). (**B**) Bar plots show fold changes after 3 h and 9 h of re-illumination following a treatment of MT+ cells with Sex Inducing Pheromone (SIP) containing spent medium. Fold changes represent the expression after SIP treatment relative to the matched untreated control.

Stars indicate the significance of differential expression between SIP treated and control

conditions for each gene in each time point (false discovery rate-adjusted p-values, * p<0.05, **

p<0.01, *** p<0.001).

The presence of three closely located genes with a shared MT+ specific and SIP-responsive

properties suggests a shared function during early mate finding. While functional gene

clusters are commonly found in other eukaryotic taxa such as plants, animals and fungi

(Nützmann *et al.*, 2018), where they play a role in the same cellular process, often a metabolic

pathway, the only clear example known in diatoms has been described for fucoxanthin

chlorophyll proteins (Devaki & Grossman, 1993). Co-expression of clustered genes is

typically a result of co-regulation by the same transcription factors, through shared binding

sites or common chromatin-level regulation (Nützmann *et al.*, 2018). Further studies are

warranted to elucidate the functional significance and regulatory elements associated with this

gene cluster.

Mating type differentiation and size sensing genes are associated with pennate-

and *Cylindrotheca*-aged genes

Phylostratigraphic analysis, a method used to date the origin of genes or gene families by

looking at their homologs across different species, revealed that size- and mating type-

responsive DE genes were on average evolutionarily young compared to the entire expressed

transcriptome. Indeed, genes upregulated in small cells, in one or both mating types, were

significantly enriched in genes for which we could not detect homologs outside the

C. closterium genome (**Figure 5A**), and the same was true in the size-dependent genes from

the additional paired analysis (adjusted p-value $< 1e^{-5}$, hypergeometric test). These genes are

most likely involved in the species-specific pheromone signaling and mate finding behavior,

which become only apparent below the SST (Klapper *et al.*, 2021).

Size-sensing and size-dependent sexual maturation have been observed for most diatom

species from both centric and pennate clades (Chepurnov *et al.*, 2004; Poulíčková *et al.*,

2019). Nevertheless, none of the 21 purely size-responsive genes we found in *C. closterium*

were of diatom age, i.e., whose gene family was restricted to pennate and centric diatoms

(**Figure 5A**). While such diatom-aged size genes may have been missed due to the large

variability between genotypes limiting the power to detect DE genes, it may also suggest that

size-sensing in diatoms does not share a common master regulator. In other eukaryotic

organisms, physicochemical signals such as the scaling of the nucleus and membrane-free

structures, i.e., nucleolus, spindle, and microtubules, have been used to measure their size in

comparison to the cell size, the cell volume or the surface area (Marshall, 2015; Reber &

Goehring, 2015). Hence, mating type differentiation and sexual maturation may proceed

directly downstream of such physical signals through intracellular scaling without a

transcriptionally regulated master regulator.

Genes that were differentially expressed between mating types but only in small cells were

significantly more likely to be of pennate age compared to all other *C. closterium* genes

(**Figure 5A**). Indeed, many (30%) of the genes involved in mating type differentiation below

the SST were rooted within the pennate diatoms. Given that heterothally is a common

reproductive strategy in pennate diatoms while centric diatoms typically display homothallic

reproduction without mating types (Poulíčková *et al.*, 2019), these results suggest that the

molecular mechanisms underlying mating type determination have a shared origin in the

pennate diatoms around 137 million years ago (Nakov *et al.*, 2018).

Figure 5: Phylostratigraphic analysis of size and mating type-responsive genes and comparative analysis of gene families containing genes with a mating type- or size-biased expression in two pennate diatoms: *Cylindrotheca closterium* **and** *Seminavis robusta***.** (**A**) Bar plots showing the proportion of different phylostrata (phylogenetic gene ages) among sets of differentially expressed (DE) genes as well as the background (all expressed genes). Stars indicate the enrichment significance level of a given phylostratum relative to the background (*: p < 0.05, **: p < 0.01, hypergeometric test), MT: mating type. (**B**) For each gene family, the number of differentially expressed genes in each contrast is shown by circles, with genes that were differentially expressed in more than one contrast connected by a horizontal line. Pie charts show

the taxonomic composition of gene families, while bar charts show the size (number of genes) of

each family. LRR: leucine-rich repeat, vWF-D: Von Willebrand factor type D domain.

Leucine-rich repeat proteins are involved in mating type differentiation in

pennate diatoms

We next tried to pinpoint which common genes drive mating type differentiation in pennate

diatoms by performing a comparative analysis using existing transcriptomic libraries from

another heterothallic pennate diatom, *Seminavis robusta* (Osuna-Cruz *et al.*, 2020). We took

advantage of the presence of 36 and 12, vegetative RNA-seq samples from small untreated

MT+ and MT- cells respectively, to perform a differential expression analysis. This resulted

in a set of 183 genes that showed a mating type-specific expression response. We intersected

the gene families of these *S. robusta* mating type-specific genes with families to which the

112 *C. closterium* size- and mating type-biased genes belong. Seven gene families contained

DE genes in both species (**Figure 5B**).

Although the seven shared gene families range from small (11 genes) to large (992 genes), all

of them are predominantly composed of genes belonging to pennate diatoms, further

corroborating how mating type-dependent genes have evolved simultaneously with

heterothally at the origin of pennate diatoms. Genes from two families containing Leucine-

rich repeat (LRR) InterPro domain proteins showed a mating type biased response in both *C.*

closterium and *S. robusta* (**Figure 5B**). Since the LRR domain is typically involved in

protein-protein interaction or signaling, their conserved mating type dependent response

suggests a role in communication between the opposite mating types in pennate diatoms.

Although the first LRR family HOM02SEM000188 includes one *S. robusta* and one *C.*

closterium gene with a MT+ specific expression, phylogenetic analysis showed that these two

genes are not orthologous (**Supplementary Figure 9**). Hence, the functional and evolutionary

significance of their shared expression pattern is unclear.

On the other hand, HOM02SEM009752, a pennate-specific family with only 11 genes in 8

species, comprises two *C. closterium* genes, both of which are specific for small cells, but

with a contrasting expression pattern between mating types: gene Ccl_4437 was upregulated

in MT+ cells, whereas Ccl_5484 was upregulated in MT- cells (**Figure 5B**). The *S. robusta*

homolog (Sro702_g189950) showed a strong MT+ specific expression. Importantly, this

family encodes the Mating-type Related Minus 2 (*MRM2*) gene that was previously identified

in the raphid pennate diatom *P. multistriata*. *MRM2* is a component of the mating type

determination system of *P. multistriata*, where it is downstream of the sex-determining gene

MRP3 (Russo *et al.*, 2018).

Importantly, a paired analysis identified three LRR genes (Ccl_1352, Ccl_11832, Ccl_7129)

that are upregulated in small cells irrespective of the mating type, suggesting that some LRR

proteins may be involved in bidirectional recognition between mating types below the SST.

Mating-type Related Minus 2 (MRM2) is widely conserved in pennate diatoms

Multiple sequence alignment showed that proteins from the MRM2 family are conserved

through their LRR domain. Therefore, we set out to identify additional homologs of MRM2

by constructing a profile hidden Markov model (HMM) of the LRR domain in the 11 proteins

affiliated with HOM02SEM009752. Subsequently, the profile HMM was used to search the

PLAZA Diatoms reference dataset for additional proteins that are homologous to MRM2

through their LRR domain (Osuna-Cruz *et al.*, 2020). Phylogenetic analysis based on the full-

length protein sequence of the top-50 hits revealed a well-resolved clade containing the

original MRM2 gene family as well as two genes belonging to other gene families (**Figure

6A, Supplementary Figure 10**). The proteins in this MRM2 clade share structural

similarities, consisting of a predicted transmembrane domain on the C terminus and the LRR domain displayed extracellularly (**Figure 6A**). Motif discovery in the LRR domains of MRM2 homologs revealed the NxLt/sGxIPx motif typical to the plant specific (PS) family of diatom LRR proteins (**Figure 6A**) (Schulze *et al.*, 2015). The short (~50AA) N-terminal intracellular domain does not contain a kinase domain or other known signaling functionality, similar to the plant LRR-RLPs (leucine-rich repeat receptor-like proteins), which associate with co-receptors to transduce immune signals to the plant cell (Snoeck *et al.*, 2023). Together, MRM2 homologs were found in the genome of seven raphid pennate diatoms, as well as the araphid *Synedra acus*, suggesting a common origin at the root of the pennate diatoms.

Duplicated homologs of *MRM2* show an inverted mating type identity

The *MRM2* homologs of *C. closterium* and *S. robusta* have seemingly undergone gene duplication, with Ccl_4437 and Sro702_g189950 being closest related to the *P. multistriata* *MRM2* gene, whereas Ccl_5484 and Sro56_g032980 are more distant (**Figure 6A**). These distant out-paralogs of *MRM2* are lost or missing in most other pennate diatoms that encode only one copy of *MRM2*, including the original gene identified in *P. multistriata* (Russo *et al.*, 2018).

Our data suggest that both Ccl_4437 and Ccl_5484 are conditionally expressed only in cells below the SST (**Figure 6B**). Ccl_4437 was specifically expressed in MT+ strains, while Ccl_5484 showed the opposite expression pattern, being biased towards MT-. We propose calling these homologs *MRM2h+* (*MRM2* homolog expressed in MT+) and *MRM2h-* respectively. Expression profiles of the small cells from the *S. robusta* expression atlas are congruent with what we observed in *C. closterium*: one homolog (Sro702_g189950, *MRM2h+*) is MT+ specific, while the other (Sro56_g032980, *MRM2h-*) is MT- specific

(**Figure 6C**). To explore the variability of this response in other genotypes beyond the ones

included in the *S. robusta* expression atlas (MT-: 84A, 85B, MT+: 85A), we performed RNA

sequencing of six additional *S. robusta* genotypes below the SST, three for each mating type

(MT+: 85A-9, P5, P7 and MT-: 85A-1, 85A-6, P4). While this experiment confirmed the

MT- specific expression of *MRM2h-* in two out of three strains, *MRM2h+* was expressed at a

very low level in all six strains (average expression of 0.76 counts per million). Hence, the

mating type specific expression pattern of *MRM2h+* could not be confirmed in these

additional strains (**Supplementary Figure 11**).

Combined, the mating type specific expression of *MRM2* homologs in three species (*P.*

multistriata, C. closterium and *S. robusta*) and its distribution throughout the pennates point to

a role for *MRM2* as a key gene for mating type identity. As transcription of *MRM2* homologs

was activated only when the cell size decreased below the SST, the function of MRM2

appears to be specific for sexually competent cells. Similar studies on the transcriptomic

response to cells passing the SST should be undertaken in other life cycle model organisms

such as *S. robusta* and *P. multistriata* to determine if MRM2's size-dependency is conserved

throughout the pennate diatoms. The duplication and inversion of mating type preference in *S.*

robusta and *C. closterium* suggests that MRM2 homologs have evolved to play similar roles,

yet in opposite mating types. Finally, our data encourages discussion about how mating type

identity is assigned. Mating types are currently distinguished based on cellular behavior

during mating: MT- represents passive cells, while either the MT+ gametangia or gametes are

motile (Kaczmarska *et al.*, 2013). Our data suggests that distinguishing mating types by their

behavior may not reflect molecular evolution. To illustrate, *MRM2h+* genes Ccl_4437 and

Sro702_g189950 exhibited MT+ expression but are phylogenetically closest to the original *P.*

multistriata MRM2, which was expressed in MT- (**Figure 6**).

Figure 6: Phylogeny and expression of MRM2 homologs in pennate diatoms. (**A**) Bootstrap-consensus phylogenetic tree of MRM2 proteins in diatoms. Bootstrap support is shown as node labels. The full phylogenetic tree of the top-50 best MRM2 hits is available as Supplementary Figure 10. Next to the gene labels, their mating type-dependent expression pattern, gene family (GF), taxonomy and protein structures (InterPro domains and transmembrane domains predicted by Phobius) are shown. The inset shows the MEME sequence logo of the leucine-rich repeat domains of these proteins. (**B**) Bar plots showing expression in counts per million (CPM) of MRM2 homologs in large and small cells of different *Cylindrotheca closterium* strains (left) and during three stages of sexual reproduction (right). For the sexual reproduction, bar plots show the

average expression while dots show the individual RNA-seq samples. SST: sexual size threshold,

MRM2h+: MT+ biased homolog of MRM2, MRM2h-: MT- biased homolog of MRM2. **(C)**

Expression of MRM2 homologs in the *S. robusta* expression atlas (Osuna-Cruz *et al.*, 2020). Left:

boxplots of expression in MT+ (n = 108) and MT- (n = 21) samples in a range of non-sexual

conditions, spanning biotic interactions, environmental conditions, sexual reproduction, and diel

cycles. Right: expression during sexual reproduction covering three time points after crossing. Bar

plots show average expression while dots show individual RNA-seq samples.

Upregulation of *MRM2* homologs during sex suggests a function in recognition

between compatible mating types

The differentiation between mating types in raphid pennate diatoms is most apparent during

sexual reproduction, where attraction and cell-cell recognition ensure the correct pairing of

compatible gametangia and fusion of a MT+ with a MT- gamete (reviewed in Bilcke et al.,

2022). Therefore, we inspected the expression of *MRM2* homologs during different stages of

sexual reproduction (Audoor *et al.*, 2024; Osuna-Cruz *et al.*, 2020) (**Figure 6B,C**). Both *C.*

closterium MRM2 homologs as well as the *S. robusta MRM2h+* were significantly

transcriptionally upregulated in at least one stage of sexual reproduction. Strikingly, the *C.*

closterium gene *MRM2h+* is only upregulated during the early stage whereas *MRM2h-* was

most strongly upregulated in the last stage. Hence, not only are *MRM2* homologs in *C.*

closterium specific for the opposing mating types but they also appear to play a role in

different stages of sexual reproduction: gametangia pairing (*MRM2h+*) versus the

differentiation of gametes or auxospore development (*MRM2h-*).

Conclusions and Outlook

In this study, we leveraged experimental cell size manipulation to decipher the transcriptional reprogramming associated with size-based sexual maturation in diatoms. By simultaneously harvesting the same six genotypes in large and small cells, we avoided potential confounding due to the use of different genotypes in the two size classes. Furthermore, the use of full biological replication (i.e. independent genotypes) further ensures that the small set of DE genes we identified are robust and valid for a range of *C. closterium* strains. In general, size and mating type were not associated with large transcriptional differences in our experiment, which resulted in relatively few DE genes (< 100 genes per contrast). This may suggest that only few genes are involved in the differentiation of size and mating type classes or may alternatively be caused by the large variability between genotypes, making it harder to detect true DE genes. Indeed, correcting for the genotype effect in a paired analysis increased the number of identified genes by about 10-fold (from 21 to 182 genes), but this was only possible for the "Size" contrast.

A set of 29 mating type-specific genes were transcriptionally activated in cells that have passed the SST. We argue that such genes must underlie the chemical and behavioral differentiation between mating types, which only becomes apparent in small cells. Three such co-expressed genes were located nearby on the same contig. Their confined expression in small MT+ cells and coordinated induction in response to sex pheromones suggest an important role during the pairing process, in which MT+ cells are actively gliding towards an attracting MT- cell.

Furthermore, we demonstrated that the *P. multistriata* gene *MRM2* not only has a wider distribution in pennate diatoms than was previously reported (Russo *et al.*, 2018), but also displays mating type-specificity in at least three raphid pennate diatoms from diverse taxonomic clades (*C. closterium, P. multistriata, S. robusta*). Judging from the fact that these

plant-like LRR receptor proteins are most strongly expressed during cell pairing and gamete

fusion, we argue that MRM2 is part of an ancient partner recognition system that dates back

to the origin of the pennate diatoms. Consequently, the conservation of MRM2 is the first

molecular evidence for heterothally being an ancestral feature in pennate diatoms. While most

pennate diatoms indeed display a heterothallic mating system, some pennate species display

homothallic reproduction or facultative heterothally (Poulíčková *et al.*, 2019). Heterothally

would consequently have been secondarily lost in these species. Additionally, in accordance

with Russo and colleagues (2018) we did not find any homologs of the *P. multistriata* mating

locus *MRP3* outside the *Pseudo-nitzschia/Fragilariopsis* clade (Russo *et al.*, 2018),

suggesting that different clades use different mating type determination systems. In future

studies, assessing the conservation of mating type dependent genes such as *MRM2* in a

diverse set of heterothallic and homothallic pennate diatoms will be instrumental to

understand their evolutionary history. In that respect, the 100 Diatom Genomes project

(jgi.doe.gov/csp-2021-100-diatom-genomes) will soon make it possible to investigate life

cycle genes at an unprecedented scale.

While the earliest reports of the unique size reduction-restitution life cycle of diatoms date as

far back as 1869 (Pfitzer, 1869; Macdonald, 1869), the mechanism that cells use to accurately

measure their own size is still completely unknown. The fact that the size is constantly

monitored by the cell can be deduced from the sexual maturation after the experimental cell

size manipulation we performed using a glass Pasteur pipette. Such microsurgery-based size

manipulation, which was previously executed using a razor blade by van Stosch in 1965 (von

Stosch, 1965) and later repeated by Roshchin (Roshchin, 1994) and Chepurnov (Chepurnov *et

al*., 2004), represents a time and cost-efficient way to significantly reduce the generation time

of experimental models such as *C. closterium*. Using this approach, we identified a set of 21

genes that are upregulated in small cells (<SST) relative to large cells (>SST) in both mating

types. These size markers include several nucleus-localized and DNA-interacting genes,

suggesting a function as size-dependent switches, activating downstream sexual maturation

genes only below the SST, but need to be further validated by functional experiments.

However, none of these genes could be clearly linked to a physical function in cell size

sensing, and neither did any of them occur specifically in the genome of both pennate and

centric diatoms. Hence, it remains unexplained how cells measure their own size and whether

a diatom-wide transcriptional master regulator exists, which our analysis may have lacked the

power to detect. The identification of the first molecular markers for cell size dependent

sexual maturation in diatoms - including conserved key genes such as *MRM2* - paves the road

for future research into the physicochemical and molecular changes taking place when cells

pass the SST.

Acknowledgements

G.B. is a postdoctoral fellow supported by Fonds Wetenschappelijk Onderzoek (FWO,

1228423N). W.V. acknowledges partial financial support from FWO project G001521N,

BOF/GOA projects No. 01G01715 and 01G01323, as well as infrastructure funded by

EMBRC Belgium-FWO project GOH3817N. The BCCM/DCG culture collection is supported

by the Belgian Science Policy (Belspo).

We sincerely thank Dr. Koen Van den Berge for helpful discussion about experimental

design. We are grateful towards Olga Chepurnova for help with maintaining and manipulation

of diatom cultures in cryopreservation tanks. We would further like to thank Dr. Katerina

Pargana for her help with bioinformatic analysis and Dr. Petra Bulánková for assisting with

molecular biology protocols.

Competing Interests

The authors declare that they have no conflict of interest.

Authors Contribution

DB, GB, and WV conceived the study with input from KV and LDV. DB performed microsurgery and transcriptomic experiments. DB and SD jointly performed RNA extraction and quality control, while SA and DB were responsible for RT-qPCR of selected genes. Differential expression and comparative transcriptomic analyses were performed by GB with the support of DB. DB, GB, WV, and KV wrote the manuscript. All authors commented on and approved of the manuscript.

References

Audoor S, Bilcke G, Pargana K, Belišová D, Thierens S, Van Bel M, Sterck L, Rijsdijk N, Annunziata R, Ferrante MI, *et al.* 2024. Transcriptional chronology reveals conserved genes involved in pennate diatom sexual reproduction. *Molecular Ecology* **33**: e17320.

Bachtrog D. **2006**. A dynamic view of sex chromosome evolution. *Current Opinion in Genetics and Development* **16**: 578–585.

Basu S, Patil S, Mapleson D, Russo MT, Vitale L, Fevola C, Maumus F, Casotti R, Mock T, Caccamo M, *et al*. 2017. Finding a partner in the ocean: molecular and evolutionary bases of the response to sexual cues in a planktonic diatom. *New Phytologist* **215**: 140–156.

Bilcke G, Van Den Berge K, De Decker S, Bonneure E, Poulsen N, Bulankova P, Osuna-Cruz CM, Dickenson J, Sabbe K, Pohnert G, *et al*. 2021. Mating type specific transcriptomic response to sex inducing pheromone in the pennate diatom Seminavis robusta. *The ISME Journal* **15**: 562–576.

Bilcke G, Ferrante MI, Montresor M, De Decker SD, Veylder L De, Vyverman W. **2022**. Life Cycle Regulation. *The Molecular Life of Diatoms*: 205–228.

Bucchini F, Del Cortona A, Kreft Ł, Botzki A, Van Bel M, Vandepoele K. **2021**. TRAPID

2.0: A web application for taxonomic and functional analysis of de novo transcriptomes.

Nucleic Acids Research **49**: e101–e101.

Chepurnov VA, Mann DG, Sabbe K, Vyverman W. **2004**. Experimental studies on sexual

reproduction in diatoms. *International Review of Cytology* **237**: 91–154.

Davidovich, NA . **2002**. Sexual heterogeneity of the clones of Nitzschia longissima (Breb.)

Ralfs (Bacillariophyta). *Algologiya* **12**: 279–289.

Davidovich NA, Davidovich OI, Podunai YA, Shorenko KI, Kulikovskii MS. **2015**.

Reproductive properties of diatoms significant for their cultivation and biotechnology.

Russian Journal of Plant Physiology **62**: 153–160.

Devaki B, Grossman AR. **1993**. Characterization of gene clusters encoding the fucoxanthin

chlorophyll proteins of the diatom Phaeodactylum tricornutum. *Nucleic acids research*

21(19): 4458-4466

Field CB, Behrenfeld MJ, Randerson JT, Falkowski P. **1998**. Primary production of the

biosphere: Integrating terrestrial and oceanic components. *Science* **281**: 237–240.

Gillard J, Frenkel J, Devos V, Sabbe K, Paul C, Rempt M, Inzé D, Pohnert G, Vuylsteke

M, Vyverman W. **2013**. Metabolomics Enables the Structure Elucidation of a Diatom Sex

Pheromone. *Angewandte Chemie International Edition* **52**: 854–857.

Hobza R, Cegan R, Jesionek W, Kejnovsky E, Vyskot B, Kubat Z. **2017**. Impact of

repetitive elements on the Y chromosome formation in plants. *Genes* **8** (11): 302.

Kaczmarska I, Ehrman JM, Mills KE, Sutcliffe SG, Samanta B. **2022**. Vegetative cell

enlargement in selected centric diatom species–an alternative way to propagate an

individual genotype. *European Journal of Phycology* **58**(3), 315-332.

Kaczmarska I, Poulíčková A, Sato S, Edlund MB, Idei M, Watanabe T, Mann DG. **2013**.

Proposals for a terminology for diatom sexual reproduction, auxospores and resting stages.

Diatom Research **28**: 263–294.

Katoh K, Standley DM. **2013**. MAFFT multiple sequence alignment software version 7:

Improvements in performance and usability. *Molecular Biology and Evolution* **30**: 772–

780.

Kent T V., Uzunović J, Wright SI. **2017**. Coevolution between transposable elements and

recombination. *Philosophical Transactions of the Royal Society B: Biological Sciences* **372**

(1736), 20160458.

Klapper F, Audoor S, Vyverman W, Pohnert G. **2021**. Pheromone Mediated Sexual

Reproduction of Pennate Diatom Cylindrotheca closterium. *Journal of Chemical Ecology*

47: 504–512.

Klapper FA, Kiel C, Bellstedt P, Vyverman W, Pohnert G. **2023**. Structure Elucidation of

the First Sex-Inducing Pheromone of a Diatom. *Angewandte Chemie - International*

Edition **62** (43), e202307165.

Macdonald JD. **1869**. I.— On the structure of the Diatomaceous frustule, and its genetic

cycle . *Annals and Magazine of Natural History* **3**: 1–8.

Marshall WF. **2015**. How Cells Measure Length on Subcellular Scales. *Trends in Cell*

Biology **25**: 760–768.

McQuaid JB, Kustka AB, Oborník M, Horák A, McCrow JP, Karas BJ, Zheng H,

Kindeberg T, Andersson AJ, Barbeau KA, *et al.* **2018**. Carbonate-sensitive

phytotransferrin controls high-affinity iron uptake in diatoms. *Nature* **555**: 534–537.

Moeys S, Frenkel J, Lembke C, Gillard JTF, Devos V, Van Den Berge K, Bouillon B,

Huysman MJJ, De Decker S, Scharf J, *et al.* **2016**. A sex-inducing pheromone triggers

cell cycle arrest and mate attraction in the diatom Seminavis robusta. *Scientific Reports* **6**:

1–13.

Nagai S, Hori Y, Manabe T, Imai I. **1995**. Restoration of cell size by vegetative cell

enlargement in Coscinodiscus wailesii (Bacillariophyceae). *Phycologia* **34**: 533–535.

Nakov T, Beaulieu JM, Alverson AJ. **2018**. Accelerated diversification is related to life

history and locomotion in a hyperdiverse lineage of microbial eukaryotes (Diatoms,

Bacillariophyta). *New Phytologist* **219**: 462–473.

Nützmann HW, Scazzocchio C, Osbourn A. **2018**. Metabolic gene clusters in eukaryotes.

Annual Review of Genetics **52**: 159–183.

Osuna-Cruz CM, Bilcke G, Vancaester E, De Decker S, Bones AM, Winge P, Poulsen N,

Bulankova P, Verhelst B, Audoor S, *et al.* 2020. The Seminavis robusta genome provides

insights into the evolutionary adaptations of benthic diatoms. *Nature Communications* **11**

(1), 3320.

Patro R, Duggal G, Love MI, Irizarry RA, Kingsford C. **2017**. Salmon provides fast and

bias-aware quantification of transcript expression. *Nature methods* **14**: 417–419.

Pfitzer E. 1869. Über den Bau und die Zellteilung der Diatomeen. Botanische Zeitung.

27(46):774-6

Pinseel E, Nakov T, Van den Berge K, Downey KM, Judy KJ, Kourtchenko O, Kremp

A, Ruck EC, Sjöqvist C, Töpel M, *et al.* 2022. Strain-specific transcriptional responses

overshadow salinity effects in a marine diatom sampled along the Baltic Sea salinity cline.

ISME Journal **16**: 1776–1787.

Poulíčková A, Mann DG, Mann DG. **2019**. Diatom Sexual Reproduction and Life Cycles.

Diatoms: Fundamentals and Applications. 245–272.

Reber S, Goehring NW. **2015**. Intracellular scaling mechanisms. *Cold Spring Harbor*

Perspectives in Biology **7** (12), a019067.

Robinson MD, McCarthy DJ, Smyth GK. **2009**. edgeR: A Bioconductor package for

differential expression analysis of digital gene expression data. *Bioinformatics* **26**: 139–

140.

Roshchin, AM. 1994. Zhiznennye tsiklydiatomovykh vodoroslej. *Naukova Dumka, Kiev*. 170

Ruggiero A, Mager S, Manfellotto F, Tuccio V Di, Nishimura T, Pan L, Russo MT,

Sanges R, Ferrante MI. **2023**. DNA methylation is linked to the monoallelic expression of

MRP3, a diatom mating type determining gene. *bioRxiv*, 2023-10.

Russo MT, Vitale L, Entrambasaguas L, Anestis K, Fattorini N, Romano F, Minucci C,

De Luca P, Biffali E, Vyverman W, *et al.* **2018**. MRP3 is a sex determining gene in the

diatom Pseudo-nitzschia multistriata. *Nature Communications* **9**: 1–10.

Schulze B, Buhmann MT, Río Bártulos C, Kroth PG. **2015**. Comprehensive computational

analysis of leucine-rich repeat (LRR) proteins encoded in the genome of the diatom

Phaeodactylum tricornutum. *Marine Genomics* **21**: 43–51.

Slotkin RK, Martienssen R. **2007**. Transposable elements and the epigenetic regulation of

the genome. *Nature Reviews Genetics* **8**: 272–285.

Snoeck S, Garcia AG, Steinbrenner AD. **2023**. Plant Receptor-like proteins (RLPs):

Structural features enabling versatile immune recognition. *Physiological and Molecular*

Plant Pathology **125**: 102004.

Soneson C, Love MI, Robinson MD. **2016**. Differential analyses for RNA-seq: Transcript-

level estimates improve gene-level inferences. *F1000Research,* **4**.

Stock W, Vanelslander B, Rüdiger F, Sabbe K, Vyverman W, Karsten U. **2019**. Thermal

niche differentiation in the benthic diatom Cylindrotheca closterium (Bacillariophyceae)

complex. *Frontiers in Microbiology* **10**: 1395.

von Stosch HA. **1965**. Manipulierung der Zellgrosse von Diatomeen im Experiment.

Phycologia **5**: 21–44.

Turnšek J, Brunson JK, Martinez Viedma MDP, Deerinck TJ, Horák A, Oborník M,

Bielinski VA, Allen AE. **2021**. Proximity proteomics in a marine diatom reveals a putative

cell surface-to-chloroplast iron trafficking pathway. *eLife* **10**: 1–113.

Vanormelingen P, Vanelslander B, Sato S, Gillard J, Trobajo R, Sabbe K, Vyverman W.

2013. Heterothallic sexual reproduction in the model diatom Cylindrotheca. *European*

Journal of Phycology **48**: 93–105

Vanstechelman I, Sabbe K, Vyverman W, Vanormelingen P, Vuylsteke M. **2013**. Linkage

Mapping Identifies the Sex Determining Region as a Single Locus in the Pennate Diatom

Seminavis robusta (T Yin, Ed.). *PLoS ONE* **8**: e60132.