Unveiling the influence of tumor and immune signatures on immune checkpoint therapy in advanced lung cancer

Authors and affiliations

Nayoung Kim[1,2,10], Sehhoon Park[3,10], Areum Jo[1,2], Hye Hyeon Eum[1,2], Hong Kwan Kim[4], Kyungjong Lee[5], Jong Ho Cho[4], Bo Mi Ku[6], Hyun Ae Jung[3], Jong-Mu Sun[3], Se-Hoon Lee[3], Jin Seok Ahn[3], Jung-Il Lee[7], Jung Won Choi[7], Dasom Jeong[1,2], Minsu Na[1,2], Huiram Kang[1,2], Jeong Yeon Kim[8], Jung Kyoon Choi[8], Hae-Ock Lee[1,2,9,*], and Myung-Ju Ahn[3,*]

[1] Department of Microbiology, College of Medicine, The Catholic University of Korea, Seoul, 06591, Korea.

[2] Department of Biomedicine and Health Sciences, Graduate School, The Catholic University of Korea, Seoul, 06591, Korea.

[3] Division of Haematology-Oncology, Department of Medicine, Samsung Medical Center, Sungkyunkwan University School of Medicine, Seoul, 06351, Korea.

[4] Department of Thoracic and Cardiovascular Surgery, Samsung Medical Center, Sungkyunkwan University School of Medicine, Seoul, 06351, Korea.

[5] Division of Pulmonary and Critical Care Medicine, Department of Medicine, Samsung Medical Center, Sungkyunkwan University School of Medicine, 06351, Seoul, Korea.

[6] Research Institute for Future Medicine, Samsung Medical Center, Sungkyunkwan University School of Medicine, Seoul, 06351, Korea.

[7] Department of Neurosurgery, Samsung Medical Center, Sungkyunkwan University School of Medicine, Seoul, 06351, Republic of Korea

[8] Department of Bio and Brain Engineering, KAIST, Daejeon, Korea.

[9] Precision Medicine Research Center, College of Medicine, The Catholic University of

Korea, Seoul, 06591, Korea.

[10] These authors contributed equally: Nayoung Kim, Sehhoon Park.

[*] Corresponding authors.

Running title: Tumor and immune signatures in advanced lung cancer therapy

Corresponding Author: Myung-Ju Ahn, Division of Haematology-Oncology, Department of

Medicine, Samsung Medical Center, Sungkyunkwan University School of Medicine, 81

Irwon-ro, Gangnam-gu, Seoul 06351, Korea. Phone: +82-2-3410-3438; Fax: +82-2-3410-

1754; E-mail: silk.ahn@skku.edu; Hae-Ock Lee, Department of Microbiology, College of

Medicine, The Catholic University of Korea, 222 Banpo-daero, Seocho-gu, Seoul 06591,

Korea. Phone: +82-2-2258-8155; Fax: +82-2-2258-7744; E-mail: haeocklee@catholic.ac.kr

Conflicts of interest: The authors declare no potential conflicts of interest.

Keywords

Immune checkpoint inhibitors, Responsiveness, Advanced lung cancer, Single-cell,

Transcriptome

Abstract

This study investigates the variability among patients with non-small cell lung cancer

(NSCLC) in their responses to immune checkpoint inhibitors (ICI). Recognizing that patients

with advanced-stage NSCLC rarely qualify for surgical interventions, it becomes crucial to

identify biomarkers that influence responses to ICI therapy. We conducted an analysis of

single-cell transcriptomes from 33 lung cancer biopsy samples, with a particular focus on 14

core samples taken before the initiation of palliative ICI treatment. Our objective was to link

tumor and immune cell profiles with patient responses to ICI. We discovered that ICI non-

responders exhibited a higher presence of CD4+ regulatory T cells, resident memory T cells,

and TH17 cells. This contrasts with the diverse activated CD8+ T cells found in responders.

Furthermore, tumor cells in non-responders frequently showed heightened transcriptional

activity in the NF-kB and STAT3 pathways, suggesting a potential inherent resistance to ICI

therapy. Through the integration of immune cell profiles and tumor molecular signatures, we

achieved an discriminative power (AUC) exceeding 95% in identifying patient responses to

ICI treatment. These results underscore the crucial importance of the interplay between tumor

and immune microenvironment, including within metastatic sites, in affecting the

effectiveness of ICIs in NSCLC.

Introduction

Treatment landscape in cancer has rapidly evolved with the introduction of immune

checkpoint inhibitors (ICI). Among various immune checkpoints, antibody-targeting

programmed cell death-1 (PD-1) and its ligand (PD-L1) have demonstrated clinical benefits

over conventional systemic chemotherapy in patients with non-small cell lung cancer

(NSCLC). They have been approved as either monotherapy in patients with high PD-L1

expression [1] or combined with cytotoxic chemotherapy regardless of PD-L1 expression [2,3].

Moreover, the clinical benefits were validated in unresectable stage III NSCLC as

consolidation therapy after definitive chemoradiotherapy or early stage NSCLC (Ib-IIIA) as

adjuvant therapy after curative surgery [4].

There have been many efforts to elucidate the predictive biomarkers of PD-(L)1 inhibitors.

The PD-L1 expression in tumor tissue evaluated via immunohistochemistry has been

incorporated as a companion diagnostic biomarker from the early clinical trials, which

enhanced the response rate up to 46% in patients with PD-L1 ≥50% and showed an overall

survival rate of up to 26.3 months [5]. Other biomarkers, such as tumor mutation burden or

gene expression profile, also demonstrated a positive predictive value [6,7]. Recent large-scale

meta-analysis of clinico-immunogenomics shows that both tumor- and T cell intrinsic factors

exert a substantial impact on ICI response [8], supporting the necessity of in-depth investigation

of high-throughput profiles of tumor and microenvironment.

ICI treatment modifies systemic immune responses, which can be monitored by alterations in

the proportion of specific immune cell populations. An increase in PD-1+Ki67+CD8+ T cells

in peripheral blood after PD-1 inhibitor treatment is associated with better outcomes in

patients with NSCLC [9,10]. This cell type has also been identified in tumor tissues prior to the

ICI treatment, where it is linked to clinical outcomes, and shows impaired production of

classical effector cytokines [11]. More specifically, PD-1 expression is upregulated upon

antigen recognition [12], indicating that certain T cells in the tumor microenvironment are

actively engaged as tumor-specific T cells. Beyond CD8+ T cells, other immune cell types,

such as myeloid-derived suppressor cells or regulatory T cells, which regulate tumor-specific

T cell immunity may also influence therapeutic outcomes [13-15]. Overall, the multicellular

regulation of the tumor-immune microenvironment emphasizes the importance of profiling

systemic tumor and immune cells at single cell resolution to investigate factors associated

with responses to ICI treatment.

Results

Variability and features of the lung cancer samples. We conducted scRNA-seq on 33 lung cancer samples from 26 patients treated with immune checkpoint inhibitors (ICI) between August 2017 and December 2019 to understand how cellular dynamics in lung cancer affect treatment sensitivity to PD-(L)1 inhibitors, used alone or in combination (Fig. 1a and Table S1). Immune checkpoint therapy provides clinical benefit in advanced metastatic NSCLC across different treatment lines [16]. Notably, our samples have been collected from various tissue sites. In the scRNA-seq analysis, all specimens were used for the cell type profiling in an unbiased manner. For the evaluation of clinical outcomes, only refined 14 core samples from 11 patients were used to minimize sample specific variations. Exclusion criteria from the core group encompass samples with treatment applied as adjuvant therapy, acquired after ICI treatment, no tumor content, non-evaluable for the clinical response, or histology other than non-small cell lung cancer such as nuclear protein in testis (NUT) and small cell lung cancer (SCLC) (Table 1 and Fig. 1a). Of the 11 core patients, 8 had adenocarcinoma (ADC) and 3 had squamous cell carcinoma (SQ). Clinical outcomes of ICI were partial response (PR) in four patients, stable disease (SD) in two patients, and progressive disease (PD) in five patients. Patients were classified as responders (PR) and non-responders (SD and PD) according to ICI response.

Due to the diversity of sample collection sites, our data may be influenced by varied immune cell composition at different sample collection sites. Therefore, we analyzed immune and stromal cell subsets across early-stage (tLung) and late-stage (tL/B) lung tumors, and metastatic lymph nodes (mLN), comparing them to normal lung (nLung) and lymph node (nLN) tissues. This analysis was conducted on public scRNA-seq data from 43 samples from 33 lung adenocarcinoma (LUAD) patients [17] (Fig. S1a-c). Although there were differences in

tissue-specific resident populations, we found that the immune cell profiles, especially T/NK

cells of mLN were similar to those of primary tumor tissues (Fig. S1d-g).

Classification of immune cell subset in lung cancer. Global cell-type profiling (Fig. 1b,c

and Table S2) illustrates the cellular composition of each sample as epithelial/tumor cells,

fibroblasts, endothelial cells, T/natural killer (NK) cells, B/plasma cells, myeloid immune

cells, and mast cells. Individual samples show variations in epithelial/tumor content as well

as in immune cell composition (Fig. 1d,e).

For further analysis of immune cell subtypes, we applied sequential subclustering on global

immune cell clusters. As scRNA-seq shows limited performance in separating CD4+ and

CD8+ T cell subsets, antibody-derived tag (ADT) information [18] was used to complement the

transcriptome data and to predict CD4+ T cells, CD8+ T cells, and NK cells (Fig. 2a). Finally,

fourteen CD4+ and fourteen CD8+ T cell subclusters were identified excluding <5%

ambiguous cells (Fig. 2b,c and Table S2). In the CD4+ T cell compartment, naïve-like T cells

(TN, CD4_cluster0) and central memory T cells (TCM, CD4_cluster1) expressing *SELL,*

TCF7, LEF1, and *CCR7* genes or tissue-resident memory T cells (TRM, CD4_clusters3, 5, 6)

expressing *NR4A1, MYADM,* and *PTGER4* genes were abundant in most samples. Regulatory

T cells (Treg, CD4_cluster2) with *FOXP3, CTLA4, ICOS,* and *BATF* expression were also

abundant, which has been demonstrated as tumor-specific alterations in the tissue

microenvironment [17,19,20]. In the CD8+ T cell compartment, effector memory T cells (TEM),

effector T cells (TEFF), effector memory CD45RA positive cells (TEMRA) (CD8_clusters0,

2, 3, 8) expressing *PRF1* and *IFNG* were dominant over TN/TCM (CD8_clusters4, 5) types.

Exhausted T cells (TEX, CD8_clusters1, 12) expressed multiple checkpoint genes (*HAVCR2*

and *PDCD1*) along with high levels of *PRF1*, *IFNG*, *CXCR3,* and *CXCL13*. Co-expression of

cytotoxic effectors and checkpoint molecules in TEX clusters indicates that cluster

populations may retain functional capacity as cytotoxic effector T cells [21]. Further, clonotype

analysis of TCR supported the T cell subset classification demonstrating higher clonal

expansion in the CD8+ T cell compartment than that in CD4+ T cells, with the highest levels

within the TEX subclass (Fig. S2a-c).

Natural killer cells can be subclassified as CD56bright, transitional, active, and mature types

[22] (Fig. S3a,b and Table S2). Active NK cells expressed the highest level of *PRF1*, *TNF,* and

IFNG, reflecting a cytotoxic effector function.

Compared to T cell clusters, fewer B/plasma cells were detected as follicular (B_clusters 0, 1,

2, 4, 6, 11), germinal center (B_cluster 10), and plasma/mucosa-associated lymphoid tissue

(MALT) B cells (B_clusters 3, 7, 12) (Fig. S3a,b and Table S2). Plasma/MALT B cells

manifested higher levels of clonal expansion of BCR than follicular B cells (Fig. S3c,d).

Identical clonality of some follicular B and plasma cells suggests in situ maturation and

differentiation of B cells to plasma cells in tumor tissues (Fig. S3c, clonotype 16).

Myeloid cells were composed of monocytes, dendritic cells, and a large number of

macrophages   (Fig.   S3a,b   and   Table   S2).   CD14+CD16-   classical   monocytes

(Myeloid_clusters1,6) were predominantly found over non-classical CD14^{lo}CD16+ types

(Myeloid_cluster15). Alveolar macrophages (Alveolar Mac, Myeloid_cluster0) expressed

well-defined marker genes such as *MARCO*, *FABP4*, and *MCEMP1* along with anti-

inflammatory genes such as *CD163, APOE,* and *C1QA/B/C*. Monocyte-derived macrophages

represent heterogeneous populations with a similar gene expression profile to alveolar

macrophages (Mo-Mac, Myeloid_clusters2, 3, 5, 9, 11, 13), with an elevated chemokine gene

expression (CXCL10+ Mo-Mac, Myeloid_clusters4, 14, 16), or with active cell cycle

progression (Proliferating Mac, Myeloid_cluster7). Dendritic cells were categorized as

CD1c+ (Myeloid_cluster8, *CD1C* and *ITGAX*), activated (Myeloid_cluster17, *CCR7* and

LAMP3), and CD141+ (Myeloid_cluster19, *CLEC9A* and *XCR1*) subclusters.

Overall immune cell composition is comparable to those reported in previous studies [17,19,20].

Immune cell landscape fostering the ICI response. In global cell-type profiling (Fig. 1b,c),

abundance in T, NK, or myeloid cell types shows no difference between responders and non-

responders (Fig. 1d, e). Nonetheless, as specific differentiation features within the cell types

may influence the response to ICI treatment, we compared the response groups using the

proportion of subclusters within CD4+ T, CD8+ T, NK, B/plasma, and myeloid cells. After

subclustering (Fig. 2b and Fig. S3a), three subsets of CD4+ T cells, i.e., Treg, TRM, and

CD4+ T helper 17 (TH17), were significantly (p<0.01) overrepresented in the non-responder

group (Fig. 2d). In contrast, among CD8+ T cell populations, TEM subsets demonstrated a

modest level of association with the patients who responded well against progressive disease

(Fig. 2d, p=0.06). TCR clonotype analysis supported the cellular dynamics such that clonal

expansion was more prominent in cytotoxic CD8+ T cells over CD4+ Tregs in the responder

group (Fig. 2e and Fig. S2c). Overall landscape in each cell type (Fig. 2f) suggests that CD4+

Treg and TRM as well as follicular B cells may interfere with the ICI response, whereas

CD8+ T cell activation (TEM, TEMRA/TEFF, and TEX), mature NK cells, and CXCL10+

Mo-Mac cells support the ICI response. The balance between separate immune cell types

informs immune regulatory axes that may be targeted to favor the activation of tumor-

reactive immunity.

Systemic evaluation of the immune microenvironment associated with ICI response.

Next, we evaluated the immune microenvironment as an entity by using all immune cells as a

denominator in the subtype proportions. In this setting, we used diverse clinical group

comparisons and identified the immune cell blocks separated by clinical outcomes (Fig. 3a).

The immune cell blocks overrepresented in the non-responder groups consisted of CD4+

Treg, follicular B cells, and CD4+ TH17/TRM/ T helper 1 (TH1)-like cells. In the immune

cell blocks of the responder groups, CD8+ TEM cells showed the strongest enrichment along

with the other CD8+ TEX/TEMRA/TEFF/Mitochondria (MT) high cells as well as

CXCL10+ Mo-Mac. Despite the immune footprints of the ICI responders, extensive

variations among individual patients (Fig. 3b) hamper patient stratification solely based on

the immune profiles.

Tumor cell signatures associated with ICI response. We investigated the associations

between genomic characteristics in tumor and ICI response. The constraints of mutation

analysis with 10x chromium data complicated the direct correlation between tumor mutation

burden and ICI outcome. Rather, we assessed copy number alterations (CNA) indirectly, via

chromosomal gene expression patterns. These analyses revealed a moderate correlation

between low levels of CNA, including both gain and loss of heterozygosity, and positive

responses to ICI (Fig. S4). This finding is consistent with the result from previous genetic

association studies [23].

To assess gene expression characteristics of tumors influencing ICI response, we separated

malignant tumor cell clusters from normal epithelial cell types (Fig. S5). Subsequent DEG

analysis (Fig. 4a and Table S3) identified genes in poor response groups linked to the

regulation of cell death, cell motility, and cell activation (Fig. S6 and Table S4). The DEGs

were refined later by combinations of various tumor signatures separating responder and non-

responder groups (Table S5).

Next, to explore the existence of gene programs and modules influencing the ICI response,

we applied factorization using non-negative matrix factorization (NMF) and scINSIGHT [24].

Among 30 factors from NMF across all malignant cells, we identified factors showing high

loadings for a specific RECIST group as NMF programs p1~4 (Fig. 4b, c). There were clear

distinction among RECIST groups according to the gene expression levels associated with

these NMF programs (Fig. 4d and Table S5). To identify gene modules consistent across

different patients, we examined RECIST-specific modules though scINSIGHT analysis [24]

(Fig. 4e). Unfortunately, we found that contributions to these gene modules varied

significantly among patients. To mitigate this variability, we adjusted the RECIST-specific

modules by combining genes from the original modules (Table S5). The refined gene

modules showed a specific gene expression pattern for each RECIST group, similar to the

NMF programs (Fig. 4f). Overall, both genes and their functional categories segregated

depending on the selection techniques used (Fig. 4g, h). However, transcription factors

governing the signatures derived from DEG, NMF, and scINSIGHT analyses consistently

delineated between responders and non-responders (Fig. 4i). Responder-specific gene

signatures showed associations with the transcription factors Regulatory Factor X Associated

Ankyrin Containing Protein (RFXANK), Regulatory Factor X Associated Protein (RFXAP),

and Regulatory Factor X5 (RFX5). This RFX protein complex has emerged as a positive

biomarker for the immune response in diverse cancer types [25]. Non-responder-specific gene

signatures were regulated by Activator Of Transcription 3 (STAT3) and Nuclear Factor

Kappa B Subunit 1 (NFKB1), known to play roles in PD-L1 regulation and T cell activation

in cancer [26].

We also adopted principal component analysis (PCA) to isolate correlated gene signatures

variably expressed in tumor cells (Fig. S7, and Table S4, S5). Among the top 10 PCs,

negatively correlated genes in PC2, PC7, and PC8 distinguished the tumor cells in the poor

response groups, whereas positively correlated genes in PC6 and PC9 were upregulated in the

better response groups (Fig. S8a-c). Tumor cells from the PR group had low PC2.neg scores,

suggesting low growth factor/type I interferon response signaling as a tumor cell-specific

positive predictor of the ICI response. Conversely, high levels of growth factor/type I

interferon response signaling in tumor cells may present intrinsic resistance to PD-(L)1

inhibitor alone or in combination. Type I interferon is known to drive anti-tumor effect

directly or indirectly on tumor and surrounding immune cells, but also acts to counter the

anti-tumor effect by inducing CD8+ T cell exhaustion and up-regulating immune-suppressive

genes on tumor cells [27]. We assessed whether tumor cell signatures are applicable in

association with ICI response in other tumors. They had a modest influence on the response

to ICI treatment of melanoma (Fig. S8d) in bulk gene expression data [28,29].

These tumor cell signatures were specifically linked to elucidating the ICI response, but did

not demonstrate any prognostic value for LUAD or lung squamous cell carcinoma (LUSC) in

the TCGA RNA sequencing data (Table 2).

Combination of tumor signatures and immune cell dynamics classify ICI response.

Immune cell dynamics or tumor signature alone has a limited capacity to profile the

therapeutic outcome of PD-(L)1 inhibitor alone or in combination (Fig. 5a). Similar to the

immune cell blocks and tumor signatures that were over-represented in the poor response

group, each of CD4+ Treg, CD4+ TH17, PC7.neg, INT.down, and UNION.down was

significantly associated with ICI response in univariate regression analysis (Fig. 5b). PC7.neg

denotes genes negatively correlated with PC7, a principal component extracted from PCA

that distinguishes tumor cells in poor response groups. INT.down and UNION.down

represent the intersection and union of down-regulated genes in the responder group,

respectively. The variation in ICI response was not affected by clinical variables of tissue

origin, cancer subtype, pathological stage, and smoking status. When we performed a

combined analysis of the top tumor-immune features to classify response, the discriminative

power (AUC) was improved to over 95% (Fig. 5c). Overall, features of the non-responders,

especially CD4+ Treg, B/Plasma cells, INT.down, and UNION.down, showed a higher

estimate than those of responders. These non-responder features suggest heterogeneous

mechanisms of resistance conferred by tumor and immune regulatory axes.

Discussion

ICI alone, or in combination with chemotherapy, are considered standard first-line therapy for

patients with NSCLC. NSCLC may harbor large numbers of genetic perturbations due to

genotoxic environmental exposure, which likely generate high mutation burden or

neoantigens [30]. The neoantigen-directed T cell response is hampered by diverse immune

suppressive mechanisms exerted by tumor cells and the immune regulatory network [31].

Current ICIs targeting PD-(L)1 aim one angle of many suppressive mechanisms, and

identifying the features of non-responders will reveal additional regulatory angles to improve

ICI response.

Immune regulatory network would determine the balance between activation and suppression

of tumor-directed immunity. In previous studies, prediction of the response to ICI highlighted

CD8+ cytotoxic effector T cells and CD4+ Tregs [32]. The involvement of effector CD8+ T

cells is consistent in most studies regardless of the cellular origin (blood or tissues) or tumor

type (melanoma, lung cancer) [33]. Their phenotypes slightly differ depending on the cellular

resolution of the study. Our data provided the highest resolution cell types, and CD8+ TEM

cells (*GZMK*, *CXCR4* expression) were overrepresented in the responders. By comparison,

Tregs were underrepresented in the responder group. Previously, our group reported a

contrasting result demonstrating an increase in Tregs in the posttreatment blood samples (not

baseline) in the responder group [34]. This discrepancy can be explained by the differences in

the measurements, sites, and timing of sampling, and the resolution of subpopulations. In

mouse preclinical models, PD-L1 inhibitor treatment induces T cell expansion of all

phenotypes including CD4+/CD8+ TEFF and Tregs [35]. Thus, Treg expansion captured in the

posttreatment blood samples may represent overall immune activation in human patients.

Alternatively, heterogeneity of Tregs and complex effects of PD-(L)1 inhibition on this cell

type may contribute to variable results in response prediction.

The identification of an abundance of CD4+ TRM cells as a negative predictor of ICI

response is an unexpected finding, considering that higher frequencies of TRM cells in lung

tumor tissues are generally associated with better clinical outcomes in NSCLC [36]. This is

largely due to their role in sustaining high densities of tumor-infiltrating lymphocytes and

promoting anti-tumor responses. Additionally, previous studies have demonstrated that TRM

cell subsets coexpressing PD-1 and TIM-3 are relatively enriched in patients who respond to

PD-1 inhibitors [37]. However, recent findings suggest that pre-existing TRM-like cells in lung

cancer may promote immune evasion mechanisms, contributing to resistance to immune

checkpoint blockade therapies [38]. These observations suggest that the roles of TRM subsets in

tumor immunity are highly context-dependent.

Similarly, CD4+ TH17 cells, which were overrepresented in the non-responder groups,

exhibit context-dependent roles in tumor immunity and may be associated with both

unfavorable and favorable outcomes [39,40]. In exploring tumor cell signatures linked to ICI

response, non-responder attributes were regulated by STAT3 and NFKB1. The STAT3 and

NF-κB pathways are crucial for Th17 cell differentiation and T cell activation [41,42]. Notably,

STAT3 activation in lung cancer orchestrates immunosuppressive characteristics by

inhibiting T-cell mediated cytotoxicity [43]. The combined influence of the Th17/STAT3 axis

and TRM cell activity in predicting ICI response underscores the complexity of these

pathways and suggests that their roles in tumor immunity and therapy response warrants

further investigation.

In search of tumor cell signatures associated with the response to ICI, we adopted two

approaches, an individual gene level comparison between the responder and non-responder

group and a feature extraction approach to decompose data using the NMF and PCA. Both

approaches highlighted attributes of non-responders governed by key transcription factors,

which play a significant role in immune response regulation. The ability to predict ICI

response based on tumor signatures was as accurate as predictions based on immune cell

behavior. Integrating data from both immune and tumor cells enhanced the discriminative

power (AUC) for identifying responder, suggesting the presence of both interactive and

distinct mechanisms of resistance.

Our study has limitations. Primarily, most samples were obtained from metastatic lymph

nodes rather than original tumor tissues, potentially not reflecting the tumor

microenvironment accurately. However, prior study [17] and Fig. S1 have shown that the

immune microenvironment within metastatic lymph nodes closely resembles that of lung

tumor tissues, rather than normal lymph nodes. On a positive note, our findings indicate that

the immune landscape of metastatic lymph nodes can predict ICI response. Another challenge

is the small sample size and the issue of gene expression drop-out, necessitating further

studies with a larger patient cohort. Despite these limitations, our study stands out by

employing high-throughput scRNA-seq to both tumor cells and immune microenvironment,

offering a comprehensive analysis of the multicellular factors that affect ICI response in

patients with advanced NSCLC.

Materials and Methods

Human specimens. This study was approved by the Institutional Review Board (IRB) of Samsung Medical Center (IRB no. 2010-04-039-052). Informed written consent was obtained from all patients enrolled in the study. The study participants included 26 patients diagnosed with lung cancer (Table S1). The study population (n=26) has been treated with the investigator's choice either as a clinical trial (n=5) or as standard clinical practice (n=21). Regardless of the treatment selection, the specimens were prospectively collected based on the study protocol. A total of 33 samples were collected and immediately transferred on ice for tissue preparation. Metastatic lymph nodes, metastatic liver tissues, and lung/bronchus tumor tissues from patients with lung cancer were collected using endobronchial ultrasound bronchoscopy, neck lymph node ultrasound and biopsy, liver biopsy, and percutaneous transthoracic cutting needle biopsy. Tumors, normal lungs, normal lymph nodes, and normal brain tissues were obtained during resection surgery. Pleural fluid was collected from patients with malignant pleural effusion.

Clinical outcomes. The clinical outcomes of ICI were evaluated based on the Response Evaluation Criteria in Solid Tumor (RECIST) 1.1 [44]. In this study, we described non-responders as patients with a stable or progressive disease. Responders were considered as patients with a partial response. None of the patients showed a complete response.

Sample preparation. Single-cell isolation was performed differently depending on the samples. (1) Biopsy samples and normal lymph node tissues were chopped into 2-4 mm pieces and dissociated in an enzyme solution containing collagenase/hyaluronidase (STEMCELL Technologies, Vancouver, Canada) and DNase I, RNase-Free (lyophilized)

(QIAGEN, Hilden, Germany) at 37°C for 1 h. Tissue pieces were re-mixed by gentle

pipetting at 20-min intervals during incubation. (2) Tumor and normal lung tissue

dissociation was performed using a tumor dissociation kit (Miltenyi Biotech, Germany)

following the manufacturer's instructions. Briefly, tissue was cut into 2–4 mm pieces and

transferred to a C tube containing the enzyme mix (enzymes H, R, and A in RPMI1640

medium). The GentleMACS programs h_tumor_01, h_tumor_02, and h_tumor_02 were run

with two 30-min incubations on a MACSmix tube rotator at 37°C. (3) Brain tissue was

chopped into 2–4 mm pieces and incubated in an enzyme solution (collagenase (Gibco,

Waltham, MA, USA), DNase I (Roche, Basel, Switzerland), and Dispase I (Gibco) in DMEM)

at 37°C for 1 h. Tissue pieces were re-mixed by gentle pipetting at 15-min intervals during

incubation. (4) Pleural fluids were transferred to a 50-ml tube, and the cells were spun down

at 300g.

Each cell suspension was transferred to a new 50-ml (15-ml for biopsy samples) tube through

a 70-µm strainer. The volume in the tube was readjusted to 50-ml (or 15-ml) with RPMI1640

medium, and spun down to remove the enzymes. The supernatant was aspirated, the cell

pellet was resuspended in 4 ml of RPMI1640 medium, and dead cells were removed using

Ficoll-Paque PLUS (GE Healthcare, Chicago, IL, USA) separation.

For samples subjected to multiplexing, dissociated cells were cryopreserved in

CELLBANKER1 (Zenogen, Fukushima, Japan) and thawed for pooling.

Single-cell RNA sequencing (scRNA-seq) and read processing. Single-cell suspensions

were loaded into a Chromium system (10x Genomics, Pleasanton, CA, USA). Following the

manufacturer's instructions, 3' scRNA-seq libraries for the 14 samples were generated using

Chromium Single Cell 3′ v2 Reagent Kits. The 3' library preparation for EBUS_119 used

Chromium Single Cell 3ʹ v3 Reagent Kits. The 5' scRNA-seq libraries for twelve individual

and two pooled samples were generated using Chromium Single Cell 5ʹ v2 Reagent Kit.

Libraries were then sequenced on an Illumina HiSeq 2500 for 3' scRNA-seq and an Illumina

NovaSeq 6000 for 5' scRNA-seq. Sequencing reads were mapped to the GRCh38 human

reference genome using Cell Ranger toolkit (v5.0.0).

SNP genotyping array. Genomic DNA was extracted from the peripheral blood of six

patients and subjected to sample multiplexing (DNeasy Blood & Tissue Kit, QIAGEN). The

766,221 single nucleotide polymorphisms (SNPs) was genotyped using Illumina Global

Screening Array MG v2, following the manufacturer's instructions. Normalized signal

intensity and genotype were processed using Illumina's GenomeStudio v.2 software.

Demultiplexing of pooled samples. The individuals in sample multiplexing were assigned

by a software tool *freemuxlet*, which is an extension of *demuxlet* [45]

(https://github.com/statgen/popscle). First, the popscle tool *dsc-pileup* was run with the bam

file generated by Cell Ranger toolkit and reference vcf file. The reference was assembled

after a lift-over process with GRCh38 from 1000 Genomes Project phase 1 data and the

variant allele frequency in East Asian >0.01 were discarded. Next, *freemuxlet* was used to

determine the sample identity with default parameters. The individuals were matched based

on the similarity between freemuxlet-annotated genotypes and SNP array-detected genotypes.

Acquisition of scRNA-seq data from lung adenocarcinoma (LUAD) patients. We

obtained raw 3' scRNA-seq from 43 specimens acquired from 33 LUAD patients including

early-stage (tLung) and late-stage (tL/B) lung tumor tissues, metastatic lymph nodes (mLN),

normal lung tissues (nLung) and lymph nodes (nLN) [17]. Sequencing reads were mapped to

the GRCh38 human reference genome using Cell Ranger toolkit (v5.0.0).

scRNA-seq data analysis. The raw gene-cell-barcode matrix from Cell Ranger pipeline was

processed using Seurat v3.2.2 R package [46]. Cells were selected using two quality criteria:

mitochondrial genes (<20%) and gene count (>200). Cell multiplets predicted by Scrublet [47]

were filtered out. From the filtered cells, the unique molecular identifier (UMI) count matrix

was log-normalized and scaled by z-transform while regressing out the effects of cell-cycle

variations for subsequent analysis. For batch correction, we used Harmony v1.0 R package [48]

interfacing with Seurat as the *RunHarmony* function. A total of 2,000 variably expressed

genes were selected using *FindVariableFeatures* with a parameter selection.method="vst". A

subset of principal components (PCs) was selected based on *ElbowPlot* function. Uniform

Manifold Approximation and Projection (UMAP) for dimension reduction and cell clustering

was performed using *RunUMAP, FindNeighbors*, and *FindClusters* functions with the

selected PCs and resolutions [Advanced lung cancer patients (Total cells, 33 PCs and

resolution=0.3; CD4+ T cells, 28 PCs and resolution=0.9; CD8+ T cells, 28 PCs and

resolution=0.9; NK cells, 26 PCs and resolution=0.3; B/Plasma cells, 28 PCs and

resolution=0.3; Myeloid cells, 30 PCs and resolution=1.2), LUAD patients (Total cells, 23

PCs and resolution=0.3; T/NK cells, 24 PCs and resolution=0.9)]. We applied the

FindAllMarkers function to identify differentially expressed genes (DEGs) for each cell

cluster. Significance was determined using Wilcoxon Rank Sum test. Genes were selected

according to the following statistical thresholds; log fold change>0.25, p-value<0.01,

adjusted p-value (bonferroni correction)<0.01, and percentage of cells (pct)>0.25. Cell

identity was determined by comparing the expression of known marker genes and DEGs for

each cluster.

Principal component analysis (PCA) analysis using the proportion of cell lineages and

T/NK cell subsets. PCA analysis was performed for the % proportion of cell lineages and

T/NK cell subsets in individual LUAD samples using *prcomp* function of stats v3.6.3 R

package. For total cells, the percentages of immune and stromal cells were calculated except

for epithelial, cycling, and AMB (ambiguous) cells. For T/NK cells, unknown cells annotated

as MT high and AMB cells were excluded.

In silico classification of CD4+ T, CD8+ T, and NK cells. We characterized CD4+ T,

CD8+ T, and NK cell populations by combined analysis of gene and protein expression using

Cellular Indexing of Transcriptomes and Epitopes by Sequencing data from primary tumor

and normal lungs. Among the cells in clusters annotated as T/NK cells, we identified CD3-

expressing cells with CD3D or CD3E or CD3G >0 at the RNA level. CD4 and CD8 positive

cells were then identified with a cutoff at 55th percentile of antibody-derived tags (ADT)

level. NK cells were identified based on the RNA expression level of NK cell markers (*XCL1*,

NCAM1, *KLRD1*, and *KLRF1*) in CD3 negative cells. The gene expression matrix with cell

identity of CD4 positive, CD8 positive, and NK was applied as reference data for supervised

cell-type classification using *getFeatureSpace* and *trainModel* functions of scPred v1.9.0 R

package [49]. Finally, we classified T/NK cells in Fig. 1b into CD4+ T, CD8+ T, and NK cells

using scPred *scPredict* function.

Analysis of TCR/BCR repertoires in CD4+ T, CD8+ T, and B/Plasma cells. The data

derived from Cell Ranger pipeline for T cell receptor (TCR) and B cell receptor (BCR)

sequencing data were processed using scRepertoire v1.2.0 R package [50] in R v4.1.1. We

selected contigs that generated alpha-beta chain pairs for TCR and heavy-light chain pairs for

BCR for subsequent analysis. We called clonotypes based on V(D)JC genes and CDR3

nucleotide sequence with the parameter clonecall="gene+nt". The set of clone types was

classified by total frequency using the parameter cloneTypes defined as Single=1, Small=5,

Medium=10, Large=20, and Hyperexpanded=Inf.

Scoring of T cell functional features. Scores for T cell functional features were calculated

as the mean expression of regulatory (*ICOS*, *FOXP3*, *IKZF2*, *LAYN*, *TNFRSF18*, *CTLA4*,

IL21R, *BATF*, *CCR8*, *IL2RA*, and *TNFRSF4*) and cytotoxic (*CX3CR1*, *PRF1*, *GZMA*, *GZMB*,

GZMH, *GNLY*, *KLRG1*, and *NKG7*) genes at the log-normalized level.

Identification of malignant cells based on inferred copy number variation (CNV) from

scRNA-seq   **data.**   Two   computational   tools,   inferCNV   v1.2.1

(https://github.com/broadinstitute/inferCNV) and CopyKAT v1.0.5 R packages [51], were used

to infer genomic copy numbers from scRNA-seq. In a run with inferCNV, the UMI count

matrix of each tumor sample was loaded into inferCNV *CreateInfercnvObject* function along

with cell lineage annotations. The reference (normal) cells were selected as cells annotated

with T/NK, B/Plasma, myeloid, and mast cells. We maintained the proportion of epithelial

cells below 20% in each tumor sample using the expression profiles of the normal lung and

lymph node tissues. Inferred CNV signals were analyzed using inferCNV *run* function using

the parameters: cutoff=0.1, denoise=TRUE, HMM=TRUE, and HMM_type="i6". The

signals were then summarized as standard deviations (s.d.) for all windows and the

correlation between the CNV in each cell and the mean of the top 5% cells [52]. Cancer cells

showing CNV perturbation (>0.03 s.d. or >0.3 CNV correlation) were classified as malignant

cells, otherwise as non-malignant cells. The UMI count matrix of each tumor sample was

loaded into CopyKAT *copykat* function along with cell lineage annotations using the

following parameters: ngene.chr=3, KS.cut=0.05, and norm.cell.names. Cancer cells

predicted as aneuploid cells by CopyKAT were classified as malignant cells. Finally, we

identified malignant cells, which are cancer cells classified as malignant cells in either

inferCNV or CopyKAT.

Single-cell DEGs between response groups in malignant cells. A total of 12,975 malignant

cells were used to identify DEGs in pairwise comparisons according to responder versus non-

responder, PR versus PD, and PR versus SD. Differential expression levels were calculated

using Seurat *FindMarkers* function with the Wilcoxon Rank Sum test. Genes were selected

according to the following statistical thresholds: log fold change>0.25, p-value<0.01,

adjusted p-value (bonferroni correction)<0.01, and pct>0.25. We reconstructed DEGs as four

groups: INT.up, INT.down, UNION,up, and UNION.down, based on with the intersection

(INT) and union (UNION) of up- or down-regulated genes for pairwise comparisons between

responder versus non-responder, PR versus PD, and PR versus SD. INT.up and INT.down

represent the intersection of up- and down-regulated genes in the responder group,

respectively. UNION.up and UNION.down represent the union of up- and down-regulated

genes in the responder group, respectively.

Non-negative matrix factorization (NMF) programs of the malignant cells. The UMI

count matrix for malignant cells was loaded into *nmf* function of RcppML v0.5.6 R package.

A NMF model was learned with a rank of 30 using all genes. For each of the 30 NMF factors,

the top-ranked 50 genes in the NMF score were defined as signatures. RECIST-enriched

NMF program consisted of selected factors based on their relative sum of loadings. We

aggregated and redefined gene signatures of factors included in each NMF program. The

uniqueness of each NMF program for RECIST groups was evaluated as an odds ratio using

fisher.test function of stats v3.6.3 R package. Annotations of NMF programs were assigned

using Metascape [53].

RECIST-specific gene modules in malignant cells. The RECIST-specific gene modules

were analyzed with a matrix factorization named scINSIGHT [24] using log-normalized count

and 2,000 highly variable genes for each sample. For each module, we selected the 100 genes

with the highest coefficients. Combinations of the top 100 genes for modules specific to each

RECIST group were defined as module genes. The uniqueness of each module for RECIST

groups was evaluated as an odds ratio using *fisher.test* function of stats v3.6.3 R package.

Annotations of gene modules were assigned using Metascape [53].

Principal component signatures of the malignant cells. The UMI count matrix for

malignant cells was log-normalized and scaled by z-transform while regressing out the effects

of cell-cycle variations for PCA. A total of 2,000 variably expressed genes selected using

FindVariableFeatures with selection.method="vst" were used for PCA. PCs were calculated

by Seurat *RunPCA* function. PC signatures were selected for 30 genes with + (pos) and −

(neg) scores that highly contributed to each PC from PC1 to PC10.

Functional category analysis. Functional categories representing the enriched gene

expression in comparisons for responder vs. non-responder, PR vs. PD, and PR vs. SD as

well as in the PCs were identified using fgsea v1.12.0 [54] R package with parameters:

minSize=10, maxSize=600, and nperm=10000. Gene sets for Gene Ontology (GO) Biological

Process were collected from the MSigDB database using msigdbr v7.1.1 R package [55,56]. The

gene list was ranked by the log fold change for each comparison and feature loadings for each

PC. Significant GO terms were selected after collapsing redundant terms using fgsea

collapsePathways function with a statistical threshold for Benjamini-Hochberg adjusted p-

value<0.05.

Multivariate overall survival analysis of tumor signatures. To evaluate the prognostic

potential of tumor signatures, RNA sequencing data from LUAD and LUSC samples were

retrieved from The Cancer Genome Atlas (TCGA) data portal (https://portal.gdc.cancer.gov/)

[57]. The dataset comprised 533 primary tumors from TCGA LUAD and 502 primary tumors

from TCGA LUSC. Gene expression levels were quantified as (log2 FPKM-UQ + 1), where

FPKM-UQ represents the upper quartile fragments per kilobase per million mapped reads for

each sample.

Clinical variables were first categorized, including age (below and above median age), gender

(male and female), pathological stage (I/II and III/IV), distant metastasis (M0 and M1), and

nodal status (N0 and Ns). Samples were then classified into high and low groups based on the

upper quantile of the mean expression for each tumor signature group. Multivariate survival

analysis was conducted using the *analyse_multivariate* function of the survivalAnalysis R

package.

Evaluation of discriminative power of identifying responders for tumor signatures and

combinatorial indexes. Classification models of responders and non-responders for PC

signatures and combinatorial indexes between tumor and/or immune cells were generated

based on in-sample performance and tested by Receiver Operating Characteristic (ROC)

curve. Relative numbers between the observed and expected cells (*Ro/e*) for each sample

were obtained from the chi-square test [19]. To describe the separability, area under the curve

(AUC) was calculated using *ROC* function of Epi v2.44 R package with *Ro/e* scores as input.

Significance was calculated by Wilcoxon Rank Sum test and confirmed by adjusting with the

Benjamini-Hochberg correction.

Univariate regression analysis for ICI response. Univariate regression was performed

using the *lm* function of stats v3.6.3 with *Ro/e* scores for each sample as input. We evaluated

the relationship between the target variable ICI response, classified as responders and non-

responders, and one predictor variable of immune cell types, tumor signatures, and clinical

factors such as tissue origin, cancer subtype, pathological stage, and smoking status. The

significance of predictor was calculated using *Anova* function of car v3.0-9 R package.

Validation of PC signatures in melanoma cohorts. We used Riaz et al.'s [29] and Van Allen

al.'s [28] RNA sequencing data from melanoma patients receiving PD-1 and CTLA-4 immune

checkpoint therapy to assess expressional changes of PC signatures along RECIST. The mean

expression of each PC signature in each RECIST group was calculated as the log2

normalized level.

Data availability. Raw single-cell RNA sequencing data generated during the current study

are available in the European Genome-phenome Archive (EGA) database (accession code

EGAD00001008703), and processed data can be accessed from the NCBI Gene Expression

Omnibus (GEO) database (accession code GSE205335).

Single-cell RNA sequencing data for LUAD patients analyzed in this study are available in

the EGA database at EGAD00001005054. Bulk RNA sequencing data analyzed in this study

were obtained from GEO at GSE91061 and database of Genotypes and Phenotypes (dbGap)

at phs000452.v2.p1.

Acknowledgements

This study was supported by the Collaborative Genome Program for Fostering New Post-

Genome Industry (NRF-2017M3C9A6044633 and NRF-2017M3C9A6044636), Mid-Career

Researcher Program (NRF-2022R1A2C1091451), and Basic Research Laboratory Program

(RS-2023-00220840) of the National Research Foundation of Korea funded by the Korea

government. We also acknowledge the Basic Medical Science Facilitation Program, through

the Catholic Medical Center of the Catholic University of Korea funded by the Catholic

Education Foundation and the KREONET/GLORIAD service provided by KISTI (Korea

Institute of Science and Technology Information).

References

1   Reck, M. *et al.* Pembrolizumab versus Chemotherapy for PD-L1-Positive Non-Small-

Cell Lung Cancer. *N Engl J Med* **375**, 1823-1833, doi:10.1056/NEJMoa1606774

(2016).

2   Gandhi, L. *et al.* Pembrolizumab plus Chemotherapy in Metastatic Non-Small-Cell

Lung Cancer. *N Engl J Med* **378**, 2078-2092, doi:10.1056/NEJMoa1801005 (2018).

3   Paz-Ares, L. *et al.* Pembrolizumab plus Chemotherapy for Squamous Non-Small-Cell

Lung Cancer. *N Engl J Med* **379**, 2040-2051, doi:10.1056/NEJMoa1810865 (2018).

4   Wakelee H, A. N., Zhou C, Csőszi T, Vynnychenko IO, Goloborodko O, Luft A,

Andrey Akopov, Martinez-Marti A, Kenmotsu H, Chen Y, Chella A, Sugawara S,

Gitlitz BJ, Bennett E, Wu F, Yi J, Deng Y, McCleland M, Felip E. IMpower010:

Primary results of a phase III global study of atezolizumab versus best supportive care

after adjuvant chemotherapy in resected stage IB-IIIA non-small cell lung cancer

(NSCLC). *J Clin Oncol* **39**, 8500, doi:10.1200/JCO.2021.39.15_suppl.8500 (2021).

5   Reck, M. *et al.* Five-Year Outcomes With Pembrolizumab Versus Chemotherapy for

Metastatic Non-Small-Cell Lung Cancer With PD-L1 Tumor Proportion Score >/= 50.

J Clin Oncol **39**, 2339-2349, doi:10.1200/JCO.21.00174 (2021).

6   Fehrenbacher, L. *et al.* Atezolizumab versus docetaxel for patients with previously

treated non-small-cell lung cancer (POPLAR): a multicentre, open-label, phase 2

randomised controlled trial. *Lancet* **387**, 1837-1846, doi:10.1016/S0140-

6736(16)00587-0 (2016).

7   Hellmann, M. D. *et al.* Nivolumab plus Ipilimumab in Lung Cancer with a High

Tumor Mutational Burden. *N Engl J Med* **378**, 2093-2104,

doi:10.1056/NEJMoa1801946 (2018).

8   Litchfield, K. *et al.* Meta-analysis of tumor- and T cell-intrinsic mechanisms of

sensitization to checkpoint inhibition. *Cell* **184**, 596-614 e514,

doi:10.1016/j.cell.2021.01.002 (2021).

9   Kamphorst, A. O. *et al.* Proliferation of PD-1+ CD8 T cells in peripheral blood after

PD-1-targeted therapy in lung cancer patients. *Proc Natl Acad Sci U S A* **114**, 4993-

4998, doi:10.1073/pnas.1705327114 (2017).

10   Kim, K. H. *et al.* The First-week Proliferative Response of Peripheral Blood PD-

1(+)CD8(+) T Cells Predicts the Response to Anti-PD-1 Therapy in Solid Tumors.

Clin Cancer Res **25**, 2144-2154, doi:10.1158/1078-0432.CCR-18-1449 (2019).

11   Thommen, D. S. *et al.* A transcriptionally and functionally distinct PD-1(+) CD8(+) T

cell pool with predictive potential in non-small-cell lung cancer treated with PD-1

blockade. *Nat Med* **24**, 994-1004, doi:10.1038/s41591-018-0057-z (2018).

12   Simon, S. & Labarriere, N. PD-1 expression on tumor-specific T cells: Friend or foe

for   immunotherapy?   *Oncoimmunology*   **7**,   e1364828,

doi:10.1080/2162402X.2017.1364828 (2017).

13   Arce Vargas, F. *et al.* Fc-Optimized Anti-CD25 Depletes Tumor-Infiltrating

Regulatory T Cells and Synergizes with PD-1 Blockade to Eradicate Established

Tumors. *Immunity* **46**, 577-586, doi:10.1016/j.immuni.2017.03.013 (2017).

14   Krieg, C. *et al.* Author Correction: High-dimensional single-cell analysis predicts

response to anti-PD-1 immunotherapy. *Nat Med* **24**, 1773-1775, doi:10.1038/s41591-

018-0094-7 (2018).

15   Kumagai, S. *et al.* The PD-1 expression balance between effector and regulatory T

cells predicts the clinical efficacy of PD-1 blockade therapies. *Nat Immunol* **21**, 1346-

1358, doi:10.1038/s41590-020-0769-3 (2020).

16   Ruiz-Patino, A. *et al.* Immunotherapy at any line of treatment improves survival in

patients with advanced metastatic non-small cell lung cancer (NSCLC) compared

with chemotherapy (Quijote-CLICaP). *Thorac Cancer* **11**, 353-361,

doi:10.1111/1759-7714.13272 (2020).

17   Kim, N. *et al.* Single-cell RNA sequencing demonstrates the molecular and cellular

reprogramming of metastatic lung adenocarcinoma. *Nat Commun* **11**, 2285,

doi:10.1038/s41467-020-16164-1 (2020).

18   Stoeckius, M. *et al.* Simultaneous epitope and transcriptome measurement in single

cells. *Nat Methods* **14**, 865-868, doi:10.1038/nmeth.4380 (2017).

19   Guo, X. *et al.* Global characterization of T cells in non-small-cell lung cancer by

single-cell sequencing. *Nat Med* **24**, 978-985, doi:10.1038/s41591-018-0045-3 (2018).

20   Gueguen, P. *et al.* Contribution of resident and circulating precursors to tumor-

infiltrating CD8(+) T cell populations in lung cancer. *Sci Immunol* **6**,

doi:10.1126/sciimmunol.abd5778 (2021).

21   Groom, J. R. & Luster, A. D. CXCR3 in T cell function. *Exp Cell Res* **317**, 620-631,

doi:10.1016/j.yexcr.2010.12.017 (2011).

22   Yang, C. *et al.* Heterogeneity of human bone marrow and blood natural killer cells

defined by single-cell transcriptome. *Nat Commun* **10**, 3931, doi:10.1038/s41467-019-

11947-7 (2019).

23   Liu, L. *et al.* Combination of TMB and CNA Stratifies Prognostic and Predictive

Responses to Immunotherapy Across Metastatic Cancer. *Clin Cancer Res* **25**, 7413-

7423, doi:10.1158/1078-0432.CCR-19-0558 (2019).

24   Qian, K., Fu, S., Li, H. & Li, W. V. scINSIGHT for interpreting single-cell gene

expression from biologically heterogeneous data. *Genome Biol* **23**, 82,

doi:10.1186/s13059-022-02649-3 (2022).

25   Lapuente-Santana, O., van Genderen, M., Hilbers, P. A. J., Finotello, F. & Eduati, F.

Interpretable systems biomarkers predict response to immune-checkpoint inhibitors.

Patterns (N Y) **2**, 100293, doi:10.1016/j.patter.2021.100293 (2021).

26   Betzler, A. C. *et al.* NF-kappaB and Its Role in Checkpoint Control. *Int J Mol Sci* **21**,

doi:10.3390/ijms21113949 (2020).

27   Fenton, S. E., Saleiro, D. & Platanias, L. C. Type I and II Interferons in the Anti-

Tumor Immune Response. *Cancers (Basel)* **13**, doi:10.3390/cancers13051037 (2021).

28   Van Allen, E. M. *et al.* Genomic correlates of response to CTLA-4 blockade in

metastatic melanoma. *Science* **350**, 207-211, doi:10.1126/science.aad0095 (2015).

29   Riaz, N. *et al.* Tumor and Microenvironment Evolution during Immunotherapy with

Nivolumab. *Cell* **171**, 934-949 e916, doi:10.1016/j.cell.2017.09.028 (2017).

30   Lawrence, M. S. *et al.* Mutational heterogeneity in cancer and the search for new

cancer-associated genes. *Nature* **499**, 214-218, doi:10.1038/nature12213 (2013).

31   Sharma, P., Hu-Lieskovan, S., Wargo, J. A. & Ribas, A. Primary, Adaptive, and

Acquired   Resistance   to   Cancer   Immunotherapy.   *Cell*   **168**,   707-723,

doi:10.1016/j.cell.2017.01.017 (2017).

32   Gibellini, L. *et al.* Single-Cell Approaches to Profile the Response to Immune

Checkpoint Inhibitors. *Front Immunol* **11**, 490, doi:10.3389/fimmu.2020.00490

(2020).

33   Zheng, L. *et al.* Pan-cancer single-cell landscape of tumor-infiltrating T cells. *Science*

374, abe6474, doi:10.1126/science.abe6474 (2021).

34   Koh, J. *et al.* Regulatory (FoxP3(+)) T cells and TGF-beta predict the response to

anti-PD-1 immunotherapy in patients with non-small cell lung cancer. *Sci Rep* **10**,

18994, doi:10.1038/s41598-020-76130-1 (2020).

35   Wei, S. C. *et al.* Combination anti-CTLA-4 plus anti-PD-1 checkpoint blockade

utilizes cellular mechanisms partially distinct from monotherapies. *Proc Natl Acad*

Sci U S A **116**, 22699-22709, doi:10.1073/pnas.1821218116 (2019).

36   Ganesan, A. P. *et al.* Tissue-resident memory features are linked to the magnitude of

cytotoxic T cell responses in human lung cancer. *Nat Immunol* **18**, 940-950,

doi:10.1038/ni.3775 (2017).

37   Clarke, J. *et al.* Single-cell transcriptomic analysis of tissue-resident memory T cells

in human lung cancer. *J Exp Med* **216**, 2128-2149, doi:10.1084/jem.20190249 (2019).

38   Weeden, C. E. *et al.* Early immune pressure initiated by tissue-resident memory T

cells sculpts tumor evolution in non-small cell lung cancer. *Cancer Cell* **41**, 837-852

e836, doi:10.1016/j.ccell.2023.03.019 (2023).

39   Marques, H. S. *et al.* Relationship between Th17 immune response and cancer. *World*

J Clin Oncol **12**, 845-867, doi:10.5306/wjco.v12.i10.845 (2021).

40   Chang, S. H. T helper 17 (Th17) cells and interleukin-17 (IL-17) in cancer. *Arch*

Pharm Res **42**, 549-559, doi:10.1007/s12272-019-01146-9 (2019).

41   Park, S. H., Cho, G. & Park, S. G. NF-kappaB Activation in T Helper 17 Cell

Differentiation. *Immune Netw* **14**, 14-20, doi:10.4110/in.2014.14.1.14 (2014).

42   Poholek, C. H. *et al.* Noncanonical STAT3 activity sustains pathogenic Th17

proliferation and cytokine response to antigen. *J Exp Med* **217**,

doi:10.1084/jem.20191761 (2020).

43   Jing, B. *et al.* IL6/STAT3 Signaling Orchestrates Premetastatic Niche Formation and

Immunosuppressive Traits in Lung. *Cancer Res* **80**, 784-797, doi:10.1158/0008-

5472.CAN-19-2013 (2020).

44   Eisenhauer, E. A. *et al.* New response evaluation criteria in solid tumours: revised

RECIST guideline (version 1.1). *Eur J Cancer* **45**, 228-247,

doi:10.1016/j.ejca.2008.10.026 (2009).

45   Kang, H. M. *et al.* Multiplexed droplet single-cell RNA-sequencing using natural

genetic variation. *Nat Biotechnol* **36**, 89-94, doi:10.1038/nbt.4042 (2018).

46   Stuart, T. *et al.* Comprehensive Integration of Single-Cell Data. *Cell* **177**, 1888-1902

e1821, doi:10.1016/j.cell.2019.05.031 (2019).

47   Wolock, S. L., Lopez, R. & Klein, A. M. Scrublet: Computational Identification of

Cell Doublets in Single-Cell Transcriptomic Data. *Cell Syst* **8**, 281-291 e289,

doi:10.1016/j.cels.2018.11.005 (2019).

48   Korsunsky, I. *et al.* Fast, sensitive and accurate integration of single-cell data with

Harmony. *Nat Methods* **16**, 1289-1296, doi:10.1038/s41592-019-0619-0 (2019).

49   Alquicira-Hernandez, J., Sathe, A., Ji, H. P., Nguyen, Q. & Powell, J. E. scPred:

accurate supervised method for cell-type classification from single-cell RNA-seq data.

Genome Biol **20**, 264, doi:10.1186/s13059-019-1862-5 (2019).

50   Borcherding, N., Bormann, N. L. & Kraus, G. scRepertoire: An R-based toolkit for

single-cell immune receptor analysis. *F1000Res* **9**, 47,

doi:10.12688/f1000research.22139.2 (2020).

51   Gao, R. *et al.* Delineating copy number and clonal substructure in human tumors from

single-cell transcriptomes. *Nat Biotechnol* **39**, 599-608, doi:10.1038/s41587-020-

00795-2 (2021).

52   Puram, S. V. *et al.* Single-Cell Transcriptomic Analysis of Primary and Metastatic

Tumor Ecosystems in Head and Neck Cancer. *Cell* **171**, 1611-1624 e1624,

doi:10.1016/j.cell.2017.10.044 (2017).

53   Zhou, Y. *et al.* Metascape provides a biologist-oriented resource for the analysis of

744   systems-level datasets. *Nat Commun* **10**, 1523, doi:10.1038/s41467-019-09234-6

745   (2019).

746   54   Korotkevich G, S. V., Sergushichev A. Fast gene set enrichment analysis. *bioRxiv*,

747   doi:10.1101/060012 (2019).

748   55   Liberzon, A. *et al.* Molecular signatures database (MSigDB) 3.0. *Bioinformatics* **27**,

749   1739-1740, doi:10.1093/bioinformatics/btr260 (2011).

750   56   Subramanian, A. *et al.* Gene set enrichment analysis: a knowledge-based approach for

751   interpreting genome-wide expression profiles. *Proc Natl Acad Sci U S A* **102**, 15545-

752   15550, doi:10.1073/pnas.0506580102 (2005).

753   57   Grossman, R. L. *et al.* Toward a Shared Vision for Cancer Genomic Data. *N Engl J*

754   *Med* **375**, 1109-1112, doi:10.1056/NEJMp1607591 (2016).

 Figure legends

 Figure 1. Cell lineage identification of 96,505 single cells from 26 patients with lung

 cancer treated with ICI. a, Workflow of sample collection and single-cell analysis of lung

 cancer patients treated with ICI. **b,** UMAP plot of 96,505 single cells from 33 samples

 acquired from 26 advanced lung cancer patients, colored by clusters. AMB cells, Ambiguous

 cells. **c,** Dot plot of mean expression of canonical marker genes for cell lineages. **d,**

 Proportions of the cell lineages in NSCLC tissue from core patients shown by individual

 samples aligned with clinical data. Labels for origins indicate LN, Metastatic lymph node;

 Lung, Tumor lung; PE, Malignant pleural effusion; Liver, Metastatic liver. **e,** Box plot of the

 percentage of cell lineages in responder and non-responder groups. Label represents p-value

 calculated via two-tailed Student's t-test. Each box represents the median and the

 interquartile range (IQR, the range between the 25th and 75th percentile), whiskers indicate

 the 1.5 times of IQR.

 Figure 2. Classification and characterization of CD4+ and CD8+ T cell subtypes. a,

 Prediction strategy to classify CD4+, CD8+ T, and NK cells by applying ADT data from lung

 cancer. **b,** UMAP plot of CD4+ and CD8+ T cells, colored by clusters. **c,** Dot plot of mean

 expression of selected CD4+ (left) and CD8+ (right) T cell marker genes in each cell cluster.

 d, Box plot of the percentage of CD4+ and CD8+ T cell types within total CD4+ plus CD8+

 cells for sample groups representing responses to ICI. Label represents p-value calculated via

 two-tailed Student's t-test. Each box represents the median and the IQR, whiskers indicate

 the 1.5 times of IQR. **e,** Association of T cell functional features with clonal expansion. Dot

 size depicts the relative clone size of each cell, which is divided by the total number of CD4+

 and CD8+ T cells, respectively. Color indicates the cell lineage. **f,** Comparisons of

proportional changes in cell subtypes along ICI responses within each immune cell lineage.

The quantitative values shown on the axis represent the mean difference in % cell proportions

between sample groups. Dot size and color represent -log (p-value) for responder vs. non-

responder and non-PD vs. PD, respectively. The lower left quadrant shows cell types

overrepresented in the poor responder groups, while the upper right quadrant indicates cell

types overrepresented in the better responder groups. p-value, two-tailed Student's t-test.

Figure 3. Systemic evaluation of immune cell dynamics associated with response to ICI.

a, Heat map with unsupervised hierarchical clustering (left) and depicting significance (right)

of proportional changes in cell subtypes within total immune cells. Proportional changes were

compared for multiple ICI response groups. Color represents the -log (p-value) determined

using two-tailed Student's t-test. **b,** Distribution map for each cell type across individual

samples aligned with clinical data. Color represents *Ro/e* score calculated using the chi-

square test.

Figure 4. Single-cell tumor signatures associated with response to ICI. a, Volcano plot of

expression difference for responder vs. non-responder, PR vs. PD, and PR vs. SD in 12,975

malignant cells from 11 core patients. The log fold change indicates the difference in the

mean expression level for each gene. The significance level was determined using two-tailed

Wilcoxon Rank Sum test. **b,** Relative sum of loadings for all NMF factors contributed to

malignant cells from 11 core patients across RECIST, tissue origins, and cancer subtypes,

respectively. **c,** Selection of RECIST-enriched NMF programs. **d,** Enrichment of NMF

programs for RECIST groups. Color represents the z-transformed odds ratio. **e,** Expression

map of RECIST-specific scINSIGHT modules across individual samples aligned with clinical

data. Color represents the z-transformed mean expression of genes contributing to each

module. **f,** Enrichment of RECIST-specific gene modules for RECIST groups. Color

represents the z-transformed odds ratio. **g,** Hierarchical clustering of pairwise similarities

between tumor signatures. INT and UNION, intersection and union of DEGs for responder vs.

non-responder, PR vs. PD, and PR vs. SD in Fig. 4a. **h,** Functional categories and **i,**

transcription factors of the selected tumor signatures, analyzed by Metascape.

**Figure 5. Combination of tumor signatures and immune index classifying the response
to ICI. a,** Heat map of relative contribution of tumor signatures and immune index across

individual samples aligned with clinical data. Mean expression of each tumor signature and

the percentage of each immune index are divided by the maximum value across samples. INT

and UNION, intersection and union of DEGs for responder vs. non-responder, PR vs. PD,

and PR vs. SD in Fig. 4a. **b,** Univariate regression analysis of immune and tumor signatures

for ICI response, together with clinical variables. **c,** ROC analysis of combinatorial index to

classify responder and non-responder. p-value, two-tailed Wilcoxon Rank Sum test. P-values

adjusted with the Benjamini-Hochberg correction; UNION.down + B/Plasma, 0.22;

INT.down + B/Plasma, 0.22; UNION.down + CD4+ Treg, 0.26; INT.down + UNION.down,

0.26; INT.down, 0.26; UNION.down, 0.26.

Supplementary Files

Supplementary file 1: Fig. S1-S8. Supplementary figures and legends.

Supplementary file 2: Table S1. Patient information of lung cancer ICI cohorts.

**Supplementary file 3: Table S2. List of genes specific to the cell clusters in total cells and
each cell lineage.**

Supplementary file 4: Table S3. List of DEGs in comparison between ICI response

groups.

Supplementary file 5: Table S4. Details of GO terms significantly enriched in

comparisons for ICI response groups and PCs.

Supplementary file 6: Table S5. Gene list of tumor signatures.

Table 1. Clinical overview of NSCLC patients treated with ICI.

Immunotherapy	Targets	No. of samples	No. of patients
Pembrolizumab	PD-1	9	8
Atezolizumab	PD-L1	1	1
Nivolumab	PD-1	2	1
Vibostolimab+Pembrolizumab	TIGIT + PD-1	2	1

RECIST	Description	No. of samples	No. of patients
PR	Partial response	5	4
SD	Stable disease	3	2
PD	Progressive disease	6	5

839 **Table 2. Multivariate overall survival analysis of tumor signature genes.**

	TCGA LUAD				TCGA LUSC			
	HR	**Lower CI**	**Upper CI**	**p-value**	**HR**	**Lower CI**	**Upper CI**	**p-value**
NMF.p1	0.89	0.53	1.48	0.64	0.92	0.66	1.27	0.60
NMF.p2	0.87	0.53	1.43	0.59	0.93	0.67	1.28	0.66
NMF.p3	0.76	0.49	1.21	0.25	1.06	0.76	1.49	0.72
NMF.p4	1.00	0.64	1.55	0.98	1.00	0.70	1.44	0.98
scINSIGHT.PR.m	0.97	0.63	1.51	0.90	0.87	0.62	1.23	0.44
scINSIGHT.SD.m	0.81	0.53	1.23	0.32	0.94	0.67	1.33	0.74
scINSIGHT.PD.m	0.89	0.57	1.40	0.62	0.87	0.62	1.24	0.45
PC1.pos	0.48	0.27	0.84	0.01	0.78	0.55	1.12	0.17
PC1.neg	0.98	0.65	1.48	0.92	0.72	0.49	1.06	0.10
PC2.pos	0.87	0.55	1.36	0.53	0.98	0.69	1.40	0.91
PC2.neg	0.82	0.53	1.25	0.35	0.90	0.63	1.27	0.54
PC3.pos	1.14	0.68	1.90	0.63	0.87	0.63	1.19	0.38
PC3.neg	0.86	0.55	1.35	0.52	0.79	0.54	1.16	0.23
PC4.pos	0.88	0.54	1.42	0.59	0.66	0.46	0.95	0.02
PC4.neg	1.10	0.75	1.62	0.62	0.97	0.69	1.35	0.84
PC5.pos	1.09	0.71	1.68	0.68	0.76	0.53	1.10	0.15
PC5.neg	1.01	0.69	1.50	0.95	1.03	0.71	1.49	0.88
PC6.pos	0.77	0.45	1.29	0.32	1.09	0.79	1.51	0.60
PC6.neg	0.68	0.41	1.13	0.13	0.81	0.58	1.13	0.21
PC7.pos	1.10	0.71	1.72	0.67	0.75	0.51	1.10	0.14
PC7.neg	0.88	0.58	1.33	0.53	0.77	0.55	1.09	0.14
PC8.pos	0.77	0.50	1.18	0.23	1.01	0.72	1.43	0.94
PC8.neg	1.00	0.65	1.52	0.99	0.95	0.66	1.37	0.80
PC9.pos	0.91	0.60	1.38	0.66	1.02	0.73	1.43	0.91
PC9.neg	1.11	0.73	1.68	0.63	1.02	0.72	1.45	0.89
PC10.pos	1.13	0.76	1.68	0.55	1.28	0.88	1.86	0.20
PC10.neg	0.93	0.59	1.46	0.75	0.93	0.65	1.35	0.71
INT.up	0.82	0.53	1.26	0.37	1.00	0.71	1.42	0.99
UNION.up	0.80	0.52	1.21	0.28	0.77	0.53	1.13	0.18
INT.down	0.84	0.56	1.24	0.37	1.07	0.73	1.57	0.74
UNION.down	0.67	0.42	1.06	0.08	0.75	0.52	1.07	0.11

840
841

Figure 2

b

● p-value < 0.05

INT.down	0.009
UNION.down	0.009
CD4+ TH17	0.034
PC7.neg	0.047
CD4+ Treg	0.048

Smoking

Origins
Subtypes

Stages

Significance (p-value)

Immune and tumor signatures

c

UNION.down + B.Plasma
INT.down + B.Plasma
UNION.down + CD4.Treg
INT.down + UNION.down
INT.down
UNION.down

−log10 (p-value)

AUC

● Tumor−Immune ● Immune−Immune
● Tumor−Tumor ● Single