Lateral Entorhinal Cortex Dysfunction in Alzheimer's disease Mice

Radha Raghuraman[1,2], Andrew Aoun[1,2,†], Mathieu Herman[1,2,†], Oliver Shetler[1,2], Eden Nahmani[1,2] and S. Abid Hussaini[1,2,*]

1 *Taub Institute for Research on Alzheimer's disease and the Aging Brain, Columbia University Irving Medical Center, New York, NY, 10032, USA*

2 *Department of Pathology and Cell Biology, Columbia University Irving Medical Center, New York, NY, 10032, USA*

† Equal contribution

* Contact: S. Abid Hussaini (sah2149@columbia.edu)

Abstract

In Alzheimer's disease (AD), the formation of amyloid beta (Aβ) and neurofibrillary tangles (NFTs) leads to neuronal loss in entorhinal cortex (EC), a crucial brain region involved in memory and navigation. These pathological changes are concurrent with the onset of memory-related issues in AD patients with symptoms of forgetfulness such as misplacing items, disorientation in familiar environments etc. The lateral EC (LEC) is associated with non-spatial memory processing including object recognition. Since in LEC, neurons fire in response to objects (object cells) and at locations previously occupied by objects (trace cells), pathology in this region could lead to dysfunction in object location coding. In this paper we show that a transgenic mouse model, EC-App/Tau, which expresses both APP and tau primarily in the EC region, have deficits in LEC-specific memory tasks. Using in vivo single-unit electrophysiology recordings we show that the LEC neurons are hyperactive with low information content and high sparsity compared to the controls indicating poor firing fidelity. We finally show that object cells and trace cells fire less precisely in the EC-App/Tau mice compared to controls indicating poor encoding of objects. Overall, we show that AD pathology causes erratic firing of LEC neurons and object coding defects leading to LEC-specific memory impairment.

Introduction   1

Detecting early signs of neuronal vulnerability in Alzheimer's disease (AD) prior to the onset of   2
cognitive symptoms is crucial to intervene and prevent further progression. In AD, the   3
entorhinal cortex (EC) is shown to have selective vulnerability due to the accumulation of   4
amyloid beta (Aβ) and tau, especially in the lateral EC (LEC) [1–3]. Therefore, early AD   5
symptoms such as misplacing objects, anosmia and forgetting events could be due to the   6
distinct role of LEC in object-, odor- and episodic memories [4–11]. Anatomical studies have   7
shown that while hippocampal place cells receive spatial information from the MEC, the   8
non-spatial information is mainly provided by the LEC [12–15]. The LEC neurons show very   9
little spatial selectivity in an open field even with several cues and landmarks but fire in   10
response to discrete objects [13,16–18]. Two distinct types of specialized cells viz., object cells 11

and object-trace cells have been identified in the LEC of rats thus far. While an object cell fires in response to an object, an object-trace cell fires at a location previously occupied by an object, indicating that there are two independent neural correlates for object-place memory in the LEC [9]. These cells collect, remember and process object-place information as part of non-spatial memory to be integrated with spatial memory in the hippocampus.

The LEC also processes olfactory information and receives input from both olfactory bulb (OB) and piriform cortex (PCX). The LEC neurons, unlike the PCX neurons, are more narrowly tuned and process complex odors [19]. The dominant input to LEC is from the afferent fibers from olfactory areas [20] which then projects back to both the aforementioned areas [21,22]. Odor-discrimination-trained animals display EC signaling to OB prior to the odor onset which prepares the system for odor sampling [23]. This indicates LEC to be instrumental in providing highly odor-specific, expectation-dependent feedback to the olfactory bulb [24,25]. Any loss or deterrence of this feedback results in the interference with odor memory, thereby contributing to early impairment in olfactory perception that is commonly observed in AD [26–29]. In addition, the LEC is known to coalesce multisensory information. Several studies, including LEC lesions, have shown that object-context or odor-context memory is impaired while the animal's ability to discern new objects, odors and contexts individually is intact, indicating that LEC is critical in complex, association memory and plays an important role in the integration of episodic information [5,6,30–32].

Several studies including ours have shown that both amyloid beta and tau pathologies affect MEC neurons and impair the grid cell and speed cell function, yet not much is known about the impact on LEC neurons [33–36]. In vitro studies have implicated the LEC neurons in App knockin mice to be specifically hyperexcitable [8]. Studies using functional imaging have shown that expressing App and tau in the EC of mice affects the LEC function [3]. To our knowledge, this is the first in vivo electrophysiology study that investigates the impact of AD pathology on LEC neurons. Further, we are the first to report the object and trace cells in mice and how they are affected by AD pathologies. In this paper we show that AD pathology in the LEC of mice causes them to become hyperactive with loss of firing fidelity and poor encoding of objects possibly leading to object-context memory deficits.

Results

EC-App/Tau mice are impaired in LEC-specific memory tasks

EC-App/Tau mice accumulate Aβ and tau in the entire hippocampal formation but only the LEC region appears to be vulnerable in functional imaging studies ([3]). Using the same mouse model, we confirmed that Aβ and tau accumulates in the hippocampal formation of EC-App/Tau mice (Figure S1) and wanted to test if they had LEC-specific memory deficits. We subjected 20-month-old EC-App/Tau mice and age-matched Neuropsin controls (NON) to three behavioral tasks; Object context recognition (OCR), Odor discrimination (OD) and T-maze alternation tasks. Among these, the OCR and OR are LEC-specific memory tasks while T-maze is a hippocampal-dependent memory task ([4,5,32,37,38]). In the OCR task, the mice were allowed to explore two identical sets of objects (two cubes or two triangular pyramids) in two different contexts (black cylinder and gray striped cylinder) during training for 2 days. During test on day 3, one of the objects in one context was replaced with an object belonging to the other context (e.g. a triangular pyramid replaced with a cube in the gray striped

context). The expectation was that the mice with intact memory would spend more time exploring the out-of-context object (blue cube) (Figure 1, Left). EC-APP/tau mice did not explore the out-of-context objects as much as the controls did evident from the Fisher's exact test given by $p < 0.001$ wherein the controls show a significant difference with the EC-App/Tau in being able to recognize the objects in the context. In the OD task, mice were trained to turn left or right to receive food pellets based on the odor they were presented with in the stem of the T-maze (Figure 1, Right). EC-APP/tau mice made many errors compared to the controls, indicating poor odor discrimination (evident from the contingency table representing the frequencies of corresponding arm choices made by the mice belonging to respective cohorts of EC-App/Tau: incorrect arm choice = 68.89% and correct arm choice = 31.11%) while the NON group shows 40% incorrect arm choice and 60% correct arm choice from Fig1 right panel representing odor discrimination task). Finally, we tested mice in a T-maze spontaneous alternation task where mice were trained to turn left and right alternatively by blocking one arm in the sample trial and allowing them to choose between two arms in the choice trial. While EC-App/Tau mice made more errors in this task with incorrect alternations compared to the control mice, the effect was not significant (p = 0.1178) (Figure S2). This is likely due to the fact that T-maze spontaneous alternation is a hippocampal memory task while OCR and OD are LEC-specific tasks.

Figure 1. Object context Recognition and Odor Discrimination The left panel shows the set up of Object Context Recognition (OCR) with context 1 showing a certain arena with two objects of cube shape and context 2, another arena showing two other objects of the triangular shape with the test phase showing two objects one each from both the contexts in the training phase; the adjacent histogram represents the frequencies (in percentage) indicating the time spent by the mouse with each of the objects. The right panel shows the set up of Odor Disrimination (OD) task where the mouse is introduced to a puff of apple odor upon taking left and a puff of banana odor upon taking right; the histogram on the right shows the contingency table representing the frequencies corresponding to the correct (in plain) and incorrect arm chocies (represented in hatched lines) made by the mouse from the respective cohort

LEC neurons of EC-App/Tau mice show increased firing and reduced information

Since LEC-specific memory is affected, we focused on understanding how Aβ and tau impacts neurons in the LEC. We collected single-unit activity from neurons in the LEC of 22-month-old EC-App/Tau mice, age-matched NON mice, and 8-month-old C57BL/6J (B6) mice while they explored an open field cylindrical arena with or without an object for 10 minutes (see next section and methods for details). We found that the LEC neurons of EC-App/Tau mice had a higher firing rate compared to the NON and B6 mice (Figure 2, left), and NON mice had higher firing rate compared to the B6 mice evident from the $p < 0.0001$ between the EC-App/Tau and the respective control groups. The Skaggs information content in EC-App/Tau mice was much lower compared to the control mice indicating poor encoding of spatial information (Figure 2, right) EC-App/Tau has a significantly lower information content compared to NON evident from $p < 0.0001$; as is the case with EC-App/Tau showing a significantly lower information content with B6 given by $p < 0.0001$ while it is noteworthy that there is no significant difference between the cohorts NON and B6. Consistent with this, the sparsity of LEC neurons in EC-App/Tau mice was higher compared to the controls (Figure S3). However, both selectivity and coherence of EC-App/Tau mice were similar to B6 mice (Figure S3).

Figure 2. Firing Rate and Skaggs Information Content The left panel represents the histogram showing comparisons of firing rate amongst the three cohorts - B6 (blue), NON (green) and Ec-App/Tau (red). The right panel represents the histogram showing comparisons of Skagg's information content amongst the three cohorts - B6 (blue), NON (green), and EC-App/Tau (red). P values are represnted by asterisks and error bars are SEM

Object and Trace cells in EC-App/Tau mice are poorly tuned

Studies in rats have shown that LEC neurons fire in response to objects (object cells) and also fire at locations occupied by previous objects (object-trace or trace cells) ([9, 17]). Since LEC neurons of EC-App/Tau mice are impaired (figure 2 and figure S2), their responses towards object locations could also be affected, explaining the object memory impairment in these mice (1). We collected single-unit responses from LEC neurons in the three groups of mice; B6, NON

and EC-App/Tau, as they explored a cylindrical arena for 3 or more sessions for 10 minutes each. In the first session, the mice explored the arena for 10 minutes without an object and in subsequent sessions, an object was positioned at the edge of the cylinder at one of the following locations: 0, 90, 180 or 270 degrees (Figure 3, bottom).

We quantified LEC neuronal responses towards objects in each session using the EMD (Earth Mover's Distance) metric ([39]). The distances between firing fields and the object locations were computed using a single point Wasserstein distance (see methods). For no object sessions, the center of the arena was used as a reference point. A cell whose firing field was closest to an object location above other nearby random locations for 2 consecutive sessions was defined as an object cell. Representative firing fields for each group; B6, NON and EC-App/Tau, for No object and objects at 4 possible locations (black circles in columns 2-5) are shown in figure 3, middle. A cell whose firing field was closest to a previous object location above other nearby random locations was defined as a trace cell. Representative firing fields for each group for objects previously placed at 4 possible locations (white circles in columns 2-5) are shown in figure 3. We plotted the average population firing response of all neurons towards all 4 object locations combined (figure 3) and individually (Figure S4) for each groups. The population response of LEC neurons revealed that the firing fields in B6 mice were more uniformly spread across 4 object locations, with high firing (warm colors) at the edge of the circle and low firing (cold colors) in the center, compared to NON and EC-Tau/App mice indicating better tuning towards objects.

Object and Trace cells in EC-App/Tau mice are less stable

We found about 5%-10% object cells (figure 4, top left) with significantly more cells in NON mice compared to the other two groups (B6: 5%, NON: 10% and EC-App/Tau:5%). We found 20%-30% trace cells (figure 4, top right) with significantly more cells in NON compared to B6 (B6: 24%, NON: 30% and EC-App/Tau: 26%).

To interpret group stability however, it was important to choose one reference point to remain consistent across groups. Given that we expected object and trace behavior, and that our tested locations were along the edges, we chose the center of the arena as our reference and hypothesized that greater object/trace content would shift stability away from the center. To compare stability, we fit regression models to the distribution of EMD quantiles associated with each group pair and for a given remapping metric. This model incorporated both random animal effects as well as fixed effects for group and session number. For both metrics, we observed the expected trend in stability with B6 showing the largest averaged quantile followed by NON and subsequently EC-App/Tau (Figure S5, top panel). We found that the fitted coefficient for the quantiles computed relative to the center of the arena indicated that the EC-App/Tau group was significantly less stable than B6 for the whole map EMD metric while both EC-App/Tau and NON were significantly less stable than B6 for the field-restricted EMD metric.

In effect, many object and trace cell sessions in the EC-App/Tau mice were of low quality (figure 4, bottom panel), i.e. fields were farther away from current or previous object locations compared to NON and B6 mice. In about 23% of sessions, the fields of EC-App/Tau mice were closer to the center of the arena rather than any of the 4 corners compared to the fields of B6 (10%) and NON mice (20%). In about 40% of the sessions, the fields of EC-App/Tau mice were in between two object positions (for example at 45 degree position instead of 0 or 90 degrees) compared to the fields of B6 (25%) and NON mice (30%). Similarly, cell sessions in NON mice

Figure 3. Object Cells: Representative object cell responses for B6, NON and EC-App/Tau mice for no object (left column) and 4 possible current object locations; 0, 90, 180 and 270 (black circles in columns 2-5). Columns 1-3 are part of the same session and rest from another session. **Trace Cells:** Representative trace cell responses for 4 possible previous object locations; 0, 90, 180 and 270 (while circles in columns 2-5). Columns 1 and 2 are part of the same session. Black circles indicate current object location. Average population response of neurons towards objects for the 3 groups.

Figure 4. Cell Counts. Cell counts are compared using Fisher's exact test. The percentage of cells which show object and trace behavior across their sessions compared for the 3 groups (top panel). The first quality metric shows the proportion of cases where the EMD quantile is smaller relative to the center point than to any of the 4 object locations (left). The second shows the proportion of cases between two object locations such that the reference point of the minimum quantile is different for the field-restricted EMD and the field centroid distance (right). Raw cell and cell-session counts are provided above each proportion.

were of lower quality compared to B6 mice. The remaining 60%-75% of cells were not tuned to an object (unassigned cells).

Importantly, unlike NON and EC-APP/Tau mice, we had few B6 recordings with more than 3 sessions, and therefore limited the net cell type counts across groups to 3 sessions (figure 4). However, given that LEC responses are shaped by learning and experience, we repeated the comparison between NON and the EC-APP/Tau mice for up to 6 consecutive sessions (Figure S6, bottom panel). This validates the use of our quality metrics as we found that the proportion of fields between two angles remained stable across sessions for both groups. Additionally, we show that with more experience, NON mice increase their tuning by reducing the proportion of cells closer to the center while ANT mice increase it. To confirm this result, we show that the trend in EMD quantiles is significantly positive for the NON mice while that of the EC-APP/Tau is significantly negative or close to zero (Figure S5). We therefore repeated

the object and trace counts after filtering out ambiguous cases based on our quality metrics. While this reduced the raw count of object and trace cells, EC-APP/Tau mice demonstrated a significantly reduced proportion of both cell types compared to the NON mice (Figure S6, top panel). This shows how learning and memory deficits in EC/App-Tau mice are specifically tied to the reduced quality of object and trace cell content.

Discussion

This manuscript studies the impact of Alzheimer's disease (AD) pathology on the lateral entorhinal cortex (LEC), a region of the brain associated with non-spatial memory processing. The study uses a transgenic mouse model, EC-App/Tau, which expresses both amyloid precursor protein (APP) and tau primarily in the entorhinal cortex (EC) region. It demonstrates that EC-App/Tau mice exhibit deficits in LEC-specific memory tasks. Among the three behavioral tasks, viz. Object context recognition (OCR), Odor discrimination (OD), and T-maze alternation tasks, the mice were affected only in the LEC-specific, OCR and OD tasks and not in the hippocampal-dependent task. This is consistent with previous studies that have shown OCR deficits due to lesions but NOR memory was spared [5].

Previous studies have shown that Aβ and tau can cause neuronal hyperexcitability leading to network dysfunction ([40–43]) and memory deficits. We confirm that this is also the case in the LEC neurons of EC-App/Tau mice which not only show hyperexcitability, but also have low information content and high sparsity compared to the controls. This indicates poor firing fidelity, which could contribute to the observed memory deficits. The fMRI study done on EC-App/Tau mice which measured cerebral blood volume (CBV) showed hypofunction in the LEC region but we observe hyperexcitability in our data. This could be due to the differences in recording method, age of mice and brain state. While CBV measures relative blood volume in the LEC our technique directly measures neuronal firing which might be inherently different. The age used in fmRI was 12 months while we used older, 24+ old mice. Finally, fMRI was done on anesthetized mice while our mice were freely moving.

The study found that while the AD mice do have an equal or greater number of object and trace cells, their coding for object or object location was sub par. Using our sensitive EMD metric, we show that tuning and stability of object and trace cells in the EC-App/Tau mice is significantly poorer compared to the controls. Many object and trace cell sessions in the EC-App/Tau mice were of low quality, with fields farther away from current or previous object locations. This poor tuning and stability could contribute to the observed object-context memory deficits. Additional evidence for this is provided in the findings that, with experience, EC-App/Tau mice recruit fewer object and trace cells compared to healthy controls.

One of the caveats of the EC-App/Tau model used here is the known leakiness in the expression of neuropsin which could express amyloid beta and tau in other regions too [44]. While we specifically did not investigate this, our IHC data shows high amyloid and tau accumulation in the EC and other regions of hippocampal formation. However, this doesn't preclude our electrophysiology results since data was collected spefically in the LEC region and it is unlikely that the neuronal dysfunction we see in LEC is due to pathology in other brain regions.

Future research could further explore the mechanisms underlying the observed neuronal dysfunction in the LEC region. Previous studies have shown that fan cells are partially involved in object cell dysfunction and these specific cell-type vulnerability in AD could be studied. [45].

While this paper did not explore how LEC neurons coding for odors are affected, other studies have indicated that fan cells are involved in associative memory and future studies could study if in AD these cells are affected [7]. It would also be interesting to investigate whether similar patterns of neuronal dysfunction and memory deficits are observed in other regions of the brain affected by AD pathology. Additionally, the development and validation of LEC-specific memory tasks in human subjects could be a valuable avenue for early detection and intervention in AD.

Materials and Methods

Animals

We used a previously described transgenic AD mouse model, EC-APP/Tau ([3,46]), and its littermates containing neuropsin promoter without amyloid beta and tau (NON) as age-matched control. These animals were used at 20+ months of age for behavior, 22+ months for electrophysiology experiments and euthanized before 25 months of age for immunohistochemistry. We used C57BL/6J mice (B6) as a young control for electrophysiology experiments. All animals were maintained on a 12-hour light/dark cycle with food and water provided ad libitum. All animal experiments were performed in accordance with national guidelines (NIH; National Institutes of Health) and approved by the Institutional Animal Care and Use Committee at Columbia University.

Behavior

Object Context Recognition

The experiment was performed during the light phase of the circadian cycle at a 30-lux light level in a 40-centimeter diameter cyilnder with interchangeable walls. Throughout testing, the experimenter remained unaware of the animal genotypes. Mice were tested in 2 different contexts; each displayed a pair of objects in a similar setup previously used for rats ([47]). During training (sample phase), context 1 displayed white walls and presented a pair of small identical objects (objects 1A and 1B) whereas context 2 displayed gray walls and presented another pair of small identical objects (objects 2A and 2B). On habituation day (Day 1), all mice were allowed to explore both contexts without objects for 20 min in the morning and in the afternoon. During the first two days of training (Days 2 and 3), each animal was familiarized with both contexts and their pairs of identical objects. Mice were run 4 consecutive times for 3 min each time with rest intervals of 2 min in a holding cage. As a result, animals randomly experienced each context and their respective, identical objects twice per day. On testing day (Day 4), mice were randomly run in each context once before being tested in the last context to which they were exposed, only this time object 1A/B was substituted with object 2A/B or vice versa. The object that did not belong to the context was defined as the "novel object", while the other object was referred to as the "familiar object". Testing lasted 10 min and aimed to determine whether the animals spent more time exploring the novel object. A contingency table for entries into familiar object and novel object zone was tabulated and Fisher's exact test was used to calculate the statistical significance. Odds ratio was calculated using Baptista-Pike method. Number of entries were converted to % entries and plotted as stacked bar graphs.

Odor Discrimination Task

The task was carried out with the same mice habituated in the same environment described in the T-maze spontaneous alternation protocol. Prior to testing, mice were single housed and food restricted. Weights were monitored and recorded in a logbook twice per day, before and after animal performance, to make sure mice remained above 85% of their body weight. Food restriction was adjusted accordingly. Animals were trained to discriminate between 2 distinct odors; odor A (Isoamyl acetate) and odor B (Ethyl valerate). Mice had to learn the association of a specific odor to a goal arm. To accomplish the task, animals performed a repetition of 40 trials per day for 4 consecutive days as follows: Each mouse was first confined in the stem where one of the 2 odors was presented to them using a Q-tip. Mice remained in the stem for 30 seconds, after which, the door was opened and the mice were allowed to choose an arm. On correct choices (for example, left arm for Isoamyl acetate), a reward (sugar pellet) was placed at the far end of the goal arm. After the animal consumed the reward, they were placed back in the stem of the T-maze. The maze was wiped with 70% ethanol in between each trial. The number of correct and incorrect entries into the arms for the final day were recorded in a contingency table. Data was analyzed using Fisher's exact test to assess statistical significance.

T-maze Spontaneous Alternation

The experiment was conducted during the light phase of the circadian cycle at a 30-lux light level. Mice were tested to perform spontaneous alternation in a T-shaped maze as described previously ([48]). The maze measured 40 (stem) x 46 (arm) x 10 (width) cm and displayed a black cue card on the left side of the stem. During testing, experimenter was blind to the animal genotypes. On habituation day, all mice were allowed to explore both arms freely. Mice were then tested 6 times per day for 3-4 consecutive days as follows: Each mouse was first placed at the stem and offered to explore one randomly-picked open goal arm (with other arm closed) for 1 min. If the goal arm was reached, mice were placed at the stem again and the trial was repeated with both arms open. Testing was ended after the animal chose one arm or stayed in the stem for 1 min. Between trials, the maze was wiped with 70% ethanol. The number of correct and incorrect choices for all days (alternative goal-arm and same-arm/no-arm choices) were recorded in a contingency table. Data was analyzed using Fisher's exact test to assess statistical significance.

Surgery and electrode implantation

Custom-made, reusable 16-channel microdrives (Axona, UK) were constructed by attaching an inner (23 ga) and an outer (19 ga) stainless steel cannula to the microdrives ([49]). Tetrodes were built by twisting four 25 mm thick platinum-iridium wires (California wires) and heat bonding them. Four such tetrodes were inserted into the inner cannula of the microdrive and connected to the wires of the microdrive. One day prior to surgery, the tetrodes were cut to an appropriate length and plated with a platinum/gold solution until the impedance dropped to about 150 ohms. On the day of surgery, mice were anesthetized with a mixture of ketamine and xylazine (100 mg/ml and 15 mg/ml, respectively, per 10 g body weight) and monitored for depth of anesthesia before proceeding. Mice were then fixed within the stereotaxic frame with the use of zygomatic process cuff holders and an incision was made to expose the skull. About 3–4 jeweler's screws were inserted into the skull to support the microdrive implant. An

additional screw connected with wire was also inserted into the skull which served as a
ground/reference for local field potential (LFP) recordings. A 2 mm hole was made on the left
hemisphere of the skull at ML 4.3 mm and AP -3.4 mm from bregma and tetrodes were lowered
to -3.0 mm to target the LEC. Dental cement was spread onto the skull and secured with the
microdrive. Mice were allowed to recover from anesthesia in a cleaned cage placed on a warm
heating pad until awake (45 min) before finally being transported to housing. Carprofen (5
mg/kg) was administered to mice prior to surgery and post-operatively to reduce pain. Mice
usually recovered within 24 h, after which the tetrodes were lowered and recording began. All
recording depths reported are from the surface of the brain.

In vivo electrophysiology recording and cell classification

The mice explored a 40 cm diameter white cylindrical arena with a vertical black cue card (8.5
x 11 inch) attached to north-side of the cylinder. Mice underwent 3-5 recording sessions per day,
with 10 minutes to 2 hours between sessions. A session always started without an object in the
cylinder (No object) followed by an object positioned at the edge of the cylinder at either North
(0 degrees), East (90 degrees), South (180 degrees) or West (270 degrees) direction. Tetrode
positions were moved no more than 50μm at a time, and only after the last recording session of
the day, allowing 12 hr of stable electrode positioning prior to the next recording session.
Neuronal signals from experimental mice were recorded using the Axona DacqUSB system. The
signals were amplified 15,000 to 30,000 times and band pass filtered between 0.8 and 6.7 kHz.
LFP was recorded from 4 channels of the electrodes. The LFP was amplified 15,000 times,
lowpass filtered at 125 Hz and sampled at 250 Hz. Notch filter was used to eliminate 60 Hz
noise. The recording system tracked the position of the infrared LED on the head stage
(sampling rate 50 Hz) by means of an overhead video camera. Tracking artifacts were removed
by deleting samples greater than 100 cm/s and missing positions were interpolated with total
durations less than 1 s, and then smoothing the path with a 21- sample boxcar window filter
(400 ms; 10 samples on each side). Spike sorting was performed offline using MountainSort
software ([50]) followed by manually checking and validating the clusters. Quantitative
measurements of cluster quality were subsequently performed, yielding isolation Distances in
Mahalanobis space ([51]). There were no significant differences between clusters in our
experimental groups (Median Isolation distances: 8+ mo B6- 9.44, 24+ mo NON- 10.13 and
24-mo EC-App/Tau- 9.83. Kruskal-Wallis test- P = 0.334). A total of 704 neurons from 15
animals were recorded with 199 neurons in B6 mice (N=5), 169 in NON mice (N=5) and 336 in
EC-App/Tau mice (N=5). To match cells across sessions, a custom-script in Python called Unit
matcher was used which compared the waveform properties of all cells and re-assigned the cell
numbers in the clusters.

Neural coding properties

All the data for neural coding properties (e.g. information content) are expressed as the mean $\pm$
the standard error of the mean (SEM). We performed the D'Agostino & Pearson omnibus
normality test to determine if the data was normally distributed. We then chose the following
statistical tests. Kruskal-Wallis test was used to compare medians in all groups. Fisher's exact
test was used to compare the binary data from the Object context recognition, Odor
discrimination and T-maze tests. A value of $p < 0.05$ was considered statistically significant for
all measures. The N represents the number of animals in each group, unless specified otherwise.

The exact values of N are indicated in figure legends, and in the Methods section.   322

Object and trace cell analysis   323

We used our previously published EMD method [39] to calculate object and trace cell stability   324
across sessions. For a given cell's ratemap on a given session, we quantified remapping relative   325
to chosen reference points. Specifically, we computed a remapping distance between an activity   326
map and the object location. We did this for the full ratemap, to capture whole map stability,   327
and for a masked version, to isolate and track individual fields. We used the single point   328
Wasserstein distance, a strong approximation of the true distance between the firing rate map   329
and a full pseudo-rate map with all the density at one point. Alongside these EMD metrics, we   330
also compute a euclidean distance between the weighted average firing peak of a given field and   331
each of the reference points. We refer to this as the centroid distance.   332

Reference points and distribution   333

The reference points were selected as the 4 tested object locations (0, 90, 180, 270). We also   334
computed a 5th reference point set at the center of the arena (NO = no object). While an   335
object may or may not have been present on a given session, we computed remapping relative   336
to all 5 reference points for every cell ratemap on every session. This allows us to track   337
object/trace behavior and movement towards the objects throughout tested sessions.   338
  For each reference point we compute a remapping distance and convert this into a quantile   339
based on a reference distribution. For a given cell on a given session, we computed the   340
minimum and maximum 'x' and 'y' positions covered by the animal on that session. This gave   341
us an egocentric map of possible (x,y) locations covered. Given the EMD property where   342
gradients of remapping emerge across the map when localizing a field, we decided to sample in   343
a way that would give us coverage across these gradients. We therefore sampled from the   344
possible (x,y) positions using a hexagonal lattice with a spacing parameter of 2 resulting in a   345
hexagon side of $2/\sqrt{3}$ cm. Points were sampled from all 6 vertices of each hexagon as well as   346
the center. We also masked the collected points to ensure only those that would fall inside the   347
circular arena were selected. This gave us a uniform sampling across the gradients and on   348
average consisted of 776 ± 148 sample points for a given map.   349
  Using these sample points, we computed a remapping distance from the given map score to   350
each of these randomly sampled reference points. With a distribution of remapping distances to   351
random reference points and our empirically observed distances (for each of the 5 reference   352
points), we compute a quantile describing the spatial location/dispersion of our map/field   353
relative to other possible random locations.   354

Cell assignments   355

We define cells in three categories: object, trace and unassigned. Each cell appearance in a   356
session is labeled with isTrace and isObject identifiers, which are set to 1 if conditions are met,   357
otherwise 0. For object cells, if the minimum quantile during a session matches the current   358
location, isObject is set to 1. For trace cells, isTrace is 1 if the quantile matches any prior   359
tested location. After processing all cell appearances, we apply filters to ensure data integrity   360
and exclude ambiguous cells. We use three filters: 361

- Exclude cells with an isolation distance under 5.   362

- The quantile using the arena center must exceed the quantile of at least 1 object angle. In other words, the arena center quantile cannot be the minimum

- The reference point of the minimum quantile using the field-restricted EMD should align with the minimum quantile using the centroid distance. This accounts for cases where two object angles have very small and similar quantiles.

The first filter is applied across all cell appearances. The next two filters are used to generate filtered cell assignments based on the ambiguity of a field's position. These two filters are relative in that they require no threshold. We preserve cells appearing in at least two sessions post-filtering. Definitions applied for cell assignment are:

- An object cell appears in at least two sessions with the minimum quantile at the current location.

- A trace cell appears in at least one session with the quantile at a prior object location.

Object cells are prioritized over trace cells, minimizing their overlap (1-2 cells typically). This methodological rigor helps maintain the session structure and prevents loss of data from low-quality recordings.

We performed Fisher's exact test to determine if there were differences between the total population cell assignments (object, trace, unassigned). We then performed pairwise 2x2 Fisher's exact test to compare specific group pairs within a given assignment of object or trace. The 2x2 Fisher's exact tests were also applied separately in each group to group comparison for each of the quality metrics.

Spatial stability

To explore spatial stability over time (relative to the arena center), we grouped our data by session number from 1 to 6 and computed a mean quantile for each group. There were 6 means for ANT and NON and 3 means for B6. The Mann-Kendall test was applied to obtain an empirical test statistic for each. The slope was computed as the Theil-Sen estimator. To assess significance given the low number of sessions and high fluctuations within a given session we applied a bootstrap Mann Kendall test where, for a given session number, we shuffled data points across the 3 groups and recomputed a new set of mean values from which a Mann Kendall test statistic was computed. This was done 1000 times to create a distribution to which the empirical test statistic was compared and a p-value obtained. Benjamini-Hochberg corrections were applied to collected p-values.

To look at overall group stability, we used the standardized EMD quantiles as part of a beta regression model to compare quantiles across two groups. For each possible group to group comparison, we fit a model with the group id and session id as fixed effects, the animal id as a random effect and a session id to group id interaction term.

This gives the formulas:

$$\text{Obj\_q\_NO} \sim (1 \mid \text{name}) + \text{group} + \text{session\_id} + \text{session\_id} : \text{group}$$

DAB Immunohistochemistry and Imaging

After final recording, all mice were anesthetized with a mix of ketamine/xylazine and were transcardially perfused with ice-cold 100 mM phosphate-buffered saline (PBS, pH 7.4), followed by 10% formalin (Fisher Scientific, USA). For mice implanted with microdrive, the electrodes were not moved until after perfusion. Brains were then harvested and left in 10% formalin overnight, then incubated in 30% sucrose until the brains sank to the bottom of the conical tube (all at 4°C). Horizontal brain sections were sliced (30 μm) using a Leica CM3050 S cryostat and stored in cryoprotectant at -20°C until immunostaining procedures. EC-APP/Tau mice and NON mice that appear in Fig 1 were euthanized immediately for immunostaining. For both immunoperoxidase staining and immunofluorescence imaging, EC-APP/Tau mice brain sections were processed in parallel with sections from NON mice where appropriate.

Immunoperoxidase staining was performed using a Mouse-on-Mouse kit (Vector Laboratories, USA) and modified from previous methodology [11, 39, 55]. Briefly, cryoprotectant was washed from the free-floating tissue sections with PBS before quenching endogenous peroxidases with 3% H_2O_2. Sections were then blocked with mouse IgG–blocking reagent for 1.5 hours at room temperature, followed by overnight incubation at 4°C with either anti-beta amyloid 6E10 (mouse, 1:1,000 dilution of 1 mg/mL stock) (Biolegend, USA), anti-tau MC1 (mouse, 1:500; courtesy of Peter Davies, Feinstein Institute for Medical Research, North Shore LIJ), biotinylated anti-phospho tauSer202/Thr205 AT8 (mouse, 1:500) (Thermo Scientific, USA) or anti-human tau CP27 (mouse, 1:500; courtesy of Peter Davies). Tissue sections were then rinsed in PBS and incubated for approximately 20 minutes at room temperature in a working solution of biotinylated anti-mouse IgG reagent (excluding the biotinylated AT8-labeled sections). After several PBS rinses, sections were then incubated in an avidin-biotin conjugate for 10 minutes before being developed in H_2O containing 3,3'-diaminobenzidine (DAB) hydrochloride and urea hydrogen peroxide (Sigma Aldrich, USA). After staining was completed, tissue sections were mounted onto glass Superfrost Plus slides (Fisher Scientific, USA), allowed to air dry completely, and then dehydrated in ethanol and cleared with xylenes before being coverslipped.

Code

Python code from our custom toolkit NeuroScikit was used to match units across sessions, quantify neural coding, assess spatial stability and identify cell types. Post-hoc analyses and visualizations were done on Jupyter notebooks and using Matplotlib or Prism 9 software (GraphPad, San Diego, CA, USA).

Author contributions

S.A.H. performed the surgeries and implanted the electrodes. M.H., E.N. and S.A.H. collected the behavioral and electrophysiology data. M.H. and E.N. performed immunohistochemistry. R.R. and S.A.H. sorted the spike data. R.R. and S.A.H. analysed the behavioral and immunohistochemistry data. A.A. and O.S. developed the analytical code in Python. A.A. and R.R. processed the electrophysiology data. A.A. conducted post-hoc analyses, with help from O.S., and data visualizations. S.A.H. conceived the idea, discussed the methods and results with R.R, A.A. and M.H.. R.R., A.A., M.H. and S.A.H. wrote the manuscript. S.A.H. supervised the project and obtained the funding.

Supplementary Figure 1. Tau and Amyloid beta accumulation in EC-App/Tau mice.
Left: EC-App/Tau mice at 24 months with more pronounced tau accumulation (AT8 staining) in the EC region and some in the neighboring hippocampal regions. **RIght**: Aβ accumulation in the hippocampal formation of same mouse as left with more deposition in the EC region

Acknowledgements

We would like to thank Dr. Karen Duff for providing us with the EC-App/Tau line along with neuropsin controls. We thank Aisha Kazeem, Lisa Shah, Adithi Jayaraman and Sandhya Senthilkumar for help with spike sorting earlier dataset that first identified object and trace cells. We thank Dr. Gustavo Rodriguez for help with immunohistochemistry techniques provided to M.H. and for the valuable feedback on the manuscript. These tools were supported by NIH/NIA grants R01AG050425 and R01AG064066.

Supplementary Figure 2. EC-app/Tau mice are not impaired in T-maze alternation task. **Left**: Schematic of two trial blocks with sample (left) and choice (right) runs. **Right**: Stacked bars showing relative % entries for NON and EC-App/Tau mice into left or right arm of the T-maze. EC-App/Tau mice made fewer correct entries and more errors compared to the NON mice but this was not significant.

Supplementary Figure 3. Selectivity, Sparsity and Coherence The histogram on the left represents comparisons amongst the B6, NON and EC-App/Tau cohorts for selectivity; the panel in the middle shows sparsity and that on the right shows coherence. statistical significance is represented by asterisks wherein ns indicates $P>0.05$, ** indicates $P<0.01$, *** indicates $P<0.001$ and **** indicates $P<0.0001$

Supplementary Figure 4. Average population responses. All valid cells are considered across 3 groups (B6, NON, EC-App/Tau) and 5 possible locations (0, 90, 180, 270, NO = no object). Ratemaps averaged by mouse group ('B6', 'NON', 'EC-App/Tau') (A). Ratemaps grouped by current object location on a given session and averaged (B). The top row displays a representative sequence with current (black) and previous (white) object locations displayed at each step.

Supplementary Figure 5. Group stability. Stability metrics are displayed as quantiles for the whole map EMD (left column) and the field-restricted EMD (right column). The 3 groups (B6, NON, ANT) are compared using boxplots with significance values from the regression group coefficient (top row). EMD quantiles are separated by session number and trends are compared using a bootstrapped Mann-Kendall slope test (bottom row).

Supplementary Figure 6. Cell Counts Over Time. Cell counts are compared using Fisher's exact test. The percentage of filtered cells which show object and trace behavior with an increasing session limit compared for the 3 groups (A). Later sessions are shaded in progressively lighter colors. The percentage of cell-session appearances (individual ratemaps) failing a quality metric are displayed (B). The first quality metric shows the proportion of cases where the EMD quantile is smaller relative to the center point than to any of the 4 object locations (left). The second shows the proportion of cases between two object locations such that the reference point of the minimum quantile is different for the field-restricted EMD and the field centroid distance (right). Raw cell and cell-session counts are provided above each proportion.

References

1. H. Braak and E. Braak. Neuropathological stageing of Alzheimer-related changes. *Acta Neuropathologica*, 82(4):239–259, 1991.

2. Teresa Gómez-Isla, Joseph L. Price, Daniel W. McKeel Jr., John C. Morris, John H. Growdon, and Bradley T. Hyman. Profound Loss of Layer II Entorhinal Cortex Neurons Occurs in Very Mild Alzheimer's Disease. *The Journal of Neuroscience*, 16(14):4491–4500, July 1996.

3. Usman A. Khan, Li Liu, Frank A. Provenzano, Diego E. Berman, Caterina P. Profaci, Richard Sloan, Richard Mayeux, Karen E. Duff, and Scott A. Small. Molecular drivers and cortical spread of lateral entorhinal cortex dysfunction in preclinical Alzheimer's disease. *Nature Neuroscience*, 17(2):304–311, February 2014. Number: 2 Publisher: Nature Publishing Group.

4. Julie Chapuis, Yaniv Cohen, Xiaobin He, Zhijan Zhang, Sen Jin, Fuqiang Xu, and Donald A. Wilson. Lateral entorhinal modulation of piriform cortical activity and fine odor discrimination. *The Journal of Neuroscience: The Official Journal of the Society for Neuroscience*, 33(33):13449–13459, August 2013.

5. David I.G. Wilson, Sakurako Watanabe, Helen Milner, and James A. Ainge. Lateral entorhinal cortex is necessary for associative but not nonassociative recognition memory. *Hippocampus*, 23(12):1280–1290, 2013. _eprint: https://onlinelibrary.wiley.com/doi/pdf/10.1002/hipo.22165.

6. David I.G. Wilson, Rosamund F. Langston, Magdalene I. Schlesiger, Monica Wagner, Sakurako Watanabe, and James A. Ainge. Lateral entorhinal cortex is critical for novel object-context recognition. *Hippocampus*, 23(5):352–366, 2013. _eprint: https://onlinelibrary.wiley.com/doi/pdf/10.1002/hipo.22095.

7. Jason Y. Lee, Heechul Jun, Shogo Soma, Tomoaki Nakazono, Kaori Shiraiwa, Ananya Dasgupta, Tatsuki Nakagawa, Jiayun L. Xie, Jasmine Chavez, Rodrigo Romo, Sandra Yungblut, Meiko Hagihara, Koshi Murata, and Kei M. Igarashi. Dopamine facilitates associative memory encoding in the entorhinal cortex. *Nature*, 598(7880):321–326, October 2021.

8. Alexandra L. Petrache, Aarib Rajulawalla, Anqi Shi, Andrea Wetzel, Takashi Saito, Takaomi C. Saido, Kirsten Harvey, and Afia B. Ali. Aberrant Excitatory-Inhibitory Synaptic Mechanisms in Entorhinal Cortex Microcircuits During the Pathogenesis of Alzheimer's Disease. *Cerebral Cortex (New York, N.Y.: 1991)*, 29(4):1834–1850, April 2019.

9. Albert Tsao, May-Britt Moser, and Edvard I. Moser. Traces of Experience in the Lateral Entorhinal Cortex. *Current Biology*, 23(5):399–405, March 2013.

10. Albert Tsao, Jørgen Sugar, Li Lu, Cheng Wang, James J. Knierim, May-Britt Moser, and Edvard I. Moser. Integrating time from experience in the lateral entorhinal cortex. *Nature*, 561(7721):57–62, September 2018.

11. Maria E. Montchal, Zachariah M. Reagh, and Michael A. Yassa. Precise temporal memories are supported by the lateral entorhinal cortex in humans. *Nature Neuroscience*, 22(2):284–288, February 2019.

12. Howard Eichenbaum, Magdalena Sauvage, Norbert Fortin, Robert Komorowski, and Paul Lipton. Towards a functional organization of episodic memory in the medial temporal lobe. *Neuroscience and Biobehavioral Reviews*, 36(7):1597–1608, August 2012.

13. Eric L. Hargreaves, Geeta Rao, Inah Lee, and James J. Knierim. Major dissociation between medial and lateral entorhinal input to dorsal hippocampus. *Science (New York, N.Y.)*, 308(5729):1792–1794, June 2005.

14. Robin M. Hayman and Kathryn J. Jeffery. How heterogeneous place cell responding arises from homogeneous grids–a contextual gating hypothesis. *Hippocampus*, 18(12):1301–1313, 2008.

15. James J. Knierim, Inah Lee, and Eric L. Hargreaves. Hippocampal place cells: Parallel input streams, subregional processing, and implications for episodic memory. *Hippocampus*, 16(9):755–764, 2006. _eprint: https://onlinelibrary.wiley.com/doi/pdf/10.1002/hipo.20203.

16. D. Yoganarasimha, Geeta Rao, and James J. Knierim. Lateral entorhinal neurons are not spatially selective in cue-rich environments. *Hippocampus*, 21(12):1363–1374, December 2011.

17. Sachin Deshmukh and James Knierim. Representation of Non-Spatial and Spatial Information in the Lateral Entorhinal Cortex. *Frontiers in Behavioral Neuroscience*, 5, 2011.

18. Joshua P. Neunuebel, D. Yoganarasimha, Geeta Rao, and James J. Knierim. Conflicts between local and global spatial frameworks dissociate neural representations of the lateral and medial entorhinal cortex. *The Journal of Neuroscience: The Official Journal of the Society for Neuroscience*, 33(22):9246–9258, May 2013.

19. W. Xu and D. A. Wilson. Odor-evoked activity in the mouse lateral entorhinal cortex. *Neuroscience*, 223:12–20, October 2012.

20. Kristin M. Kerr, Kara L. Agster, Sharon C. Furtak, and Rebecca D. Burwell. Functional neuroanatomy of the parahippocampal region: the lateral and medial entorhinal areas. *Hippocampus*, 17(9):697–708, 2007.

21. Kara L. Agster and Rebecca D. Burwell. Cortical efferents of the perirhinal, postrhinal, and entorhinal cortices of the rat. *Hippocampus*, 19(12):1159–1186, 2009. _eprint: https://onlinelibrary.wiley.com/doi/pdf/10.1002/hipo.20578.

22. L. B. Haberly and J. L. Price. Association and commissural fiber systems of the olfactory cortex of the rat. II. Systems originating in the olfactory peduncle. *The Journal of Comparative Neurology*, 181(4):781–807, October 1978.

23. L. M. Kay, L. R. Lancaster, and W. J. Freeman. Reafference and attractors in the olfactory system during odor recognition. *International Journal of Neural Systems*, 7(4):489–495, September 1996.

24. Engel Ak, Fries P, and Singer W. Dynamic predictions: oscillations and synchrony in top-down processing. *Nature reviews. Neuroscience*, 2(10), October 2001. Publisher: Nat Rev Neurosci.

25. G. Schoenbaum and H. Eichenbaum. Information coding in the rodent prefrontal cortex. I. Single-neuron activity in orbitofrontal cortex compared with that in pyriform cortex. *Journal of Neurophysiology*, 74(2):733–750, August 1995. Publisher: American Physiological Society.

26. D. P. Devanand, K. S. Michaels-Marston, X. Liu, G. H. Pelton, M. Padilla, K. Marder, K. Bell, Y. Stern, and R. Mayeux. Olfactory deficits in patients with mild cognitive impairment predict Alzheimer's disease at follow-up. *The American Journal of Psychiatry*, 157(9):1399–1405, September 2000.

27. R. I. Mesholam, P. J. Moberg, R. N. Mahr, and R. L. Doty. Olfaction in neurodegenerative disease: a meta-analysis of olfactory functioning in Alzheimer's and Parkinson's diseases. *Archives of Neurology*, 55(1):84–90, January 1998.

28. C. Murphy. Loss of olfactory function in dementing disease. *Physiology & Behavior*, 66(2):177–182, April 1999.

29. Bàrbara Segura, Hugo César Baggio, Elisabeth Solana, Eva M. Palacios, Pere Vendrell, Núria Bargalló, and Carme Junqué. Neuroanatomical correlates of olfactory loss in normal aged subjects. *Behavioural Brain Research*, 246:148–153, June 2013.

30. Maneesh V. Kuruvilla and James A. Ainge. Lateral Entorhinal Cortex Lesions Impair Local Spatial Frameworks. *Frontiers in Systems Neuroscience*, 11:30, 2017.

31. Lise Boisselier, Barbara Ferry, and Rémi Gervais. Involvement of the lateral entorhinal cortex for the formation of cross-modal olfactory-tactile associations in the rat. *Hippocampus*, 24(7):877–891, July 2014.

32. Bjorn M. Persson, Veronika Ambrozova, Stephen Duncan, Emma R. Wood, Akira R. O'Connor, and James A. Ainge. Lateral entorhinal cortex lesions impair odor-context associative memory in male rats. *Journal of Neuroscience Research*, 100(4):1030–1046, April 2022.

33. Hongjun Fu, Gustavo A. Rodriguez, Mathieu Herman, Sheina Emrani, Eden Nahmani, Geoffrey Barrett, Helen Y. Figueroa, Eliana Goldberg, S. Abid Hussaini, and Karen E. Duff. Tau Pathology Induces Excitatory Neuron Loss, Grid Cell Dysfunction and Spatial Memory Deficits Reminiscent of Early Alzheimer's Disease. *Neuron*, 93(3):533–541.e5, February 2017.

34. Heechul Jun, Allen Bramian, Shogo Soma, Takashi Saito, Takaomi C. Saido, and Kei M. Igarashi. Disrupted Place Cell Remapping and Impaired Grid Cells in a Knockin Model of Alzheimer's Disease. *Neuron*, 107(6):1095–1112.e6, September 2020.

35. Thomas Ridler, Jonathan Witton, Keith G Phillips, Andrew D Randall, and Jonathan T Brown. Impaired speed encoding and grid cell periodicity in a mouse model of tauopathy. *eLife*, 9:e59045, November 2020. Publisher: eLife Sciences Publications, Ltd.

36. Johnson Ying, Alexandra T. Keinath, Raphael Lavoie, Erika Vigneault, Salah El Mestikawy, and Mark P. Brandon. Disruption of the grid cell network in a mouse model of early Alzheimer's disease. *Nature Communications*, 13(1):886, February 2022.

37. D. Reisel, D. M. Bannerman, W. B. Schmitt, R. M. J. Deacon, J. Flint, T. Borchardt, P. H. Seeburg, and J. N. P. Rawlins. Spatial memory dissociations in mice lacking GluR1. *Nature Neuroscience*, 5(9):868–873, September 2002. Publisher: Nature Publishing Group.

38. J. N. P. Rawlins and D. S. Olton. The septo-hippocampal system and cognitive mapping. *Behavioural Brain Research*, 5(4):331–358, August 1982.

39. Andrew Aoun, Oliver Shetler, Radha Raghuraman, Gustavo A. Rodriguez, and S. Abid Hussaini. Beyond correlation: optimal transport metrics for characterizing representational stability and remapping in neurons encoding spatial memory. *Frontiers in Cellular Neuroscience*, 17, 2024.

40. Marc Aurel Busche, Gerhard Eichhoff, Helmuth Adelsberger, Dorothee Abramowski, Karl-Heinz Wiederhold, Christian Haass, Matthias Staufenbiel, Arthur Konnerth, and Olga Garaschuk. Clusters of Hyperactive Neurons Near Amyloid Plaques in a Mouse Model of Alzheimer's Disease. *Science*, 321(5896):1686–1689, September 2008. Publisher: American Association for the Advancement of Science.

41. Jorge J. Palop, Jeannie Chin, Erik D. Roberson, Jun Wang, Myo T. Thwin, Nga Bien-Ly, Jong Yoo, Kaitlyn O. Ho, Gui-Qiu Yu, Anatol Kreitzer, Steven Finkbeiner, Jeffrey L. Noebels, and Lennart Mucke. Aberrant Excitatory Neuronal Activity and Compensatory Remodeling of Inhibitory Hippocampal Circuits in Mouse Models of Alzheimer's Disease. *Neuron*, 55(5):697–711, September 2007. Publisher: Elsevier.

42. Jessica Brown, Elena Camporesi, Juan Lantero-Rodriguez, Maria Olsson, Alice Wang, Blanca Medem, Henrik Zetterberg, Kaj Blennow, Thomas K. Karikari, Mark Wall, and Emily Hill. Tau in cerebrospinal fluid induces neuronal hyperexcitability and alters hippocampal theta oscillations. *Acta Neuropathologica Communications*, 11(1):67, April 2023.

43. Gustavo A. Rodriguez, Geoffrey M. Barrett, Karen E. Duff, and S. Abid Hussaini. Chemogenetic attenuation of neuronal activity in the entorhinal cortex reduces A and tau pathology in the hippocampus. *PLOS Biology*, 18(8):e3000851, August 2020. Publisher: Public Library of Science.

44. Michael J. Yetman, Sveinung Lillehaug, Jan G. Bjaalie, Trygve B. Leergaard, and Joanna L. Jankowsky. Transgene expression in the Nop-tTA driver line is not inherently restricted to the entorhinal cortex. *Brain Structure & Function*, 221(4):2231–2249, May 2016.

45. Brianna Vandrey, Derek L. F. Garden, Veronika Ambrozova, Christina McClure, Matthew F. Nolan, and James A. Ainge. Fan Cells in Layer 2 of the Lateral Entorhinal Cortex Are Critical for Episodic-like Memory. *Current biology: CB*, 30(1):169–175.e5, January 2020.

46. Li Liu, Valerie Drouet, Jessica W. Wu, Menno P. Witter, Scott A. Small, Catherine Clelland, and Karen Duff. Trans-Synaptic Spread of Tau Pathology In Vivo. *PLOS ONE*, 7(2):e31302, February 2012. Publisher: Public Library of Science.

47. S. L. Dix and J. P. Aggleton. Extending the spontaneous preference test of recognition: evidence of object-location and object-context recognition. *Behavioural Brain Research*, 99(2):191–200, March 1999.

48. Robert M. J. Deacon and J. Nicholas P. Rawlins. T-maze alternation in the rodent. *Nature Protocols*, 1(1):7–12, 2006.

49. Syed A. Hussaini, Kimberly A. Kempadoo, Sébastien J. Thuault, Steven A. Siegelbaum, and Eric R. Kandel. Increased Size and Stability of CA1 and CA3 Place Fields in HCN1 Knockout Mice. *Neuron*, 72(4):643–653, November 2011. Publisher: Elsevier.

50. Jason E. Chung, Jeremy F. Magland, Alex H. Barnett, Vanessa M. Tolosa, Angela C. Tooker, Kye Y. Lee, Kedar G. Shah, Sarah H. Felix, Loren M. Frank, and Leslie F. Greengard. A Fully Automated Approach to Spike Sorting. *Neuron*, 95(6):1381–1394.e6, September 2017.

51. Christoph Schmidt-Hieber and Michael Häusser. Cellular mechanisms of spatial navigation in the medial entorhinal cortex. *Nature Neuroscience*, 16(3):325–331, March 2013.