A nutrient-dependent foraging phenotype of *S. coelicolor* exhibits enhanced antimicrobial activity
and is common across the genus

Niklas Söderholm[a,b], Hugh Tanner[a,b], Linda Sandblad[a,b c, #]

[a]Department of Chemistry, Umeå University, 901 87 Umeå, Sweden
[b]The Laboratory for Molecular Infection Medicine Sweden (MIMS), 901 87 Umeå, Sweden

[c]SciLifeLab at Umeå University, 901 87 Umeå, Sweden

Address correspondence to Linda Sandblad, linda.sandblad@umu.se

SUMMARY

Species of the genus *Streptomyces* are filamentous bacteria mainly residing in soils. The capacity
of streptomycetes to adapt to various environments is reflected in the sheer size of their genomes
and the number of encoded regulatory proteins. The nutrient-rich laboratory conditions
significantly differ from the natural habitat of streptomycetes and fail to stimulate their full range
of adaptive behaviors; as a result, the majority of biosynthetic gene clusters remain silent.
Moreover, the use of rich media has led to the assumption that nutrient depletion ultimately
promotes sporulation. However, we demonstrate that *S. coelicolor* colonies can respond to
nutrient depletion by undergoing a previously unidentified phenotypic transition termed
"foraging" due to its submerged and continuous growth on depleted media. The foraging
phenotype is distinctly different from conventional phenotypes in terms of colony morphology,
genomic stability, and metabolomic profile. This adaptation to nutrient deprivation through
foraging is found throughout the *Streptomyces* phylogeny, indicating that the phenotype is highly
conserved. Furthermore, foraging *S. coelicolor* gains the ability to inhibit molds and enhanced
competitive activity against both gram-negative and gram-positive bacteria is detected in other
species. These findings highlight how morphological adaptations of streptomycetes in nutrient-
limited environments alter secondary metabolite production, enabling the screening of novel
antimicrobial activities. These discoveries have implications ranging from the basic biology of
streptomycetes to drug discovery and microbial ecology.

INTRODUCTION

The gram-positive genus *Streptomyces*, known for its filamentous morphology and production of bioactive
secondary metabolites, plays a crucial role in soil as saprophytes that hydrolyze a wide array of
polysaccharides and macromolecules [1]. Streptomycetes undergo a complex life cycle, beginning with
spore germination, progressing through vegetative hyphal growth with branching to form a compact
mycelial network, giving rise to the convex colony appearance. A differentiation process leads to the
development of aerial hyphae and spores, which often coincides with production of many secondary
metabolites. The differentiation relies on the *bld*-genes to control the aerial development phase and the
whi-genes for spore maturation [2].

One of the *bld* genes, *bldA* encodes a Leu-tRNA^{UUA} necessary for translation of mRNA UUA codons [3-6]
and hence regulates the expression of genes containing this rare leucine codon. One of the genes under
this regulation is *adpA*, a global regulator of development and secondary metabolites [7-11]. Interestingly,
AdpA also controls the transcription of *bldA* [12]. The life cycle of streptomycetes has been extensively
studied on rich media, where subsequent decreasing nutrient levels are believed to promote mycelial
differentiation and secondary metabolite production. In a study using rich media, it was shown that *S.*

coelicolor metabolite production was enhanced because of genomic instability leading to a genomic

heterogeneity and a consequent lower fitness reduction and eventual extinction of these cells [13, 14].

While the importance of optimized growth media for metabolite production and life cycle progression is

widely recognized [15, 16], little is known about streptomyces growth and metabolite production under

nutrient poor conditions. An observation that aligns with its adaptation to a life as a saprophyte in the soil,

is the abundance of gene regulatory proteins, two-component systems, transporters, and substrate-

binding proteins, as first identified in the *S. coelicolor* genome, suggesting a significant capacity for

adaptation and the utilization of varied nutrient sources [17, 18]. Co-incubation of certain streptomycetes

with yeast lead to the discovery of the "exploratory phenotype", which in turn led to the understanding that

specific media components alone could induce this phenotype [19, 20].

The exploratory growth phenotype, like conventional growth, is connected to metabolite production and it

was discovered that production of cryptic metabolites was activated [19]. Identification of biosynthetic

gene clusters (BGCs), the genes coding for proteins synthesizing the metabolites, has revealed that each

Streptomyces species could have the capacity to produce over 20 secondary metabolites [17, 21-23].

However, current standard cultivation methods often fail to trigger the regulatory pathways responsible for

their synthesis [24, 25] and only a fraction of possible metabolites have consequently been studied for

biological activities. Rapidly consumed carbon sources often repress production of secondary metabolites

[26]. However, after primary substrates have been exhausted, carbon catabolite repression adapts growth

and metabolism to utilize less prioritized substrates. Therefore, altering environmental stimuli to activate

metabolite production from cryptic BGCs is a promising strategy in the search for new bioactive

compounds [27].

Considering the saprophyte niche of streptomycetes in soil environments, we sought to determine how

growth is affected when degradation of polysaccharides becomes a necessity because it is the main

source of carbon. This represents an environment that, in terms of carbon sources, would better reflect

their *in situ* growth.

We demonstrate that global nutrient depletion in growth media coincides with a morphological transition

of *S. coelicolor* colonies to a phenotype termed "foraging". Observations throughout the phylogeny

suggest that foraging is common to all streptomycetes and present in a closely related genus. We explore

this novel growth morphology, investigate antimicrobial properties, and compare metabolomic profiles of

conventional and foraging growth. This study indicates that nutrient-deprived growth may be key to both

discovering novel bioactive secondary metabolites and providing a more complete understanding of the

Streptomyces life cycle.

RESULTS

Phenotype transition correlate with nutrient depletion

To investigate how *Streptomyces* colonies respond to nutrient depletion, *S. coelicolor* was incubated on

different diluted agar media. A unique growth morphology of *S. coelicolor* was observed after 60 days of

incubation on 0.5x diluted TSA. The colonies had developed a thin, veil-like, outer growth zone, distinct

from the conventional *S. coelicolor* morphology (Figure 1A). We further investigated this morphology,

termed "foraging", using time-lapse imaging over 75 days, which revealed the transition around day 50

(Video S1).

It became clear that foraging growth was linked to nutrient availability, as it occurred earlier on diluted

media, on smaller media volumes and was delayed on richer media. The foraging transition was evident

already after 21 days when grown in smaller ⌀ 4 cm dishes with 0.5xTSA. Using these smaller dishes, we

monitored nutrient depletion by inoculating plates with spores and sampled the agar every three days for

27 days to obtain time point both pre- and post-transition. The agar was sampled 0.5 cm away from the

colony, to ensure that the samples would not be contaminated with hyphae, and the very edge of the dish. The

nutrient levels in the agar samples were analyzed using mass spectrometry. Altogether, 51 metabolites

were annotated, including sugars, amino acids, and fatty acids, most of which depleted over time (Figure

1B). Nutrient levels proximal to the colony mirrored those at the dish-edge (Figure S1A). These

observations are consistent with calculations based on agar diffusion coefficients suggest that nutrients

like e.g., glucose can diffuse 1.5 cm^2 over three days [28]. This explains the persistence of the foraging

phenotype even as it expands into unexplored areas of the agar.

Although the levels of certain metabolites, such as glucose and fructose, were decreasing from the start

and depleted at the onset of foraging, others, such as glutamine and lactic acid, initially increased before

decreasing. Some metabolites, such as leucine and phenylalanine, remained stable until day 12 before

decreasing (Figure 1C, Figure 1, Figure S1B). The levels of certain metabolites, such as sucrose,

remained constant throughout, and only five metabolites were observed to increase more than one-fold

(Figure 1C). Supplementation of 0.5xTSA and MS agar with these accumulating metabolites did not

induce foraging growth, suggesting they do not act as signals for phenotypic transition.

We conclude that the observed phenotypic transition is nutrient dependent and that the presence of

polysaccharides in the form of the agar matrix does not interfere with the foraging phenotype.

Absence of easily accessible nutrients enables foraging *S. coelicolor* growth.

The transition to the foraging phenotype coincided with media depletion; hence the growth of *S. coelicolor*

was next investigated in the absence of excessive nutrients. Even the nutrient-depleted plates clearly

contained complex carbohydrates in the form of agar at the onset of foraging. Hence, an agar medium

without additional carbon sources was sought. *Streptomyces* minimal medium contains only three carbon

sources: agar, L-asparagine, and glucose [29]. Substituting L-asparagine with ammonium sulfate and

excluding glucose from the medium made agar the sole carbon source; this medium is hereafter referred

to as foraging agar medium (FAM). Colonies on FAM germinated and displayed similar characteristics to

those on 0.5xTSA after the phenotypic transition, exhibiting a thin, "veil-like submerged expansion" that

eventually covered the entire plate (Figure 2A). Over time, lineages developed within the colony, some of

which became pigmented (Figure 2B). Aerial hyphae formation and pigment production at the inoculation

site were influenced by the presence of metabolizable nutrients such as glycerol present in the spore

storage buffer (Figure 2C), further demonstrating the effect of nutrients on the foraging phenotype. FAM

supplemented with glucose or fructose effectively inhibited foraging growth, leading to conventional raised

colonies (Figure S2A). Notably, sucrose supplementation of FAM did not prevent foraging growth,

indicating the inability of *S. coelicolor* to metabolize sucrose.

The progressive expansion and sector formation of foraging colonies on FAM were studied using time-

lapse imaging (Figure 2D, Video S2). Introduction of glucose or tryptone to a foraging colony induced

raised colony growth and strong pigment production, typical for growth in nutrient-rich conditions (Figure

2E, 2F, Video S3 and Video S4).

Agar was utilized as the main carbon source when growing on a deprived media such as FAM, and the

agar degradation was visualized by staining the plates with iodine-based Lugol's solution which revealed

the distinct halo around the colonies indicating substrate degradation (Figure 2G). Proteins were

extracted from the zones outside of foraging colonies and analyzed using mass spectrometry. Signal

peptide prediction using SignalP 6.0 [30] was performed on all *S. coelicolor* coding sequences, identifying

696 proteins with predicted signal peptides. Among the 203 proteins identified from the foraging

secretome, 56 were predicted to have a Sec translocon signal peptide, 18 had a Tat translocon signal

peptide, and 123 lacked any signal peptide (Figure 2H, Table S1). PANTHER classification of signal

peptide-containing proteins categorized them into four categories (Figure 2, Table S2). The metabolite

interconversion category featured 13 hydrolases and one oxidoreductase, while the protein modifying

enzyme category included ten proteases and a ubiquitin-protein ligase. Among the unclassified proteins

were additional hydrolases and solute binding proteins. The transporter category included the ABC

transporter BldKB, known for its role in carbon catabolite repression and mycelium formation [31, 32].

Four proteins in the foraging secretome, encoded by genes containing the *bldA*-specific TTA codon,

suggest that *bldA* is active during foraging growth: an uncharacterized protein without a signal peptide

(Q9L2F9, SCO2524), a carbohydrate-binding module (CBM) domain-containing hydrolase (Q9RKF0,

SCO3487) with a Sec-signal peptide, and two uncharacterized secreted proteins with Sec-signal peptides

(Q9RKE0, SCO3498 and Q9K452, SCO7233) (Table S3). However, a *bldA* mutant still displayed the

foraging phenotype, suggesting that BldA is active but not essential for the phenotype. The classical

regulatory proteins Whi, downstream if *bldA* in the regulatory cascade, also displayed the foraging

phenotype (Figure 4, Figure S2B). In total, 24 secretome proteins were identified in the Carbohydrate-

Active enZYmes (CAZy) Database [33] and are associated with hydrolytic and non-hydrolytic cleavage of
glycosidic bond and carbohydrate adhesion (Figure 2J).

Distinct behavioral differences between foraging and conventional phenotypes

Colonies grown on FAM and TSA were imaged in high resolution using scanning electron microscopy
(SEM) imaging to clearly visualize the two types of growths. First, the predominantly submerged growth in
FAM was confirmed. However, on FAM, patches of hyphae on the surface were found at the site of
inoculation and occasionally at other sites on the surface, while growth on TSA showed a thick layer of
hyphae piling on top of each other (Figure 3A, B). The piled-up surface growth continued to the outskirts
of the colonies (Figure 3C). In contrast, few hyphae were seen at the surface on FAM except a few
displaying a "dolphin-like" behavior (Figure 3D). Cross sections of colonies confirmed that nutrient-rich
growth resulted in limited number of hyphae inside the agar, while the majority of growth occurred inside
the agar on FAM (Figure 3E, F), with numerous of clearly distinguishable hyphae within the colony in
contrast to the agar material alone (Figure 3G, H). The high-resolution view demonstrates that, in an
environment with low levels of low-molecular-weight organic compounds and abundant polysaccharides,
the colony infiltrates the substrate, a strategy that would allow efficient colonisation and possibilities to
degrade e.g., plant debris in the soil.

To further investigate potential differences between growth at different nutrient availability, we turned the
attention to the sectors formed during foraging growth (Figure 1A, 2A, 2B, Video S1 and S2). Since
significant genomic instability in *S. coelicolor* colonies grown on rich media has been reported [13, 14], we
isolated and sequenced foraging sectors to investigate whether they are formed as a result from the
genomic instability observed on rich media. A lineage from the initial inoculation, a subsequent sector 'A',
and its subsequent sectors 'I' and 'IV', were sequenced using the Illumina platform (Figure S3A). Analysis
using breseq [34] showed no evidence of terminal deletions in foraging sectors. Point-mutations were
found in three genes but did not influence growth on rich or nutrient-deprived agar (Figure S3A). These
results emphasize that the adaptation strategies significantly differ between nutrient-rich and nutrient-
deprived environments. The absence of major genomic deletions on FAM suggests an importance to
maintain the integrity of the genome for survival in nutrient limited conditions. Altogether, these results
demonstrates that the survival strategies of *S. coelicolor* differ between nutrient-rich and nutrient-depleted
growth.

Foraging phenotype is found across the phylogeny of *Streptomyces*

To assess whether foraging growth was unique to *S. coelicolor*, the study was extended to include other
Streptomyces species representing different branches of the phylogenetic tree, including a species from

the closely related genus *Kitasatospora*. All 12 strains tested displayed the foraging growth phenotype on
FAM (Figure 4A, B), and agar degradation was confirmed by Lugol's solution staining (Figure S3B).
Similar to *S. coelicolor*, foraging growth was inhibited in all strains with the addition of 1% glucose or
tryptone to FAM (Figure S3C, D).

To investigate whether foraging was a characteristic that could be used to distinguish a streptomycetes,
local soil samples were incubated on FAM, and colonies displaying the foraging phenotype were analyzed
using MALDI-TOF to determine the species. One colony displaying the foraging phenotype was picked at
random from each of the seven soil samples. All seven colony samples were confirmed to be
streptomycetes (Table S4). These findings further strengthened the ubiquity of foraging growth across the
Streptomyces phylogeny. Until recently, it was uncertain whether *Kitasatospora* was a separate genus
from *Streptomyces but* has been elucidated since [35-37]. The fact that *K. setae* also displayed foraging
demonstrates that the phenotype is not confined to the *Streptomyces* genus but also occurs in at least
one closely related genus.

Foraging streptomycetes demonstrates increased antibacterial activity.

Since streptomycetes are known for their production of antimicrobial metabolites and also encode
numerous BGCs that are not expressed during conventional growth, the potential of foraging colonies to
inhibit bacterial growth was investigated. Due to the nutrient-depleted state of FAM, *B. subtilis* and *E. coli*
were unable to proliferate directly on the agar surface. Therefore, to test for antibacterial activity, FAM
with foraging *S. coelicolor* colonies was overlaid with diluted LA and *B. subtilis* 168 and *E. coli* MG1655
then added on top of the LA surface. The growth was evaluated after incubation using a stereo
microscope because of the smaller colonies formed dure to the slower proliferation of the indicator
bacteria on the diluted LA. This method was applied to the 12 additional *Streptomyces* species to also
assess their antibacterial properties during foraging. Most species showed increased capacity to inhibit
both indicator bacteria on nutrient-depleted FAM (11/13) compared to FAM supplemented with glucose
(FAM+G, 3/13) and tryptone (FAM+T, 1/13) (Figure 5A). However, *S. coelicolor* did not exhibit enhanced
antibacterial activity on FAM compared to the nutrient-supplemented media. Performing the same overlay
assay on *S. coelicolor* developmental mutants showed that the *bldA* mutant lost its ability to inhibit *B.*
subtilis growth, whereas the *adpA* mutant acquired the ability to inhibit *E. coli* (Figure 5B).

S. coelicolor colonies grown on FAM were found to inhibit the growth of a contaminating mold (Figure
5C), identified as *Penicillium citrinum* by 18S/ITS sequencing. This inhibition was replicated by growing *S.*
coelicolor on FAM for five days before inoculating *P. citrinum* spores at a distance of 2 cm, resulting in an
average 4.3 mm inhibition zone (±0.76 mm, n = 96). Inhibition was not observed when glucose was
added to the agar or on any other nutrient-rich media, nor on FAM with pH below 7. Interestingly, the *bldA*
mutant, typically unable to produce certain antimicrobial metabolites, inhibited *P. citrinum* growth on FAM.

However, the *adpA*-mutant strain M851 showed reduced inhibition compared to the wild-type (Figure 5D).

All tested *whi*-mutants *(-A, -B, -G, -H,* and *-I)* exhibited mold-inhibitory activity (Figure S4A). Inhibition of a

local *Aspergillus spp.* isolate was also demonstrated using an overlay assay with a 100x diluted TSA-

overlay, since the *Aspergillus spp.* did not grow on FAM alone (Figure S4B). However, no inhibition of

Saccharomyces cerevisiae or *Candida albicans* was detected using the same assay. To our knowledge,

there are no reports of antifungal activity for *S. coelicolor* A3(2), despite it being extensively studied for its

pigment and metabolite production since the 1950s [38]. To investigate whether mold inhibition was a

common property during foraging, the 12 *Streptomyces* species were tested for mold inhibition, but the

mold inhibitory activity was specific to *S. coelicolor*, as none of the other species inhibited *P. citrinum.* The

altered antimicrobial activates suggest that the foraging characteristics extend beyond genomic stability

and morphology and include significant changes to metabolite production.

Foraging *S. coelicolor* displays an altered metabolomic profile

To determine whether there were any differences in the metabolomic profiles of *S. coelicolor* grown on

FAM and FAM supplemented with 1% glucose (FAM+G), agar was sampled around the colonies for

untargeted mass spectrometric analysis. The metabolomic profiles differed between FAM and FAM+G,

both in negative (anions, R2(X)=0.75, Q2(X)=0.97) and positive detection modes (cations, R2(X)=0.63,

Q2(X)=0.47) (Figure 5E, F). Some features with relatively higher concentrations on FAM-agar (highlighted

in blue in Figure 5G, H) could be matched by monoisotopic molecular weight with known *S. coelicolor*

metabolites from the StreptomeDB [39]. Among these annotated features, the signaling molecule γ-

butyrolactone was found at higher concentrations in FAM+G. Wailupemycin G was the only annotated

metabolite found at considerably higher concentrations on FAM+G (Figure S4C, Table S5). There were

an additional 30 features, with higher levels in the FAM condition, that could not be annotated using their

monoisotopic mass and StreptomeDB (Figure S4D, E, Table S6). Overall, more metabolites features

were detected in the FAM+G samples. Twelve of these features were unique to FAM and therefore were

either present below detection limits or not produced on FAM+G. Genetic analysis of BGCs has revealed

that each *Streptomyces* species could have the capacity to produce over 20 secondary metabolites [17,

21-23]. However, standard cultivation often fails to stimulate the activation of the regulatory pathways

responsible for their synthesis [24, 25]. The mass spectrometry results demonstrate a significant

alteration in the metabolomic profile of foraging *S. coelicolor* as a consequence of the different

environmental stimuli. The gained ability of *S. coelicolor* to inhibit mold-growth together with the overall

enhanced antibacterial activities of other *Streptomyces* suggests that there are also alterations of their

metabolomes. This shows potential for using FAM growth in screening for active secondary metabolites not

produced on rich media.

DISCUSSION

The primary habitat of streptomycetes, the soil, is a dynamic environment with many variable
environmental factors. Traditional laboratory cultivation using a limited number of rich media is unlikely to
represent the full repertoire of adaptations in a diverse and competitive environment such as the soil. In
this study, we demonstrate that global nutrient depletion in growth media coincides with a morphological
transition of *S. coelicolor* colonies to a previously undescribed phenotype termed "foraging".

Media composition is known to be crucial for life-cycle progression, with nutrient depletion often referred
to as one of the triggers of differentiation and sporulation, though the process is acknowledged to be
complex [2, 40-42]. Observations of *S. coelicolor* colonies over time, combined with metabolomics
analysis, showed that the transition to the foraging phenotype on 0.5xTSA coincided with nutrient
depletion and not differentiation and subsequent sporulation. This demonstrates that nutrient-depleted
sporulation is conditional, and that growth on certain polysaccharides does not necessarily lead to
sporulation. Furthermore, it underscores that the use of conventional rich media has not revealed the full
repertoire of *S. coelicolor* adaptive regulation. This ability to continue growth in a nutrient-poor
environment allows *S. coelicolor* to avoid entering a dormant state as spores awaiting a rich supply of
nutrients, enabling continuous growth. It was recently demonstrated that secondary metabolite production
is significantly increased as a result from genetic instability on nutrient rich agar [13]. However, we could
not find any support for major mutational deletions under nutrient limited foraging growth when
investigating the sectors formed during foraging growth. This further emphasizes that the survival
strategies of *S. coelicolor* differ between nutrient-depleted and nutrient-rich agar growth.

Fluctuations in metabolite levels during growth results from the breakdown and uptake of external
sources and the subsequent release into the surroundings [43]. Four distinct trends in metabolite
dynamics were distinguishable throughout the colony growth: depletion, accumulation, initial
accumulation followed by depletion, and unchanged levels. These fluctuations of metabolite levels are the
result of breakdown, uptake, and release into the surroundings [43]. The trend of initial accumulation
followed by depletion is likely due to the breakdown of macromolecules from the TSA media and
subsequent consumption. Accumulation of metabolites may occur due to by-product overflow or overflow-
independent mechanisms, such as strong transport affinity [44]. Supplementation of the accumulated
metabolites to media did not trigger foraging growth, demonstrating that their accumulation is not
promoting the phenotype. Instead, the release of metabolites might be a deliberate strategy to interact
with other organisms in the natural habitat of *S. coelicolor*. Two accumulated metabolites, β-alanine and
pantothenic acid, are crucial in various biochemical pathways. β-alanine, essential for pantothenic acid
synthesis – a precursor of Coenzyme A – enhances respiration in alkaline soils [45].

In contrast to rich media, where low-molecular-weight organic compounds are abundant [29, 46], soil
environments typically have transient availability of such nutrients [47, 48]. Fertile soils are rich in complex
carbohydrates and plant debris and have a flux of available nutrients throughout the year to which soil

inhabitants must adapt [49]. FAM, our minimal defined agar medium, where agar acts as the sole carbon
source, supports spore germination and foraging growth, but the addition of easily metabolized nutrient
(e.g., glucose) promotes conventional colony phenotypes. This indicates that the complex
polysaccharides in agar are not preferred over simpler organic compounds. Nitrogen and phosphate also
play a crucial role in cycle progression and metabolite production. The FAM media used here, with higher
nitrogen levels than typically found in soils [47], may influence the phenotype. Elevated levels of nitrogen
in rich media have been associated with repression of secondary metabolite production [50].
Furthermore, elevated levels of inorganic phosphate have been shown to inhibit the production of several
secondary metabolites [51-53], and there has been shown that there is a significant flux of organic
nitrogen in soils using microdiffusion experiments [54-56]. Supplementation of FAM with an organic
nitrogen source (i.e., asparagine), inhibited foraging growth and promoted conventional colony growth.
Therefore, the levels and types of phosphate and nitrogen sources are also contributing to the foraging
phenotype on FAM.

The prospect of finding new metabolites produced by novel colony morphologies has been previously
demonstrated for the exploratory phenotype that certain streptomycetes exhibit when co-incubated with
yeast or when grown with specific media components alone [20, 57]. The gained ability of foraging *S.*
coelicolor to inhibit mold-growth, together with the overall enhanced potential of other *Streptomyces*
species to inhibit Gram-positive and Gram-negative bacteria, underscores a competitive advantage for
foraging colonies in nutrient-deprived conditions and suggests that the altered metabolomic profile
detected for *S. coelicolor* might be a general phenomenon. It is well established that growth conditions,
such as the choice of carbon source, affect secondary metabolism of streptomycetes [58, 59] but there
are also other strategies to activate silent BGCs [25, 60]. Therefore, there is potential for using FAM
growth in screening for biologically active secondary metabolites not produced on rich media. The low
nutrient levels used for the overlay assays, to avoid interfering with the foraging colonies, will naturally
also influence the target bacteria rendering them either more or less susceptible to the antimicrobial
effects of foraging colonies [61-63], a factor which should be considered in future studies.

The foraging secretome contained 74 proteins with predicted signal peptides. Despite the presence of
BldA-dependent proteins in the foraging secretome, a *bldA* mutant still displayed foraging growth,
demonstrating that *bldA* regulation is not essential for this phenotype. The regulatory mechanism of BldA-
dependent *adpA* translation [6] and the AdpA regulation of *bldA* [12] in combination with the activation of
downstream regulatory genes by both factors appear to be of importance for the antimicrobial activity of
the foraging phenotype. The observation that a *bldA* mutant loses the ability to inhibit *B. subtilis* growth,
while an *adpA* mutant gains the ability to inhibit *E. coli* during foraging, suggests that *S. coelicolor*
inhibition of *B. subtilis* is *bldA*-dependent, and its inhibition of *E. coli* is prevented by *adpA,* either directly
or indirectly. Conversely, mold inhibition seems to depend on *bldA* and is prevented by *adpA*. Therefore,
although *bldA* is not required for the foraging colony morphology nor mold inhibition, it is required for the

growth inhibition of *B. subtilis*. These observations highlight the complex regulatory roles of *bldA* and
adpA in antimicrobial regulating the antimicrobial activities during foraging growth.

Unlike exploratory growth, which so far has only been observed in some species [20], foraging growth
was observed in all tested *Streptomyces* species from different parts of the phylogeny, indicating it may
be a common feature across the genus. Further, *K. setae* from the closely related genus *Kitasatospora*
also displayed the foraging phenotype, demonstrating that the phenotype is not limited to streptomycetes.
The mechanism by which foraging growth is regulated remains to be elucidated. Identifying the regulatory
components would result in understanding the required nutrient deprived conditions. Importantly,
knowledge about its regulation could also be utilized as a tool for manipulating the production of the
secondary metabolites.

In summary, this study demonstrates a previously overlooked growth phenotype of *Streptomyces* with
distinctly different colony morphology and metabolite profiles compared to conventional growth on rich
media. The nutrient-deprived growth leads to a distinct phenotype and an altered metabolome, with
enhanced antimicrobial properties. The "problem of rediscovery" of known antimicrobial compounds [64],
despite the presence of numerous silent BGCs, is a significant challenge to overcome in combating
antimicrobial resistance. This novel growth behavior offers new insights into this issue and paves the way
for discovering new regulatory mechanisms underlying the foraging growth phenotype. Taken together,
this novel growth phenotype provides a new perspective on the environmental adaptations and
competitiveness of streptomycetes under nutrient-deprivation, with potential implications for antimicrobial
drug discovery.

Limitations of the study

This study highlights the potential of exploring growth conditions that better represent *in situ* growth,
specifically conditions with fewer available nutrients and dependency on the degradation of complex
carbohydrates to observe phenotypes not present on rich media.

The streptomycetes were grown on FAM, a modified version of minimal media used for *Streptomyces*
cultivation. While FAM represents a poorer growth environment compared to rich media, it remains an
artificial medium where agar is the main carbon source. In a natural setting, *Streptomyces* species
degrade a diverse array of polysaccharides, which could lead to different gene regulation and metabolic
responses. The concentrations, availability, and sources of other media components (ammonium,
phosphate, and iron) would vary in a natural setting and fluctuate over time. Field studies are essential to
confirm the ecological role of the foraging phenotype and to understand how it is affected by
environmental factors and how it affects its surroundings.

To further study phenotypes occurring in natural environments, the study of biotic and abiotic factors
would be desirable. Soil is not only different from rich media in terms of nutrient levels but also hosts a
multitude of interacting organisms, both prokaryotic and eukaryotic, origins [60, 65-67]. These biotic
interactions and abiotic factors such as pH, temperature, moisture will influence microbial development
and metabolite production and were not within the scope of this study.

Nutrient availability is the central mechanisms controlling the foraging phenotype, and even though *bldA*
and *adpA* affect the inhibitory activity of foraging colonies, the exact gene regulatory mechanisms are yet
to be determined. A comprehensive genetic and molecular analysis is necessary to elucidate the
pathways involved.

RESOURCE AVAILABILITY

Lead contact

Requests for further information and resources should be directed to and will be fulfilled by the lead
contact, Linda Sandblad (linda.sandblad@umu.se).

Materials availability

This study did not generate new unique reagents.

Data and code availability

The Illumina sequencing data, the metabolomics and nutrient mass spectrometry data has been
deposited in the SciLifeLab Data Repository (Sandblad, Linda; Söderholm, Niklas, 2024) under DOI:
https://doi.org/10.1101/2024.04.15.589617

ACKNOWLEDGMENT

We thank National Microscopy Infrastructure, NMI (VR-RFI 2016-00968) at Umeå Centre for Electron
Microscopy (UCEM). Swedish Metabolomics Centre, Umeå, Sweden
(www.swedishmetabolomicscentre.se) is acknowledged for metabolic profiling by LC-MS and GC-MS.
Illumina sequencing was provided by MicrobesNG (http://www.microbesng.com). Postdoc fellowship
funding from the Kempe foundations and Sven och Lilly Lawskis foundation. Project granted by Umeå
Centre for Microbial research (UCMR) and Swedish Foundation for Strategic Research (RIF21-0067).
Anders Sjöstedt for MALDI-TOF analysis of streptomycetes and Ala Javadi for sample preparation. We
are grateful to Cecilia Drakskog for her thorough and constructive feedback on the manuscript.

AUTHOR CONTRIBUTIONS

N.S. and L.S. conceived and designed the study. N.S. and H.T. conducted experiments. N.S. analyzed data and drafted the manuscript, and all authors revised and edited the manuscript. L.S. secured funding and supervised the project.

DECLERATION OF INTEREST

The authors declare no competing interests.

Supplemental information

Figure S1. Nutrient levels during phenotype transition on 0.5xTSA.

Figure S2. Influence of nutrients and developmental mutations on foraging phenotypes.

Figure S3. Foraging by *Streptomyces* species.

Figure S4. Mold inhibition and metabolomics of foraging *S. coelicolor*.

Table S1. Prediction of signal peptides in the foraging secretome using SignalP 6.0.

Table S2. PANTHER classification of signal peptide containing proteins with regard to protein class.

Table S3. TTA containing genes in *S. coelicolor* A3(2).

Table S4. Mass spectrometric identification of locally isolates streptomycetes.

Table S5. Metabolomic features annotated using monoisotopic mass and StreptomeDB.

Table S6. Monoisotopic mass of unannotated features significantly higher levels in FAM conditions.

Table S7. Strain list.

Video S1. *S. coelicolor* phenotype transition.

Video S2. *S. coelicolor* growing on FAM-agar.

Video S3. Addition of glucose to foraging *S. coelicolor.*

Video S4. Addition of tryptone to foraging *S. coelicolor.*

METHODS

Strains, media, and culture conditions

The strains used in this study are listed in Table S7. *Streptomyces* strains were grown on foraging agar medium (FAM) (1 g $(NH_4)_2SO_4$, 0.5 g $K_2PO_4x7H_2O$, 0.1 g $FeSO_4x7H_2O$, 10 g agar, in 1L pH 7.5) [29]. FAM were supplemented with 1% of various sugars: glucose (FAM+G), sucrose (FAM+S), fructose

(FAM+F) or tryptone as an amino acid source (FAM+T) when indicated. To investigate the effect of pH on

mold-inhibition the pH of FAM was set using HEPES-buffer and HCl/NaOH. Tryptone soy broth (Thermo

Fisher Scientific) was prepared according to instructions and diluted 0.5x when indicated with water and

1.5% agar added before autoclaving. MS-agar (20 g mannitol, 20 g Soy flour, 15 g agar in 1L) agar were

used for sporulation and petri dish time-lapse imaging. Effects of accumulated metabolites were

conducted by adding 20mg/ml: beta-alanine, meso-erythriol (Merck), Potassium-citramalate monohydrate

(Merck), urea (Merck), and pantothenic acid (Merck). All incubations were carried out at 30°C.

Isolation and identification of local isolates

The inhibited mold was isolated from the FAM dish and recultivated on TSA. The agar plate was sent to

Eurofins for 18S/ITS sequencing. For isolation of local streptomycetes, soil samples from potatoes and

plant soil were dried at room temperature before being deposited on FAM. The plates were incubated at

37°C until foraging colonies were observed after approx. 5–10 days. Colonies with characteristic foraging

phenotypes were re-streaked on TSA and characterized using MALDI-TOF (Bruker). The local Aspergillus

spp. was isolated from a soil sample and assessed morphologically.

Colony imaging

For all imaging of small colonies spores were inoculated on agar media and visualized using a Nikon

SMZ1500 dissection microscope and a Nikon Digital sight DS-Fi1 camera, with either surface illumination

using LEDs or oblique sub-stage illumination. Whole petri dishes were imagined using a Nikon D7000

DSLR. Agar time-lapses were recorded using a Raspberry Pi 3 model B with camera module v2 mounted

in a 10x lens (Olympus). The device was placed in a 30°C dark room with an LED lamp programmed to

illuminate the plate only during image capture, one frame per hour. ImageJ distribution Fiji [68] was used

to process single frames into videos and Adobe Photoshop CC software for still images.

Electron microscopy

Colonies of *S. coelicolor* on agar were cut, fixed overnight at 4°C in 2.5% glutaraldehyde/0.1M sodium

cacodylate buffer, dehydrated in series of ethanol, critical point dried and coated with a 5 nm Au/Pd

before being analyzed with a Merlin Field Emission SEM (Zeiss) using SE-HE2 and in-lens detectors.

Images were processed using Adobe Photoshop CC software.

Inhibition assays

For mold inhibition, *Streptomyces* spores were inoculated on FAM and incubated at 30°C for 5 days.

Spores from *P. citrinum* were inoculated 2 cm from the foraging colony. The plates were incubated for 14

days and observed daily for inhibition zones and imaged as described above. For bacterial inhibition,

plates with *Streptomyces* spp. were incubated for 20 days before a thin layer of 100x diluted TSA (1.5%

agar) was placed on top of agar and incubated overnight at 4°C. Overnight cultures of *E. coli* and *B.*

subtilis were diluted and $2x10^7$ CFU in 100 µl was added to the agar overlays and incubated overnight at

37°C. Inhibition zones were assessed and imaged using a Nikon SMZ1500 dissection microscope and a

Nikon Digital sight DS-Fi1 camera.

Foraging secretome protein analysis

The agar outside of foraging *S. coelicolor* on FAM incubated for 20 days was extracted. The agar was

crushed and incubated overnight with water in a 50 ml Falcon tube. Proteins in the supernatant were

precipitated by addition of TCA and incubated on ice for 4 minutes before centrifugation at 20,000g for 5

min. The pellet was washed twice with 200 μl ice-cold acetone. The pellet was dried, dissolved in protein

sample buffer and loaded on an SDS-PAGE. Protein dense regions were cut and sent for mass

spectroscopy identification to Linköping University, where general in-gel digestion was performed as

described by Shevchenko et al. (Shevchenko, Tomas, Havlis, Olsen, & Mann, 2006).

Metabolomic analysis

Metabolic profiling of media nutrients by GC-MS was performed at the Swedish Metabolomics Center in

Umeå, Sweden. For the nutrient analysis of agar, six replicates of agar plugs were collected from the

edge of the dish and 0.5 cm away from the *S. coelicolor* colonies on 0.5xTSA medium. This sampling was

performed on six different ⌀ 4 cm plates. Three control samples were taken at every time point from three

sterile 0.5xTSA dishes. Sample preparation was performed according to A et al. (A et al., 2005). The

samples were stored at -80 °C until analysis. Small aliquots of the remaining supernatants were pooled

and used to create quality control (QC) samples. The samples were analyzed in batches according to a

randomized run order on GC-MS. Derivatization and GCMS analysis were performed as described

previously A et al. (A et al., 2005). Non-processed MS-files from the metabolic analysis were exported

from the ChromaTOF software in NetCDF format to MATLAB-R2020a (Mathworks, Natick, MA, USA),

where all data pre-treatment procedures (base-line correction, chromatogram alignment, data

compression and Multivariate Curve Resolution) were performed. The extracted mass spectra were

identified by comparisons of their retention index and mass spectra with libraries of retention time indices

and mass spectra (Schauer et al., 2005). Mass spectra and retention index comparison was performed

using NIST MS 2.2 software. Annotation of mass spectra was based on reverse and forward searches in

the library. Masses and ratio between masses indicative of a derivatized metabolite were especially

notified. The mass spectrum with the highest probability indicative of a metabolite and the retention index

between the sample and library for the suggested metabolite was ± 5 the deconvoluted "peak" was

annotated as an identification of a metabolite. The annotated nutrients were plotted using GraphPad

Prism (version 9.0.2) to show trends over time and ridge plots were generated using R version 4.2 and

packages ggplot2 3.4.4 and ggridges 0.5.4.

For untargeted metabolite analysis, agar stubs were punched out from just outside of *S. coelicolor*

colonies on FAM and FAM+G. For each condition, twelve samples were isolated from plates inoculated

with *S. coelicolor*, in addition to two control samples each from sterile plates. The weight of the samples

was adjusted to 100 mg before the addition of 1 mL extraction buffer (90/10 v/v methanol:water). Internal

standards (13C9-Phenylalanine, 13C3-Caffeine was, D4-Cholic acid, D8-Arachidonic Acid, 13C9-Caffeic

Acid) were added to each sample. The sample was shaken at 30 Hz for 3 min in a mixer mill, and proteins

were precipitated at +4 °C for 2h on ice. The sample was centrifuged at +4 °C, 14 000 rpm, for 10 min.

The supernatant was transferred to a microvial and solvents evaporated. Before analysis, the sample was

re-suspended in 10 + 10 µL methanol and water. The set of samples were analyzed in positive and

negative mode. The chromatographic separation was performed on an Agilent 1290 Infinity UHPLC-

system (Agilent Technologies, Waldbronn, Germany). Re-suspended aliquots (2 µL) of the agar extracts

were injected onto an Acquity UPLC HSS T3, 2.1 x 50 mm, 1.8 µm C18 column in combination with a 2.1

mm x 5 mm, 1.8 µm VanGuard precolumn (Waters Corporation, Milford, MA, USA) held at 40°C. The

gradient elution buffers were A (H_2O, 0.1 % formic acid) and B (75/25 acetonitrile:2-propanol, 0.1 %

formic acid). The compounds were detected with an Agilent 6550 Q-TOF mass spectrometer equipped

with a jet stream electrospray ion source. The settings were kept identical between the modes, except the

capillary voltage. The data processing was performed using the Recursive Feature Extraction algorithm

within Agilent Masshunter Profinder version B.08.00 (Agilent Technologies Inc., Santa Clara, CA, USA).

All multivariate statistical investigations (PCA) were performed using the software package SIMCA®-P+

version 15.0.2 (Umetrics, Umeå, Sweden). Annotation of unknown metabolites were performed by

comparing monoisotopic molecular weights with *S. coelicolor* metabolites in the StreptomeDB (Moumbock

et al., 2021) using R software (4.2). Metabolomic features were plotted using GraphPad Prism (version

9.0.2). All raw mass spectrometry files used in this study have been deposited to ScieLifeLab Data

Repository with the dataset identifier DOI: https://doi.org/10.1101/2024.04.15.589617

Phylogenetic analysis

Phylogenetic analysis was conducted to investigate the evolutionary relationships among *Streptomyces*

species exhibiting the foraging phenotype. Phylogenomic data were obtained from the Genome

Taxonomy Database Toolkit (GTDB-Tk) release 220.0, accessed in April 2024. The phylogenetic tree file

(bac120.tree) and associated taxonomy and metadata files (bac120_taxonomy.tsv and

bac120_metadata.tsv) were imported into R version 4.2 and processed using the ape 5.8, tidyverse 2.0.0,

readr 2.1.5, and ggtree 3.12.0 packages [69-72]. Taxonomic data were processed to filter species

belonging to the genus *Streptomyces* and the species *Kitasatospora*. To ensure non-redundancy, only

one representative accession per species was included by selecting distinct species entries from the

dataset. The phylogenetic tree was pruned to include only the selected genus using the "drop.tip" function

from the ape package. The final pruned tree was visualized using the ggtree package [69]. The positions

of each species tested for foraging were annotated to visualize the phylogenetic relationships among the

selected species.

DNA sequencing of foraging sectors

Two levels of sectors formed during foraging on FAM were grown in TSB-media and 600 mg of pellet was

sent to MicrobesNG for Illumina sequencing. Quality controlled and trimmed reads with 30x coverage

were obtained and analyzed using breseq software [34] with default settings, using the *S. coelicolor* A(3)2

reference genome (Genbank: GCA_000203835.1, [17]) updated using wild-type sequence data to

accommodate for genomic variation in lab strain. The reads of descendent sectors were mapped against

this updated reference genome. Illumina read data used in this study have been deposited to SciLifeLab

Data Repository with the dataset identifier DOI: https://doi.org/10.1101/2024.04.15.589617

Statistical analysis

In this study, certain measurements in the untargeted metabolomics analysis fell below the detection

limits, resulting in missing values. To ensure an accurate representation of our findings and maintain the

integrity of the statistical analysis, these missing values were omitted from the dataset. Consequently, all

values presented in the violin plots depict observable data trends and variations above the detection

threshold. All statistical analyses were conducted using GraphPad Prism (version 9.0.2). Details on the

specific type of statistical analysis and the criteria for significance testing are provided in the legends of

the respective figures.

FIGURE LEGENDS

Figure 1. *S. coelicolor* growth phenotype transition correlates with nutrient depletion. (**A**) Colony

appearance after 75 days growth on 0.5xTSA. The point of phenotype transition is indicated by the dotted

circle. (**B**) Line plot visualization comparing nutrient proximal nutrient levels during *S. coelicolor* growth on

0.5xTSA. Each nutrient is categorized by color to indicate its trend throughout the time series. The

measurements proximal to the colony were taken in hexaduplicate for each time point and normalized

against the controls. Day 21 when the transition to foraging growth was completed is indicated with the

black dotted line. (**C**) Examples of metabolites levels throughout the study. The time point where the

transition to foraging growth was observed is indicated with the dotted lines on day 21. Each

measurement proximal to the colony and at the edge of the dish hexatuplicate were taken for each time

point and normalized against the controls.

Figure 2. Nutrient poor agar media promote the *S. coelicolor* foraging phenotype. (**A**) *S. coelicolor* spores

in glycerol storage buffer inoculated on FAM. The foraging phenotype covering the petri dish (⌀ 8.8 cm)

was photographed after 15 weeks of incubation at 30°C. Transparent pigment-free and lightly pigmented

sectors are visible. (**B**) A photograph of a pigmented sector extending towards the edge of the dish. (**C**)

Formation of aerial hyphae were only observed at the site of spore inoculation. (**D**) Time-lapse stills

showing the foraging growth progression at day 12 and 52. (**E**) Time-lapse stills after addition of glucose

(+G) to a foraging colony at day 0 and 12. (**F**) Time-lapse stills of addition of tryptone (+T) to a foraging

colony at day 0 and 12. (**G**) Micrograph of foraging *S. coelicolor* grown on FAM, stained with Lugol's

solution. (**H**) Number of proteins containing a signal sequence in the foraging secretome, predicted using

SignalP 6.0 [30]. Categories include general secretory (Sec) substrates cleaved by SPase I (Sec/SPI) or

SPase II (Sec/SPII, lipoproteins) and twin-arginine substrates cleaved by SPase I (Tat/SPI) or SPase II

(Tat/SPII, lipoprotein), and "No SP" for no predicted signal peptide. (**I**) PANTHER functional classification

of proteins from the foraging secretome with predicted signal peptides. The graph shows the number of

proteins assigned to each protein class. (**J**) CAZy classification of the foraging secretome: Glycoside

Hydrolases (GH), Polysaccharide Lyases (PLs) and Carbohydrate-Binding Modules (CBMs).

Figure 3. Foraging hyphae are primarily found in the agar substrate. SEM imaging of *S. coelicolor* grown

on FAM and 0.5xTSA for 7 days. The glycerol storage buffer was washed away and replaced with water.

(**A**) Top view of the center of a colony grown on TSA. (**B**) Top view of the center of a foraging colony. (**C**)

Top view of the edge of a TSA colony. (**D**) Top view of the edge of a foraging colony shows few hyphae at

the surface of the FAM. (**E**) Cross section of a TSA-colony show hyphae predominantly grow on top on

0.5xTSA. (**F**) Cross section of a foraging colony shows that hyphae extending through the agar are mainly

observed on the FAM-plates. Black dotted lines indicate the agar surface in the cross sections. White

dotted lines encompass where hyphae are found. (**G**) A magnified view shows hyphae inside the FAM.

(**H**) A magnified view of FAM showing absence of hyphae outside of the foraging colony.

Figure 4. Phylogenetic tree of the Streptomyces and Kitasatospora genera. (**A**) The phylogenetic tree,

displayed as a half-circle, was generated using GTDB-Tk and pruned to include only the *Streptomyces*

(blue) and *Kitasatospora* (green) genera. The positions of the species tested for foraging phenotype are

indicated with a point on the branch and the species name. Species confirmed to exhibit the foraging

phenotype are indicated by points on their respective branches, along with their NCBI species names. (**B**)

Microscopy images showing the foraging phenotype of each *Streptomyces spp.* cultivated for three weeks

on FAM on the right side of the tree. Abbreviations: SANT; *S. antibioticus*, SANU; *S. anulatus*, SCO; *S.*

coelicolor, SFI; *S. filipinensis*, SGR; *S. griseus*, SGRI; *S. griseolus*, SHIR; *S. hiroshimensis*, SHYG; *S.*

hygroscopicus, SKAN; *S. kanamyceticus*, SLIV; *S. lividans*, SNOR; *S. norsei*, SVEN; *S. venezuelae*,

KSET; *Kitasatospora setae*.

Figure 5. Metabolome and antimicrobial activity of foraging colonies. (**A**) Bactericidal activity of foraging

streptomycetes determined by overlay-assays with indicator bacteria (Ec=*E. coli*, Bs=*Bacillus subtilis*) on

FAM, FAM+G and FAM+T. (**B**) Bactericidal activity of *S. coelicolor* developmental mutants as determined

by overlay assays on FAM. Dark purple indicates inhibition of indicator bacteria, light purple indicates

partial growth inhibition, and grey indicates that there was no detectable effect. (**C**) Inhibition zones

around *S. coelicolor* on FAM contaminated by *P. citrinum* mold. (**D**) Inhibition zones around *S. coelicolor*

wild-type (WT), bldA and M851 (*adpA*-mutant) inoculated with *P. citrinum*. (**E**) Scores plot of LC-MS

analysis in negative detection mode of FAM and FAM+G-agar inoculated with *S. coelicolor*. (**F**) Loadings

plot from negative detection mode. Features with relatively elevated levels on FAM are highlighted in blue

and features with relatively elevated levels in FAM+G-agar are highlighted in red. (**G**) Scores plot of LC-

MS analysis in positive detection mode of FAM and FAM+G-agar inoculated with *S. coelicolor*. (**H**)

Loadings plot from positive detection mode. Features with relatively elevated levels on FAM-agar are

highlighted in blue and features with relatively elevated levels in FAM+G-agar are highlighted in red.

Supplemental information

Figure S1. Nutrient levels during phenotype transition on 0.5xTSA. (**A**) The fold change for every nutrient

at every time point is plotted for both sampled positions on the dish. The plot reveals a highly positive

correlation, as indicated by the R-value of >0.96 suggesting a significant linear relationship. (**B**) The fold

change for each annotated metabolite during phenotype transition on 0.5xTSA is overlayed for both

positions on the dish. The time point where the transition was observed is indicated with the dotted lines

at day 21.

Figure S2. Influence of nutrients and developmental mutations on foraging phenotypes. (**A**) Micrograph

showing the phenotypic growth on WT *S. coelicolor* inoculated on FAM, FAM+sucrose (+S),

FAM+glucose (+G) and FAM+fructose (+F) for 15 weeks at 30°C. (**B**) Growth of *S. coelicolor* wild-types

(M145 and M600) and developmental mutants: *bldA*, *whiA* (J2401), *whiB* (J2402), *whiG* (J2400), *whiH*

(J2408), *whiI* (J2450), and M851 (Δ*adpA*) on FAM.

Figure S3. Foraging by *Streptomyces* species. (**A**) Sector point mutations of foraging colonies. Spores of

S. coelicolor were inoculated on FAM-agar and the sectors were isolated and subjected to Illumina

sequencing. The inoculum was compared to the first level sector formed (α) and two succeeding sectors

(I and IV) originating from lineage α. Using the breseq software, two point-mutations were detected in

sector α: DnaA (SCO3879:E520K) and uncharacterized DUF349-containing protein (SCO1511:+G). One

mutation was found in SCO5585 $(CG)_{5\rightarrow4}$ in linage IV. (**B**) Streptomycetes on FAM stained with Lugol's

solution after 5 days of incubation. Abbreviations: SANT; *S. antibioticus*, SANU; *S. anulatus*, SCO; *S.*

coelicolor, SFI; *S. filipinensis*, SGR; *S. griseus*, SGRI; *S. griseolus*, SHIR; *S. hiroshimensis*, SHYG; *S.*

hygroscopicus, SKAN; *S. kanamyceticus*, SLIV; *S. lividans*, SNOR; *S. norsei*, SVEN; *S. venezuelae*,

KSET; *Kitasatospora setae*. (**C**) Micrographs of the tested streptomycetes grown on FAM+G for 14 days.

(**D**) Micrographs of the tested streptomycetes grown on FAM+T for 14 days.

Figure S4. Mold inhibition and metabolomics of foraging *S. coelicolor*. (**A**) Inhibition of *P. citrinum by S.*

coelicolor developmental mutants: *whiA* (J2401), *whiB* (J2402), *whiG* (J2400), *whiH* (J2408), *whiI* (J2450)

and wild-type strain M600. (**B**) Inhibition of local *Aspergillus*-isolate in an overlay-assay on foraging *S.*

coelicolor. Pictures were taken of foraging colonies using a dissection microscope. (**C**) Graphs showing

the features annotated by monoisotopic mass and StreptomeDB. All graphs depict statistically significant

outcomes comparing FAM and FAM+G as determined by t-tests, with p-values <=0.0002. 7-Acetyl-3.6-

dihydroxy-8-methyl-tetralone is abbreviated as 7A36D8MT in the plot. (**D**) Venn-diagram showing the

number of shared and unique metabolites detected during growth on FAM and FAM+G. **(E)** Plots of the

unannotated metabolomic features unique to growth on FAM. The top right corner of each graph indicates

the molecular weight of corresponding metabolite. All graphs depict statistically significant outcomes as

determined by t-tests, with p-values <0.0001.

REFERENCES

1.   Chater, K.F., Biro, S., Lee, K.J., Palmer, T., and Schrempf, H. (2010). The complex extracellular
biology of Streptomyces. FEMS Microbiol Rev *34*, 171-198.
2.   McCormick, J.R., and Flardh, K. (2012). Signals and regulators that govern Streptomyces
development. FEMS Microbiol Rev *36*, 206-231.
3.   Kalan, L., Gessner, A., Thaker, M.N., Waglechner, N., Zhu, X., Szawiola, A., Bechthold, A.,
Wright, G.D., and Zechel, D.L. (2013). A cryptic polyene biosynthetic gene cluster in
Streptomyces calvus is expressed upon complementation with a functional bldA gene. Chem Biol
20, 1214-1224.
4.   Leskiw, B.K., Bibb, M.J., and Chater, K.F. (1991). The use of a rare codon specifically during
development? Mol Microbiol *5*, 2861-2867.
5.   Wolanski, M., Donczew, R., Kois-Ostrowska, A., Masiewicz, P., Jakimowicz, D., and Zakrzewska-
Czerwinska, J. (2011). The level of AdpA directly affects expression of developmental genes in
Streptomyces coelicolor. J Bacteriol *193*, 6358-6365.
6.   Takano, E., Tao, M., Long, F., Bibb, M.J., Wang, L., Li, W., Buttner, M.J., Bibb, M.J., Deng, Z.X.,
and Chater, K.F. (2003). A rare leucine codon in adpA is implicated in the morphological defect of
bldA mutants of Streptomyces coelicolor. Mol Microbiol *50*, 475-486.
7.   Guyet, A., Benaroudj, N., Proux, C., Gominet, M., Coppee, J.Y., and Mazodier, P. (2014).
Identified members of the Streptomyces lividans AdpA regulon involved in differentiation and
secondary metabolism. BMC Microbiol *14*, 81.
8.   Higo, A., Hara, H., Horinouchi, S., and Ohnishi, Y. (2012). Genome-wide distribution of AdpA, a
global regulator for secondary metabolism and morphological differentiation in Streptomyces,
revealed the extent and complexity of the AdpA regulatory network. DNA Res *19*, 259-273.
9.   Ohnishi, Y., Yamazaki, H., Kato, J.Y., Tomono, A., and Horinouchi, S. (2005). AdpA, a central
transcriptional regulator in the A-factor regulatory cascade that leads to morphological
development and secondary metabolism in Streptomyces griseus. Biosci Biotechnol Biochem *69*,
431-439.
10.   Yamazaki, H., Tomono, A., Ohnishi, Y., and Horinouchi, S. (2004). DNA-binding specificity of
AdpA, a transcriptional activator in the A-factor regulatory cascade in Streptomyces griseus. Mol
Microbiol *53*, 555-572.

11. Nguyen, K.T., Tenor, J., Stettler, H., Nguyen, L.T., Nguyen, L.D., and Thompson, C.J. (2003).
Colonial differentiation in Streptomyces coelicolor depends on translation of a specific codon
within the adpA gene. J Bacteriol *185*, 7291-7296.
12. Higo, A., Horinouchi, S., and Ohnishi, Y. (2011). Strict regulation of morphological differentiation
and secondary metabolism by a positive feedback loop between two global regulators AdpA and
BldA in Streptomyces griseus. Mol Microbiol *81*, 1607-1622.
13. Zhang, Z., Du, C., de Barsy, F., Liem, M., Liakopoulos, A., van Wezel, G.P., Choi, Y.H.,
Claessen, D., and Rozen, D.E. (2020). Antibiotic production in Streptomyces is organized by a
division of labor through terminal genomic differentiation. Sci Adv *6*, eaay5781.
14. Zhang, Z., Shitut, S., Claushuis, B., Claessen, D., and Rozen, D.E. (2022). Mutational meltdown
of putative microbial altruists in Streptomyces coelicolor colonies. Nat Commun *13*, 2266.
15. van Wezel, G.P., and McDowall, K.J. (2011). The regulation of the secondary metabolism of
Streptomyces: new links and experimental advances. Nat Prod Rep *28*, 1311-1333.
16. Singh, V., Haque, S., Niwas, R., Srivastava, A., Pasupuleti, M., and Tripathi, C.K. (2016).
Strategies for Fermentation Medium Optimization: An In-Depth Review. Front Microbiol *7*, 2087.
17. Bentley, S.D., Chater, K.F., Cerdeno-Tarraga, A.M., Challis, G.L., Thomson, N.R., James, K.D.,
Harris, D.E., Quail, M.A., Kieser, H., Harper, D., et al. (2002). Complete genome sequence of the
model actinomycete Streptomyces coelicolor A3(2). Nature *417*, 141-147.
18. Nikolaidis, M., Hesketh, A., Frangou, N., Mossialos, D., Van de Peer, Y., Oliver, S.G., and
Amoutzias, G.D. (2023). A panoramic view of the genomic landscape of the genus Streptomyces.
Microb Genom *9*.
19. Shepherdson, E.M.F., and Elliot, M.A. (2022). Cryptic specialized metabolites drive Streptomyces
exploration and provide a competitive advantage during growth with other microbes. Proc Natl
Acad Sci U S A *119*, e2211052119.
20. Jones, S.E., Ho, L., Rees, C.A., Hill, J.E., Nodwell, J.R., and Elliot, M.A. (2017). Streptomyces
exploration is triggered by fungal interactions and volatile signals. Elife *6*.
21. Ikeda, H., Ishikawa, J., Hanamoto, A., Shinose, M., Kikuchi, H., Shiba, T., Sakaki, Y., Hattori, M.,
and Omura, S. (2003). Complete genome sequence and comparative analysis of the industrial
microorganism Streptomyces avermitilis. Nat Biotechnol *21*, 526-531.
22. Ohnishi, Y., Ishikawa, J., Hara, H., Suzuki, H., Ikenoya, M., Ikeda, H., Yamashita, A., Hattori, M.,
and Horinouchi, S. (2008). Genome sequence of the streptomycin-producing microorganism
Streptomyces griseus IFO 13350. J Bacteriol *190*, 4050-4060.
23. Qin, Z., Munnoch, J.T., Devine, R., Holmes, N.A., Seipke, R.F., Wilkinson, K.A., Wilkinson, B.,
and Hutchings, M.I. (2017). Formicamycins, antibacterial polyketides produced by Streptomyces
formicae isolated from African Tetraponera plant-ants. Chem Sci *8*, 3218-3227.
24. Nett, M., Ikeda, H., and Moore, B.S. (2009). Genomic basis for natural product biosynthetic
diversity in the actinomycetes. Nat Prod Rep *26*, 1362-1384.
25. Ochi, K., and Hosaka, T. (2013). New strategies for drug discovery: activation of silent or weakly
expressed microbial gene clusters. Appl Microbiol Biotechnol *97*, 87-98.
26. Romero-Rodriguez, A., Rocha, D., Ruiz-Villafan, B., Guzman-Trampe, S., Maldonado-Carmona,
N., Vazquez-Hernandez, M., Zelarayan, A., Rodriguez-Sanoja, R., and Sanchez, S. (2017).
Carbon catabolite regulation in Streptomyces: new insights and lessons learned. World J
Microbiol Biotechnol *33*, 162.
27. Zhang, X., Hindra, and Elliot, M.A. (2019). Unlocking the trove of metabolic treasures: activating
silent biosynthetic gene clusters in bacteria and fungi. Curr Opin Microbiol *51*, 9-15.
28. Miyamoto, S., Atsuyama, K., Ekino, K., and Shin, T. (2018). Estimating the Diffusion Coefficients
of Sugars Using Diffusion Experiments in Agar-Gel and Computer Simulations. Chem Pharm Bull
(Tokyo) *66*, 632-636.
29. Kieser, T., Bibb, M.J., Buttner, M.J., Chater, K.F., and Hopwood, D.A. (2000). Practical
Streptomyces Genetics, (John Innes Foundation).
30. Teufel, F., Almagro Armenteros, J.J., Johansen, A.R., Gislason, M.H., Pihl, S.I., Tsirigos, K.D.,
Winther, O., Brunak, S., von Heijne, G., and Nielsen, H. (2022). SignalP 6.0 predicts all five types
of signal peptides using protein language models. Nat Biotechnol *40*, 1023-1025.
31. Chavez, A., Forero, A., Sanchez, M., Rodriguez-Sanoja, R., Mendoza-Hernandez, G., Servin-
Gonzalez, L., Sanchez, B., Garcia-Huante, Y., Rocha, D., Langley, E., et al. (2011). Interaction of

SCO2127 with BldKB and its possible connection to carbon catabolite regulation of morphological
differentiation in Streptomyces coelicolor. Appl Microbiol Biotechnol *89*, 799-806.
32. Nodwell, J.R., and Losick, R. (1998). Purification of an extracellular signaling molecule involved in
production of aerial mycelium by Streptomyces coelicolor. J Bacteriol *180*, 1334-1337.
33. Drula, E., Garron, M.L., Dogan, S., Lombard, V., Henrissat, B., and Terrapon, N. (2022). The
carbohydrate-active enzyme database: functions and literature. Nucleic Acids Res *50*, D571-
D577.
34. Deatherage, D.E., and Barrick, J.E. (2014). Identification of mutations in laboratory-evolved
microbes from next-generation sequencing data using breseq. Methods Mol Biol *1151*, 165-188.
35. Labeda, D.P., Goodfellow, M., Brown, R., Ward, A.C., Lanoot, B., Vanncanneyt, M., Swings, J.,
Kim, S.B., Liu, Z., Chun, J., et al. (2012). Phylogenetic study of the species within the family
Streptomycetaceae. Antonie Van Leeuwenhoek *101*, 73-104.
36. Labeda, D.P., Dunlap, C.A., Rong, X., Huang, Y., Doroghazi, J.R., Ju, K.S., and Metcalf, W.W.
(2017). Phylogenetic relationships in the family Streptomycetaceae using multi-locus sequence
analysis. Antonie Van Leeuwenhoek *110*, 563-583.
37. Komaki, H. (2023). Recent Progress of Reclassification of the Genus Streptomyces.
Microorganisms *11*.
38. Hopwood, D.A. (2019). Highlights of Streptomyces genetics. Heredity (Edinb) *123*, 23-32.
39. Moumbock, A.F.A., Gao, M., Qaseem, A., Li, J., Kirchner, P.A., Ndingkokhar, B., Bekono, B.D.,
Simoben, C.V., Babiaka, S.B., Malange, Y.I., et al. (2021). StreptomeDB 3.0: an updated
compendium of streptomycetes natural products. Nucleic Acids Res *49*, D600-D604.
40. Bush, M.J., Tschowri, N., Schlimpert, S., Flardh, K., and Buttner, M.J. (2015). c-di-GMP signalling
and the regulation of developmental transitions in streptomycetes. Nat Rev Microbiol *13*, 749-760.
41. Flardh, K., and Buttner, M.J. (2009). Streptomyces morphogenetics: dissecting differentiation in a
filamentous bacterium. Nat Rev Microbiol *7*, 36-49.
42. Barka, E.A., Vatsa, P., Sanchez, L., Gaveau-Vaillant, N., Jacquard, C., Meier-Kolthoff, J.P.,
Klenk, H.P., Clement, C., Ouhdouch, Y., and van Wezel, G.P. (2016). Taxonomy, Physiology,
and Natural Products of Actinobacteria. Microbiol Mol Biol Rev *80*, 1-43.
43. Douglas, A.E. (2020). The microbial exometabolome: ecological resource and architect of
microbial communities. Philos Trans R Soc Lond B Biol Sci *375*, 20190250.
44. Pinu, F.R., Granucci, N., Daniell, J., Han, T.L., Carneiro, S., Rocha, I., Nielsen, J., and Villas-
Boas, S.G. (2018). Metabolite secretion in microorganisms: the theory of metabolic overflow put
to the test. Metabolomics *14*, 43.
45. Datta, R., Baraniya, D., Wang, Y.-F., Kelkar, A., Meena, R.S., Yadav, G.S., Teresa Ceccherini,
M., and Formanek, P. (2017). Amino Acid: Its Dual Role as Nutrient and Scavenger of Free
Radicals in Soil. Sustainability *9*, 1402.
46. Shirling, E.B., and Gottlieb, D. (1966). Methods for characterization of Streptomyces species.
International journal of systematic bacteriology *16*, 313-340.
47. Rousk, J., and Jones, D.L. (2010). Loss of low molecular weight dissolved organic carbon (DOC)
and nitrogen (DON) in H2O and 0.5 M K2SO4 soil extracts. Soil biology & biochemistry *42*, 2331-
2335.
48. Jones, D.L., and Murphy, D.V. (2007). Microbial response time to sugar and amino acid additions
to soil. Soil biology & biochemistry *39*, 2178-2182.
49. Randewig, D., Marshall, J.D., Näsholm, T., and Jämtgård, S. (2019). Combining microdialysis
with metabolomics to characterize the in situ composition of dissolved organic compounds in
boreal forest soil. Soil Biology and Biochemistry *136*, 107530.
50. Hodgson, D.A. (2000). Primary metabolism and its control in streptomycetes: A most unusual
group of bacteria. In Advances in Microbial Physiology, Volume 42. (Academic Press), pp. 47-
238.
51. Bibb, M.J. (2005). Regulation of secondary metabolism in streptomycetes. Curr Opin Microbiol *8*,
208-215.
52. Hobbs, G., Obanye, A.I., Petty, J., Mason, J.C., Barratt, E., Gardner, D.C., Flett, F., Smith, C.P.,
Broda, P., and Oliver, S.G. (1992). An integrated approach to studying regulation of production of
the antibiotic methylenomycin by Streptomyces coelicolor A3(2). J Bacteriol *174*, 1487-1494.

53.   Martin, J.F. (2004). Phosphate control of the biosynthesis of antibiotics and other secondary
798   metabolites is mediated by the PhoR-PhoP system: an unfinished story. J Bacteriol *186*, 5197-
799   5201.
800   54.   Inselsbacher, E., and Nasholm, T. (2012). The below-ground perspective of forest plants: soil
801   provides mainly organic nitrogen for plants and mycorrhizal fungi. The New phytologist *195*, 329-
802   334.
803   55.   Inselsbacher, E., Öhlund, J., Jämtgård, S., Huss-Danell, K., and Näsholm, T. (2011). The
804   potential of microdialysis to monitor organic and inorganic nitrogen compounds in soil. Soil
805   biology & biochemistry *43*, 1321-1332.
806   56.   Leitner, S., Minixhofer, P., Inselsbacher, E., Keiblinger, K.M., Zimmermann, M., and Zechmeister-
807   Boltenstern, S. (2017). Short-term soil mineral and organic nitrogen fluxes during moderate and
808   severe drying–rewetting events. Applied soil ecology : a section of Agriculture, ecosystems &
809   environment *114*, 28-33.
810   57.   Shepherdson, E.M.F., Netzker, T., Stoyanov, Y., and Elliot, M.A. (2022). Exploratory Growth in
811   Streptomyces venezuelae Involves a Unique Transcriptional Program, Enhanced Oxidative
812   Stress Response, and Profound Acceleration in Response to Glycerol. J Bacteriol *204*,
813   e0062321.
814   58.   Sanchez, S., Chavez, A., Forero, A., Garcia-Huante, Y., Romero, A., Sanchez, M., Rocha, D.,
815   Sanchez, B., Avalos, M., Guzman-Trampe, S., et al. (2010). Carbon source regulation of
816   antibiotic production. J Antibiot (Tokyo) *63*, 442-459.
817   59.   van Keulen, G., and Dyson, P.J. (2014). Production of Specialized Metabolites by Streptomyces
818   coelicolor A3(2). Advances in Applied Microbiology *89*, 217-266.
819   60.   van der Meij, A., Worsley, S.F., Hutchings, M.I., and van Wezel, G.P. (2017). Chemical ecology of
820   antibiotic production by actinomycetes. FEMS Microbiol Rev *41*, 392-416.
821   61.   Liu, H., Wang, Y., Zhang, Z., Qi, H., Zhang, Y., Li, W., Shi, Q., and Xie, X. (2023). Nutrient
822   condition modulates the antibiotic tolerance of Pseudomonas aeruginosa. Sci Total Environ *904*,
823   166749.
824   62.   Windels, E.M., Cool, L., Persy, E., Swinnen, J., Matthay, P., Van den Bergh, B., Wenseleers, T.,
825   and Michiels, J. (2024). Antibiotic dose and nutrient availability differentially drive the evolution of
826   antibiotic resistance and persistence. ISME J *18*.
827   63.   Nguyen, D., Joshi-Datar, A., Lepine, F., Bauerle, E., Olakanmi, O., Beer, K., McKay, G., Siehnel,
828   R., Schafhauser, J., Wang, Y., et al. (2011). Active starvation responses mediate antibiotic
829   tolerance in biofilms and nutrient-limited bacteria. Science *334*, 982-986.
830   64.   Hutchings, M.I., Truman, A.W., and Wilkinson, B. (2019). Antibiotics: past, present and future.
831   Curr Opin Microbiol *51*, 72-80.
832   65.   Seipke, R.F., Barke, J., Heavens, D., Yu, D.W., and Hutchings, M.I. (2013). Analysis of the
833   bacterial communities associated with two ant-plant symbioses. Microbiologyopen *2*, 276-283.
834   66.   Worsley, S.F., Newitt, J., Rassbach, J., Batey, S.F.D., Holmes, N.A., Murrell, J.C., Wilkinson, B.,
835   and Hutchings, M.I. (2020). Streptomyces Endophytes Promote Host Health and Enhance
836   Growth across Plant Species. Appl Environ Microbiol *86*.
837   67.   Seipke, R.F., Kaltenpoth, M., and Hutchings, M.I. (2012). Streptomyces as symbionts: an
838   emerging and widespread theme? FEMS Microbiol Rev *36*, 862-876.
839   68.   Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T., Preibisch, S.,
840   Rueden, C., Saalfeld, S., Schmid, B., et al. (2012). Fiji: an open-source platform for biological-
841   image analysis. Nat Methods *9*, 676-682.
842   69.   Yu, G., Smith, D.K., Zhu, H., Guan, Y., and Lam, T.T.-Y. (2017). ggtree: an r package for
843   visualization and annotation of phylogenetic trees with their covariates and other associated data.
844   Methods in Ecology and Evolution *8*, 28-36.
845   70.   Wickham, H., Hester, J., and Bryan, J. (2024). readr: Read Rectangular Text Data.
846   71.   Wickham, H., Averick, M., Bryan, J., Chang, W., McGowan, L., François, R., Grolemund, G.,
847   Hayes, A., Henry, L., Hester, J., et al. (2019). Welcome to the Tidyverse. Journal of open source
848   software *4*, 1686.
849   72.   Paradis, E., and Schliep, K. (2019). ape 5.0: an environment for modern phylogenetics and
850   evolutionary analyses in R. Bioinformatics *35*, 526-528.

Figure 1. *S. coelicolor* growth phenotype transition correlates with nutrient depletion.

Figure 2. Absence of easily accessible nutrients enables *S. coelicolor* foraging growth.

Figure 3. Foraging hyphae are primarily found in the agar substrate.

Figure 4. Phylogenetic tree of the *Streptomyces* and *Kitasatospora* genera.

Figure 5. Metabolome and antimicrobial activity of foraging colonies.