Nutrient-deprived growth of streptomycetes reveals a novel growth phenotype with

enhanced antimicrobial activity

Niklas Söderholm[a,b], Hugh Tanner[a,b], Linda Sandblad[a,b]*#

[a]Department of Chemistry, Umeå University, 901 87 Umeå, Sweden

[b]The Laboratory for Molecular Infection Medicine Sweden (MIMS), 901 87 Umeå, Sweden

Address correspondence to Linda Sandblad, linda.sandblad@umu.se

Abstract

The capacity of gram-positive streptomycetes to adapt to various environments is reflected in the sheer size of their genomes and the number of encoded regulatory proteins. However, typical studies of Streptomycetes are conducted using media rich in available nutrients. Here, we demonstrate that *S. coelicolor* colonies respond to nutrient depletion with a phenotype transition. We refer to this phenotype as "foraging growth" for its submerged and continuous growth on depleted media. The foraging phenotype differ distinctly from colonies on nutrient-rich media, in terms of colony morphology, genomic stability and their secretome. The results demonstrate that adaption to nutrient deprivation through foraging is found throughout the phylogeny, indicating it is highly conserved. Furthermore, foraging *S. coelicolor* gains the ability to inhibit mold on deprived media and exhibits an altered metabolomic profile. An enhanced competitive activity against gram-negative and gram-positive bacteria was also observed among the tested species. These findings highlight the morphological adaptions of streptomycetes during nutrient deprived growth and demonstrate altered secondary metabolite production and that novel antimicrobial activities can be detected.

Introduction

The gram-positive genus *Streptomyces*, known for its filamentous morphology and production of bioactive secondary metabolites, plays a crucial role in soil as saprophytes hydrolyzing a wide array of polysaccharides and macromolecules (Chater, Biro, Lee, Palmer, & Schrempf, 2010). Streptomycetes undergo a complex life cycle, beginning with spore germination, progressing through vegetative hyphal growth and branching to form a mycelial network. A differentiation process leads to the development of aerial hyphae and spores, which often coincide with activation of secondary metabolites. The differentiation relies on the *bld*-genes to control aerial development and the *whi*-genes for spore maturation (McCormick & Flardh, 2012). The life cycle of streptomycetes has

extensively been studied on rich media where subsequent nutrient depletion is thought to contribute

to mycelial differentiation and secondary metabolite production.

While the importance of optimized growth medium for metabolite production and life cycle

progression is widely recognized, few have examined nutrient poor conditions. The abundance of gene

regulatory proteins, two-component systems, transporters, and substrate-binding proteins identified

in the *S. coelicolor* genome suggests a significant capacity for adaptation and the utilization of varied

nutrient sources (Bentley et al., 2002; Nikolaidis et al., 2023). A recent study on rich media showed

that *S. coelicolor* metabolite production was enhanced because of genomic heterogeneity (Z. Zhang

et al., 2020). Additionally, certain streptomycetes exhibit an "exploratory phenotype" when co-

incubated with yeast or by specific media components alone (S. E. Jones et al., 2017; Shepherdson &

Elliot, 2022). As for conventional growth, exploratory growth is closely connected to metabolite

production and activates cryptic metabolites (Shepherdson & Elliot, 2022).

Genetic identification of biosynthetic gene clusters (BGCs) has revealed that each *Streptomyces spp.*

could have the capacity to produce over 20 secondary metabolites (Bentley et al., 2002; Ikeda et al.,

2003; Ohnishi et al., 2008; Qin et al., 2017). However, standard cultivation often fails to stimulate the

activation of the regulatory pathways responsible for their synthesis (Nett, Ikeda, & Moore, 2009; Ochi

& Hosaka, 2013). After exhausting primary substrates, carbon catabolite repression enables the

adaptation of growth and metabolism to utilize to less prioritized substrates. Rapidly consumed

carbon sources are often found to repress production of secondary metabolites (Romero-Rodriguez

et al., 2017). Therefore, altering environmental stimuli to activate silent secondary metabolite

production is a promising strategy in the search for new bioactive compounds (X. Zhang, Hindra, &

Elliot, 2019).

In this study, we demonstrate that global nutrient depletion in growth media coincides with a

morphological transition of *S. coelicolor* colonies to a phenotype termed "foraging". Our findings

reveal that *bldA* is not essential for foraging growth, although TTA codon-containing gene products

were detected in the foraging secretome. Additionally*, S. coelicolor* exhibited mold-inhibitory activity

under nutrient-depleted conditions. Metabolomic analysis showed significant changes in the

metabolome, and growth inhibition assays suggested roles for *bldA* and *adpA* in antimicrobial activity

during nutrient-deprived growth. These results underscore a competitive advantage for foraging

colonies on nutrient-deprived media. Furthermore, all strains cultured under these conditions

exhibited foraging growth, suggesting this behavior is evolutionarily conserved. This study indicates

that nutrient-deprived growth may be a key method to both generate novel bioactive secondary

metabolites in addition to providing a more complete understanding of the life cycle.

Results

Phenotype transition from conventional *S. coelicolor* colony morphology correlate with

nutrient depletion.

In a phenotypic screening of different media formulation, we observed a unique growth morphology

of *S. coelicolor* after 60 days on 0.5x diluted TSA medium. The colonies had developed an unusual

thinner, veil-like, outer growth zone, distinct from the conventional *S. coelicolor* morphology (Fig. 1A).

This morphology, termed "foraging growth", was further investigated using time-lapse imaging over

75 days and revealed a morphological transition around day 50 (Video S1).

This transition appears linked to nutrient availability, as it occurred earlier on smaller media volumes

and later on undiluted TSA. Foraging growth was evident already after 21 days when grown in ⌀ 4 cm

dishes with 0.5xTSA. Using these dishes, we monitored nutrient depletion by inoculating plates with

spores and sampled the agar every three days for 27 days and analyzed nutrient content using mass

spectrometry. Samples were taken 0.5 cm away from the colony, to ensure that the samples would

not be contaminated with hyphae, and the very edge of the dish. Altogether, 51 metabolites were

annotated, including sugars, amino acids, and fatty acids, most of which depleted over time (Fig. 1B).

Nutrient levels proximal to the colony mirrored those at the dish-edge (Figure 1 – figure supplement

1A). Although the levels of e.g., glucose and fructose were depleted at foraging onset, others e.g.,

levels of glutamine and lactic acid initially increased before decreasing. Some, e.g., leucine and

phenylalanine, remained stable until day 12, before decreasing (Fig. 1C, Figure 1 – figure supplement

1B). The levels of some, e.g., sucrose, stayed constant throughout, and only five metabolites increased

more than one-fold (Fig. 1C). However, supplementing these metabolites in 0.5xTSA and MS agar did

not induce foraging growth. These findings demonstrate that the morphological transition coincides

with media nutrient depletion.

Nutrient depletion promotes foraging growth.

Modified *Streptomyces*-minimal medium only contain two carbon sources: agar and 55 mM glucose

(Kieser et al., 2000). Observing that foraging growth commenced upon carbon source depletion, we

excluded glucose, making agar the sole carbon source in the modified medium, hereafter referred to

as foraging agar media (FAM). Colonies on FAM displayed similar characteristics to those on 0.5xTSA

post-transition, exhibiting a thin, veil-like submerged expansion eventually covering the entire plate

(Fig. 2A). Over time, lineages developed within the colony, some of which became pigmented (Fig.

2B). The amount of aerial hyphae and pigment production at the inoculation site was influenced by

the presence of metabolizable nutrients, such as glycerol, in the spore storage buffer (Fig. 2C),

underscoring the effect of nutrients on the growth characteristics of the foraging phenotype. FAM

supplemented with glucose or fructose effectively inhibited foraging growth, leading to conventional

raised colonies (Fig. 2D). Notably, sucrose supplementation of FAM did not prevent foraging growth,

indicating the inability of *S. coelicolor* to metabolize sucrose.

Time-lapse imaging captured the progressive expansion and sector formation of a foraging colony on

FAM (Fig. 2E, Supplementary Video S2). Introduction of glucose or tryptone to a foraging colony

induced raised colony growth and strong pigment production, typical for growth in nutrient-rich

conditions (Fig. 2F, 2G, Supplementary Video S3 and S4). Scanning Electron Microscopy (SEM) imaging

of FAM revealed predominantly submerged hyphal growth, with limited hyphae at the surface (Fig.

2H). Aerial hyphae were found both at the site of inoculation and uncommonly at other patches

throughout the colony. In contrast, growth on nutrient-rich media such as 0.5x TSA revealed that

vegetative hyphae were primarily piled up on the surface, with only a few extending further into the

agar.

Sector formation was observed during foraging growth (Fig. 1, 2A, 2B, Video S1 and S2). To investigate

these sectors for potential genomic instability as reported for *S. coelicolor* grown on nutrient rich

media (Z. Zhang et al., 2020), a lineage from the initial inoculation, a subsequent sector 'A', and its

subsequent sectors 'I' and 'IV', were sequenced using the Illumnia platform. Analysis using breseq

software (Deatherage & Barrick, 2014) showed no significant genomic instability compared to growth

on nutrient-rich media. Only three mutations were identified: two mutations in sector A, in the

chromosomal replication initiator protein DnaA (SCO3879, E520K) and in a hypothetical protein

(SCO1511, Q8CK26). Sector IV had an additional mutation in a putative regulator (SCO5582, Q9ZBP7)

(Figure 2 – figure supplement 1). These mutations did not visibly affect colony morphology on rich

media. Altogether, these results demonstrate that *S. coelicolor* behaves distinctly during nutrient

deprived growth.

***S. coelicolor* foraging growth secretome is both Sec and Tat dependent.**

S. coelicolor grown on a deprived media such as FAM must utilize the agar as the main carbon source.

Agar degradation was visualized by staining the plates with iodine-based Lugol's solution, which

revealed a distinct halo around the colonies indicating substrate degradation (Fig. 3A). Proteins were

extracted from the zones outside of foraging colonies and analyzed using mass spectrometry. The

resulting protein composition was visualized using proteomaps (Liebermeister et al., 2014). Of the 203

detected proteins, 52 were annotated into six categories: membrane transport, biosynthesis, central

carbon metabolism, signal transduction, other enzymes, and

folding/sorting/degradation/transcription, as shown in the treemap (Fig. 3B) and the proteins

represented in the gene name mosaic map (Fig. 3C).

Signal peptide prediction on all *S. coelicolor* coding sequences using SignalP 6.0 (Teufel et al., 2022)

identified 696 proteins with signal sequences. Prediction of the 203 proteins in the foraging secretome

showed: 56 with the Sec translocon, 18 with the Tat translocon and 123 without any signal peptide

(Fig. 3D, Supplementary table S1). Proteomaps of these three groups indicated functional overlaps

among proteins with Sec, Tat and without signal peptides (Fig. 3E). Comparing our findings with

previous studies of *S. coelicolor* secretomes on rich media (Kim, Chater, Lee, & Hesketh, 2005; Widdick

et al., 2006) showed that 24% of the proteins identified in foraging conditions were also detected in

nutrient-rich conditions, while 75% were unique to foraging, including the glucose-repressible

extracellular agarose DagA (Figure 3 – figure supplement 1A, Supplementary table S2). Comparing the

proteins lacking signal peptides from foraging (n=203) and rich conditions (n=401) revealed an overlap

of 60 proteins (Supplementary table S4).

PANTHER classification of signal peptide-containing proteins categorized them into four groups (Fig.

3F, Supplementary table S4). Notably, the transporter category included the ABC transporter BldKB,

known for its role in carbon catabolite repression and mycelium formation (Chavez et al., 2011;

Nodwell & Losick, 1998). The metabolite interconversion category featured 13 hydrolases and one

oxidoreductase, while the protein modifying enzyme category included ten proteases and a ubiquitin-

protein ligase. Among the unclassified proteins were additional hydrolases and solute binding

proteins. The non-signal peptide proteins were classified into 11 categories (Figure 3 – figure

supplement 1B, Supplementary table S5).

Four proteins in the foraging secretome, encoded by genes containing the *bldA*-specific TTA codon,

suggest that *bldA* is active during foraging growth: an uncharacterized protein without a signal peptide

(Q9L2F9, SCO2524), a carbohydrate-binding module (CBM) domain-containing hydrolase (Q9RKF0,

SCO3487) with a Sec-signal peptide, and two uncharacterized secreted proteins with Sec-signal

peptides (Q9RKE0, SCO3498 and Q9K452, SCO7233) (Supplementary table S6). In total, 24 secretome

proteins identified in the Carbohydrate-Active enZYmes (CAZy) Database (Drula et al., 2022) are

associated with hydrolytic and non-hydrolytic cleavage of glycosidic bond and carbohydrate adhesion

(FIG 3G).

In summary, these results uncover the secretome components crucial for foraging growth of *S.*

coelicolor on depleted media, highlighting the role of both Sec and Tat transport systems in nutrient

degradation and uptake of agar components.

Foraging phenotype is found across the phylogeny of *Streptomyces*.

To investigate the influence of classical developmental regulators (*bld* and *whi*) on foraging growth,

we cultured wild-types and developmental-mutants on FAM and found that all tested mutants

displayed the foraging phenotype (Figure 4 – figure supplement 1A). Notably, the *bldA* mutant

exhibited foraging growth despite the presence of TTA-codon containing gene products in the foraging

secretome.

We then extended the study to various *Streptomyces spp.*, representing different branches of the

phylogenetic tree (Table 1), to assess if foraging growth was unique to *S. coelicolor*. Pairwise average

nucleotide identity (ANI) values were utilized to create a heatmap that groups strains according to

their genomic similarity, showing that only *S. coelicolor* and *S. lividans* shared an ANI >99 (Fig. 4A). All

tested strains displayed the foraging growth phenotype on FAM (Fig. 4B), with little to no

pigmentation and agar degradation confirmed by Lugol's solution staining (Figure 4 – figure

supplement 1B). Similar to *S. coelicolor*, the foraging growth was inhibited in all strains with the

addition of 1% glucose or tryptone to FAM (Figure 4 – figure supplement 1C and 1D). Additionally,

from local soil samples grown on FAM, we selected seven colonies exhibiting the foraging morphology.

These were all confirmed to be *Streptomyces* spp. using MALDI-TOF analysis (Supplementary table S7).

This observation suggests the ubiquity of foraging growth across the Streptomycetes phylogenetic

spectrum. Furthermore, the growth behavior of developmental mutants on FAM indicates that these

traditional developmental regulators do not control foraging growth.

Table 1. Phylogenetic clades of *streptomycetes* according to Labeda *et al.*, 2012.

Species	Clade
S. antibioticus	4
S. filipiniensis	10
S. griseus	*
S. anulatus	*
S. griseolus	36
S. venezuelae	40
S. kanamyceticus	*
S. noursei	67
S. hygroscopicus	75
S. hiroshimensis	84
S. coelicolor	112
S. lividans	**
K. setae	***

211 * Unsupported and single member clusters

212 ** Not included in Labeda *et al.*, 2012

213 *** Clusters outside of Streptomyces according to Labeda *et al.*, 2017

Foraging *S. coelicolor* displays an altered metabolomic profile.

S. coelicolor colonies grown on FAM were found to inhibit the growth of a contaminating mold (Fig.

5A), identified as *Penicillium citrinum* through 18S/ITS sequencing. This inhibition was replicated by

growing *S. coelicolor* on FAM for five days before inoculating *P. citrinum* spores at a distance of 2 cm,

resulting in an average 4.3 mm inhibition zone (±0.76 mm, n = 96). Inhibition was not observed when

glucose was added to the agar or on any other nutrient-rich media, nor on FAM with pH below 7.

Interestingly, the *bldA* mutant, is typically unable to produce certain antimicrobial metabolites,

inhibited *P. citrinum* growth on FAM. However, the *adpA*-mutant strain M851 showed less inhibition

compared to the wild-type (Fig. 5B). All tested *whi*-mutants (-A, -B, -G, -H, and -I) exhibited mold-

inhibitory activity (Figure 5 – figure supplement 1A). Inhibition of a local *Aspergillus spp*. isolate was

also demonstrated using an overlay assay with a 100x diluted TSA-overlay, since the *Aspergillus spp.*

did not grow on FAM (Figure 5 – figure supplement 1B). However, no inhibition of *Saccharomyces*

cerevisiae or *Candida albicans* was detected using the same assay.

To explore the metabolomic profile of *S. coelicolor* grown on FAM and FAM supplemented with 1%

glucose (FAM+G), agar was sampled around the colonies for mass spectrometric analysis. The

metabolomic profiles differed between FAM and FAM+G, both in negative (R2(X)=0.75, Q2(X)=0.97)

and positive detection modes (R2(X)=0.63, Q2(X)=0.47) (Fig. 5C, 5E). Some features with relatively

higher concentrations on FAM-agar (highlighted in green in Fig. 5D, 5F) could be matched with known

S. coelicolor metabolites based on monoisotopic molecular weight using the StreptomeDB

(Moumbock et al., 2021). Among these annotated features, the signaling molecule γ-butyrolactone

was found at higher concentrations in FAM+G. Wailupemycin G was the only annotated metabolite

found at considerably higher concentrations on FAM+G (Fig. 5G, Table 2). There were 30 features,

with higher levels in the FAM condition, that could not be annotated using their monoisotopic mass

and StreptomeDB (Supplementary table S8). Twelve of these features were unique to FAM and not

detected in any other tested condition (Figure 5 – figure supplement 1C and 1D).

The mass spectrometry results demonstrate a significant alteration in the metabolomic profile of

foraging *S. coelicolor*. The mold inhibitory assay using mutants suggests that mold inhibition involves

adpA and other genes dependent on *bldA*.

Table 2 Metabolomic features annotated using monoisotopic mass and StreptomeDB.

Annotation	Detection mode/RT	ppm error
7-Acetyl-3.6-dihydroxy-8-methyl-tetralone	+/3.44	3.4
7-Acetyl-3.6-dihydroxy-8-methyl-tetralone	-/3.03	0.9
Acetylglucosamine	+/0.39	0.3
Aloesaponarin II	+/3.16	0.4
Dehydromutacin	+/3.26	1.9
Fluostatin C	+/3.06	1.2
Fluostatin C	+/3.38	1.2
Fluostatin C	+/7.27	7.4
Fluostatin F	+/3.93	0.1
Fluostatin K	+/3.16	1.5
γ-butyrolactone	-/1.42	2.6
Gentisic acid	-/3.76	2.5
Homogentisic acid / Orsellic acid	-/2.02	1.5
Litmomycin	+/2.85	1.5
Mycothiol	+/1.14	12.5
Tryptophan	-/2.26	0.6
Wailupemycin G	+/0.4	11.2

247   *RT = Retention time

249   **Foraging *Streptomycetes* demonstrate increased antibacterial activity.**

250   To investigate whether mold-inhibition was a common behavior during foraging, all *Streptomyces spp.*

251   tested for foraging growth above, were tested for mold inhibition. However, none of the other species

inhibited *P. citrinum*. The altered metabolome of *S. coelicolor* during foraging, combined with its ability

to hinder mold growth, led us to investigate its potential to inhibit bacterial growth. Due to the

nutrient-depleted state of FAM, *B. subtilis* and *E. coli* were unable to proliferate. Therefore, to test for

antibacterial activity, we overlaid FAM with foraging *S. coelicolor* colonies with diluted LA and then

added *B. subtilis* 168 and *E. coli* MG1655 on top of the LA layers. The growth was evaluated after

incubation using a stereo microscope. This method was applied to all previously studied *Streptomyces*

spp. to assess their antibacterial properties. Most species showed increased inhibitory activity on

nutrient-depleted FAM-agar compared to FAM supplemented with glucose (FAM+G) and tryptone

(FAM+T) (Fig. 6A). However, *S. coelicolor* did not exhibit enhanced antibacterial activity on FAM

compared to the nutrient-supplemented media.

Analysis of bacterial inhibition by developmental mutants indicated that the *bldA*-mutant lost its

ability to inhibit *B. subtilis* growth, whereas the *adpA* mutant acquired the ability to inhibit *E. coli* (Fig.

6B). In summary, these findings suggest that streptomycetes are more competitive on FAM than on

rich agar. The distinct responses of the *bldA* and *adpA* mutants further emphasize their roles in the

antimicrobial activities of foraging *S. coelicolor*.

Methods

Strains, media, and culture conditions

The strains used in this study are listed in Supplementary Table S9. Streptomyces strains were grown

on FAM (1 g $(NH_4)_2SO_4$, 0.5 g $K_2PO_4x7H_2O$, 0.1 g $FeSO_4x7H_2O$, 10 g agar, in 1L pH 7.5) (Kieser et al.,

2000). FAM were supplemented with 1% of various sugars: glucose (FAM+G), sucrose (FAM+S),

fructose (FAM+F) or tryptone as an amino acid source (FAM+T) when indicated. To investigate the

effect of pH on mold-inhibition the pH of FAM was set using HEPES-buffer and HCl/NaOH. Tryptone

soy broth (Thermo Fisher Scientific) was prepared according to instructions and diluted 0.5x when

indicated with water and 1.5% agar added before autoclaving. MS-agar (20 g mannitol, 20 g Soy flour,

15 g agar in 1L) agar were used for sporulation and petri dish time-lapse imaging. Effects of

accumulated metabolites were conducted by adding 20mg/ml: beta-alanine, meso-erythriol (Merck),

Potassium-citramalate monohydrate (Merck), urea (Merck), and pantothenic acid (Merck). All

incubations were carried out at 30°C.

Isolation and identification of local isolates

The inhibited mold was isolated from the FAM dish and recultivated on TSA. The agar plate was sent

to Eurofins for 18S/ITS sequencing. For isolation of local streptomycetes, soil samples from potatoes

and plant soil were dried at room temperature before being deposited on FAM. The plates were

incubated at 37°C until foraging colonies were observed after approx. 5–10 days. Colonies with

characteristic foraging phenotypes were re-streaked on TSA and characterized using MALDI-TOF

(Bruker). The local *Aspergillus spp.* was isolated from a soil sample and assessed morphologically.

Colony imaging

For all imaging of small colonies spores were inoculated on agar media and visualized using a Nikon

SMZ1500 dissection microscope and a Nikon Digital sight DS-Fi1 camera, with either surface

illumination using LEDs or oblique sub-stage illumination. Whole petri dishes were imagined using a

Nikon D7000 DSLR. Agar time-lapses were recorded using a Raspberry Pi 3 model B with camera

module v2 mounted in a 10x lens (Olympus). The device was placed in a 30°C dark room with an LED

lamp programmed to illuminate the plate only during image capture, one frame per hour. ImageJ

distribution Fiji (Schindelin et al., 2012) was used to process single frames into videos and Adobe

Photoshop CC software for still images.

Electron microscopy

Colonies of *S. coelicolor* on agar were cut, fixed overnight at 4°C in 2.5% glutaraldehyde/0.1M sodium

cacodylate buffer, dehydrated in series of ethanol, critical point dried and coated with a 5 nm Au/Pd

before being analyzed with a Merlin Field Emission SEM (Zeiss) using SE-HE2 and in-lens detectors.

Images were processed using Adobe Photoshop CC software.

Inhibition assays

For mold inhibition, *Streptomyces* spores were inoculated on FAM and incubated at 30°C for 5 days.

Spores from *P. citrinum* were inoculated 2 cm from the foraging colony. The plates were incubated for

14 days and observed daily for inhibition zones and imaged as described above. For bacterial

inhibition, plates with *Streptomyces spp.* were incubated for 20 days before a thin layer of 100x diluted

TSA (1.5% agar) was placed on top of agar and incubated overnight at 4°C. Overnight cultures of *E. coli*

and *B. subtilis* were diluted and $2x10^7$ CFU in 100 µl was added to the agar overlays and incubated

overnight at 37°C. Inhibition zones were assessed and imaged using a Nikon SMZ1500 dissection

microscope and a Nikon Digital sight DS-Fi1 camera.

Foraging secretome protein analysis

The agar outside of foraging *S. coelicolor* on FAM incubated for 20 days was extracted. The agar was

crushed and incubated overnight with water in a 50 ml Falcon tube. Proteins in the supernatant were

precipitated by addition of TCA and incubated on ice of 4 minutes before centrifugation at 20,000g for

5 min. The pellet was washed twice with 200 µl ice-cold acetone. The pellet was dried, dissolved in

protein sample buffer and loaded on an SDS-PAGE. Protein dense regions were cut and sent for mass

spectroscopy identification to Linköping University, where general in-gel digestion was performed as

described by Shevchenko *et al.* (Shevchenko, Tomas, Havlis, Olsen, & Mann, 2006).

Metabolomic analysis

Metabolic profiling of media nutrients by GC-MS was performed at the Swedish Metabolomics Center

in Umeå, Sweden. For the nutrient analysis of agar, six replicates of agar plugs were collected from

the edge of the dish and 0.5 cm away from the *S. coelicolor* colonies on 0.5xTSA medium. This sampling

was performed on six different ⌀ 4 cm plates. Three control samples were taken at every time point

from three sterile 0.5xTSA dishes. Sample preparation was performed according to A *et al.* (A et al.,

2005). The samples were stored at -80 °C until analysis. Small aliquots of the remaining supernatants

were pooled and used to create quality control (QC) samples. The samples were analyzed in batches

according to a randomized run order on GC-MS. Derivatization and GCMS analysis were performed as

described previously A *et al*. (A et al., 2005). Non-processed MS-files from the metabolic analysis were

exported from the ChromaTOF software in NetCDF format to MATLAB-R2020a (Mathworks, Natick,

MA, USA), where all data pre-treatment procedures (base-line correction, chromatogram alignment,

data compression and Multivariate Curve Resolution) were performed. The extracted mass spectra

were identified by comparisons of their retention index and mass spectra with libraries of retention

time indices and mass spectra (Schauer et al., 2005). Mass spectra and retention index comparison

was performed using NIST MS 2.2 software. Annotation of mass spectra was based on reverse and

forward searches in the library. Masses and ratio between masses indicative of a derivatized

metabolite were especially notified. The mass spectrum with the highest probability indicative of a

metabolite and the retention index between the sample and library for the suggested metabolite was

± 5 the deconvoluted "peak" was annotated as an identification of a metabolite. The annotated

nutrients were plotted using GraphPad Prism (version 9.0.2) to show trends over time and ridge plots

were generated using R version 4.2 and packages ggplot2 3.4.4 and ggridges 0.5.4.

For untargeted metabolite analysis, agar stubs were punched out from just outside of *S. coelicolor*

colonies on FAM and FAM+G. For each condition, twelve samples were isolated from plates inoculated

with *S. coelicolor*, in addition to two control samples each from sterile plates. The weight of the

samples were adjusted to 100 mg before the addition of 1 mL extraction buffer (90/10 v/v

methanol:water). Internal standards (13C9-Phenylalanine, 13C3-Caffeine was, D4-Cholic acid, D8-

Arachidonic Acid, 13C9-Caffeic Acid) were added to each sample. The sample was shaken at 30 Hz for

3 min in a mixer mill, and proteins were precipitated at +4 °C for 2h on ice. The sample was centrifuged

at +4 °C, 14 000 rpm, for 10 min. The supernatant was transferred to a microvial and solvents

evaporated. Before analysis, the sample was re-suspended in 10 + 10 µL methanol and water. The set

of samples were analyzed in positive mode and negative mode. The chromatographic separation was

performed on an Agilent 1290 Infinity UHPLC-system (Agilent Technologies, Waldbronn, Germany).

Re-suspended aliquots (2 µL) of the agar extracts were injected onto an Acquity UPLC HSS T3, 2.1 x 50

mm, 1.8 µm C18 column in combination with a 2.1 mm x 5 mm, 1.8 µm VanGuard precolumn (Waters

Corporation, Milford, MA, USA) held at 40°C. The gradient elution buffers were A (H2O, 0.1 % formic

acid) and B (75/25 acetonitrile:2-propanol, 0.1 % formic acid). The compounds were detected with an

Agilent 6550 Q-TOF mass spectrometer equipped with a jet stream electrospray ion source. The

settings were kept identical between the modes, except the capillary voltage. The data processing was

performed using the Recursive Feature Extraction algorithm within Agilent Masshunter Profinder

version B.08.00 (Agilent Technologies Inc., Santa Clara, CA, USA). All multivariate statistical

investigations (PCA) were performed using the software package SIMCA®-P+ version 15.0.2 (Umetrics,

Umeå, Sweden). Annotation of unknown metabolites were performed by comparing monoisotopic

molecular weights with *S. coelicolor* metabolites in the StreptomeDB (Moumbock et al., 2021) using R

software (4.2). Metabolomic features were plotted using GraphPad Prism (version 9.0.2). All raw mass

spectrometry files used in this study have been deposited to ScieLifeLab Data Repository with the

dataset identifier DOI: https://doi.org/10.17044/scilifelab.c.7171122.

Average nucleotide identity of streptomycetes

FastANI (Jain, Rodriguez, Phillippy, Konstantinidis, & Aluru, 2018) version 1.33 was employed to

calculate the Average Nucleotide Identity (ANI) values, assessing the genomic similarity among the 13

species investigated in this study (GenBank assembly IDs: GCA_000010605.1, GCA_000203835.1,

GCA_000269985.1, GCA_000721185.1, GCA_000739105.1, GCA_001514065.1, GCA_001886595.1,

GCA_008704495.1, GCA_014649495.1, GCA_014650335.1, GCA_016917755.1, GCA_019933235.1,

GCA_020783455.1), were analyzed using FastANI with its default settings. Pairwise comparisons were

conducted across all genomes, and the resulting ANI values were utilized to generate a heatmap using

R version 4.2.

376 **DNA sequencing of foraging sectors**

377 Two levels of sectors formed during foraging on FAM were grown in TSB-media and 600 mg of pellet

378 was sent to MicrobesNG for Illumnia sequencing. Quality controlled and trimmed reads with 30x

379 coverage were obtained and analyzed using breseq software (Deatherage & Barrick, 2014) with

380 default settings, using the S. coelicolor A(3)2 reference genome (Genbank: GCA_000203835.1,

381 (Bentley et al., 2002)) updated using wild-type sequence data to accommodate for genomic variation

382 in lab strain. The reads of descendent sectors were mapped against this updated reference genome.

383 Illumnia read data used in this study have been deposited to SciLifeLab Data Respository with the

384 dataset identifier DOI: https://doi.org/10.17044/scilifelab.c.7171122.

385 **Statistical analysis**

386 In this study, certain measurements in the untargeted metabolomics analysis fell below the detection

387 limits, resulting in missing values. To ensure an accurate representation of our findings and maintain

388 the integrity of the statistical analysis, these missing values were omitted from the dataset.

389 Consequently, all values presented in the violin plots depict observable data trends and variations

390 above the detection threshold. All statistical analyses were conducted using GraphPad Prism (version

391 9.0.2). Details on the specific type of statistical analysis and the criteria for significance testing are

392 provided in the legends of the respective figures.

393 # Discussion

394 The primary habitat of streptomycetes, the soil, is a dynamic environment with variable

395 environmental factors. Laboratory cultivation on a narrow range of rich media may not fully represent

396 adaptations occurring in the diverse, competitive soil environment. This study reports a novel

397 phenotype-transition of *S. coelicolor* colonies upon nutrient depletion.

398

Metabolomics analysis showed that the phenotype transition on 0.5xTSA coincided with nutrient

depletion, leading to the "foraging phenotype". Media composition is known to be crucial for life-cycle

progression, with nutrient depletion often referred to as the trigger of differentiation and sporulation,

though the process is acknowledged to be complex (Barka et al., 2016; Bush, Tschowri, Schlimpert,

Flardh, & Buttner, 2015; Flardh & Buttner, 2009; McCormick & Flardh, 2012). Instead of a sporulation,

S. coelicolor responded to nutrient depletion with a phenotype transition. Four distinct trends in

metabolite dynamics were distinguishable: depletion, accumulation, initial accumulation followed by

depletion, and unchanged levels. The trend of initial accumulation followed by depletion is likely due

to the breakdown of macromolecules in the TSA-media. It is established that fluctuations of

metabolite levels during growth are a result of the breakdown and uptake of external sources and

subsequent release into the surroundings (Douglas, 2020). Steady accumulation may occur from by-

product export or independently of metabolite overflow (Pinu et al., 2018). Supplementation of the

accumulated metabolites to media did not trigger foraging growth, suggesting they do not act as

signals for phenotype transition. However, two accumulated metabolites, β-alanine and pantothenic

acid, are crucial in various biochemical pathways. β-alanine, essential for pantothenic acid synthesis –

a precursor of Coenzyme A – enhances respiration in alkaline soils (Datta et al., 2017). Their release

might influence the microbial landscape in the natural habitat of *S. coelicolor*.

Nutrient levels proximal to the colony and at the edge of the dish closely followed each other,

explaining the persistence of the foraging phenotype even as it expands into unexplored areas.

Calculations based on agar diffusion coefficients suggest that nutrients like glucose can diffuse 1.5 cm^2

over three days (Miyamoto, Atsuyama, Ekino, & Shin, 2018).

In contrast to rich media, where low molecular weight organic compounds are abundant (Kieser et al.,

2000; Shirling & Gottlieb, 1966), soil environments typically have transient availability of such

nutrients (D. L. Jones & Murphy, 2007; Rousk & Jones, 2010). The amount of complex carbohydrates

and plant debris of fertile soils are considered high, but there is also a flux of available nutrients

throughout the year that the soil inhabitants need to accommodate for (Randewig, Marshall,

Näsholm, & Jämtgård, 2019). FAM, our minimal defined agar medium without glucose, supports spore

germination and foraging growth, but growth reverted to conventional phenotype upon the addition

of nutrients. This indicates that the complex polysaccharides in agar are not preferred over simpler

organic compounds. Nitrogen and phosphate also play a crucial role in cycle progression and

metabolite production. Our FAM media, with higher nitrogen levels than typically found in soil, may

influence these processes (Rousk & Jones, 2010). Nitrogen-concentrations higher than 1 mM allow

synthesis of glutamate independent of ATP, but low levels require the activity of glutamate synthetase

(Reuther & Wohlleben, 2007). Glutamate synthetase II activity its mRNA levels correlate with mycelial

differentiation on solid media demonstrating its importance in mycelial development (Fink,

Weissschuh, Reuther, Wohlleben, & Engels, 2002). High levels of nitrogen in rich media cause

repression of secondary metabolite production (Hodgson, 2000). Further, high levels of inorganic

phosphate have been shown to inhibit the production of several secondary metabolites (Bibb, 2005;

Hobbs et al., 1992; Martin, 2004). Accordingly, we detected an increase of some metabolites during

FAM-growth, but we also detected a decreased production of several metabolites. Microdiffusion

experiments in soils have revealed that there is a significant flux of organic nitrogen (Inselsbacher &

Nasholm, 2012; Inselsbacher, Öhlund, Jämtgård, Huss-Danell, & Näsholm, 2011; Leitner et al., 2017).

A nitrogen source in the form of organic ammonium, as e.g., asparagine prevents foraging growth on

FAM-plates. It will therefore be of future interest to evaluate the effect of altered nitrogen and

phosphate levels as well as organic versus inorganic sources, on foraging growth and metabolome.

The foraging secretome contained 74 proteins with predicted signal peptides. Comparing the foraging

secretome with rich growth secretome (Widdick et al., 2006) showed that 75% of foraging secretome

was not found during rich growth. This underscores the adaptation that takes place upon nutrient

depletion that results in the foraging phenotype. Despite the presence of *bldA*-dependent proteins in

the secretome, a *bldA*-mutant still displayed foraging growth, suggesting *bldA*-regulation is not

essential for this phenotype. The developmental protein BldKB was identified in the secretome,

suggesting a role in foraging. Additionally, the foraging behavior was observed in various *Streptomyces*

species, indicating it may be a common feature across the genus (Labeda et al., 2017; Labeda et al.,

2012). Multi locus sequence analysis suggests that *Kitasatospora* is a distinct genus, and suggests that

foraging is not limited to *Streptomycetes*, since foraging growth was also observed for *K. setae*.

The study demonstrates that foraging growth conditions affect secondary metabolism. The gained

ability of *S. coelicolor* to inhibit mold-growth together with the overall enhanced potential of other

Streptomyces spp. to inhibit gram-positive and gram-negative bacteria suggests that there are also

significant alterations of their metabolomes. This shows potential for using FAM growth in screening

for active secondary metabolites not produced on rich media. However, the low nutrient levels used

for the overlay assays could also influence the target bacteria rendering them more susceptible to the

antimicrobial effects of foraging colonies. It is well established that growth conditions, e.g., choice of

carbon source, affect secondary metabolism of streptomycetes (Sanchez et al., 2010; van Keulen &

Dyson, 2014) and other strategies are also being employed to activate silent BGC (Ochi & Hosaka,

2013; van der Meij, Worsley, Hutchings, & van Wezel, 2017). A recent discovery shows that secondary

metabolite production is significantly increased as a result from genetic instability on nutrient rich MS-

agar (Z. Zhang et al., 2020). We could however not find any support for major mutational deletions

during foraging growth when investigating the sectors formed during foraging growth. This further

emphasizes that the survival strategies of *S. coelicolor* differ between depleted and rich growth

conditions.

The mechanism by which foraging growth is regulated remains to be elucidated. Identifying the

regulatory components would result in understanding the required nutrient deprived conditions.

Importantly, knowledge about its regulation could also be utilized as a tool for manipulating the

production of the secondary metabolites. In summary, this study demonstrates that nutrient-deprived

growth leads to a distinct growth phenotype and altered metabolome, with enhanced antimicrobial

properties. This novel growth behavior and its metabolic implications offer new insights into

environmental adaptation and competitiveness of streptomycetes under nutrient deprivation with

potential implications in anti-microbial drug discovery.

Additional files

•   Figure 1 – figure supplement 1. Nutrient levels during phenotype transition on 0.5xTSA.

•   Figure 2 – figure supplement 1. Sector mutations of foraging colonies.

•   Figure 3 – figure supplement 1. Secretome of foraging colonies.

•   Figure 4 – figure supplement 1. Growth of *Streptomyces spp.* on FAM.

•   Figure 5 – figure supplement 1. Mold inhibition and unannotated metabolomic features.

•   Supplementary table 1. Prediction of signal peptides in the foraging secretome using SignalP

6.0.

•   Supplementary table 2. Comparison of the foraging secretome with proteins secreted on rich

media (Widdick et al 2006).

•   Supplementary table 3. Comparison of cytosolic protein detected during foraging growth and

growth on rich media by Widdick et al (2006).

•   Supplementary table 4. PANTHER classification of signal peptide containing proteins with

regard to protein class.

•   Supplementary table 5. PANTHER classification of cytosolic protein with regard to protein

class.

•   Supplementary table 6. TTA containing genes in *S. coelicolor* A3(2).

•   Supplementary table 7. Locally isolates streptomcyetes.

• Supplementary table 8. Monoisotopic mass of features significantly higher levels in FAM

conditions.

• Supplementary table 9. Strain list.

• Video 1. *S. coelicolor* phenotype transition.

• Video 2. *S. coelicolor* growing on FAM-agar.

• Video 3. Addition of glucose to foraging *S. coelicolor.*

• Video 4. Addition of tryptone to foraging *S. coelicolor.*

• Figure 1 - source data 1. Nutrient fold change values over time used for the ridge plots.

• Figure 1 – source data 2. Nutrient fold change values for edge and proximal samples used for

plotting single nutrients over time.

• Figure 1 – figure supplement 1B – source data. Nutrient fold change values for edge and

proximal samples used for plotting single nutrients over time.

• Figure 4A – source data. ANI values for each species used for generating the heatmap.

• Figure 5G – source data. Values used for plotting the annotated features detected using

untargeted mass spectrometry.

• Figure 5 – figure supplement 1D – source data. Values used for plotting the unannotated

features specific to foraging growth on FAM detected using untargeted mass spectrometry.

Data availability

520 The Illumina sequencing data and the metabolomics mass spectrometry data has been deposited in

521 the SciLifeLab Data Repository under DOI: https://doi.org/10.17044/scilifelab.c.7171122

523 The following datasets were generated

524 Sandblad, Linda; Söderholm, Niklas (2024). Foraging growth of Streptomyces - Datasets. SciLifeLab.

525 Collection. https://doi.org/10.17044/scilifelab.c.7171122

Acknowledgments

We thank National Microscopy Infrastructure, NMI (VR-RFI 2016-00968) at Umeå Centre for Electron

Microscopy (UCEM). Swedish Metabolomics Centre, Umeå, Sweden

(www.swedishmetabolomicscentre.se) is acknowledged for metabolic profiling by LC-MS and GC-MS.

Illumnia sequencing was provided by MicrobesNG (http://www.microbesng.com). Postdoc fellowship

funding from the Kempe foundations and Sven och Lilly Lawskis foundation. Project granted by Umeå

Centre for Microbial research (UCMR) and Swedish Foundation for Strategic Research (RIF21-0067).

Anders Sjöstedt for MALDI-TOF analysis of streptomycetes and Ala Javadi for sample preparation.

Figure legends

Figure 1. *S. coelicolor* growth phenotype transition correlates with nutrient depletion. (**A**) Colony

appearance after 75 days growth on 0.5xTSA. The point of phenotype transition is indicated by the

dotted circle. (**B**) Visualization comparing nutrient levels during *S. coelicolor* growth on 0.5xTSA. The

distribution within each ridge plot shows the variation in fold change across the time points. The

amplitude of each peak indicates how common the fold changes are for that nutrient. Each nutrient

is accompanied by a colored square indicating its status at the end of the time series. The

measurements proximal to the colony were taken in sextuplicate for each time point and normalized

against the controls. Source file of all fold change over time used for the ridge plots is available in the

Figure 1 – Source data 1. (**C**) Examples of metabolites levels throughout the study. The time point

where the transition to foraging growth was observed is indicated with the dotted lines on day 21.

Each measurement proximal to the colony and at the edge of the dish were taken in sextuplicate for

each time point and normalized against the controls. Source file of nutrient fold change over time in

edge and proximal samples used for the plots is available in the Figure 1 – Source data 2.

Figure 2. Nutrient poor agar media promote the *S. coelicor* foraging phenotype. (**A**) *S. coelicor* spores in glycerol storage buffer inoculated on FAM. The foraging phenotype covering the petri dish was photographed after 15 weeks of incubation at 30°C. Transparent pigment-free and lightly pigmented sectors are visible. (**B**) A photograph of a pigmented sector extending towards the edge of the dish. (**C**) Formation of aerial hyphae were only observed at the site of spore inoculation. (**C**) A photograph of aerial hyphae at the site of spore inoculation. (**D**) Micrograph showing the phenotypic growth on WT *S. coelicor* inoculated on FAM, FAM+sucrose (+S), FAM+glucose (+G) and FAM+fructose (+F) for 15 weeks at 30°C. (**E**) Time-lapse stills showing the foraging growth progression at day 12, 26, 40, 52. (**F**) Time-lapse stills after addition of glucose (+G) to a foraging colony at day 0, 4, 8, 12. (**G**) Time-lapse stills of addition of tryptone (+T) to a foraging colony. (**H**) SEM imaging of *S. coelicor* grown on FAM and 0.5xTSA for 7 days. The glycerol storage buffer was washed away and replaced with water. A top view of the center of the colonies and at the edge show few hyphae at the surface of the FAM. Hyphae extending through the agar are mainly observed on the FAM-plates, while they predominantly grow on top on 0.5xTSA.

Figure 3. Characterization of the foraging phenotype secretome. (**A**) Micrograph of foraging *S. coelicor* grown on FAM, stained with lugol solution. (**B**) Treemap displaying the functional categories of subproteomes from the foraging secretome. Out of 203 detected proteins, 52 were assigned to categories. The category area size represents the number of assigned proteins in each category, not protein abundance. (**C**) Mosaic showing the gene names within each category. Area size does not represent protein abundance. (**D**) Number of proteins containing a signal sequence in the foraging secretome, predicted using SignalP 6.0 (Teufel et al., 2022). Categories include general secretory (Sec) substrates cleaved by SPase I (Sec/SPI) or SPase II (Sec/SPII, lipoproteins) and twin-arginine substrates cleaved by SPase I (Tat/SPI) or SPase II (Tat/SPII, lipoprotein), and "No SP" for no predicted signal peptide. (**E**) Treemap showing the functional categories when separated into the groups: No SP (left, 22/129 proteins assigned), Sec (middle, 17/56 proteins assigned) and Tat-signal peptide (right, 7/18 proteins assigned). (**F**) PANTHER functional classification of proteins from the foraging secretome with

predicted signal peptides. The graph shows the number of proteins assigned to each protein class. (**G**)

CAZy classification of the foraging secretome: Glycoside Hydrolases (GH), Polysaccharide Lyases (PLs)

and Carbohydrate-Binding Modules (CBMs).

Figure 4. Phylogenetically separated streptomycetes display the foraging phenotype. (**A**) Heatmap

visualizing pairwise ANI values, calculated using fastANI (Jain et al., 2018), among the genomes of

tested species as a quantitative measure of their genomic similarity. Each cell in the heatmap

represents the ANI percentage between a pair of genomes. The branches of a dendrogram represent

how each strain is grouped based on hierarchical clustering with euclidean distance as the metric for

dissimilarity. Source file of ANI values used for the heatmap is available in the Figure 4 – Source data

1. (**B**) Microscopy images showing the foraging phenotype of each *Streptomyces spp.* cultivated for

three weeks on FAM. Abbreviations: SANT; *S. antibioticus*, SANU; *S. anulatus*, SCO; *S. coelicolor*, SFI;

S. filipinensis, SGR; *S. griseus*, SGRI; *S. griseolus*, SHIR; *S. hiroshimensis*, SHYG; *S. hygroscopicus*, SKAN;

S. kanamyceticus, SLIV; *S. lividans*, SNOR; *S. norsei*, SVEN; *S. venezuelae*, KSET; *Kitasatospora setae*.

Figure 5. Mold inhibitory activity and metabolomic analysis of foraging *S. coelicolor*. (**A**) Inhibition

zones around *S. coelicolor* on FAM contaminated by *P. citrinum* mold. (**B**) Inhibition zones around *S.*

coelicolor wild-type (WT), bldA and M851 (*adpA*-mutant) inoculated with *P. citrinum*. (**C**) Scores plot

of LC-MS analysis in negative detection mode of FAM and FAM+G-agar inoculated with *S. coelicolor*.

(**D**) Loadings plot from negative detection mode. Features with relatively elevated levels on FAM are

highlighted in blue and features with relatively elevated levels in FAM+G-agar are highlighted in red.

(**E**) Scores plot of LC-MS analysis in positive detection mode of FAM and FAM+G-agar inoculated with

S. coelicolor. (**F**) Loadings plot from positive detection mode. Features with relatively elevated levels

on FAM-agar are highlighted in blue and features with relatively elevated levels in FAM+G-agar are

highlighted in red. (**G**) Graphs showing the features annotated by monoisotopic mass and

StreptomeDB. All graphs depict statistically significant outcomes comparing FAM and FAM+G as

determined by t-tests, with p-values <=0.0002. 7-Acetyl-3.6-dihydroxy-8-methyl-tetralone is

abbreviated as 7A36D8MT in the plot. Source file with plotted values is available in the Figure 5 – Source data 1.

Figure 6. Antibacterial activity of foraging streptomycetes. (**A**) Bactericidal activity of foraging streptomycetes determined by overlay-assays with indicator bacteria (Ec=*E. coli*, Bs=*Bacillus subtilis*) on FAM, FAM+G and FAM+T. (**B**) Bactericidal activity of *S. coelicolor* developmental mutants as determined by overlay assays on FAM. Dark purple indicates inhibition of indicator bacteria, light purple indicates partial growth inhibition, and grey indicates that there was no detectable effect.

Figure supplement legends

Figure 1 - figure supplement 1. Nutrient levels during phenotype transition on 0.5xTSA. (**A**) The fold change for every nutrient at every time point is plotted for both sampled positions on the dish. The plot reveals a highly positive correlation, as indicated by the R-value of >0.96 suggesting a significant linear relationship. (**B**) The fold change for each annotated metabolite during phenotype transition on 0.5xTSA is overlayed for both positions on the dish. The time point where the transition was observed is indicated with the dotted lines at day 21. Source file of nutrient fold change over time in edge and proximal samples used for the plots is available in the Figure 1 – Source data 2.

Figure 2 - figure supplement 1. Sector mutations of foraging colonies. Spores of *S. coelicolor* were inoculated on FAM-agar and the sectors were isolated and subjected to Illumnia sequencing. The inoculum was compared to the first level sector formed (α) and two succeeding sectors (I and IV) originating from lineage α. Using the breseq software, two point mutations were detected in sector α: DnaA (SCO3879:E520K) and uncharacterized DUF349-containing protein (SCO1511:+G). One mutation was found in SCO5585 $(CG)_{5\rightarrow4}$ in linage IV.

Figure 3 - figure supplement 1. Secretome of foraging colonies. (**A**) Venn diagram of secretome from nutrient deprived and rich conditions highlighting the shared and distinct protein profiles of each. (**B**) PANTHER functional classification of proteins from the foraging secretomes that were predicted to not have signal peptides. The plot shows the number of proteins assigned to each protein class.

Figure 4 - figure supplement 1. Growth of Streptomyces on FAM. (**A**) Growth of *S. coelicolor* wild-types (M145 and M600) and developmental mutants: *bldA*, *whiA* (J2401), *whiB* (J2402), *whiG* (J2400), *whiH* (J2408), *whiI* (J2450), and M851 (Δ*adpA*) on FAM. (**B**) Streptomycetes on FAM stained with lugol solution after 5 days of incubation. Abbrevations: SANT; *S. antibioticus*, SANU; *S. anulatus*, SCO; *S. coelicolor*, SFI; *S. filipinensis*, SGR; *S. griseus*, SGRI; *S. griseolus*, SHIR; *S. hiroshimensis*, SHYG; *S. hygroscopicus*, SKAN; *S. kanamyceticus*, SLIV; *S. lividans*, SNOR; *S. norsei*, SVEN; *S. venezuelae*, KSET; *Kitasatospora setae*. (**C**) Micrographs of the tested streptomycetes grown on FAM+G for 14 days. (**D**) Micrographs of the tested streptomycetes grown on FAM+T for 14 days.

Figure 5 – figure supplement 1. Mold inhibition and significant unannotated metabolomic features. (**A**) Inhibition of *P. citrinum by S. coelicolor* developmental mutants: *whiA* (J2401), *whiB* (J2402), *whiG* (J2400), *whiH* (J2408), *whiI* (J2450) and wild-type M600. (**B**) Inhibition of local *Aspergillus*-isolate in an overlay-assay on foraging S. coelicolor. Pictures were taken of foraging colonies using a dissection microscope. (**C**) Plots of the unannotated metabolomic features unique to growth on FAM. The top right corner of each graph indicates the molecular weight of corresponding metabolite. All graphs depict statistically significant outcomes as determined by t-tests, with p-values <0.0001. Source file with plotted values is available in the Figure 5 – figure supplement 1 - Source data 1.

References

A, J., Trygg, J., Gullberg, J., Johansson, A. I., Jonsson, P., Antti, H., . . . Moritz, T. (2005). Extraction and GC/MS analysis of the human blood plasma metabolome. *Anal Chem, 77*(24), 8086-8094. doi:10.1021/ac051211v

Barka, E. A., Vatsa, P., Sanchez, L., Gaveau-Vaillant, N., Jacquard, C., Meier-Kolthoff, J. P., . . . van Wezel, G. P. (2016). Taxonomy, Physiology, and Natural Products of Actinobacteria. *Microbiol Mol Biol Rev, 80*(1), 1-43. doi:10.1128/MMBR.00019-15

Bentley, S. D., Chater, K. F., Cerdeno-Tarraga, A. M., Challis, G. L., Thomson, N. R., James, K. D., . . . Hopwood, D. A. (2002). Complete genome sequence of the model actinomycete Streptomyces coelicolor A3(2). *Nature, 417*(6885), 141-147. doi:10.1038/417141a

Bibb, M. J. (2005). Regulation of secondary metabolism in streptomycetes. *Curr Opin Microbiol, 8*(2), 208-215. doi:10.1016/j.mib.2005.02.016

Bush, M. J., Tschowri, N., Schlimpert, S., Flardh, K., & Buttner, M. J. (2015). c-di-GMP signalling and
the regulation of developmental transitions in streptomycetes. *Nat Rev Microbiol, 13*(12),
749-760. doi:10.1038/nrmicro3546
Chater, K. F., Biro, S., Lee, K. J., Palmer, T., & Schrempf, H. (2010). The complex extracellular biology
of Streptomyces. *FEMS Microbiol Rev, 34*(2), 171-198. doi:10.1111/j.1574-
6976.2009.00206.x
Chavez, A., Forero, A., Sanchez, M., Rodriguez-Sanoja, R., Mendoza-Hernandez, G., Servin-Gonzalez,
L., . . . Sanchez, S. (2011). Interaction of SCO2127 with BldKB and its possible connection to
carbon catabolite regulation of morphological differentiation in Streptomyces coelicolor.
Appl Microbiol Biotechnol, 89(3), 799-806. doi:10.1007/s00253-010-2905-8
Datta, R., Baraniya, D., Wang, Y.-F., Kelkar, A., Meena, R. S., Yadav, G. S., . . . Formanek, P. (2017).
Amino Acid: Its Dual Role as Nutrient and Scavenger of Free Radicals in Soil. *Sustainability,*
9(8), 1402. Retrieved from https://www.mdpi.com/2071-1050/9/8/1402
Deatherage, D. E., & Barrick, J. E. (2014). Identification of mutations in laboratory-evolved microbes
from next-generation sequencing data using breseq. *Methods Mol Biol, 1151*, 165-188.
doi:10.1007/978-1-4939-0554-6_12
Douglas, A. E. (2020). The microbial exometabolome: ecological resource and architect of microbial
communities. *Philos Trans R Soc Lond B Biol Sci, 375*(1798), 20190250.
doi:10.1098/rstb.2019.0250
Drula, E., Garron, M. L., Dogan, S., Lombard, V., Henrissat, B., & Terrapon, N. (2022). The
carbohydrate-active enzyme database: functions and literature. *Nucleic Acids Res, 50*(D1),
D571-D577. doi:10.1093/nar/gkab1045
Fink, D., Weissschuh, N., Reuther, J., Wohlleben, W., & Engels, A. (2002). Two transcriptional
regulators GlnR and GlnRII are involved in regulation of nitrogen metabolism in
Streptomyces coelicolor A3(2). *Mol Microbiol, 46*(2), 331-347. doi:10.1046/j.1365-
2958.2002.03150.x
Flardh, K., & Buttner, M. J. (2009). Streptomyces morphogenetics: dissecting differentiation in a
filamentous bacterium. *Nat Rev Microbiol, 7*(1), 36-49. doi:10.1038/nrmicro1968
Hobbs, G., Obanye, A. I., Petty, J., Mason, J. C., Barratt, E., Gardner, D. C., . . . Oliver, S. G. (1992). An
integrated approach to studying regulation of production of the antibiotic methylenomycin
by Streptomyces coelicolor A3(2). *J Bacteriol, 174*(5), 1487-1494. doi:10.1128/jb.174.5.1487-
1494.1992
Hodgson, D. A. (2000). Primary metabolism and its control in streptomycetes: A most unusual group
of bacteria. In *Advances in Microbial Physiology* (Vol. 42, pp. 47-238): Academic Press.
Ikeda, H., Ishikawa, J., Hanamoto, A., Shinose, M., Kikuchi, H., Shiba, T., . . . Omura, S. (2003).
Complete genome sequence and comparative analysis of the industrial microorganism
Streptomyces avermitilis. *Nat Biotechnol, 21*(5), 526-531. doi:10.1038/nbt820
Inselsbacher, E., & Nasholm, T. (2012). The below-ground perspective of forest plants: soil provides
mainly organic nitrogen for plants and mycorrhizal fungi. *The New phytologist, 195*(2), 329-
334. doi:10.1111/j.1469-8137.2012.04169.x
Inselsbacher, E., Öhlund, J., Jämtgård, S., Huss-Danell, K., & Näsholm, T. (2011). The potential of
microdialysis to monitor organic and inorganic nitrogen compounds in soil. *Soil biology &*
biochemistry, 43(6), 1321-1332. doi:10.1016/j.soilbio.2011.03.003
Jain, C., Rodriguez, R. L., Phillippy, A. M., Konstantinidis, K. T., & Aluru, S. (2018). High throughput
ANI analysis of 90K prokaryotic genomes reveals clear species boundaries. *Nat Commun,*
9(1), 5114. doi:10.1038/s41467-018-07641-9
Jones, D. L., & Murphy, D. V. (2007). Microbial response time to sugar and amino acid additions to
soil. *Soil biology & biochemistry, 39*(8), 2178-2182. doi:10.1016/j.soilbio.2007.03.017
Jones, S. E., Ho, L., Rees, C. A., Hill, J. E., Nodwell, J. R., & Elliot, M. A. (2017). Streptomyces
exploration is triggered by fungal interactions and volatile signals. *Elife, 6*.
doi:10.7554/eLife.21738

Kieser, T., Bibb, M. J., Buttner, M. J., Chater, K. F., Hopwood, D. A., & John Innes, F. (2000). *Practical*
Streptomyces genetics. Norwich: John Innes Foundation Norwich.
Kim, D. W., Chater, K. F., Lee, K. J., & Hesketh, A. (2005). Effects of growth phase and the
developmentally significant bldA-specified tRNA on the membrane-associated proteome of
Streptomyces coelicolor. *Microbiology (Reading), 151*(Pt 8), 2707-2720.
doi:10.1099/mic.0.28000-0
Labeda, D. P., Dunlap, C. A., Rong, X., Huang, Y., Doroghazi, J. R., Ju, K. S., & Metcalf, W. W. (2017).
Phylogenetic relationships in the family Streptomycetaceae using multi-locus sequence
analysis. *Antonie Van Leeuwenhoek, 110*(4), 563-583. doi:10.1007/s10482-016-0824-0
Labeda, D. P., Goodfellow, M., Brown, R., Ward, A. C., Lanoot, B., Vanncanneyt, M., . . . Hatano, K.
(2012). Phylogenetic study of the species within the family Streptomycetaceae. *Antonie Van*
Leeuwenhoek, 101(1), 73-104. doi:10.1007/s10482-011-9656-0
Leitner, S., Minixhofer, P., Inselsbacher, E., Keiblinger, K. M., Zimmermann, M., & Zechmeister-
Boltenstern, S. (2017). Short-term soil mineral and organic nitrogen fluxes during moderate
and severe drying–rewetting events. *Applied soil ecology : a section of Agriculture,*
ecosystems & environment, 114, 28-33. doi:10.1016/j.apsoil.2017.02.014
Liebermeister, W., Noor, E., Flamholz, A., Davidi, D., Bernhardt, J., & Milo, R. (2014). Visual account
of protein investment in cellular functions. *Proc Natl Acad Sci U S A, 111*(23), 8488-8493.
doi:10.1073/pnas.1314810111
Martin, J. F. (2004). Phosphate control of the biosynthesis of antibiotics and other secondary
metabolites is mediated by the PhoR-PhoP system: an unfinished story. *J Bacteriol, 186*(16),
5197-5201. doi:10.1128/JB.186.16.5197-5201.2004
McCormick, J. R., & Flardh, K. (2012). Signals and regulators that govern Streptomyces development.
FEMS Microbiol Rev, 36(1), 206-231. doi:10.1111/j.1574-6976.2011.00317.x
Miyamoto, S., Atsuyama, K., Ekino, K., & Shin, T. (2018). Estimating the Diffusion Coefficients of
Sugars Using Diffusion Experiments in Agar-Gel and Computer Simulations. *Chem Pharm Bull*
(Tokyo), 66(6), 632-636. doi:10.1248/cpb.c18-00071
Moumbock, A. F. A., Gao, M., Qaseem, A., Li, J., Kirchner, P. A., Ndingkokhar, B., . . . Gunther, S.
(2021). StreptomeDB 3.0: an updated compendium of streptomycetes natural products.
Nucleic Acids Res, 49(D1), D600-D604. doi:10.1093/nar/gkaa868
Nett, M., Ikeda, H., & Moore, B. S. (2009). Genomic basis for natural product biosynthetic diversity in
the actinomycetes. *Nat Prod Rep, 26*(11), 1362-1384. doi:10.1039/b817069j
Nikolaidis, M., Hesketh, A., Frangou, N., Mossialos, D., Van de Peer, Y., Oliver, S. G., & Amoutzias, G.
D. (2023). A panoramic view of the genomic landscape of the genus Streptomyces. *Microb*
Genom, 9(6). doi:10.1099/mgen.0.001028
Nodwell, J. R., & Losick, R. (1998). Purification of an extracellular signaling molecule involved in
production of aerial mycelium by Streptomyces coelicolor. *J Bacteriol, 180*(5), 1334-1337.
doi:10.1128/JB.180.5.1334-1337.1998
Ochi, K., & Hosaka, T. (2013). New strategies for drug discovery: activation of silent or weakly
expressed microbial gene clusters. *Appl Microbiol Biotechnol, 97*(1), 87-98.
doi:10.1007/s00253-012-4551-9
Ohnishi, Y., Ishikawa, J., Hara, H., Suzuki, H., Ikenoya, M., Ikeda, H., . . . Horinouchi, S. (2008).
Genome sequence of the streptomycin-producing microorganism Streptomyces griseus IFO
13350. *J Bacteriol, 190*(11), 4050-4060. doi:10.1128/JB.00204-08
Pinu, F. R., Granucci, N., Daniell, J., Han, T. L., Carneiro, S., Rocha, I., . . . Villas-Boas, S. G. (2018).
Metabolite secretion in microorganisms: the theory of metabolic overflow put to the test.
Metabolomics, 14(4), 43. doi:10.1007/s11306-018-1339-7
Qin, Z., Munnoch, J. T., Devine, R., Holmes, N. A., Seipke, R. F., Wilkinson, K. A., . . . Hutchings, M. I.
(2017). Formicamycins, antibacterial polyketides produced by Streptomyces formicae
isolated from African Tetraponera plant-ants. *Chem Sci, 8*(4), 3218-3227.
doi:10.1039/c6sc04265a

Randewig, D., Marshall, J. D., Näsholm, T., & Jämtgård, S. (2019). Combining microdialysis with
metabolomics to characterize the in situ composition of dissolved organic compounds in
boreal forest soil. *Soil Biology and Biochemistry, 136*, 107530.
doi:https://doi.org/10.1016/j.soilbio.2019.107530
Reuther, J., & Wohlleben, W. (2007). Nitrogen metabolism in Streptomyces coelicolor:
transcriptional and post-translational regulation. *J Mol Microbiol Biotechnol, 12*(1-2), 139-
146. doi:10.1159/000096469
Romero-Rodriguez, A., Rocha, D., Ruiz-Villafan, B., Guzman-Trampe, S., Maldonado-Carmona, N.,
Vazquez-Hernandez, M., . . . Sanchez, S. (2017). Carbon catabolite regulation in
Streptomyces: new insights and lessons learned. *World J Microbiol Biotechnol, 33*(9), 162.
doi:10.1007/s11274-017-2328-0
Rousk, J., & Jones, D. L. (2010). Loss of low molecular weight dissolved organic carbon (DOC) and
nitrogen (DON) in H2O and 0.5 M K2SO4 soil extracts. *Soil biology & biochemistry, 42*(12),
2331-2335. doi:10.1016/j.soilbio.2010.08.017
Sanchez, S., Chavez, A., Forero, A., Garcia-Huante, Y., Romero, A., Sanchez, M., . . . Ruiz, B. (2010).
Carbon source regulation of antibiotic production. *J Antibiot (Tokyo), 63*(8), 442-459.
doi:10.1038/ja.2010.78
Schauer, N., Steinhauser, D., Strelkov, S., Schomburg, D., Allison, G., Moritz, T., . . . Kopka, J. (2005).
GC-MS libraries for the rapid identification of metabolites in complex biological samples.
FEBS Lett, 579(6), 1332-1337. doi:10.1016/j.febslet.2005.01.029
Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T., . . . Cardona, A.
(2012). Fiji: an open-source platform for biological-image analysis. *Nat Methods, 9*(7), 676-
682. doi:10.1038/nmeth.2019
Shepherdson, E. M. F., & Elliot, M. A. (2022). Cryptic specialized metabolites drive Streptomyces
exploration and provide a competitive advantage during growth with other microbes. *Proc
Natl Acad Sci U S A, 119*(40), e2211052119. doi:10.1073/pnas.2211052119
Shevchenko, A., Tomas, H., Havlis, J., Olsen, J. V., & Mann, M. (2006). In-gel digestion for mass
spectrometric characterization of proteins and proteomes. *Nat Protoc, 1*(6), 2856-2860.
doi:10.1038/nprot.2006.468
Shirling, E. B., & Gottlieb, D. (1966). Methods for characterization of Streptomyces species.
International journal of systematic bacteriology, 16(3), 313-340. doi:10.1099/00207713-16-
3-313
Teufel, F., Almagro Armenteros, J. J., Johansen, A. R., Gislason, M. H., Pihl, S. I., Tsirigos, K. D., . . .
Nielsen, H. (2022). SignalP 6.0 predicts all five types of signal peptides using protein
language models. *Nat Biotechnol, 40*(7), 1023-1025. doi:10.1038/s41587-021-01156-3
van der Meij, A., Worsley, S. F., Hutchings, M. I., & van Wezel, G. P. (2017). Chemical ecology of
antibiotic production by actinomycetes. *FEMS Microbiol Rev, 41*(3), 392-416.
doi:10.1093/femsre/fux005
van Keulen, G., & Dyson, P. J. (2014). Production of Specialized Metabolites by Streptomyces
coelicolor A3(2). *Advances in Applied Microbiology, 89*, 217-266. doi:10.1016/B978-0-12-
800259-9.00006-8
Widdick, D. A., Dilks, K., Chandra, G., Bottrill, A., Naldrett, M., Pohlschroder, M., & Palmer, T. (2006).
The twin-arginine translocation pathway is a major route of protein export in Streptomyces
coelicolor. *Proc Natl Acad Sci U S A, 103*(47), 17927-17932. doi:10.1073/pnas.0607025103
Zhang, X., Hindra, & Elliot, M. A. (2019). Unlocking the trove of metabolic treasures: activating silent
biosynthetic gene clusters in bacteria and fungi. *Curr Opin Microbiol, 51*, 9-15.
doi:10.1016/j.mib.2019.03.003
Zhang, Z., Du, C., de Barsy, F., Liem, M., Liakopoulos, A., van Wezel, G. P., . . . Rozen, D. E. (2020).
Antibiotic production in Streptomyces is organized by a division of labor through terminal
genomic differentiation. *Sci Adv, 6*(3), eaay5781. doi:10.1126/sciadv.aay5781

Figure 1. *S. coelicolor* growth phenotype transition correlates with nutrient depletion.

Figure 2. Nutrient poor agar media promote the *S. coelicolor* foraging phenotype.

Figure 3. Characterization of the foraging phenotype secretome.

Figure 4. Phylogenetically separated streptomycetes display the foraging phenotype.

Figure 5. Mold inhibitory activity and metabolomic analysis of foraging *S. coelicolor*.

Figure 6. Antibacterial activity of foraging streptomycetes.

Figure 1 - figure supplement 1

Organic acids

Carbohydrates and carbohydrate conjugates

Fatty acids

Others

Figure 2 - figure supplement 1

Figure 3 - figure supplement 1

Figure 4 - figure supplement 1

Figure 5 - figure supplement 1