Title: Habitat fragmentation mediates the mechanisms underlying long-term

climate-driven thermophilization in birds

Authors: Juan Liu[1], Morgan W. Tingley[2], Qiang Wu[1], Peng Ren[1], Tinghao Jin[1], Ping

Ding[1*], Xingfeng Si[3*]

Emails:

Juan Liu (juanliu@zju.edu.cn); Morgan W. Tingley (mtingley@ucla.edu);

Qiang Wu (qiangwu.eco@gmail.com); Peng Ren (renpeng@zju.edu.cn);

Tinghao Jin (tinghaojin2020@gmail.com); Ping Ding (dingping@zju.edu.cn);

Xingfeng Si (sixf@des.ecnu.edu.cn)

Institutional affiliations:

[1]MOE Key Laboratory of Biosystems Homeostasis & Protection, College of Life Sciences, Zhejiang

University, Hangzhou, Zhejiang 310058, China;

[2]Department of Ecology and Evolutionary Biology, University of California, Los Angeles, Los

Angeles, California, USA;

[3]Zhejiang Zhoushan Island Ecosystem Observation and Research Station, Institute of Eco-

Chongming, Zhejiang Tiantong Forest Ecosystem National Observation and Research Station,

School of Ecological and Environmental Sciences, East China Normal University, Shanghai 200241,

China.

Corresponding authors:

Ping Ding: Tel: 86-571-88206468; Email: dingping@zju.edu.cn

Xingfeng Si: Tel: 86-21-54341138; Email: sixf@des.ecnu.edu.cn

Abstract

Climatic warming can shift community composition driven by the colonization-extinction

dynamics of species with different thermal preferences; but simultaneously, habitat

fragmentation can mediate species' responses to warming. As this potential interactive effect

has proven difficult to test empirically, we collected data on birds over 10 years of climate

warming in a reservoir subtropical island system that was formed 65 years ago. We

investigated how the mechanisms underlying climate-driven directional change in

community composition were mediated by habitat fragmentation. We found

thermophilization driven by increasing warm-adapted species and decreasing cold-adapted

species in terms of trends in colonization rate, extinction rate, occupancy rate and population

size. Critically, colonization rates of warm-adapted species increased faster temporally on

smaller or less isolated islands; cold-adapted species generally were lost more quickly

temporally on closer islands. This provides support for dispersal limitation and microclimate

buffering as primary proxies by which habitat fragmentation mediates species range shift.

Overall, this study advances our understanding of biodiversity responses to interacting

global change drivers.

1 Introduction

One widespread consequence of anthropogenic-caused climate warming is the shift in species' distributions toward cooler regions (Chen et al., 2011; La Sorte & Thompson, 2007; Thomas & Lennon, 1999). As species move across the globe, the composition of community changes, with species both colonizing and departing (Menéndez et al., 2006; Tingley & Beissinger, 2013); yet colonizing and departing species are unlikely to be randomly drawn from species pools, but instead will be drawn depending on their traits, such as being warm-adapted versus cold-adapted. As a result, a warming climate should lead to compositional shifts toward relatively more warm-adapted species in local communities (Thomas et al., 2006), a phenomenon termed "thermophilization" (Feeley et al., 2020; Gottfried et al., 2012; Zellweger et al., 2020).

In practice, thermophilization can be measured by a temporal increase in the Community Temperature Index (CTI) (Devictor et al., 2008) – an estimation of the average temperature experienced by all species in an assemblage throughout their respective ranges. While thermophilization of communities is usually accompanied by an increase in warm-adapted species and a decrease in cold-adapted species (Curley et al., 2022; Prince & Zuckerberg, 2015; Tayleur et al., 2016) (**Figure 1a**), this process is usually spatially heterogeneous and depends on the availability of current habitat and the ability to disperse into newly suitable habitats (Gaüzère et al., 2017; Richard et al., 2021), both of which will be directly or indirectly affected by other global change stressors. Habitat fragmentation, usually defined as the shifts of continuous habitat into spatially isolated and small patches (Fahrig, 2003), in particular, has been hypothesized to have interactive effects with climate change on community dynamics. Specifically, habitat fragmentation usually alters both habitat spatial structure and microclimate characteristics (Ewers & Banks-Leite, 2013;

Fahrig, 2003), either of which can influence thermophilization processes (Gaüzère et al.,

2017; Platts et al., 2019; Richard et al., 2021; Zellweger et al., 2020). Understanding the

interaction between climate change and habitat fragmentation on community composition is

essential to predicting biodiversity outcomes but we generally lack empirical examples with

which to study this phenomenon (Opdam & Wascher, 2004).

Theoretically, habitat fragmentation could mediate differential dynamics of warm-

adapted versus cold-adapted species either by changing existing habitats' suitability or by

limiting dispersal into newly suitable habitats (Newson et al., 2014; Opdam & Wascher,

2004) (**Figure 1**). The overall effect of fragmentation on climatic niche tracking of

communities can be decomposed into three different mechanisms, each of which has been

proposed and supported with different degrees of evidence. First, fragmented habitats may

more weakly buffer macroclimate than intact habitats (Ewers & Banks-Leite, 2013),

resulting in fragmented patches experiencing relatively higher temperatures. Increased

temperature, especially in small patches, may boost the colonization or population growth

rate of warm-adapted species but cause higher extinction risk or demographic declines of

cold-adapted species (**Figure 1b**). This microclimate mechanism has been proposed as an

important factor driving thermophilization in plant communities (Zellweger et al., 2020).

Second, larger patches generally contain higher habitat heterogeneity and therefore also

encompass higher food availability and microrefugia (Liu et al., 2020). A diversity of

habitats and resources is closely related to a species' successful establishment and can act as

a buffer against extinction under climate change (Luoto & Heikkinen, 2008). For instance,

higher habitat diversity can boost the ability of niche tracking in bird communities (Gaüzère

et al., 2017), mainly by fostering species' dispersal into new, climatically suitable sites.

Similarly, in areas where topography generates greater microclimate variation, population

losses of plants and insects due to climate change are substantially reduced (Suggitt et al.,

2018). We thus expect that the reduction in habitat heterogeneity associated with higher

fragmentation will intensify the loss of cold-adapted species and weaken the colonization of

warm-adapted species (**Figure 1c**).

Third, habitat fragmentation could prevent or interrupt species' dispersal which is

essential for them to track suitable climatic niches (McGuire et al., 2016; Opdam & Wascher,

2004; Thomas et al., 2006). This limitation may come from higher mortality rate with

increasing dispersal distance during searching for suitable patches (Brooker et al., 1999),

and will influence the colonization-extinction dynamics underlying climate response

(Fourcade et al., 2021; Hylander & Ehrlén, 2013). For example, habitat fragmentation

renders habitats to be too isolated to be colonized, causing sedentary butterflies to lag more

behind climate warming in Britain than mobile ones (Warren et al., 2001). All things being

equal, less isolated patches should experience increasing rate in colonization of warm-

adapted species under warming, benefitting from near sources of species; meanwhile, cold-

adapted species should experience higher extinction rates on less isolated patches because

they can emigrate easily from these patches under climate warming (**Figure 1d**).

Although the overall effect of habitat fragmentation on climate-induced community

dynamics has been established, few empirical studies have ever considered these multiple

mechanisms simultaneously. Here, we use 10 years of bird community data in a subtropical

land-bridge island system (Thousand Island Lake, TIL, China, **Figure 2**) during a period of

consistent climatic warming (**Appendix 1—figure 1**) to explore whether and how

community thermal niche composition responds to climate change and how habitat

fragmentation mediates the process. This reservoir island system was formed 65 years ago

due to dam construction. Specifically, we focused on the following three predictions:

1. We predict the existence of thermophilization of bird communities in our fragmented

land-bridge island system, characterized by an increasing temporal trend of CTI.

2. We predict that thermophilization of bird communities should result from both increasing

warm-adapted species and abundances and decreasing cold-adapted species and

abundances. We also predict that, over time with a warming climate, warm-adapted

species should increase their colonization rate while cold-adapted species should

increase their extinction rate (**Figure 1a**).

3. We predict that habitat fragmentation – measured as island area and distance to the

mainland – will mediate the two main driving processes of thermophilization: colonization

of warm-adapted species and extinction of cold-adapted species. Following our theoretical

framework, if the climate buffering effect dominates (**Figure 1b**) area effects, we expect the

differential trends of warm-adapted species versus cold-adapted species to be more extreme

on smaller islands (i.e., even faster colonization of warm-adapted, even faster extinction of

cold-adapted) due to weaker microclimate buffering. Alternatively, if the effects of habitat

heterogeneity dominate (**Figure 1c**) area effects, cold-adapted species will again show

greater extinction, but warm-adapted species will alternatively experience lower

colonization on smaller islands (relative to larger islands) due to less abundant resources (Si

et al., 2017). Finally, we predict that on more isolated islands (**Figure 1d**), the colonization

of warm-adapted species and the extinction of cold-adapted species will increase more

slowly because isolation hinders both immigration and emigration.

2　Results

The number of species detected in surveys on each island across the study period averaged

13.37 ± 6.26 (mean $\pm$ SD) species, ranging from 2 to 40 species, with an observed gamma

diversity of 60 species. The STI of all 60 birds averaged 19.94 ± 3.58 □ (mean $\pm$ SD) and

ranged from 9.30 □ (*Cuculus canorus*) to 27.20 □ (*Prinia inornate*), with a median of

20.63 □ (**Appendix 1—figure 2**; **Appendix 1—figure 3**). STI of resident species (n = 47)

and summer visitors (n = 13) did not show a significant difference (t-test: $t = 0.23$, df =

17.82, $p = 0.82$). No significant correlation was found between STI and species' ecological

traits; specifically, the continuous variables of dispersal ability, body size, body mass and

clutch size (Pearson correlations for each, $|r| < 0.22$), and the categorial variables of diet

(carnivorous/omnivorous/herbivory), active layer (canopy/mid/low), and residence type

(resident species/summer visitor).

2.1   Thermophilization of bird communities

At the landscape scale, considering species detected across the study area, occurrence-based

CTI (CTI_{occur}; see section 4.4) showed no trend (posterior mean temporal trend = 0.414; 95%

CrI: -12.751, 13.554) but abundance-based CTI (CTI_{abun}; see section 4.4) showed a

significant increasing trend (temporal trend [mean ± SE] = 0.327 ± 0.041, t = 7.989, $p <$

0.0001). When measuring CTI trends for individual islands (expressed as °/ unit year), we

found significant increases in CTI for both occurrence- (mean temporal trend = 0.124; 95%

CrI: 0.108, 0.137) and abundance-based indices (temporal trend = 0.320 ± 0.021, $t_{493} =$

15.534, $p < 0.0001$; **Figure 3**).

2.2   Mechanisms underlying CTI trends

Comparing across species with different thermal affinities, we found a weak positive linear

relationship between STI and species-specific temporal occupancy trends (t-test: slope =

0.206, $t_{58} = 1.766$, $p = 0.082$; **Figure 4a**), indicating warm-adapted species were marginally

more likely to increase in occurrence over time. Decomposing occurrence change to

dynamic parameters; however, we found a significant positive relationship between STI and

species' temporal trends in colonization (t-test: slope = 0.259, $p = 0.043$) and a significant

negative relationship between STI and species' temporal trends in extinction (t-test: slope = -

0.414, $p = 0.015$; **Appendix 1—figure 4**). Thus, warm-adapted species generally increased

in occupancy over time, which was driven by an increase in colonization probability and a

decrease in extinction probability; in contrast, cold-adapted species generally decreased in

occupancy over time, which was driven by an increase in extinction probability and a

decrease in colonization probability (**Figure 4b).** Considering all species, the influence of a

thermal-association gradient underlying occurrence dynamics resulted in a particularly

strong correlation between species' trends in colonization and extinction probabilities

(Pearson correlation r = -0.77; **Figure 4b**).

Similar patterns were found for abundance, where we detected a strong positive

relationship (t-test: slope = 0.153, $t_{58} = 3.047$, $p = 0.003$) between STI and each species'

temporal trend in abundance: warm-adapted species generally increased abundance while

cold-adapted species decreased over 10 years (**Figure 4c**). At the island scale, both total

abundance and total species richness increased significantly for the warm-adapted species

group while both responses significantly decreased for the cold-adapted species (richness

trend, $p < 0.0001$; total abundance trend, $p < 0.0001$; **Appendix 1—figure 5**).

2.3   The mediating effects of fragmentation on colonization and extinction processes

Both colonization and extinction probabilities showed evidence of being moderated by

fragmentation (**Figure 5**). Of particular interest was how the temporal trend in dynamic

parameters varied as a function of area or isolation (i.e., Year:Area or Year:Isolation effects).

Although species-specific interaction terms were not generally significant for most species,

overall trends could be assessed across species. In particular, the effect of isolation on

temporal dynamics of thermophilization was relatively consistent across cold- (**Figure 5a**)

and warm-adapted species (**Figure 5b**); specifically, on islands nearer to the mainland,

warm-adapted species (15 out of 15 investigated species) increased their colonization

probability at a higher rate over time, while most cold-adapted species (21 out of 23 species)

increased their extinction probability at a higher rate.

Island area, in contrast, had a more mixed role in moderating colonization/extinction

rates across species. For most warm-adapted species (11 out of 15 investigated species;

Figure 5b), colonization rates increased faster over time on smaller islands (i.e., negative

area:year interaction term); however, the colonization rates of the other four species

increased faster on larger islands, including the only significant positive interactive effect

(*Dendrocitta formosae*). For cold-adapted species, extinction rates were approximately

equally split between increasing faster on smaller islands (12 out of 23 species) and

increasing faster on larger islands (11 out of 23 species), including two species, *Spizixos*

semitorques and *Pericrocotus cantonensis*, which showed significantly increasing rates of

extinction on larger islands.

3 Discussion

Our study explored the driving processes underlying a decade of thermophilization of a bird

community breeding in a model fragmented system, specifically testing whether and how

habitat fragmentation may mediate the colonization-extinction dynamics which underly

shifts in mean community traits. Our results confirm that community thermal niche

composition was changing directionally, mainly through gains in warm-adapted species and

individuals and losses in cold-adapted species and individuals. We further found that habitat

fragmentation influences two processes of thermophilization: colonization rates of most

warm-adapted species tended to increase faster on smaller and less isolated islands, while

the loss rates of most cold-adapted species tended to be exacerbated on less isolated islands.

3.1 Thermophilization of bird communities

Although thermophilization has been found in many taxa (Lajeunesse & Fourcade, 2022) and is more apparent in the tropics than in temperate zones (Freeman et al., 2021), there are few empirical demonstrations of this process in subtropical fragmented systems. We found decadal-scale thermophilization evident in both occurrence- and abundance-based metrics, although the response was considerably stronger for abundance. This result is consistent with work in other systems, where abundance-weighted CTI generally responds more quickly than occurrence-based CTI given climatic warming (Devictor et al., 2008; Lindström et al., 2013; Oliver et al., 2017; Tayleur et al., 2016). Notably, when tested on the landscape scale (versus on individual island communities), only the abundance-based thermophilization trend was significant, indicating thermophilization of bird communities was mostly due to occurrence dynamics within the region, rather than exogenous community turnover outside the region.

3.2 Mechanisms underlying CTI trends

Consistent with our expectation, we found an overall increase in occupancy and abundance of warm-adapted species and a decrease in occupancy and abundance of cold-adapted species, which together contributed to thermophilization. This pattern is consistent with many previous studies conducted in continuous habitats in Europe or North America both for breeding and non-breeding bird communities (Curley et al., 2022; Prince & Zuckerberg, 2015; Tayleur et al., 2016). Also consistent with our expectation was that occupancy change was the product of temporal trends in both colonization and extinction dynamics; namely, warm-adapted species showed increasing colonization and decreasing extinction rates, while, cold-adapted species showed decreasing colonization and increasing extinction rates. Following that STI is predictably related to temporal trends in colonization and extinction

rate (**Figure 4**), we can conclude that STI is an effective broad indicator of relative climatic

vulnerability within a community, where species with relatively higher thermal preferences

are generally poised to benefit from local climate warming and vice versa. This conclusion is

supported by studies that have established relationships between STI and species' population

trend (Pearce-Higgins et al., 2015; Rigal et al., 2023), species' contribution analysis (Tayleur

et al., 2016), or spatial occupancy (Anderson et al., 2023). Decomposing average community

trait shifts into colonization and extinction dynamics of individual species and their

associated traits provides a comprehensive and more nuanced insight into community

response to climate change.

There are numerous reasons why we would expect temperature to strongly drive

community change, as temperature is related to species' survival through food resources,

foraging time, and thermoregulatory requirements (Dunn & Møller, 2019). The local

population growth of warm-adapted species may mirror their range expansion under climate

warming. For example, thermophilization of avian communities in North America is mainly

owed to the species that have increased in both range and population size (Curley et al.,

2022). Contemporary latitudinal range shifts are thought to be primarily resultant of the

expansion of polar margins of equatorial-distributed species (La Sorte & Thompson, 2007;

Thomas & Lennon, 1999), possibly due to extinction lags at equatorial margins(La Sorte &

Jetz, 2012). TIL is near the cold edge of most warm-adapted species like *Hemixos*

castanonotus, *Pericrocotus solaris*, and *Lonchura punctulate*. Combined with the prediction

that most birds in China should move upslope or poleward under climate change (Hu et al.,

2020), we infer that local climate will continue to shift these warm-adapted species toward a

thermal optimum, predicting continued population growth in the near future for these

species (Sagarin & Gaines, 2002).

Meanwhile, most cold-adapted species in our system demonstrated increasing

extinction rates over time. Extinctions are more difficult to detect than range expansions,

partly due to temporal lags (Thomas et al., 2006; Thomas & Lennon, 1999) but also due to

pervasive bias of false absences in time series data (Tingley & Beissinger, 2009). Moreover,

species' extinctions are more likely to be mediated by factors such as rainfall, species

interactions, and changes in vegetation (Paquette & Hargreaves, 2021; Thomas & Lennon,

1999), which will further lead to unequal rates of extinction relative to colonization. In our

community, although there are many cold-adapted species – such as *Streptopelia orientalis*,

Phasianus colchicus, *Passer montanus*, and *Emberiza cioides* – that have northern

distributional limits much farther northern than our site, our site is not close to the current

southern boundary for these species either. Even so, they still showed considerable loss in

TIL, suggesting fragmented landscape exacerbated the negative effect of warming on these

cold-adapted species.

Despite the general trend, not all warm-adapted species increased (in abundance or

occupancy) and not all cold-adapted species decreased. In other words, not all species

contributed equally to the process of thermophilization, as has been seen elsewhere (Tayleur

et al., 2016). Variation in species' response relates to their true thermal tolerance in the study

region, which would require additional physiological data to measure and could also be

influenced by habitat usage and biotic interactions.

3.3   The mediating effect of fragmentation on colonization and extinction processes

It has long been thought that habitat fragmentation can impact species' climate tracking

(Opdam & Wascher, 2004) but there are only a few empirical studies to date (Fourcade et al.,

2021; Warren et al., 2001). Our research examining colonization-extinction dynamics

provides empirical evidence of the possible mechanisms in a land-bridge island system.

As expected, the mediating effect of isolation on the two main processes of

thermophilization was relatively consistent, with warm-adapted species colonizing faster and

cold-adapted species being extirpated faster on islands nearer to the mainland. This indicates

that isolation from population sources can limit the dispersal that underlies range shifts,

supporting the general perception that habitat fragmentation is thought to impede climate-

driven range expansions (Opdam & Wascher, 2004) or even block them (Warren et al., 2001).

While we cannot truly distinguish in our system between local extinction and emigration, we

suspect that given two islands equal except in isolation, if both lose suitability due to climate

change, individuals can easily emigrate from the island nearer to the mainland, while

individuals on the more isolated island would be more likely to be trapped in place until the

species went locally extinct due to a lack of rescue.

As a caveat, we only consider the distance to the nearest mainland as a measure of

fragmentation, consistent with previous work in this system (Si et al., 2014), but we

acknowledge that other distance-based metrics of isolation that incorporate inter-island

connections could reveal additional insights on fragmentation effects. The spatial

arrangement of islands, like the arrangement of habitat, can influence niche tracking of

species (Fourcade et al., 2021). Future studies should take these metrics into account to

thoroughly understand the influence of isolation and spatial arrangement of patches in

mediating the effect of climate warming on species.

Island size also had pervasive effects on underlying thermophilization dynamics. In our

study, the colonization rate of a large proportion of warm-adapted species (11 out of 15) and

half of cold-adapted species (12 out of 23) were increasing more rapidly on smaller islands.

Similarly, a lower proportion of habitats enhances the colonization of warm-adapted species

under climate change in the butterfly assemblage (Fourcade et al., 2021). The observed

colonization-extinction patterns on smaller islands support the hypothesis that warm-adapted

species benefit from the lowered thermal buffering of ambient temperature on smaller

islands, while the same mechanism may speed up extinction (or emigration) of cold-adapted

species (**Figure 1b**). To investigate this mechanistic hypothesis more thoroughly, using

understory air temperature data monitored on 20 islands in the breeding season in 2022, we

found that temperature buffering ability indeed was diminished on smaller islands

(**Appendix 1—figure 6**), further supporting microclimate as a primary driver of

fragmentation's mediation of thermophilization (Zellweger et al., 2020).

The increased extinction rate of some cold-adapted species on smaller islands is also

consistent with the hypothesized impacts of reduced habitat heterogeneity (**Figure 1c**).

Previous studies found that areas with greater amounts of habitat featured lower extinction

rates of cold-adapted butterfly species (Fourcade et al., 2021), possibly by facilitating

movement to colder microclimates (Suggitt et al., 2018). We thus suppose that habitat

heterogeneity could also mitigate the loss of these relatively cold-adapted species as

expected. Habitat diversity, including the observed number of species, the rarefied species

richness per island, species density and the rarefied species richness per unit area, all

increased significantly with island area instead of isolation in our system (Liu et al., 2020).

Thus, as habitat heterogeneity was only consistent with our observed response of cold-

adapted species (**Figure 1c**), while microclimate buffering is consistent with observed

responses of both warm- and cold-adapted species (**Figure 1b**), our evidence more

consistently supports the latter than the former.

Contrary to any expected mechanisms (**Figure 1**), the extinction rate of some cold-

adapted species – such as *Spizixos semitorques* and *Pericrocotus cantonensis* – was

increased faster on larger islands. Although post hoc, we speculate that these species face

stronger predation pressures on larger islands. For example, predators like snakes and felids

may also increase activity under climate warming (DeGregorio et al., 2015), but these

predators only exist in our system on the larger islands. Besides, *Garrulax canorus*, which is

a warm-adapted species, increased faster on larger islands. This species may depend more on

habitat heterogeneity than microclimate alone, including food resources and microrefugia,

which is related to faster climate tracking (Gaüzère et al., 2017). Overall, these idiosyncratic

responses reveal several possible mechanisms in regulating species' climate responses,

including resource demands and biological interactions like competition and predation.

Future studies are needed to take these factors into account to understand the complex

mechanisms by which habitat loss meditates species range shifts.

Overall, our findings have important implications for conservation practices. Firstly, we

confirmed the role of isolation in limiting range shifting. Better connected landscapes should

be developed to remove dispersal constraints and facilitate species' relocation to the best

suitable microclimate. Second, small patches can foster the establishment of newly adapted

warm-adapted species while large patches can act as refugia for cold-adapted species.

Therefore, preserving patches of diverse sizes can act as stepping stones or shelters in a

warming climate depending on the thermal affinity of species. These insights are important

supplement to the previous emphasis on the role of habitat diversity in fostering (Richard et

al., 2021) or reducing (Gaüzère et al., 2017) community-level climate debt.

4   Materials and Methods

4.1   Study area and islands selection

The Thousand Island Lake (TIL), located in eastern China, was formed in 1959 following

the construction of the Xin'anjiang Dam for hydroelectricity (**Figure 2**). When the lake is at

its highest, there are 1078 islands with an area larger than 0.25 ha. Currently, about 90% of

the forested areas are dominated by Masson pine (*Pinus massoniana*) in the canopy and

broad□leaved species (e.g. *Loropetalum chinensei*, *Vaccinium carlesii* and *Rhododendron*

simsii) in the sub☐canopy and understory (Liu et al., 2020). The climatic zone is a typical

monsoon climate. The precipitation is mainly concentrated between April and June with an

average yearly rainfall of 1430 mm. The average annual temperature is 17°C (hottest from

June to August) and the average daily temperature ranges from –7.6 °C in January to 41.8 °C

in July (Si et al., 2017).

We selected 36 islands according to a gradient of island area and isolation with a

guarantee of no significant correlation between island area and isolation (Pearson r = -0.21,

$p = 0.21$). For each island, we calculated island area and isolation (measured in the nearest

Euclidean distance to the mainland) to represent the degree of habitat fragmentation (**Figure**

2). Distance to the mainland is the best distance-based measure in fitting species'

colonization rate and extinction rate in TIL (Si et al., 2014). Since lake formation, the islands

have been protected by forbidding logging, allowing natural succession pathways to occur.

4.2   Bird data

We established 53 survey transects on 36 islands with the sampling effort on each island

roughly proportional to the logarithm of the island area (Schoereder et al., 2004). As a result,

53 transects on 36 islands were sampled, including eight transect trails on the largest study

island (area = 1058 ha), four transects on two islands between 100 and 500 ha, two transects

on four islands between 10 and 100 ha, and one on each of the remaining small islands (c. 1

ha for most islands) (Si et al., 2018). Breeding bird communities were surveyed on each

transect nine times annually (three times per month from April to June) from 2012 to 2021

(Si et al., 2017). In each survey, observers walked along each transect at a constant speed

(2.0 km/h) and recorded all the birds seen or heard on the survey islands. To minimize the

bias, the order and direction of each island surveyed were randomized. We based our

abundance estimate on the maximum number of individuals recorded across the nine annual

surveys. We excluded non-breeding species, nocturnal and crepuscular species, high-flying

species passing over the islands (e.g., raptors, swallows) and strongly water-associated birds

(e.g., cormorants) from our record. First, our surveys were conducted during the day, so

some nocturnal and crepuscular species, such as the owls and nightjars were excluded for

inadequate survey design. Second, wagtail, kingfisher, and water birds such as ducks and

herons were excluded because we were only interested in forest birds. Third, birds like

swallows, and eagles who were usually flying or soaring in the air rather than staying on

islands, were also excluded as it was difficult to determine their definite belonging islands.

Following these operations, 60 species were finally retained.

4.3   Climate data

We obtained climate data (monthly average temperature from 2012 to 2021) from the

Meteorological Bureau of Chun'an County in Zhejiang Province, China. We calculated

breeding season temperature as the average of the mean monthly temperature from April to

June in order to represent the thermal conditions birds experience at this site each year

(Lindström et al., 2013). Breeding season temperature increased significantly over 10 years

(slope = 0.078, r^2 = 34.53%, p = 0.04, **Appendix 1—figure 1**).

4.4   STI and CTI

We followed the methods of Devictor et al. (2008) to calculate a species temperature index

(STI) using ArcMap 10.3. STI of a given species is defined as the average temperature of

breeding months (April – June) across its distributional range (restricted to the Northern

hemisphere) averaged over 1970–2000. Monthly temperature data were obtained from

WorldClim at a resolution of 30-arc seconds (http://www.worldclim.org). Distributional

maps were extracted from Birdlife International 2019

(http://datazone.birdlife.org/species/requestdis), in which we selected only the distributional

regions where the species is either resident or breeding. All species were divided into two

groups indicating their relative thermal preference: species with STI higher than the median

STI were labeled as warm-adapted species, and species lower than the median STI were

labeled as cold-adapted species (A. E. Bates et al., 2017). The STI of a species can differ

depending on distributional and climate data but the rank order of species STI is highly

correlated across methods (Barnagaud et al., 2013). We verified the robustness of our STI

calculations using different distributional ranges and annual time windows (**Appendix 1—**

figure 2).

CTI is a community-level index representing the average species thermal niche of all

species or individuals in the community (Devictor et al., 2008). Accordingly, for each

community in each year, CTI was calculated as the average STI of all occurring species

(hereafter: $\mathrm{CTI_{occur}}$) or counted individuals (hereafter: $\mathrm{CTI_{abun}}$):

415
$$\mathrm{CTI}_{\mathrm{occur},j,t} = \frac{\sum_{i=1}^{N_{j,t}} STI_i}{N_{j,t}}$$

416
$$\mathrm{CTI}_{\mathrm{abun},j,t} = \frac{\sum_{i=1}^{N_{j,t}} (STI_i \times A_{i,j,t})}{\sum_{i=1}^{N_{t,j}} A_{i,j,t}}$$

where $N_{j,t}$ is the total number of species surveyed in the community j in year t, $A_{i,j,t}$ denoted

the maximum abundance among nine surveys of the i^{th} species in community j in year t.

$\mathrm{CTI_{occur}}$ was calculated using occurrence data corrected for imperfect detection (see

next section), while $\mathrm{CTI_{abun}}$ was estimated using the maximum annual count across 9

surveys. Given that our survey effort was so high (i.e., 9 repeat surveys per year), true

abundance should be highly correlated with maximum observed abundance (MacKenzie et

al., 2006). We also used these occurrence and abundance metrics from all islands to compute

each year's regional CTI at the landscape level.

4.5   Multispecies dynamic occupancy model

Presence-absence data can be highly sensitive to false negatives, so to explore the driving

processes of change in CTI_{occur}, and to explore how habitat fragmentation mediated the main

processes while accounting for the imperfect detection, we developed a spatially hierarchical

dynamic multi-species occupancy model (MSOM) in a Bayesian framework based on the

model by Royle and Kéry (2007). We denote y_{nijt} as the observation (detected = 1;

undetected = 0) for species n (1–60 species) in survey visit j (1–9 surveys) at transect i (1–53

transects) in year t (1–10 years). Observation, y_{nijt} is assumed to be the result of imperfect

detection of the true occurrence status, z_{nit} (1 or 0), and is thus modeled as a Bernoulli-

distributed variable with a probability of $p_{nijt} \times z_{nit}$, where p_{nijt} is the probability of detection

for a given survey along a transect:

$$y_{nijt} \sim Bernoulli\left(z_{nit} * p_{nijt}\right)$$

where z_{nit} is assumed to be of fixed presence/absence status for a given species across all j

survey intervals within year t. However, given that individual birds move dynamically across

territories within a year, during 9 surveys each year, we broadly interpret our estimates of

"occupancy" as "use" (Si et al., 2018) to relax the closure assumption (MacKenzie et al.,

2004).

Each site has an initial value of occupancy (ψ_{ni1}) which can be given a uniform prior

distribution from 0 to 1, and occupancy (ψ_{nit}) in subsequent years ($t > 1$) is determined based

on whether sites become colonized or remain occupied through persistence (MacKenzie et

al., 2003):

$$Z_{nit} \sim Bernoulli(\psi_{nit})$$

$$447 \qquad \psi_{nit} = Z_{ni,t-1}\big(1 - {}_{ni,t-1}\big) + (1 - Z_{ni,t-1})\gamma_{ni,t-1} \text{ for } t > 1$$

448 where ψ_{nit}, $\varepsilon_{ni,t-1}$, $\gamma_{ni,t-1}$ are transect-level probabilities of occupancy, extinction, and

449 colonization, respectively.

450   We modeled the probability of detection, p_{nijt}, as a function of the length of the transect,

451 $length_i$ and the ordinal day of year, day_{ijt}. Island identity was included as a random effect to

452 account for the nonindependence of transects within the same islands.

$$453 \qquad logit\big(p_{nijt}\big) = p0_n + p1_n \text{length}_i + p2_n \text{day}_{ijt} + island_{REni}$$

To explore how colonization rate and extinction rate change across 10 years and how

habitat fragmentation mediated these dynamics processes, we modeled site-level extinction

and colonization ($\varepsilon_{ni,t}$, $\gamma_{ni,t}$) each as a logit-linear function of 5 covariates: year, island area,

isolation, the interaction between year and area and between isolation and year. Year was

added as a random slope, thus allowing the temporal trend in the colonization or extinction

to vary with island area and isolation:

$$460 \qquad logit\big(\varepsilon_{ni,t}\big) = \alpha0_n + \alpha1_n \text{area}_i + \alpha2_n \text{isolation}_i + (\alpha3_n + RE_i)\text{year}_t + \alpha4_n \text{area}_i \text{year}_t$$

$$461 \qquad\qquad + \alpha5_n \text{isolation}_i \text{year}_t + island_{REni}$$

$$462 \qquad logit\big(\gamma_{ni,t}\big) = \beta0_n + \beta1_n \text{area}_i + \beta2_n \text{isolation}_i + (\beta3_n + RE_i)\text{year}_t + \beta4_n \text{area}_i \text{year}_t$$

$$463 \qquad\qquad + \beta5_n \text{isolation}_i \text{year}_t + island_{REni}$$

In all cases, continuous covariates were z-transformed (mean of 0 and a standard

deviation of 1). There is no strong correlation between island area and isolation (Pearson

correlation r = -0.214) so these effects can be considered independent. We fit the MSOM

with JAGS (Plummer, 2003) using the package rjags (Plummer, 2023) in R v4.3.1 (R Core

Team, 2023). We used vague priors (e.g., normal with $\mu = 0$, $\tau = 0.01$). We ran three chains

for 60,000 iterations, discarded the first 40,000 as burn-in and thinned by 20, yielding a

combined posterior sample of 3000. We extracted 600 iterations across three chains for Z to

calculate CTI_{occur}. Convergence was checked visually with trace plots and confirmed with a

Gelman–Rubin statistic < 1.1 (Gelman & Rubin, 1992). Inference on parameters was made

using 95% and 80% Bayesian credible intervals. We checked the posterior predictive ability

of the model fit by calculating Bayesian p-values ($p = 0.488$, indicating an unbiased

estimation of our model) (Gelman et al., 1996).

4.6   Statistical analysis

4.6.1   Thermophilization of bird communities

To test for occurrence-based thermophilization at the island level, Bayesian linear regression

using R2jags (Su & Yajima, 2021) was used to derive a posterior estimate of the temporal

trend in CTI_{occur} while propagating error in the estimation of CTI_{occur}. The continuous

variable *year* (z-transformed) was incorporated as the only predictor. To account for the

nonindependence of data within islands, we included island identity as a random intercept,

thus allowing intercepts to vary across islands. We used vague priors (e.g., normal with $\mu = 0$,

$\tau = 0.001$) for intercept and year effect. We ran three chains for 10,000 iterations, discarded

the first 5,000 as burn-in and thinned by 5, yielding a combined posterior sample of 3000.

We used the function *MCMCsummary* to get posterior intervals of the intercept and year

effect, which were then used for plotting.

To test for abundance-based thermophilization, linear mixed effect models (LMM) with

restricted maximum likelihood (REML) were used to model temporal trends in CTI_{abun} using

function *lme* in R package nlme (Pinheiro et al., 2017). As with the occurrence analysis, we

used the continuous variable *year* (z-transformed) as the only fixed effect and island as a

random intercept. To account for the temporal autocorrelation of CTI from the same island

through time, we included the first-order correlation structure (e.g. the corAR1 structure)

because observations sampled from the nearby years may be more similar.

We also tested whether the overall regional bird composition (i.e. gamma diversity)

experienced thermophilization. Bayesian linear regression was used for modeling temporal

trends in regional CTI_{occur}. For abundance, a generalized least square (GLS) model was used

to test the relationship between CTI_{abun} and continuous variable *year* while accounting for

potential temporal autocorrelation. GLS was conducted using *gls* function in nlme R

package (Pinheiro et al., 2023).

4.6.2   *Mechanisms underlying CTI trends*

To explore the dynamics underlying trends in CTI_{occur}, specifically, whether

thermophilization was accompanied by increasing occupancy of warm-adapted species and

decreasing occupancy of cold-adapted species, we extracted 600 posterior samples of

occupancy rate per species, and summarized ψ_{nt} by its mean and variation (logit-transformed

to meet normality). We then modeled each species' temporal trend in occupancy (ψ_{nt}) over

time using Bayesian linear regression in R2jags (Su & Yajima, 2021) to account for

propagated uncertainty in $\psi_{nt.}$ Similarly, to decompose the CTI_{abun} trend, we modeled each

species' temporal trend in maximum abundance using a generalized linear mixed effect

model with Poisson error structure while accounting for potential overdispersion – via an

observation-specific random effect (Harrison, 2014). Weighted linear regression models

tested for a relationship between STI and occupancy trend and between STI and abundance

trend. This analysis was conducted using function *lmer* in package lme4 (D. Bates et al.,

2015). Temporal correlations between occupancy and colonization or extinction rates were

measured with Pearson correlations, using colonization and extinction rates derived directly

from the fitted MSOM.

As a complementary analysis, we also investigated the trend in species richness or total

abundance of the two thermal preference groups, separately. The total richness or abundance

in each group was calculated on each transect and in each year. We then modeled temporal

trends of natural logarithm-transformed richness or abundance using LMM with island

identity as the random effect. Partial regression plots from LMM were produced using

ggeffects R package (Lüdecke, 2018). For each model, we checked for normality of

residuals and computed goodness-of-fit metrics (R^2) including conditional and marginal R^2

(Nakagawa & Schielzeth, 2013) using performance R package (Lüdecke et al., 2021). All

continuous fixed effects in the models were standardized (mean = 0 and SD = 1) to ease

computation and facilitate the interpretation and comparison of coefficients within and

across models (Schielzeth, 2010).

4.6.3 The mediating effects of habitat fragmentation on colonization and extinction
processes

An increasing colonization trend of warm-adapted species and increasing extinction trend of

cold-adapted species are two main expected processes that cause thermophilization

(Fourcade et al., 2021). To test our third prediction about the mediating effect of habitat

fragmentation, we selected warm-adapted species that had an increasing trend in

colonization rate (positive year effect in colonization rate) and cold-adapted species that had

an increasing extinction rate (positive year effect in extinction rate). For each of these

species, we extracted their posterior effect of year, area, isolation, and the interaction terms

from previously fit models. The interaction terms reveal if habitat fragmentation positively

or negatively affects either colonization or extinction processes.

Acknowledgements

540 We sincerely thank numerous graduate students in our group for the bird surveys in the field,

and the Xin'an River Ecological Development Group Corporation, Chun'an Forestry Bureau, and the Thousand Island Lake National Forest Park for research permits. We also thank Dr. Di Zeng and Dr. Yuhao Zhao from East China Normal University for their insightful comments on the research. Zhe Liu from Ludong University helped a lot in learning ArcMap.

Funding

This study is supported by the National Natural Science Foundation of China (#32030066 to P.D.; #32071545 to Xingfeng Si), Natural Science Foundation of Zhejiang Province (#LD21C030002 to Ping Ding). The funders had roles in study design, data collection and interpretation, and the decision to submit the work for publication.

Author contributions

Juan Liu: Conceptualization; data curation; formal analysis; methodology; visualization; writing – original draft; writing – review and editing. **Morgan W. Tingley:** formal analysis; methodology; visualization; writing – review and editing. **Qiang Wu:** data curation; visualization; writing – original draft (supporting); writing – review and editing. **Peng Ren:** data curation; visualization; writing – original draft (supporting); writing – review and editing. **Tinghao Jin:** data curation; formal analysis; methodology; visualization; writing – original draft (supporting); writing – review and editing. **Ping Ding**: Supervision; conceptualization; data curation; funding acquisition; methodology; writing – review and editing. **Xingfeng Si**: Supervision; conceptualization; methodology; data curation; funding acquisition; writing – review and editing.

Competing interests

The authors declare no conflicts of interests.

Data availability

The data that support the findings of this study are openly available in Figshare at

https://figshare.com/s/7a16974114262d280ef7.

Code availability

The code that supports the findings of this study is openly available in Figshare at

https://figshare.com/s/7a16974114262d280ef7.

References

Anderson, C. M., Fahrig, L., Rausch, J., Martin, J.-L., Daufresne, T., & Smith, P. A. (2023). Climate-related range shifts in Arctic-breeding shorebirds. *Ecology and Evolution, 13*(2), e9797. doi:10.1002/ece3.9797

Barnagaud, J. Y., Barbaro, L., Hampe, A., Jiguet, F., & Archaux, F. (2013). Species' thermal preferences affect forest bird communities along landscape and local scale habitat gradients. *Ecography, 36*(11), 1218-1226. doi:10.1111/j.1600-0587.2012.00227.x

Bates, A. E., Stuart-Smith, R. D., Barrett, N. S., & Edgar, G. J. (2017). Biological interactions both facilitate and resist climate-related functional change in temperate reef communities. *Proceedings of the Royal Society B: Biological Sciences, 284*(1856), 20170484. doi:10.1098/rspb.2017.0484

Bates, D., Maechler, M., Bolker, B., & Walker, S. (2015). Fitting Linear Mixed-Effects Models Using lme4. Journal of Statistical Software, 67(1), 1-48. In.

Brooker, L., Brooker, M., & Cale, P. (1999). Animal dispersal in fragmented habitat: measuring habitat connectivity, corridor use, and dispersal mortality. *Conservation Ecology, 3*(1), 4. doi:10.5751/ES-00109-030104

Chen, I.-C., Hill, J. K., Ohlemüller, R., Roy, D. B., & Thomas, C. D. (2011). Rapid range shifts of species associated with high levels of climate warming. *Science, 333*(6045), 1024-1026. doi:10.1126/science.1206432

Curley, S. R., Manne, L. L., Ramírez-Garofalo, J. R., & Veit, R. R. (2022). Evaluating compositional changes in the avian communities of eastern North America using temperature and precipitation indices. *Journal of Biogeography, 49*(4), 739-752. doi:10.1111/jbi.14340

DeGregorio, B. A., Westervelt, J. D., Weatherhead, P. J., & Sperry, J. H. (2015). Indirect

effect of climate change: Shifts in ratsnake behavior alter intensity and timing of

avian nest predation. *Ecological Modelling, 312*, 239-246.

doi:10.1016/j.ecolmodel.2015.05.031

Devictor, V., Julliard, R., Couvet, D., & Jiguet, F. (2008). Birds are tracking climate

warming, but not fast enough. *Proceedings of the Royal Society B: Biological*

Sciences, 275(1652), 2743-2748. doi:10.1098/rspb.2008.0878

Dunn, P. O., & Møller, A. P. (2019). *Effects of climate change on birds*. Oxford, UK: Oxford

University Press.

Ewers, R. M., & Banks-Leite, C. (2013). Fragmentation Impairs the Microclimate Buffering

Effect of Tropical Forests. *PLOS ONE, 8*(3), e58093.

doi:10.1371/journal.pone.0058093

Fahrig, L. (2003). Effects of habitat fragmentation on biodiversity. *Annual review of ecology,*

evolution, and systematics, 34(1), 487-515.

doi:0.1146/annurev.ecolsys.34.011802.132419

Feeley, K., Bravo-Avila, C., Fadrique, B., Perez, T., & Zuleta, D. (2020). Climate-driven

changes in the composition of New World plant communities. *Nature Climate*

Change, 10(10), 965-970. doi:10.1038/s41558-020-0873-2

Fourcade, Y., WallisDeVries, M. F., Kuussaari, M., van Swaay, C. A., Heliölä, J., &

Öckinger, E. (2021). Habitat amount and distribution modify community dynamics

under climate change. *Ecology Letters, 24*(5), 950-957. doi:10.1111/ele.13691

Freeman, B. G., Song, Y., Feeley, K. J., & Zhu, K. (2021). Montane species track rising

temperatures better in the tropics than in the temperate zone. *Ecology Letters, 24*(8),

1697-1708. doi:10.1111/ele.13762

Gaüzère, P., Princé, K., & Devictor, V. (2017). Where do they go? The effects of topography

and habitat diversity on reducing climatic debt in birds. *Global Change Biology,*

23(6), 2218-2229. doi:10.1111/gcb.13500

Gelman, A., Meng, X.-L., & Stern, H. (1996). Posterior predictive assessment of model

fitness via realized discrepancies. *Statistica sinica, 6*(1996), 733-760.

doi:10.2307/24306036

Gelman, A., & Rubin, D. B. (1992). Inference from iterative simulation using multiple

sequences. *Statistical science, 7*(4), 457-472. doi:10.1214/ss/1177011136

Gottfried, M., Pauli, H., Futschik, A., Akhalkatsi, M., Barančok, P., Alonso, J. L. B., . . .

Kazakis, G. (2012). Continent-wide response of mountain vegetation to climate

change. *Nature Climate Change, 2*(2), 111-115. doi:10.1038/NCLIMATE1329

Harrison, X. A. (2014). Using observation-level random effects to model overdispersion in

count data in ecology and evolution. *PeerJ, 2*, e616. doi:10.7717/peerj.616

Hu, R., Gu, Y., Luo, M., Lu, Z., Wei, M., & Zhong, J. (2020). Shifts in bird ranges and

conservation priorities in China under climate change. *PLOS ONE, 15*(10), e0240225.

doi:10.1371/journal.pone.0240225

Hylander, K., & Ehrlén, J. (2013). The mechanisms causing extinction debts. *Trends in*

ecology & evolution, 28(6), 341-346. doi:10.1016/j.tree.2013.01.010

La Sorte, F. A., & Jetz, W. (2012). Tracking of climatic niche boundaries under recent

climate change. *Journal of Animal Ecology, 81*(4), 914-925. doi:10.1111/j.1365-

2656.2012.01958.x

La Sorte, F. A., & Thompson, F. R. (2007). Poleward shifts in winter ranges of North

American birds. *Ecology, 88*(7), 1803-1812. doi:10.1890/06-1072.1

Lajeunesse, A., & Fourcade, Y. (2022). Temporal analysis of GBIF data reveals the

restructuring of communities following climate change. *Journal of Animal Ecology,*

92(2), 391-402. doi:10.1111/1365-2656.13854

Lindström, Å., Green, M., Paulson, G., Smith, H. G., & Devictor, V. (2013). Rapid changes

in bird community composition at multiple temporal and spatial scales in response to

recent climate change. *Ecography, 36*(3), 313-322. doi:10.1111/j.1600-

0587.2012.07799.x

Liu, J., Matthews, T. J., Zhong, L., Liu, J., Wu, D., & Yu, M. (2020). Environmental filtering

underpins the island species—area relationship in a subtropical anthropogenic

archipelago. *Journal of Ecology, 108*(2), 424-432. doi:10.1111/1365-2745.13272

Lüdecke, D. (2018). ggeffects: Tidy data frames of marginal effects from regression models.

Journal of Open Source Software, 3(26), 772. doi:10.21105/joss.00772

Lüdecke, D., Ben-Shachar, M. S., Patil, I., Waggoner, P., & Makowski, D. (2021).

performance: An R Package for Assessment, Comparison and Testing of Statistical

Models. *Journal of Open Source Software, 6*(60), 3139.

Luoto, M., & Heikkinen, R. (2008). Disregarding topographical heterogeneity biases species

turnover assessments based on bioclimatic models. *Global Change Biology, 14*(3),

483-494. doi:10.1111/j.1365-2486.2007.01527.x

MacKenzie, D. I., Bailey, L. L., & Nichols, J. D. (2004). Investigating species co-occurrence

patterns when species are detected imperfectly. *Journal of Animal Ecology, 73*(3),

546-555. doi:10.1111/j.0021-8790.2004.00828.x

MacKenzie, D. I., Nichols, J. D., Hines, J. E., Knutson, M. G., & Franklin, A. B. (2003).

Estimating site occupancy, colonization, and local extinction when a species is

detected imperfectly. *Ecology, 84*(8), 2200-2207. doi:10.1890/02-3090

MacKenzie, D. I., Nichols, J. D., Royle, J. A., Pollock, K. H., & Hines, J. E. (2006).

Occupancy estimation and modeling: inferring patterns and dynamics of species

occurrence: Academic Press.

McGuire, J. L., Lawler, J. J., McRae, B. H., Nuñez, T. A., & Theobald, D. M. (2016).

Achieving climate connectivity in a fragmented landscape. *Proceedings of the*

National Academy of Sciences, 113(26), 7195-7200. doi:10.1073/pnas.1602817113

Menéndez, R., Megías, A. G., Hill, J. K., Braschler, B., Willis, S. G., Collingham, Y., . . .

Thomas, C. D. (2006). Species richness changes lag behind climate change.

Proceedings of the Royal Society B: Biological Sciences, 273(1593), 1465-1470.

Nakagawa, S., & Schielzeth, H. (2013). A general and simple method for obtaining R2 from

generalized linear mixed-effects models. *Methods in Ecology and Evolution 4*(2),

133-142. doi:10.1111/j.2041-210x.2012.00261.x

Newson, S. E., Oliver, T. H., Gillings, S., Crick, H. Q., Morecroft, M. D., Duffield, S. J., . . .

Pearce-Higgins, J. W. (2014). Can site and landscape-scale environmental attributes

buffer bird populations against weather events? *Ecography, 37*(9), 872-882.

doi:10.1111/ecog.00575

Oliver, T. H., Gillings, S., Pearce-Higgins, J. W., Brereton, T., Crick, H. Q., Duffield, S.

J., . . . Roy, D. B. (2017). Large extents of intensive land use limit community

reorganization during climate warming. *Global Change Biology, 23*(6), 2272-2283.

doi:10.1111/gcb.13587

Opdam, P., & Wascher, D. (2004). Climate change meets habitat fragmentation: linking

landscape and biogeographical scale levels in research and conservation. *Biological*

Conservation, 117(3), 285-297. doi:10.1016/j.biocon.2003.12.008

Paquette, A., & Hargreaves, A. L. (2021). Biotic interactions are more often important at

species' warm versus cool range edges. *Ecology Letters, 24*(11), 2427-2438.

doi:10.1111/ele.13864

Pearce-Higgins, J. W., Eglington, S. M., Martay, B., & Chamberlain, D. E. (2015). Drivers

of climate change impacts on bird communities. *Journal of Animal Ecology, 84*(4),

943-954. doi:10.1111/1365-2656.12364

Pinheiro, J., Bates, D., DebRoy, S., Sarkar, D., Heisterkamp, S., Van Willigen, B., &

Maintainer, R. (2017). nlme: Linear and Nonlinear Mixed Effects Models. R package

version 3.1-162.

Pinheiro, J., Bates, D., & Team, R. C. (2023). nlme: Linear and Nonlinear Mixed Effects

Models. R package version 3.1-162. In.

Platts, P. J., Mason, S. C., Palmer, G., Hill, J. K., Oliver, T. H., Powney, G. D., . . . Thomas,

C. D. (2019). Habitat availability explains variation in climate-driven range shifts

across multiple taxonomic groups. *Scientific Reports, 9*(1), 1-10.

doi:10.1038/s41598-019-51582-2

Plummer, M. (2003). *JAGS: A program for analysis of Bayesian graphical models using*

Gibbs sampling. Paper presented at the Proceedings of the 3rd international

workshop on distributed statistical computing.

Plummer, M. (2023). rjags: Bayesian graphical models using MCMC. R package version 4-

14. In.

Prince, K., & Zuckerberg, B. (2015). Climate change in our backyards: the reshuffling of

North America's winter bird communities. *Global Change Biology, 21*(2), 572-585.

doi:10.1111/gcb.12740

Richard, B., Dupouey, J. l., Corcket, E., Alard, D., Archaux, F., Aubert, M., . . . Macé, S.

(2021). The climatic debt is growing in the understorey of temperate forests: Stand

characteristics matter. *Global Ecology and Biogeography, 30*(7), 1474-1487.

doi:10.1111/geb.13312

Rigal, S., Dakos, V., Alonso, H., Auniņš, A., Benkő, Z., Brotons, L., . . . Del Moral, J. C.

(2023). Farmland practices are driving bird population decline across Europe.

Proceedings of the National Academy of Sciences, 120(21), e2216573120.

doi:10.1073/pnas.2216573120

Royle, J. A., & Kéry, M. (2007). A Bayesian state-space formulation of dynamic occupancy

models. *Ecology, 88*(7), 1813-1823. doi:10.1890/06-0669.1

Sagarin, R. D., & Gaines, S. D. (2002). The 'abundant centre' distribution: to what extent is

it a biogeographical rule? *Ecology Letters, 5*(1), 137-147. doi:10.1046/j.1461-

0248.2002.00297.x

Schielzeth, H. (2010). Simple means to improve the interpretability of regression

coefficients. *Methods in Ecology and Evolution 1*(2), 103-113. doi:10.1111/j.2041-

210X.2010.00012.x

Schoereder, J. H., Galbiati, C., Ribas, C. R., Sobrinho, T. G., Sperber, C. F., DeSouza, O., &

Lopes-Andrade, C. (2004). Should we use proportional sampling for species-area

studies? *Journal of Biogeography, 31*(8), 1219-1226. doi:10.1111/j.1365-

2699.2004.01113.x

Si, X., Cadotte, M. W., Zeng, D., Baselga, A., Zhao, Y., Li, J., . . . Ding, P. (2017).

Functional and phylogenetic structure of island bird communities. *Journal of Animal*

Ecology, 86(3), 532-542. doi:10.1111/1365-2656.12650

Si, X., Cadotte, M. W., Zhao, Y., Zhou, H., Zeng, D., Li, J., . . . Tingley, M. W. (2018). The

importance of accounting for imperfect detection when estimating functional and

phylogenetic community structure. *Ecology, 99*(9), 2103-2112. doi:10.1002/ecy.2438

Si, X., Pimm, S. L., Russell, G. J., & Ding, P. (2014). Turnover of breeding bird

communities on islands in an inundated lake. *Journal of Biogeography, 41*(12),

2283-2292. doi:10.1111/jbi.12379

Su, Y.-S., & Yajima, M. (2021). R2jags: Using R to Run 'JAGS'. R package version 0.7-1. In.

Suggitt, A. J., Wilson, R. J., Isaac, N. J., Beale, C. M., Auffret, A. G., August, T., . . . Fox, R.

(2018). Extinction risk from climate change is reduced by microclimatic buffering.

Nature Climate Change, 8(8), 713-717. doi:10.1038/s41558-018-0231-9

Tayleur, C. M., Devictor, V., Gaüzère, P., Jonzén, N., Smith, H. G., & Lindström, Å. (2016).

Regional variation in climate change winners and losers highlights the rapid loss of

cold-dwelling species. *Diversity and Distributions, 22*(4), 468-480.

doi:10.1111/ddi.12412

Team, R. C. (2023). R: A Language and Environment for Statistical Computing. R

Foundation for Statistical Computing. In. Vienna, Austria.

Thomas, C. D., Franco, A. M., & Hill, J. K. (2006). Range retractions and extinction in the

face of climate warming. *Trends in Ecology & Evolution 21*(8), 415-416.

doi:10.1016/j.tree.2006.05.012

Thomas, C. D., & Lennon, J. J. (1999). Birds extend their ranges northwards. *Nature,*

399(6733), 213-213. doi:10.1038/20335

Tingley, M. W., & Beissinger, S. R. (2009). Detecting range shifts from historical species

occurrences: new perspectives on old data. *Trends in ecology & evolution, 24*(11),

625-633. doi:10.1016/j.tree.2009.05.009

Tingley, M. W., & Beissinger, S. R. (2013). Cryptic loss of montane avian richness and high

community turnover over 100 years. *Ecology, 94*(3), 598-609. doi:10.1890/12-0928.1

Warren, M., Hill, J., Thomas, J., Asher, J., Fox, R., Huntley, B., . . . Harding, P. (2001).

Rapid responses of British butterflies to opposing forces of climate and habitat

change. *Nature, 414*(6859), 65-69. doi:10.1038/35102054

Zellweger, F., Frenne, P. D., Lenoir, J., Vangansbeke, P., & Coomes, D. (2020). Forest

microclimate dynamics drive plant responses to warming. *Science, 368*(6492), 772-

775. doi:10.1126/science.aba6880

Figures

Figure 1. The general process of thermophilization and the hypothesized framework of thermophilization in fragmented habitats. Two general processes of thermophilization are shown on the top: increasing colonization rate of warm-adapted species, and increasing extinction rate of cold-adapted species over time (**a**). Compared to continuous habitats, a hotter microclimate caused by lower buffering ability on fragmented patches may attract more warm-adapted species to colonize while causing cold-adapted species to extirpate or emigrate faster (**b**). Loss of cold-adapted species can also be exacerbated when highly fragmented patches harbor lower habitat heterogeneity (e.g. resource, microrefugia), which will also reduce colonization of warm-adapted species (**c**). Isolated patches due to habitat fragmentation will block warm-adapted species' colonization and cold-adapted species' emigration under warming. The three distinct patches signify a fragmented background and the community in the middle of the three patches was selected to exhibit colonization-extinction dynamics in fragmented habitats. Relative species richness is shown by the number of bird silhouettes in the community (**d**). Note that extinction here may include both

the emigration of species and then the local extinction of species and that CTI can also

change simply due to differential colonization-extinction rates by thermal affinity if the

system is not at equilibrium prior to the study. In our study system, we have no way of

knowing whether our island system was at equilibrium at onset of the study, thus, focusing

on changing rates of colonization-extinction over time presents a much stronger tests of

thermophilization.

Figure 2. The location of Thousand Island Lake and the experimental islands. The map

was created using ESRI (Environmental Systems Resource Institute) ArcMap software

(version 10.3). The base map sources include Esri, Maxar, Earthstar Geographics, and the

GIS User Community.

Figure 3. Temporal trend in CTI$_{occur}$ and CTI$_{abun}$ in the TIL. Vertical lines in **(a)** are

posterior estimated CTI (mean ± SD); solid black line and shaded area are predictions and

95% credible intervals extracted from JAGS modeling posterior mean CTI$_{occur}$ as a function

of year, and using island identity as a random effect while accounting for the variation in

posterior CTI$_{occur}$. Points and error bars in **(b)** are observed CTI$_{abun}$ (mean ± 2SD); solid

black line and shaded area are predicted values and 95% confidence intervals estimated from

LMM modeling observed CTI$_{abun}$ as a function of year, and using island identity as a random

effect.

Figure 4. Relationship between STI and occupancy trend, abundance trend and the

relationship among species thermal preference and trend in occupancy, colonization,

and extinction rates. Each point and error bar in **(a)** represents the temporal trend of

occupancy rate (posterior mean and 95% credible interval of year effect on occupancy rate)

for each species, with the filled dots indicating a significant year effect while the hollow dots

indicating nonsignificant year effect. The dotted vertical line indicates the median of STI

values. Cold-adapted species are plotted in blue and warm-adapted species are plotted in

orange. The number of species with significant occupancy trends in each quadrant was

added to the plot. The black dashed line and shaded area are the predicted values and 95%

confidence intervals of the weighted linear regression model. Each point in **(b)** represents

812   the posterior mean estimate of year in colonization, extinction or occupancy rate for each

813   species. The color of the point indicates the temporal trend in occupancy. Similar to (a), each

814   point and error bar in **(c)** represents the temporal trend of abundance (year effect and 95%

815   confidence interval of year effect on abundance) for each species.

Figure 5. Posterior estimates of logit-scale parameters related to cold-adapted species'

extinction rates and warm-adapted species' colonization rates. Points are species-

specific posterior means on the logit-scale, where parameters > 0 indicate positive effects

(on extinction [a] or colonization [b]) and parameters < 0 indicate negative effects. Only

cold-adapted species with positive trends in extinction **(a)** and warm-adapted species with

positive trends in colonization **(b)** are shown, which are two main processes contributing

positively to thermophilization. Points in blue/orange indicate significant effects (95% or 80%

credible intervals) and points in white indicate non-significant effects. All points were

jittered slightly by 0.22 in width.

Appendix 1

Appendix 1—figure 1. Breeding season temperature from the year 2012 to 2021 in the

Thousand Island Lake, China. Breeding season temperature was calculated as the average

of the mean monthly temperature for breeding seasons (April – June). Black lines and grey

shade indicate the regression line and confidence interval of the linear regression model

(slope $= 0.078$, $r^2 = 34.53\%$, $p = 0.04$).

Appendix 1—figure 2. Robustness of STI to changes in data resources. The units on the

horizontal and vertical axes are degrees Celsius. Pearson correlation coefficients between

STI used in our study and STI calculated with different time-windows: March to August (a)

and January to December (b), were significantly high. Pearson correlation coefficients

between STI used in our study and STI calculated with different distributional ranges: the

distributional range in China (c) and distributional range across the world (d), were

significantly high.

Appendix 1—figure 3. Species Temperature Index of 60 species in the Thousand Islands Lake, China. The dotted horizontal line indicates the median of STI values; the dashed horizontal line indicates the mean of STI values. Grey points are Oriental species, red points are Palearctic species and purple points are Eurytopic species. The bottom right presents the frequency of STI values (right) and the overview relationship between total abundance (natural logarithm transversion of total abundance over 10 years and on all islands) and STI.

Appendix 1—figure 4. Relationship between STI and mean estimates and 95%, and 80%

credible intervals of posterior distribution for temporal trends of species extinction rate

(a) and species colonization rate (b). Points and error bars in blue or orange indicate

significant posterior estimates and error bars in grey indicate non-significant effects. Results

of linear regression models modeling posterior mean of temporal trends in colonization or

extinction weighted by standard deviations were added to plots.

Appendix 1—figure 5. Partial regression effects of year effect on total species richness (a) and total abundance (b) for warm versus cold-adapted species groups. The solid lines indicate significant effects. The shaded bands are 95% confidence intervals of fitted values with other variables at their mean values. Total richness was calculated as the mean of the posterior estimate of species absence/presence, while total abundance was calculated from raw maximum abundance data.

Appendix 1—figure 6. Relationship between island area and understory breeding season air temperature of 20 islands monitored in the year 2022. The understory air temperature was monitored in 20 selected study islands differing in island area. Each plot represents the average air temperature (April – June) of a site within the islands. Lines and shaded areas are predicted values and 95% confidence intervals from LM models.