Classification: Biological Sciences. Ecology

Title: Estimation of Species Abundance Based on the Number of Segregating Sites using

Environmental DNA (eDNA)

Short Title: Quantifying Species Abundance Using eDNA

Authors: Qiaoyun Ai*, Hao Yuan†,‡, Ying Wang§, Chenhong Li+

Affiliations:

[1] Shanghai Universities Key Laboratory of Marine Animal Taxonomy and Evolution, Shanghai

Ocean University, Shanghai 201306, China

[2] Engineering Research Center of Environmental DNA and Ecological Water Health

Assessment, Shanghai Ocean University, Shanghai 201306, China

Current address: *Department of Biochemistry and Molecular Biology, Michigan State

University, East Lansing, MI 48823, USA, †Genetics and Genome Sciences Program, Michigan

State University, East Lansing, MI 48823, USA, Ecology, ‡Evolution and Behavior Program,

Michigan State University, East Lansing, MI 48823, USA, §Amoydx, Xiamen, China

+**Correspondence to:** Chenhong Li, chli@shou.edu.cn

Keywords: Environmental DNA (eDNA), Biodiversity Monitoring, Genetic Variation, Aquatic

Organisms

Abstract: The advance of environmental DNA (eDNA) has enabled rapid and non-invasive

species detection in aquatic environments. Although most studies focus on species detections,

some recent studies explored the potential of using eDNA concentration to quantify species

abundance. However, the differential individual DNA contribution to eDNA samples could

easily obscure the concentration-species abundance relationship. We propose using the number

of segregating sites as a proxy for estimating species abundance. Since segregating sites reflects

the genetic diversity of the population, which is less sensitive to differential individual DNA

contribution than eDNA concentration. We examined the relationship between the number of

segregating sites and species abundance in silico, in vitro, and in situ using two brackish goby

species, *Acanthogobius hasta* and *Tridentiger bifasciatus*. Analyses of the simulated data and in

vitro data with DNA mixed from a known number of individuals showed a strong correlation

between the number of segregating sites and species abundance ($R^2 > 0.9$; $P < 0.01$). Results

from the in situ experiment further validated the correlation ($R^2 = 0.70$, $P < 0.01$), and such

correlation was not affected by biotic factors, including body size and feeding behavior ($P >$

0.05). Results of the cross-validation test also showed that the number of segregating sites

predicted species abundance with less bias and variability than the eDNA concentration. Overall,

the number of segregating sites correlates stronger with species abundance and also provides a

better estimate than eDNA concentration. This advancement can significantly enhance the

quantitative capabilities of eDNA technology.

INTRODUCTION

The environmental DNA (eDNA) technique has emerged as a prevalent approach in ecological

studies due to its cost-effectiveness, enhanced efficiency, and non-invasive nature compared to

traditional approaches (Furlan and Gleeson 2016; Hosler 2017; Sigsgaard et al. 2015). By

sampling environment samples from various ecosystems, such as tropical freshwater (Robson et

al. 2016), intermittent river (Bylemans et al. 2016), brackish water (Forsström and Vasemägi

2016), kelp forest (Port et al. 2016), and even terrestrial habitat (Bohmann and Lynggaard 2023),

the eDNA technique can capture ambient DNA released by organisms, which is subsequently

analyzed to detect species. The eDNA approach has been widely applied in biomonitoring,

including endangered (Fukumoto, Ushimaru, and Minamoto 2015) and invasive species (Uchii,

Doi, and Minamoto 2016), allowing for monitoring single or multiple species simultaneously

(DiBattista et al. 2017). Additionally, the eDNA technique has also been expanded into

population studies, which provide a practical and nondestructive approach to determining species

composition (Madduppa et al. 2021), symbiotic relationships (Shinzato et al. 2018), complex

species structures (DiBattista et al. 2017) and breeding habits (Sakata et al. 2017).

Another promising frontier in eDNA research is the estimation of species abundance.

This application, while challenging, holds significant value for conservation ecology and

population genetics. eDNA offers a non-invasive method to monitor changes in species

abundance, particularly for invasive species, thereby informing more effective management

strategies (Matthews and Whittaker 2015). Furthermore, species abundance is a key parameter in

understanding population dynamics. Fluctuations in abundance are closely related to changes in

various population attributes, including genetic diversity (Hughes et al. 2008) and selection

pressures (Lankau and Strauss 2011). Several studies have explored the potential of using eDNA

concentration as a proxy by relating it to species abundance (Chambert et al. 2018; Dunn et al.

2017:201; Hänfling et al. 2016; Yusishen et al. 2020). Some studies reported strong positive

relationships between eDNA concentration and species abundance (Karlsson et al. 2022; Yates,

Fraser, and Derry 2019), while others showed non-significant relationships (Brys et al. 2021).

Those contradictory findings likely resulted from the influences of various biotic and abiotic

factors on eDNA dynamics - shedding, decaying, and transporting, which collectively affect

eDNA concentration (Barnes and Turner 2016). For example, Biological characteristics such as

species (Ficetola et al. 2008; Jerde et al. 2011), tissue source (Merkes et al. 2014), life stage

(Maruyama et al. 2014), and weight (Mizumoto et al. 2018) could affect the shedding rate of

eDNA. Environmental conditions including temperature (Strickler, Fremier, and Goldberg

2015), UV intensity (Strickler et al. 2015), and microbial community (Poté, Ackermann, and

Wildi 2009) play a role in determining the decaying rate of eDNA. Additionally, physical

processes like water flow (Foppen et al. 2011) and precipitation/resuspension (Turner, Uy, and

Everhart 2015) contribute to the transport of eDNA in the ecosystem. These factors combined

lead to the uneven contribution of DNA from individuals among the population. Some

individuals significantly influence the measured concentration, potentially leading to

overestimation of their numbers, while others contribute less to the overall eDNA concentration

and potentially are underrepresented in species abundance estimates. Thus, the differential

contribution of individuals to the collected eDNA samples skews the species abundance

estimates from eDNA concentration (Andres et al. 2023). Consequently, developing a metric

resilient to differential individual contribution to eDNA samples is imperative for achieving an

accurate and precise estimate of species abundance.

To establish a resilient metric against the differential individual contribution to eDNA

samples, we propose using the number of segregating sites - polymorphic sites within the

population - as a proxy for species abundance. As an indicator of genetic variation, the number

of segregating sites rises with increasing sample size in equilibrium populations (Tajima 1989;

Takahashi and Tajima 2017). The inherent redundancy nature of segregating sites ensures that

each individual is accessed only once regardless of the number of copies of DNA detected in the

samples. We investigated the relationship between the number of segregating sites and species

abundance through a comprehensive, three-tiered experimental design. Our study began with an

in silico analysis, where we tested the relationship using completely simulated data generated at

various mutation rates (Fig. 1a). The in silico study was then complemented by an in vitro

analysis, in which we simulated eDNA samples by subsampling sequences from tissues of two

brackish water goby species, *Acanthogobius hasta* and *Tridentiger bifasciatus* (Fig. 1b). Finally,

to evaluate our findings in a more natural context, we conducted in situ (mesocosm) experiments

to analyze eDNA collected from aquariums (Fig. 1c). Results revealed that the number of

segregating sites extracted from enriched sequences showed a stronger correlation and a better

estimate of species abundance than eDNA concentration.

2 MATERIAL AND METHODS

2.1 In Silico Experiment

We first assessed the relationship between the number of segregating sites and species

abundance by entirely simulated sequences. The length of simulated sequences was set at 17,000

bp. The number of simulated sequences/individuals was 1,000, and sequences were generated at

the mutation rate of 10^{-6} /bp/gen. To account for mutation rate variation among different species,

we also generated another two datasets at the mutation rate of 10^{-7} /bp/gen and 10^{-8} /bp/gen.

These mutation rates represent a biologically meaningful range for vertebrates (Allio et al. 2017).

All data were generated using the software Fastsimcoal2 (Excoffier and Foll 2011). A subset of

sequences were randomly chosen from the simulated data, ranging from 20 to 980 sequences

with intervals of 20. Selected sequences were aligned using MUSCLE v1.0 (Edgar 2004), and

then the number of segregating sites was counted from alignments. The simulation process was

repeated three times at each specified number of sequences. The correlation between the number

of segregating sites and the number of individuals/sequences was estimated by regression

analysis using Microsoft Excel, in which the number of segregating sites was the dependent

variable (y), the number of individuals/sequences was the independent variable (x) and the

significance of the correlation was estimated using R^2 and p-value. The simulated sequences are

available in Dryad at https://doi.org/10.5061/dryad.w3r2280zz.

2.2 Target selection

In our study, we employ target enrichment rather than metabarcoding to collect DNA from

eDNA samples. Target enrichment offers superior coverage to PCR, allowing us to extract more

segregating sites from wider genomic regions for species abundance estimates (Li et al. 2023).

Our goal is to design a single set of RNA baits for enrichment that can be applied to a wide range

of species. To achieve this versatility, we focus on identifying mtDNA regions that are highly

conserved across diverse species. This strategy ensures better enrichment across multiple species

while utilizing variable flanking regions to extract segregating sites.

We conducted a comprehensive alignment of mitochondrial genome sequences from a

broad spectrum of vertebrates, including *Danio rerio*, *Cyprinus carpio*, *Oryzias latipes*,

Tetraodon nigroviridis, *Gasterosteus aculeatus*, *Lepisosteus oculatus*, *Anguilla japonica*, *Gadus*

morhua, *Oreochromis niloticus*, *Homo sapiens*, *Xenopus tropicalis*, *Anolis carolinensis*,

Callorhinchus milii, and *Gallus gallus* (Table S1). We selected regions with an average pairwise

similarity above 0.8 and a minimum length of 120 bp to ensure sufficient conservation. This

rigorous selection process yielded 11 segments from the mitochondrial genome that met our

criteria. These segments were subsequently employed as target markers for both our in vitro and

in situ experiments.

2.3 In Vitro Experiment

Before using 11 markers to enrich targets from eDNA samples, we need to test the performance

of using those markers to enrich DNA extracted from tissue samples. Thus, we examined the

correlation between the number of segregating sites and species abundance using enriched

sequences to get a sense of correlation from actual populations. We selected two species of fish

for the in vitro experiment: *A. hasta* (92 samples) were bought from fishermen who caught fish

from the East China Sea in August 2016, and *T. bifasciatus* (78 samples) caught from Dishui

Lake in Shanghai. The DNA of two species was extracted from fin clips using an Ezup Column

Animal Genomic DNA Purification Kit (Sangon, Shanghai, China). The purified DNA was

quantified with a NanoDrop 3300 Fluorospectrometer (Thermo Fisher Scientific, Wilmington,

DE, U.S.A.) and visualized by agarose gel electrophoresis. The primers were designed according

to the mitochondrial genome sequence of *A. hasta* (NCBI accession: AY486321) and *T.*

trigonocephalus (NCBI accession: NC_029738) (Table S1). We evaluated the specificity of the

primers using the software Primer3.

RNA baits used for target enrichment were RT-PCR products of the 11 target segments.

TakaRa Ex Taq® (RR001 Takara) was used to amplify 11 target DNA segments using the DNA

from an *A. hasta* sample *(*CL1176_43) as a template. Eleven pairs of primers used to amplify

target segments are listed in Table S4. The PCR products were checked using agarose gel

electrophoresis and mixed as template transcription. Then, we followed the manual of

MEGAshortscript™ kit (Thermo Fisher Scientific, Wilmington, DE, U.S.A.) to transcribe PCR

products to RNA transcripts. The concentration of purified transcribed product was measured

and then stored at -80 °C until the gene enrichment.

The process of gene enrichment followed Li et al.'s protocol (Li et al. 2013). We sheared

the 1,000 ng purified DNA into around 250 bp using a Covaris M220 Focused-ultrasonicator

(Woburn, Massachusetts, U.S.A.). Sheared DNA was used to prepare libraries with two different

indexes incorporated in both ends. All indexed libraries were mixed into one tube equimolarly

for gene enrichment and Illumina sequencing on a HiSeq 2500 sequencer (Illumina, Inc, San

Diego, CA, U.S.A.).

The enriched sequences were mapped to mtDNA reference sequences of *A. hasta* and *T.*

bifasciatus to get potential target sequences, which were further assembled using the software

Trinity2.06 (Grabherr et al. 2011). We evaluated the enrichment quality based on raw read

numbers, mapped read numbers, enriched loci numbers, and sequencing depth.

We constructed a phylogenetic tree using the assembled sequences to validate the species

identity of our collected samples. Assembled sequences of 92 *A. hasta* and 78 *T. bifasciatus* were

used to build maximum likelihood trees using GTRGAMMA model in RAxML v8.0.0

(Stamatakis 2014) with *Mugilogobius abei* (NCBI accession: NC_023353.1) as outgroup. We

used 92 sequences enriched from *A. hasta* and 78 sequences enriched from *T. bifasciatus* to

evaluate the correlation between the number of segregating sites and the number of individuals.

The sequences were randomly selected across diverse group sizes, ranging from 5 to 90

for *A. hasta*, and 5 to 70 for *T. bifasciatus*. We generated three repeats for each group size. The

sequences in each group were aligned using MUSCLE v1.0 (Edgar 2004), then the number of

segregating sites was counted using MEGA7 (Kumar, Stecher, and Tamura 2016). The

correlation between the number of segregating sites and the number of individuals was evaluated

by regression analysis using Microsoft Excel. The number of segregating sites was set as the

dependent variable (y), while the number of individuals was the independent variable (x) and the

significance of the correlation was estimated by R^2 and p-value.

2.3 In Situ Experiment (Mesocosm experiment)

Finally, we conducted a mesocosm experiment to evaluate the relationship between the number

segregating sites and species abundance in water samples. *A. hasta* used for the mesocosm

experiment were captured from the East China Sea in September 2017. The fish were kept in

dechlorinated water with 5‰ salinity and fed with frozen shrimp before the experiment. The

water in the aquaria was changed at least four times, and the dead fish were removed in a timely

manner until all the fish were accustomed to the aquaria. All equipment, including containers, air

stones, and pipes were sterilized one week before the experiment. To reduce the mortality of fish,

we did not weigh the individuals for size grouping, but divided them into four groups according

to the body length (< 10 cm, 10-12.5 cm, 12.5-15 cm, > 15 cm). The average weight of fish was

measured after the experiment (Table 1). Twenty 65 L plastic containers were prepared to keep

A. hasta (Table 1), and one aquarium without fish was set as negative control. We set five fish

number gradients, including 7, 14, 21, 28, 35. To account for the effect of feeding behavior in the

estimation of species abundance, *Penaeus vannamei* was cut into pieces to feed the fish, at

approximately 5% of the estimated total weight of fish in each aquarium. One liter of water in

every aquarium was collected after three days. The collected water was filtered using 47-mm

GF/F glass filters (pore size c. o.7 μm; Whatman International, Ltd), then the filters were soaked

in pure ethanol and stored at -80 °C until DNA extraction. DNA samples were extracted using an

Ezup Column Animal Genomic DNA Purification Kit (Sangon, Shanghai, China). The purified

eDNA was checked using agarose gel electrophoresis and stored at -20 °C until the target gene

enrichment and qPCR.

For better target enrichment efficiency, we redesigned primers for the in situ experiment

according to sequences we obtained from the in vitro experiment (Table S4). PCR products were

further transcribed and prepared as baits following instructions in section 2.3. By checking

agarose gel electrophoresis, we found that the maximum length of purified eDNA was too large

for next-generation sequencing. Thus, we sheared DNA to around 250 bp using a Covaris M220

Focused-ultrasonicator (Woburn, Massachusetts, U.S.A.). Libraries preparation, target gene

enrichment and sequencing followed the above procedure. After preprocessing raw reads

including trimming adaptors, removing low-quantity reads, and PCR duplicates, the reads of

each sample were mapped to the reference of *A. hasta* using BWA-MEM (Li and Durbin 2010).

The number of segregating sites was counted from the targeted regions as well as flanking

sequences, spanning 50 bp upstream and downstream of the targeted regions. We removed

polymorphic sites with sequencing depth below seven and minor allele frequencies below 3% to

filter out potentially erroneous segregating sites stemming from PCR stutter and sequencing

errors (Fumagalli 2013).

We also accessed the eDNA concentration of each sample. Among the 11 target

segments, T2, the second segment, was chosen as the target sequence for qPCR due to its length

(140 bp) falling within the recommended range for qPCR amplicons (Van Holm et al. 2021). To

create standard samples, we amplified the T2 target region by TakaRa Ex Taq® (RR001 Takara)

using the genomic DNA of CL1176_43 as the template. The amplified product was purified by

gel extraction using DNA gel extraction kit (Qiagen, Hilden). The purified product was further

diluted to create six standard samples for qPCR, with concentrations ranging from 1.75 ng/L to

1.75×10^6 ng/L. The qPCR procedure followed the protocol outlined in KAPA SYBR FAST

Universal qPCR Kit (KAPABIOSYSTEMS, U.S.A). We assessed the specificity of the eDNA

amplified products through a melting curve analysis using the online BIO-RAD melting curve

generator. The copy number was estimated using the online BIO-RAD copy number calculator.

The number of DNA copies was estimated as the mean value of three repeats for the same water

sample (Fig. S2). We averaged the number of DNA copies across replicates and converted this

value to eDNA concentration (copies/L).

Statistical analysis was performed using Stata (https://www.stata.com/). Fixed effect

models were used to determine if fish number, body size, and feeding behavior affected the

number of segregating sites or eDNA concentration. We also constructed another simpler fix

effect model, which only considered the fish number as the variable. We used the F-test, Akaike

Information Criterion (AIC), and Bayesian information criterion (BIC) to discern whether

species abundance is the only factor attributed to the impact on the number of segregating sites

or eDNA concentration.

To take a step further from correlation analysis, we conducted an exhaustive leave-p-out

cross-validation test to compare the performance of estimating species abundance using

segregating sites and eDNA. Specifically, p samples were randomly selected from the mesocosm

dataset as testing samples, while the rest served as training samples. p ranges from 1 to 10.

Training samples were used to establish regression models between the number of segregating

sites/eDNA concentration and species abundance. Then, those models were applied to estimate

the species abundance in test samples. Notably, for every value of p, we exhaustively accessed

the performance for every possible combination of samples. If p was greater than one, the

species abundance of the same sample could be estimated multiple times by models built based

on different training samples. We evaluated estimation performance using absolute bias, i.e., the

absolute difference between ground truth and estimated species abundance. We further averaged

absolute bias for the same sample when assessing performance. The precision of the estimation

of each sample was calculated using the standard deviation of predicted species abundance.

3 RESULTS

3.1 In Silico Analysis

Simulation analysis revealed a strong positive correlation between the number of segregating

sites and species abundance across all mutation rates. All regression models had R^2 above 0.9.

The regression coefficient for the number of segregating sites increased with mutation rate,

reflecting the increased occurrence of segregating sites in the mtDNA of species with higher

mutation rates (Fig. 2). Regression models fit better with log-transformed species abundance

rather than its absolute value, since the number of segregating sites reached saturation when

sampling enough of the number of individuals.

3.2 In Vitro Analysis

Before the mesocosm experiment, we used baits designed from 11 target segments to enrich

DNA from tissues of *A. hasta* and *T. bifasciatus*, including target segments and their flanking

regions (Table S1). The result revealed the successful enrichment of the 11 targets in all samples.

After quality check and PCR duplication removal, an average of 5,304,806 bp reads for *A. hasta*

and 46,355 bp for *T. bifasciatus* remained. Among those, an average of 16,354 bp reads (9%)

were mapped to the assembly of enriched sequences, with an average sequencing depth of 119

for *A. hasta* and 35 for *T. bifasciatus* (Table S2). The length of aligned sequences of *A. hasta* was

9,183 bp on average. For *T. bifasciatus*, the length of aligned sequences was 6,469 bp on

average. We observed 507 segregating sites in *A. hasta*, and 163 segregating sites in *T.*

bifasciatus respectively. The raw reads and the assembled sequences are available at NCBI

Sequence Read Archive (SRA) with accession numbers PRJNA1100443.

Using *M. abei* as an outgroup (NCBI accession: NC_023353.1), phylogenetic analyses

showed that samples were grouped into two clades. The sequences of *A. hasta* were clustered

with the sequence of *A. hasta* retrieved from GenBank (NCBI accession: AY486321). The

sequences of *T. bifasciatus* were grouped with *T. trigonocephalus* (NCBI accession:

NC_029738) as their close sister species (Fig. S1). The topology of the phylogenetic tree and the

grouping pattern of samples ensure the correct identification of the species.

We examined the relationship between the number of segregating sites and species

abundance using enriched sequences from tissues. Regression analysis revealed that the number

of segregating sites was positively correlated with the species abundance in both *A. hasta* and *T.*

bifasciatus (Fig. 3). R^2 values for regression models of both species exceeded 0.9. In contrast to

the in silico study, the absolute value of species abundance fit better with models than log

transformed form since the number of segregating sites observed in collected individuals was far

from reaching saturation. A strong correlation was observed between the number of segregating

sites and species abundance in the enrichment data from both *A. hasta* and *T. bifasciatus*.

However, *A. hasta* exhibited higher genetic variability, resulting in stronger correlations.

Consequently, only *A. hasta* was selected for the subsequent mesocosm experiments.

3.3 In Situ (Mesocosm) Experiments

We enriched target sequences from eDNA samples using baits designed based on 11 targeted

segments. After quality filtering and deduplication, 164,145,279 reads remained. Among those,

20,571 reads (0.59 %) were successfully mapped to the majority consensus assembly of *A. hasta*

from in vitro analysis, with an average sequencing depth of 197 (Table S3). Subsequently, a

depth filter was applied to remove potential false positive polymorphic sites with sequencing

depth below seven and minor allele frequency below 3% (Fumagalli 2013). The raw reads are

available at SRA with accession numbers PRJNA1100453.

Mesocosm Experiments revealed a strong and significant positive correlation between the

number of segregating sites and species abundance ($r = 4.09$, $p < 0.01$). Although we also

observed a positive relationship ($r = 3.08$, $p < 0.01$) between eDNA concentration and species

abundance, species abundance accounted for a smaller proportion of variation in the eDNA

concentration ($R^2 = 0.46$) compared to the number of segregating sites ($R^2 = 0.70$) (Fig. 4).

According to the F-test, AIC and BIC value, species abundance dominated the impact on the

number of segregating sites. At the same time, mean weight and feeding behavior showed no

significant influence (F-test $p = 0.38$). On the contrary, mean weight or feeding behavior

demonstrated a nearly significant effect on eDNA concentration (F-test $p = 0.07$) (Table 2).

The cross-validation results indicated that the number of segregating sites is a more

accurate and precise proxy for estimating species abundance. The median absolute bias of

species abundance predicted based on the number of segregating sites was slightly lower than the

one derived from eDNA concentration. This difference in median absolute bias became more

pronounced when fewer samples were used as training data. Additionally, the upper limit of bias

significantly decreased when predicting species abundance using the number of segregating sites

compared to eDNA concentration (Fig. 5A). We evaluated the precision of species abundance

estimation by examining the standard deviation of predictions for the same sample across

different training datasets. As anticipated, we observed increasing standard deviation when fewer

samples were served as training data. The standard deviation of species abundance estimates

based on segregating sites was consistently lower than those predicted using eDNA

concentration (Fig. 5B).

4 DISCUSSION

Estimating the species abundance in natural ecosystems could provide vital insights into a wide

array of evolutionary and ecological questions, and also hold implications for effective

conservation strategies. Emerging eDNA technologies, as a non-invasive and rapid approach,

represent a promising complement to traditional methods for estimating species abundance. This

study proposes using the number of segregating sites in eDNA as a proxy for estimating species

abundance. Segregating sites represent inherent genetic variability within a population, making

this metric less susceptible to differential individual DNA contributions in eDNA samples than

concentration-based methods. Thus, the segregating sites-based approach ensures a more precise

and less biased estimate of species abundance. Notably, our study has yet to estimate species

abundance in field tests, so the results from the in silico, in vitro, and mesocosm experiments

must be interpreted with caution. But still, our study is an essential step toward accurate and

precise estimates of species abundance using eDNA.

Previous studies have already leveraged genetic variability as an indicator of species

abundance in eDNA samples. Recent methods count distinct microsatellite alleles (Carreon-

Martinez et al. 2014) or mtDNA haplotypes (Yoshitake et al. 2019, 2021) as a representation of

species abundance. Another approach, the DNA mixture model, uses a probabilistic framework

to estimate the number of contributors to eDNA samples based on observed microsatellite alleles

or mtDNA haplotypes (Andres et al. 2021, 2023; Sethi et al. 2019). Both methods utilize distinct

alleles or haplotypes to infer species abundance but differ in complexity. Allele counting

provides a simple, conservative estimate of minimal possible abundance. The DNA mixture

model takes this concept further by considering the relative frequency of distinct

alleles/haplotypes and accounting for potentially unobserved alleles. In contrast, our method

employs a direct correlation between the number of segregating sites, and abundance. This

approach simplifies the analytical process, especially compared to the DNA mixture model,

while still capturing the underlying relationship between genetic variability and species

abundance. Another advantage is that our method does not require sequence assembly when

using haplotypes, allowing us to capture variation even when small pieces of sequencing data are

missing. To apply our method in the field, a pilot study is required to establish the relationship

between the number of segregating sites and species abundance, while theoretically, the allele

counting method or DNA mixture model does not need such a pilot study. On the other hand,

this initial investigation gives us a good sense of when genetic variability is exhausted, thus

helping us avoid scenarios where results might be underestimated without our knowledge.

In the in silico study, we found a strong correlation between genetic variation in

sequences and abundance estimates across all tested mutation rates. Notably, data generated

under higher mutation rates exhibited stronger correlations between segregating sites and

individual counts, with R^2 values of 0.99, 0.98, and 0.91 for mutation rates of 10^{-6}/bp/gen,

10^{-7}/bp/gen, and 10^{-8}/bp/gen. We also notice that the relationship between segregating sites and

species abundance can reach a plateau (Fig. 2), indicating that when genetic variability is

exhausted, the maximum detectable species abundance significantly depends on the population's

mutation rate. Our in silico study suggests that even at the lowest mutation rate (10^{-8}/bp/gen), we

can reliably estimate populations of up to 100 individuals, which suffices for most applications

(Fig. 2c). However, preliminary surveys are advisable for investigating larger populations.

Complementing the in silico study, the in vitro experiments also demonstrated a strong

relationship between the number of segregating sites and species abundance, particularly evident

in *A. hasta* ($R^2 = 0.98$). However, differing from in silico analysis, neither *A. hasta* nor *T.*

bifasciatus showed signs of reaching a plateau in the relationship between segregating sites and

abundance. This finding suggests a broader potential for using segregating sites to estimate

species abundance in large populations. We noticed lower sequence depth and coverage for

enriched data from *T. bifasciatus* in some samples, likely due to excessive PCR cycles during

DNA library preparation. This resulted in fewer on-target sequences for *T. bifasciatus* compared

to *A. hasta* (Table S2). Despite the relatively poor quality of the *T. bifasciatus* enrichment data,

we still identified a strong positive linear relationship between the number of segregating sites

and species abundance. This suggests that low sequencing depth did not significantly impact our

ability to extract segregating sites.

Mesocosm experiment results showed that the number of individuals predicted the

number of segregating sites ($R^2 = 0.70$) better than predicting the eDNA concentration ($R^2 =$

0.46) as expected. We observed that the overall R^2 was lower in the mesocosm experiments

compared to the in vitro and in silico studies. This difference can be attributed to variables

introduced from eDNA samples, such as DNA degradation and the influence of ambient water

chemistry on target enrichment. Counter-intuitively, both the number of segregating sites and the

number of DNA copies were not heavily affected by body size or feeding/non-feeding treatment

in this study, which was inconsistent with previous results (Klymus et al. 2015; Mizumoto et al.

2018). Moreover, eDNA concentrations measured in our study did not show overdispersed

distribution as described in the previous study (Chambert et al. 2018). One possible explanation

is that the experiment duration was too short, and the behavior of the fish was disrupted, so the

size of the fish and their feeding behavior had erratic effects on releasing DNA. Alternatively,

unforeseen factors, such as placing aquaria in proximity to corridors or disturbances, may have

led to varying stress responses and DNA release from the fish.

383   Our cross-validation analysis revealed that species abundance estimates using segregating

sites outperformed those based on eDNA concentration, showing lower absolute bias and

reduced standard deviation. The performance difference became more evident as the testing set

size increased. This is because multiple estimates per sample, derived from models built on

different training sample sets, provided a more comprehensive representation of the estimate bias

and variation distribution. While the segregating sites method showed substantially lower upper

quantile absolute mean bias, the difference in median absolute mean bias was less pronounced

but still favorable for segregating site methods. We anticipate that longer experiments, which

might introduce more confounding factors for the eDNA concentration-based method, could

further distinguish the performance of these two approaches.

393   Nevertheless, a couple of limitations need to be addressed when applying the

segregating-site method. Firstly, there are no global criteria for thresholding read depth and

minor allele frequency to distinguish polymorphism from sequencing errors. We set an ad hoc

criterion of more than seven reads and at least 3% minor-allele reads to call a segregating site.

Alternatively, duplex Sequencing (Schmitt et al. 2012) or adding unique molecular identifiers

(UMIs) (Yoshitake et al. 2021) may be applied to discriminate genuine polymorphic sites from

sequencing errors. Secondly, a pilot survey of the genetic diversity of the population has to be

done before extrapolating the correlation between the number of segregating sites and the

number of individuals in actual samples. There is often a limited number of individuals sampled

in the pilot study. Therefore, future research should explore the optimal number of individuals

for the pilot study and how much we can extrapolate the correlation between the number of

individuals and segregating sites from those pilot study samples. A recommended approach for

estimating the number of individuals needed for a pilot study is to conduct an in silico analysis

similar to the one in our study. Researchers can gather relevant parameters for the population of

target species, such as mutation rates, from existing literature or by estimating them based on

closely related species in the phylogeny. By simulating various scenarios, researchers can obtain

a reliable estimate of the sample size required for an effective pilot study.

In this study, we employed target enrichment instead of the conventional metabarcoding

approach to collect sequences from eDNA samples. Target enrichment has already been applied

in species detection, proving effective in capturing mitochondrial DNA (Jensen et al. 2021; Li et

al. 2023; Wilcox et al. 2018) and even nuclear DNA (Jensen et al. 2021) from water samples.

Our application extends the utility of target enrichment to species abundance estimates.

Compared to metabarcoding, target enrichment is particularly useful when eDNA is partially

degraded or when primer binding sites are inaccessible or partially accessible. This advantage is

crucial, as non-amplifiable sequences in eDNA could still contain novel alleles or haplotypes not

observed in amplifiable sequences, potentially leading to a loss of variability that could bias final

species abundance estimates. Moreover, unlike metabarcoding, target enrichment can extract

sequences from flanking genomic regions (Li et al. 2013), not just short amplifiable regions

suitable for PCR. In our study, this advantage enabled us to obtain segregating sites beyond

baits-targeted regions. While previous studies typically designed species-specific probes (Jensen

et al. 2021; Li et al. 2023; Wilcox et al. 2018), we designed RNA baits based on conserved

mitochondrial regions across vertebrates, allowing our probes to simultaneously capture eDNA

from multiple species, thereby reducing costs associated with species-specific baits design.

There are several challenges when applying target enrichment in the field study. eDNA

samples typically contain genetic material from a diverse range of species, resulting in the

capture of numerous non-target sequences when using the target enrichment method. While

mapping to the reference genome can filter out most non-target reads, this approach may still fail

to exclude some similar sequences from closely related species that differ by only a few base

pairs. This limitation could potentially lead to false positive segregating sites, resulting in

inflated species abundance estimates. Another challenge was the variability in capture efficiency.

Although most eDNA samples achieved satisfactory sequencing depth (over 50x) (Table S3),

some samples exhibited poor capture efficiency. For instance, one sample, FW1N5, only attained

an average sequencing depth of 18x, which might impede the detection of rare segregating sites.

These limitations underscore the need for careful consideration and potential refinement of target

enrichment protocols when applied to eDNA samples.

eDNA techniques provide a non-invasive and effective solution for biomonitoring,

providing insight into ecological and evolutionary processes in the wild. Qualitative studies, such

as species detection, are well-established using eDNA, while quantitative studies that require

accurate species abundance estimates still pose challenges. Our study proposes the use of

segregating site counts as a more precise and less biased proxy than eDNA concentration,

marking a significant advance toward accurate species abundance estimates from eDNA. The

following steps in developing this method will include (1) accurately identifying rare alleles with

low frequencies from eDNA data, (2) extended testing of different species in mesocosms and the

fields for longer durations, and (3) investigating effective sampling strategies for pilot studies.

AUTHOR CONTRIBUTIONS

Q.A. and C.L. designed the study. Q.A., H.Y., and Y.W. conducted the study and collected specimens. Q.A. and H.Y. completed laboratory work, analyzed the data, and interpreted the results. Q.A. and H.Y. wrote the manuscript. C.L. provided feedback on the manuscript. All authors approved the manuscript for publication.

ACKNOWLEDGEMENTS

This work was supported by the National Key Research and Development Program of China (2022YFC2601301). Dr Z. Zhang helped with model selection and correlation analyses.

CONFLICT OF INTEREST STATEMENT

The authors declare no conflict of interest.

DATA ACCESSIBILITY STATEMENT

Simulated sequences for in-silico study are available at https://doi.org/10.5061/dryad.w3r2280zz. Raw sequences of targeted enrichment from tissues of *A. hasta* and *T. bifasciatus* were deposited in NCBI's Sequence Read Archive (BioProject ID: PRJNA1100443). Raw sequences of enriched sequences from eDNA samples were also available at NCBI's Sequence Read Archive (BioProject ID: PRJNA1100453).

BENEFIT-SHARING STATEMENT

The study developed an accurate and precise method for estimating species abundance using eDNA data. This advancement will guide future efforts in non-invasive biomonitoring techniques for conservation ecology and population genetics. All data from the study are freely accessible through the NCBI's Sequence Read Archive and DRYAD digital repository.

REFERENCE

[dataset] Ai, Q., Yuan, H., Wang, Y., Li, C.; 2024; Estimation of Species Abundance Based on
the Number of Segregating Sites using Environmental DNA (eDNA); DRYAD;
10.5061/dryad.w3r2280zz

[dataset] Ai, Q., Yuan, H., Wang, Y., Li, C.; 2024; Estimation of Species Abundance Based on
the Number of Segregating Sites using Environmental DNA (eDNA); Sequence Read Archive;
PRJNA1100443

[dataset] Ai, Q., Yuan, H., Wang, Y., Li, C.; 2024; Estimation of Species Abundance Based on
the Number of Segregating Sites using Environmental DNA (eDNA); Sequence Read Archive;
PRJNA1100453

Allio, Remi, Stefano Donega, Nicolas Galtier, and Benoit Nabholz. 2017. "Large Variation in
479   the Ratio of Mitochondrial to Nuclear Mutation Rate across Animals: Implications for
480   Genetic Diversity and the Use of Mitochondrial DNA as a Molecular Marker." *Molecular*
481   *Biology and Evolution* 34(11):2762–72. doi: 10.1093/molbev/msx197.
482 Andres, Kara J., David M. Lodge, Suresh A. Sethi, and Jose Andrés. 2023. "Detecting and
483   Analysing Intraspecific Genetic Variation with eDNA: From Population Genetics to
484   Species Abundance." *Molecular Ecology* 32(15):4118–32. doi: 10.1111/mec.17031.
485 Andres, Kara J., Suresh A. Sethi, David M. Lodge, and Jose Andrés. 2021. "Nuclear eDNA
486   Estimates Population Allele Frequencies and Abundance in Experimental Mesocosms
487   and Field Samples." *Molecular Ecology* 30(3):685–97. doi: 10.1111/mec.15765.
488 Barnes, Matthew A., and Cameron R. Turner. 2016. "The Ecology of Environmental DNA and
489   Implications for Conservation Genetics." *Conservation Genetics* 17(1):1–17. doi:
490   10.1007/s10592-015-0775-4.
491 Bohmann, Kristine, and Christina Lynggaard. 2023. "Transforming Terrestrial Biodiversity
492   Surveys Using Airborne eDNA." *Trends in Ecology & Evolution* 38(2):119–21. doi:
493   10.1016/j.tree.2022.11.006.
494 Brys, Rein, David Halfmaerten, Sabrina Neyrinck, Quentin Mauvisseau, Johan Auwerx, Michael
495   Sweet, and Joachim Mergeay. 2021. "Reliable eDNA Detection and Quantification of the
496   European Weather Loach (Misgurnus Fossilis)." *Journal of Fish Biology* 98(2):399–414.
497   doi: 10.1111/jfb.14315.
498 Bylemans, Jonas, Elise M. Furlan, Luke Pearce, Trevor Daly, and Dianne M. Gleeson. 2016.
499   "Improving the Containment of a Freshwater Invader Using Environmental DNA
500   (eDNA) Based Monitoring." *Biological Invasions* 18(10):3081–89. doi: 10.1007/s10530-
501   016-1203-5.
502 Carreon-Martinez, Lucia B., Kyle W. Wellband, Timothy B. Johnson, Stuart A. Ludsin, and
503   Daniel D. Heath. 2014. "Novel Molecular Approach Demonstrates That Turbid River
504   Plumes Reduce Predation Mortality on Larval Fish." *Molecular Ecology* 23(21):5366–77.
505   doi: 10.1111/mec.12927.
506 Chambert, Thierry, David S. Pilliod, Caren S. Goldberg, Hideyuki Doi, and Teruhiko Takahara.
507   2018. "An Analytical Framework for Estimating Aquatic Species Density from
508   Environmental DNA." *Ecology and Evolution* 8(6):3468–77. doi: 10.1002/ece3.3764.
DiBattista, Joseph D., Darren J. Coker, Tane H. Sinclair-Taylor, Michael Stat, Michael L.

Berumen, and Michael Bunce. 2017. "Assessing the Utility of eDNA as a Tool to Survey
Reef-Fish Communities in the Red Sea." *Coral Reefs* 36(4):1245–52. doi:
10.1007/s00338-017-1618-1.
Dunn, Nicholas, Victoria Priestley, Alba Herraiz, Richard Arnold, and Vincent Savolainen.
2017. "Behavior and Season Affect Crayfish Detection and Density Inference Using
Environmental DNA." *Ecology and Evolution* 7(19):7777–85. doi: 10.1002/ece3.3316.
Edgar, Robert C. 2004. "MUSCLE: Multiple Sequence Alignment with High Accuracy and High
Throughput." *Nucleic Acids Research* 32(5):1792–97. doi: 10.1093/nar/gkh340.
Excoffier, Laurent, and Matthieu Foll. 2011. "Fastsimcoal: A Continuous-Time Coalescent
Simulator of Genomic Diversity under Arbitrarily Complex Evolutionary Scenarios."
Bioinformatics 27(9):1332–34. doi: 10.1093/bioinformatics/btr124.
Ficetola, Gentile Francesco, Claude Miaud, François Pompanon, and Pierre Taberlet. 2008.
"Species Detection Using Environmental DNA from Water Samples." *Biology Letters*
4(4):423–25. doi: 10.1098/rsbl.2008.0118.
Foppen, Jan Willem, Caesar Orup, Renaud Adell, Vincent Poulalion, and Stefan Uhlenbrook.
2011. "Using Multiple Artificial DNA Tracers in Hydrology." *Hydrological Processes*
25(19):3101–6. doi: 10.1002/hyp.8159.
Forsström, Tiia, and Anti Vasemägi. 2016. "Can Environmental DNA (eDNA) Be Used for
Detection and Monitoring of Introduced Crab Species in the Baltic Sea?" *Marine
Pollution Bulletin* 109(1):350–55. doi: 10.1016/j.marpolbul.2016.05.054.
Fukumoto, Sou, Atushi Ushimaru, and Toshifumi Minamoto. 2015. "A Basin-Scale Application
of Environmental DNA Assessment for Rare Endemic Species and Closely Related
Exotic Species in Rivers: A Case Study of Giant Salamanders in Japan." *Journal of
Applied Ecology* 52(2):358–65. doi: 10.1111/1365-2664.12392.
Fumagalli, Matteo. 2013. "Assessing the Effect of Sequencing Depth and Sample Size in
Population Genetics Inferences." *PLOS ONE* 8(11):e79667. doi:
10.1371/journal.pone.0079667.
Furlan, Elise M., and Dianne Gleeson. 2016. "Environmental DNA Detection of Redfin Perch,
Perca Fluviatilis." *Conservation Genetics Resources* 8(2):115–18. doi: 10.1007/s12686-
016-0523-1.
Grabherr, Manfred G., Brian J. Haas, Moran Yassour, Joshua Z. Levin, Dawn A. Thompson, Ido
Amit, Xian Adiconis, Lin Fan, Raktima Raychowdhury, Qiandong Zeng, Zehua Chen,
Evan Mauceli, Nir Hacohen, Andreas Gnirke, Nicholas Rhind, Federica di Palma, Bruce
W. Birren, Chad Nusbaum, Kerstin Lindblad-Toh, Nir Friedman, and Aviv Regev. 2011.
"Full-Length Transcriptome Assembly from RNA-Seq Data without a Reference
Genome." *Nature Biotechnology* 29(7):644–52. doi: 10.1038/nbt.1883.
Hänfling, Bernd, Lori Lawson Handley, Daniel S. Read, Christoph Hahn, Jianlong Li, Paul
Nichols, Rosetta C. Blackman, Anna Oliver, and Ian J. Winfield. 2016. "Environmental
DNA Metabarcoding of Lake Fish Communities Reflects Long-Term Data from
Established Survey Methods." *Molecular Ecology* 25(13):3101–19. doi:
10.1111/mec.13660.
Hosler, Denise. 2017. "Where Is the Body? Dreissenid Mussels, Raw Water Testing, and the
Real Value of Environmental DNA." *Management of Biological Invasions* 8(3):335–41.
doi: 10.3391/mbi.2017.8.3.07.
Hughes, A. Randall, Brian D. Inouye, Marc T. J. Johnson, Nora Underwood, and Mark Vellend.
2008. "Ecological Consequences of Genetic Diversity." *Ecology Letters* 11(6):609–23.

doi: 10.1111/j.1461-0248.2008.01179.x.

Jensen, Mads Reinholdt, Eva Egelyng Sigsgaard, Shenglin Liu, Andrea Manica, Steffen Sanvig Bach, Michael Møller Hansen, Peter Rask Møller, and Philip Francis Thomsen. 2021. "Genome-Scale Target Capture of Mitochondrial and Nuclear Environmental DNA from Water Samples." *Molecular Ecology Resources* 21(3):690–702. doi: 10.1111/1755-0998.13293.

Jerde, Christopher L., Andrew R. Mahon, W. Lindsay Chadderton, and David M. Lodge. 2011. "'Sight-Unseen' Detection of Rare Aquatic Species Using Environmental DNA." *Conservation Letters* 4(2):150–57. doi: 10.1111/j.1755-263X.2010.00158.x.

Karlsson, Erik, Martin Ogonowski, Göran Sundblad, Josefin Sundin, Ofir Svensson, Ilkka Nousiainen, and Anti Vasemägi. 2022. "Strong Positive Relationships between eDNA Concentrations and Biomass in Juvenile and Adult Pike (Esox Lucius) under Controlled Conditions: Implications for Monitoring." *Environmental DNA* 4(4):881–93. doi: 10.1002/edn3.298.

Klymus, Katy E., Catherine A. Richter, Duane C. Chapman, and Craig Paukert. 2015. "Quantification of eDNA Shedding Rates from Invasive Bighead Carp Hypophthalmichthys Nobilis and Silver Carp Hypophthalmichthys Molitrix." *Biological Conservation* 183:77–84. doi: 10.1016/j.biocon.2014.11.020.

Kumar, Sudhir, Glen Stecher, and Koichiro Tamura. 2016. "MEGA7: Molecular Evolutionary Genetics Analysis Version 7.0 for Bigger Datasets." *Molecular Biology and Evolution* 33(7):1870–74. doi: 10.1093/molbev/msw054.

Lankau, Richard A., and Sharon Y. Strauss. 2011. "Newly Rare or Newly Common: Evolutionary Feedbacks through Changes in Population Density and Relative Species Abundance, and Their Management Implications." *Evolutionary Applications* 4(2):338–53. doi: 10.1111/j.1752-4571.2010.00173.x.

Li, Chenhong, Michael Hofreiter, Nicolas Straube, Shannon Corrigan, and Gavin J. P. Naylor. 2013. "Capturing Protein-Coding Genes across Highly Divergent Species." *BioTechniques* 54(6):321–26. doi: 10.2144/000114039.

Li, Heng, and Richard Durbin. 2010. "Fast and Accurate Long-Read Alignment with Burrows–Wheeler Transform." *Bioinformatics* 26(5):589–95. doi: 10.1093/bioinformatics/btp698.

Li, Juan, Peter Seeber, Jan Axtner, Rachel Crouthers, Milou Groenenberg, Arnulf Koehncke, Alexandre Courtiol, Pin Chanratana, and Alex D. Greenwood. 2023. "Monitoring Terrestrial Wildlife by Combining Hybridization Capture and Metabarcoding Data from Waterhole Environmental DNA." *Biological Conservation* 284:110168. doi: 10.1016/j.biocon.2023.110168.

Madduppa, Hawis, Ni Kadek Dita Cahyani, Aji Wahyu Anggoro, Beginer Subhan, Edwin Jefri, Lalu M. Iqbal Sani, Dondy Arafat, Nebuchadnezzar Akbar, and Dietriech Geoffrey Bengen. 2021. "eDNA Metabarcoding Illuminates Species Diversity and Composition of Three Phyla (Chordata, Mollusca and Echinodermata) across Indonesian Coral Reefs." *Biodiversity and Conservation* 30(11):3087–3114. doi: 10.1007/s10531-021-02237-0.

Maruyama, Atsushi, Keisuke Nakamura, Hiroki Yamanaka, Michio Kondoh, and Toshifumi Minamoto. 2014. "The Release Rate of Environmental DNA from Juvenile and Adult Fish." *PLOS ONE* 9(12):e114639. doi: 10.1371/journal.pone.0114639.

Matthews, Thomas J., and Robert J. Whittaker. 2015. "REVIEW: On the Species Abundance Distribution in Applied Ecology and Biodiversity Management." *Journal of Applied Ecology* 52(2):443–54. doi: 10.1111/1365-2664.12380.

Merkes, Christopher M., S. Grace McCalla, Nathan R. Jensen, Mark P. Gaikowski, and Jon J.
        Amberg. 2014. "Persistence of DNA in Carcasses, Slime and Avian Feces May Affect
        Interpretation of Environmental DNA Data." *PLOS ONE* 9(11):e113346. doi:
        10.1371/journal.pone.0113346.

Mizumoto, Hiroki, Hirokazu Urabe, Takashi Kanbe, Michio Fukushima, and Hitoshi Araki.
        2018. "Establishing an Environmental DNA Method to Detect and Estimate the Biomass
        of Sakhalin Taimen, a Critically Endangered Asian Salmonid." *Limnology* 19(2):219–27.
        doi: 10.1007/s10201-017-0535-x.

Port, Jesse A., James L. O'Donnell, Ofelia C. Romero-Maraccini, Paul R. Leary, Steven Y.
        Litvin, Kerry J. Nickols, Kevan M. Yamahara, and Ryan P. Kelly. 2016. "Assessing
        Vertebrate Biodiversity in a Kelp Forest Ecosystem Using Environmental DNA."
        Molecular Ecology 25(2):527–41. doi: 10.1111/mec.13481.

Poté, John, Rafaël Ackermann, and Walter Wildi. 2009. "Plant Leaf Mass Loss and DNA
        Release in Freshwater Sediments." *Ecotoxicology and Environmental Safety* 72(5):1378–
        83. doi: 10.1016/j.ecoenv.2009.04.010.

Robson, Heather L. A., Tansyn H. Noble, Richard J. Saunders, Simon K. A. Robson, Damien W.
        Burrows, and Dean R. Jerry. 2016. "Fine-Tuning for the Tropics: Application of eDNA
        Technology for Invasive Fish Detection in Tropical Freshwater Ecosystems." *Molecular
        Ecology Resources* 16(4):922–32. doi: 10.1111/1755-0998.12505.

Sakata, Masayuki K., Nobutaka Maki, Hideki Sugiyama, and Toshifumi Minamoto. 2017.
        "Identifying a Breeding Habitat of a Critically Endangered Fish, Acheilognathus Typus,
        in a Natural River in Japan." *The Science of Nature* 104(11):100. doi: 10.1007/s00114-
        017-1521-1.

Schmitt, Michael W., Scott R. Kennedy, Jesse J. Salk, Edward J. Fox, Joseph B. Hiatt, and
        Lawrence A. Loeb. 2012. "Detection of Ultra-Rare Mutations by next-Generation
        Sequencing." *Proceedings of the National Academy of Sciences* 109(36):14508–13. doi:
        10.1073/pnas.1208715109.

Sethi, Suresh A., Wesley Larson, Keith Turnquist, and Dan Isermann. 2019. "Estimating the
        Number of Contributors to DNA Mixtures Provides a Novel Tool for Ecology." *Methods
        in Ecology and Evolution* 10(1):109–19. doi: 10.1111/2041-210X.13079.

Shinzato, Chuya, Yuna Zayasu, Miyuki Kanda, Mayumi Kawamitsu, Noriyuki Satoh, Hiroshi
        Yamashita, and Go Suzuki. 2018. "Using Seawater to Document Coral-Zoothanthella
        Diversity: A New Approach to Coral Reef Monitoring Using Environmental DNA."
        Frontiers in Marine Science 5.

Sigsgaard, Eva Egelyng, Henrik Carl, Peter Rask Møller, and Philip Francis Thomsen. 2015.
        "Monitoring the Near-Extinct European Weather Loach in Denmark Based on
        Environmental DNA from Water Samples." *Biological Conservation* 183:46–52. doi:
        10.1016/j.biocon.2014.11.023.

Stamatakis, Alexandros. 2014. "RAxML Version 8: A Tool for Phylogenetic Analysis and Post-
        Analysis of Large Phylogenies." *Bioinformatics* 30(9):1312–13. doi:
        10.1093/bioinformatics/btu033.

Strickler, Katherine M., Alexander K. Fremier, and Caren S. Goldberg. 2015. "Quantifying
        Effects of UV-B, Temperature, and pH on eDNA Degradation in Aquatic Microcosms."
        Biological Conservation 183:85–92. doi: 10.1016/j.biocon.2014.11.038.

Tajima, F. 1989. "The Effect of Change in Population Size on DNA Polymorphism." *Genetics*
        123(3):597–601. doi: 10.1093/genetics/123.3.597.

Takahashi, Takuya, and Fumio Tajima. 2017. "The Amount of DNA Polymorphism When
Population Size Changes Linearly." *Genes & Genetic Systems* 92(1):55–57. doi:
10.1266/ggs.16-00055.
Turner, Cameron R., Karen L. Uy, and Robert C. Everhart. 2015. "Fish Environmental DNA Is
More Concentrated in Aquatic Sediments than Surface Water." *Biological Conservation*
183:93–102. doi: 10.1016/j.biocon.2014.11.017.
Uchii, Kimiko, Hideyuki Doi, and Toshifumi Minamoto. 2016. "A Novel Environmental DNA
Approach to Quantify the Cryptic Invasion of Non-Native Genotypes." *Molecular
Ecology Resources* 16(2):415–22. doi: 10.1111/1755-0998.12460.
Van Holm, Wannes, Justien Ghesquière, Nico Boon, Tim Verspecht, Kristel Bernaerts, Naiera
Zayed, Ioanna Chatzigiannidou, and Wim Teughels. 2021. "A Viability Quantitative PCR
Dilemma: Are Longer Amplicons Better?" *Applied and Environmental Microbiology*
87(5):e02653-20. doi: 10.1128/AEM.02653-20.
Wilcox, Taylor M., Katherine E. Zarn, Maxine P. Piggott, Michael K. Young, Kevin S.
McKelvey, and Michael K. Schwartz. 2018. "Capture Enrichment of Aquatic
Environmental DNA: A First Proof of Concept." *Molecular Ecology Resources*
18(6):1392–1401. doi: 10.1111/1755-0998.12928.
Yates, Matthew C., Dylan J. Fraser, and Alison M. Derry. 2019. "Meta-Analysis Supports
Further Refinement of eDNA for Monitoring Aquatic Species-Specific Abundance in
Nature." *Environmental DNA* 1(1):5–13. doi: 10.1002/edn3.7.
Yoshitake, Kazutoshi, Atushi Fujiwara, Aiko Matsuura, Masashi Sekino, Motoshige Yasuike,
Yoji Nakamura, Reiichiro Nakamichi, Masaaki Kodama, Yumiko Takahama, Akinori
Takasuka, Shuichi Asakawa, Kazuomi Nishikiori, Takanori Kobayashi, and Shugo
Watabe. 2021. "Estimation of Tuna Population by the Improved Analytical Pipeline of
Unique Molecular Identifier-Assisted HaCeD-Seq (Haplotype Count from eDNA)."
Scientific Reports 11(1):7031. doi: 10.1038/s41598-021-86190-6.
Yoshitake, Kazutoshi, Tatsuki Yoshinaga, Chikaya Tanaka, Nanami Mizusawa, Md. Shaheed
Reza, Atsumi Tsujimoto, Takanori Kobayashi, and Shugo Watabe. 2019. "HaCeD-Seq:
A Novel Method for Reliable and Easy Estimation About the Fish Population Using
Haplotype Count from eDNA." *Marine Biotechnology* 21(6):813–20. doi:
10.1007/s10126-019-09926-6.
Yusishen, Michael E., Frances-Claire Eichorn, W. Gary Anderson, and Margaret F. Docker.
2020. "Development of Quantitative PCR Assays for the Detection and Quantification of
Lake Sturgeon (Acipenser Fulvescens) Environmental DNA." *Conservation Genetics
Resources* 12(1):17–19. doi: 10.1007/s12686-018-1054-8.
683

TABLES AND FIGURES

Fig. 1. Three main components of this study. (A) In silico study. We tested the correlation

between the number of segregating sites and the species abundance using simulated data. (B) In

vitro study. We designed markers and synthesized baits to capture target genes from tissues of *A.*

hasta and *T. bifasciatus* individuals. Datasets mixed from captured sequences were used to

assess the correlation between the number of segregating sites and the species abundance. (C) In

situ (mesocosm) study. We conducted controlled mesocosm experiments using *A. hasta*.

Segregating sites were extracted from captured sequences in eDNA samples, and used to

evaluate their correlation to species abundance and their performance in estimating species

abundance.

Fig. 2. Relationship between the number of segregating sites and the number of simulated

individuals/sequences from in-silico analysis. Sequences of 1,000 individuals, each with a length

of 17,000 bp, were simulated with a mutation rate of 10^{-6} /bp/gen (A), 10^{-7} /bp/gen (B) and 10^{-8}

/bp/gen (C) respectively. For each mutation rate, 20 to 980 samples were randomly drawn from

the population, and the number of segregating sites was counted. The sampling process was

repeated three times for each number of individuals. Each dot in the plots represents one random

sampling. The fitted line represents the relationship between the number of segregating sites and

the species abundance.

Fig. 3. Relationship between the number of individuals and the number of segregating sites

observed in the assembled sequences of *A. hasta* and *T. bifasciatus*. For *A. hasta*, 5 to 90

individuals were randomly sampled from the population, and the number of segregating sites was

counted (A). Similarly, for *T. bifasciatus*, 5 to 70 were randomly sampled from the population,

then the number of segregating sites was counted (B). The sampling process was repeated three

times for each number of individuals. Each dot in the plots represents one random sampling.

Fitted lines represent the relationship between the number of individuals and the species

abundance.

Fig. 4. Relationship between the number of individuals and the number of segregating sites in

eDNA samples (A), or between the number of individuals and the eDNA concentration (B). The

eDNA concentration was estimated based on the number of DNA copies at locus T2 using

qPCR. The colors of the dots depict the weight range of average weight for individuals in each

aquarium: blue (6-11 g), green (12-15 g), yellow (16-24 g), and red (38-49 g). The solid dots

indicate that the individuals in these aquariums were fed, and the circles indicate that the

individuals were not fed. The number of individuals stocked in each tank ranged from 7,14, 21,

28 to 35. Four replicates were set up at each stocking density level, with different average

weights and feeding treatments.

Fig. 5. Cross-validation test results for predicting species abundance. Boxplots show (A)

absolute mean bias and (B) standard deviation of predicted species abundance based on

segregating sites (blue) and eDNA concentration (orange) with increasing size of testing sets (top

to bottom, 1 to 10 samples). Lower bias and standard deviation indicate better model accuracy

and precision.

729   **Table 1.** Design of the mesocomic experiments.

Feeding\No. fish	7[a]	14	21	28	35
F[b]	$W_1(6)$[c]	$W_2(12)$	$W_3(18)$	$W_4(39)$	$W_1(11)$
F_0	$W_2(13)$	$W_1(7)$	$W_4(38)$	$W_3(24)$	$W_2(15)$
F	$W_3(19)$	$W_4(44)$	$W_1(7)$	$W_2(13)$	$W_3(20)$
F_0	$W_4(49)$	$W_3(16)$	$W_2(14)$	$W_1(8)$	$W_4(42)$

[a]The number (7, 14, 21, 28 and 35) indicates the number of fish kept in each aquarium.

[b]F means the fish were fed during the experiment. F_0 means the fish were not fed.

[c]The number in parentheses is the average weight of fish in this aquarium. The number in

subscript denotes the rank of the average weight of fish in the treatment group. Smaller rank in

subscript means lower average fish weight in this aquarium.

Table 2. Comparison of linear models for predicting the number of segregating sites or eDNA

concentration using fish number, mean weight and feeding behavior.

Variable	Segregating sites		eDNA concentration	
	Model 1	Model 2	Model 1	Model 2
Fish number (Coef.)	4.09**	4.08**	3.08**	3.06**
Mean weight (Coef.)		2.54		-25.95
Feeding (Coef.)		0.63		0.66
R^2	0.70	0.73	0.46	0.60
AIC	191.6	193.5	200.7	198.9
BIC	193.6	197.5	202.7	202.9
F-test (p-value)[a]		0.38		0.07

737 * $P < 0.05$; ** $P < 0.01$.

738 [a] Comparing model 1 vs. model 2 using F-test statistics.