1
2
3
Relaxin Modulates the Genomic Actions and Biological Effects of Estrogen in the Myometrium

6
7
8
<u>Short Title</u>:
9
Effects of Relaxin on Estrogen Signaling in the Myometrium
10
11

Sudeshna Tripathy[1, 2, 3,], Anusha Nagari[1, 4], Shu-Ping Chiu[1], Tulip Nandu[1, 4],
Cristel V. Camacho[1, 3], Mala Mahendroo[2, 3, 5], and W. Lee Kraus[1, 3, 5, 6]

14
[1] Laboratory of Signaling and Gene Regulation, Cecil H. and Ida Green Center for Reproductive Biology Sciences, University of Texas Southwestern Medical Center, Dallas, TX 75390, USA.

[2] Laboratory of Cervical Remodeling and Preterm Birth, Cecil H. and Ida Green Center for Reproductive Biology Sciences, University of Texas Southwestern Medical Center, Dallas, TX 75390, USA.

[3] Section of Laboratory Research, Department of Obstetrics and Gynecology, University of Texas Southwestern Medical Center, Dallas, TX 75390, USA.

[4] Computational Core Facility, Cecil H. and Ida Green Center for Reproductive Biology Sciences, University of Texas Southwestern Medical Center, Dallas, TX 75390, USA.

[5] Address correspondence to: W. Lee Kraus (Lee.Kraus@utsouthwestern.edu) and Mala Mahendroo (Mala.Mahendroo@utsouthwestern.edu).

27
28
29
DISCLOSURE SUMMARY
30
The authors having no conflicts to disclose.
31
32

33
[6] <u>Lead Contact</u>:
34
W. Lee Kraus, Ph.D.
35
Cecil H. and Ida Green Center for Reproductive Biology Sciences
36
The University of Texas Southwestern Medical Center at Dallas
37
5323 Harry Hines Boulevard
38
Dallas, TX 75390-8511
39
40
Phone: 214-648-2388
41
Fax: 214-648-0383
42
E-mail: Lee.Kraus@utsouthwestern.edu
43
44
https://orcid.org/: 0000-0002-8786-2986

ABSTRACT

Estradiol (E2) and relaxin (Rln) are steroid and polypeptide hormones, respectively, with

important roles in the female reproductive tract, including myometrium. Some actions of Rln,

which are mediated by its membrane receptor RXFP1, require or are augmented by E2 signaling

through its cognate nuclear steroid receptor, estrogen receptor alpha (ERα). In contrast, other

actions of Rln act in opposition to the effects of E2. Here we explored the molecular and

genomic mechanisms that underlie the functional interplay between E2 and Rln in the

myometrium. We used both ovariectomized female mice and immortalized human myometrial

cells expressing wild-type or mutant ERα (hTERT-HM-ERα cells). Our results indicate that Rln

modulates the genomic actions and biological effects of estrogen in the myometrium and

myometrial cells by reducing phosphorylation of ERα on serine 118 (S118), as well as by

reducing the E2-dependent binding of ERα across the genome. These effects were associated

with changes in the hormone-regulated transcriptome, including a decrease in the E2-dependent

expression of some genes and enhanced expression of others. The inhibitory effects of Rln

cotreatment on the E2-dependent phosphorylation of ERα required the nuclear dual-specificity

phosphatases DUSP1 and DUSP5. Moreover, the inhibitory effects of Rln were reflected in a

concomitant inhibition of the E2-dependent contraction of myometrial cells. Collectively, our

results identify a pathway that integrates Rln/RXFP1 and E2/ERα signaling, resulting in a

convergence of membrane and nuclear signaling pathways to control genomic and biological

outcomes.

KEYWORDS

Estrogen, estradiol, relaxin, dual-specificity phosphatase, myometrium, estrogen receptor,

relaxin receptor, contraction.

ABBREVIATIONS

E2, estradiol; Rln, relaxin; RXFP1, relaxin family peptide receptor 1 (relaxin receptor);

DUSP, dual-specificity phosphatase; ChIP, chromatin immunoprecipitation; RT-qPCR, reverse

transcription- quantitative PCR.

INTRODUCTION

The uterus must provide and maintain an environment that is conducive to the growth and

development of the fetus throughout pregnancy, yet also gradually prepare for parturition, the

process of labor and delivery of the fetus at term. Uterine myometrial smooth muscle cell

(SMC) plasticity ensures transition between the non-contractile and contractile states in response

to physiological signaling cues, at term (1,2). Misregulation of myometrium function can result

in numerous obstetrical complications, including preterm birth or Cesarean sections (3,4).

Maintenance of myometrial quiescence and the transition to contractility is regulated

through hormones such as progesterone (P4), estradiol (E2), and relaxin (Rln) (5,6). As the

actions of P4 decline in late pregnancy, the actions and levels of E2 are enhanced (7,8). The

effects of E2 are mediated through nuclear estrogen receptors (ERs) (9,10). ERα is generally

expressed at higher levels than ERβ and plays a dominant role in the biology of the female

reproductive tract (9,11). The effects of E2 in the myometrium include increased responsiveness

to contraction-promoting molecules (e.g., oxytocin and prostaglandins), increased gap junction

formation, decrease nitric oxide activity, and increased Ca^{+2} influx (12-14). The peptide

hormone Rln has numerous roles in the physiology of pregnancy in humans and other species

(15-17). Rln, which is secreted from the corpus luteum and the placenta, mediates its actions

through a G-protein coupled receptor, relaxin family peptide receptor 1 (RXFP1) (16). Although

Rln is best known for its actions in the cervix (18,19), numerous studies have uncovered diverse

roles for Rln in the myometrium (16,17,20,21).

In the uterus, Rln induces uterine growth and development, inhibits apoptosis, promotes

smooth muscle cell relaxation via cAMP induction, and promotes vasodilation of the uterine

arteries (15,17,22). Pharmacological doses of Rln diminish myometrial contraction induced by

either oxytocin or prostaglandins (15,23). A critical role of Rln signaling during parturition is

supported by incomplete cervical remodeling and parturition defects in mice with targeted

mutation in the *Rln1* (Rln) or *Rxfp1* (Rln receptor) genes (24,25). Interestingly, many of the

effects of Rln on reproductive tissues, such as cell proliferation, ECM remodeling, and

angiogenic actions, require or are augmented by E2/ERα signaling (16,26,27). Conversely, in

some cases, the two hormones have opposing actions. For example, Rln promotes smooth

muscle quiescence in the myometrium, while E2/ERα promotes contraction (2,28-30). The

molecular mechanisms by which Rln/RXFP1 signaling can promote or oppose the actions of

E2/ERα have not been defined.

Many of the biological actions of estrogens are mediated by ERα, which functions

primarily as a ligand-regulated, DNA-binding transcription factor (TF) in the nuclei of estrogen-

responsive cells. Nuclear ERα dimerizes in response to E2 and binds to many thousands of ERα

binding sites (ERBSs) across the genome, collectively called the ERα 'cistrome' (31-35). Most

ERα binding sites are pre-bound by "pioneer" transcription factors (e.g., FoxA1, AP-2γ) prior to

estrogen exposure, which facilitate the binding of ERα to a repertoire of predetermined sites

(36,37). Many ERα binding sites contain a DNA sequence motif called the estrogen response

element (ERE) (10).   The binding of ERα to genomic DNA promotes the coordinated

recruitment of coregulator proteins that establish an active 'enhancer,' leading to chromatin

looping and target gene transcription (38-42).

Importantly, ERα actions can be modulated by post-translational modifications (PTMs).

For example, ERα can be phosphorylated on multiple sites by a wide range of kinases, which

regulate various functions such as hormone sensitivity, nuclear localization, DNA binding,

protein and chromatin interactions, and gene transcription (43,44). Phosphorylation at serine 118

has been reported to be important for ERα dimerization, interactions with coactivators such as

CBP/p300 and SRC, receptor activation, and ERα enhancer formation (45-47). Although serine

118 phosphorylation may promote the progression of uterine leiomyomas, its effects on

myometrium function have not been characterized (48,49).

In the studies described herein, we examine the extent and nature of functional

interactions between the E2 and Rln signaling pathways in the myometrium. In addition, we

delineate the mechanisms underlying functional interplay between the E2/ERα and Rln/RXFP1

signaling pathways at the molecular and physiological levels in myometrial tissue and cells

during quiescence and contractility.

METHODS AND MATERIALS

Antibodies, hormones, and inhibitors

The antibodies used were as follows: ERα, rabbit polyclonal generated in house (50);

phospho-ERα (S118), mouse monoclonal (Cell Signaling Technology, 2511S; RRID:

AB_331289); phospho-ERα (S104/106) (Abcam, ab75753; RRID: AB_1310197); phospho-ERα

(S167), rabbit monoclonal (Cell Signaling Technology, 64508; RRID: AB_2799660); H3K27ac,

rabbit polyclonal (Abcam, ab4729; RRID: AB_2118291); H3K4me1, rabbit polyclonal (Abcam,

ab8895; RRID: AB_306847); ERK1/2 (Cell Signaling Technology, 4695; RRID: AB_390779);

phospho-ERK1/2 (Cell Signaling Technology, 4370; RRID: AB_2315112); DUSP1, rabbit

polyclonal (Cell Signaling Technology, 35217; RRID: AB_3371713); DUSP5, rabbit polyclonal

(Thermo Fisher Scientific, PA5-85961; RRID: AB_2802762); β-actin, mouse monoclonal (Cell

Signaling Technology, 3700; RRID: AB_2242334); goat anti-rabbit IgG (H+L)-HRP conjugate

(BioRad, 1706515; RRID: AB_11125142); and goat anti-mouse IgG (H+L)-HRP conjugate

(BioRad, 1706516; RRID: AB_2921252). The hormones used were as follows: 17β-estradiol

(E2; Sigma) and recombinant human relaxin-2 (Rln; Peprotech). The MEK inhibitor Trametinib

(GSK1120212) (51) was from MedChemExpress (HY-10999).

Mouse models, ovariectomy, and hormone administration

All animal studies were conducted in accordance with the standards of humane animal

care as described in the NIH Guide for the Care and Use of Laboratory Animals. The research

protocols were approved by the IACUC office at the University of Texas Southwestern Medical

Center. The mouse colony was maintained in a barrier facility and the health status of the mice

was assessed on a quarterly basis. We used the following mouse strains: Wild-type ($Rxfp1^{+/+}$)

and relaxin receptor knockout ($Rxfp1^{-/-}$) (52) on a C57BL/6 background housed under a 12 h-

light/12 h-dark cycle at 22°C. Female mice at 7 to 8 weeks of age were ovariectomized and

rested for 2 weeks to deplete endogenous ovarian hormones. For experiments, the

ovariectomized mice were injected i.p. with 0.1 mL of saline (vehicle), 2.5 μg/mL 17β-estradiol

(E2, E; Sigma), 80 μg/mL 0.1% BSA-PBS recombinant human relaxin-2 (Rln, R; Peprotech) and

a combination of E+R for the specified amount of time (five mice per treatment per biological

replicate). Mice were sacrificed under 2,2,2-tribromomethanol (Sigma) anesthesia. The uteri

were harvested and further enriched for myometrium by sterile scraping, washing in cold 1x PBS

and blotting with a paper towel.   The enriched myometrial tissues were flash frozen for

subsequent genomic and biochemical assays.

Cell culture and treatments

Telomerase-immortalized human myometrial (hTERT-HM) cells (53) were cultured in

phenol-red free DMEM/F12 (ThermoFisher Scientific) with 10% fetal bovine serum (FBS, v/v),

1% penicillin/streptomycin at 37°C in 5% CO_2. hTERT-HM cells with doxycycline (Dox)-

inducible expression of ERα (wild-type, WT) and ERα mutants (S118A and S118E) were

generated by lentivirus-mediated transduction using the pINDUCER20 vector (RRID:

Addgene_44012), as described below.

For experiments, the cells were treated with vehicle (DMSO), 100 nM 17β-estradiol (E2,

E; Sigma, St. Louis, MO, USA), 0.2 μg/mL recombinant human relaxin-2 in 0.1% BSA-PBS

(Rln, R; Peprotech, Rocky Hill, NJ, USA) and a combination of E+R for the specified amount of

time. Prior to treatment, the media was supplemented with 5% charcoal-dextran-treated calf

serum (CDCS) for 3 days. The cells were treated with doxycycline for 24 hours prior to the

different treatments.

Generation of recombinant ERα cDNAs and expression constructs

Plasmids containing gene of interest, ERα (WT, S118A and S118E), were used as a

template for PCR amplification using following primers:

• S118A Forward: 5'-GCAGGAAAGGCGCCAGCTGCGGCGG-3'

• S118A Reverse: 5'-CCGCCGCAGCTGGCGCCTTTCCTGC-3'

• S118E Forward: 5'-CTGCAGGAAAGGCTCCAGCTGCGGCGGC-3'

• S118E Reverse: 5'GCCGCCGCAGCTGGAGCCTTCCTGCAG-3'

PCR-amplified fragments were cloned into pINDUCER20 using a ligation-independent

cloning protocol with 5x In-Fusion HD Enzyme Premix (Takara Bio USA, Inc.) per

manufacturer's protocol. The recombinant constructs were confirmed by DNA sequencing.

Generation of hTERT-HM cells stably expressing ERα

hTERT-HM cells with Dox-inducible expression of ERα (WT, S118A and S118E) were

generated by lentivirus-mediated transduction. The lentiviral ERα expression constructs or the

empty vector were co-transfected with additional virus constructs (pRSV-Rev, pMDLg, pCMV-

VSV-G) into HEK-293 FT cells using Polyjet reagent (SignaGen Laboratories). The medium

containing the transfection reagent was replaced with fresh medium (DMEM) 7 hours post-

transfection. The lentivirus containing supernatant was collected and filtered for transduction.

hTERT-HM cells were infected with recombinant lentivirus at a multiplicity of infection (MOI)

or approximately 1. One day after infection, the cells were allowed to recover in fresh medium

for 24 hours and were then subjected to drug selection using G418. The expression of ERα

under Dox induction was confirmed by Western blotting.

Knockdown of *DUSP1* and *DUSP5* mRNAs using siRNAs

Commercially available siRNAs targeting *DUSP1* mRNA (Sigma,

SASI_Hs01_00098748) and *DUSP5* mRNA (Sigma, SASI_Hs02_00337568) were transfected

into hTERT-HM- ERα cells at a final concentration of 30 pmol using Lipofectamine RNAi

MAX reagent (Invitrogen, 13778-150) according to the manufacturer's instructions. All

experiments were performed 72 hours after siRNA transfection.

Western blotting

For determination of protein levels and phosphorylation by Western blotting, extracts

were prepared from mouse myometrial tissue or cultured hTERT-HM-ERα cells from a

minimum of three biological replicates. Samples were collected for each of the four treatment

time points (vehicle, E2, Rln, and E+R) from ovariectomized mice or hTERT-HM cells (1 hour

and 30 min treatment times, respectively). Frozen myometrial tissue was powdered using a

metal pulverizer. Whole cell lysates of the powdered myometrial tissues and hTERT-HM-ERα

grown on 10 cm diameter plates were prepared in ice cold Whole Cell Extract Buffer (10 mM

Tris-HCl pH 7.5, 0.5 M NaCl, 2.5 mM EDTA, 1% Nonidet P-40), supplemented with Halt

Protease and Phosphatase Inhibitors (Thermo Scientific). The cells were resuspended by

pipetting, lysed, and incubated on ice for 30 min. The lysates were centrifuged at maximum

speed in a microcentrifuge for 10 min at 4°C and the supernatants were collected as the whole

cell extracts. The extracts were subjected to colorimetric bicinchoninic acid (BCA) protein assay

(Pierce). Fifteen to 20 μg of total protein were analyzed on 8% polyacrylamide-SDS gel and

transferred to a nitrocellulose membrane. Membranes were blocked at room temperature for 1

hour in 5% nonfat dry milk (blocking grade, Bio-Rad) dissolved in Tris-buffered saline

containing 0.1% Tween 20 (TBST). The blots were incubated with primary antibody (listed

above) in blocking solution overnight at 4°C, washed in TBST, and incubated with horseradish

peroxidase (HRP)-labeled secondary antibodies for 1 hour at room temperature. The membrane

was imaged with a Bio-Rad ChemiDoc MP Imaging System (BioRad).

Isolation of total RNA

Total RNA was isolated from myometrial tissue or cultured hTERT-HM-ERα cells using

an RNeasy Plus Mini Kit (Qiagen), according to manufacturer's instructions. Myometrial tissue

was homogenized in RLT buffer with a cooled blade homogenizer (Power Gen25; Fisher).

hTERT-HM-ERα cells were collected and lysed in RLT buffer. The samples were centrifuged

to remove debris and passed through a gDNA eliminator column, followed by RNA binding

column. After multiple washes, total RNA was eluted with RNase-free water. The eluted RNA

was assayed for quality by electrophoresis (Agilent Technologies, Santa Clara, CA, USA). Only

samples with RIN values greater than 8 were included in subsequent analyses.

Reverse transcription-quantitative PCR (RT-qPCR)

Analysis of the levels of specific mRNAs were assessed by RT-qPCR. Total RNA was

reverse transcribed using MMLV reverse transcriptase (Promega, M150B) with oligo(dT)

primers (Sigma-Aldrich). The resulting cDNA was analyzed by qPCR using the primer sets

listed below using a LightCycler 480 real-time PCR thermocycler (Roche):

• *DUSP1* Forward: 5'-ACCACCACCGTGTTCAACTTC-3'

• *DUSP1* Reverse: 5'-TGGGAGAGGTCGTAATGGGG-3'

• *DUSP5* Forward: 5'-TGTCGTCCTCACCTCGCTA-3'

• *DUSP5* Reverse: 5'-GGGCTCTCTCACTCTCAATCTTC-3'

RNA-seq library preparation and sequencing

RNA-seq libraries from mouse myometrial tissue and hTERT-HM-ERα cells (WT and

S118E) were prepared as follows using a protocol described previously (54). Briefly, biological

replicates were generated by pooling 1.5 μg of total RNA from five animals for a total of 7.5 μg

of total RNA per replicate. A total of ten animals were used for each treatment time point.

Polyadenylated RNA was isolated from total RNA using Oligo(dT)25 Dynabeads (Invitrogen)

and reverse transcribed using SuperScript III Reverse Transcriptase (Invitrogen). Second strand

synthesis was performed using DNA polymerase I. Complementary DNA ends were repaired,

tailed with dATP, and ligated with barcode adapters for Illumina sequencing platform. Ligated

libraries were selected for average size 270 bp, PCR amplified for 9 cycles, and sequenced on an

Illumina NextSeq 500 in a 75 bp single-end format in the McDermott Center Next Generation

Sequencing Core at UT Southwestern Medical Center to a total read count of ~40 M per

condition.

Analysis of RNA-seq data

Quality check, alignment, and differential expression. The raw data were subjected to

QC analyses using the FastQC tool (55). The reads were aligned to the mouse genome (mm10)

or the human genome (hg38) using TopHat v.2.0.12 (56). Using aligned reads as input, we

employed cufflinks v.2.2.1 (57) to assemble the reads into transcripts using Gencode v.M5

annotations. We then used cuffdiff v.2.2.1 (57) to call differentially regulated transcripts. All

programs were run with default parameters. The expression of differentially expressed genes

was visualized as heatmaps using Java Treeview (58). To categorize treatment-regulated genes,

we filtered for genes with: (1) a p-value < 0.01 and fold change cutoff ≥ 2.0 or ≤ 0.5 (log2 ≥ 1.0

or ≤ -1.0) for mouse myometrial cells and (1) a p-value < 0.01 and fold change cutoff ≥ 1.5 or ≤

0.67 (log2 ≥ 0.58 or ≤ -0.79) for hTERT-HM-ERα cells.

Gene ontology (GO) analyses. Genes associated with ERα peaks depleted by Rln

treatment were evaluated with WEB-based GEne SeT AnaLysis Toolkit (WebGestalt) (59).

Multiple gene ontology terms relating to collagen metabolism and extracellular matrix (ECM)

organization were identified. Additional gene and pathways analyses were performed using

Gene Ontology (GO) annotations (60) for GO biological process gene sets and the PANTHER

enrichment analysis tool (61).

Chromatin immunoprecipitation (ChIP) in mouse myometrial tissue

Tissue preparation and crosslinking. Pools of frozen myometrial tissue (5 myometrium

per pool; 60 to 80 mg of tissue) were powdered using a metal pulverizer. The powdered

myometrial pool was immediately resuspended in 1% formaldehyde in phosphate-buffered saline

(PBS) with a volume of 10 mL per pool, then incubated on a rotating platform at room

temperature for 10 min. Crosslinking was quenched by adding 5 mL per pool of 2.5 M glycine,

with a subsequent 5 min incubation at room temperature. The crosslinked tissue was collected

by centrifugation at 1000 x *g* for 3 min at room temperature. The supernatant was decanted, and

the tissue was resuspended in 4°C PBS with added protease inhibitor cocktail (Roche Molecular

Biochemicals) at 10 mL per pool, with incubation on ice for 2 min. The samples were

centrifuged at 1000 x *g* for 3 min at 4°C. The PBS wash was repeated, and the tissue was

resuspended in 2.5 mL per pool of homogenization buffer (50 mM Tris-HCl pH 7.5, 1% Nonidet

P-40, 0.25% deoxycholic acid, 1 mM EDTA) supplemented with proteinase inhibitor cocktail.

Preparation of soluble chromatin. The crosslinked tissue was homogenized using a

cooled blade homogenizer (Power Gen25). The pellet was collected by centrifugation at 1000 x

g for 5 min at 4°C. The supernatant was decanted and the pellet was resuspended in 1 mL per

pool of Farnham Buffer (5 mM PIPES, pH 8.0, 85 mM KCl, 0.5% Nonidet P-40, 1 mM DTT, 0.1

mM PMSF) supplemented with proteinase inhibitor cocktail by mixing five times with a pipettor

and then incubated on ice for 10 min. The chromatin was pelleted by centrifugation for 1000 x *g*

for 10 min at 4°C and resuspended in SDS Lysis Buffer (50 mM Tris-HCl pH 7.9, 10 mM

EDTA, 1% SDS, 1 mM DTT) supplemented with proteinase inhibitor cocktail and histone

deacetylases inhibitors (10 mM sodium butyrate, 5 mM nicotinamide). Resuspended chromatin

was incubated on ice for 10 min, split into 300 µL aliquots, and sonicated on ice using a

Bioruptor UCD-200 (Diagenode) on the high setting for 9 cycles, 30 sec on and 30 sec off with 1

min rest, 4 times, to obtain genomic DNA fragments of approximately 200 to 400 bp, as

determined by gel electrophoresis. Sonicated chromatin was centrifuged at maximum speed in a

microcentrifuge for 1 min at 4°C to remove insoluble debris, and the soluble chromatin

supernatant was flash frozen in liquid N_2 and stored at -80°C until use.

Immunoprecipitation (IP). Chromatin was diluted 1:9 with Dilution Buffer (20 mM

Tris-HCl pH 7.9, 300 mM NaCl, 2 mM EDTA, 0.5% Triton X-100, 1 mM DTT, 0.2 mM PMSF)

supplemented with proteinase inhibitor cocktail and histone deacetylases inhibitors. The diluted

chromatin was pre-cleared with 20 µL/mL protein A-agarose beads (ThermoFisher Scientific;

50% slurry in 10 mM Tris-HCl pH 8.1; 1 mM EDTA) on a nutator for 1 hour at 4°C. The pre-

cleared chromatin was centrifuged in a microcentrifuge at 1000 x *g* for 3 min at 4°C. The pre-

cleared supernatant was divided into input (2.5%) and a 1 mL aliquot for immunoprecipitation.

Immunoprecipitation was performed overnight at 4°C using the polyclonal antibodies for ERα,

H3K27ac, or H3K4me1 described above. For a control, rabbit IgG (BioRad; PRABP01) was

added at 1:200.

Immune complexes were collected by the addition of 80 µL of protein A-agarose beads

(Cell Signaling Technology; 9863) with an additional 2 hour incubation at 4°C with mixing. The

samples were centrifuged in a microcentrifuge at 500 x *g* at 4°C for 1 min to collect the resin.

The resin was washed four times for 10 min each in Low Salt Wash Buffer (20 mM Tris-HCl pH

7.9, 2 mM EDTA, 125 mM NaCl, 0.05% SDS, 1% Triton X-100), four times for 10 min each in

High Salt Wash Buffer (20 mM Tris-HCl pH 7.9, 2 mM EDTA, 500 mM NaCl, 0.05% SDS, 1%

Triton X-100), one time with LiCl Wash Buffer (10 mM Tris-HCl pH 7.9, 1 mM EDTA, 250

mM LiCl, 1% Nonidet P-40, 1% sodium deoxycholate), and one time in TE. All buffers were

supplemented with proteinase inhibitor cocktail and histone deacetylases inhibitors. The cross-

linked chromatin was then eluted twice using Elution Buffer (1% SDS and 100 mM sodium

bicarbonate) at room temperature for 15 min. The eluted chromatin was heated at 65°C

overnight in Elution Buffer supplemented with 100 mM NaCl to reverse the formaldehyde

crosslinking. RNase was added to 3 µg/mL final concentration was added and incubated at 37°C

for 30 min, followed by inactivation with Tris-HCl pH 7.9, 5 mM EDTA, 50 µg/mL proteinase

K at 55°C for 2 hours. The immunoprecipitated DNA fragments were purified by

phenol:chloroform extraction, followed by ethanol precipitation. The purified ChIPed DNA was

stored at -20°C until it was analyzed by quantitative real time PCR or ChIP-seq as described

below.

Chromatin immunoprecipitation reverse transcription-quantitative PCR (ChIP-qPCR)

Purified ChIPed DNA from mouse myometrial tissue was analyzed by reverse

transcription-quantitative PCR using the following primers:

• *Igf1* Forward: 5'-GACACACACACAGACCCTAAA-3'

• *Igf1* Reverse: 5'-GTTGTGCTGGAGACCAAGT-3'

• *Pgr* Forward: 5'-TTGTAAAGAGGTGGGATAATGCT-3'

• *Pgr* Reverse: 5'-GTCCAAGGGTGTGTGGTAAA-3'

• *Nrip1* Forward: 5'-GTACTCCTCACTGACACCAAATAG-3'

• *Nrip1* Reverse: 5'-CCTTGCTGAGTGTCGTTTCT-3'

ChIP-seq library preparation and sequencing

ChIP-seq libraries were generated from two biological replicates for each condition as

described previously (62,63). Immunoprecipitated (ChIPed) DNA was purified further by

polyethylene glycol (PEG)-mediated precipitation (64) using PEG Precipitation Solution (20% in

20 mM Tris-HCl pH 7.9, 4 mM EDTA, 2.5 M NaCl). After purification, 5 ng (for ERα) and 10

ng (for H3K27ac and H3K4me1) of ChIPed DNA for each treatment condition was used to

generate libraries for sequencing, as previously described with some minor modifications (63).

Briefly, the ChIPed DNA was end-repaired and a single "A"-base overhang was added using the

Klenow fragment of *E.coli* DNA polymerase. The A-modified DNA was ligated with Illumina

sequencing adapters using the Illumina TruSeq DNA Sample Prep Kit. The ligated DNA (250 ±

25 bp) was size-selected using a bead-based method with magnetic beads (AmPure XP beads,

Beckman Coulter) isolate DNA fragment sizes of interest. The size-selected DNA was amplified

by PCR (9 to 11 cycles as determined empirically) and purified using AmPure XP beads. The

final libraries were run on agarose gel, and extracted using QIAquick gel extraction kit (Qiagen).

The final libraries were subjected to QC (size, purity, adapter contamination). The quality of

libraries was assessed using a High Sensitivity D1000 screen tape on a 2200 TapeStation

(Agilent) and quantified using a Qubit dsDNA HS assay kit (ThermoFisher Scientific). The

libraries with unique adapter barcodes were multiplexed and sequenced on an Illumina HiSeq

2500 (single-end, 75 base reads) to a depth of ~40 M reads per condition.

Analysis of ChIP-seq data

Quality check, alignment, and peak calling. The raw data were subjected to QC

analyses using the FastQC tool (55). The ChIP-seq reads were trimmed using Cutadapt v.1.9.1

(65) and mapped to the mouse reference genome (mm10) using Bowtie v.1.0.0 (66). Uniquely

mappable reads were converted into bigWig files using BEDTools v.2.25 (67) for visualization

in the UCSC genome browser (68). The aligned reads were used to measure library complexity

using BEDTools (v 2.17.0) (67) and met minimum ENCODE data quality standards (69). ERα

peaks under all the treatment conditions were called versus input as control using HOMER

software (70) and an FDR cutoff 0.05.

Peak Annotation and clustering. In order to study the effects of the treatments on the

ERα cistromes identified by HOMER, the peaks identified upon a single treatment were

combined with the peaks identified upon co-treatment (E+R). The peaks that were gained,

maintained, and depleted upon co-treatment were identified as described previously (62,71). A

cut-off of 2MAD (median absolute deviation) was used for the ERα peaks to categorize them

into gained, maintained, and depleted peaks. The normalized read counts in the 2 kb window (±

1 kb) around the peak categories was visualized as boxplots using the box plot function in R.

Wilcoxon rank sum tests were performed to determine the statistical significance of all

comparisons.

Heatmaps. The read densities surrounding the gained, maintained, and lost ERα peaks

[5 kb (± 2.5 kb) for ERα and 10 kb (± 5 kb) for H3K27ac and H3K4me1] were determined using

the annotatePeaks.pl module of HOMER software (70) and visualized as heatmaps using Java

Treeview (58).

Collagen-based cell contraction assays

Collagen-based cell contraction assays were performed as described previously (53).

Briefly, hTERT-HM-ERα cells (WT, S118A and S118E), with or without siRNA-mediated

knockdown of *DUSP1* or *DUSP5*, were plated in collagen gel. Subconfluent cells were collected

with trypsin, centrifuged at 1,000 x *g* for 5 min, and resuspended in serum-free DMEM/F12

medium. The cells were treated with Dox for 24 hours prior to setting up the assay to induce the

expression of ERα. Cells without Dox treatment served as control. The number of cells was

quantified, and their viability was ascertained by staining with trypan blue. A solution of type I

rat tail collagen (Cultrex rat collagen I, 3440-100-01, R and D Systems) was adjusted to pH 7.2

with 0.1% glacial acetic acid and 1 M NaOH and a final concentration of 3 mg/mL. An amount

of hTERT-HM-ERα cells (WT, S118A and S118E) was added to the neutralized collagen

solution to achieve 1×10^5 cells per well in a 12-well plate, mixed, and incubated on ice for 5

min. One half mL of collagen gel-cell suspension was plated per well in a 12-well culture plate,

with subsequent incubation for 1 hour at 37°C to allow gelling. After incubation, 1 mL of fresh

DMEM/F12 medium containing DMSO (vehicle), 100 nM E2, 0.2 μg/mL Rln, or a combination

of E+R was added to the wells. After 20 hours, the gel matrices were gently detached from the

sides and lifted off the bottom of the well. The area of each gel matrix was measured

periodically for up to 48 hr. Images of the floating gels were captured using Bio-Rad ChemiDoc

MP Imaging System (Bio-Rad). The area of each gel was determined using in ImageJ (v.1.52k)

(72). The percentage of gel contraction per well was calculated as:

$$\left(\frac{\text{Area of gel at start of assay} - \text{Area of gel at end of assay}}{\text{Area of the gel at the start of assay}} \right) \text{ x } 100$$

Data availability

The new RNA-seq and ChIP-seq data sets generated for this study are available from the

NCBI's Gene Expression Omnibus (GEO) database (http://www.ncbi.nlm.nih.gov/geo/) using

accession number GSE244843.

RESULTS

Rln signaling inhibits the E2-induced genomic binding of ERα in mouse myometrium

To better understand the genomic and molecular mechanisms underlying interplay

between the E2/ERα and Rln/RXFP1 signaling pathways in the myometrium, we used 7- to 8-

week old ovariectomized female C57BL/6 mice injected i.p. with 0.1 mL of vehicle (V; saline),

E2 (E; 2.5 µg/mL), recombinant human relaxin-2 (R, 80 µg/mL) or both (E+R) for 1 or 2 hours

[for chromatin immunoprecipitation-sequencing (ChIP-seq) or RNA-sequencing (RNA-seq),

respectively] (Fig. 1A). We then collected the myometrium and subjected the tissue to Western

blotting for ERα and various genomic analyses. As shown in Fig. 1B, the treatments had no

discernable effect on the levels of nuclear ERα by Western blotting.

To explore the effects of Rln signaling on the genomic actions of E2 in the mouse

myometrium, we determined the ERα cistrome by ChIP-seq in myometrium isolated from mice

subjected to the four treatments listed above. With E2 treatment, we observed a robust induction

of ERα binding across the genome as indicated by a large increase in the number of statistically

significant ERα binding sites, as expected (4,983 sites; Fig. 1C), as determined by peak calls in

ChIP versus input from HOMER (70). Remarkably, we observed a dramatic reduction of E2-

induced ERα binding across the genome with Rln cotreatment as indicated by a dramatic

reduction in the number of statistically significant E2-induced ERα binding sites (880 sites; Fig.

1C), with 775 sites overlapping those observed in the E2 only condition.

To quantify in more detail the effects of Rln treatment on E2-induced ERα binding, we

used read counts and the universe of ERα binding sites to determine median absolute deviation

(MAD) using methods described by Rao et al. (71), which allowed us to determine "gained,"

"maintained," and "depleted" ERα binding sites. Again, we observed a dramatic reduction in

E2-induced ERα binding sites upon cotreatment with Rln, as shown in a heatmap (Fig. 1D), box

plots (Fig. 1E), and metaplots (Supplemental Fig. S1). We also examined the enrichment of two

histone modifications associated with enhancer activity, histone H3 lysine 4 monomethylation

(H3K4me1) and histone H3 lysine 27 acetylation (H3K27ac) (73,74), at the ERα binding sites.

We observed that in spite of the significant reduction in ERα binding at the "depleted" sites upon

cotreatment with Rln, significant enrichment of H3K27ac and H3K4me1 was maintained (Figs.

1, D and E; Supplemental Fig. S2). Together, these analyses indicate that Rln signaling broadly

inhibits the E2-induced binding of ERα in mouse myometrium from ovariectomized animals.

But, the maintenance of H3K4me1 and H3K27ac enrichment indicates that these enhancers may

remain open, accessible, and active.

Rln signaling alters the E2-regulated transcriptome in mouse myometrium

457   To relate the changes in the E2-dependent ERα cistrome in response to Rln signaling to

gene expression outcomes in the mouse myometrium, we first performed RNA-seq under the

conditions described above. We expressed the results in a heatmap, Venn diagram, and tabular

representation to illustrate differences in gene expression (Fig. 2, A-C).   These results

demonstrate that E2 is a potent regulator of gene expression in the myometrium, with >1,000

genes upregulated and >1,000 genes downregulated, as expected (Fig. 2C). Interestingly, Rln

also regulated the expression of many genes (>400 upregulated and >500 downregulated) (Fig.

2C). Alterations in the E2-regulated transcriptome by cotreatment with Rln were also evident;

372 E2-regulated genes lost significant regulation and 673 E2-regulated genes gained significant

regulation upon cotreatment with Rln (Fig. 2, A and B). The response of these gene sets in

aggregate to the various treatments are evident in quantitative box plots (Fig. 2D).

468   Gene ontology (GO) analysis revealed an enrichment of terms related to signaling and

cell biology for the 372 E2-regulated genes that lost significant regulation upon cotreatment with

Rln (Supplemental Fig. S3A). In contrast, the GO analysis revealed an enrichment of terms

related to transcription and translation for the 673 E2-regulated genes that gained significant

regulation upon cotreatment with Rln (Supplemental Fig. S3B). Together, these results indicate

that cotreatment with Rln can alter the E2-regulated transcriptome in mouse myometrium – both

by inhibiting the expression of some E2-regulated genes (e.g., *Aoc1* and *Gcat*; Fig. 3A,

Supplemental Fig. S4A) and by expanding the transcriptome through increased expression of

genes not regulated by E2 or Rln alone (e.g., *Sik1* and *Hapln4*; Fig. 3B, Supplemental Fig. S4B).

These changes in the E2-regulated transcriptome lead to changes in downstream biological

processes, as illustrated by the GO analysis (Supplemental Fig. S3).

Linking the effects of Rln signaling on the ERα cistrome and the E2-regulated

transcriptome in mouse myometrium

To better understand expected links between the effects of Rln signaling on the ERα

cistrome and the E2-regulated transcriptome in mouse myometrium, we integrated the ChIP-seq

data with the RNA-seq data to link ERα peaks depleted by Rln cotreatment with the expression

of the nearest neighboring genes (Fig. 4A). We collected Rln-depleted ERα binding sites

located within 50 kb of the nearest neighboring protein-coding gene promoter from the ChIP-seq

data and then determined the expression of the neighboring genes using the RNA-seq data. For

these analyses, we focused on genes upregulated by E2 and downregulated by cotreatment with

Rln with a fold change (FC) difference ($\log_2$FC_E2 – $\log_2$FC_E+R) > 0.5. We observed a

significant reduction in the E2-dependent expression of the nearest neighboring genes in

response to Rln cotreatment (Fig. 4B). We then determined the ontologies for the genes

associated with the Rln-depleted ERα peaks. We identified multiple GO terms related to

collagen metabolism and extracellular matrix (ECM) organization (Fig. 4C; Supplemental Fig.

S5A). We also identified pathways related to Rln signaling, as well as signaling via the ECM

(Supplemental Fig. S5B).

Collectively, the ChIP-seq and RNA-seq results demonstrate that the Rln-dependent

changes that we observed in the E2-dependent ERα cistrome are reflected in the gene expression

outcomes and are connected to biological outcomes related to Rln signaling. This is evident for

genes encoding amine oxidase copper containing 1 (*Aoc1*) and glycine C-acetyltransferase

(*Gcat*), both of which exhibit reduced E2-dependent ERα binding at a nearby enhancer, as well

reduced expression, upon cotreatment with Rln (Fig. 3A, Supplemental Fig. S4A). However,

Rln-mediated modulation of the E2-dependent ERα cistrome (depletion of >5,000 ERα binding

sites; Fig. 1D) was more extensive than Rln-mediated modulation of the E2-dependent

transcriptome (reduced expression of ~370 genes; Fig. 2, A and B). In this regard, classical

E2/ERα-regulated genes, such as *Pgr*, *Nrip1*, and *Igf1* exhibited a loss of E2-dependent ERα

binding at a nearby enhancer upon cotreatment with Rln without showing an observable effect on

expression (Fig. 5; Supplemental Fig. S6). Importantly, the inhibitory effects of Rln on the E2-

dependent binding of ERα at enhancers was abrogated by genetic knockout of *Rxfp1* (52), the

gene encoding the Rln receptor (Supplemental Fig. S7), reinforcing the role of Rln acting

through its receptor for the modulation of E2/ERα signaling, as well as the resulting downstream

transcriptional effects and biological outcomes.

The Rln receptor and ERα phosphorylation mediate the inhibitory effects of Rln on

estrogen signaling in myometrial tissue and cells

We considered the possibility that receptor-mediated Rln signaling might ultimately

connect to the E2 signaling pathway through phosphorylation of ERα, which has been shown to

regulate ERα-dependent gene expression (43,44,75). ERα is phosphorylated on multiple amino

acid residues, including three prominent sites in its unstructured amino-terminal region: Ser118,

Ser104/106, and Ser167 (43). Ser118 and Ser104/106 are the predominant E2-dependent

phosphorylation sites, with phosphorylation of Ser118 thought to occur in response to activation

of mitogen-activated protein kinases (MAPKs) (43). The latter was determined in breast cancer

cells, but has not be explored in detail in myometrium. We observed a dramatic increase in

Ser118 phosphorylation in response to E2 treatment in wild-type mouse myometrium, as

indicated by Western blotting (Fig. 6A). We also observed a more modest enhancement of

Ser104/106 phosphorylation upon E2 treatment, but we did not observe a significant change in

Ser167 phosphorylation (Fig. 6A; Supplemental Fig. S8). Cotreatment with Rln abrogated the

E2-mediated increase in Ser118 phosphorylation and enhanced the E2-mediated increase in

Ser104/106 phosphorylation (Fig. 6A; Supplemental Fig. S8). The effects of Rln were

dependent on the Rln receptor, as demonstrated in studies using $Rxfp1^{-/-}$ mice (Fig. 6A).

To determine if similar effects of Rln cotreatment on ERα Ser118 phosphorylation occur

in human myometrial cells, we used telomerase-immortalized human myometrial (hTERT-HM)

cells (53), which we engineered with doxycycline (Dox)-inducible expression of ERα (wild-

type, WT) and ERα mutants (S118A and S118E) by lentivirus-mediated transduction. These

cells show similarities to mouse myometrial cells in their responses to E2 and Rln based on gene

expression (Supplemental Fig. S9) and GO analysis (Supplemental Fig. S10). Moreover,

hTERT-HM cells expressing the ERα S118E mutant exhibited increased basal (i.e., without E2)

expression of an E2-regulated gene set and a loss of Rln-mediated inhibition of the same gene set

(Supplemental Fig. S11).

As with the mouse myometrium, we observed a dramatic increase in Ser118

phosphorylation in response to E2 treatment in the presence of wild-type ERα (Fig. 6B). The

effect of E2 on S118 phosphorylation was blocked by cotreatment with Rln, as well as mutation

of S118 to glutamate (S118E) or alanine (S118A) (Fig. 6B; Supplemental Fig. S12A).

Importantly, although S118E blocks the epitope recognized by the phosphorylated ERα Ser118
antibody, this mutant served as a useful phosphomimetic in the studies described below, in
contrast to the S118A mutant, due to the negatively charged glutamate. Importantly, an inhibitor
of the mitogen-activated protein kinase kinase (MEK), Trametinib (GSK1120212), which blocks
signaling through the MAPK/ERK pathway (51), had no observable effect on E2-induced ERα
S118 phosphorylation or its inhibition by cotreatment with Rln (Supplemental Fig. S12B).
Together, these studies link ERα S118 phosphorylation and global gene expression outcomes,
but indicate the phosphorylation event is not mediated by a MAPK/ERK pathway in myometrial
cells.

The inhibitory effects of Rln on ERα Ser118 phosphorylation blocks the E2-induced
contraction of myometrial cells

To explore the downstream biology of the inhibitory effects of Rln on ERα Ser118
phosphorylation, we used a collagen gel contraction assay with hTERT-HM-ERα cells
expressing wild-type or S118E mutant ERα. This assay measures the ability of the myometrial
cells to contract *in vitro*. Briefly, hTERT-HM-ERα cells (WT, S118A and S118E) with Dox-
induced ERα expression were plated in collagen with subsequent incubation for 1 hour at 37°C
to allow gelling. The cells were then treated in fresh medium with 100 nM E2, 0.2 μg/mL Rln,
or a combination of E+R. After 20 hours, the gel matrices were gently detached from the wells
and the areas of the gel matrices were measured periodically for up to 48 hr. Using this assay,
we observed a potent effect of E2 on the induction of contraction in cells expressing wild-type
ERα [Percent contraction: 24 ± 0.8 (V) versus 55 ± 1.5 (E2)] , which was completely blocked by
cotreatment with Rln [Percent contraction: 55 ± 1.5 (E2) versus 24 ± 0.5 (E+R)] (Fig. 7). We
observed a similar stimulatory effect of E2 in cells expressing ERα S118E, which was
unaffected by cotreatment with Rln [Percent contraction: 58 ± 1.3 (E2) versus 58 ± 1.0 (E+R)]
(Fig. 7). In contrast, no contraction was observed in cells expressing ERα S118A (Supplemental
Fig. S12C). These results indicate that E2-induced contraction of myometrial cells requires
phosphorylation of ERα at Ser118 and that Rln blocks this effect by inhibiting Ser118
phosphorylation.

DUSP1 and DUSP5 are required for Rln-mediated inhibition of E2-induced ERα Ser118

phosphorylation and contraction in myometrial cells

We sought to connect the effects of Rln treatment to a molecular mechanism that could

inhibit E2-induced phosphorylation of ERα at Ser118. We considered the possibility that a Rln-

regulated phosphatase might act to dephosphorylate Ser118, with nuclear dual-specificity

phosphatases (DUSPs) being good candidates. DUSPs dephosphorylate many key signaling

molecules, thereby regulating the magnitude and duration of phosphorylation-dependent events

(76,77). In this regard, a previous study demonstrated that E2-induced phosphorylation and

activation of ERα was inhibited by catalytically active DUSP22, but not catalytically inactive

mutants, in breast cancer cells (78). In our studies in myometrial cells, we focused on two

nuclear DUSPs, DUSP1 and DUSP5 (2,79,80), which are expressed in hTERT-HM-ERα cells

and whose expression is unaffected by the treatments used in this study (E2, Rln, E+R)

(Supplemental Fig. S13A).

We used siRNA-mediated knockdown to deplete DUSP1, DUSP5, or both in hTERT-

HM-ERα cells (Supplemental Fig. S13, B and C), then examined the effect on E2-induced

phosphorylation of ERα at Ser118, as well as its reversal by cotreatment with Rln. Depletion of

DUSP1, DUSP5, or both in combination alone had little effect on E2-induced Ser118

phosphorylation, but completely blocked the inhibition of Ser118 phosphorylation by Rln (Fig.

8A). In the contraction assay, treatment of hTERT-HM cells expressing wild-type ERα and

control siRNA (siControl) again showed a robust induction of contraction [Percent contraction:

24 ± 0.7 (V) versus 54 ± 0.4 (E2)] and inhibition by cotreatment with Rln [Percent contraction:

54 ± 0.4 (E2) versus 23 ± 0.6 (E+R)] (Fig. 8, B and C). Depletion of DUSP1 alone had little

effect on E2-induced contraction or inhibition by cotreatment with Rln [Percent contraction: $23 \pm$

0.5 (V), 55 ± 0.6 (E2), 27 ± 0.1 (E+R)], whereas depletion of DUSP5 alone enhanced E2-

induced contraction and partially reversed inhibition by cotreatment Rln [Percent contraction: 30

± 0.3 (V), 64 ± 0.4 (E2), 40 ± 0.9 (E+R)] (Fig. 8, B and C). Depletion of both DUSP1 and

DUSP5 enhanced E2-induced contraction and reversed inhibition by cotreatment with Rln

[Percent contraction: 28 ± 0.4 (V), 68 ± 0.3 (E2), 55 ± 0.5 (E+R)] (Fig. 8, B and C). Depletion

of DUSP1, DUSP5, or both in combination also modestly enhanced the effects of Rln treatment

alone (Fig. 8, B and C). These results implicate both DUSP1 and DUSP5 as key mediators of

the inhibitory effects of Rln on ERα, with perhaps a more prominent role for DUSP5.

DISCUSSION

In this study, we explored the molecular mechanisms by which Rln/RXFP1 signaling can

promote or oppose the actions of E2/ERα in myometrial cells and tissue, which is a poorly

studied aspect of reproductive tract biology. Our results indicate that Rln attenuates the genomic

actions and biological effects of estrogen in the myometrium by reducing ERα S118

phosphorylation. Interestingly, we observed a potent inhibitory effect of Rln/RXFP1 on ERα

binding across the genome. The reduction in ERα binding was associated with changes in the

hormone-regulated transcriptome, including a decrease in the E2-dependent expression of some

neighboring genes, but not selected classical E2-regulated genes (Fig. 9). Cotreatment with E2

and Rln also promoted an expansion of the transcriptome by further enhancing the expression of

E2-regulated genes beyond that observed with E2 or Rln alone (Fig. 9). The effects of Rln

cotreatment were manifested in a reduction in the E2-dependent phosphorylation of ERα on Ser

118, as well the E2-dependent contraction of myometrial cells. Importantly, we observed a

requirement for DUSP1 and DUSP5 for the inhibitory effects of Rln cotreatment on estrogen

signaling. Collectively, our results identify a pathway that integrates Rln/RXFP1 and E2/ERα

signaling, resulting in a convergence of membrane and nuclear signaling pathways to control

genomic and biological outcomes (Fig. 9).

Molecular mechanisms underlying the unique regulatory effects of Rln on estrogen

signaling in the myometrium

Previous studies have characterized both stimulatory and inhibitory crosstalk between

Rln and E2 signaling on biological outcomes in various reproductive tissues. For example, many

of the effects of Rln on reproductive tissues, such as cell proliferation, ECM remodeling, and

angiogenic actions, require or are augmented by E2/ERα signaling (16,26,27). In contrast, Rln

promotes smooth muscle quiescence in the myometrium, while E2/ERα promotes contraction

(2,28-30). Our studies suggest a potential mechanism for the opposing actions, namely the

control of E2-dependent gene regulation by altering ERα phosphorylation status and binding

across the genome. In this regard, a phosphomimetic ERα mutant S118E enhanced basal ERα-

dependent gene expression and conferred resistance to the inhibitory effects of Rln. We also

found that DUSP1 and DUSP5 mediate the reduction in E2-dependent phosphorylation of ERα

635 on Ser118 with Rln cotreatment (Fig. 9). Ultimately, this leads to reduced contraction of

636 myometrial cells. Interestingly, another DUSP, DUSP22, was previously shown to regulate

637 ERα-mediated signaling via direct dephosphorylation of Ser118 (78). Previous studies in breast

638 cancer cells showed that phosphorylation of ERα on Ser118 occurs predominantly in response to

639 activation of MAPKs (43). In contrast, our studies herein using a MEK inhibitor did not

640 recapitulate these results in myometrial cells, leaving open the question of the kinase that

641 phosphorylates ERα in these cells. Nonetheless, as we show herein, inhibition of E2-dependent

642 phosphorylation of ERα by cotreatment with Rln can be reversed by the actions of DUSP1 and

643 DUSP5 (Fig. 9).

645 **Limitations of the study and future directions**

646  Some of the best characterized effects of Rln are during pregnancy, where it acts to relax

647 the uterine muscle and soften the pelvic ligaments to facilitate parturition (15-17). In our

648 ovariectomized, non-pregnant mouse model, we were unable to investigate the functional

649 interplay between Rln/RXFP1 and E2/ERα signaling during pregnancy and parturition. This

650 will be an interesting direction to pursue in future studies. Regarding the molecular mechanisms

651 by which DUSP1 and DUSP5 inhibit the phosphorylation of ERα on Ser118, our results and

652 previous studies (78) suggest that this might occur through the direct actions of the DUSPs on

653 phosphorylated ERα, but we have not shown this directly in biochemical assays. In addition, as

654 noted above, we have identified the enzyme that mediates ERα phosphorylation. These will be

655 additional interesting questions to address in future studies.

ACKNOWLEDGMENTS

The authors would like to acknowledge and thank the following: Dr. Carole Mendelson

for the hTERT-HM cells; Dr. Alexander Agoulnik for the *Rxfp1*$^{-/-}$ mice; Asha Varghese, Cosette

Taggart, and Carla De Cassia Villela for assistance with mouse genotyping; ShanmugaPriyaa

Madhukaran for assistance with ovariectomy surgery and members of the Kraus and Mahendroo

laboratories for input and feedback on this project. We would also like to acknowledge the UT

Southwestern Next Generation Sequencing Core, under the direction of Ralf Kittler.

DEDICATION

The authors would like to dedicate this manuscript to the memory of Dr. Carole

Mendelson, a beloved colleague and friend: https://www.utsouthwestern.edu/education/medical-

school/departments/green-center/assets/in-memoriam-carole-mendelson.pdf

FUNDING

This work was supported by a grant from the NIH/NICHD (P01 HD087150) to M.M., a

grant from the NIH/NIDDK (R01 DK058110) to W.L.K., and funds from the Cecil H. and Ida

Green Center for Reproductive Biology Sciences Endowment to W.L.K.

CONFLICT OF INTEREST

The authors have nothing to declare.

AUTHOR CONTRIBUTIONS

ST - Conceptualization, Methodology, Validation, Formal analysis, Investigation, Writing –

original draft preparation, Visualization.

AN - Software, Formal analysis, Data Curation, Writing – original draft preparation,

Visualization.

SPC - Formal analysis, Investigation.

TN - Formal analysis, Writing – review and editing, Visualization.

CVC - Methodology, Writing – review and editing, Supervision.

MM - Conceptualization, Methodology, Writing – review and editing, Visualization,

Supervision, Project administration, Funding acquisition.

WLK - Conceptualization, Methodology, Writing – review and editing, Visualization,
Supervision, Project administration, Funding acquisition.
690

DATA AVAILABILITY

692   The genomic datasets generated for this study can be accessed from the NCBI's Gene
693   Expression Omnibus (GEO) repository (www.ncbi.nlm.nih.gov/geo/) using superseries accession
694   number GSE244843. All other data generated in this study are available within the article and its
695   supplementary data files. The pipelines and scripts used for the computational analyses are
696   available on request.
697

SUPPLEMENTARY DATA FILES

699   This paper contains Supplementary Figures S1-S13.
700

FIGURE LEGENDS

Figure 1. Rln reorganizes the ERα cistrome in the mouse myometrium by promoting the

depletion of ERα from its enhancers.

(**A**) Experimental schematic for surgical preparation and treatment of female mice, sample

processing, and genomic and downstream bioinformatic analyses. C57BL/6 ovariectomized

female mice were treated with vehicle (V, 0.1 mL saline), 17β-estradiol (E2; E, 2.5 µg/mL),

recombinant human relaxin-2 (R, 80 µg/mL 0.1% BSA-PBS) or both (E+R) for 1 or 2 hours.

(**B**) Western blots *(top)* and densitometric quantification *(bottom)* of ERα protein in mouse

myometrium samples collected at 1 hour after the indicated treatments. β-actin is a loading

control. Each bar represents the mean ± SEM, n = 3. No significant differences were observed.

(**C**) Venn diagram showing the number of ERα enhancer peaks under each treatment condition

as revealed by ChIP-seq.

(**D**) Heatmaps of ERα ChIP-seq data *(left)* from myometrium obtained from mice treated with

vehicle (V), E2 (E), Rln (R), or both (E+R). The changes shown in the heatmap were centered

relative to the control (vehicle, V) condition. Heatmaps of H3K27ac ChIP-seq data *(right)* at the

genomic loci where ERα was bound under the same treatment conditions.

(**E**) Box plots of normalized read counts for ERα and H3K27ac ChIP-seq data at three distinct

sets of ERα binding sites (gained, maintained, depleted). Bars marked with different letters are

significantly different (Wilcoxon rank sum test, $p < 0.001$).

Figure 2. Genomic analyses of the transcriptional response to E2 and Rln mouse

myometrium.

(**A through C**) Three representations of the RNA-seq data analyses: (A) Heatmap, (B) Venn

diagram, and (C) tabular representation showing differential gene expression from RNA-seq

analyses in ovariectomized mouse myometrium after 2 hours of the indicated treatments. The

changes in expression shown in the heatmap were centered relative to the control (vehicle, V)

condition and scaled relative to the maximum absolute value of the expression change for each

gene (range between +1 and -1, where +1 or -1 denote the maximum or minimum values,

respectively). In (B) and (C), we categorized the treatment-regulated genes with a p-value filter

[p value < 0.01 (q value < 0.05)] and fold change cutoff filter (≥ 2.0 or ≤ 0.5) versus vehicle.

Only genes passing both filters were included in the category.

(D) Box plots showing the effects of the treatments on (*left*) genes uniquely regulated by E2 only

(372 genes) and (*right*) genes uniquely regulated by E2+Rln (673 genes) from the Venn diagram

in (B). The data are expressed relative to control, which is set to 1. Dotted red lines indicate

fold change cutoff ≥ 2.0 or ≤ 0.5 ($\log 2 \geq 1.0$ or ≤ -1.0). Boxes marked with different letters are

significantly different from each other and the control (Wilcoxon rank sum test, p-value < 2.2 x

10^{-16}). Genes listed below the "Upregulated" categories are representative of that category (see

browser tracks in Fig. 3).

Figure 3. Genomic analysis of the effects of Rln treatment on genomic signaling by

E2/ERα.

(A and B) Genome browser views of RNA-seq and ERα, H3K27ac, and H3K4me1 ChIP-seq

data for representative genes from ovariectomized mouse myometrium after treatment with

vehicle (V), E2 (E), Rln (R), or both (E+R). Highlight (box with magenta dashed line) indicate

regions of ERα enrichment from ChIP-seq. Gene schematic and scale bars are shown. The

genes shown are (A) *Aoc1* (upregulated by E2 and inhibited by Rln) and (B) *Sik1* (further

upregulated by E2+Rln).

Figure 4. Linking the effects of Rln treatment on the ERα cistrome and E2-regulated

transcriptome.

(A) Overview of the pipeline for integrating the ChIP-seq data from Figure 1 with the RNA-seq

data to link ERα peaks depleted by Rln (R) treatment with the expression of the nearest

neighboring genes in mouse myometrium after treatment with vehicle (V), E2 (E), Rln (R), or

both (E+R).

(B) Box plots showing the $\log_2$(fold change) difference for the nearest neighboring gene

expression by RNA-seq at Rln-depleted ERα binding sites. Bars marked with different letters

are significantly different (Wilcoxon rank sum test, $p < 0.001$).

(C) Gene ontologies for the genes associated with Rln-depleted ERα peaks were evaluated using

WEB-based GEne SeT AnaLysis Toolkit (WebGestalt) (59). Multiple gene ontology terms

related to collagen metabolism and extracellular matrix (ECM) organization were identified.

Figure 5. Genomic analysis of the effects of Rln treatment on genomic signaling by E2/ERα

at a classical estrogen-regulated gene.

Genome browser views of RNA-seq and ERα, H3K27ac, and H3K4me1 ChIP-seq data for the

classical E2-regulated gene *Igf1* from ovariectomized mouse myometrium after treatment with

vehicle (V), E2 (E), Rln (R), or both (E+R). Highlight (box with magenta dashed line) indicates

regions of ERα enrichment from ChIP-seq. Gene schematic and scale bars are shown.

Figure 6. The effects of Rln on estrogen signaling in myometrial cells requires inhibition of

ERα phosphorylation at serine 118.

(A) Western blotting *(left)* and densitometric quantification *(right)* of ERα phosphorylation

levels at Ser118, Ser104/106, and Ser167 in myometrium from ovariectomized wild-type and

Rln receptor knockout (*Rxfp1*$^{-/-}$) mice treated for 1 hour with the treatments indicated: vehicle

(V), E2 (E), Rln (R), or both (E+R). β-actin serves as a loading control. Each bar represents the

mean + SEM, n = 3, 5 mice/treatment). The asterisk indicates significant difference from the

corresponding control (V) (One-way ANOVA, * $p < 0.05$).

(B) Western blotting *(left)* and densitometric quantification *(right)* of ERα phosphorylation

levels at Ser118 in Dox-inducible hTERT-HM-ERα cells with wild-type or phosphomimetic

S118E mutant ERα subjected to the treatments indicated for 30 minutes: vehicle (V), E2 (E), Rln

(R), or both (E+R). β-actin serves as a loading control. Each bar represents the mean + SEM, n

= 3. The asterisks indicate significant difference from the corresponding control (V) (One-way

ANOVA, **** $p < 0.0001$).

Figure 7. The effects of Rln on an estrogen-dependent biological outcome in myometrial

cells requires inhibition of ERα phosphorylation at serine 118.

(A and B) Representative images (A) and quantification (B) of collagen gel contraction assays

conducted in hTERT-HM-ERα cells with wild-type or S118E mutant ERα. The cells were

plated in collagen with subsequent incubation for 1 hour at 37°C to allow gelling. The cells were

then treated in fresh medium with vehicle (V), E2 (E), Rln (R), or both (E+R). After 20 hours,

the gel matrices were gently detached from the wells and the areas of the gel matrices were

measured periodically for up to 48 hr. The area of each gel was measured after 48 hours and

plotted as percent contraction compared to the area of untreated well at the beginning of

experiment. Each bar represents the mean + SEM, n = 3, 6 wells/treatment. The asterisks

indicate significant difference from the corresponding control (V) (One-way ANOVA, [****] p <

0.0001).

Figure 8. DUSP1 and DUSP5 are required for Rln-mediated inhibition of E2-induced ERα

Ser118 phosphorylation and contraction in myometrial cells.

(A) Western blotting of phospho ERα Ser118 and total ERα in hTERT-HM-ERα cells with

siRNA-mediated knockdown of *DUSP1* mRNA, *DUSP5* mRNA or both. The cells were

subjected to the treatments indicated for 30 minutes: vehicle (V), E2 (E), Rln (R), or both (E+R).

β-actin serves as a loading control.

(B) Representative images from collagen gel contraction assays conducted in hTERT-HM-ERα

cells with siRNA-mediated knockdown of *DUSP1* mRNA, *DUSP5* mRNA or both. The cells

were subjected to the treatments indicated as described in the Methods: vehicle (V), E2 (E), Rln

(R), or both (E+R).

(C) Quantification of the collagen gel contraction assays shown in (B). The area of each gel was

measured after 48 hours and plotted as percent contraction compared to the area of untreated

well at the beginning of experiment. Each bar represents the mean + SEM, n = 3, 6

wells/treatment. The asterisks indicate significant difference from the corresponding control (V)

(One-way ANOVA, [*] p < 0.05, [**] p < 0.01, [****] p < 0.0001).

Figure 9. Summary of the observations regarding the effects of Rln on ERα-dependent

gene regulation in myometrial cells.

Additional components (e.g., DUSPs 1 and 5) are involved in E2/ERα signaling upon

cotreatment with Rln. Cotreatment with Rln alters and expands the E2-regulated transcriptome,

as indicated in the bottom panel. See the text for details.

REFERENCES

1.   Kota SK, Gayatri K, Jammula S, Kota SK, Krishna SV, Meher LK, Modi KD.
Endocrinology of parturition. *Indian J Endocrinol Metab*. 2013;17(1):50-59.

2.   Renthal NE, Williams KC, Montalbano AP, Chen CC, Gao L, Mendelson CR. Molecular
regulation of parturition: A myometrial perspective. *Cold Spring Harb Perspect Med*.
2015;5(11).

3.   Hertelendy F, Zakar T. Regulation of myometrial smooth muscle functions. *Curr Pharm*
Des. 2004;10(20):2499-2517.

4.   Mesiano S, DeFranco E, Muglia LJ. Parturition. In: Plant TM, Zeleznik AJ, eds. Knobil
and Neill's Physiology of Reproduction. Vol Vol. 2. 4th ed. London, U.K.: Elsevier, Inc.;
2015:1875-1925.

5.   Aguilar HN, Mitchell BF. Physiological pathways and molecular mechanisms regulating
uterine contractility. *Hum Reprod Update*. 2010;16(6):725-744.

6.   Shynlova O, Tsui P, Jaffer S, Lye SJ. Integration of endocrine and mechanical signals in
the regulation of myometrial functions during pregnancy and labour. *Eur J Obstet*
Gynecol Reprod Biol. 2009;144 Suppl 1:S2-10.

7.   Di WL, Lachelin GC, McGarrigle HH, Thomas NS, Becker DL. Oestriol and oestradiol
increase cell to cell communication and connexin43 protein expression in human
myometrium. *Mol Hum Reprod*. 2001;7(7):671-679.

8.   Mesiano S, Chan EC, Fitter JT, Kwek K, Yeo G, Smith R. Progesterone withdrawal and
estrogen activation in human parturition are coordinated by progesterone receptor A
expression in the myometrium. *J Clin Endocrinol Metab*. 2002;87(6):2924-2930.

9.   Couse JF, Korach KS. Estrogen receptor null mice: what have we learned and where will
they lead us? *Endocr Rev*. 1999;20(3):358-417.

10.   Welboren WJ, Sweep FC, Span PN, Stunnenberg HG. Genomic actions of estrogen
receptor alpha: what are the targets and how are they regulated? *Endocrine-related*
cancer. 2009;16(4):1073-1089.

11.   Hewitt SC, Korach KS. Oestrogen receptor knockout mice: roles for oestrogen receptors
alpha and beta in reproductive tissues. *Reproduction*. 2003;125(2):143-149.

12.   Bossmar T, Osman N, Zilahi E, Haj MA, Nowotny N, Conlon JM. Expression of the
oxytocin gene, but not the vasopressin gene, in the rat uterus during pregnancy: influence
of oestradiol and progesterone. *J Endocrinol*. 2007;193(1):121-126.

13.   Ratajczak CK, Muglia LJ. Insights into parturition biology from genetically altered mice.
Pediatr Res. 2008;64(6):581-589.

14.   Wakle-Prabagaran M, Lorca RA, Ma X, Stamnes SJ, Amazu C, Hsiao JJ, Karch CM,
Hyrc KL, Wright ME, England SK. BKCa channel regulates calcium oscillations induced
by alpha-2-macroglobulin in human myometrial smooth muscle cells. *Proc Natl Acad Sci
U S A*. 2016;113(16):E2335-2344.

15.   Sherwood OD. Relaxin's physiological roles and other diverse actions. *Endocr Rev*.
2004;25(2):205-234.

16.   Bathgate RA, Halls ML, van der Westhuizen ET, Callander GE, Kocan M, Summers RJ.
Relaxin family peptides and their receptors. *Physiol Rev*. 2013;93(1):405-480.

17.   Jelinic M, Marshall SA, Stewart D, Unemori E, Parry LJ, Leo CH. Peptide hormone
relaxin: from bench to bedside. *Am J Physiol Regul Integr Comp Physiol*.
2018;314(6):R753-R760.

18.   Soh YM, Tiwari A, Mahendroo M, Conrad KP, Parry LJ. Relaxin regulates hyaluronan
synthesis and aquaporins in the cervix of late pregnant mice. *Endocrinology*.
2012;153(12):6054-6064.

19.   Marshall SA, McGuane JT, Soh YM, Gehring HM, Simpson E, Parry LJ. Abnormal
extracellular matrix remodelling in the cervix of pregnant relaxin-deficient mice is not
associated with reduced matrix metalloproteinase expression or activity. *Reprod Fertil
Dev*. 2018.

20.   Valkovic AL, Bathgate RA, Samuel CS, Kocan M. Understanding relaxin signalling at
the cellular level. *Mol Cell Endocrinol*. 2018.

21.   Yegorov S, Bogerd J, Good SV. The relaxin family peptide receptors and their ligands:
new developments and paradigms in the evolution from jawless fish to mammals. *Gen*
Comp Endocrinol. 2014;209:93-105.

22.   Vodstrcil LA, Tare M, Novak J, Dragomir N, Ramirez RJ, Wlodek ME, Conrad KP,
Parry LJ. Relaxin mediates uterine artery compliance during pregnancy and increases
uterine blood flow. *FASEB J*. 2012;26(10):4035-4044.

23.   Vodstrcil LA, Shynlova O, Verlander JW, Wlodek ME, Parry LJ. Decreased expression
of the rat myometrial relaxin receptor (RXFP1) in late pregnancy is partially mediated by
the presence of the conceptus. *Biol Reprod*. 2010;83(5):818-824.

24.   Kaftanovskaya EM, Huang Z, Lopez C, Conrad K, Agoulnik AI. Conditional deletion of
the relaxin receptor gene in cells of smooth muscle lineage affects lower reproductive
tract in pregnant mice. *Biol Reprod*. 2015;92(4):91.

25.   Krajnc-Franken MA, van Disseldorp AJ, Koenders JE, Mosselman S, van Duin M,
Gossen JA. Impaired nipple development and parturition in LGR7 knockout mice. *Mol*
Cell Biol. 2004;24(2):687-696.

26.   Sarwar M, Du XJ, Dschietzig TB, Summers RJ. The actions of relaxin on the human
cardiovascular system. *Br J Pharmacol*. 2017;174(10):933-949.

27.   Yao L, Cooke PS, Meling DD, Shanks RD, Jameson JL, Sherwood OD. The effect of
relaxin on cell proliferation in mouse cervix requires estrogen receptor {alpha} binding to
estrogen response elements in stromal cells. *Endocrinology*. 2010;151(6):2811-2818.

28.   Bani D, Baccari MC, Nistri S, Calamai F, Bigazzi M, Sacchi TB. Relaxin up-regulates
the nitric oxide biosynthetic pathway in the mouse uterus: involvement in the inhibition
of myometrial contractility. *Endocrinology*. 1999;140(10):4434-4441.

29.   Friebe-Hoffmann U, Baston DM, Chiao JP, Winebrenner LD, Krussel JS, Hoffmann TK,
Hirchenhain J, Rauk PN. The effect of relaxin on the oxytocin receptor in human uterine
smooth muscle cells. *Regul Pept*. 2007;138(2-3):74-81.

30.   Meera P, Anwer K, Monga M, Oberti C, Stefani E, Toro L, Sanborn BM. Relaxin
901   stimulates myometrial calcium-activated potassium channel activity via protein kinase A.
902   *Am J Physiol*. 1995;269(2 Pt 1):C312-317.

903   31.   Carroll JS, Liu XS, Brodsky AS, Li W, Meyer CA, Szary AJ, Eeckhoute J, Shao W,
904   Hestermann EV, Geistlinger TR, Fox EA, Silver PA, Brown M. Chromosome-wide
905   mapping of estrogen receptor binding reveals long-range regulation requiring the
906   forkhead protein FoxA1. *Cell*. 2005;122(1):33-43.

907   32.   Carroll JS, Meyer CA, Song J, Li W, Geistlinger TR, Eeckhoute J, Brodsky AS, Keeton
908   EK, Fertuck KC, Hall GF, Wang Q, Bekiranov S, Sementchenko V, Fox EA, Silver PA,
909   Gingeras TR, Liu XS, Brown M. Genome-wide analysis of estrogen receptor binding
910   sites. *Nat Genet*. 2006;38(11):1289-1297.

911   33.   Lin CY, Vega VB, Thomsen JS, Zhang T, Kong SL, Xie M, Chiu KP, Lipovich L,
912   Barnett DH, Stossi F, Yeo A, George J, Kuznetsov VA, Lee YK, Charn TH, Palanisamy
913   N, Miller LD, Cheung E, Katzenellenbogen BS, Ruan Y, Bourque G, Wei CL, Liu ET.
914   Whole-genome cartography of estrogen receptor alpha binding sites. *PLoS genetics*.
915   2007;3(6):e87.

916   34.   Hewitt SC, Li L, Grimm SA, Chen Y, Liu L, Li Y, Bushel PR, Fargo D, Korach KS.
917   Research resource: whole-genome estrogen receptor alpha binding in mouse uterine
918   tissue revealed by ChIP-seq. *Molecular endocrinology*. 2012;26(5):887-898.

919   35.   Welboren WJ, van Driel MA, Janssen-Megens EM, van Heeringen SJ, Sweep FC, Span
920   PN, Stunnenberg HG. ChIP-Seq of ERalpha and RNA polymerase II defines genes
921   differentially responding to ligands. *The EMBO journal*. 2009;28(10):1418-1428.

922   36.   Tan SK, Lin ZH, Chang CW, Varang V, Chng KR, Pan YF, Yong EL, Sung WK,
923   Cheung E. AP-2gamma regulates oestrogen receptor-mediated long-range chromatin
924   interaction and gene transcription. *The EMBO journal*. 2011;30(13):2569-2581.

925   37.   Hurtado A, Holmes KA, Ross-Innes CS, Schmidt D, Carroll JS. FOXA1 is a key
926   determinant of estrogen receptor function and endocrine response. *Nat Genet*.
927   2011;43(1):27-33.

38.   Foulds CE, Feng Q, Ding C, Bailey S, Hunsaker TL, Malovannaya A, Hamilton RA,
Gates LA, Zhang Z, Li C, Chan D, Bajaj A, Callaway CG, Edwards DP, Lonard DM,
Tsai SY, Tsai MJ, Qin J, O'Malley BW. Proteomic analysis of coregulators bound to ERα
on DNA and nucleosomes reveals coregulator dynamics. *Mol Cell*. 2013;51(2):185-199.

39.   Hah N, Murakami S, Nagari A, Danko CG, Kraus WL. Enhancer transcripts mark active
estrogen receptor binding sites. *Genome Res*. 2013;23(8):1210-1223.

40.   Fullwood MJ, Liu MH, Pan YF, Liu J, Xu H, Mohamed YB, Orlov YL, Velkov S, Ho A,
Mei PH, Chew EG, Huang PY, Welboren WJ, Han Y, Ooi HS, Ariyaratne PN, Vega VB,
Luo Y, Tan PY, Choy PY, Wansa KD, Zhao B, Lim KS, Leow SC, Yow JS, Joseph R, Li
H, Desai KV, Thomsen JS, Lee YK, Karuturi RK, Herve T, Bourque G, Stunnenberg HG,
Ruan X, Cacheux-Rataboul V, Sung WK, Liu ET, Wei CL, Cheung E, Ruan Y. An
oestrogen-receptor-alpha-bound human chromatin interactome. *Nature*.
2009;462(7269):58-64.

41.   Shlyueva D, Stampfel G, Stark A. Transcriptional enhancers: from properties to genome-
wide predictions. *Nature reviews Genetics*. 2014;15(4):272-286.

42.   He HH, Meyer CA, Chen MW, Jordan VC, Brown M, Liu XS. Differential DNase I
hypersensitivity reveals factor-dependent chromatin dynamics. *Genome Res*.
2012;22(6):1015-1025.

43.   Lannigan DA. Estrogen receptor phosphorylation. *Steroids*. 2003;68(1):1-9.

44.   Murphy LC, Seekallu SV, Watson PH. Clinical significance of estrogen receptor
phosphorylation. *Endocrine-related cancer*. 2011;18(1):R1-14.

45.   Duplessis TT, Williams CC, Hill SM, Rowan BG. Phosphorylation of Estrogen Receptor
alpha at serine 118 directs recruitment of promoter complexes and gene-specific
transcription. *Endocrinology*. 2011;152(6):2517-2526.

46.   Murakami S, Nagari A, Kraus WL. Dynamic assembly and activation of estrogen
receptor alpha enhancers through coregulator switching. *Genes Dev*. 2017;31(15):1535-
1548.

47. Helzer KT, Ozers MS, Meyer MB, Benkusky NA, Solodin N, Reese RM, Warren CL,
Pike JW, Alarid ET. The Phosphorylated Estrogen Receptor Cistrome Identifies a Subset
of Active Enhancers Enriched for Direct ER-DNA Binding and the Transcription Factor
GRHL2. *Mol Cell Biol*. 2018.

48. Hermon TL, Moore AB, Yu L, Kissling GE, Castora FJ, Dixon D. Estrogen receptor
alpha (ERalpha) phospho-serine-118 is highly expressed in human uterine leiomyomas
compared to matched myometrium. *Virchows Arch*. 2008;453(6):557-569.

49. Yu L, Moore AB, Castro L, Gao X, Huynh HL, Klippel M, Flagler ND, Lu Y, Kissling
GE, Dixon D. Estrogen Regulates MAPK-Related Genes through Genomic and
Nongenomic Interactions between IGF-I Receptor Tyrosine Kinase and Estrogen
Receptor-Alpha Signaling Pathways in Human Uterine Leiomyoma Cells. *J Signal*
Transduct. 2012;2012:204236.

50. Kraus WL, Kadonaga JT. p300 and estrogen receptor cooperatively activate transcription
via differential enhancement of initiation and reinitiation. *Genes Dev*. 1998;12(3):331-
342.

51. Abe H, Kikuchi S, Hayakawa K, Iida T, Nagahashi N, Maeda K, Sakamoto J, Matsumoto
N, Miura T, Matsumura K, Seki N, Inaba T, Kawasaki H, Yamaguchi T, Kakefuda R,
Nanayama T, Kurachi H, Hori Y, Yoshida T, Kakegawa J, Watanabe Y, Gilmartin AG,
Richter MC, Moss KG, Laquerre SG. Discovery of a Highly Potent and Selective MEK
Inhibitor: GSK1120212 (JTP-74057 DMSO Solvate). *ACS Med Chem Lett*.
2011;2(4):320-324.

52. Kamat AA, Feng S, Bogatcheva NV, Truong A, Bishop CE, Agoulnik AI. Genetic
targeting of relaxin and insulin-like factor 3 receptors in mice. *Endocrinology*.
2004;145(10):4712-4720.

53. Condon J, Yin S, Mayhew B, Word RA, Wright WE, Shay JW, Rainey WE. Telomerase
immortalization of human myometrial cells. *Biol Reprod*. 2002;67(2):506-514.

54. Zhong S, Joung JG, Zheng Y, Chen YR, Liu B, Shao Y, Xiang JZ, Fei Z, Giovannoni JJ.
High-throughput illumina strand-specific RNA sequencing library preparation. *Cold*
Spring Harb Protoc. 2011;2011(8):940-949.

55.   Andrews S. FastQC: A quality control tool for high throughput sequence data. Vol
985   20152015.

986   56.   Kim D, Pertea G, Trapnell C, Pimentel H, Kelley R, Salzberg SL. TopHat2: accurate
987   alignment of transcriptomes in the presence of insertions, deletions and gene fusions.
988   *Genome Biol*. 2013;14(4):R36.

989   57.   Trapnell C, Roberts A, Goff L, Pertea G, Kim D, Kelley DR, Pimentel H, Salzberg SL,
990   Rinn JL, Pachter L. Differential gene and transcript expression analysis of RNA-seq
991   experiments with TopHat and Cufflinks. *Nat Protoc*. 2012;7(3):562-578.

992   58.   Saldanha AJ. Java Treeview - extensible visualization of microarray data. *Bioinformatics*.
993   2004;20(17):3246-3248.

994   59.   Wang J, Vasaikar S, Shi Z, Greer M, Zhang B. WebGestalt 2017: a more comprehensive,
995   powerful, flexible and interactive gene set enrichment analysis toolkit. *Nucleic Acids Res*.
996   2017;45(W1):W130-W137.

997   60.   Ashburner M, Ball CA, Blake JA, Botstein D, Butler H, Cherry JM, Davis AP, Dolinski
998   K, Dwight SS, Eppig JT, Harris MA, Hill DP, Issel-Tarver L, Kasarskis A, Lewis S,
999   Matese JC, Richardson JE, Ringwald M, Rubin GM, Sherlock G. Gene ontology: tool for
1000   the unification of biology. The Gene Ontology Consortium. *Nat Genet*. 2000;25(1):25-29.

1001   61.   Mi H, Muruganujan A, Ebert D, Huang X, Thomas PD. PANTHER version 14: more
1002   genomes, a new PANTHER GO-slim and improvements in enrichment analysis tools.
1003   *Nucleic Acids Res*. 2019;47(D1):D419-D426.

1004   62.   Franco HL, Nagari A, Kraus WL. TNFalpha signaling exposes latent estrogen receptor
1005   binding sites to alter the breast cancer cell transcriptome. *Mol Cell*. 2015;58(1):21-34.

1006   63.   Luo X, Chae M, Krishnakumar R, Danko CG, Kraus WL. Dynamic reorganization of the
1007   AC16 cardiomyocyte transcriptome in response to TNFalpha signaling revealed by
1008   integrated genomic analyses. *BMC Genomics*. 2014;15:155.

1009   64.   Lis JT, Schleif R. Size fractionation of double-stranded DNA by precipitation with
1010   polyethylene glycol. *Nucleic Acids Res*. 1975;2(3):383-389.

1011   65.   Martin M. Cutadapt removes adapter sequences from high-throughput sequencing reads.
1012   *EMBnetjournal*. 2011;17.

1013   66.   Langmead B, Trapnell C, Pop M, Salzberg SL. Ultrafast and memory-efficient alignment
1014   of short DNA sequences to the human genome. *Genome Biol*. 2009;10(3):R25.

1015   67.   Quinlan AR, Hall IM. BEDTools: a flexible suite of utilities for comparing genomic
1016   features. *Bioinformatics*. 2010;26(6):841-842.

1017   68.   Kent WJ, Zweig AS, Barber G, Hinrichs AS, Karolchik D. BigWig and BigBed: enabling
1018   browsing of large distributed datasets. *Bioinformatics*. 2010;26(17):2204-2207.

1019   69.   Landt SG, Marinov GK, Kundaje A, Kheradpour P, Pauli F, Batzoglou S, Bernstein BE,
1020   Bickel P, Brown JB, Cayting P, Chen Y, DeSalvo G, Epstein C, Fisher-Aylor KI,
1021   Euskirchen G, Gerstein M, Gertz J, Hartemink AJ, Hoffman MM, Iyer VR, Jung YL,
1022   Karmakar S, Kellis M, Kharchenko PV, Li Q, Liu T, Liu XS, Ma L, Milosavljevic A,
1023   Myers RM, Park PJ, Pazin MJ, Perry MD, Raha D, Reddy TE, Rozowsky J, Shoresh N,
1024   Sidow A, Slattery M, Stamatoyannopoulos JA, Tolstorukov MY, White KP, Xi S,
1025   Farnham PJ, Lieb JD, Wold BJ, Snyder M. ChIP-seq guidelines and practices of the
1026   ENCODE and modENCODE consortia. *Genome Res*. 2012;22(9):1813-1831.

1027   70.   Heinz S, Benner C, Spann N, Bertolino E, Lin YC, Laslo P, Cheng JX, Murre C, Singh H,
1028   Glass CK. Simple combinations of lineage-determining transcription factors prime cis-
1029   regulatory elements required for macrophage and B cell identities. *Mol Cell*.
1030   2010;38(4):576-589.

1031   71.   Rao NA, McCalman MT, Moulos P, Francoijs KJ, Chatziioannou A, Kolisis FN, Alexis
1032   MN, Mitsiou DJ, Stunnenberg HG. Coactivation of GR and NFKB alters the repertoire of
1033   their binding sites and target genes. *Genome Res*. 2011;21(9):1404-1416.

1034   72.   Abramoff MD, Magalhaes, P.J., Ram, S.J. Image Processing with ImageJ. *Biophotonics*
1035   *International*. 2004;11(7):36-42.

1036   73.   Andersson R, Sandelin A. Determinants of enhancer and promoter activities of regulatory
1037   elements. *Nature reviews Genetics*. 2020;21(2):71-87.

1038   74.   Calo E, Wysocka J. Modification of enhancer chromatin: what, how, and why? *Mol Cell.*
1039   2013;49(5):825-837.

1040   75.   Anbalagan M, Rowan BG. Estrogen receptor alpha phosphorylation and its functional
1041   impact in human breast cancer. *Mol Cell Endocrinol*. 2015;418 Pt 3:264-272.

1042   76.   Caunt CJ, Keyse SM. Dual-specificity MAP kinase phosphatases (MKPs): shaping the
1043   outcome of MAP kinase signalling. *FEBS J*. 2013;280(2):489-504.

1044   77.   Huang CY, Tan TH. DUSPs, to MAP kinases and beyond. *Cell Biosci*. 2012;2(1):24.

1045   78.   Sekine Y, Ikeda O, Hayakawa Y, Tsuji S, Imoto S, Aoki N, Sugiyama K, Matsuda T.
1046   DUSP22/LMW-DSP2 regulates estrogen receptor-alpha-mediated signaling through
1047   dephosphorylation of Ser-118. *Oncogene*. 2007;26(41):6038-6049.

1048   79.   Kidger AM, Rushworth LK, Stellzig J, Davidson J, Bryant CJ, Bayley C, Caddye E,
1049   Rogers T, Keyse SM, Caunt CJ. Dual-specificity phosphatase 5 controls the localized
1050   inhibition, propagation, and transforming potential of ERK signaling. *Proc Natl Acad Sci*
1051   *U S A*. 2017;114(3):E317-E326.

1052   80.   Seo H, Cho YC, Ju A, Lee S, Park BC, Park SG, Kim JH, Kim K, Cho S. Dual-
1053   specificity phosphatase 5 acts as an anti-inflammatory regulator by inhibiting the ERK
1054   and NF-kappaB signaling pathways. *Sci Rep*. 2017;7(1):17348.

1055

A

V E R E+R

Reduced Expression

Enhanced Expression

B

Differential Gene Expression from RNA-Seq

E2   372 ⎡137 Up
            ⎣235 Down

547   1,477

Rln

172   183

E+R   673 ⎡311 Up
              ⎣362 Down

C

Treatment	Up	Down
E2	1,109	1,332
Rln	412	535
E+R	1,396	1,484

D

Uniquely Regulated by E2 Only (372 Genes)

Upregulated — *137 Genes*

Log_2(Fold Change)

a *b* *c*

V E R E+R

Downregulated — *235 Genes*

Log_2(Fold Change)

a *b* *c*

V E R E+R

e.g., Aoc1

Uniquely Regulated by E2+Rln (673 Genes)

Upregulated — *311 Genes*

Log_2(Fold Change)

a *b* *c*

V E R E+R

Downregulated — *362 Genes*

Log_2(Fold Change)

a *b* *c*

V E R E+R

e.g., Sik1

Figure 2 – Tripathy *et al.* (2024)

Figure 3 – Tripathy *et al.* (2024)

Figure 4 – Tripathy *et al.* (2024)

Classical E2-Upregulated Gene
Unaffected by Cotreatment with Rln

Figure 7 – Tripathy *et al.* (2024)

Figure 8 – Tripathy *et al.* (2024)

Figure 9 – Tripathy *et al.* (2024)