Title: Loss of circulating CD8α[+] NK cells during human *Mycobacterium tuberculosis* infection

Authors: Nezar Mehanna[1,†], Atul Pradhan[1,†], Rimanpreet Kaur[1,†], Theodota Kontopoulos[2], Barbara Rosati[1], David Carlson[1], Nai-Kong Cheung[2], Hong Xu[2], James Bean[2], Katherine Hsu[2], Jean-Benoit Le Luduec[2, ‡], and Charles Kyriakos Vorkas[1,3,4‡]

Affiliations:

[1]Renaissance School of Medicine at Stony Brook University, Stony Brook, NY, 11794

[2]Sloan Kettering Institute, Memorial Sloan Kettering Cancer Center, New York, NY 10065

[3]Department of Microbiology and Immunology, Stony Brook University, Stony Brook, NY 11794

[4]Center for Infectious Diseases, Stony Brook University, Stony Brook, NY, 11794

†, ‡: These authors contributed equally to this manuscript.

*Corresponding author Charles Kyriakos Vorkas, MD, charles.vorkas@stonybrookmedicine.edu,

101 Nicolls Road, Health Sciences Center 15060-I, Stony Brook, NY 11794.

Abstract: Natural Killer (NK) cells can recognize and kill *Mtb*-infected cells in vitro, however their role after natural human exposure has not been well-studied. To identify *Mtb*-responsive NK cell populations, we analyzed the peripheral blood of healthy household contacts of active Tuberculosis (TB) cases and source community donors in an endemic region of Port-au-Prince, Haiti by flow cytometry. We observed higher CD8α expression on NK cells in putative resistors (IGRA- contacts) with a progressive loss of these circulating cells during household-associated latent infection and disease. In vitro assays and CITE-seq analysis of CD8α[+] NK cells demonstrated enhanced maturity, cytotoxic gene expression, and response to cytokine stimulation

relative to CD8α^- NK cells. CD8α^+ NK cells also displayed dynamic surface expression dependent

on MHC I in contrast to conventional CD8$^+$ T cells. Together, these results support a specialized

role for CD8α^+ NK cell populations during *Mtb* infection correlating with disease resistance.

INTRODUCTION

Tuberculosis (TB) remains the leading global cause of death from a single bacterial infection and

is a significant public health concern (*1, 2*). Exposure to *Mycobacterium tuberculosis* (*Mtb*) bacilli

can result in a broad spectrum of clinical outcomes ranging from clearance of primary infection,

asymptomatic latent TB infection (LTBI)—measured by *Mtb* peptide-specific interferon γ release

assay (IGRA)—or symptomatic active pulmonary TB disease (*2*). It is estimated that 25% of the

global population has evidence of latent infection and 10% will progress to active pulmonary

disease (*3*). The host immunologic determinants of these clinical outcomes remain poorly defined

and are a major obstacle to designing effective TB vaccines and host-directed therapies to prevent

disease. Most TB immunology studies focus on conventional peptide-specific T cells (*4-8*), but

the presence of these cells have not been shown to correlate with protection (*9-14*). In contrast,

few studies have examined *Mtb*-reactive innate lymphocytes that contribute to *Mtb* clearance (*15-

18*).

Here, we focus on the most abundant innate lymphocytes in mammals: Natural Killer (NK) cells

(*19*). NK cells are prevalent in the peripheral blood (0.1-18% of leukocytes) and populate tissue

sites such as liver, lung, and mucosal barriers (*20*). They express a broad range of germ-line

encoded pattern recognition receptors enabling rapid responses to infection without the need for

antigen-specific expansion (*21*). Prior work provides mechanistic evidence for NK cell recognition

of intracellular and extracellular *Mtb* in vitro through natural cytotoxicity receptor engagement of

Mtb cell wall components or stress-induced self-ligands (*22-33*). However, few published studies

directly examine NK cell biology during natural *Mtb* transmission in humans (*27, 28, 34*).

In this study, we interrogate peripheral blood NK cells in a cohort of household TB contacts in an

endemic region in Port-au-Prince, Haiti. We analyzed samples from active TB cases (n=50) and

their contacts from the same households (n=50) in addition to interferon-gamma (IFNγ) release

assay (IGRA)-matched unrelated donors from the source community without household exposure

(n=50). We used spectral flow cytometry to test our hypothesis that specific NK cell

subpopulations are activated and expand during household-associated *Mtb* exposure and infection.

We identify preserved frequency of CD8α$^+$ NK cells in blood of uninfected household contacts

(IGRA$^-$) with progressive loss of this population during household latent infection and active TB

disease. CD8α$^+$ NK cells demonstrated enhanced responsiveness to cytokine stimulation in vitro

compared to CD8α$^-$ NK cells and exhibited dynamic surface expression in culture and after in

vitro stimulation. Cellular indexing of transcriptomes and epitopes (CITEseq) revealed largely

overlapping transcriptional programs between CD8α$^{+/-}$ NK cells with enhanced expression of

select cytotoxicity-associated genes in the CD8α$^+$ subset. We also observed ubiquitous

intracellular NK cell CD8α protein expression. Taken together, our results identify a novel

subpopulation of *Mtb*-reactive CD8α$^+$ NK cells associated with resistance to primary tuberculosis

infection.

RESULTS

Clinical cohort description:

To study early events during the immune response to *Mtb*, we used biobanked samples from a

cross-sectional study performed at GHESKIO Centers, Port-au-Prince, Haiti to assemble a clinical

cohort of household contacts of active pulmonary TB cases between 2015-2018 (NIH U19

AI111143, completed). Out of the total 92 household contacts and 591 community controls

enrolled, we randomly sampled an age-, sex-, and IGRA-matched subset of 50 TB household

contacts and 50 active TB cases from the same households, as well as 50 source community

controls without household exposure, based on peripheral blood mononuclear cell (PBMC)

availability and quality control (total N=150). All available specimens from IGRA- contacts were

analyzed.

Age, sex, IGRA status, and relevant sociodemographic variables are summarized in Table 1. No

significant variations were observed in the number of household members, household income,

smoking status, or alcohol consumption between groups, consistent with previous work (*15*). Fifty

percent of contacts slept in the same room as the active pulmonary TB index case. In the overall

study cohort, there were a 50% prevalence of IGRA-positivity in the community controls relative

to 75% in contacts, consistent with increased transmission within the household. Among the eleven

IGRA- contacts enrolled, only 2 IGRA conversions were detected during 6-month follow-up IGRA

testing.

IGRA+ community controls have elevated frequency of mature CD56$^-$ NK cells:

We first assessed the frequency of NK cell subpopulations in the clinical cohorts using flow

cytometry. Human NK cells were defined by the absence of lineage markers for T cells (CD3),

monocytes (CD14), and B cells (CD19), and gated on surface expression of neural cell adhesion

molecule (NCAM), CD56, and Fc receptor γRIIIA, CD16 (**Fig. 1A**). We stratified NK cells into

three subsets based on expression of CD56: CD56$^-$, CD56^{dim}, and CD56^{bright} (*35, 36*). There was

no difference in frequency of total NK cells or CD56^{dim} NK cells between clinical groups (**Fig. 1B**

and 1C center). IGRA+ community controls had fewer CD56^{bright} NK cells (3.11% vs 5.45%,

p=0.006) and more CD56⁻ NK cells (19.1% vs 12.6%, p=0.002) compared to IGRA- controls or

IGRA+ contacts, indicating an accumulation of mature NK cells during latent TB infection within

the source community **(Fig. 1C, right and left)**.

IGRA+ contacts demonstrate a loss of circulating CD16⁺ NK cells:

NK cell activity against *Mtb* depends upon complex combinatory signaling transduced by

activating and inhibiting receptors (*37-39*). We next investigated markers of NK cell activation ex

vivo. We analyzed representative receptors, including CD16a (FcγRIIIA), which binds the F_c

region of antibodies and can induce antibody-dependent cellular cytotoxicity (ADCC). Within the

major CD56^{dim} subset, IGRA+ contacts had decreased circulating CD16⁺ NK cells compared to

IGRA+ controls (60% vs 80%, p < 0.0001) with a reciprocal increase in CD16⁻ population,

representing downregulation or shedding of CD16 surface expression during household-associated

LTBI **(Fig. 2A)** (*40*). We found no significant differences in other canonical markers of NK cell

activation ex vivo including activation markers CD69 or CD25, or HLA-E restricted receptor

NKG2C. We observed decreased staining for degranulation marker CD107a in CD56^{bright} NK cells

of IGRA- contacts, potentially representing recent activation events during household exposure.

(**Fig. 2B - D**).

NK cells of IGRA- contacts have a depressed functional phenotype:

To assess functional activity of NK cells in vitro, we cultured PBMCs in media alone or with K562

cells—a leukemia cell line with low levels of HLA I expression known to activate NK cells—co-

incubated at 1:20 K562:PBMC for 15 hours. Cells were then stained, and percent difference was

calculated for CD69, CD107a, CD16, and NKG2C. Our positive control analyses in **Fig. 3A**

display results from the major CD56^{dim} subset and demonstrate CD107a, CD69, and NKG2C were

significantly increased with K562, while CD16 was significantly decreased, across all clinical groups.

Next, we directly compared NK cell response to K562 between clinical groups. IGRA+ contacts demonstrated significantly more degranulation (CD107a) and NKG2C expression after stimulation relative to IGRA- contacts (**Fig. 3B, left**). IGRA+ contacts also demonstrated depressed CD69 responses relative to IGRA+ controls (**Fig. 3B, top right**). No significant differences in CD16 expression were detected. The same analyses applied to CD56$^-$ and CD56^{bright} NK cells demonstrated similar responses to stimulation, but no significant differences between clinical groups (**Fig. S1, A - D**). Taken together with our ex vivo data, these in vitro restimulation results indicate that CD56^{dim} NK cells of putative *Mtb* resistors (IGRA- contacts) have depressed function, potentially representing sequelae of recent household-associated *Mtb* exposure.

Loss of circulating CD8α^+ NK cells during household-associated Mtb infection:

Despite their canonical classification as short-lived innate lymphocytes, NK cells are increasingly recognized for their marked diversity and ability to differentiate into longer-lived effectors (*16, 18, 27, 41-44*), including after mycobacterial stimuli (*45*). The most extensively studied is an NK cell memory population co-expressing NKG2C and CD57 that expands in blood during human cytomegalovirus (CMV) infection (*27, 46*). Multiple putative markers of NK cell memory during *Mtb* infection have been suggested, including CD57, CD8α, CD161, and CD27 (*28, 34, 45*). We compared expression of these markers across clinical groups and found that household-associated LTBI (IGRA+ contacts) and active TB disease donors had progressively decreasing frequency of CD8α^+CD56^{dim} NK cells (**Fig. 4A**). A similar trend was observed with the marker of terminal differentiation, CD57. There was significantly fewer CD57$^+$CD56^{dim} NK cells in household-associated LTBI (IGRA+ contacts) compared to healthy household contacts (**Fig. 4B**). There was

no association between CD8α expression and age or sex, and loss of CD8α$^+$ was specific to NK

cells and was not observed among T cells during active TB (**Fig. S2A-B**). We did not observe any

significant differences in the expression of CD27, CD127, or CD161 between clinical groups (**Fig**

S3). Together, these results indicate that donors who develop household-associated LTBI and

active TB disease experience a loss of circulating CD8α$^+$ and CD57$^+$ CD56^{dim} NK cells. In turn,

CD8α$^+$ NK cell frequency significantly correlates with resistance to primary household-associated

infection.

CD8α$^+$ NK cells represent a mature population with high functional potential:

Despite the identification of cytotoxic CD8α$^+$ NK cells over 30 years ago (*47-49*), the function of

CD8α on NK cells during health and disease is not well-characterized and is of emerging interest

as a cellular target against immune-mediated diseases (*47, 49, 50*). We next used flow cytometry

to define co-expressed receptors of CD8α$^+$ NK cells in our cohorts. In the Haitian cohorts, a mean

of 27% of NK cells expressed CD8α on their surface. Among NK cell subsets, CD56^{dim} NK cells

had the largest proportion of CD8 positivity (median: 36%, range: 1-74%) (**Fig. S4A**). We then

defined co-expressed receptors on CD8α$^+$ NK cells among all healthy Haitian donors, excluding

active TB cases with chronic symptomatic disease. CD8α$^+$ NK cells expressed significantly higher

levels of CD57 and CD16, and lower levels of CD25 and CD127 (**Fig. 4C**). This indicates that

CD8α$^+$ NK cells represent more mature effectors compared to their CD8α$^-$ counterparts. When

compared to T cells, median fluorescence intensity (MFI) of CD8α surface expression was

significantly lower in NK cells (**Fig. S4B**).

To interrogate function, PBMCs were co-incubated with K562 cells for 15 hours and compared to

resting conditions as described above, stratified by CD8α. CD8α$^+$ NK cells were more responsive

to K562 induction measured by CD69 upregulation compared to CD8α⁻ NK cells but demonstrated

lower levels of degranulation as measured by CD107a staining (**Fig. 4D**). The propensity for

CD8α⁺ NK cells to upregulate CD69 upon K562 stimulation was more pronounced among the

CD56⁻ compartment, where the CD8α⁻ fraction had little to no response (**Fig. 4E**). Overall, this

suggests that CD8α expression among NK cells marks a population with higher capacity for

activation, specifically among terminally differentiated CD56⁻ NK cells, that are reported to have

depressed effector functions (*51*).

CD8α is ubiquitously expressed in NK cells:

We next interrogated the immune phenotype and function of CD8α⁺ NK cells in a distinct

biorepository of healthy human PBMCs (IRB2021-00478; PI: Vorkas) to validate our findings and

perform additional functional assays. First, we confirmed the presence of CD8α expression in NK

cells by extracellular staining with CD8α and CD8β antibodies. About 60% of NK cells did not

express surface CD8, CD8αα was found in ~40% of NK cells, while rare CD8αβ NK cells were

detected (**Fig. 5A**). In blood, CD8α was expressed in all three compartments of NK cells, and most

frequently among CD56ᵈⁱᵐ cells, with median expression comparable to Haitian donors (**Fig. 5B**).

We also validated our prior result from the contact cohort (**Fig. 4C**) demonstrating CD16 and

CD8α were significantly co-expressed (**Fig. 5C).** As our TB contact studies only assessed surface

CD8α on NK cells, we next analyzed intracellular CD8α expression by flow cytometry. We

observed that nearly all CD56ᵈⁱᵐ NK cells expressed CD8α protein intracellularly, contrasting with

T cells whose CD8 expression did not significantly differ between surface and intracellular

staining (**Fig. 5D**). CD56ᵇʳⁱᵍʰᵗ and CD56⁻ NK cells also demonstrated ubiquitous expression of

CD8α protein intracellularly (**Fig. S5**). To test the stability of surface CD8α on NK cells over time,

we incubated PBMCs for 1 or 7 days in pro-survival cytokine IL-2. We observed significantly

increased CD8α surface expression following prolonged incubation, indicating that surface

presentation of NK cell CD8α is more dynamic than in T cells where it is a constitutively expressed

surface receptor (**Fig. 5E**).

CD8α$^+$ NK cells have enhanced responses to cytokine stimulation:

We next hypothesized that surface CD8α was associated with enhanced responsiveness to

inflammatory signals, following prior reports of increased cytotoxicity against leukemic cells

relative to CD8α$^-$ NK cells (*48*). We stimulated PBMCs with either the potent IL15-IL15Rα sushi

domain fusion construct (IL15-15RαSu) (*52*), IL15-15RαSu/IL12/IL18, IL12/IL18 alone, or

virulent *Mtb* Erdman whole cell lysate and assessed intracellular IFNγ expression by flow

cytometry. Cytokine stimulation led to decreased CD8α$^+$ surface staining while *Mtb* stimulation

did not alter CD8 surface expression, suggesting that cytokine stimulation induces downregulation

of the CD8α receptor or cell death (**Fig. 6A**). Both IL15/IL12/IL18 and to a lesser degree *Mtb*

lysate conditions induced IFNγ responses in CD56^{dim} NK cells compared to resting (**Fig. 6B**).

When stratified by CD8α, we observed increased intracellular IFNγ with cytokine stimulation in

both CD16$^{-/+}$ CD8α$^+$ NK cells relative to CD8α$^-$ cells, but no differences upon co-incubation with

Mtb lysates (**Fig. 6C**). Percentage of CD25, CD69, and intracellular Granzyme B responses were

also enhanced in CD8α$^+$ NK cells across CD56-stratified subsets (**Fig. S6**).

CD8α$^+$ surface expression on NK cells is MHC I-dependent:

To determine a mechanism by which CD8α may modulate NK cell function, we hypothesized that

CD8α may engage MHC I molecules like CD8$^+$ T cells (*53, 54*). To test this, we stimulated whole

PBMCs with MHC I blocking antibody (clone: W6/32) and measured surface CD8α expression.

Strikingly, surface CD8α expression on NK cells—but not T cells—was significantly decreased

in all anti-MHC I conditions (**Fig. 6D**). Our results indicate that CD8α surface expression is

maintained through engagement of MHC I molecules.

CD8α⁺ NK cells have overlapping transcriptomes with CD8α⁻ NK cells:

While CD8α-expressing NK cells have previously been described (*47-49, 55*), the transcriptional

programs associated with surface expression of CD8α and whether these relate to homologous

function in CD8α⁺ T cells during disease is an area of emerging interest. To interrogate the

transcriptional profile of CD8α⁺ NK cells, we sorted innate lymphocyte subsets from PBMCs

including NK cells, γδ T cells, MAIT cells, and invariant Natural Killer T cells (iNKTs) by

fluorescence-activated cell sorting (FACS) from the peripheral blood of two healthy donors. We

performed CITE-seq (*7*) of these populations directly after sorting or after coincubation for seven

days in IL2/IL7 supplemented media +/- *Mtb* lysates in order to define surface protein and

transcript expression in CD8α NK cells. We discriminated NK cells from surrounding innate

lymphocytes based on surface antibody barcodes for CD56 and CD16 together with expression of

canonical NK cell gene transcripts *CD3E*, *NCR1*, and *NCAM1* (CD56) (**Fig. S7**) (*47, 56, 57*).

Dimensionality reduction with UMAP identified four clusters of NK cells with distinct

transcriptional signatures (**Fig. 7, A to D**). To assess pathogen-specific responses, we first

performed differential expression between the 'rest' and '*Mtb* lysate' conditions and display the

top 8 most differentially expression genes in **Fig. 7E**. The cytotoxic serine protease *GZMB*

*(*Granzyme B*)* and killer immunoglobulin-like receptor, *KIR3DL1*, were the most upregulated

genes in the *Mtb* lysate condition.

We next analyzed the distribution of *CD8A* transcript and CD8α protein expression among NK

cells using both transcript and surface antibody barcodes for CD8α. NK cells did not cluster

independently based on the presence of *CD8A/*CD8α (**Fig. 8, F and G**) and *CD8A*$^+$/CD8α$^+$ NK

cells were distributed throughout all clusters (**Fig. 8I**). Low amplification of surface antibody

barcodes for CD8α was observed in NK cells, in contrast with other CD8α -expressing immune

subsets sequenced (e.g., MAIT cells; **Fig. S8**) in line with CD8α MFI differences detected by flow

cytometry (**Fig. S4B**).

To define transcriptional signatures associated with *CD8A* transcript expression, we stratified NK

cells expressing greater than zero *CD8A* transcripts and performed differential expression relative

to *CD8A*$^-$ cells. Half of all NK cells expressed *CD8A* transcripts (**Fig. 8J**). Differential gene

expression revealed few statistically significant differences between *CD8A*$^{+/-}$ cells. Notable

differentially expressed genes in *CD8A*$^+$ NK cells include cytotoxicity-associated proteins natural

cytotoxicity receptor 3 (*NCR3*) and the serine protease granzyme M (*GZMM*) (**Fig. 8K**).

DISCUSSION

Identifying the correlates of protection against natural *Mtb* infection is an area of critical

importance that will inform the next generation of TB vaccines and host-directed therapies (*5, 37,*

45, 58-61). In this study, we hypothesized that investigation of household-associated *Mtb* exposure

would reveal specialized NK cell populations and represent early events following high-risk

transmission. Our results identify a CD8α-expressing NK cell subpopulation with preserved

frequency in the blood of contacts who remain uninfected (IGRA-) that is progressively depleted

during household LTBI and active TB disease. Few studies have interrogated CD8α$^+$ NK cells, as

this receptor is most commonly understood as a lineage marker for T cells (*62*) and is absent in

specific pathogen free murine NK cells (*63*). Recent work reporting an association of CD8α$^+$ NK

cells with protection against human infectious, autoinflammatory and malignant disease has

generated interest in targeting this subpopulation for host-directed therapeutics (*39, 47, 48*).

However, little is known how CD8α receptors may modulate NK cell activity, namely, enhanced

cytotoxic functions previously reported in vitro against leukemia cells and strong associations with

protection against both infectious and non-infectious diseases. (*48, 50, 55*). Here, we show that the

majority of peripheral blood NK cells can express CD8α protein, but neither *CD8A* transcript nor

CD8α protein marked a distinct transcriptional state in healthy donors. Importantly, we

demonstrate that surface expression of CD8α on NK cells depends upon MHC I expression and

marks a population with enhanced responsiveness to cytokine stimuli in vitro. Together, our study

highlights a novel innate effector against human tuberculosis and provides important insights into

NK cell biology and innate TB immunology.

To our knowledge, there have been only two published household TB contact studies that

specifically examined NK cell biology during early *Mtb* exposure and infection. Our study expands

upon the sample size of this prior work and provides additional analyses of NK cell subpopulations

stratified by CD56 not previously addressed (*42, 64, 65*). Specifically, we observed decreased

frequency of immature CD56[bright] NK cells with reciprocal increase in mature CD56[-] NK cells

during community LTBI, in line with findings during chronic viral infections such hepatitis C and

HIV (*66-68*). We also observed that CD16 expression was increased during community LTBI

relative to household-associated infection, which may represent a late effect of latency within the

CD56[dim] subset (*61*). An equally valid interpretation of these results is that loss of surface CD16

is an early event detected after household-associated exposure and may represent downregulation

or shedding of the $F_c\gamma$ receptor after engagement (*40*). Together, these changes in frequency of

CD16 and CD56 expression indicate a gradual expansion of mature NK cells in peripheral blood

after asymptomatic infection (*29, 40, 64, 65, 67, 69*), accompanied by both markers of maturity

and dysfunction (loss of CD56), but also enhanced potential for ADCC (gain of CD16).

Two phenotypic markers whose surface expression were both independently and inversely

associated with infection in households, were CD8α and CD57. CD8α has long been appreciated

to be expressed on the surface of human (*47-50, 70*) and non-human primate NK cells (*63, 71*).

CD8α expression on human NK cells has been associated with enhanced cytotoxicity (*48, 55*) and

is referred to in one study as a putative *Mtb* memory marker (*45*). Recent studies in prospective

longitudinal human cohorts report that loss of CD8α expression at the transcript (*CD8A*) or protein

level correlate with HIV progression (*39, 49*), relapsing multiple sclerosis (*47*), and recurrence of

acute myeloid leukemia after stem cell transplantation (*50*), where CD8α^+ NK cells are

hypothesized to exercise combined cytotoxic and regulatory roles to prevent disease. Sorted

surface CD8α^+ NK cells also demonstrate enhanced anti-leukemia cytotoxicity in vitro compared

to sorted CD8α^- cells (*50*). We also report that CD8α^+ NK cells were prevalent in healthy US

donors with ubiquitous intracellular CD8α protein expression. This finding suggests that most NK

cells are capable of presenting CD8α to the cell surface to enhance effector function, in contrast

to constitutive surface CD8α expression on T cells. Moreover, CD8α^+ NK cells were exquisitely

sensitive to cytokine stimulation, suggesting complex roles during inflammatory immune

responses against disease. Further, we show that CD8α surface expression was strongly dependent

on MHC I expression (*50, 54, 72*) raising critical questions for future studies to define the

molecular mechanisms of CD8α engagement with MHC I and to determine whether this may be

modulated by different peptide-MHC I complexes. Taken together with previously published work

(*47-49*), CD8α⁺ NK cell subpopulations emerge as an attractive target in vaccination and host-

directed approaches against infectious and noninfectious diseases (*73*).

In addition to these mechanistic insights, we present the first multi-omic analysis of CD8α⁺ NK

cells and show that in contrast to T cells, CD8α does not mark a distinct transcriptional state of

NK cells in healthy donors assayed, though *CD8A⁺* NK cells did differentially express cytotoxic

genes including *NCR3* and *GZMM*. These results will need to be validated in other healthy donors

as well as compared to future CITE-seq experiments in high-risk *Mtb* exposure cohorts or other

disease states.

There are certain limitations inherent to our human immunology study design. While household

contacts experienced high risk exposures to pre-treatment smear+ active TB cases, this does not

exclude the possibility they may have been exposed or infected in the community. We control for

this using source community donors to account for background transmission. Similarly, IGRA-

community controls may have also been recently exposed and resisted primary infection, though

this group represents the most rigorous control for household exposure. Our study was conducted

in one TB endemic population in Haiti and will need to be validated in other cohorts. Due to the

largely cross-sectional study design, we were unable to follow study volunteers beyond 6 months

for IGRA- contacts, of whom only two converted to IGRA+ providing insufficient sample size to

analyze correlates of IGRA conversion prospectively. Finally, all our biological samples in TB

cohorts were derived from human blood and may not represent the abundance or function of NK

cells within infected lung tissue. Future studies in *Mtb*-infected donors with multi-omic validation

are needed to understand the role of CD8α⁺ NK cells at primary sites of infection.

In sum, our study adds to the growing literature highlighting a protective role for CD8α⁺ NK cells

against both infections and non-infectious immune-mediated disease. Ongoing work will define

how surface CD8αα⁺ NK cells may enhance targeting of *Mtb*-infected cells and will inform future

cellular therapies and vaccination approaches to prevent or treat TB disease.

MATERIALS AND METHODS

Study design and participant recruitment:

Donor recruitment and protection of human subjects. Haitian donors were enrolled through the

Tri-Institutional Tuberculosis Research Unit (TBRU) at the Groupe Haitien d'etude du Sarcome

de Kaposi et des Infections Opportunistes (GHESKIO) Centers under Institutional Review Board

Approval at Weill Cornell Medicine and GHESKIO Centers in Haiti. All participants provided

written informed consent, including consent for future use of biobanked samples for research. A

dedicated clinical field team at the GHESKIO Centers in Port-au-Prince, Haiti, recruited as part of

the NIH-funded TBRU (AI111143). LTBI was detected using QuantiFERON-TB® Gold

(QIAGEN), and active TB was excluded by clinical screening for signs and symptoms of active

TB including fever, cough, night sweats, weight loss, and abnormal findings on Chest X-ray. Most

contacts were recruited close to the time of diagnosis of the index case and were living in the same

house as the active TB case for at least 1 month in the 6 months prior to diagnosis (median time to

contact recruitment from TB case diagnosis, 2.4 months; range, 0.2-26 months). To select for high

transmission settings, each household was required to have a history of two active pulmonary TB

cases. Healthy donors without reported TB exposure from the same community were recruited as

controls. All cases with active pulmonary TB received periodic follow-up appointments while on

treatment, and anyone with known contact with an active TB patient who remained IGRA-

received 6-month follow-up and was rescreened with IGRA. The median smear grade of the active

pulmonary TB cases in the households was 2-3+, conferring high risk for person-to-person

transmission (*74*). All donors provided informed consent prior to peripheral blood donation for

PBMC isolation. All donor samples were de-identified on site using a barcode system before they

were shipped to Stony Brook University for analysis. Healthy donors were also recruited at Stony

Brook University (SBU) under IRB2021-00478 (PI: Vorkas) and New York Blood Center

(NYBC), which were used for stimulation assays and CITE-seq.

PBMC isolation:

PBMCs were isolated from peripheral blood at GHESKIO (Port-au-Prince, Haiti) using the Ficoll-

Paque (GE Healthcare) density centrifugation, frozen in 5×10^6 cells/ml aliquots in 90% FBS/10%

DMSO (Thermo Fisher Scientific) and stored at –80°C prior to being shipped on dry ice to New

York. PBMCs from healthy donors recruited at SBU and New York Blood Center (NYBC) were

isolated using Ficoll-Paque, BD Vacutainer CPT cell preparation tube (Becton, Dickinson and

Company) or SepMate tubes (Stem Cell Technologies) using the manufacturer protocol. PBMCs

isolated at SBU were cryopreserved in serum free Bambanker medium (GC Lymphotec) and stored

in liquid N_2.

Flow cytometry acquisition and analysis:

A complete list of fluorescent antibodies is listed in Supplementary Table 1: Reagents and

resources. All antibodies were commercially acquired, and multicolor panels were designed and

validated using fluorescent minus one (FMO) controls. MR1 tetramers were obtained from the

NIH Tetramer Core Facility (Atlanta, GA). In brief, cryopreserved PBMCs were thawed, plated,

Fc receptor blocked (eBioscience), and underwent direct ex vivo extracellular staining for 15

minutes at room temperature (RT). For intracellular staining, PBMCs were incubated with 1X

Brefeldin A (BioLegend) for final hour of incubation prior to staining with intracellular antibodies.

Samples were then washed twice with FACS buffer and acquired using an Aurora Spectral

Analyzer (Cytek). Flow cytometry analysis, including manual gating, was performed using FCS

Express 7 (De Novo Software) or R (v4.3.1) with the following library packages downloaded from

Bioconductor: *flowWorkspace, ggcyto, and flowCore.*

***Mtb* whole cell lysate:**

Mtb whole cell lysate was prepared in Biosafety Level 3 facility at SBU. Fifty mL of *Mtb* Erdman

7H9 culture was grown to OD_{600} 0.6, centrifuged at 4000 × g for 10 min at RT and pellet was

resuspended in 2.5 ml of 10mM Tris-HCl buffer (pH 7.5) and rotated for 2 hr at RT. The mixture

was centrifuged at 500 × g for 15 min at RT and aqueous phase was collected and filtered twice

through 0.22 µM spin filter (Corning Costar Spin-X) at 6000 × g for 5 min. The aliquots were

stored at -80°C for further use.

NK cell co-culture with K562 cell line

K562 cells (ATCC) were incubated in Iscove's Modified Dulbecco's Medium (IMDM)

supplemented with 10% FBS. Cryopreserved K562 cells were thawed at 37°C water bath and cells

were cultured at 1 x 10^5 viable cells/ml in T-75 flask as per manufacturers protocol. Once the cell

density reached 1 x 10^6 cells/ml, K562 was co-incubated with PBMCs at 1:20 K562 to PBMC in

a round bottom clear plate (Greiner) for 15 hrs, at 37°C with 5% CO_2. At the beginning of

incubation period anti-CD107a antibody was added to the culture. After incubation was complete,

cells were washed, stained and acquired by flow cytometry.

NK cell stimulation assay:

Cryopreserved healthy donor PBMCs from SBU and NYBC were used for NK cell stimulation

assays. Thawed PBMCs were cultured 1 x 10^4 cells per well in round bottom clear plate (Greiner)

for 15 hr, at 37°C with 5% CO_2. Samples were stimulated and incubated overnight in complete

RPMI media with 10% heat-inactivated FBS and supplemented with IL-2 (50 U/ml; PeproTech).

The cytokine stimulation conditions were performed with commercially available recombinant IL-

12 (100 ng/mL; BioLegend), recombinant IL-18 (100 ng/mL; Biolegend) and IL15-Su construct

(1 µg/ml) containing a truncated Sushi domain of IL15Rα synthesized by the Cheung lab, that is

demonstrated to be a more potent and stable compound with longer half-life than recombinant

IL15 (*52*). For *Mtb* stimulation condition we used 1:100 and 1:200 dilution of *Mtb* whole cell

lysate.

MHC I blockade:

Cryopreserved healthy donor PBMCs from SBU and NYBC were used for MHC I blockade.

PBMCs were thawed at 37°C and incubated in complete RPMI media with 10% heat-inactivated

FBS and supplemented with IL-2 (50 U/ml; PeproTech). 1×10^4 PBMCs were plated per well in

round bottom clear plate (Greiner) and incubated with or without 5 µg/ml *Invivo*MAb anti-human

MHC I (clone W6/32; Bio X cell) for 15 hr, at 37°C with 5% CO_2 before being stained and

acquired.

scRNA-seq and CITE-seq library generation and sequencing:

PBMCs were stained with fluorescent antibodies for cell sorting of MAIT cells (CD3[+], CD161[+],

MR1-5-OP-RU tetramer[+]), NK cells (CD3[-], CD19[-], CD14[-], CD16[+] CD56[dim]), γδ T cells (CD3[+],

TCRγδ[+]) and iNKT cells (CD161[+] iNKT[+]) using FACS. The sorted cells were blocked with human

Trustain, FcX blocking reagent (BioLegend) for 10 minutes at 4°C and then stained with TotalSeq-

C barcoded tagged and hashtags antibodies for 30 minutes at 4°C in a total volume of 100ul. The

residual TotalSeq-C and hashtags antibodies were removed by centrifugation at 400 *g* for 5

minutes at 4°C. A complete hashtag and TotalSeq-C antibody list is provided in **Supplementary**

Table 1: Reagent and Resources.

Gene expression and surface antigen library preparation was conducted at the Stony Brook University Single Cell Genomics Facility. Briefly, cell viability and concentrations were determined on a Countess 3FL with trypan blue staining. Approximately 5,000 live cells from each condition were loaded in a single 10X Genomics 5' NextGEM Single Cell assay reaction (30,000 total cells per reaction), to perform 5' single cell immune profiling assay using a Chromium iX instrument (10X Genomics). Gene expression (GEX) and surface antigen (SA) libraries were prepared, and quality controlled using Chromium NextGEM 5' Single Cell with Feature Barcode assay and a TapeStation 4200, according to the manufacturer's instructions. The libraries were sequenced at a depth of approx. 25,000 (GEX) and 10,000 (SA) paired reads/cell on an Illumina NovaSeqX sequencer (Novogene, Inc).

The gene expression and surface feature barcode raw FASTQ files of single cell RNA-seq were analyzed on Cell Ranger (10x Genomics, v.7.0) (https://www.10xgenomics.com/analysis-guides/demultiplexing-and-analyzing-5%E2%80%99-immune-profiling-libraries-pooled-with-hashtags), multi-pipeline with human reference transcriptome (GRCh38) downloaded from 10x Genomics website and hashtag configuration file, which contained the barcodes related to each different samples.

Single-cell RNAseq analysis:

The generated gene expression matrix files were analyzed with R (4.3.2) using the *Seurat* (5.0) package and cells were filtered based on following criteria: 1.) the number of genes was equal to or greater than 200 and below than 6000 and 2.) the mitochondrial percentage was less than 5. Using the FindVariableFeatures() function we selected the top 3000 features with highest variance. Then, different samples were integrated using the harmony algorithm and scaled. Log-normalization and principal component analysis (PCA) was done on integrated values. The top 20

principal components were selected for clustering and visualization with UMAP using the default

Louvain algorithm. Differential gene expression was performed using FindAllMarkers() with a

minimum threshold of 25% cells expressing.

Statistical Analysis:

Statistical analysis was performed using R (4.3.2) and GraphPad Prism (10.1.2). Non-parametric

testing with Wilcoxon tests was used to compare the means between groups for flow markers. P-

values were not adjusted because cross-wise comparisons were not performed between all six

clinical groups. Only comparisons a priori hypothesized to represent biological features were

tested, these included: IGRA- contacts vs controls, IGRA+ contacts vs controls, IGRA- contact vs

IGRA+ control, and IGRA+ control vs Active TB.

Supplementary Materials

<u>Supplementary Table 1: Reagents and resources</u>. See attached file.

References and Notes:

1.   R. M. Houben, P. J. Dodd, The Global Burden of Latent Tuberculosis Infection: A Re-

estimation Using Mathematical Modelling. *PLoS Med* **13**, e1002152 (2016).

2.   S. Bagcchi, WHO's Global Tuberculosis Report 2022. *Lancet Microbe* **4**, e20 (2023).

3.   K. M. Shea, J. S. Kammerer, C. A. Winston, T. R. Navin, C. R. Horsburgh, Jr., Estimated

rate of reactivation of latent tuberculosis infection in the United States, overall and by

population subgroup. *Am J Epidemiol* **179**, 216-225 (2014).

4.   K. B. Urdahl, S. Shafiani, J. D. Ernst, Initiation and regulation of T-cell responses in

tuberculosis. *Mucosal Immunol* **4**, 288-293 (2011).

5.   E. Nemes *et al.*, Prevention of M. tuberculosis Infection with H4:IC31 Vaccine or BCG

Revaccination. *N Engl J Med* **379**, 138-149 (2018).

6.   D. R. Tait *et al.*, Final Analysis of a Trial of M72/AS01E Vaccine to Prevent Tuberculosis.

N Engl J Med **381**, 2429-2439 (2019).

7.   A. Nathan *et al.*, Multimodally profiling memory T cells from a tuberculosis cohort

identifies cell state associations with demographics, environment and disease. *Nat Immunol*

22, 781-793 (2021).

8.   A. Pradhan, C. K. Vorkas, T cells target TB. *Cell Host Microbe* **31**, 329-330 (2023).

9.   A. M. Gallegos *et al.*, A gamma interferon independent mechanism of CD4 T cell mediated

control of M. tuberculosis infection in vivo. *PLoS Pathog.* **7**, e1002052 (2011).

10.   S. Sakai *et al.*, CD4 T Cell-Derived IFN-gamma Plays a Minimal Role in Control of

Pulmonary Mycobacterium tuberculosis Infection and Must Be Actively Repressed by PD-

1 to Prevent Lethal Disease. *PLoS Pathog.* **12**, (2016).

11.   H. A. Fletcher, L. Schrager, TB vaccine development and the End TB Strategy: importance

and current status. *Trans R Soc Trop Med Hyg* **110**, 212-218 (2016).

12.   G. Voss *et al.*, Progress and challenges in TB vaccine development. *F1000Res* **7**, 199

(2018).

13.   J. D. Ernst, Mechanisms of M. tuberculosis Immune Evasion as Challenges to TB Vaccine

Design. *Cell Host Microbe* **24**, 34-42 (2018).

14.   D. Roordink *et al.*, The TB vaccine development pathway - An innovative approach to

accelerating global TB vaccine development. *Tuberculosis (Edinb)* **126**, 102040 (2021).

15.   C. K. Vorkas *et al.*, Mucosal-associated invariant and gammadelta T cell subsets respond

to initial Mycobacterium tuberculosis infection. *JCI Insight* **3**, (2018).

16.   J. A. Choreno-Parra, L. I. Weinstein, E. J. Yunis, J. Zuniga, R. Hernandez-Pando, Thinking

Outside the Box: Innate- and B Cell-Memory Responses as Novel Protective Mechanisms

Against Tuberculosis. *Front Immunol* **11**, 226 (2020).

17.   S. A. Khader *et al.*, Targeting innate immunity for tuberculosis vaccination. *J Clin Invest*

129, 3482-3491 (2019).

18.   J. A. Choreño Parra *et al.*, Memory of Natural Killer Cells: A New Chance against

Mycobacterium tuberculosis? *Front Immunol* **8**, 967 (2017).

19.   D. Artis, H. Spits, The biology of innate lymphoid cells. *Nature* **517**, 293-301 (2015).

20.   P. Dogra *et al.*, Tissue Determinants of Human NK Cell Development, Function, and

Residence. *Cell* **180**, 749-763.e713 (2020).

21.   L. L. Lanier, Natural killer cell receptor signaling. *Current opinion in immunology* **15**, 308-

314 (2003).

22.   P. Schierloh *et al.*, Mycobacterium tuberculosis-induced gamma interferon production by

natural killer cells requires cross talk with antigen-presenting cells involving Toll-like

receptors 2 and 4 and the mannose receptor in tuberculous pleurisy. *Infect Immun* **75**, 5325-

5337 (2007).

23.   S. Venkatasubramanian *et al.*, IL-21-dependent expansion of memory-like NK cells

enhances protective immune responses against Mycobacterium tuberculosis. *Mucosal*

Immunol **10**, 1031-1042 (2017).

24.   S. Esin, G. Batoni, Natural killer cells: a coherent model for their functional role in

Mycobacterium tuberculosis infection. *J Innate Immun* **7**, 11-24 (2015).

25.   C. B. W. Giacoia-Gripp *et al.*, Changes in the NK Cell Repertoire Related to Initiation of

TB Treatment and Onset of Immune Reconstitution Inflammatory Syndrome in TB/HIV

Co-infected Patients in Rio de Janeiro, Brazil-ANRS 12274. *Front Immunol* **10**, 1800
(2019).

26. M. Allen *et al.*, Mechanisms of Control of Mycobacterium tuberculosis by NK Cells: Role
of Glutathione. *Front Immunol* **6**, 508 (2015).

27. M. Garand *et al.*, Functional and Phenotypic Changes of Natural Killer Cells in Whole
Blood during Mycobacterium tuberculosis Infection and Disease. *Front Immunol* **9**, 257
(2018).

28. L. D. Harris *et al.*, Distinct Human NK Cell Phenotypes and Functional Responses to
Mycobacterium tuberculosis in Adults From TB Endemic and Non-endemic Regions.
Front Cell Infect Microbiol **10**, 120 (2020).

29. G. Batoni *et al.*, Human CD56bright and CD56dim natural killer cell subsets respond
differentially to direct stimulation with Mycobacterium bovis bacillus Calmette-Guérin.
Scand J Immunol **62**, 498-506 (2005).

30. S. Esin *et al.*, Interaction of Mycobacterium tuberculosis cell wall components with the
human natural killer cell receptors NKp44 and Toll-like receptor 2. *Scand J Immunol* **77**,
460-469 (2013).

31. S. Esin *et al.*, Functional characterization of human natural killer cells responding to
Mycobacterium bovis bacille Calmette-Guérin. *Immunology* **112**, 143-152 (2004).

32. R. Vankayalapati *et al.*, Role of NK cell-activating receptors and their ligands in the lysis
of mononuclear phagocytes infected with an intracellular bacterium. *J Immunol* **175**, 4611-
4617 (2005).

33. R. Vankayalapati *et al.*, The NKp46 receptor contributes to NK cell lysis of mononuclear
phagocytes infected with an intracellular bacterium. *J Immunol* **168**, 3451-3457 (2002).

34.   D. Tripathi *et al.*, Metabolites enhance innate resistance to human Mycobacterium

tuberculosis infection. *JCI Insight* **7**, (2022).

35.   M. E. Ross, M. A. Caligiuri, Cytokine-induced apoptosis of human natural killer cells

identifies a novel mechanism to regulate the innate immune response. *Blood* **89**, 910-918

(1997).

36.   W. E. Carson *et al.*, Interleukin (IL) 15 is a novel cytokine that activates human natural

killer cells via components of the IL-2 receptor. *J Exp Med* **180**, 1395-1403 (1994).

37.   S. Suliman *et al.*, Peripheral Blood Mucosal-Associated Invariant T Cells in Tuberculosis

Patients and Healthy Mycobacterium tuberculosis-Exposed Controls. *J Infect Dis* **222**, 995-

1007 (2020).

38.   P. Schierloh *et al.*, Increased susceptibility to apoptosis of CD56dimCD16+ NK cells

induces the enrichment of IFN-gamma-producing CD56bright cells in tuberculous

pleurisy. *J Immunol* **175**, 6852-6860 (2005).

39.   B. Lucia, C. Jennings, R. Cauda, L. Ortona, A. L. Landay, Evidence of a selective depletion

of a CD16+ CD56+ CD8+ natural killer cell subset during HIV infection. *Cytometry* **22**,

10-15 (1995).

40.   K. Srpan *et al.*, Shedding of CD16 disassembles the NK cell immune synapse and boosts

serial engagement of target cells. *J Cell Biol* **217**, 3267-3283 (2018).

41.   I. Terren *et al.*, Cytokine-Induced Memory-Like NK Cells: From the Basics to Clinical

Applications. *Front Immunol* **13**, 884648 (2022).

42.   D. Brownlie *et al.*, Expansions of adaptive-like NK cells with a tissue-resident phenotype

in human lung and blood. *Proc Natl Acad Sci U S A* **118**, (2021).

43.   R. Nikzad *et al.*, Human natural killer cells mediate adaptive immunity to viral antigens.

Sci Immunol **4**, (2019).

44.   N. M. Adams *et al.*, Cytomegalovirus Infection Drives Avidity Selection of Natural Killer

Cells. *Immunity* **50**, 1381-1390 e1385 (2019).

45.   S. Suliman *et al.*, Bacillus Calmette-Guérin (BCG) Revaccination of Adults with Latent

Mycobacterium tuberculosis Infection Induces Long-Lived BCG-Reactive NK Cell

Responses. *J Immunol* **197**, 1100-1110 (2016).

46.   K. Van Der Ploeg *et al.*, Emergence of Human CMV-induced NKG2C+ NK cells is

Associated with CD8+ T cell Recovery following Allogeneic HCT. *Blood Adv*, (2023).

47.   E. F. McKinney *et al.*, A CD8(+) NK cell transcriptomic signature associated with clinical

outcome in relapsing remitting multiple sclerosis. *Nat Commun* **12**, 635 (2021).

48.   E. G. Addison *et al.*, Ligation of CD8alpha on human natural killer cells prevents

activation-induced apoptosis and enhances cytolytic activity. *Immunology* **116**, 354-361

(2005).

49.   F. Ahmad *et al.*, High frequencies of polyfunctional CD8+ NK cells in chronic HIV-1

infection are associated with slower disease progression. *J Virol* **88**, 12397-12408 (2014).

50.   M. W. Lowdell *et al.*, Evidence that continued remission in patients treated for acute

leukaemia is dependent upon autologous natural killer cells. *Br J Haematol* **117**, 821-827

(2002).

51.   N. K. Björkström, H.-G. Ljunggren, J. K. Sandberg, CD56 negative NK cells: origin,

function, and role in chronic viral disease. *Trends in Immunology* **31**, 401-406 (2010).

52.   H. Xu, I. N. Buhtoiarov, H. Guo, N. V. Cheung, A novel multimeric IL15/IL15Rα-Fc

complex to enhance cancer immunotherapy. *Oncoimmunology* **10**, 1893500 (2021).

53.   L. J. Walker *et al.*, Human MAIT and CD8alphaalpha cells develop from a pool of type-
17 precommitted CD8+ T cells. *Blood* **119**, 422-433 (2012).

54.   H. Cheroutre, F. Lambolez, Doubting the TCR coreceptor function of CD8alphaalpha.
Immunity **28**, 149-159 (2008).

55.   K. Kushima *et al.*, Flow cytometric analysis of chicken NK activity and its use on the effect
of restraint stress. *J Vet Med Sci* **65**, 995-1000 (2003).

56.   M. Gutierrez-Arcelus *et al.*, Lymphocyte innateness defined by transcriptional states
reflects a balance between proliferation and effector functions. *Nat Commun.* **10**, 687
(2019).

57.   C. Yang *et al.*, Heterogeneity of human bone marrow and blood natural killer cells defined
by single-cell transcriptome. *Nat Commun* **10**, 3931 (2019).

58.   M. Musvosvi *et al.*, T cell receptor repertoires associated with control and disease
progression following Mycobacterium tuberculosis infection. *Nat Med*, (2023).

59.   S. Suliman *et al.*, MR1-Independent Activation of Human Mucosal-Associated Invariant
T Cells by Mycobacteria. *J Immunol.* **203**, 2917-2927 (2019).

60.   P. Andersen, T. J. Scriba, Moving tuberculosis vaccines from theory to practice. *Nat Rev*
Immunol **19**, 550-562 (2019).

61.   R. Roy Chowdhury *et al.*, A multi-cohort study of the immune factors associated with M.
tuberculosis infection outcomes. *Nature* **560**, 644-648 (2018).

62.   D. Masopust, V. Vezys, E. J. Wherry, R. Ahmed, A brief history of CD8 T cells. *Eur J*
Immunol **37 Suppl 1**, S103-110 (2007).

63.   L. J. Kieffer *et al.*, Human CD8 alpha expression in NK cells but not cytotoxic T cells of
transgenic mice. *Int Immunol* **8**, 1617-1626 (1996).

64.   B. Muller-Durovic, J. Grahlert, O. P. Devine, A. N. Akbar, C. Hess, CD56-negative NK

cells with impaired effector function expand in CMV and EBV co-infected healthy donors

with age. *Aging (Albany NY)* **11**, 724-740 (2019).

65.   A. Poli *et al.*, CD56bright natural killer (NK) cells: an important NK cell subset.

Immunology **126**, 458-465 (2009).

66.   A. S. Fauci, D. Mavilio, S. Kottilil, NK cells in HIV infection: paradigm for protection or

targets for ambush. *Nat Rev Immunol* **5**, 835-843 (2005).

67.   N. K. Björkström, H. G. Ljunggren, J. K. Sandberg, CD56 negative NK cells: origin,

function, and role in chronic viral disease. *Trends Immunol* **31**, 401-406 (2010).

68.   B. Rehermann, M. Nascimbeni, Immunology of hepatitis B virus and hepatitis C virus

infection. *Nat Rev Immunol* **5**, 215-229 (2005).

69.   L. L. Lanier, R. Testi, J. Bindl, J. H. Phillips, Identity of Leu-19 (CD56) leukocyte

differentiation antigen and neural cell adhesion molecule. *J Exp Med* **169**, 2233-2238

(1989).

70.   L. A. Terry, J. P. DiSanto, T. N. Small, N. Flomenberg, Differential expression and

regulation of the human CD8 alpha and CD8 beta chains. *Tissue Antigens* **35**, 82-91 (1990).

71.   R. L. Webster, R. P. Johnson, Delineation of multiple subpopulations of natural killer cells

in rhesus macaques. *Immunology* **115**, 206-214 (2005).

72.   J. Geng, M. Raghavan, CD8alphaalpha homodimers function as a coreceptor for

KIR3DL1. *Proc Natl Acad Sci U S A* **116**, 17951-17956 (2019).

73.   S. Schmidt, L. Tramsen, B. Rais, E. Ullrich, T. Lehrnbecher, Natural killer cells as a

therapeutic tool for infectious diseases - current status and future perspectives. *Oncotarget*

9, 20891-20907 (2018).

74.   H. del Corral *et al.*, IFNgamma response to Mycobacterium tuberculosis, risk of infection

and disease in household contacts of tuberculosis patients in Colombia. *PLoS One* **4**, e8257

(2009).

Acknowledgments: We are thankful for the support of the Stony Brook Foundation, the

Department of Medicine of the Renaissance School of Medicine and the Office of the Vice

President of Research of Stony Brook University. We thank Daniel W. Fitzgerald, MD

(Weill Cornell Medicine, Center for Global Health) and Michael S. Glickman (Memorial

Sloan Kettering Cancer Center) for generously providing clinical samples from TB contact

cohorts. We thank Joseph C. Sun (MSKCC) for valuable discussions on adaptive NK cells.

We are also grateful to Thomas MacCarthy, PhD (Stony Brook University, Applied

Mathematics) for supportive advice regarding CITE-seq analyses, who passed away during

the preparation of this manuscript.

Funding: This study was supported by the Tri-I TBRU, part of the TBRU Network (U19

AI11143). CKV acknowledges support from the NIAID T32 Pathogenesis of Infectious

Diseases Training Program (T32AI007613-18; Roy Gulick, WCM), the IDSA Education

and Research Foundation Postdoctoral Fellowship Award in Infectious Diseases, a Potts

Memorial Foundation Award, NIAID K08AI132739, R21AI171578, and R21AI183259.

Author contributions: The following authors contributed to the manuscript in various ways. Each

author's contribution is listed below.

Conceptualization: KH, JBL, CKV

Methodology: NKC, AP, RK, KCH, JBL, CKV

Investigation: NM, AP, RK, TK, BR, NKC, HX, JBL and CKV

627   Visualization: NM, AP, RK, JB, BR, DEC, JBL, and CKV

628   Funding acquisition: CKV

629   Writing – original draft: NM, AP, CKV

630   Writing – review & editing: all authors

631 **Competing interests:** Authors declare that they have no competing interests.

632 **Data and materials availability:** All data, code, and materials used in the analysis will be

633   available to any researcher for purposes of reproducing or extending the analysis.

635 **Figures:**

Variable	Control (n=50)	Contact (n=50)	Active TB (n=50)
Age – yr	30.1 ± 9.3	33.9 ± 17.8	20.2 ± 4.26
≤ 18 yr	2 (4)	12 (24)	18 (36)
>18 and <65	48 (96)	35 (76)	32 (64)
≥ 65 yr	0 (0)	3	0 (0)
Female sex – no. (%)	26 (52)	30 (60)	26 (52)
IGRA positive – no. (%)	25 (50)	34 (68)	38 (76)

636 **Table 1. Demographics of study participants in Haitian Cohort (N=150)**

Fig. 1. Frequencies of NK cell subpopulations are altered after *Mtb* infection. (A) Gating strategy employed for

NK cells from a representative sample. NK cells were defined as CD19⁻, CD3⁻, CD14⁻ lymphocytes who

expressed any level of either CD16 or CD56. NK cells were further subdivided into three subsets based on

CD56 expression. (B) Change in frequency of total NK cells as a proportion of total live population. (C)

Frequency of each NK cell subset in clinical groups. Frequency calculated as percent of each subset among

total NK cells. Brackets represent Wilcoxon statistical tests, with unadjusted p-values. *p ≤ 0.05, **p ≤

0.01.

644 **Fig. 2. NK cells demonstrate loss of CD16 expression during household-associated latent infection. (A)**

645 Frequency of CD16 in CD56^{dim} NK cells. Brackets represent Wilcoxon tests with unadjusted p-values.

646 Percent positive for 5 activation markers in **(B)** CD56^{bright} NK cells, **(C)** CD56^{dim} NK cells, and **(D)**

647 CD56^{-} NK cells. Brackets represent Wilcoxon tests with unadjusted p-values. *p ≤ 0.05, ****p ≤

648 0.0001.

Fig. 3. *Mtb* resistors demonstrate depressed NK cell function in vitro. NK cells stimulated overnight by co-culture with K562 cell line compared to plain media alone. **(A)** Overall results of activation assay for CD56^{dim} NK cells for four activation markers. Y-axis represents mean percent positive for each activation marker. P values calculated as a global paired Wilcoxon test including all clinical groups. Each clinical group is displayed separately indicated by line color. **(B)** Percent difference of four activation markers (stimulated minus resting) comparing between clinical groups. Brackets represent Wilcoxon tests with unadjusted p-values. *p ≤ 0.05, **p ≤ 0.01, ***p ≤ 0.001, ****p ≤ 0.0001.

Fig. 4. CD8α⁺ NK cells are depleted in the blood of cases with household-associated *Mtb* infection. (A) Representative gating of CD8α⁺ in CD56^{dim} NK cells and frequency of CD8α⁺ within CD56^{dim} NK cells. **(B)** Representative gating of CD57 in CD56^{dim} NK cells and boxplots of percent CD57⁺ within CD56^{dim} NK cells. **(C)** Expression of select surface markers between CD8α⁺ vs CD8α⁻ CD56^{dim} NK cells, active TB are excluded. **(D)** Activation assay with overnight K562 co-culture versus plain media comparing CD8α⁺ vs CD8α⁻ CD56^{dim} NK cells. Y-values represent percent difference for CD69 and CD107a. Asterisks represent paired Wilcoxon tests. **(E)** As in panel D, instead displaying CD56⁻ NK cells. P-values represent unadjusted Wilcoxon testing. *p ≤ 0.05, **p ≤ 0.01, ***p ≤ 0.001, ****p ≤ 0.0001.

Fig. 5. CD8α⁺ NK cells are prevalent in healthy US donors with ubiquitous intracellular CD8α expression.

(A) Flow cytometry plot and percentage of CD8αα⁻, CD8αα⁺, CD8αβ⁺, and CD8ββ⁺ among total NK (N=3). (B) Percentage of CD8α⁺ NK cells in different subsets based on CD56 expression (N=10). (C) Frequency of CD16 expression in CD8α⁺ CD56^{dim} NK cells. (D) Frequency and representative flow cytometry plots of CD8α expression comparing CD56^{dim} NK cells and T cells with extracellular (EC), IC (Intracellular), and EC + IC CD8α staining. (E) Percent CD8α⁺ in EC, IC, and EC + IC staining at day 1 and day 7 in CD56^{dim} NK cells and T cells. Statistical significance was measured using unpaired t tests, ns – no significance, *$p<0.05$, **$p<0.005$, ***$p<0.001$, ****$p<0.0001$.

Fig. 6. CD8α⁺ CD56^{dim} NK cells have enhanced response to cytokine stimulation. (**A**) Pairwise analysis of

percent CD8α⁺ NK cells upon stimulation with IL15/12/18 and MtbLys 1:200 (n=9). (**B**) Pairwise analysis

and representative flow plots of intracellular IFNγ in NK CD56^{dim} upon stimulation with IL15/12/18 and

MtbLys at dilution 1:200 (n=9). (**C**) Percent intracellular IFNγ⁺ in CD8α⁺ versus CD8α⁻ CD56^{dim} NK cells

comparing IL-15-Su, IL15-Su/12/18, IL-12-18, or MtbLys conditions (n=9). Both the CD16⁺ (right) and

CD16⁻ (left) CD56^{dim} NK subpopulations are displayed. (**D**) Representative flow cytometry plots and paired

analysis of percent change CD8α⁺ with anti-MHC I for 15 hours among CD56^{dim} NK cells (left) and CD8⁺

T cells (right) (n=8). Statistical significance was measured using paired t-test, with threshold of *p<0.05,

**p<0.005, ***p<0.001. IFNγ: Interferon γ, Mtb Lys: *Mtb* whole cell lysate.

Fig. 7. CITE-seq analysis of CD8α⁺ vs CD8α⁻ NK cells from two healthy donors reveals overlapping

transcriptomes. (A) Results of clustering and uniform manifold approximation (UMAP) projection for all

NK cells. **(B)** UMAP reduction colored by Donor. **(C)** UMAP reduction colored by condition. **(D)** Violin

plots representing top six genes that were differentially expressed between the four NK cell clusters. **(E)**

Violin plots of the eight differentially expressed genes between resting and *Mtb* lysate condition. **(F)**

Distribution of NK cell expression of CD8 based on barcoded antibodies against CD8α and *CD8A*

transcript. **(G)** Violin plot of the expression of *CD8A* and *CD8B* transcripts, as well as CD8α surface

expression between the four NK cell clusters. **(H)** Correlation of CD8α expression between transcript and

protein measurement. **(I)** Annotated UMAP based on transcript expression of *CD8A* into CD8α⁺ and

CD8α⁻ NK cells. **(J)** Percent of NK cells expressing CD8A transcript between stimulation conditions. **(K)**

Dot-plot of genes differentially expressed between CD8α⁺ and CD8α⁻ NK cells.