Cytosolic S100A8/A9 promotes Ca^{2+} supply at LFA-1 adhesion clusters during neutrophil recruitment

Matteo Napoli[1], Roland Immler[1], Ina Rohwedder[1], Valerio Lupperger[2], Johannes Pfabe[3], Mariano Gonzalez Pisfil[1], Anna Yevtushenko[1], Thomas Vogl[4], Johannes Roth[4], Melanie Salvermoser[1], Steffen Dietzel[1], Marjan Slak Rupnik[3], Carsten Marr[2], Barbara Walzog[1], Markus Sperandio[1] and Monika Pruenster[1]*

[1]Walter Brendel Center of Experimental Medicine, Biomedical Center, Institute of Cardiovascular Physiology and Pathophysiology, Ludwig-Maximilians-University, Planegg-Martinsried, Germany

[2]Institute of AI for Health, Helmholtz Zentrum München - German Research Center for Environmental Health, Neuherberg, Germany

[3]Center for Physiology and Pharmacology, Medical University of Vienna, Vienna, Austria

[4]Institute of Immunology, University of Muenster, Muenster, Germany

*Corresponding author:

Monika Pruenster, PhD

Walter Brendel Center of Experimental Medicine

Institute of Cardiovascular Physiology and Pathophysiology

Biomedical Center Munich

Ludwig-Maximilians-Universität München (LMU)

Großhaderner Str. 9

82152 Planegg-Martinsried

GERMANY

Voice: +49 (0)89 2180 71526

Fax: +49 (0)89 2180 71513

Email: monika.pruenster@med.uni-muenchen.de

ABSTRACT

S100A8/A9 is an endogenous alarmin secreted by myeloid cells during many acute and chronic

inflammatory disorders. Despite increasing evidence of the proinflammatory effects of

extracellular S100A8/A9, little is known about its intracellular function. Here, we show that

cytosolic S100A8/A9 is indispensable for neutrophil post-arrest modifications during outside-

in signaling under flow conditions in vitro and neutrophil recruitment in vivo, independent of

its extracellular functions. Mechanistically, genetic deletion of S100A9 in mice (*Mrp14*[-/-],

functional *S100a8/a9*[-/-]) caused dysregulated Ca^{2+} signatures in activated neutrophils resulting

in reduced Ca^{2+} availability at the formed LFA-1/F-actin clusters with defective β_2 integrin

outside-in signaling during post-arrest modifications. Consequently, we observed impaired

cytoskeletal rearrangement, cell polarization and spreading, as well as cell protrusion

formation in *Mrp14*[-/-] compared to WT neutrophils, making *Mrp14*[-/-] cells more susceptible to

detach under flow, thereby preventing efficient neutrophil recruitment and extravasation into

inflamed tissue.

One-sentence summary: intracellular S100A8/A9 is indispensable for firm leukocyte

adhesion under flow

INTRODUCTION

Neutrophils are the most abundant circulating leukocyte subpopulation in humans and are rapidly mobilized from the bone marrow to the circulation upon sterile inflammation and/or bacterial/viral infection [1]. The interplay between activated endothelial cells and circulating neutrophils leads to a tightly regulated series of events described as leukocyte recruitment cascade [2]. Tissue-derived proinflammatory signals provoke expression of selectins on the inflamed endothelium that capture free floating neutrophils from the bloodstream by triggering tethering and rolling through interaction with selectin ligands on the neutrophil surface [3]. Selectin mediated rolling allows neutrophils to engage with immobilized chemokines and other proinflammatory mediators such as leucotriene B4 (LTB4), N-formylmethionyl-leucyl-phenylalanine (fMLF) and various agonists for Toll-like receptors (TLRs) like TLR2, TLR4 and TLR5, presented on the endothelial surface and resulting in the activation of β_2 integrins on neutrophils [4-7]. High affinity β_2 integrin interaction with their corresponding receptors on the endothelium induces downstream outside-in signaling leading to post-arrest modifications such as cell spreading, adhesion strengthening and neutrophil crawling, critical requirements for successful recruitment of neutrophils into inflamed tissue [8, 9]. Accordingly, impairment in those steps favors neutrophil detachment under shear flow and re-entry of neutrophils into the blood stream [5].

Local regulation of intracellular calcium (Ca^{2+}) levels is critical to synchronize rolling, arrest and polarization [8, 10]. During rolling, neutrophils show only minor Ca^{2+} activity, but a rapid increase in intracellular Ca^{2+} signaling is registered during transition from slow rolling to firm adhesion and subsequent polarization of neutrophils [11].

Neutrophil transition from rolling into firm arrest involves conformational changes of the

integrin lymphocyte function-associated antigen (LFA-1) into a high affinity state allowing

bond formation with intercellular adhesion molecule-1 (ICAM-1) expressed on inflamed

endothelium. Tension on focal clusters of LFA-1/ICAM-1 bonds mediated by shear stress

promotes the assembly of cytoskeletal adaptor proteins to integrin tails and mediates Ca^{2+}-

release activated (CRAC) channel ORAI-1 recruitment to focal adhesion clusters ensuring high

Ca^{2+} concentrations at the "inflammatory synapse" [10]. Finally, shear stress mediated local

bursts of Ca^{2+} promote assembly of the F-actin cytoskeleton allowing pseudopod formation

and transendothelial migration (TEM) [8, 10, 12-14].

S100A8/A9, also known as MRP8/14 or calprotectin, is a member of the EF-hand family of

proteins and the most abundant cytosolic protein complex in neutrophils [15]. Secretion of

S100A8/A9 can occur via passive release of the cytosolic protein due to cellular necrosis or

neutrophil extracellular trap (NET) formation [16]. Active release of S100A8/A9 without cell

death can be induced by the interaction of L-selectin/PSGL-1 with E-selectin during neutrophil

rolling on inflamed endothelium [6, 17, 18]. We have recently shown that E-selectin induced

S100A8/A9 release occurs through a NLRP3 inflammasome dependent pathway involving

GSDMD pore formation. Pore formation is a time-limited and transient process, which is

reversed by the activation of the ESCRT-III machinery membrane repair mechanism [19]. Once

released, the protein acts as an alarmin, exerting its proinflammatory effects on different cell

types like endothelial cells, lymphocytes and neutrophils [16, 20].

In the present study, we focused on the cytosolic function of S100A8/A9 in neutrophils. We

demonstrate its unique role in supplying Ca^{2+} at LFA-1 adhesion clusters during neutrophil

recruitment thereby orchestrating Ca^{2+} dependent post-arrest modifications, which are

critical steps for subsequent transmigration and extravasation of these cells into inflamed

tissues.

RESULTS

Cytosolic S100A8/A9 promotes leukocyte recruitment in vivo regardless of extracellular

S100A8/A9 functions

As demonstrated previously by our group, rolling of neutrophils on inflamed endothelium

leads to E-selectin mediated, NLRP3 inflammasome dependent, secretion of S100A8/A9 via

transient GSDMD pores [19]. Released S100A8/A9 heterodimer in turn binds to TLR4 on

neutrophils in an autocrine manner, leading to β_2 integrin activation, slow leukocyte rolling

and firm neutrophil adhesion [6]. Interestingly, E-selectin-triggered S100A9/A9 release does

not substantially affect the cytosolic S100A8/A9 content. Analysis of S100A8/A9 levels in the

supernatants of E-selectin versus Triton X-100 treated neutrophils demonstrated that only

about 1-2% of the cytosolic S100A8/A9 content was secreted to the extracellular

compartment (Fig. 1A). In addition, immunofluorescence analysis of the inflamed cremaster

muscle tissue confirmed no major difference in the amount of cytosolic S100A8/A9 between

intravascular and extravasated neutrophils (Fig S1A and S1B). Given the abundance of

cytosolic S100A8/A9 even after its active release during neutrophil rolling, we wanted to

investigate a putative role of intracellular S100A8/A9 in leukocyte recruitment independently

of its extracellular function.

To investigate this, we made use of WT and *Mrp14*$^{-/-}$ mice, which are functional double

knockout mice for MRP8 and MRP14 (S100A8 and S100A9) at the protein level [21], and

studied neutrophil recruitment in mouse cremaster muscle venules upon TNF-α treatment

(Fig. 1B), a well-established model to assess neutrophil recruitment into inflamed tissue in vivo

[22]. Two hours after onset of inflammation, we exteriorized the cremaster muscle and

investigated the number of rolling and adherent cells by intravital microscopy. While rolling

was not affected by the absence of S100A8/A9 (Fig. 1C), we detected a reduced number of

adherent neutrophils in postcapillary cremaster muscle venules of $Mrp14^{-/-}$ compared to WT

mice (Fig. 1D). We found a significant negative correlation between increasing shear rates and

the number of adherent leukocytes in $Mrp14^{-/-}$ animals while this correlation could not be

detected in WT mice (Fig. 1E). These findings indicate that lack of cytosolic S100A8/A9 impairs

shear stress resistance of adherent neutrophils in vivo. To exclude differences in surface

expression of rolling and adhesion relevant molecules on neutrophils, we performed FACS

analysis and could not detect differences in the baseline expression of CD11a, CD11b, CD18,

CD62L, PSGL1, CXCR2 and CD44 in WT and $Mrp14^{-/-}$ neutrophils (Fig. S1C - S1J). In order to test

whether the observed phenotype of decreased neutrophil adhesion in $Mrp14^{-/-}$ mice was

simply a consequence of the lack of extracellular S100A8/A9 induced β_2 integrin activation, we

again performed intravital microscopy in the exteriorized but otherwise unstimulated mouse

cremaster muscles. In this scenario, only a mild inflammation is induced which leads to the

mobilization of pre-stored P-selectin from Weibel-Pallade bodies, but no upregulation of E-

selectin and therefore no E-selectin induced S100A8/A9 release [6]. After exteriorization and

trauma-induced induction of inflammation in the cremaster muscle tissue, we systemically

injected soluble S100A8/A9 via a carotid artery catheter to induce TLR4 mediated integrin

activation and firm leukocyte adhesion in exteriorized cremaster muscle venules (Fig. 1F) [6].

To prevent S100A8/A9 tetramerization in plasma, which would abolish binding of S100A8/A9

to TLR4 [23], we took advantage of a mutant S100A8/A9 protein (S100A8/A9^{mut}, aa exchange

N70A and E79A) which is unable to tetramerize upon Ca^{2+} binding thereby inducing substantial

TLR4 downstream signaling [24, 25]. Injection of S100A8/A9^{mut} induced a significant increase

in leukocyte adhesion in WT mice (Fig. 1G), whereas induction of adhesion was completely

absent in *Mrp14$^{-/-}$* mice (Fig. 1G), suggesting that loss of S100A8/A9 causes an intrinsic

adhesion defect, which cannot be rescued by application of extracellular S100A8/A9 and

subsequent TLR4 mediated β$_2$ integrin activation. In addition, similar results were obtained in

the TNF-α stimulated cremaster muscles model (Fig. S1K) where S100A8/A9^{mut} increased

leukocyte adhesion in WT mice, but again could not induce an increase in leukocyte adhesion

in *Mrp14$^{-/-}$* mice (Fig. S1L). In addition, microvascular parameters were quantified in order to

compare different vessels in every in vivo experiment and no difference was detected (Table

S1).

Further, we wanted to investigate whether reduced adhesion results in impaired leukocyte

extravasation in *Mrp14$^{-/-}$* mice and stained TNF-α stimulated cremaster muscles of WT and

Mrp14$^{-/-}$ mice with Giemsa and analyzed number of perivascular neutrophils. Indeed, we

observed a reduced number of transmigrated neutrophils in *Mrp14$^{-/-}$* compared to WT mice

(Fig. 1H and 1I). Taken together, these data indicate that cytosolic S100A8/A9 regulates key

processes during neutrophil recruitment into inflamed tissue in vivo.

Loss of cytosolic S100A8/A9 impairs neutrophil adhesion under flow conditions without

affecting β$_2$ integrin activation

Next, we focused on the adhesion defect of S100A8/A9 deficient neutrophils. For this purpose,

we used an autoperfused microflow chamber system as described earlier [26]. Flow chambers

were coated with E-selectin, ICAM-1 and CXCL1 (Fig. 2A). This combination of recombinant

proteins mimics the inflamed endothelium and allows studying leukocyte adhesion under flow

conditions. In line with our in vivo findings, lack of S100A8/A9 did not affect leukocyte rolling

(Fig. 2B), but resulted in a lower number of adherent $Mrp14^{-/-}$ leukocytes compared to WT

leukocytes (Fig. 2C), without affecting white blood cell count (WBC) (Table S2). Reduced

neutrophil adhesion could be a consequence of defective β_2 integrin activation induced by

chemokines or other inflammatory mediators [5-7, 27]. In order to study the effect of

S100A8/A9 deficiency on rapid β_2 integrin activation via $G_{\alpha i}$ coupled signaling (inside-out

signaling), we investigated the capacity of WT and $Mrp14^{-/-}$ neutrophils to bind soluble ICAM-

1 upon CXCL1 stimulation using flow cytometry (Fig. 2D). CXCL1 induced a significant and

similar increase in soluble ICAM-1 binding in both, WT and $Mrp14^{-/-}$ neutrophils (Fig. 2E),

suggesting that $G_{\alpha i}$ coupled integrin activation is independent of cytosolic S100A8/A9. To

corroborate this finding, we performed a static adhesion assay where we plated WT and

$Mrp14^{-/-}$ neutrophils on ICAM-1 coated plates, stimulated them with PBS or CXCL1 and

quantified the number of adherent cells. As expected, CXCL1 stimulated WT cells displayed

increased adhesion to ICAM-1 coated plates compared to PBS treatment (Fig. 2F). In line with

the findings from the soluble ICAM-1 binding assay, this increase was also detected in $Mrp14^{-}$

$^{/-}$ cells indicating that chemokine-induced β_2 integrin activation is not dependent on cytosolic

S100A8/A9.

Cytosolic S100A8/A9 is crucial for neutrophil spreading, crawling and post-arrest

modifications under flow

Activated and ligand bound β_2 integrins start to assemble focal clusters thereby transmitting

signals into the inner cell compartment [28]. This process named outside-in signaling is

required to strengthen adhesion and to induce cell shape changes, fundamental for neutrophil

spreading, crawling and finally transmigration [29]. Since $Mrp14^{-/-}$ neutrophils displayed a

defect in leukocyte adhesion in vivo and ex vivo, although their inside-out signaling is fully

functional, we started to study a putative role of cytosolic S100A8/A9 in β_2 integrin dependent

outside-in signaling. Therefore, isolated WT and $Mrp14^{-/-}$ bone marrow neutrophils were

introduced into E-selectin, ICAM-1 and CXCL1 coated microflow chambers and changes in cell

shape were monitored over 10min (Fig. 3A). WT neutrophils displayed normal spreading

properties as depicted by the gradual increase in area and perimeter over time (Fig. 3B). In

line with these findings, circularity and solidity, parameters reflecting the polarization

capability of the cells and the amount of protrusions the cell developed, respectively,

decreased over time (Fig. 3C). In contrast, increment of area and perimeter was significantly

less pronounced in $Mrp14^{-/-}$ cells (Fig. 3B). Circularity and solidity did only marginally decrease

over time in $Mrp14^{-/-}$ cells, suggesting that neutrophils are unable to polarize properly and to

extend protrusions (Fig. 3C). These results imply a substantial role of cytosolic S100A8/A9 in

β_2 integrin outside-in signaling.

Next, we wanted to examine consequences of impaired neutrophil spreading in absence of

S100A8/A9 by analyzing neutrophil crawling under flow. Therefore, we introduced isolated

neutrophils into E-selectin, ICAM-1 and CXCL1 coated microflow chambers and allowed them

to adhere for 3min to the substrates. Thereafter, we applied physiological shear stress (2dyne

cm^{-2}) and analyzed crawling behavior. WT neutrophils resisted shear forces and slowly crawled

in the direction of the flow, whereas $Mrp14^{-/-}$ neutrophils crawled in an intermittent and jerky

manner (Fig. 3D and Movie S1). In line, $Mrp14^{-/-}$ neutrophils covered significantly longer

distances (Fig. 3E), with an increased directionality toward flow direction (Fig. 3F) and

displayed an increased crawling velocity compared to WT cells (Fig. 3G).

To confirm impaired crawling and defective outside-in signaling dependent adhesion

strengthening in neutrophils lacking cytosolic S100A8/A9, we conducted a neutrophil

detachment assay using E-selectin, ICAM-1 and CXCL1 coated microflow chambers and

applied increasing shear stress. We found lower numbers of adherent $Mrp14^{-/-}$ neutrophils

compared to WT neutrophils with increasing shear stress (Fig. S2A). This is in line with our in

vivo findings where we detected a negative correlation between the number of adherent cells

and increasing shear stress in *Mrp14$^{-/-}$* animals (Fig. 1E). Together, these findings suggest a

critical role of intracellular S100A8/A9 in adherent neutrophils to resist high shear stress

conditions.

Following engagement of the ligand ICAM-1 to activated β$_2$ integrins in neutrophils, the

proline-rich tyrosine kinase Pyk2 and the focal adhesion adaptor protein paxillin are, among

other proteins, rapidly tyrosine phosphorylated thereby being critical events for cell adhesion,

migration and podosome formation [27, 30, 31]. To test the role of cytosolic S100A8/A9 in

mediating outside-in signaling events on the mechanistic level, we seeded WT and *Mrp14$^{-/-}$*

neutrophils on ICAM-1 coated plates, stimulated the cells with CXCL1 and determined Pyk2

and paxillin phosphorylation by western blot analysis. We found increased abundance of Pyk2

and paxillin phosphorylation in CXCL1 stimulated WT cells, while no increase was detectable

in *Mrp14$^{-/-}$* neutrophils (Fig. 3H and 3I). Taken together, these data indicate that cytosolic

S100A8/A9 is essential during ICAM-1 induced integrin outside-in signaling events and

therefore indispensable for post arrest modifications including cell polarization and the

formation of cell protrusions.

Cytosolic S100A8/A9 drives neutrophil cytoskeletal rearrangement by regulating LFA-1

nanocluster formation and Ca^{2+} availability within the clusters

Integrin outside-in signaling strongly depends on focal cluster formation of high-affinity LFA-1

and high Ca^{2+} concentrations within these clusters [10, 12-14]. Since S100A8/A9 is a Ca^{2+}

binding protein, we studied LFA-1 clustering and Ca^{2+} signatures during neutrophil adhesion

under flow conditions. For this approach, we isolated neutrophils from Ca^{2+} reporter mice (WT

Lyz2xGCaMP5) and S100A8/A9 deficient Ca^{2+} reporter mice (*Mrp14$^{-/-}$ ^{Lyz2xGCaMP5}*) and fluorescently

labelled the cells with an LFA-1 antibody (Fig.4a). Neutrophils were then introduced into E-

selectin, ICAM-1 and CXCL1 coated flow chambers, allowed to settle for 3min before shear

was applied (2dyne cm^{-2}). Time-lapse movies of fluorescence LFA-1 and Ca^{2+} signals were

recorded for 10min by confocal microscopy. First, LFA-1 signals from single cell analysis (Fig.

4A) were segmented through automatic thresholding in order to generate a binary image of

the LFA-1 signals (LFA-1 mask) (Fig. 4B). Then, LFA-1 nanoclusters were considered as such if

they spanned a minimum area of 0.15µm^2 (Fig. 4C), according to literature [32]. We found that

Mrp14$^{-/-}$ ^{Lyz2xGCaMP5} neutrophils formed significantly less LFA-1 nanoclusters compared to WT

^{Lyz2xGCaMP5} neutrophils suggesting an involvement of cytosolic S100A8/A9 in LFA-1 nanocluster

formation (Fig. 4D and Movie S2). Next, we investigated Ca^{2+} intensities within LFA-1

nanoclusters (Fig. 4E) to determine Ca^{2+} levels at the LFA-1 focal adhesion spots (Fig. 4F). We

found a significant reduction of Ca^{2+} levels in LFA-1 nanocluster areas of *Mrp14$^{-/-}$ ^{Lyz2xGCaMP5}*

neutrophils compared to WT ^{Lyz2xGCaMP5} neutrophils (Fig. 4G and Movie S3) suggesting an

impaired availability of free intracellular Ca^{2+} at LFA-1 nanocluster sites in absence of cytosolic

S100A8/A9. Strikingly, Ca^{2+} levels in the cytoplasm (outside of LFA-1 nanoclusters, Fig. 4H and

4I) did not differ between WT ^{Lyz2xGCaMP5} and *Mrp14$^{-/-}$ ^{Lyz2xGCaMP5}* neutrophils (Fig. 4J), suggesting

that cytosolic S100A8/A9 plays an important role especially in supplying Ca^{2+} at LFA-1 adhesion

spots. To investigate localization of S100A8/A9 during neutrophil post-arrest modification, we

isolated neutrophils from WT mice and labeled them with the cell tracker green CMFDA and

an LFA-1 antibody. The cells were introduced into flow chambers coated with E-selectin,

ICAM-1, and CXCL1, allowed to settle for 3min, and then subjected to continuous shear stress

(2dyne cm^{-2}) for 10min. After fixation and permeabilization, the cells were stained for

intracellular S100A9. LFA-1 nanoclusters were identified, and S100A9 intensity in these

clusters was compared to that in cytoplasmic areas outside the nanoclusters. We observed

higher S100A9 intensity at LFA-1 nanoclusters compared to the rest of the cytoplasm (non

LFA-1 nanoclusters) in stimulated WT neutrophils (**Fig. 4K and 4L**), indicating that S100A8/A9

localizes at LFA-1 nanocluster sites, supplying them with Ca^{2+} locally.

In line, overall Ca^{2+} levels under basal conditions (poly-L-lysine coating, static conditions) were

similar between WT ^{Lyz2xGCaMP5} and $Mrp14^{-/- \ Lyz2xGCaMP5}$ neutrophils (Fig. S3A). Calmodulin levels

did not differ between WT ^{Lyz2xGCaMP5} and $Mrp14^{-/- \ Lyz2xGCaMP5}$ cells as analyzed by western blot

(Fig. S3B).

LFA-1 is known to be rapidly recycled and to spatially redistribute to form a ring like structure

that co-clusters with endothelial ICAM-1 during neutrophil migration [33]. To study spatial

distribution of LFA-1 nanoclusters (Fig. 4M), we used Ripley's K function in WT ^{Lyz2xGCaMP5} and

$Mrp14^{-/- \ Lyz2xGCaMP5}$ neutrophils (Fig. S3C). Ripley's K is a spatial statistic that compares a given

point distribution with a random distribution [34]. WT ^{Lyz2xGCaMP5} neutrophils showed

significantly more aggregated LFA-1 nanoclusters within $10\mu m^2$ area, suitable for LFA-1

enriched pseudopods, compared to $Mrp14^{-/- \ Lyz2xGCaMP5}$ neutrophils (Fig. 4N), independent

from the total LFA-1 nanocluster number. These results show that in the absence of cytosolic

S100A8/A9, LFA-1 nanoclusters are more randomly distributed compared to control and

indicate that subcellular redistribution of LFA-1 during migration requires cytosolic

S100A8/A9.

Recent work has shown that Ca^{2+} signaling promotes F-actin polymerization at the uropod of

polarized neutrophils [13]. Actin waves in turn are known to be important for membrane

protrusion formation, neutrophil polarization and firm arrest [35]. Therefore, we examined F-

actin dynamics in the presence or absence of cytosolic S100A8/A9. For this, we used the same

experimental setting as for the LFA-1 cluster analysis but this time we fluorescently labelled

WT ^{Lyz2xGCaMP5} and $Mrp14^{-/- \ Lyz2xGCaMP5}$ neutrophils for F-actin. We generated a mask using the

myeloid cell marker $Lyz2$ (Fig. 4O) and applied the mask to the F-actin channel (Fig. 4P). In line

with our previous results on reduced Ca^{2+} levels within LFA-1 adhesion clusters in the absence

of S100A8/A9, we found a strongly reduced F-actin signal in $Mrp14^{-/- \ Lyz2xGCaMP5}$ neutrophils

compared to WT ^{Lyz2xGCaMP5} neutrophils (Fig. 4Q and Movie S4). Total actin levels as determined

by western blot analysis did not differ between WT ^{Lyz2xGCaMP5} and $Mrp14^{-/- \ Lyz2xGCaMP5}$

neutrophils (Fig. S3D).

Finally, we analyzed the frequency of Ca^{2+} flickers in WT ^{Lyz2xGCaMP5} and $Mrp14^{-/- \ Lyz2xGCaMP5}$

neutrophils induced by E-selectin, ICAM-1 and CXCL1 stimulation using high throughput

computational analysis. We found an increased number of Ca^{2+} flickers/min in the absence of

S100A8/A9 (Fig. S3E and S3F), going along with a shorter duration of the Ca^{2+} event compared

to control cells (Fig. S3G and S3H). This finding suggests that cytosolic S100A8/A9 is not only

important for local Ca^{2+} supply at focal LFA-1 nanocluster sites, but in addition "stabilizes" Ca^{2+}

signaling, preventing fast and uncontrolled Ca^{2+} flickering.

Taken together, these data show that cytosolic S100A8/A9 is indispensable for LFA-1

nanocluster formation and actin-dependent cytoskeletal rearrangements by providing and/or

promoting Ca^{2+} supply at the LFA-1 nanocluster sites.

Cytosolic S100A8/A9 is dispensable for chemokine induced ER store Ca^{2+} release and for the

initial phase of SOCE.

Our data suggest that intracellular S100A8/A9 is a fundamental regulator of cytosolic Ca^{2+}

availability within neutrophils during the recruitment process thereby affecting subcellular

LFA-1 and actin dynamics and distribution. Finally, we wanted to study any potential impact

of cytosolic S100A8/A9 on Ca^{2+} store release and on Store-Operated Ca^{2+} Entry (SOCE) during

neutrophil activation by investigating GPCR induced Ca^{2+} signaling using flow cytometry. First,

we investigated Ca^{2+} release from the ER and therefore performed the experiments in absence

of extracellular Ca^{2+}. We could not detect any differences in CXCL1 induced ER store Ca^{2+}

release between WT and $Mrp14^{-/-}$ cells, indicating that GPCR induced downstream signaling

leading to ER store depletion is not affected by the absence of cytosolic S100A8/A9 (Fig. 5A

and 5B). In addition, overall basal Ca^{2+} levels (prior to chemokine stimulation) were similar

between WT and $Mrp14^{-/-}$ neutrophils (Fig. 5C).

Next, we wanted to investigate whether the absence of cytosolic S100A8/A9 might modify

chemokine induced SOCE. Therefore, we stimulated isolated WT and $Mrp14^{-/-}$ neutrophils

with CXCL1 in the presence of extracellular Ca^{2+} (Fig. 5D). Again, basal Ca^{2+} levels were not

different between WT and $Mrp14^{-/-}$ cells (Fig. 5E). Also Ca^{2+} release-activated channels (CRAC)

functionality was intact as shown by an identical increase in cytosolic Ca^{2+} amount upon CXCL1

stimulation in WT and $Mrp14^{-/-}$ neutrophils (Fig. 5F). However, we detected different decay

kinetics between WT and $Mrp14^{-/-}$ neutrophils as $Mrp14^{-/-}$ neutrophils displayed a steeper

decay (Fig. 5G). Taken together, these data suggest that the presence of cytosolic S100A8/A9

is not a prerequisite for chemokine/GPCR induced Ca^{2+} release from ER stores and for the

initialization of SOCE via CRAC channels. However, absence of cytosolic S100A8/A9 might

disturb Ca^{2+} signaling in a temporal manner.

DISCUSSION

S100A8/A9 is a Ca^{2+} binding protein, mainly located within the cytosolic compartment of

myeloid cells [6, 16]. Once secreted, S100A8/A9 heterodimers exhibit pro-inflammatory

effects by engagement with its respective receptors including TLR4 and RAGE on a broad

spectrum of effector cells, among them phagocytes, lymphocytes and endothelial cells [16,

20]. In addition, extracellular S100A8/A9 is a well-established biomarker for many acute and

chronic inflammatory disorders, including cardiovascular diseases, autoimmune diseases and

infections [16, 20, 36]. The tetrameric form of S100A8/A9 was recently shown to have an anti-

inflammatory effects during an inflammatory process potentially protecting the organism

from overwhelming immune responses [23]. Despite increasing evidence of the pro- and anti-

inflammatory effects of secreted S100A8/A9, little is known about its intracellular role in

myeloid cells. Here, we show that S100A8/A9 is still abundantly present in the cytosolic

compartment of neutrophils even after its active release during inflammation. In addition, we

demonstrate that cytosolic S100A8/A9 has a functional impact on neutrophil recruitment

during β_2 integrin outside-in signaling events by ensuring high Ca^{2+} levels at LFA-1 cluster sites

independent of its extracellular functions. Neutrophil β_2 integrin outside-in signaling is known

to mediate post-arrest modifications including cytoskeletal rearrangements [22, 37]. *Mrp14*$^{-/-}$

cells, which also lack MRP8 in mature cells of the myeloid lineage (functional S100A8/A9

deficient cells) [21, 38], were unable to properly spread, polarize and crawl. This resulted in a

marked impairment of adherent neutrophils to withstand physiological shear forces exerted

by the circulating blood. Defective outside-in signaling in absence of S100A8/A9 was

accompanied by reduced phosphorylation of paxillin and Pyk2, two critical factors involved in

the regulation of β_2 integrin mediated cytoskeletal rearrangements [27]. Of note, *Mrp14*$^{-/-}$

myeloid cells have been shown to comprise alterations of cytoskeletal function before [21, 39-

43]. In the original publication describing the phenotype of *Mrp14*$^{-/-}$ mice, an abnormally

polarized cell shape of MRP14 deficient cells was described and therefore a potential role of

S100A8/A9 in cytoskeletal reorganization was already hypothesized [21]. In 2014, Vogl et al.

demonstrated that cytosolic S100A8/A9 had an impact on the stabilization of microtubules

(MTs) via direct interaction of S100A8/A9 with tubulin in resting phagocytes. Upon p38 MAPK

and concomitant Ca^{2+} signaling, S100A8/A9 was shown to dissociate from MTs, leading to de-

polymerization of MTs thereby allowing neutrophils to transmigrate into inflamed tissue. This

might also explain decreased migration of *Mrp14$^{-/-}$* granulocytes in a mouse wound healing

model [39]. Additional studies described cytosolic S100A8/A9 to translocate to the membrane

and colocalize with vimentin in monocytes upon activation [40], to interact with keratin in

epithelial cells [41] and to associate with F-actin localized to lamellipodia in fMLF stimulated

neutrophils [42]. Those findings led us to investigate a potential role of cytosolic S100A8/A9

in the Ca^{2+} dependent interplay of plasma membrane located adhesion sites and the

cytoskeleton during neutrophil recruitment.

Neutrophil activation during leukocyte recruitment goes along with Ca^{2+} flux initiated e.g. by

the engagement of chemokines with G-protein coupled receptors (GPCRs). Subsequently,

phospholipase C beta (PLC-β) is activated and leads to the production of Inositol-1,4,5-

triphosphate (IP3), which in turn elicits the IP3-receptor in the endoplasmic reticulum (ER),

resulting in a rapid Ca^{2+} release from ER stores into the cytoplasm. The decrease in Ca^{2+}

concentration in the ER in turn activates the Ca^{2+} sensor stromal interaction molecules (STIM1

and STIM2) triggering the entry of extracellular Ca^{2+} through SOCE mainly via the Ca^{2+}-release

activated (CRAC) channel ORAI-1 and transient receptor potential (TRP) channels [8, 10, 44].

ORAI-1 is recruited to adhesion cluster sites ensuring high Ca^{2+} levels at the "inflammatory

synapse" and rapid rise in intracellular Ca^{2+} concentration, which mediates the assembly of

cytoskeletal adaptor proteins to integrin tails and allows the onset of pseudopod formation

[14, 45]. The importance of localized Ca^{2+} availability in subcellular domains has also been

shown in T-cells during the engagement with antigen presenting cells within the

immunological synapse. In T-cells, mitochondria play a central role as Ca^{2+} buffers and as Ca^{2+}

conductors that collect cytosolic Ca^{2+} at the entry site, (i.e. through open CRAC channels

located at the immunological synapse) and distribute it throughout the cytosol. [46]. Here we

show that in neutrophils cytosolic S100A8/A9 colocalizes with LFA-1 during intravascular

adhesion and increases and stabilizes Ca^{2+} availability at the LFA-1 nanocluster sites, mediating

spatial clustering of LFA-1 and sustained polymerization of F-actin, both essential steps for

efficient neutrophil adhesion strengthening. In addition, presence of cytosolic S100A8/A9

stabilizes duration of Ca^{2+} signals within the cells, as WT cells displayed longer frequencies of

Ca^{2+} events with less flickers min^{-1}, which might in addition be important for the stability of

the inflammatory synapse.

As reported earlier, chemokine induced Ca^{2+} influx through SOCE at the plasma membrane is

indispensable for the activation of high affinity β_2 integrins [11]. This early step during

leukocyte recruitment (inside-out signaling) was not affected in absence of cytosolic

S100A8/A9. In line, $Mrp14^{-/-}$ cells displayed similar CXCL1 induced Ca^{2+} fluxes compared to WT

cells. Ca^{2+} release from intracellular stores and initial phases of SOCE were fully functional, as

shown by flow cytometry of Indo-1 dye loaded neutrophils. These findings are in accordance

with a study by Hobbs et al., which also described normal Ca^{2+} influx in S100A8/A9 deficient

neutrophils induced by the chemokine MIP-2 [47]. However, we found an impact of cytosolic

S100A8/A9 in sustaining high Ca^{2+} concentrations, as Ca^{2+} fluxes decreased faster in the

absence of S100A8/A9. Whether this faster decrease is mediated through a direct effect of

cytosolic S100A8/A9 on SOCE or through a potential buffer capacity of cytosolic S100A8/A9

needs to be further investigated. Hobbs et al. proposed no impact of S100A9 deletion in the

recruitment of neutrophils by using a thioglycolate induced peritonitis model. However,

peritoneal neutrophil emigration was shown to be rather independent of LFA-1 [48, 49],

whereas extravasation into cremaster muscle tissue strongly relies on the β_2 integrin LFA-1

and integrin clustering [50].

Taken together, we identified a critical role of cytosolic S100A8/A9 in neutrophil recruitment.

We show that its absence leads to reduced Ca^{2+} signaling and impaired sustained Ca^{2+} supply

at LFA-1 nanocluster sites. Attenuated Ca^{2+} signatures in turn affect β_2 integrin dependent

cytoskeletal rearrangements and substantially compromises neutrophil recruitment during

the inflammatory response. These findings uncover cytosolic S100A8/A9 as a potentially

interesting therapeutic target to reduce neutrophil recruitment during inflammatory

disorders with unwanted overwhelming neutrophil influx.

MATERIALS AND METHODS

Mice

C57BL/6 wildtype (WT) mice were purchased from Charles Rivers Laboratories (Sulzfeld, Germany). *Mrp14*[-/-] (functional double S100A8 and S100A9 ko animals) mice were kindly provided by Johannes Roth (Institute for Immunology, Muenster, Germany). *B6;129S6-Polr2atm1(CAG-GCaMP5g-tdTomato)* crossbred with Lyz2[Cre] (*GCaMP5xWT*) were kindly provided by Konstantin Stark (LMU, Munich, Germany) and crossbred with *Mrp14*[-/-] mice (GCaMP5x*Mrp14*[-/-]). All mice were housed at the Biomedical Center, LMU Munich, Planegg-Martinsried, Germany. Male and/or female mice (8–25 weeks old) were used for all experiments. Animal experiments were approved by the Regierung von Oberbayern (AZ.: ROB-55.2-2532.Vet_02-17-102 and ROB-55.2-2532.Vet_02-18-22) and carried out in accordance with the guidelines from Directive 2010/63/EU. For in vivo experiments, mice were anaesthetized via i.p. injection using a combination of ketamine/xylazine (125mg kg^{-1} and 12.5mg kg^{-1} body weight, respectively in a volume of 0.1mL NaCl per 8g body weight). All mice were sacrificed at the end of the experiment by cervical dislocation.

Neutrophil isolation

Bone marrow neutrophils were isolated using the EasySep Mouse Neutrophil Enrichment Kit according to the manufacturer's instructions (STEMCELL Technologies). Isolated neutrophils were then resuspended in HBSS buffer [containing 0.1% of glucose, 1mM $CaCl_2$, 1mM $MgCl_2$, 0.25% BSA, and 10mM HEPES (Sigma-Aldrich), pH7.4, complete HBSS].

S100A8/A9 ELISA

In vitro release of S100A8/A9 was performed as described before [51]. Briefly, bone marrow neutrophils were isolated from WT mice. 24 well-plates were coated with recombinant murine

(rm) E-selectin (rmCD62E-Fc chimera, 10μg mL^{-1}, R&D Systems) or PBS/0.1% BSA at 4°C

overnight, blocked with PBS/5% casein (Sigma-Aldrich) and washed twice with PBS. 5x10^5

neutrophils were reconstituted in complete HBSS buffer and incubated under shaking

conditions on the coated slides for 10min at 37°C and 5% CO2. To assess the total intracellular

S100A8/A9 levels, cells were lysed in 2% Triton X-100 (Applichem). Finally, cellular

supernatants were analyzed by Enzyme-Linked Immunosorbent Assay (ELISA) to determine

the concentrations of S100A8/A9.

Murine cremaster muscle models

Leukocyte recruitment was investigated by intravital microscopy in inflamed cremaster

muscle venules as reported previously [52]. Shortly, intrascrotal (i.s.) injection of rmTNF-α

(500ng, R&D Systems) was applied to WT and *Mrp14*$^{-/-}$ mice in order to induce an acute

inflammation in the cremaster muscle. Two hours after injection, the carotid artery of

anaesthetized mice was catheterized for later blood sampling (ProCyte Dx; IDEXX

Laboratories) or intra-arterial (i.a.) injection. Thereafter, the cremaster muscle was

exteriorized and intravital microscopy was conducted on an OlympusBX51 WI microscope,

equipped with a 40x objective (Olympus, 0.8NA, water immersion objective) and a CCD

camera (KAPPA CF 8 HS). Post-capillary venules were recorded using VirtualDub software for

later analysis. Rolling flux fraction, number of adherent cells mm^{-2}, vessel diameter and vessel

length were analyzed using FIJI software [53]. During the entire experiment, the cremaster

muscle was superfused with thermo-controlled bicarbonate buffer as described earlier [54].

Centerline blood flow velocity in each venule was measured with a dual photodiode (Circusoft

Instrumentation). Subsequently, cremaster muscles were removed, fixed in 4% PFA solution

O.N. at 4°C and the next day stained with Giemsa (Merck) to assess the number of perivascular

neutrophils. The tissues were mounted in Eukytt mounting medium and covered with a 170μm

coverslip. Neutrophils were discriminated from other leukocyte subpopulations based on

nuclear shape and granularity of the cytosol. The analysis of transmigrated leukocytes was

carried out at the Core Facility Bioimaging of the Biomedical Center with a Leica DM2500

transmission bright field microscope, equipped with a 100x, 1.4 NA, oil immersion objective

and a Leica DMC2900 CMOS camera. Resulting images had 2048x1536 pixels and a pixel size

of 58nm.

For rescue experiments, we adopted either the TNF-α-induced inflammation model as

described above or the trauma-induced inflammation model of the mouse cremaster muscle.

In the trauma model, sterile inflammation was induced by opening and exteriorizing the

cremaster muscle without application of any stimulus. Intravital microscopy was conducted

as described above. After finding an appropriate spot, the same vessel was recorded before

and after injection of mutant murine S100A8/S100A9N70AE79A (S100A8/A9^{mut}, aa exchange

N70A and E79A, 50µg mouse^{-1} in 100µL, provided by Thomas Vogl, University of Muenster,

Germany) and the number of adherent cells mm^{-2} were counted pre and post injection in WT

and $Mrp14^{-/-}$ mice.

S100A8/A9 intracellular staining

For the analysis of cytosolic S100A8/A9 levels, TNF-α stimulation of the mouse cremaster

muscle was carried out as described above. Subsequently, cremaster muscles were removed,

fixed in 4% PFA solution, and immunofluorescence staining for PECAM-1 (AlexaFluor488

labelled primary monoclonal rat antibody, 5µg mL^{-1}, MEC13.3, BioLegend) and S100A9 (Cy5.5

directly labelled, 5µg mL^{-1}, clone 322, provided by Thomas Vogl) was conducted. Stained

samples were mounted in Vectashield mounting medium, covered with a 0.17µm coverslip

and imaged by confocal microscopy at the Core Facility Bioimaging of the Biomedical Center,

LMU Munich, with an upright Leica SP8X WLL microscope, equipped with a HC PL APO

40x/1.30 NA oil immersion objective. AF488 was excited with 488nm, Cy5.5 with 543nm.

Detection windows were 500 – 568 and 550 – 640 nm, respectively. Both channels were

recorded sequentially. Hybrid photodetectors were used to record images with 512x512 pixels

with a pixel size of 0.427μm. Single cell analysis was carried out by FIJI software using macros

as follows: MAX projection of Z-stacks were created and neutrophils were segmented by

thresholding using S100A8/A9 signal. Then, cell masks were applied back to the original

images and S100A8/A9 mean fluorescence intensity (MFI) averaged on stack slices. Finally,

S100A8/A9 MFIs were analyzed from intravascular and extravasated neutrophils.

Neutrophil surface marker staining

Peripheral blood from WT and *Mrp14$^{-/-}$* mice was harvested and erythrocytes were lysed with

lysing solution (BD FACS™). Samples were stained for CD18-FITC (5μg mL^{-1}; C71/16;

Pharmigen), CD11a-APC (2μg mL^{-1}; M17/4; eBioscience), CD11b-BV510 (0.3μg mL^{-1}; M1/70;

BioLegend), CD62L-FITC (5μg mL^{-1}; MEL-14; BioLegend), PSGL1-PE (2μg mL^{-1}; 2PH1;

Pharmigen), CXCR2-APC (5μg mL^{-1}; 242216; R&D Systems), CD44-BV570 (0.3μg mL^{-1}; IM7;

BioLegend). Respective isotype controls were used: IgG2a-FITC (5μg mL^{-1}; RTK2759;

BioLegend), IgG2a-APC (2μg mL^{-1}; RTK2758; BioLegend), IgG2b-BV510 (0.3μg mL^{-1}; RTK4530;

BioLegend), IgG1-PE (2μg mL^{-1}; eBRG1; eBioscience), IgG2b-BV570 (0.3μg mL^{-1}; RTK4530;

BioLegend). Neutrophils were defined as Ly6G$^+$ cells (0.8μg mL^{-1}; 1A8; BioLegend).

Neutrophil adhesion ex vivo

Flow chamber assays were carried out as previously described [22]. Briefly, rectangular

borosilicate glass capillaries (0.04x0.4mm; VitroCom) were coated with a combination of rmE-

selectin (CD62E Fc chimera; 20μg mL^{-1}; R&D Systems), rmICAM-1 (ICAM-1 Fc chimera; 15μg

mL^{-1}; R&D Systems), and rmCXCL1 (15μg mL^{-1}; Peprotech) for 3h at RT and blocked with

PBS/5% casein (Sigma-Aldrich) over night at 4°C. WT and *Mrp14$^{-/-}$* whole blood was perfused

through the microflow chamber via a carotid artery catheter of anesthetized mice at varying

shear stress levels. Movies were recorded on an OlympusBX51 WI microscope with a 20x,

0.95NA, water immersion objective and a CCD camera (KAPPA CF 8 HS) with VirtualDub

software [55]. Resulting images had 768x576 pixels and a pixel size of 0.33μm .Number of

rolling and adherent leukocytes/field of view (FOV) were counted using Fiji software, over

one-minute time window after 6min of blood infusion.

β$_2$ integrin activation assay

β$_2$ integrin activation was determined through a modified soluble ICAM-1 binding assay [6].

Bone marrow murine neutrophils were isolated as described above. Enriched neutrophils

(1.5×10^6) were incubated and stained with rmICAM-1 Fc chimera (40μg mL^{-1}, R&D Systems),

IgG-Fc-biotin (12.5μg mL^{-1}; eBioscience), and streptavidin-PerCP-Cy5.5 (2μg mL^{-1}; BioLegend).

Then, cells were stimulated with rmCXCL1 (10nM) or PBS (control) in complete HBSS buffer

for 5min at 37°C. The amount of bound rmICAM-1 to the β$_2$ integrin was assessed by flow

cytometry (CytoFlex S, Beckmann Coulter) and the median shift relative to the control was

analyzed by FlowJo software.

Static adhesion assay

Neutrophil static adhesion assay was performed as previously described [56]. Shortly, 96-well

plates were coated with rmICAM-1 (3μg mL^{-1}) over night at 4°C and washed with PBS.

Neutrophils were resuspended in complete HBSS and seeded at 1x10^5 cells per well. Cells were

allowed to settle for 5min at 37°C and stimulated with 10nM rmCXCL1 or PBS (control) for

10min at 37°C. Using a standard curve, adherent neutrophils were calculated as percentage

of total cells added. Standard curve preparation was done by adding 100%, 80%, 60%, 40%,

20%, and 10% of the cell suspension on poly-L-lysine coated wells (100µg mL^{-1}) in triplicates.

Non adherent cells were washed away while adherent cells were fixed with 1% glutaraldehyde

and stained with 0.1% crystal violet solution (Sigma-Aldrich). Absorption at 590nm was

measured with a microplate reader (PowerWave HT, Biotek, USA) after lysis of cells with 10%

acetic acid solution, as previously described [57].

Spreading assay

To study neutrophil spreading, rectangular borosilicate glass capillaries (0.04x0.40mm;

VitroCom) were coated with rmE-selectin (CD62E Fc chimera; 20µg mL^{-1}), rmICAM-1 (15µg

mL^{-1}), and rmCXCL1 (15µg mL^{-1}) for 3 hours at RT and blocked with PBS/5% casein over night

at 4°C. Bone marrow neutrophils were matured in RPMI 1640 (Sigma-Aldrich) containing FCS

(10%, Sigma-Aldrich), GlutaMAX (1%, ThermoFisher), Penicillin-Streptomycin solution (1%,

Corning®) and supplemented with 20% WEHI-3B-conditioned medium over night at 37°C and

applied into the flow chamber at a shear stress level of 1dyne cm^{-2} using a high-precision

syringe pump (Harvard Apparatus, Holliston, Massachusetts, USA). Cells were incubated with

Fc-block (murine TruStain FcX; BioLegend) for 5min at RT before being introduced into the

chambers. Spreading behavior of the cells was observed and recorded on a Zeiss Axioskop2

with a 20x, 0.5NA water immersion objectibve and a Hitachi KP-M1AP camera with VirtualDub.

Resulting images had 1360x1024 pixels and a pixel size of 600nm. Cell shape changes were

quantified using FIJI software, analyzing cell area, perimeter, circularity ($4\pi \frac{[area]}{[perimeter]^2}$) and

solidity ($\frac{[area]}{[convex\ area]}$).

Crawling assay

15μ-Slides VI$^{0.1}$ (Ibidi) were coated with a combination of rmE-selectin (20μg mL^{-1}), rmICAM-

1 (15μg mL^{-1}), and rmCXCL1 (15μg mL^{-1}) for 3h at RT and blocked with PBS/5% casein over

night at 4°C. Overnight matured bone marrow neutrophils from WT and *Mrp14$^{-/-}$* mice were

resuspended in complete HBSS at 1x10^6 mL^{-1}, introduced into the chambers and allowed to

settle and adhere for 3min until flow was applied (2dyne cm^{-2}) using a high-precision perfusion

pump. Experiments were conducted on a ZEISS, AXIOVERT 200 microscope, provided with a

ZEISS 20x objective [0.25NA], and a SPOT RT ST Camera. MetaMorph software was used to

generate time-lapse movies for later analysis. 20min of neutrophil crawling under flow were

analyzed using FIJI software [53] and chemotaxis tool plugin (Ibidi).

Paxillin and Pyk2 phosphorylation

Paxillin and Pyk2 phosphorylation was investigated as previously described [22]. Briefly, 2×10^6

WT or *Mrp14$^{-/-}$* bone marrow murine neutrophils were seeded on rmICAM-1 coated wells

(15μg mL^{-1}) for 5min and stimulated with rmCXCL1 (10nM) for 5min at 37°C. Cells were then

lysed with lysis buffer [containing 150mM NaCl, 1% Triton X-100, 0.5% Sodium deoxycholate

(Sigma-Aldrich), 50mM Tris–HCl pH7.3 (Merck), 2mM EDTA (Merck) supplemented with

protease (Roche), phosphatase inhibitors (Sigma-Aldrich) and 1xLaemmli sample buffer] and

boiled (95°C, 5min). Cell lysates were resolved by SDS–PAGE and electrophoretically

transferred onto PVDF membranes. After subsequent blocking (LI-COR blocking solution),

membranes were incubated with the following antibodies for later detection and analysis

using the Odyssey® CLx Imaging System and Image Studio software: rabbit α-mouse phospho-

Paxillin (Tyr118) or rabbit α-mouse Paxillin and rabbit α-mouse phospho-Pyk2 (Tyr402) or

rabbit α-mouse Pyk2 (all Cell Signaling). Goat-α-rabbit IRDye 800RD was used as secondary

antibody (Licor).

Detachment assays

To investigate shear resistance, rectangular borosilicate glass capillaries (0.04x0.40mm; VitroCom) were coated with rmE-selectin (CD62E Fc chimera; 20µg mL^{-1}), rmICAM-1 (15µg mL^{-1}), and rmCXCL1 (15µg mL^{-1}) for 3 hours at RT and blocked with 5% casein over night at 4°C. Whole blood from WT and *Mrp14$^{-/-}$* mice was perfused in the coated flow chambers via the cannulated carotid artery, where neutrophils were allowed to attach for 3min. Then, flow was applied through a high-precision perfusion pump and detachment assays performed over 10min with increasing shear stress (34 – 272 dyne cm^{-2}) every 30s. Experiments were recorded by time-lapse movies using the upright Zeiss Axioskop2 with the 20x, 0.5 NA water immersion objective as described above. Number of attached cells was counted at the end of each step.

LFA-1 clustering, S100A8/A9 distribution, Ca^{2+} localization and F-actin signature during neutrophil crawling under flow

15µ-Slides VI$^{0.1}$ (Ibidi) were used to study LFA-1 clustering, Ca^{2+} localization and F-actin signature during neutrophil crawling. Flow chambers were coated and blocked as described above. 2x10^6 isolated neutrophils from WT ^{Lyz2xGCaMP5} or *Mrp14$^{-/-Lyz2xGCaMP5}$* were stained with in-house AlexaFluor647 labelled (Antibody Labeling Kit, Invitrogen™) monoclonal anti LFA-1 rat antibody (5µg mL^{-1}, 2D7, BD Pharmingen) for 10min prior to the experiment or SiR-actin (200nM, Spirochrome™) O.N., respectively. Cells were seeded in the chambers and allowed to settle for 2min before flow was applied (2dyne cm^{-2}) using a high-precision perfusion pump. Samples were imaged by confocal microscopy at the core facility Bioimaging of the Biomedical Center with an inverted Leica SP8X WLL microscope, equipped with a HC PL APO 40x/1.30 NA oil immersion objective. Observation was at 37°C. Hybrid photodetectors were used to record images with 512x512 pixels and a pixel size of 0.284µm. GCaMP5-GFP was excited with

488nm, AF647 or SiR-Actin with 633nm. Detection windows were 498 – 540 and 649 – 710nm,

respectively. For movies, one image was recorded every 0.44 seconds or every 2 seconds, over

10min. Automated single cell analysis was performed using macros with Fiji software, for

minute 0-1, minute 5-6 and minute 9-10 of each recording. For the LFA-1 nanocluster analysis,

the LFA-1 channel was automatically segmented and ROIs of a minimum size of $0.15\mu m^2$ were

considered as LFA-1 nanosclusters, as reported earlier [32]. This represented a minimum size

of 2 pixels in our analysis. The number of clusters was averaged for each analyzed time point

(min 0-1, min 5-6 and min 9-10). For the subcellular Ca^{2+} analysis at the LFA-1 cluster sites, the

LFA-1 segmented channel was applied to the Ca^{2+} channel and Ca^{2+} events in the selected ROIs

were determined, normalized to the LFA-1 areas, and averaged over each minute of analysis.

For the Ca^{2+} analysis in the negative LFA-1 area, we again adopted semi-automated single cell

analysis and subtracted the LFA-1 mask from the *Lyz2* mask in order to obtain "LFA-1 cluster

negative masks". Later, the "LFA-1 cluster negative masks" were applied to the Ca^{2+} channel

and Ca^{2+} intensities were measured, normalized to the "LFA-1 cluster negative masks" and

averaged over each minute of analysis. For the analysis of S100A9 distribution at LFA-1

nanocluster areas, WT neutrophils were stained with CellTracker Green CMFDA (10μM,

Invitrogen™) for 45min and in-house AlexaFluor647-labeled monoclonal anti-LFA-1 rat

antibody (5μg/mL, 2D7, BD Pharmingen) for 10min prior to the experiment. The cells were

then seeded in chambers and allowed to settle for 3min, before applying continuous flow (2

dyne cm^{-2}) using a high-precision perfusion pump for 10min. After the flow, the cells were

fixed, permeabilized, and stained overnight at 4°C for intracellular S100A9, followed by

counterstaining with DAPI. A semi-automated single-cell analysis was performed to measure

S100A9 intensity in the LFA-1 nanocluster areas (obtained as described above) and in the

negative LFA-1 nanocluster areas (determined using the same procedure but with CellTracker

Green as the cell mask).

For the F-actin analysis, the *Lyz2* channel was automatically segmented to obtain a cell mask

and applied to the F-actin channel. F-actin intensities were measured and averaged over each

minute of analysis as described above.

Ca^{2+} store release and Ca^{2+} influx measurement – flow cytometry

Ca^{2+} store release and Ca^{2+} influx was analyzed by flow cytometry through an adapted protocol

[58]. WT and *Mrp14*$^{-/-}$ bone marrow neutrophils (2.5×10^6 mL^{-1}) were resuspended in PBS and

loaded with 3 µM Indo-1 AM (Invitrogen™) for 45min at 37°C. Cells were washed, resuspended

in complete HBSS buffer (2.5×10^6 mL^{-1}) and stained with an anti Ly6G-APC antibody (1µg mL^{-}

^{1}, 1A8, BioLegend) and with the Fixable Viability Dye eFluor™ 780 (1:1000; eBioscience™). Cells

($2x10^5$) were incubated for 2min at 37°C and 10nM CXCL1 was placed on the side of the FACS

tube in a 2µL droplet form. The cells were analyzed at the flow cytometry core facility of the

biomedical center with a BD LSRFortessa^{TM} flow cytometer. Samples were recorded for 45

seconds to establish a baseline. Afterwards, CXCL1 stimulation was initiated by tapping the

tube with subsequent fall of the drop into the cell suspension while continuously recording

Indo-1 AM signals from neutrophils over time. Data were analyzed using FlowJo software.

Calcium levels are expressed as relative ratios of fluorescence emission at 375nm/525nm

(calcium bound/calcium unbound) and Ca^{2+} signatures quantified as AUC of kinetic averages.

To measure Ca^{2+} store release only, Ca^{2+} free medium was used.

Spatial distribution analysis of LFA-1 nanoclusters

To evaluate the spatial distribution of LFA-1 nanoclusters in neutrophils, Ripley's K statistics

[34] was calculated for every time point in every experiment with radii between 0.5μm and

5.5μm. For every radius r, we calculated the K(r) value as follows:

$$K(r) = N^{-1} \sum_{i} \sum_{j \neq i} I\left(dist(i,j) \leq r\right)$$

Where *i* and *j* are two different LFA-1 nanocluster locations, *I* is the indicator function which

is 1 if the content within the parentheses is "True" and 0 if the content is "False", and N is a

normalization constant. For 'dist', the Euclidian distance was chosen and calculated via the

"pairwise_distances" from sklearn [59]. The sampling part of Ripley's K statistic was done by

drawing random locations as LFA-1 nanocluster events from the cell surface. To make Ripley's

K results comparable between different experiments, we normalized K(r) values such that the

random sampling upper bound, calculated for every experiment, was set to 1, and the random

sampling lower bound was set to -1. Thus, every normalized value between -1 and 1 is within

random borders, i.e. not distinguishable from a random spatial distribution. Values above 1

indicate aggregated LFA-1 nanoclusters and values below -1 indicate dispersed LFA-1

nanoclusters. At least 10 LFA-1 nanoclusters were considered for the spatial aggregation

analysis. The number of identified aggregated LFA-1 nanoclusters (values above 1) was

counted and averaged for every condition, resulting in an aggregation index.

Frequency and duration of Ca^{2+} oscillations

After the recording, Ca^{2+} mean intensities of the cells were calculated over time, counting each

cell as an individual ROI. Data was imported into a previously described custom analysis

pipeline for Ca^{2+} imaging data [60]. Briefly, the Ca^{2+} mean intensities were sequentially filtered

according to the standard values of the pipeline, considering only events with a z-score of at

least 3 (p<0.01). From those events, a graph was constructed to detect superimposed events.

Properties of the events, AUC or half-width, were used in the calculations afterwards. The

code of the analysis pipeline can be accessed in the corresponding repository at

https://github.com/szarma/Physio_Ca.

Calmodulin and β-Actin Western Blotting

WT and *Mrp14*$^{-/-}$ bone marrow murine neutrophils (1×10^6) were isolated as described above

and lysed with lysis buffer and boiled (95°C, 5min). Cell lysates were resolved by SDS–PAGE

and electrophoretically transferred onto PVDF membranes. After subsequent blocking (LI-COR

blocking solution), membranes were incubated with the following antibodies for later

detection and analysis using the Odyssey® CLx Imaging System and Image Studio software.

Rabbit α-mouse Calmodulin (5µg mL^{-1}, CellSignaling), rabbit α-mouse β-actin (1µg mL^{-1},

CellSignaling) and mouse α-mouse GAPDH (1µg mL^{-1}, Merck/Millipore), goat-α-mouse IRDye

680RD, and goat-α-rabbit IRDye800CW-coupled secondary antibodies (1µg mL^{-1}, LI-COR).

Statistics

Data are presented as mean +SEM, as cumulative distribution or representative images, as

depicted in the figure legends. Group sizes were selected based on experimental setup. Data

were analyzed and illustrated using GraphPad Prism 9 software. Statistical tests were

performed according to the number of groups being compared. For pairwise comparison of

experimental groups, a paired/unpaired Student's t-test and for more than two groups, a one-

way or two-way analysis of variance (ANOVA) with either Tukey's (one-way ANOVA) or Sidak's

(two-way ANOVA) *post-hoc* test with repeated measurements were performed, respectively.

p-values <0.05 were considered statistically significant and indicated as follows: ∗p<0.05;

∗∗p<0.01; ∗∗∗p<0.001.

Supplementary Materials

Supplementary Figures:

Fig. S1: Cytosolic S100A8/A9 is indispensable for neutrophil recruitment in vivo

Fig. S2: S100A8/A9 deficient cells are more susceptible to increasing shear stress compared

to WT cells

Fig. S3: S100A8/A9 deficient cells display higher frequencies but shorter duration of Ca^{2+}

waves compared to WT cells

Supplementary Tables:

Table S1: Microvascular parameters in vivo

Table S2: Microvascular parameters ex vivo

Supplementary Movies:

Movie S1: Neutrophil crawling depends on S100A8/A9

Movie S2: S100A8/A9 is essential for LFA-1 nanocluster formation and turnover

Movie S3: S100A8/A9 increases Ca^{2+} levels at the LFA-1 nanocluster sites

Movie S4: S100A8/A9 induces F-actin polymerization

References and notes

1.   Németh, T., M. Sperandio, and A. Mócsai, *Neutrophils as emerging therapeutic targets.* Nature Reviews Drug Discovery, 2020. **19**(4): p. 253-275.

2.   Ley, K., et al., *Neutrophils: New insights and open questions.* Science Immunology, 2018. **3**(30): p. eaat4579.

3.   Schmidt, S., M. Moser, and M. Sperandio, *The molecular basis of leukocyte recruitment and its deficiencies.* Mol Immunol, 2013. **55**(1): p. 49-58.

4.   Patcha, V., et al., *Differential inside-out activation of β2-integrins by leukotriene B4 and fMLP in human neutrophils.* Experimental Cell Research, 2004. **300**(2): p. 308-319.

5.   Chung, K.-J., et al., *A novel pathway of rapid TLR-triggered activation of integrin-dependent leukocyte adhesion that requires Rap1 GTPase.* Molecular Biology of the Cell, 2014. **25**(19): p. 2948-2955.

6.   Pruenster, M., et al., *Extracellular MRP8/14 is a regulator of β2 integrin-dependent neutrophil slow rolling and adhesion.* Nature Communications, 2015. **6**(1): p. 6915.

7.   Uhl, B., et al., *Aged neutrophils contribute to the first line of defense in the acute inflammatory response.* Blood, 2016. **128**(19): p. 2327-2337.

8.   Immler, R., S.I. Simon, and M. Sperandio, *Calcium signalling and related ion channels in neutrophil recruitment and function.* European Journal of Clinical Investigation, 2018. **48**(S2): p. e12964.

9.   Begandt, D., et al., *How neutrophils resist shear stress at blood vessel walls: molecular mechanisms, subcellular structures, and cell–cell interactions.* Journal of Leukocyte Biology, 2017. **102**(3): p. 699-709.

10. Dixit, N. and S.I. Simon, *Chemokines, selectins and intracellular calcium flux: temporal and spatial cues for leukocyte arrest.* Front Immunol, 2012. **3**: p. 188.

11. Schaff, U.Y., et al., *Calcium flux in neutrophils synchronizes beta2 integrin adhesive and signaling events that guide inflammatory recruitment.* Ann Biomed Eng, 2008. **36**(4): p. 632-46.

12. Dixit, N., et al., *Leukocyte function antigen-1, kindlin-3, and calcium flux orchestrate neutrophil recruitment during inflammation.* J Immunol, 2012. **189**(12): p. 5954-64.

13. Dixit, N., et al., *Migrational guidance of neutrophils is mechanotransduced via high-affinity LFA-1 and calcium flux.* J Immunol, 2011. **187**(1): p. 472-81.

14. Schaff, U.Y., et al., *Orai1 regulates intracellular calcium, arrest, and shape polarization during neutrophil recruitment in shear flow.* Blood, 2010. **115**(3): p. 657-666.

15. Edgeworth, J., et al., *Identification of p8,14 as a highly abundant heterodimeric calcium binding protein complex of myeloid cells.* J Biol Chem, 1991. **266**(12): p. 7706-13.

16. Pruenster, M., et al., *S100A8/A9: From basic science to clinical application.* Pharmacol Ther, 2016. **167**: p. 120-131.

17. Morikis, V.A., et al., *Selectin catch-bonds mechanotransduce integrin activation and neutrophil arrest on inflamed endothelium under shear flow.* Blood, 2017. **130**(19): p. 2101-2110.

18. Jorch, S.K., et al., *Complex regulation of alarmins S100A8/A9 and secretion via gasdermin D pores exacerbates autoinflammation in familial Mediterranean fever.* J Allergy Clin Immunol, 2023. **152**(1): p. 230-243.

19. Pruenster, M., et al., *E-selectin-mediated rapid NLRP3 inflammasome activation regulates S100A8/S100A9 release from neutrophils via transient gasdermin D pore formation.* Nature Immunology, 2023. **24**(12): p. 2021-2031.

20. Wang, S., et al., *S100A8/A9 in Inflammation.* Front Immunol, 2018. **9**: p. 1298.

21. Manitz, M.-P., et al., *Loss of S100A9 (MRP14) Results in Reduced Interleukin-8-Induced CD11b Surface Expression, a Polarized Microfilament System, and Diminished Responsiveness to Chemoattractants In Vitro.* Molecular and Cellular Biology, 2003. **23**(3): p. 1034-1043.

22. Immler, R., et al., *The voltage-gated potassium channel KV1.3 regulates neutrophil
recruitment during inflammation.* Cardiovascular Research, 2022. **118**(5): p. 1289-1302.
23. Russo, A., et al., *Alarming and Calming: Opposing Roles of S100A8/S100A9 Dimers and
Tetramers on Monocytes.* Advanced Science, 2022. **n/a**(n/a): p. 2201505.
24. Vogl, T., et al., *Autoinhibitory regulation of S100A8/S100A9 alarmin activity locally restricts
sterile inflammation.* The Journal of Clinical Investigation, 2018. **128**(5): p. 1852-1866.
25. Leukert, N., et al., *Calcium-dependent tetramer formation of S100A8 and S100A9 is essential
for biological activity.* J Mol Biol, 2006. **359**(4): p. 961-72.
26. Frommhold, D., et al., *Sialyltransferase ST3Gal-IV controls CXCR2-mediated firm leukocyte
arrest during inflammation.* J Exp Med, 2008. **205**(6): p. 1435-46.
27. Abram, C.L. and C.A. Lowell, *The ins and outs of leukocyte integrin signaling.* Annu Rev
Immunol, 2009. **27**: p. 339-62.
28. Legate, K.R., S.A. Wickström, and R. Fässler, *Genetic and cell biological analysis of integrin
outside-in signaling.* Genes & Development, 2009. **23**(4): p. 397-418.
29. Zarbock, A. and K. Ley, *Neutrophil Adhesion and Activation under Flow.* Microcirculation,
2009. **16**(1): p. 31-42.
30. Lu, F., et al., *Mechanism of integrin activation by talin and its cooperation with kindlin.*
Nature Communications, 2022. **13**(1): p. 2362.
31. Bouti, P., et al., *β2 Integrin Signaling Cascade in Neutrophils: More Than a Single Function.*
Front Immunol, 2020. **11**: p. 619925.
32. Fan, Z., et al., *Neutrophil recruitment limited by high-affinity bent β2 integrin binding ligand
in cis.* Nat Commun, 2016. **7**: p. 12658.
33. Shaw, S.K., et al., *Coordinated redistribution of leukocyte LFA-1 and endothelial cell ICAM-1
accompany neutrophil transmigration.* J Exp Med, 2004. **200**(12): p. 1571-80.
34. Dixon, P.M., *Ripley's K Function*, in *Encyclopedia of Environmetrics*. 2001.
35. Inagaki, N. and H. Katsuno, *Actin Waves: Origin of Cell Polarization and Migration?* Trends in
Cell Biology, 2017. **27**(7): p. 515-526.
36. Jukic, A., et al., *Calprotectin: from biomarker to biological function.* Gut, 2021. **70**(10): p.
1978-1988.
37. Rohwedder, I., et al., *Src family kinase-mediated vesicle trafficking is critical for neutrophil
basement membrane penetration.* Haematologica, 2020. **105**(7): p. 1845-1856.
38. Ehrchen, J.M., et al., *The endogenous Toll–like receptor 4 agonist S100A8/S100A9
(calprotectin) as innate amplifier of infection, autoimmunity, and cancer.* Journal of
Leukocyte Biology, 2009. **86**(3): p. 557-566.
39. Vogl, T., et al., *MRP8 and MRP14 control microtubule reorganization during transendothelial
migration of phagocytes.* Blood, 2004. **104**(13): p. 4260-4268.
40. Roth, J., et al., *MRP8 and MRP14, S-100-like proteins associated with myeloid differentiation,
are translocated to plasma membrane and intermediate filaments in a calcium-dependent
manner.* Blood, 1993. **82**(6): p. 1875-1883.
41. Goebeler, M., et al., *Increase of calcium levels in epithelial cells induces translocation of
calcium-binding proteins migration inhibitory factor-related protein 8 (MRP8) and MRP14 to
keratin intermediate filaments.* Biochem J, 1995. **309 (Pt 2)**(Pt 2): p. 419-24.
42. Lominadze, G., et al., *Myeloid-Related Protein-14 Is a p38 MAPK Substrate in Human
Neutrophils1.* The Journal of Immunology, 2005. **174**(11): p. 7257-7267.
43. Wolf, M., et al., *S100A8/S100A9 Integrates F-Actin and Microtubule Dynamics to Prevent
Uncontrolled Extravasation of Leukocytes.* Biomedicines, 2023. **11**(3): p. 835.
44. Clemens, R.A. and C.A. Lowell, *Store-operated calcium signaling in neutrophils.* Journal of
Leukocyte Biology, 2015. **98**(4): p. 497-502.
45. Simon, S.I., et al., *Leucocyte recruitment under fluid shear: mechanical and molecular
regulation within the inflammatory synapse.* Clinical and Experimental Pharmacology and
Physiology, 2009. **36**(2): p. 217-224.

46. Kummerow, C., et al., *The immunological synapse controls local and global calcium signals in*
T lymphocytes. Immunol Rev, 2009. **231**(1): p. 132-47.
47. Hobbs Josie, A.R., et al., *Myeloid Cell Function in MRP-14 (S100A9) Null Mice.* Molecular and
Cellular Biology, 2003. **23**(7): p. 2564-2576.
48. Mizgerd, J.P., et al., *Neutrophil Emigration in the Skin, Lungs, and Peritoneum: Different*
Requirements for CD11/CD18 Revealed by CD18-deficient Mice. Journal of Experimental
Medicine, 1997. **186**(8): p. 1357-1364.
49. Buscher, K., et al., *Protection from septic peritonitis by rapid neutrophil recruitment through*
omental high endothelial venules. Nature Communications, 2016. **7**(1): p. 10828.
50. Wen, L., M. Moser, and K. Ley, *Molecular mechanisms of leukocyte β2 integrin activation.*
Blood, 2022. **139**(24): p. 3480-3492.
51. Vogl, T., et al., *Alarmin S100A8/S100A9 as a biomarker for molecular imaging of local*
inflammatory activity. Nature Communications, 2014. **5**(1): p. 4593.
52. Immler, R., et al., *Extratubular Polymerized Uromodulin Induces Leukocyte Recruitment and*
Inflammation In Vivo. Front Immunol, 2020. **11**: p. 588245.
53. Schindelin, J., et al., *Fiji: an open-source platform for biological-image analysis.* Nat Methods,
2012. **9**(7): p. 676-82.
54. Ley, K. and P. Gaehtgens, *Endothelial, not hemodynamic, differences are responsible for*
preferential leukocyte rolling in rat mesenteric venules. Circ Res, 1991. **69**(4): p. 1034-41.
55. Sperandio, M., et al., *Analysis of Leukocyte Rolling In Vivo and In Vitro*, in *Methods in*
Enzymology. 2006, Academic Press. p. 346-371.
56. Zehrer, A., et al., *A Fundamental Role of Myh9 for Neutrophil Migration in Innate Immunity.*
The Journal of Immunology, 2018. **201**(6): p. 1748.
57. Schymeinsky, J., et al., *A fundamental role of mAbp1 in neutrophils: impact on β2 integrin–*
mediated phagocytosis and adhesion in vivo. Blood, 2009. **114**(19): p. 4209-4220.
58. Grimes, D., et al., *ORAI1 and ORAI2 modulate murine neutrophil calcium signaling, cellular*
activation, and host defense. Proceedings of the National Academy of Sciences of the United
States of America, 2020. **117**(39): p. 24403-24414.
59. Pedregosa, F., et al., *Scikit-learn: Machine Learning in Python.* Journal of Machine Learning
Research, 2011.
60. Postić, S., et al., *High-resolution analysis of the cytosolic Ca2+ events in β cell collectives in*
situ. American Journal of Physiology-Endocrinology and Metabolism, 2023. **324**(1): p. E42-
E55.

Acknowledgements

We thank Dorothee Gössel, Anke Lübeck, Sabine D'Avis, Susanne Bierschenk and Jennifer

Troung for excellent technical assistance, and the core facility Flow Cytometry at the

Biomedical Center, LMU, Planegg-Martinsried, Germany.

Funding

This work was supported by the German Research Foundation (DFG) collaborative research

grants: SFB914, projects A02 (B.W.), B01 (M.S.), B11 (M.P.), TRR-332 projects C2 (M.S.), C3

(B.W.), C7 (T.V.), B5 (J.R.) and the TRR-359 project B2 (M.S. and R.I.).

Author contributions

M.N. designed and conducted experiments, analyzed and interpreted data, and wrote the

manuscript. R.I. and I.R. designed experiments, acquired and analyzed data V.L. performed

the integrin spatial cluster analysis. J.P. performed the calcium frequency distribution analysis.

M.Sa and A.Y. analyzed results. T.V. provided critical reagents (S100A8/A9^{mut}, anti-S100A9-

Cy5.5) and analyzed ELISA samples. J.R., M.S.R, C.M., and B.W. provided their expertise and

conceptual advice. M.S. designed experiments and wrote the manuscript. M.P. designed

experiments, wrote the manuscript and supervised the work. All authors discussed the results,

commented on and approved the manuscript.

Competing interests

The authors declare no competing interests.

Data and materials availability

All data and materials that support the findings of this study are available from the

corresponding author upon request. Custom codes developed for data analysis and

visualization   are   available   at   https://github.com/Napo93/AG-Sperandio-MACROS,

https://github.com/marrlab/Spatial_CA#spatial_ca   and

https://github.com/szarma/Physio_Ca. Software and parameters used are stated in the

Methods with further details.

FIGURE LEGENDS

Figure 1: Cytosolic S100A8/A9 regulates leukocyte recruitment in vivo regardless of extracellular S100A8/A9

(A) ELISA measurements of S100A8/A9 levels in supernatants of WT bone marrow neutrophils stimulated for 10min with PBS, E-selectin or lysed with Triton X-100(mean+SEM, *n*=6 mice per group, RM one-way ANOVA, Holm-Sidak's multiple comparison). (B) Schematic model of the mouse cremaster muscle preparation for intravital microscopy and representative picture of a vessel showing rolling and adherent cells. WT and *Mrp14*$^{-/-}$ mice were stimulated i.s. with TNF-α 2h prior to cremaster muscle post-capillary venules imaging by intravital microscopy. Quantification of (C) number or rolling (rolling flux fraction) and (D) number of adherent neutrophils per vessel surface of WT and *Mrp14*$^{-/-}$ mice [mean+SEM, *n*=5 mice per group, 25 (WT) and 30 (*Mrp14*$^{-/-}$) vessels, unpaired Student's *t*-test]. (E) Correlation between physiological vessel shear rates and number of adherent neutrophils in WT and *Mrp14*$^{-/-}$ mice [*n*=25 (WT) and 30 (*Mrp14*$^{-/-}$) vessels of 5 mice per group, Pearson correlation]. (F) Schematic model of sterile inflammation induced by exteriorizing WT and *Mrp14*$^{-/-}$ cremaster muscles. (G) Analysis of number of adherent leukocytes by intravital microscopy before and after S100A8/A9^{mut} intra-arterial injection [mean+SEM, *n*=3 mice per group, 3 (WT) and 3 (*Mrp14*$^{-/-}$) vessels, 2way ANOVA, Sidak's multiple comparison]. (H) Representative Giemsa staining micrographs of TNF-α stimulated WT and *Mrp14*$^{-/-}$ cremaster muscles (representative micrographs, scale bar =30µm, arrows: transmigrated neutrophils) and (I) quantification of number of perivascular neutrophils [mean+SEM, *n*=5 mice per group, 56 (WT) and 55 (*Mrp14*$^{-/-}$) vessels, unpaired Student's *t*-test]. ns, not significant; *p≤0.05, **p≤0.01, ***p≤0.001.

Figure 2 Loss of cytosolic S100A8/A9 impairs neutrophil adhesion under flow conditions without affecting β₂ integrin activation

(A) Schematic representation of blood harvesting from WT and $Mrp14^{-/-}$ mice via a carotid artery catheter and perfusion into self-made flow cambers coated with E-selectin, ICAM-1, and CXCL1. Analysis of (B) number of rolling and (C) number of adherent leukocytes FOV^{-1}[mean+SEM, n=4 mice per group, 10 (WT) and 12 ($Mrp14^{-/-}$) flow chambers, paired Student's t-test]. (D) Schematic representation of the soluble ICAM-1 binding assay using bone marrow neutrophils stimulated with PBS control or CXCL1 (10nM) assessed by (E) flow cytometry (MFI=median fluorescence intensity, mean+SEM, n=5 mice per group, 2way ANOVA, Sidak's multiple comparison). (F) Spectroscopy fluorescence intensity analysis of percentage of adherent WT and $Mrp14^{-/-}$ neutrophils, seeded for 5min on ICAM-1 coated plates and stimulated with PBS or CXCL1 (10nM) for 10min (mean+SEM, n=4 mice per group, 2way ANOVA, Sidak's multiple comparison). ns, not significant; *p≤0.05, **p≤0.01, ***p≤0.001.

Figure 3: Cytosolic S100A8/A9 is crucial for neutrophil spreading, crawling and post-arrest modifications under flow

(A) Representative bright-field pictures of WT and $Mrp14^{-/-}$ neutrophils spreading over E-selectin, ICAM-1, and CXCL1 coated glass capillaries (scale bar=10µm). Analysis of cell shape parameters (B) area, perimeter, (C) circularity [4π * (Area/Perimeter)] and solidity [Area/Convex area] over time [mean+SEM, n=103 (WT) and 96 ($Mrp14^{-/-}$) neutrophils of 4 mice per group, unpaired Student's t-test]. (D) Rose plot diagrams representative of migratory crawling trajectories of WT and $Mrp14^{-/-}$ neutrophils in flow chambers coated with E-selectin, ICAM-1, and CXCL1 under flow (2dyne cm^{-2}). Analysis of (E) crawling distance, (F) directionality of migration and (G) crawling velocity of WT and $Mrp14^{-/-}$ neutrophils [mean+SEM, n=5 mice per group, 113 (WT) and 109 ($Mrp14^{-/-}$) cells, paired Student's t-test]. Western blot analysis of ICAM-1 induced (H) Pyk2 and (I) Paxillin phosphorylation of WT and $Mrp14^{-/-}$ neutrophils upon CXCL1 stimulation (10nM) (mean+SEM, representative western blot of n≥4 mice per group,

2way ANOVA, Sidak's multiple comparison). ns, not significant; *p≤0.05, **p≤0.01, ***p≤0.001.

Figure 4: Cytosolic S100A8/A9 drives neutrophil cytoskeletal rearrangement by regulating LFA-1 nanocluster formation and Ca²⁺ availability within the clusters

(A) Representative confocal images of LFA-1 staining in WT ^{Lyz2xGCaMP5} and $Mrp14^{-/-}$ ^{Lyz2xGCaMP5} crawling neutrophils on E-selectin, ICAM-1, and CXCL1 coated flow chambers (scale bar=10µm). (B) Segmentation of LFA-1 signals through automatic thresholding (scale bar=10µm). (C) Size-excluded LFA-1 nanoclusters of 0.15µm² minimum size from previously thresholded images (scale bar=10µm). (D) Single cell analysis of average number of LFA-1 nanoclusters in min 0-1, 5-6 and 9-10 of analysis of WT ^{Lyz2xGCaMP5} and $Mrp14^{-/-}$ ^{Lyz2xGCaMP5} neutrophils [mean+SEM, n=5 mice per group, 56 (WT) and 54 ($Mrp14^{-/-}$) neutrophils, 2way ANOVA, Sidak's multiple comparison]. (E) Representative confocal images of Ca²⁺ signals in WT ^{Lyz2xGCaMP5} and $Mrp14^{-/-}$ ^{Lyz2xGCaMP5} neutrophils (scale bar=10µm) and (F) Ca²⁺ signals in the previously segmented LFA-1 nanoclusters (scale bar=10µm). (G) Quantification of subcellular Ca²⁺ levels in the LFA-1 nanocluster area in min 0-1, 5-6 and 9-10 in WT ^{Lyz2xGCaMP5} and $Mrp14^{-/-}$ ^{Lyz2xGCaMP5} neutrophils [mean+SEM, n=5 mice per group, 56 (WT) and 54 ($Mrp14^{-/-}$) cells, 2way ANOVA, Sidak's multiple comparison]. (H) Segmented LFA-1 cluster negative areas (scale bar=10µm) and (I) representative confocal images of Ca²⁺ signals in the LFA-1 cluster negative areas (scale bar=10µm) [Fig. 4E-I scale bar color code: 0= black, 255=white]. (J) Analysis of cytosolic Ca²⁺ levels in the LFA-1 cluster negative areas in min 0-1, 5-6 and 9-10 of WT ^{Lyz2xGCaMP5} and $Mrp14^{-/-}$ ^{Lyz2xGCaMP5} neutrophils [mean+SEM, n=5 mice per group, 56 (WT) and 54 ($Mrp14^{-/-}$) neutrophils, 2way ANOVA, Sidak's multiple comparison]. (K) Representative confocal images showing S100A9 localization at LFA-1 nanocluster areas in stimulated WT neutrophils (scale bar = 10 µm). (L) Quantitative analysis of S100A9 levels in positive LFA-1

nanocluster areas compared to non LFA-1 nanocluster areas in stimulated WT neutrophils.

[mean+SEM, n=3 mice, 26 (WT) neutrophils, paired Student's t-test]. (M) Representative

confocal micrographs of LFA-1 nanocluster spatial aggregation in WT ^{Lyz2xGCaMP5} and $Mrp14^{-/-}$

^{Lyz2xGCaMP5} neutrophils, within 10µm^2 area and minimum 10 LFA-1 nanoclusters considered ($\geq$

10 LFA-1 nanoclusters within 10µm^2, yellow circles=spatial aggregation area, scale bar=10µm). (N)

Analysis of spatially aggregated LFA-1 nanoclusters of WT ^{Lyz2xGCaMP5} and $Mrp14^{-/-}$ ^{Lyz2xGCaMP5}

neutrophils [mean+SEM, n=5 mice per group, 56 (WT) and 54 ($Mrp14^{-/-}$) cells, unpaired

Student's t-test]. (O) Segmentation of WT ^{Lyz2xGCaMP5} and $Mrp14^{-/-}$ ^{Lyz2xGCaMP5} neutrophil area

through $Lyz2$ channel automatic thresholding and (P) representative confocal images of

respective F-actin signals. (Q) Analysis of F-actin intensity normalized to the cell area in min 0-

1, 5-6 and 9-10 of WT ^{Lyz2xGCaMP5} and $Mrp14^{-/-}$ ^{Lyz2xGCaMP5} neutrophils [mean+SEM, n=5 mice per

group, 74 (WT) and 66 ($Mrp14^{-/-}$) cells, 2way ANOVA, Sidak's multiple comparison]. ns, not

significant; *p$\leq$0.05, **p$\leq$0.01, ***p$\leq$0.001.

Figure 5: Cytosolic S100A8/A9 is dispensable for chemokine induced ER store Ca^{2+} release

and for the initial phase of SOCE (A) Average flow cytometry kinetic graphs of Ca^{2+} store

release in the absence of extracellular Ca^{2+} (Ca^{2+} free medium) in WT and $Mrp14^{-/-}$ neutrophils

upon CXCL1 stimulation (traces are shown as mean+SEM, n=5 mice per group). (B) Rapid ER

store Ca^{2+} release (MFI $_{peak}$/ MFI $_{0-30s}$) of WT and $Mrp14^{-/-}$ neutrophils [mean+SEM, n=5 mice

per group, paired Student's t-test]. (C) Quantification of Ca^{2+} levels under baseline conditions

(MFI $_{0-30s}$) [mean+SEM, n=5 mice per group, paired Student's t-test]. (D) Average flow

cytometry kinetic graphs of Ca^{2+} influx in the presence of extracellular Ca^{2+} (HBSS medium,

1.5mM Ca^{2+}) of WT and $Mrp14^{-/-}$ neutrophils upon CXCL1 stimulation (traces are shown as

mean+SEM, n=5 mice per group, double-headed arrow represents the time points of

quantification). (E) Ca^{2+} levels before CXCL1 stimulation (MFI $_{0-30s}$) [mean+SEM, n=5 mice per

group, paired Student's *t*-test]. (F) Quantification of ER store Ca^{2+} release and calcium released

activated channel (CRAC) store-operated Ca^{2+} entry (MFI_{peak}/ MFI_{0-30s}) [mean+SEM, *n*=5 mice

per group, paired Student's *t*-test]. (G) Ca^{2+} influx after CXCL1 stimulation, from peak to peak

half-life ($AUC_{peak - \frac{1}{2} peak}$) of WT and *Mrp14*$^{-/-}$ neutrophils [mean+SEM, *n*=5 mice per group,

paired Student's *t*-test]. ns, not significant; *p≤0.05, **p≤0.01, ***p≤0.001.

FIGURE 1

FIGURE 2

A

blood flow

rolling

adhesion

B

Leukocyte rolling *ex vivo*

ns

[Rolling cells FOV⁻¹] / WBC

WT   *Mrp14⁻/⁻*

C

Leukocyte adhesion *ex vivo*

*

[Adherent cells FOV⁻¹] / WBC

WT   *Mrp14⁻/⁻*

D

High-affinity LFA-1
Inactive LFA-1
ICAM-1
α-ICAM-1 Ab

PBS   CXCL1

E

ICAM-1 binding

WT
Mrp14⁻/⁻

MFI / Isotype control

**   **

CXCL1   -   +   -   +

F

Leukocyte static adhesion

WT
Mrp14⁻/⁻

Adhesion [% total inserted cells]

***   ***

CXCL1   -   +   -   +

FIGURE 3

1052

1053

1054

1055

1056

1057

1058

1059

1060

1061

1062

1063

1064

1065

1066

1067

1068

1069

1070

1071

 FIGURE 4

1096

K

1097

1098

1099

1100

1101

L

S100A9 spatial distribution upon stimulation

1102

M

WT *Lyz2xGCaMP5*   *Mrp14⁻/⁻ Lyz2xGCaMP5*

LFA-1 nanoclusters

1103

1104

1105

N

LFA-1 spatial aggregation

1106

1107

O   **P**

Cell mask   F-actin

1108

WT *Lyz2xGCaMP5*

1109

1110

Mrp14⁻/⁻ Lyz2xGCaMP5

1111

1112

WT *Lyz2xGCaMP5*
Mrp14⁻/⁻ Lyz2xGCaMP5

Q

F-Actin

1113

1114   **FIGURE 5**

1115

1140

1141