Cytoadhesion of *Plasmodium falciparum*-infected red blood cells changes the expression of cytokine-, histone- and antiviral protein-encoding genes in brain endothelial cells

Johannes Allweier[1,2†], Michael Bartels[1†], Hanifeh Torabi[1], Maria del Pilar Martinez Tauler[1], Nahla Galal Metwally[1], Thomas Roeder[3, 4], Thomas Gutsmann[2], Iris Bruchhaus[1,5*]

[1]Research group Host-Parasite Interaction, Bernhard Nocht Institute for Tropical Medicine, Hamburg Germany

[2]Division of Biophysics, Research Center Borstel, Leibniz-Center for Medicine and Biosciences, Borstel, Germany

[3]Kiel University, Department Molecular Physiology, Zoology, Kiel, Germany

[4]DZL, German Center for Lung Research, ARCN, Airway Research Center North, Kiel, Germany

[5]Department of Biology, University of Hamburg, Hamburg, Germany

[†] These authors contributed equally to this work and share first authorship.

* Correspondence:
Iris Bruchhaus
bruchhaus@bnitm.de

Keywords: *Plasmodium falciparum*, malaria, endothelial cells, cytoadhesion, transcriptome.

Abstract

Introduction: Malaria remains a significant global health problem, particularly due to the human malaria parasite *Plasmodium falciparum*, which is responsible for most fatal infections. Infected red blood cells (iRBCs) evade spleen clearance by adhering to endothelial cells (ECs), triggering capillary blockage, inflammatory cytokine release, endothelial dysfunction, and altered vascular permeability, prompting an endothelial transcriptional response.

Methods: The iRBCIT4var04/HBEC-5i model, where iRBCs present IT4var04 (VAR2CSA) on their surface was employed to analyse the effects of iRBC binding on ECs. We used this model to investigate how cytoadhesion of iRBCs to ECs influences their expression profile depending on the temperature (37°C vs 40°C).

Results: Binding of non-infected RBCs (niRBCs) and fever alone significantly changes expression of hundreds of genes in ECs. Comparing the expression profile of HBEC-5i cultured either in the presence of iRBCs or in the presence of niRBCs, genes encoding proteins assigned to the GO terms immune response, nucleosome assembly, NF-kappa B signaling, angiogenesis, and antiviral immune response/interferon-alpha/beta signaling pathway were significantly up-regulated. If the cultivation temperature is increased from 37°C to 40°C, which simulates fever, a further significant increase in expression can be observed for most regulated genes, especially for genes coding for cytokines and proteins involved in angiogenesis.

Conclusion: The presence of iRBCs leads to the stimulation of ECs, activating several immunological signaling pathways and affecting antiviral (-parasitic) mechanisms and angiogenesis. Furthermore, to our knowledge, the induction of the interferon-alpha/beta signaling pathway in ECs in response to iRBCs has been described for the first time.

1   Introduction

Despite recent success stories in vaccine development and roll-outs, malaria remains one of the world's most dangerous infectious diseases. In 2022, there were about 249 million malaria cases, including 608,000 deaths (WHO, 2023). Most fatal malaria infections are caused by *Plasmodium falciparum*. Parasite and host responses contribute to the development of complications characteristic of severe malaria infections. A vital aspect of the pathogenesis of *P. falciparum* infection is the cytoadhesion of infected red blood cells (iRBCs) to the vascular bed of vital organs such as the brain, heart, lungs, stomach, skin, and kidneys (Taylor et al., 2004; Milner et al., 2013; Milner et al., 2015). In addition to capillary blockage, iRBC cytoadhesion induces inflammatory cytokine production, endothelial dysfunction, and vascular permeability in the affected tissue (Lyke et al., 2004). As a result of the immune responses triggered by parasite growth and endothelial cytoadhesion, patients may develop fever, headache, myalgia, and muscle stiffness (Hasday et al., 2001; Oakley et al., 2007; Oakley et al., 2011; Cunnington et al., 2013; Phillips et al., 2017). Depending on the age and immune status of the patient, severe to life-threatening complications may develop, including cerebral malaria, organ failure, acidosis, and severe anemia (Cunnington et al., 2013; Gazzinelli et al., 2014; Phillips et al., 2017).

Cytoadhesion is primarily mediated by members of the *P. falciparum* erythrocyte membrane protein 1 (*Pf*EMP1) family, which is encoded by approximately 45-90 *var* genes per parasite genome (Otto et al., 2019). The expression of *var* genes is mutually exclusive in ring-stage parasites, so only one *Pf*EMP1 variant is present on the surface of trophozoite- and schizont-stage iRBCs at any one time. The *var* genes encoding *Pf*EMP1 are highly variable from parasite to parasite, resulting in a vast repertoire of *var* genes in nature (Kraemer and Smith, 2006; Voss et al., 2006; Kyes et al., 2007; Pasternak and Dzikowski, 2009; Deitsch and Dzikowski, 2017). *Pf*EMP1s consist of an intracellular and a transmembrane domain, several extracellular Duffy binding-like (DBL) domains (α, β, γ, δ, ζ, and ε), and cysteine-rich interdomain regions (CIDR; α, β, and γ) (Smith et al., 2000a; Smith et al., 2000b; Gardner et al., 2002; Rask et al., 2010) and can be classified into groups A, B, C, and E (Kraemer and Smith, 2003; Lavstsen et al., 2003; Kraemer et al., 2007; Rask et al., 2010). *Pf*EMP1s are concentrated in electron-dense protrusions of the erythrocyte plasma membrane called knobs. Knobs are composed of several submembrane structural proteins, including the major protein of this structure, the knob-associated histidine-rich protein (KAHRP). Moreover, other proteins are required for knob formation, including *Pf*EMP3, ring-infected red cell antigen (RESA), mature parasite-infected red cell surface antigen (MESA)/*Pf*EMP2, and *Pf*332 (Gruenberg et al., 1983; Maier et al., 2009; Tilly et al., 2015; Alampalli et al., 2018). Knobs play an essential role in binding iRBCs to the endothelium (Crabb et al., 1997; Horrocks et al., 2005; Cutts et al., 2017; Dorpinghaus et al., 2020; Lubiana et al., 2020). If iRBCs do not form knobs, e.g. due to loss of genes encoding knob structural proteins, adhesion under flow conditions is significantly impaired. Especially at febrile temperatures, they play a crucial role by stabilizing the binding interaction. Under these conditions, knob-negative iRBCs are no longer able to bind to various endothelial cell receptors (ECRs) (Lubiana et al., 2020).

Cytoadhesion of iRBCs also leads to a change in gene expression in endothelial cells (ECs) (Tripathi et al., 2006; Tripathi et al., 2009). Stimulation of primary human brain microvascular endothelial cells (HBMECs) through cytoadhesion of iRBCs resulted in increased ICAM-1 expression on the surface of HBMECs, which in turn resulted in increased cytoadhesion. This observation was specific to HBMECs, in contrast to human umbilical vein endothelial cells (HUVECs), where no significant changes in gene expression were observed (Tripathi et al., 2006). Cytoadhesion induced nuclear translocation of NF-κB in HBMECs, which was associated with ICAM-1 expression (Tripathi et al., 2006). In another study, the influence of cytoadhesion on HBMEC gene expression was investigated by microarray analysis (Tripathi et al., 2009). The effect of binding of iRBCs with enhanced binding

on HBMECs induced by panning was compared with the effects of binding of unpanned iRBCs. For
both populations, a 4-fold increase in the amount of ICAM-1 presented on the HBMEC surface was
detected after 6 h of co-incubation. Interestingly, there were no significant differences in gene
expression between the two parasite populations. In total, the authors identified 231 up-regulated and
155 down-regulated genes. Using GO term analysis, the proteins encoded by the regulated genes were
shown to be involved in inflammatory response, apoptosis, anti-apoptosis, cell-cell signaling and
transduction, and NF-κB activation cascade. The expression of some genes encoding pro-inflammatory
cytokines such as IL-6, CXCL8, CXCL1, CXCL2, and CCL20 was increased more than 100-fold, and
for some of them, the increase was also detected at the protein level (Tripathi et al., 2009). Recently,
the transcriptome profile of HBMECs co-cultured for different time points with iRBCs presenting
IT4_var14 (strong binding to ICAM-1) or IT4_var37 (strong binding to CD36) on their surface was
analysed. Genes coding for proteins involved in many of the pathways described above, such as
inflammation, cell-cell interaction, and apoptosis, were identified as differentially expressed. However,
significant differences were also found between iRBC^{IT4var14} and iRBC^{IT4var37}. Among others, atonal
bHLH transcription factor 8 (*ATOH8*), *CXCL8*, *CXCL10*, *VCAM-1*, radical s-adenosyl methionine
domain containing 2 (*RSAD2*), and IL6 were found to be up-regulated in the presence of iRBC^{IT4var14}
after 6 h of co-cultivation, whereas the expression of these genes was down-regulated in iRBC^{IT4var37}
(Othman et al., 2023).

Chakravorty and colleagues also investigated the influence of cytoadhesion on HUVEC gene
expression (Chakravorty et al., 2007). They confirmed the observation of Tripathi and colleagues that
cytoadhesion of iRBCs alone is not sufficient to stimulate HUVEC cells (Tripathi et al., 2006).
However, when co-incubated in the presence of 5 pg/mL of TNF, there is an increase in the presentation
of ICAM-1 on the cell membrane. Using microarray analysis, they showed that genes encoding proteins
involved in cell communication, cell adhesion, signal transduction, and immune response were
regulated. Thus, HUVEC can mobilize immune and adhesion responses, but only when co-incubated
with low concentrations of TNF in addition to iRBCs cytoadhesion (Chakravorty et al., 2007).
However, there is another study in this context that has shown stimulation of HUVECs even without
the addition of TNF (Viebig et al., 2005). They were able to show that stimulation of HUVECs with
iRBCs of the *P. falciparum* isolate IT4 results in more ICAM-1 being presented on the surface of the
ECs and increased amounts of IL-6 being secreted, whereas this is not the case for VCAM-1.
Furthermore, significantly increased expression of various cytokines such as CXCL8, CCL20, CXCL1,
and CCL2 was detected (Viebig et al., 2005).

In previous studies, an immortalized human cerebral microvascular endothelial cell line (HBEC-5i)
was used to characterize the cytoadhesion of iRBCs (Dorpinghaus et al., 2020; Lubiana et al., 2020).
HBEC-5i cells are derived from the microvasculature of the human cerebral cortex and therefore have
essential characteristics of cerebral ECs. HBEC-5i cells present Willebrand factor (vWF), VE-
cadherin, occludin, VCAM-1, ICAM-1, CD54, CD40, and chondroitin-sulfate A (CSA) on the cell
surface. However, the binding of iRBCs of the *P. falciparum* isolate IT4 to HBEC-5i cells was shown
to occur mainly via CSA. Parasites enriched for binding to HBEC-5i cells present IT4var04 (var2CSA)
on the surface of iRBCs (iRBCs^{IT4var04}), of which CSA is known to be the ligand. Thus, CSA seems to
be dominant on HBEC-5i cells and that binding of iRBCs to other receptors such as ICAM-1 is shielded
(Dorpinghaus et al., 2020). Using this model (iRBC^{IT4var04}/HBEC-5i), it was shown that binding to
CSA occurs only at a low shear stress of 0.9 dyne/cm^2 with a static binding characteristic. iRBC
adhesion to CD36 on the other hand displayed a rolling behavior. Additionally, it could be shown that
binding at febrile temperatures can only occurs when the iRBCs have formed knobs (Lubiana et al.,
2020).

In the present study, the iRBC^IT4var04/HBEC-5i model was used to investigate whether stimulation of ECs by binding of iRBCs to CSA is possible and, if so, whether and how this stimulation differs from that of HBMEC and HUVEC and what happens in the presence of fever. The aim was to determine whether cytoadhesion in general elicits similar responses in ECs, independent of the receptor. Moreover, we also studied the effects of febrile temperatures due to the co-incubation with iRBCs.

Our results show that cytoadhesion of iRBCs increases the expression of genes in HBEC-5i cells encoding proteins involved in immune response, angiogenesis, antiviral mechanisms and nucleosome assembly. In particular, under febrile conditions, the expression of some cytokines is strongly increased, suggesting the formation of a cytokine storm.

2 Materials and methods

2.1 Cell culture

P. falciparum isolate IT4 (FCR3S1.2) was cultured in human 0+ erythrocytes at a hematocrit of 5% in the presence of 10% human A+ serum according to standard procedures (Trager and Jensen, 2005). Parasites were synchronized once a week with 5% sorbitol (Lambros and Vanderberg, 1979). Every two weeks, iRBCs were enriched for the presence of knobs. IRBCs were resuspended in two volumes of pre-warmed gelatin (1%, 37°C) and incubated at 37°C for 45 min (Waterkeyn et al., 2001). The supernatant containing knob-positive iRBCs was washed with GibcoTM RPMI 1640 medium (Gibco, Thermo Fisher Scientific, Bremen, Germany) and cultured as usual.

The HBEC-5i cell line (ATCC CRL-3245) was cultured in DMEM/F12 (Gibco, Thermo Fisher Scientific, Bremen, Germany) medium supplemented with L-glutamine and sodium bicarbonate (2.438 g/l) at 37°C in 5% CO_2 and passaged every 2-4 days at 70-90% confluence.

2.2 Enrichment of iRBCs that bind to HBEC-5i

To enrich iRBCs that bind to HBEC-5i cells, synchronized iRBCs at the trophozoite stage (5-10% parasitemia, 1% hematocrit) were resuspended in binding medium (RPMI 1640 medium, 2% glucose, pH 7.2; Gibco, Thermo Fisher Scientific, Bremen, Germany) and co-incubated with HBEC-5i cells (70-90% confluence) for 90 min at 37°C and 5% CO_2. After 60, 75, and 90 min, the cultures were gently shaken orbitally. This was followed by five washes with a binding medium to remove any unbound iRBCs. The remaining parasites were further cultured with 10 ml of RPMI 1640 medium and erythrocytes (5% hematocrit) to the ring stage and harvested. Any detached HBEC-5i cells were removed with Biocoll separation solution (Merck Biochrom, Berlin, Germany). The parasites were then grown under standard conditions to a parasitemia of 5-10%, and the whole procedure was repeated seven times. Parasites enriched for binding to HBEC-5i cells were shown to express Var2CSA (IT4_var04), which binds to CSA presented on HBEC-5i cells (Dorpinghaus et al., 2020).

2.3 Infected RBC-HBEC-5i co-incubation, RNA isolation, and RNAseq

To analyse the influence of cytoadhesion on HBEC-5i cell metabolism, HBEC-5i cells (70-90% confluence) were co-incubated with HBEC-5i-enriched trophozoite-stage iRBCs at a parasitemia of 5% (hematocrit 1%) in binding medium (Dorpinghaus et al., 2020). Co-incubation was performed in T-25 flasks at either 37°C or 40°C (iRBC^37°C/iRBC^40°C). HBEC-5i cells cultured at 37°C or 40°C +/- niRBCs (hematocrit 1%) served as controls. After seven h of co-incubation, cells were collected in 4 ml of pre-warmed TRIzol (Thermo Fisher Scientific, Bremen, Germany), thoroughly resuspended, transferred to a 15 ml reaction tube, and incubated in a water bath at 37°C for 5 min. Samples could

then be stored at -80°C or processed immediately. For RNA isolation, samples of 1 ml each were divided into 4 RNase-free reaction tubes, supplemented with 200 µl of chloroform cooled to 4°C, shaken well, and incubated for 3 min at room temperature (RT). Samples were then centrifuged at 4°C for 30 min at 12,000 $x\ g$, and the supernatant was transferred to a new reaction tube. RNA was precipitated by the addition of an equivalent volume of 70% ethanol. RNA was then isolated using the PureLink RNA Mini Kit (Thermo Fisher Scientific, Bremen, Germany) according to the manufacturer's instructions. Possible contamination with genomic DNA was removed using the TURBO DNA-free kit (Fisher Scientific, Thermo Fisher, Hampton, NH, USA). RNA was then purified using the Agencourt RNAClean XP Kit (Beckham Coulter, Brea, CA, USA). Library preparation and subsequent RNA sequencing was performed on an Illumina HiSeq 4000 PE100 platform from BGI (BGI Group, Shenzhen, China).

2.4 Bioinformatic analyses/Statistics

The RNAseq data were mapped to the human reference genome (GRCh38.109) using CLC Genomics Workbench 21.0 (QIAGEN, Aarhus, Denmark). The Voroni diagram was created using Python 3.11 and matplotkib 3.7.2. The interaction network analysis was performed with the database STRING, version 12.0 (Szklarczyk et al., 2023). The following setting was used: Active interaction sources: text mining, experiments, databases, co-expression; Meaning of network edges: high confidence 0.700. Network display mode: Interactive; Network display options: Hide disconnected nodes in the network; Clustering Options: MCL clustering, inflation parameter 3; Edges between clusters: Solid lines.

3 Results

3.1 Effects of niRBCs, iRBCs, and fever on the expression of HBEC-5i cells

By performing a full transcriptome analysis of the ECs, we examined the impact of niRBCs, iRBCs, and fever on the expression profile of HEBEC-5i cells. For these studies, as mentioned above, plasmodia of isolate IT4 were used, which were enriched on HBEC-5i cells over several rounds, with adhesion taking place via CSA, thus presenting IT4var04 (var2CSA) on the surface of the iRBCs (Dorpinghaus et al., 2020). A Venn diagram was constructed to illustrate the differential expression (true vs. false) of the three parameters niRBC, iRBC, and febrile (Figure 1). Specifically, genes exhibiting a differential expression of a fold change of ≥ 2.0, a false discovery rate (FDR) of ≤ 0.05, and a minimum transcript per million (TPM) threshold of ≥ 2.0 were included in this analysis. Our findings reveal distinct regulatory effects: 60 genes were solely modulated by the presence of niRBCs, while elevating the temperature from 37°C to 40°C alone resulted in altered expression patterns in 98 genes. Similarly, the presence of iRBCs alone influenced the expression of 318 genes. Notably, 285 genes exhibited regulation in response to either niRBCs or iRBCs, with 80 genes showing such regulation specifically in the presence of iRBCs[37°C] and iRBCs[40°C]. Additionally, ten genes were differentially expressed either by the febrile stimulus or presences of niRBCs. Seventy-nine genes were found that overlap in all three comparisons (Figure 1, Supplementary Table 1).
Of the 59 genes that are significantly differentially expressed by the presence of niRBCs, 38 are up-regulated and 21 are down-regulated. With minor exceptions, the majority of these genes demonstrate a 2- to 3-fold regulation. Notably, the most prominently up-regulated gene, BOLA2, exhibits a striking 79-fold increase (from TPM 2.5 to 198.3) in response to niRBCs presence (Figure 1, Supplementary Table 1).
Of the 98 genes that are significantly differentially expressed by the temperature elevation from 37°C to 40°C, 50 display up-regulation, while 48 undergo down-regulation. Once again, most of these genes

show a 2- to 3-fold regulation. The increase in temperature upregulates the expression of 10 genes associated with the Gene Ontology Biological Process (GO-BP) term "protein folding" (FDR 4.82e-16) (Figure 1, Figure 2A, Supplementary Table 1).

Analysing the 318 genes significantly differentially expressed by the presence of iRBCs, 121 are up-regulated, while 197 are down-regulated (Supplementary Table 1). Approximately 90% of these genes undergo a 2- to 3-fold regulation, while the remaining 10% (22 genes) exhibit a 3- to 6-fold regulation. String analysis reveals the up-regulation of GO-BP terms such as "defense response to virus" (FDR 7.75e-09), "nucleosome assembly" (FDR 0.00013), "NF-κB complex" (FDR 0.0018), and "ribosome biogenesis" (FDR 0.00086) (Figure 1, Figure 2B, Supplementary Table 1).

Within the intersection of genes derived from the analyses of niRBC true vs. false and iRBCs true vs. false, a total of 285 genes exhibit regulation. Noteworthy among these are 13 genes associated with "cellular response to cytokine stimulus" (FDR 4.82e-10), 9 genes linked to "nucleosome assembly" (FDR 2.56e-15), 11 genes involved in "I-κB kinase/NF-κB signaling" (FDR 4.32e-06), 5 genes contributing to "negative regulation of I-κB kinase/NF-κB signaling" (FDR 0.0017), and 3 genes associated with "defense response to virus" (FDR 3.38e-05) GO-BP terms. Remarkably, all these pathways demonstrate up-regulation (Figure 1, Figure 2C, Supplementary Table 1).

Within the intersection encompassing genes regulated in both the iRBC true vs. false and febrile true vs. false analyses, a cohort of 80 genes is observed. Among this group, 3 genes are associated with the "vascular endothelial growth factor signaling pathway" GO-BP term (FDR 0.00025), while 4 genes are linked to the "response to unfolded protein" term (FDR 2.50e-05), all demonstrating up-regulation (Figure 1, Figure 2D, Supplementary Table 1).

In the intersection of all three comparative analyses (niRBC true vs. false; febrile true vs. false; iRBC true vs. false), a set of 80 regulated genes is identified. Notably, the pathways significantly up-regulated include the "chemokine-mediated signaling pathway" (FDR 2.26e-12), comprising 13 genes, and the "protein folding" GO-BP term (FDR 3.29e-10), comprising 7 genes (Figure 1, Figure 2E, Supplementary Table 1).

Focusing solely on the "iRBCs true vs. false" analysis, comprising a total of 762 genes, a comprehensive string analysis reveals 7 distinct clusters housing 6 to 24 of the regulated genes. These clusters correspond to the GO-BP terms "response to cytokine" (FDR 2.01e-23; 24 genes), "response to unfolded protein" (FDR 1.67e-30; 17 genes), "positive regulation of RNA metabolic process" (FDR 7.04e-10; 15 genes), "nucleosome organization" (FDR 3.83e-11; 9 genes), "defense response to virus" (FDR 7.04e-16; 12 genes), and "I-κB kinase/NF-κB signaling" (FDR 3.24e-08; 6 genes). Remarkably, all these clusters exhibit up-regulation. Conversely, the sole down-regulated cluster pertains to "nucleosome assembly" (FDR 3.64e-08; 7 genes) (Figure 1, Figure 3A, Supplementary Table 1).

Using ShinyGO analysis, the proteins encoded by the identified genes were categorized according to their biological functions. Among the 10 most significantly regulated functions are the TNF, IL-17 and NF-κB signaling pathways and cytokine-cytokine receptor interaction, as well as the response to viral (herpes simplex virus 1 infection) and bacterial (legionellosis) infections (Figure 3B).

3.2 Specific response of HBEC-5i cells to cytoadhesion of iRBCs$^{37°C}$ and iRBCs$^{40°C}$

It is important to emphasize that in the above comparisons, although the expression of genes within different intersections is altered compared to the respective controls, the magnitude of these changes in expression levels is not considered in the analyses. Hence, a more detailed investigation is warranted to delineate the specific impact of iRBC cytoadhesion on HBEC-5i cells and the role of the cultivation temperature (37°C vs. 40°C). In contrast to the previous analysis, this subsequent investigation only includes the mean values of normalized reads from three biologically independent transcriptomes. However, the focus remains exclusively on genes identified as significantly regulated within the analysis above (Supplementary Table 1).

In the first step, we identified those genes whose expression was up- or down-regulated (fold change $\geq$ 2.0, FDR $\leq$ 0.05, TPM threshold $\geq$ 2.0) in HBEC-5i cells in response to iRBCs compared to the presence of niRBCs at 37°C (iRBCs$^{37°C}$ vs niRBCs$^{37°C}$). Cytoadhesion of iRBCs$^{37°C}$ up-regulates the expression of 162 genes and down-regulates the expression of 219 genes compared to the presence of niRBCs$^{37°C}$ (Supplementary Table 1).

To get an idea of which metabolic pathways the proteins encoded by the 162 up-regulated genes are involved, a g:Profiler analysis was performed. The GO Molecular Function (GO-MF) terms "structural component of chromatin" and "chemokine activity" had the highest significance. For the GO-BP terms, these were "cellular response to organic substance" and "response to cytokine". The KEGG pathway analysis showed the highest significance for "TNF signaling pathway" and in the Reactome analysis these were "interleukin-10 signaling" and "cytokine signaling immune system" (Figure 4A). This is also reflected in the ShinyGO analysis. Here, as described in the previous paragraph, different signaling pathways of the immune system and the response to viral and bacterial infections were mainly regulated ("TNF, IL-17 and NF-κB signaling pathways", "measles", "legionellosis") (Figure 4B). The use of a string analysis yielded very similar results. Here, among the 162 up-regulated genes, the KEGG pathways exhibiting the most stringent regulation include the "TNF signaling pathway" (FDR 6.70e-09), "viral protein interaction with cytokine and cytokine receptor" (FDR 1.03e-07), "IL-17 signaling pathway" (FDR 4.42e-07), "cytokine-cytokine receptor interaction" (FDR 7.64e-07), and "NF-κB signaling pathway" (FDR 8.31e-07). In total, 8 genes are up-regulated more than 10-fold, while another 16 genes are up-regulated more than 5-fold. Notable among these are the chemokines CCL31, CCL3, CCL20, and IL-6 (Supplementary Table 1).

Furthermore, the up-regulated genes can be classified in the clusters of the following GO-BP terms: "response to cytokine" (FDR 5.50e-18, 19 genes), "nucleosome assembly" (FDR 1.99e-18, 12 genes), "defense response to virus" (FDR 6.28e-13, 10 genes), "cellular response to hormone stimulus" (FDR 0.00030, 5 genes). (Figure 5A, Supplementary Table 1). The 219 genes whose expression is down-regulated code for proteins that fall under the GO-MF term "DNA-binding transcription factor activity" (FDR 2.008×10^{-20}) and the GO-BP term "regulation of nucleobase-containing compound metabolic process" (FDR 3.664×10^{-18}) according to the g:Profiler analysis (Supplementary Table 1).

The next question was to investgate the effect of increasing the temperature from 37°C to 40°C (iRBCs$^{40°C}$), simulating fever as an additional factor besides cytoadhesion of iRBCs$^{37°C}$. Therefore, the expression profile of HBEC-5i cells following cytoadhesion iRBCs$^{40°C}$ was compared to HBEC-5i cells cultured in the presence of niRBCs$^{37°C}$. This analysis revealed 381 genes up-regulated and 213 genes down-regulated in expression (fold change $\geq$ 2.0, FDR $\leq$ 0.05, TPM threshold $\geq$ 2.0) (Supplementary Table 1). G:Profiler and ShinyGO analyses group of the 381 genes up-regulated at 40°C were also carried out here. By and large, the same pathways were identified. However, for many of the pathways, the significance was more than doubled, e.g. for the TNF signaling pathway (Figures 4C, 4D). A string analysis of this comparison also shows great agreement with the analyses mentioned above. The same KEGG pathways are regulated at 40°C, that were already addressed at 37°C but with one exception with higher significance ("TNF signaling pathway" (FDR 1. 54e-17), "IL-17 signaling pathway" (FDR 2.06e-12), "cytokine-cytokine receptor interaction" (8.20e-11) and "NF-κB signaling pathway" (FDR 8.00e-11), "viral protein interaction with cytokine and cytokine receptor" (FDR 6.70e-07)). GO-BP terms identified as significant in the iRBC$^{37°C}$ with niRBCs$^{37°C}$ comparison, such as "defence reaction to viruses" and "cellular reaction to hormone stimuli", are no longer found in the iRBC$^{40°C}$ with niRBCs$^{37°C}$ comparison. Instead of 12 genes, the GO term "nucleosome assembly" (FDR 8.01e-12) still shows 9 genes as significantly up-regulated in their expression. The significance for the GO-BP term "response to unfolded protein" increases from FDR 1.09e-09 (6 genes) to FDR 1.67e-30 (17 genes). This is also true for the GO-BP term "cytokine response" (FDR 3.62e-24), to which 23 genes can be assigned. Furthermore, the GO-BP term "I-κB kinase/NF-κB signaling" (FDR 2.36e-06, 5 genes) was identified as regulated (Figure 4B, Supplementary Table 1). In addition, significant up-

regulation of gene expression was detected, with a notable subset showing ≥10-fold increase (32 genes including *CCL26, CCL20, CCL3L1, CXCL3, CXCL8* and *Il6*) and a further 55 genes showing 5- to <10-fold increase when comparing niRBC$^{37°C}$ vs. iRBC$^{37°C}$ (Supplementary Table 1).

The 213 genes whose expression is down-regulated encode for proteins that fall under the GO-MF term "DNA-binding transcription factor activity" (FDR $1.805×10^{-12}$) and the GO-BP term "regulation of transcription by RNA polymerase II" (FDR $4.185×10^{-9}$) according to the g:Profiler analysis (Supplementary Table 1).

The next stage was to examine the genes whose expression was regulated in the presence of iRBCs at 40°C compared to those at 37°C (fold change ≥ 2.0, FDR ≤ 0.05, TPM threshold ≥ 2.0). 205 genes (18 genes fold change >10-, 25 genes fold change 5-10) were identified for which cytoadhesion at 40°C leads to a further increase in expression compared to cytoadhesion at 37°C. Down-regulation was observed for 38 genes (Supplementary Table 1). For some genes, a continuous increase in expression can be observed, starting from cultivation in the presence of niRBCs$^{37°C}$ via iRBCs$^{37°C}$ to iRBCs$^{40°C}$. For instance, this trend is evident in the case of CXCL8, where the expression rises from 11.7 TPM (no niRBCs) to 82 (niRBCs$^{37°C}$), further to 392.6 (iRBCs$^{37°C}$), and peaks at 1932 (iRBCs$^{40°C}$) (Figure 6, Supplementary Table 1). For other genes, such as HSPA1A, the expression with and without niRBCs is comparable, and an increase in expression is seen in the presence of iRBCs at 37°C and increases further at 40°C (TPM: 177 (without niRBCs); 160 (niRBCs); 537 (iRBCs$^{37°C}$), 23001 (iRBCs$^{40°C}$) (Supplementary Table 1).

The g:Profiler and ShinyGO analyses of the 205 genes whose expression was upregulated in the presence of iRBCs$^{40°C}$ compared to iRBCs$^{37°C}$, which were also carried out here, show that immunological signaling pathways are again positively influenced by the combination of iRBCs and elevated temperature (Figure 4F, G). Analysis of the 38 down-regulated genes reveals no significantly regulated pathways.

The genes whose expression is further increased in a temperature-dependent manner can be assigned to the KEGG pathways "TNF signaling pathway" (FDR 2.96e-12), "IL-17 signaling pathway" (FDR 2.76e-11), "NF-κB signaling pathway (FDR 1.09e-09) and "cytokine-cytokine receptor interaction" (FDR 2.98e-09). In particular for genes belonging to the GO-BP terms "response to cytokine" (17 genes, FDR 5.96e-19) and "protein folding" (13 genes, FDR 6.55e-23), an increase in expression can be observed with an increase in temperature from 37°C to 40°C (Figure 5C; Supplementary Table 1).

3.3 Influence of iRBCs on the expression of genes encoding proteins involved in the immune response, nucleosome assembly, angiogenesis, and interferon-alpha/beta signaling pathway

The comparative analyses described above show that cytoadhesion of iRBCs leads to an increase in the expression of genes whose encoded proteins are involved in the host's immune response. For some of these genes, the higher temperature leads to a further increase in expression, although temperature alone has little or no effect on the expression level of these genes. The encoded cytokines can be divided into four groups. 1) Genes that show an increase in expression when co-cultured with iRBCs$^{37°C}$, but maintain a similar level of expression when the temperature is raised to 40°C (*CXCL10, CXCL11, CCL5, CCL20, IL12A*); 2) Genes whose expression is up-regulated at 37°C and whose expression is further up-regulated when the temperature is raised to 40°C (*CXCL2, CXCL3, CXCL6, CXCL8, CCL3, CCL3L1, CCL26, IL1a, IL6*); 3) Genes whose expression increases only during co-cultivation at 40°C (*CXCL1, CXCL5, IL1b, IL32*); 4) Genes that show similar levels of expression under all conditions (*CXCL12, CXCL14, CXCL16, CCL2*) (Figure 6A, Supplementary Table 1).

Six genes whose encoded proteins can assume functions in angiogenesis are also differentially expressed. These include three genes (*MMP1, MMP2, MMP3*) encoding metalloproteinases as well as the genes (*NR4A1, NR4A2, NR4A3*) encoding the three members of the NR4A nuclear hormone receptor superfamily. For MMPs, the increased expression was only observed at elevated temperature in the presence of iRBCs, whereas for NR4A1, NR4A2, NR4A3, the expression increased in the

presence of iRBCs and a further increase was observed after raising the temperature from 37°C to 40°C
(Figure 6B, Supplementary Table 1).
In addition, twelve genes coding for histones or histone linker proteins were significantly up-regulated
in the presence of iRBCs. However, raising the temperature to 40°C did not lead to a further increase
in expression, but on the contrary, the expression level for four genes has the expression level
significantly decreased in the presence of niRBCs (Figure 6C, Supplementary Table 1).
Interestingly, additional 14 genes were identified coding for proteins of the antiviral immune response
or the interferon-alpha/beta signaling pathway (Reactome: FDR 2.22E-15) (Figures 6D, 6E,
Supplementary Table 1). The presence of iRBCs alone leads to a significant up-regulation in
expression. With the exception of *IL6*, there is no further increase in expression at elevated
temperatures. On the contrary, for four genes (*IFI6, ISG15, MX1, MX2*) the expression level falls back
to the baseline level of niRBCs (Figures 6D, 6E, Supplementary Table 1).
392

4 Discussion

P. falciparum malaria elicits a robust host immune response that contributes significantly to the
pathogenesis of the disease through increased systemic and local inflammation. Various studies have
shown that the cytoadhesion of iRBCs strongly influences the gene expression of the ECs. Among
other things, it has been shown that binding to HBEMCs mainly influences the signaling pathways
involved in the inflammatory response, apoptosis, cell-cell signaling, signal transduction, and the NF-
kB activation cascade (Tripathi et al., 2009; Othman et al., 2023). However, not only does cytoadhesion
seem to influence the expression, but it also seems to depend on which *Pf*EMP1 is presented on the
surface of the iRBCs (Othman et al., 2023). Similar effects have also been demonstrated for HUVECs.
However, some studies show that the stimulation of HUVECs by cytoadhesion of iRBCs is only
possible in the presence of low levels of TNF (Viebig et al., 2005; Tripathi et al., 2006; Chakravorty
et al., 2007).
406
In the present study, we analysed the influence of iRBC cytoadhesion on HBEC-5i cells. For this
purpose, an iRBC population enriched for binding to HBEC-5i cells was used. This resulted in a
homogeneous cell population presenting IT4var04 (var2CSA) on their surface, as CSA is the dominant
antigen on HBEC-5i cells (Dorpinghaus et al., 2020). This model was employed to investigate whether
stimulation of HBEC-5i cells by binding of iRBCs to CSA is possible and how cytoadhesion affects
the expression profile also at elevated temperatures, simulating fever.
413
First, the influence of the presence of niRBCs (niRBC true vs. false), iRBCs (iRBC true vs. false), and
fever (febrile true vs. false) on the expression profile of HBEC-5i cells was analysed. For these true vs.
false analyses, only one stimulus is considered. For example, in the case of iRBC true vs. false, all
HBEC-5i transcriptomes in which iRBCs were co-incubated with HBEC-5i cells were analysed,
regardless of the temperature used, and in the case of febrile true vs. false, all transcriptomes in which
HBEC-5i cells were cultured at 40°C were analysed, regardless of the presence or absence of niRBCs
or iRBCs. In this way, all three comparisons were analysed in a similar way.
The analysis showed that the presence of niRBCs alone has a strong influence on the expression profile
of HBEC-5i cells. 59 genes are differentially expressed only in the presence of niRBCs. Since the
presence of niRBCs is an organic component of EC homeostasis, the overlap of the two DEG analyses
between niRBC (true vs. false) and iRBC (true vs. false) allows the isolation of genes that are affected
by the presence of the parasite compared to those of niRBCs. However, there is a large overlap of 364
genes that are also differentially expressed in the presence of iRBCs. It is important to note that this

analysis does not provide any information about the magnitude of expression level changes.
Particularly interesting are, for example those genes showing up-regulated expression in the presence
of niRBCs and further up-regulation in the presence of iRBCs. Some examples of genes showing this
behavior are *JUNB* which encodes a subunit of the AP-1 transcription factor (TPM: 33_ no RBCs;
72_niRBCs; 227_iRBCs), *IL6* (TPM: 3_no RBCs; 13_niRBCs; 70_iRBCs), and *CXCL8* (TPM: 11_no
RBCs; 82_niRBCs; 392_iRBCs). As expected, increasing the temperature from 37°C to 40°C also has
a significant effect on the expression profile of HBEC-5i cells, mainly affecting genes encoding heat
shock proteins involved in protein folding and stress response.
BOLA2 is the most up-regulated gene in the comparison of niRBCs true vs. false, increasing from TPM
2.5 to 198 (almost 80-fold). So far, little is known concerning the function of BOLA2. However, it was
described that it could interact with glutaredoxin-3 and act as a redox-regulated regulator of Fe-S
cluster biogenesis or as a transcriptional regulator (Banci et al., 2015).
439
Looking at the iRBC true vs. false analysis, a total of 762 genes are differentially regulated. In
particular, many genes involved in immunological signaling pathways, such as the TNF signaling
pathway, the IL-17 signaling pathway, the NF-kB signaling pathway, and the cytokine-cytokine
receptor signaling pathway are up-regulated. Interestingly, also genes coding for proteins involved in
the defense response to viruses and in nucleosome organization or assembly are expressed at higher
levels.
A direct comparison of the influence of iRBCs compared to niRBCs at 37°C and 40°C shows that the
positively regulated pathways are by and large the same as found in the iRBC true vs. false analysis.
Some of these signaling pathways have already been identified as regulated in the study by Tripathi
and colleagues (Tripathi et al., 2009). In addition to the most strongly affected signaling pathways
"cytokine-cytokine receptor interaction" and "cell surface receptor-linked signal transduction", many
genes of the NFkB activation cascade were also found to be positively regulated (Tripathi et al., 2009).
On the other hand, some of the signaling pathways identified as regulated in the present study were not
identified as regulated in Tripathi *et al.*, including nucleosome assembly and the defense response to
viruses (Tripathi et al., 2009).
It is known that the complex immune response elicited by *P. falciparum* infection involves the release
of numerous cytokines, of which more than 30 have been detected at elevated levels in the serum or
plasma of malaria patients compared to healthy individuals. These cytokines include CXCL2, CXCL4,
CXCL8, CXCL9, CXCL10, CXCL11, CCL2, CCL3, CCL4, CCL5, CCL20, CCL28, IL-1b, IL-6, IL-
9, IL-10, IL-12, IL-13, IL-15, IL-17A, IL-31, IL-33, IL1-RA, TNF-a, interferon-gamma (INFg),
VEGF, and G-CSF (Lyke et al., 2004; Prakash et al., 2006; Armah et al., 2007; Jain et al., 2008;
Ayimba et al., 2011; Wilson et al., 2011; Che et al., 2015; Colborn et al., 2015; Herbert et al., 2015;
Bujarbaruah et al., 2017). Certain cytokines (such as CXCL4, CXCL8, CXCL10, CCL2, CCL3, and
CCL4) show a direct correlation with the severity of malaria infection (Armah et al., 2007; Ayimba et
al., 2011; Wilson et al., 2011) and it has been shown that patients with cerebral malaria have
significantly increased levels of CCL2, CCL4, CXCL4, CXCL8, CXCL10, IL-1RA, IL-6, TNF and G-
CSF (Jain et al., 2008; John et al., 2008a; John et al., 2008b; Wilson et al., 2011). Our study, as well
as previous studies, show that cytoadhesion of iRBCs increases the expression of many genes encoding
cytokines (for review (Dunst et al., 2017)) even at 37°C. For nine of these 14 genes, in addition to
cytoadhesion, raising the temperature from 37°C to 40°C (simulation of fever) leads to a further
increase in expression of > 2-fold. The latter group includes genes encoding the inflammatory
chemokines CXCL2, CXCL3, CXCL6, and CXCL8, which attract neutrophils, CCL26, which attracts
eosinophil granulocytes, and CCL3 and CCL3L1, which attract macrophages and other immune cells.
For genes encoding the chemokines CXCL10, CXCL11, CCL5, and CCL20, there is a significant
increase in expression due to cytoadhesion, but this is not further increased by the rise in temperature.
These chemokines attract various immune cells, including T-cells, monocytes, and dendritic cells (for

review (Hughes and Nibbs, 2018). *CXCL1* and *CXCL5* are the only chemokines encoding genes whose
expression only increases by the temperature shift to 40°C in the presence of iRBCs. These chemokines
play a role in the attraction of neutrophil granulocytes. Recently, stimulation of ECs by plasma from
malaria patients was shown to significantly increase the expression of *CXCL1* and *CXCL5* compared
to healthy controls (Raacke et al., 2021). Interestingly, the increase in expression observed here for
CXCL6 and *CCL26* has not yet been described to our knowledge. *CXCL6* (granulocyte chemotactic
protein-2 (GCP-2)) is one of the genes whose expression increases in the presence of iRBCs and is
further increased when the temperature is raised to 40°C. As described above, CXCL6 attracts
neutrophils and is a ligand of the chemokine receptors CXCR1 and CXCR2 (Fan et al., 2007). Since
neutrophils are important for the development of cerebral malaria, the expression of the gene was
analysed in the malaria mouse model *Plasmodium berghei* ANKA (Chen et al., 2000). Still, infection
of both C57BL/6 and Balb/c mice did not lead to an increase in expression in the brains of mice
compared to non-infected mice. However, this was consistent with the observation that no neutrophils
accumulated in the brains of mice with cerebral malaria (Van den Steen et al., 2008). To our knowledge,
there is only one study that has compared the plasma concentration of CCL26 in malaria patients with
that of non-infected controls. It showed that the plasma concentrations of CCL26 in the malaria-
exposed group were significantly lower than in the unexposed group. This observation correlates very
well with our results. The TMP increases only slightly from 1 to 3 in the presence of iRBCs, only by
increasing the temperature it does increase to TPM 26 (Mancebo-Perez et al., 2022). It can therefore
be postulated that CCL26 is only important during the fever attacks of malaria.
The expression of genes encoding various interleukins is also up-regulated by the presence of iRBCs.
These include the pro-inflammatory cytokines IL-1β, IL-1α, IL-6, IL-12A, and IL-32, whereby for *IL-*
1a and IL6 a more than 2-fold increase in expression is seen from co-incubation of niRBCs to
iRBC37°C to iRBC40°C, while for *IL1β* and *IL32* an increase is only observed after co-incubation
with iRBCs$^{40°C}$. A recent meta-analysis showed that serum IL1β levels are higher in patients with
severe malaria than in patients with uncomplicated malaria. However, the level in patients with
uncomplicated malaria was very similar to that of healthy controls (Mahittikorn et al., 2022). This
observation fits very well with the results presented here. Only when the temperature increases from
37°C to 40°C, does the expression of *IL1β* increases by about 2.5 times. IL-32 shows the same
expression profile as IL1β, i.e. an increase in expression only occurs in the presence of iRBCs$^{40°C}$. IL-
32 is expressed in different isoforms in immune and non-immune cells, including ECs, and its functions
include the induction of the production of several pro-inflammatory cytokines (TNF-α, IL-1β, IL-6,
and CXCL8). The function of IL-32 in *P. falciparum* infection is not yet known. However, IL-32 has
been described as a master regulator of intracellular bacterial, viral, and eukaryotic infections (for
review (Dos Santos et al., 2018). In this context, it has been shown that in *Mycobacterium tuberculosis*
(Mtb) infection, endogenous IL-32 induces the production of TNFα, IL-1β, and CXCL8, as well as the
production of antimicrobial peptides and apoptosis, so that it can be assumed that IL-32 is associated
with protection against Mtb infections (Bai et al., 2010; Montoya et al., 2014; Bai et al., 2015). As
shown for MtB infection, the production of various cytokines such as TNFα, CXCL8, IL-1R, IFNγ,
and IL-10 as well as antimicrobial peptides were induced in *Leishmania* infection (Dos Santos et al.,
2017; Dos Santos et al., 2018).
517
Our results shown here are very much in line with the findings of Tripathi and colleagues, who
investigated the response of HBMECs to the presence of iRBCs. Significantly increased expression
was detected for CXCL1, CXCL2, CXCL3, CXCL6, CXCL8, CCL20, and IL-6 (Tripathi et al., 2009).
However, we were able to identify eleven further cytokines whose expression is up-regulated in the
presence of iRBCs$^{37°C}$ and/or iRBCs$^{40°C}$. Furthermore, a number of genes encoding proteins involved
in the regulation of inflammation and immune response are also increased in their expression by the
cytoadhesion iRBCs compared to the control. These proteins include among others PTGS2 (COX-2)

(Martin-Vazquez et al., 2023) and TXNIP (thioredoxin interacting protein) (Mohamed et al., 2021).
Other genes that encode proteins involved in inflammation and immune reactions were expressed more
strongly at elevated temperatures. The corresponding encoded proteins include, among others,
TNFAIP3 (A20) (Wu et al., 2020), LGALS4 (galectin-4) (Cao and Guo, 2016), ATF3 (activating
transcription factor 3) (Hai et al., 2010), CD74 (cluster of differentiation 74) (Beswick and Reyes,
2009), CH25H (cholesterol 25-hydroxylase) (Cho et al., 2023), and CD274 (Huang et al., 2013).
The three members of the NR4A nuclear hormone receptor superfamily (NR4A1 (Nur77), NR4A2
(Nur-related factor-1), NR4A3 (Neuron-Derived Orphan Receptor 1)) are also increasingly expressed
in the presence of iRBCs, and a further increase was observed after elevating the temperature. In
inflammatory diseases, NR4A family members are rapidly expressed and control the differentiation
and activity of innate and adaptive immune cells. In addition, these proteins mediate cytokine signaling
to control angiogenesis (for review (Hamers et al., 2013; Murphy and Crean, 2022)). Furthermore,
three genes encoding members of matrix metalloproteinases (MMP1, MMP2, MMP3) are specifically
up-regulated only in the presence of iRBCs at elevated temperatures. Among other things, MMPs are
involved in the breakdown of extracellular matrix proteins and regulate functions related to
inflammation, such as the activity of inflammatory cytokines and chemokines (for review (Lee and
Kim, 2022)). MMP1 cleaves collagen type I and is involved in the modulation and remodeling of the
extracellular matrix. MMP2 cleaves collagen type IV and is involved in angiogenesis by degrading the
basement membrane of blood vessels and thus promoting the migration of endothelial cells. MMP3
can degrade various types of collagens as well as proteoglycans and fibronectin and is involved in
tissue remodeling and inflammatory reactions, among other things (for review (Lee and Kim, 2022).
Therefore, it can be speculated that the increase in the expression of MMP and NR4A encoding genes
could be a reason for the endothelial dysfunction observed in malaria.

Strikingly, 14 genes coding for proteins involved in the antiviral immune response or interferon-
alpha/beta signaling pathway are upregulated by cytoadhesive iRBCs. However, with the exception of
IL6, there is no further increase in the expression at elevated temperatures. On the contrary, for four
genes (*IFI6*, *ISG15*, *MX1*, and *MX2*) the expression level drops back to that in the presence of niRBCs.
In summary, the genes up-regulated in their expression encode the pro-inflammatory cytokine IL-6,
which can be induced by the presence of viruses, MX1 (MX dynamin-like GTPase 1), MX2 (MX
dynamin-like GTPase 2), IFI6 (interferon alpha inducible protein 6) and ISG15 (interferon-stimulated
gene 15), which inhibit viral replication, OAS2 (2'-5'-oligoadenylate synthetase 2) and OAS3 (2'-5'-
oligoadenylate synthetase 3), which synthesize secondary messengers that serve to activate RNase L,
which degrades viral and cellular RNA, and ISG20 (interferon stimulated exonuclease gene 20 kDa),
which is an endoribonuclease that supports the degradation of viral RNA (Tanaka et al., 2014), (Sun et
al., 2012; Haller et al., 2015; Zhu et al., 2015; Sajid et al., 2021; Deymier et al., 2022). In addition, the
genes *BST2* (bone marrow stroma cell antigen 2), IFI27 (interferon alpha-inducible protein 27), *DDIT4*
(DNA-damage-inducible transcript 4), *GBP4* (guanylate binding protein 4), *DDX60* (DEAD-box
helicase 60), which are also involved in the antiviral immune response, are up-regulated (Miyashita et
al., 2011; Tretina et al., 2019; Busse et al., 2020; Ullah et al., 2021; Zhao et al., 2022). It was shown
that a type I interferon (IFN) response occurs during *P. berghei* replication in the liver. A *P. berghei*
RNA was identified as a pathogen-associated molecular pattern (PAMP) that can activate a type I IFN
response. Thus, liver cells have sensor mechanisms that mediate a type I IFN-driven anti-parasite
response (Liehl et al., 2014). However, in the case of iRBCs, it is unclear how such an response is
induced in HBEC-5i cells and whether cytoadhesion is the trigger for this response or whether other
factors, such as extracellular vesicles secreted by iRBCs and containing various microRNAs, might be
involved (Wu et al., 2023).

In conclusion, the presence of iRBCs, mediated by cytoadhesion or other mechanisms such as cell-cell communication via extracellular vesicles, stimulates brain endothelial cells. In particular, as described in other studies, several immunological pathways are activated, and the stimulation of antiviral (-parasitic) mechanisms and angiogenesis was also identified in the present study. In particular, the expression of genes encoding cytokines, MMPs, and NR4As is further increased under fever conditions.

Data availability statement

Raw data available as expression browser in Supplementary Table 1

Author contributions

JA: Conceptualization, Formal analysis, Investigation, Methodology, Visualization, Writing – original draft, Writing – review & editing. MB: Conceptualization, Formal analysis, Investigation, Methodology, Writing – review & editing. HT: Writing – review & editing. MPMT: Writing – review & editing. NGM: Methodology, Writing – review & editing. TR: Writing – review & editing. TG: Funding acquisition, Writing – review & editing. IB: Conceptualization, Data curation, Formal analysis, Funding acquisition, Project administration, Supervision, Visualization, Writing – original draft, Writing – review & editing.

Funding

Acknowledgments

The authors would like to thank Susann Ofori for excellent technical assistance.

Conflict of Interest

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

References

Alampalli, S.V., Grover, M., Chandran, S., Tatu, U., and Acharya, P. (2018). Proteome and Structural Organization of the Knob Complex on the Surface of the Plasmodium Infected Red Blood Cell. *Proteomics Clin Appl* 12(4)**,** e1600177. doi: 10.1002/prca.201600177.
Armah, H.B., Wilson, N.O., Sarfo, B.Y., Powell, M.D., Bond, V.C., Anderson, W., et al. (2007). Cerebrospinal fluid and serum biomarkers of cerebral malaria mortality in Ghanaian children. *Malar J* 6**,** 147. doi: 10.1186/1475-2875-6-147.

Ayimba, E., Hegewald, J., Segbena, A.Y., Gantin, R.G., Lechner, C.J., Agosssou, A., et al. (2011).
Proinflammatory and regulatory cytokines and chemokines in infants with uncomplicated and
severe Plasmodium falciparum malaria. *Clin Exp Immunol* 166(2)**,** 218-226. doi:
10.1111/j.1365-2249.2011.04474.x.
Bai, X., Kim, S.H., Azam, T., McGibney, M.T., Huang, H., Dinarello, C.A., et al. (2010). IL-32 is a
host protective cytokine against Mycobacterium tuberculosis in differentiated THP-1 human
macrophages. *J Immunol* 184(7)**,** 3830-3840. doi: 10.4049/jimmunol.0901913.
Bai, X., Shang, S., Henao-Tamayo, M., Basaraba, R.J., Ovrutsky, A.R., Matsuda, J.L., et al. (2015).
Human IL-32 expression protects mice against a hypervirulent strain of Mycobacterium
tuberculosis. *Proc Natl Acad Sci U S A* 112(16)**,** 5111-5116. doi: 10.1073/pnas.1424302112.
Banci, L., Camponeschi, F., Ciofi-Baffoni, S., and Muzzioli, R. (2015). Elucidating the Molecular
Function of Human BOLA2 in GRX3-Dependent Anamorsin Maturation Pathway. *J Am Chem*
Soc 137(51)**,** 16133-16143. doi: 10.1021/jacs.5b10592.
Beswick, E.J., and Reyes, V.E. (2009). CD74 in antigen presentation, inflammation, and cancers of the
gastrointestinal tract. *World J Gastroenterol* 15(23)**,** 2855-2861. doi: 10.3748/wjg.15.2855.
Bujarbaruah, D., Kalita, M.P., Baruah, V., Basumatary, T.K., Hazarika, S., Begum, R.H., et al. (2017).
RANTES levels as a determinant of falciparum malaria severity or recovery. *Parasite Immunol*
39(9). doi: 10.1111/pim.12452.
Busse, D.C., Habgood-Coote, D., Clare, S., Brandt, C., Bassano, I., Kaforou, M., et al. (2020).
Interferon-Induced Protein 44 and Interferon-Induced Protein 44-Like Restrict Replication of
Respiratory Syncytial Virus. *J Virol* 94(18). doi: 10.1128/JVI.00297-20.
Cao, Z.Q., and Guo, X.L. (2016). The role of galectin-4 in physiology and diseases. *Protein Cell* 7(5)**,**
314-324. doi: 10.1007/s13238-016-0262-9.
Chakravorty, S.J., Carret, C., Nash, G.B., Ivens, A., Szestak, T., and Craig, A.G. (2007). Altered
phenotype and gene transcription in endothelial cells, induced by Plasmodium falciparum-
infected red blood cells: pathogenic or protective? *Int J Parasitol* 37(8-9)**,** 975-987. doi:
10.1016/j.ijpara.2007.02.006.
Che, J.N., Nmorsi, O.P., Nkot, B.P., Isaac, C., and Okonkwo, B.C. (2015). Chemokines responses to
Plasmodium falciparum malaria and co-infections among rural Cameroonians. *Parasitol Int*
64(2)**,** 139-144. doi: 10.1016/j.parint.2014.11.003.
Chen, L., Zhang, Z., and Sendo, F. (2000). Neutrophils play a critical role in the pathogenesis of
experimental cerebral malaria. *Clin Exp Immunol* 120(1)**,** 125-133. doi: 10.1046/j.1365-
2249.2000.01196.x.
Cho, S.J., Pronko, A., Yang, J., Pagan, K., and Stout-Delgado, H. (2023). Role of Cholesterol 25-
Hydroxylase (Ch25h) in Mediating Innate Immune Responses to Streptococcus pneumoniae
Infection. *Cells* 12(4). doi: 10.3390/cells12040570.
Colborn, J.M., Ylostalo, J.H., Koita, O.A., Cisse, O.H., and Krogstad, D.J. (2015). Human Gene
Expression in Uncomplicated Plasmodium falciparum Malaria. *J Immunol Res* 2015**,** 162639.
doi: 10.1155/2015/162639.
Crabb, B.S., Cooke, B.M., Reeder, J.C., Waller, R.F., Caruana, S.R., Davern, K.M., et al. (1997).
Targeted gene disruption shows that knobs enable malaria-infected red cells to cytoadhere
under physiological shear stress. *Cell* 89(2)**,** 287-296.
Cunnington, A.J., Riley, E.M., and Walther, M. (2013). Microvascular dysfunction in severe
Plasmodium falciparum Malaria. *J Infec Dis* 207(2)**,** 369-370. doi: 10.1093/infdis/jis681.
Cutts, E.E., Laasch, N., Reiter, D.M., Trenker, R., Slater, L.M., Stansfeld, P.J., et al. (2017). Structural
analysis of P. falciparum KAHRP and PfEMP1 complexes with host erythrocyte spectrin
suggests a model for cytoadherent knob protrusions. *PLoS Pathog* 13(8)**,** e1006552. doi:
10.1371/journal.ppat.1006552.

Deitsch, K.W., and Dzikowski, R. (2017). Variant Gene Expression and Antigenic Variation by
Malaria Parasites. *Annu Rev Microbiol* 71**,** 625-641. doi: 10.1146/annurev-micro-090816-
093841.
Deymier, S., Louvat, C., Fiorini, F., and Cimarelli, A. (2022). ISG20: an enigmatic antiviral RNase
targeting multiple viruses. *FEBS Open Bio* 12(6)**,** 1096-1111. doi: 10.1002/2211-5463.13382.
Dorpinghaus, M., Furstenwerth, F., Roth, L.K., Bouws, P., Rakotonirinalalao, M., Jordan, V., et al.
(2020). Stringent Selection of Knobby Plasmodium falciparum-Infected Erythrocytes during
Cytoadhesion at Febrile Temperature. *Microorganisms* 8(2). doi:
10.3390/microorganisms8020174.
Dos Santos, J.C., Damen, M., Joosten, L.A.B., and Ribeiro-Dias, F. (2018). Interleukin-32: An
endogenous danger signal or master regulator of intracellular pathogen infections-Focus on
leishmaniases. *Semin Immunol* 38**,** 15-23. doi: 10.1016/j.smim.2018.02.010.
Dos Santos, J.C., Heinhuis, B., Gomes, R.S., Damen, M.S., Real, F., Mortara, R.A., et al. (2017).
Cytokines and microbicidal molecules regulated by IL-32 in THP-1-derived human
macrophages infected with New World Leishmania species. *PLoS Negl Trop Dis* 11(2)**,**
e0005413. doi: 10.1371/journal.pntd.0005413.
Dunst, J., Kamena, F., and Matuschewski, K. (2017). Cytokines and Chemokines in Cerebral Malaria
Pathogenesis. *Front Cell Infect Microbiol* 7**,** 324. doi: 10.3389/fcimb.2017.00324.
Fabregat, A., Sidiropoulos, K., Viteri, G., Marin-Garcia, P., Ping, P., Stein, L., et al. (2018). Reactome
diagram viewer: data structures and strategies to boost performance. *Bioinformatics* 34(7)**,**
1208-1214. doi: 10.1093/bioinformatics/btx752.
Fan, X., Patera, A.C., Pong-Kennedy, A., Deno, G., Gonsiorek, W., Manfra, D.J., et al. (2007). Murine
CXCR1 is a functional receptor for GCP-2/CXCL6 and interleukin-8/CXCL8. *J Biol Chem*
282(16)**,** 11658-11666. doi: 10.1074/jbc.M607705200.
Gardner, M.J., Shallom, S.J., Carlton, J.M., Salzberg, S.L., Nene, V., Shoaibi, A., et al. (2002).
Sequence of Plasmodium falciparum chromosomes 2, 10, 11 and 14. *Nature* 419(6906)**,** 531-
534. doi: 10.1038/nature01094.
Gazzinelli, R.T., Kalantari, P., Fitzgerald, K.A., and Golenbock, D.T. (2014). Innate sensing of malaria
parasites. *Nat Rev Immunol* 14(11)**,** 744-757. doi: 10.1038/nri3742.
Gillespie, M., Jassal, B., Stephan, R., Milacic, M., Rothfels, K., Senff-Ribeiro, A., et al. (2022). The
reactome pathway knowledgebase 2022. *Nucleic Acids Res* 50(D1)**,** D687-D692. doi:
10.1093/nar/gkab1028.
Griss, J., Viteri, G., Sidiropoulos, K., Nguyen, V., Fabregat, A., and Hermjakob, H. (2020).
ReactomeGSA - Efficient Multi-Omics Comparative Pathway Analysis. *Mol Cell Proteomics*
19(12)**,** 2115-2125. doi: 10.1074/mcp.TIR120.002155.
Gruenberg, J., Allred, D.R., and Sherman, I.W. (1983). Scanning electron microscope-analysis of the
protrusions (knobs) present on the surface of Plasmodium falciparum-infected erythrocytes. *J*
Cell Biol 97(3)**,** 795-802.
Hai, T., Wolford, C.C., and Chang, Y.S. (2010). ATF3, a hub of the cellular adaptive-response network,
in the pathogenesis of diseases: is modulation of inflammation a unifying component? *Gene*
Expr 15(1)**,** 1-11. doi: 10.3727/105221610x12819686555015.
Haller, O., Staeheli, P., Schwemmle, M., and Kochs, G. (2015). Mx GTPases: dynamin-like antiviral
machines of innate immunity. *Trends Microbiol* 23(3)**,** 154-163. doi:
10.1016/j.tim.2014.12.003.
Hamers, A.A., Hanna, R.N., Nowyhed, H., Hedrick, C.C., and de Vries, C.J. (2013). NR4A nuclear
receptors in immunity and atherosclerosis. *Curr Opin Lipidol* 24(5)**,** 381-385. doi:
10.1097/MOL.0b013e3283643eac.

Hasday, J.D., Bannerman, D., Sakarya, S., Cross, A.S., Singh, I.S., Howard, D., et al. (2001). Exposure to febrile temperature modifies endothelial cell response to tumor necrosis factor-alpha. *J Appl Physiol (1985)* 90(1)**,** 90-98. doi: 10.1152/jappl.2001.90.1.90.

Herbert, F., Tchitchek, N., Bansal, D., Jacques, J., Pathak, S., Becavin, C., et al. (2015). Evidence of IL-17, IP-10, and IL-10 involvement in multiple-organ dysfunction and IL-17 pathway in acute renal failure associated to Plasmodium falciparum malaria. *J Transl Med* 13**,** 369. doi: 10.1186/s12967-015-0731-6.

Horrocks, P., Pinches, R.A., Chakravorty, S.J., Papakrivos, J., Christodoulou, Z., Kyes, S.A., et al. (2005). PfEMP1 expression is reduced on the surface of knobless Plasmodium falciparum infected erythrocytes. *J Cell Sci* 118(Pt 11)**,** 2507-2518.

Huang, G., Wen, Q., Zhao, Y., Gao, Q., and Bai, Y. (2013). NF-kappaB plays a key role in inducing CD274 expression in human monocytes after lipopolysaccharide treatment. *PLoS One* 8(4)**,** e61602. doi: 10.1371/journal.pone.0061602.

Hughes, C.E., and Nibbs, R.J.B. (2018). A guide to chemokines and their receptors. *FEBS J* 285(16)**,** 2944-2971. doi: 10.1111/febs.14466.

Jain, V., Armah, H.B., Tongren, J.E., Ned, R.M., Wilson, N.O., Crawford, S., et al. (2008). Plasma IP-10, apoptotic and angiogenic factors associated with fatal cerebral malaria in India. *Malar J* 7**,** 83. doi: 10.1186/1475-2875-7-83.

Jassal, B., Matthews, L., Viteri, G., Gong, C., Lorente, P., Fabregat, A., et al. (2020). The reactome pathway knowledgebase. *Nucleic Acids Res* 48(D1)**,** D498-D503. doi: 10.1093/nar/gkz1031.

John, C.C., Panoskaltsis-Mortari, A., Opoka, R.O., Park, G.S., Orchard, P.J., Jurek, A.M., et al. (2008a). Cerebrospinal fluid cytokine levels and cognitive impairment in cerebral malaria. *Am J Trop Med Hyg* 78(2)**,** 198-205.

John, C.C., Park, G.S., Sam-Agudu, N., Opoka, R.O., and Boivin, M.J. (2008b). Elevated serum levels of IL-1ra in children with Plasmodium falciparum malaria are associated with increased severity of disease. *Cytokine* 41(3)**,** 204-208. doi: 10.1016/j.cyto.2007.12.008.

Kraemer, S.M., Kyes, S.A., Aggarwal, G., Springer, A.L., Nelson, S.O., Christodoulou, Z., et al. (2007). Patterns of gene recombination shape var gene repertoires in Plasmodium falciparum: comparisons of geographically diverse isolates. *BMC Genomics* 8**,** 45.

Kraemer, S.M., and Smith, J.D. (2003). Evidence for the importance of genetic structuring to the structural and functional specialization of the Plasmodium falciparum var gene family. *Mol Microbiol* 50(5)**,** 1527-1538.

Kraemer, S.M., and Smith, J.D. (2006). A family affair: var genes, PfEMP1 binding, and malaria disease. *Curr Opin Microbiol* 9(4)**,** 374-380. doi: 10.1016/j.mib.2006.06.006.

Kyes, S.A., Kraemer, S.M., and Smith, J.D. (2007). Antigenic variation in Plasmodium falciparum: gene organization and regulation of the var multigene family. *Eukaryot Cell* 6(9)**,** 1511-1520.

Lambros, C., and Vanderberg, J.P. (1979). Synchronization of Plasmodium falciparum erythrocytic stages in culture. *J Parasitol* 65(3)**,** 418-420.

Lavstsen, T., Salanti, A., Jensen, A.T., Arnot, D.E., and Theander, T.G. (2003). Sub-grouping of Plasmodium falciparum 3D7 var genes based on sequence analysis of coding and non-coding regions. *Malar J* 2**,** 27.

Lee, H.S., and Kim, W.J. (2022). The Role of Matrix Metalloproteinase in Inflammation with a Focus on Infectious Diseases. *Int J Mol Sci* 23(18). doi: 10.3390/ijms231810546.

Liehl, P., Zuzarte-Luis, V., Chan, J., Zillinger, T., Baptista, F., Carapau, D., et al. (2014). Host-cell sensors for Plasmodium activate innate immunity against liver-stage infection. *Nat Med* 20(1)**,** 47-53. doi: 10.1038/nm.3424.

Lubiana, P., Bouws, P., Roth, L.K., Dorpinghaus, M., Rehn, T., Brehmer, J., et al. (2020). Adhesion between P. falciparum infected erythrocytes and human endothelial receptors follows

alternative binding dynamics under flow and febrile conditions. *Sci Rep* 10(1), 4548. doi:
10.1038/s41598-020-61388-2.
Lyke, K.E., Burges, R., Cissoko, Y., Sangare, L., Dao, M., Diarra, I., et al. (2004). Serum levels of the
proinflammatory cytokines interleukin-1 beta (IL-1beta), IL-6, IL-8, IL-10, tumor necrosis
factor alpha, and IL-12(p70) in Malian children with severe Plasmodium falciparum malaria
and matched uncomplicated malaria or healthy controls. *Infect Immun* 72(10), 5630-5637. doi:
10.1128/IAI.72.10.5630-5637.2004.
Mahittikorn, A., Kwankaew, P., Rattaprasert, P., Kotepui, K.U., Masangkay, F.R., and Kotepui, M.
(2022). Elevation of serum interleukin-1beta levels as a potential indicator for malarial
infection and severe malaria: a meta-analysis. *Malar J* 21(1), 308. doi: 10.1186/s12936-022-
04325-0.
Maier, A.G., Cooke, B.M., Cowman, A.F., and Tilley, L. (2009). Malaria parasite proteins that remodel
the host erythrocyte. *Nat Rev Microbiol* 7(5), 341-354.
Mancebo-Perez, C., Vidal, M., Aguilar, R., Barrios, D., Bardaji, A., Ome-Kaius, M., et al. (2022).
Eotaxin-2 and eotaxin-3 in malaria exposure and pregnancy. *Malar J* 21(1), 336. doi:
10.1186/s12936-022-04372-7.
Martin-Vazquez, E., Cobo-Vuilleumier, N., Lopez-Noriega, L., Lorenzo, P.I., and Gauthier, B.R.
(2023). The PTGS2/COX2-PGE(2) signaling cascade in inflammation: Pro or anti? A case
study with type 1 diabetes mellitus. *Int J Biol Sci* 19(13), 4157-4165. doi: 10.7150/ijbs.86492.
Milner, D., Jr., Factor, R., Whitten, R., Carr, R.A., Kamiza, S., Pinkus, G., et al. (2013). Pulmonary
pathology in pediatric cerebral malaria. *Hum Pathol* 44(12), 2719-2726. doi:
10.1016/j.humpath.2013.07.018.
Milner, D.A., Jr., Lee, J.J., Frantzreb, C., Whitten, R.O., Kamiza, S., Carr, R.A., et al. (2015).
Quantitative Assessment of Multiorgan Sequestration of Parasites in Fatal Pediatric Cerebral
Malaria. *J Infect Dis* 212(8), 1317-1321. doi: 10.1093/infdis/jiv205.
Miyashita, M., Oshiumi, H., Matsumoto, M., and Seya, T. (2011). DDX60, a DEXD/H box helicase,
is a novel antiviral factor promoting RIG-I-like receptor-mediated signaling. *Mol Cell Biol*
31(18), 3802-3819. doi: 10.1128/MCB.01368-10.
Mohamed, I.N., Li, L., Ismael, S., Ishrat, T., and El-Remessy, A.B. (2021). Thioredoxin interacting
protein, a key molecular switch between oxidative stress and sterile inflammation in cellular
response. *World J Diabetes* 12(12), 1979-1999. doi: 10.4239/wjd.v12.i12.1979.
Montoya, D., Inkeles, M.S., Liu, P.T., Realegeno, S., Teles, R.M., Vaidya, P., et al. (2014). IL-32 is a
molecular marker of a host defense network in human tuberculosis. *Sci Transl Med* 6(250),
250ra114. doi: 10.1126/scitranslmed.3009546.
Murphy, E.P., and Crean, D. (2022). NR4A1-3 nuclear receptor activity and immune cell dysregulation
in rheumatic diseases. *Front Med (Lausanne)* 9, 874182. doi: 10.3389/fmed.2022.874182.
Oakley, M.S., Gerald, N., McCutchan, T.F., Aravind, L., and Kumar, S. (2011). Clinical and molecular
aspects of malaria fever. *Trends Parasitol* 27(10), 442-449. doi: 10.1016/j.pt.2011.06.004.
Oakley, M.S., Kumar, S., Anantharaman, V., Zheng, H., Mahajan, B., Haynes, J.D., et al. (2007).
Molecular factors and biochemical pathways induced by febrile temperature in
intraerythrocytic Plasmodium falciparum parasites. *Infect Immun* 75(4), 2012-2025. doi:
10.1128/IAI.01236-06.
Othman, B., Zeef, L., Szestak, T., Rchiad, Z., Storm, J., Askonas, C., et al. (2023). Different PfEMP1-
expressing Plasmodium falciparum variants induce divergent endothelial transcriptional
responses during co-culture. *PLoS One* 18(11), e0295053. doi: 10.1371/journal.pone.0295053.
Otto, T.D., Assefa, S.A., Bohme, U., Sanders, M.J., Kwiatkowski, D., Pf3k, c., et al. (2019).
Evolutionary analysis of the most polymorphic gene family in falciparum malaria. *Wellcome*
Open Res 4, 193. doi: 10.12688/wellcomeopenres.15590.1.

Pasternak, N.D., and Dzikowski, R. (2009). PfEMP1: an antigen that plays a key role in the
pathogenicity and immune evasion of the malaria parasite Plasmodium falciparum. *Int J*
Biochem & Cell Biol 41(7), 1463-1466.

Phillips, M.A., Burrows, J.N., Manyando, C., van Huijsduijnen, R.H., Van Voorhis, W.C., and Wells,
T.N.C. (2017). Malaria. *Nat Rev Dis Primers* 3, 17050. doi: 10.1038/nrdp.2017.50.

Prakash, D., Fesel, C., Jain, R., Cazenave, P.A., Mishra, G.C., and Pied, S. (2006). Clusters of cytokines
determine malaria severity in Plasmodium falciparum-infected patients from endemic areas of
Central India. *J Infect Dis* 194(2), 198-207. doi: 10.1086/504720.

Raacke, M., Kerr, A., Dorpinghaus, M., Brehmer, J., Wu, Y., Lorenzen, S., et al. (2021). Altered
Cytokine Response of Human Brain Endothelial Cells after Stimulation with Malaria Patient
Plasma. *Cells* 10(7). doi: 10.3390/cells10071656.

Rask, T.S., Hansen, D.A., Theander, T.G., Gorm Pedersen, A., and Lavstsen, T. (2010). Plasmodium
falciparum erythrocyte membrane protein 1 diversity in seven genomes--divide and conquer.
PLoS Comput Biol 6(9). doi: 10.1371/journal.pcbi.1000933.

Sajid, M., Ullah, H., Yan, K., He, M., Feng, J., Shereen, M.A., et al. (2021). The Functional and
Antiviral Activity of Interferon Alpha-Inducible IFI6 Against Hepatitis B Virus Replication
and Gene Expression. *Front Immunol* 12, 634937. doi: 10.3389/fimmu.2021.634937.

Smith, J.D., Craig, A.G., Kriek, N., Hudson-Taylor, D., Kyes, S., Fagan, T., et al. (2000a).
Identification of a Plasmodium falciparum intercellular adhesion molecule-1 binding domain:
a parasite adhesion trait implicated in cerebral malaria. *Proc Natl Acad Sci U S A* 97(4), 1766-
1771.

Smith, J.D., Subramanian, G., Gamain, B., Baruch, D.I., and Miller, L.H. (2000b). Classification of
adhesive domains in the Plasmodium falciparum erythrocyte membrane protein 1 family. *Mol*
Biochem Parasitol 110(2), 293-310.

Sun, Z., Li, Y., Ransburgh, R., Snijder, E.J., and Fang, Y. (2012). Nonstructural protein 2 of porcine
reproductive and respiratory syndrome virus inhibits the antiviral function of interferon-
stimulated gene 15. *J Virol* 86(7), 3839-3850. doi: 10.1128/JVI.06466-11.

Szklarczyk, D., Kirsch, R., Koutrouli, M., Nastou, K., Mehryary, F., Hachilif, R., et al. (2023). The
STRING database in 2023: protein-protein association networks and functional enrichment
analyses for any sequenced genome of interest. *Nucleic Acids Res* 51(D1), D638-D646. doi:
10.1093/nar/gkac1000.

Tanaka, T., Narazaki, M., and Kishimoto, T. (2014). IL-6 in inflammation, immunity, and disease.
Cold Spring Harb Perspect Biol 6(10), a016295. doi: 10.1101/cshperspect.a016295.

Taylor, T.E., Fu, W.J., Carr, R.A., Whitten, R.O., Mueller, J.S., Fosiko, N.G., et al. (2004).
Differentiating the pathologies of cerebral malaria by postmortem parasite counts. *Nat Med*
10(2), 143-145. doi: 10.1038/nm986.

Tilly, A.K., Thiede, J., Metwally, N., Lubiana, P., Bachmann, A., Roeder, T., et al. (2015). Type of in
vitro cultivation influences cytoadhesion, knob structure, protein localization and transcriptome
profile of Plasmodium falciparum. *Sci Rep* 5, 16766. doi: 10.1038/srep16766.

Trager, W., and Jensen, J.B. (2005). Human malaria parasites in continuous culture. 1976. *J Parasitol*
91(3), 484-486. doi: 10.1645/0022-3395(2005)091[0484:HMPICC]2.0.CO;2.

Tretina, K., Park, E.S., Maminska, A., and MacMicking, J.D. (2019). Interferon-induced guanylate-
binding proteins: Guardians of host defense in health and disease. *J Exp Med* 216(3), 482-500.
doi: 10.1084/jem.20182031.

Tripathi, A.K., Sha, W., Shulaev, V., Stins, M.F., and Sullivan, D.J., Jr. (2009). Plasmodium
falciparum-infected erythrocytes induce NF-kappaB regulated inflammatory pathways in
human cerebral endothelium. *Blood* 114(19), 4243-4252. doi: 10.1182/blood-2009-06-226415.

Tripathi, A.K., Sullivan, D.J., and Stins, M.F. (2006). Plasmodium falciparum-infected erythrocytes
increase intercellular adhesion molecule 1 expression on brain endothelium through NF-
kappaB. *Infect Immun* 74(6)**,** 3262-3270. doi: 10.1128/IAI.01625-05.
Ullah, H., Sajid, M., Yan, K., Feng, J., He, M., Shereen, M.A., et al. (2021). Antiviral Activity of
Interferon Alpha-Inducible Protein 27 Against Hepatitis B Virus Gene Expression and
Replication. *Front Microbiol* 12**,** 656353. doi: 10.3389/fmicb.2021.656353.
Van den Steen, P.E., Deroost, K., Van Aelst, I., Geurts, N., Martens, E., Struyf, S., et al. (2008).
CXCR3 determines strain susceptibility to murine cerebral malaria by mediating T lymphocyte
migration toward IFN-gamma-induced chemokines. *Eur J Immunol* 38(4)**,** 1082-1095. doi:
10.1002/eji.200737906.
Viebig, N.K., Wulbrand, U., Forster, R., Andrews, K.T., Lanzer, M., and Knolle, P.A. (2005). Direct
activation of human endothelial cells by Plasmodium falciparum-infected erythrocytes. *Infect*
Immun 73(6)**,** 3271-3277. doi: 10.1128/IAI.73.6.3271-3277.2005.
Voss, T.S., Healer, J., Marty, A.J., Duffy, M.F., Thompson, J.K., Beeson, J.G., et al. (2006). A var
gene promoter controls allelic exclusion of virulence genes in Plasmodium falciparum malaria.
Nature 439(7079)**,** 1004-1008.
Waterkeyn, J.G., Cowman, A.F., and Cooke, B.M. (2001). Plasmodium falciparum: gelatin enrichment
selects for parasites with full-length chromosome 2. implications for cytoadhesion assays. *Exp*
Parasitol 97(2)**,** 115-118. doi: 10.1006/expr.2000.4593.
WHO (2023). World malaria report 2023.
Wilson, N.O., Jain, V., Roberts, C.E., Lucchi, N., Joel, P.K., Singh, M.P., et al. (2011). CXCL4 and
CXCL10 predict risk of fatal cerebral malaria. *Dis Markers* 30(1)**,** 39-49. doi: 10.3233/DMA-
2011-0763.
Wu, Y., He, X., Huang, N., Yu, J., and Shao, B. (2020). A20: a master regulator of arthritis. *Arthritis*
Res Ther 22(1)**,** 220. doi: 10.1186/s13075-020-02281-1.
Wu, Y., Leyk, S., Torabi, H., Hohn, K., Honecker, B., Tauler, M., et al. (2023). Plasmodium falciparum
infection reshapes the human microRNA profiles of red blood cells and their extracellular
vesicles. *iScience* 26(7)**,** 107119. doi: 10.1016/j.isci.2023.107119.
Zhao, Y., Zhao, K., Wang, S., and Du, J. (2022). Multi-functional BST2/tetherin against HIV-1, other
viruses and LINE-1. *Front Cell Infect Microbiol* 12**,** 979091. doi: 10.3389/fcimb.2022.979091.
Zhu, J., Ghosh, A., and Sarkar, S.N. (2015). OASL-a new player in controlling antiviral innate
immunity. *Curr Opin Virol* 12**,** 15-19. doi: 10.1016/j.coviro.2015.01.010.

Figures

Figure 1

Venn diagram showing the number of overlapping and differentially expressed genes in HBEC-5i cells
of the groups niRBC true vs. false (yellow circle), febrile true vs. false (red circle) and iRBC true vs.
false (blue circle). The niRBC true vs. false group contains the genes for which the presence of niRBC
affects the expression of HBEC-5i cells. The febrile true vs. false group contains genes for which
increasing the culture temperature from 37°C to 40°C has an effect on the expression of HBEC-5i cells.
The group iRBC true vs. false contains the genes for which the presence of iRBCs changes the
expression of HBEC-5i cells. Each circle represents one of the groups, with overlaps including genes
present in more than one group.

Figure 2

Protein-protein interaction network analysis. A STRING analysis was performed to identify protein-protein interaction networks that are regulated in HBEC-5i cells in response to the presence of niRBCs, febrile temperature and/or iRBCs. The nodes represent proteins and the connecting lines between the nodes indicate known or predicted interactions. All interaction networks that are shown are up-regulated compared to the respective controls. The thickness of the connecting lines indicates the confidence level of the interaction. Nodes marked with an asterisk are proteins whose encoding genes were identified in the respective analyses. A selection of significant networks from the GO-BP analysis is shown. (**A**) Comparison febrile true vs. false, contains 98 group-specific significantly regulated genes. (**B**) Comparison iRBC true vs. false, contains 318 group-specific significantly regulated genes. (**C**) Intersection of the groups niRBC true vs. false and iRBC true vs. false. (**D**) Intersection of the groups iRBC true vs. false and febrile true vs. false. (**E**) Intersection of the groups niRBC true vs. false, iRBC true vs. false, and febrile true vs. false. Genes exhibiting significant differential expression, meeting stringent criteria including a fold change of ≥ 2.0, a FDR of ≤ 0.05, and a minimum threshold of TPM of ≥ 2.0, were considered for inclusion in the analysis.

Figure 3

Identification of the pathways regulated by the presence of iRBCs in HBEC-5i cells. (**A**) A STRING analysis was performed to identify protein-protein interaction networks regulated in HBEC-5i cells in response to the presence of iRBCs (iRBC true vs. false including 762 genes). The nodes represent proteins and the connecting lines between the nodes indicate known or predicted interactions. The thickness of the connecting lines indicates the confidence level of the interaction. Nodes marked with an asterisk are proteins whose encoding genes were identified in the analyses. A selection of significant networks from the GO-BP analysis is shown. Genes exhibiting significant differential expression, meeting stringent criteria including a fold change of ≥ 2.0, a FDR of ≤ 0.05, and a minimum threshold of TPM of ≥ 2.0, were considered for inclusion in the analysis. (**B**) ShinyGO analysis to categorize the proteins encoded by the identified genes according to their biological functions. The 10 most significant pathways are shown.

Figure 4

Pathway analysis of the specific response of HBEC-5i cells due to the cytoadhesion of iRBCs$^{37°C}$ and iRBCs$^{40°C}$ co-cultivation. g:Profiler ((**A**), (**C**), (**E**)) and ShinyGO ((**B**), (**D**), (**F**)) analysis of the significantly up-regulated genes (fold change ≥ 2.0, FDR ≤ 0.05, and minimum threshold of TPM of ≥ 2.0) of the comparisons (**A**), (**B**): niRBC$^{37°C}$ vs. iRBC$^{37°C}$; (**C**), (**D**): niRBC$^{37°C}$ vs. iRBC$^{40°C}$; (**E**), (**F**): iRBC$^{37°C}$ vs. iRBC$^{40°C}$.

Figure 5

Identification of the pathways regulated by the presence of iRBCs in HBEC-5i cells at 37°C and 40°C using a STRING analysis. Nodes represent proteins and the connecting lines between the nodes

indicate known or predicted interactions. The thickness of the connecting lines indicates the confidence
level of the interaction. Nodes marked with an asterisk are proteins whose encoding genes were
identified in the analyses. A selection of significant networks from the GO-BP analysis is shown. Genes
exhibiting significant differential expression, meeting stringent criteria including a fold change of $\geq$
2.0, a FDR of ≤ 0.05, and a minimum threshold of TPM of ≥ 2.0, were considered for inclusion in the
analysis. STRING analysis was performed to identify protein-protein interaction networks regulated in
HBEC-5i cells after co-cultivation of (**A**) iRBCs$^{37°C}$ in comparison to niRBCs$^{37°C}$, (**B**) iRBCs$^{40°C}$ in
comparison to niRBCs$^{40°C}$, and **C** iRBCs$^{37°C}$ in comparison to iRBCs$^{40°C}$.

Figure 6

Compilation of genes whose expression is increased by the presence of iRBCs and/or an elevated
temperature compared to the control. (**A**) Cytokine encoding genes. (**B**) Genes encoding
metalloproteinases and NR4A nuclear hormone receptor proteins. (**C**) Genes encoding histones and
histone-linker proteins, (**D**) Genes encoding proteins of the interferon-alpha/beta signaling pathway.
The expression level is shown as TMP. Blue: Expression level in the presence of niRBCs$^{37°C}$; Red:
Significant increase in expression (>2-fold) due to the presence of iRBCs$^{37°C}$; Red-brown: Significant
increase in expression (>2-fold) due to the presence of iRBCs$^{40°C}$ in comparison to iRBCs$^{37°C}$. (**E**)
Reactome analyses of the genes listed in D. The pathways to which the genes can be assigned are
shown in yellow.

Supplementary Table 1

A

	niRBC$^{37°C}$	iRBC$^{37°C}$	iRBC$^{40°C}$
CXCL1	101	84	448
CXCL2	25	55	173
CXCL3	11	33	150
CXCL5	5	6	28
CXCL6	23	44	156
CXCL8	82	392	1932
CXCL10	1.3	4.2	4.9
CXCL11	1	3.4	2.8
CCL3	0.4	4.7	31
CCL3L1	0.4	10	25
CCL5	34	91	74
CCL20	0.2	3.7	6
CCL26	1	3	26
IL1A	3.4	10.7	30.5
IL1B	332	305	788
IL6	13	70	193
IL12A	10	32	53
IL32	38	33	91

B

	niRBC$^{37°C}$	iRBC$^{37°C}$	iRBC$^{40°C}$
MMP1	6	3	27
MMP2	115	173	317
MMP3	6	6	52
NR4A1	10	70	215
NR4A2	2	57	77
NR4A3	1.5	9	29

C

	niRBC$^{37°C}$	iRBC$^{37°C}$	iRBC$^{40°C}$
H1-2	73	150	116
H1-4	0.5	4.8	2.3
H2BC4	7	14	7
H2BC8	2	9	9
H2BC9	5	16	16
H2BC11	18	101	55
H2BC17	4.5	9	11
H2BC18	0.8	4.3	4.1
H2BC21	10	96	30
H3C10	3.7	11	7
H3C15	2	7	9
H4C11	3.6	12	2.5

E

	niRBC$^{37°C}$	iRBC$^{37°C}$	iRBC$^{40°C}$
BST2	43	116	74
DDIT4	202	838	806
DDX60	6	12	14
GBP4	2	9	9
IL6	13	70	193
IFI6	591	1282	814
IFI27	22	49	96
IFI44L	11	46	23
ISG15	509	1079	733
ISG20	9	20	17
MX1	78	199	95
MX2	11	37	19
OAS2	20	73	40
OAS3	32	87	62

D