Directed evolution-based discovery of ligands for in vivo restimulation of CAR-T cells

Authors: Leyuan Ma[1,7,8,10,13]*, Ranjani Ramasubramanian[12,13], Naveen K. Mehta[1], Benjamin Cossette[1], Duncan M. Morgan[1,4], Ina Sukaj[1], Elisa Bergaggio[15], Stephan Kadauke[7], Regina M. Myers[11,14], Luca Paruzzo[8,9,10], Guido Ghilardi[8,9,10], Tomasz M. Grzywa[13], Austin Cozzone[13], Stephen J. Schuster[9,10], Noelle Frey[9,10], Libin Zhang[1], Parisa Yousefpour[1], Wuhbet Abraham[1], Heikyung Suh[1], Marco Ruella[8,9,10], Stephan A. Grupp[7,11,14], Roberto Chiarle[15,16], K. Dane Wittrup[1,3,4], Darrell J. Irvine[1,2,3,5,6]*

Affiliations:
[1]David H. Koch Institute for Integrative Cancer Research, MIT, Cambridge, MA, 02139, USA.
[2]Department of Materials Science and Engineering, MIT, Cambridge, MA, 02139, USA.
[3]Department of Biological Engineering, MIT, Cambridge, MA, 02139, USA.
[4]Department of Chemical Engineering, MIT, Cambridge, MA, USA.
[5]Ragon Institute of Massachusetts General Hospital, Cambridge, MA, 02139, USA.
[6]Howard Hughes Medical Institute, Chevy Chase, MD, 20815, USA.
[7]Department of Pathology and Laboratory Medicine, Perelman School of Medicine, University of Pennsylvania, Philadelphia, PA 19104, USA.
[8]Center for Cellular Immunotherapies, University of Pennsylvania, Philadelphia, PA 19104, USA
[9]Division of Hematology/Oncology, University of Pennsylvania Perelman School of Medicine, Philadelphia, PA 19104, USA
[10]Abramson Cancer Center, University of Pennsylvania, Philadelphia, PA 19104, USA
[11]Division of Oncology, Department of Pediatrics, Perelman School of Medicine, University of Pennsylvania, Philadelphia, PA
[12]Department of Bioengineering, School of Engineering and Applied Science, University of Pennsylvania, Philadelphia, PA 19104, USA
[13]The Raymond G. Perelman Center for Cellular and Molecular Therapeutics, Children's Hospital of Philadelphia, Philadelphia, PA 19104, USA
[14]Cellular Therapy and Transplant Section and Cancer Immunotherapy Program, Children's Hospital of Philadelphia, Philadelphia, PA 19104, USA
[15]Department of Pathology, Boston Children's Hospital and Harvard Medical School, Boston, MA 02115, USA
[16]Department of Molecular Biotechnology and Health Sciences, University of Torino, 10126 Torino, Italy

* Correspondence: Leyuan.ma@pennmedicine.upenn.edu (L.M.), djirvine@mit.edu (D.J.I)

Abstract

Chimeric antigen receptor (CAR) T cell therapy targeting CD19 elicits remarkable clinical efficacy in B-cell malignancies, but many patients relapse due to failed expansion and/or progressive loss of CAR-T cells. We recently reported a strategy to potently restimulate CAR-T cells *in vivo*, enhancing their functionality by administration of a vaccine-like stimulus comprised of surrogate peptide ligands for a CAR linked to a lymph node-targeting amphiphilic PEG-lipid (termed CAR-T-vax). Here, we demonstrate a general strategy to generate and optimize peptide mimotopes enabling CAR-T-vax generation for any CAR. Using the clinical CD19 CAR FMC63 as a test case, we employed yeast surface display to identify peptide binders to soluble IgG versions of FMC63, which were subsequently affinity matured by directed evolution. CAR-T vaccines using these optimized mimotopes triggered marked expansion of both murine CD19 CAR-T cells in a syngeneic model and human CAR-T cells in a humanized mouse model of B cell acute lymphoblastic leukemia (B-ALL), and enhanced control of leukemia progression. This approach thus enables vaccine boosting to be applied to any clinically-relevant CAR-T cell product.

Introduction

CD19-targeted chimeric antigen receptor (CAR) T cell therapy elicits impressive anti-tumor activity in the treatment of relapsed or refractory B-cell acute lymphoblastic leukemia (B-ALL) and lymphoma in both pediatric and adult patients[1,2]. This clinical success led to the FDA approval of four CD19-targeted CAR-T products, namely TECARTUS, KYMRIAH, YESCARTA, and BREYANZI[1]. However, a significant fraction (30-60%) of patients still experience leukemia relapse, among which ~50% are CD19-positive relapses[3], suggesting impaired function and/or persistence of the infused CAR-T cells. Robust CAR-T cell engraftment and expansion *in vivo* is a prerequisite for antitumor efficacy[4], a feature that is affected by several characteristics of the CAR-T cell product such as T cell quality[5], CAR designs[3,4,6] and the proportion of early memory T cells in the product[7]. In current CAR-T cell therapy treatment regimens, patients usually receive lymphodepletion to promote CAR-T cell engraftment[8]. Interestingly, multiple reports have shown a correlation between the early-stage expansion of CAR-T cells and initial tumor burden[8,9]. These findings suggest the capacity of CAR-T cells to receive antigen-specific stimulation post infusion is critical for CAR-T cell expansion, engraftment, and potentially long-term persistence. However, whether the antigens presented by tumor cells are sufficient to optimally stimulate CAR-T cells remains unclear.

Vaccination is a natural process whereby T-cells receive antigen-specific stimulation leading to proliferation, differentiation, and induction of effector functions[10]– effects which could be beneficial in promoting CAR-T cell function *in vivo*. However, traditional vaccines provide antigen that must be processed into peptides by professional antigen presenting cells (APCs) and presented on MHC molecules to stimulate T-cells through the T-cell receptor (TCR). This is challenging to adapt to the setting of engineered CAR-T cells, as current CAR-T products are polyclonal and express uncharacterized TCRs. Further, recent next-generation engineering efforts such as knocking CARs into the TCR locus produce CAR-T lacking endogenous TCRs[11]. Proof of concept demonstration of CAR-T cells generated from virus-specific T cells followed by vaccine boosting via the viral-specific TCR has been shown in small clinical trials, yet the benefits observed with this approach have been limited[12,13].

To overcome these issues, we recently developed a molecularly-targeted amphiphilic vaccine (amph-vax) approach to directly stimulate CAR-T cells through the chimeric antigen receptor[14,15]. Amph-vax molecules were generated by linking a CAR-ligand to albumin-binding phospholipid polymers. Following parenteral injection, these CAR-ligand conjugates associate with endogenous albumin present in interstitial fluid and are carried into lymph and downstream draining lymph nodes (dLNs). In the dLN, the lipid tails of amph-vax ligands insert into cell membranes, including membranes of professional APCs, allowing APCs to directly "present" the CAR ligand to CAR-T cells together with additional co-stimulation and cytokine support, making them *de novo* CAR-T cell-priming APCs (**Extended Data Fig. 1**)[14]. Amph-vax stimulation led to pronounced CAR-T cell expansion with enhanced functionality, memory formation and substantially enhanced tumor clearance[14,15]. Importantly, we also found that vaccine boosting of CAR-T cells promotes antigen spreading and the induction of endogenous anti-tumor immune responses, which prevent antigen loss-mediated tumor escape[15]. We demonstrated these effects using CARs that recognize synthetic ligands (e.g., fluorescein) and a model EGFRvIII-specific CAR, where the scFv domain of the chimeric receptor recognizes a linear peptide epitope in EGFRvIII that is readily synthesized as an amph-ligand. However, to generalize this strategy and enable amph-vax boosting with any arbitrary CAR of interest, an approach is needed for *de novo* generation of surrogate peptide ligands for chimeric antigen receptors.

Here, we demonstrate a workflow to identify peptide mimotope ligands for CAR receptors using yeast surface display-based directed evolution[16], which allows for the generation of amph-vaccines that can be used with any CAR system. We demonstrate the pipeline by discovering an amph-ligand that can be used with all four FDA-approved CD19 CAR-T cell therapies. We created

a customized mimotope yeast library to identify peptides that bind to the licensed CD19 CARs.
Affinity maturation of hits from this library led to identification of a peptide with high affinity for the
CAR. An amphiphile mimotope (amph-mimotope) vaccine based on this peptide efficiently
stimulated CD19 CAR-T cells *in vitro*, triggered their expansion *in vivo*, and enhanced clearance
of tumors in humanized mouse models of leukemia. To show generality, we also generated amph-
mimotopes for a second human CAR recognizing the anaplastic lymphoma kinase (ALK) antigen
and a murine CD19 CAR. Collectively, these results show the successful development of a
clinically translatable amph-mimotope vaccine with the potential endowing currently FDA-
approved CD19 CAR-T cell therapies with enhanced anti-leukemic activity.
163
164 **Results**
165
Antigen[+] tumor cells drive transient CD19 CAR-T expansion but are insufficient to promote
long-term CAR-T persistence

CD19 positive relapse is often associated with limited CAR-T persistence and/or impaired
functionality[3,17]. To provide insight into whether the level of available antigen stimulation impacts
early CAR-T expansion/persistence, we analyzed data from several early clinical trials in children
and young adults (CAYA) with B-ALL from the Children's Hospital of Philadelphia (NCT01626495
[18] and NCT02906371[19]) and in adult patients with B-ALL (NCT02030847[20]) or B-cell lymphoma
(NCT02030834[21]) from the University of Pennsylvania. We found that CAR-T cell expansion is
generally associated with initial tumor burden, with CAR-T cells infused into patients with high
tumor burden (>40% bone marrow blasts in pre-infusion bone marrow aspirate) exhibiting more
pronounced expansion and longer persistence (**Fig. 1A-C, Extended Data Fig. 2A-B**). This
association is statistically significant in CAYA with B-ALL (**Fig. 1A-C**) and exhibits a trend towards
significance in adult B-ALL and B-cell lymphoma (**Extended Data Fig. 2A-B**), consistent with
observations from earlier studies[8,9].
Since CD19 CAR-T cells target normal B cells, continued B cell aplasia can be used as a
pharmacodynamic measure for the presence of functional CD19 CAR-T cells[22]. Nearly half of the
pediatric B-ALL patients receiving CD19 CAR-T cell therapy experienced B cell recovery after
initial B cell aplasia (**Fig. 1D**), indicating loss of functional CAR-T cells. Interestingly, there was
no significant association between the initial tumor burden and the probability of B cell recovery,
suggesting that increased early stimulation from tumor cells was not sufficient to maintain CAR-T
function. A recent long-term follow-up study found that a higher ratio of peak CAR-T expansion to
tumor burden correlates with overall survival and is a better prediction of long-term survival than
the absolute magnitude of T cell expansion or tumor burden[23]. These data suggest that if CAR-T
cells receive greater antigen stimulation following infusion, they can undergo greater expansion,
an important correlate of response to therapy[23,24]. However, sustained anti-tumor function over
time requires that T-cells receive restimulation by professional APCs, particularly dendritic cells
(DCs), which express high levels of costimulatory receptors and cytokines that are lacking on
tumor cells and resting B cells[25,26]. By mining a published dataset[27], we observed significantly
higher co-stimulatory molecule expression in DCs than healthy and likely-malignant B cells in B-
ALL patients (**Fig. 1E, Extended Data Fig. 3A-H**). We previously also observed that the
comprehensive suite of costimulatory molecules expressed by DCs is essential for promoting
memory differentiation and optimal amplification of CAR-T cells[14]. These observations suggested
to us that an efficient and safe DC-targeted vaccination approach to reinvigorate and sustain
CAR-T cell function could be impactful for enhancing current clinical CD19-targeting CAR-T cell
therapies.
201
Discovery of peptide ligands for CARs using yeast surface display

We previously demonstrated that a CAR-specific amph-vaccine could provide potent stimulation
of CAR-T cells *in vivo*[14,15]. The vaccine was created by linking a CAR ligand to an albumin-binding
PEG-lipid, which upon co-administration with adjuvant, would efficiently traffic to draining lymph
nodes and decorate the surface of APCs, allowing presentation of the ligand to CAR-T cells in
the LN (**Extended Data Fig. 1**). We demonstrated this proof of concept using CARs that
recognized arbitrary small molecule ligands (e.g., fluorescein) or linear peptide epitopes. However,
the antibody-based antigen-binding domain of the FDA-approved CD19 CARs (which employ an
antibody clone known as FMC63) bind to a conformational epitope spanning multiple domains of
the human CD19 antigen[28]. Further, the CD19 extracellular domain (ECD) is difficult to express
and prone to misfolding[29], making it challenging to use recombinant CD19 for manufacturing an
amphiphile vaccine to stimulate CD19 CAR-T cells. As it is common for CAR binding domains to
recognized complex conformational epitopes[30–32], we sought to establish a general strategy to
generate simple surrogate ligands for any CAR. We hypothesized that generation of mimotopes,
linear synthetic peptides that are also recognized by the CAR binding domain in addition to its
native target antigen, could be an ideal option (**Fig. 2A**).
To identify mimotopes for FMC63, we employed a yeast surface display library[32]
presenting ~5×10^8 randomized linear peptides of 10 amino acids (AA, 10mer) for screening
against recombinant FMC63 expressed as a full-length IgG (hereafter, FMC63^{IgG}, **Fig. 2B**). Initial
screening was performed using biotinylated FMC63^{IgG} attached to streptavidin beads for magnetic
enrichment of weakly-bound yeast clones. After a single round of screening using this enrichment
protocol, flow cytometry analysis identified two yeast populations P1 and P2, which bound with
higher or lower relative affinity to FMC63^{IgG}, respectively, but not an isotype control antibody (**Fig.
2C**). The two peptides shared a common enriched sequence motif RXCPWXCXXX. The presence
of two cysteines in the mimotope motif suggested the possibility that these mimotopes formed an
intramolecular disulfide bond, with the proline between the two cysteines likely promoting a hairpin
in the peptide structure[33]. Computational modeling of P1 (RHCPWNCSLL) and P2
(RICPWSCRAP) interactions with FMC63 using AlphaFold predicted the mimotopes take on a
curved alpha helical structure, suggesting the formation of a cyclic structure with an intra-peptide
disulfide bond (**Fig. 2D**).
To confirm the presence of a disulfide bond within the mimotope and test whether the
disulfide is necessary for peptide binding to FMC63^{IgG}, we treated the yeast clone P1 with
increasing concentrations of the reducing agent dithiothreitol (DTT, **Fig. 2E**). Treatment with 10
mM DTT lowered Aga2p levels on the yeast cells by ~10-fold, likely due to cleavage of the
disulfide linkages anchoring the protein to the yeast cell surface. However, mild reductive
conditions (i.e., 0.1mM, 1mM DTT) largely preserved Aga2p on yeast surface but eliminated
FMC63^{IgG} binding (**Fig. 2E**). To further verify that this loss in antibody binding reflected loss of a
disulfide in the mimotope itself, we chemically synthesized a cyclic version of mimotope P1 and
monitored its binding to FMC63^{IgG} using ELISA in the presence or absence of pre-treatment of
the peptide with DTT, and found that FMC63^{IgG} recognized plate-bound P1 with an estimated K_D
of 1.4 µM, but this binding was completely ablated for DTT-treated P1 (**Fig. 2F**).
Next, we created amph-mimotope molecules by conjugating P1 functionalized with an N-
terminal azide to DSPE-PEG2k-DBCO through click chemistry (**Fig. 2G**). To test whether the
amph-mimotope could insert in cell membranes and enable CD19 CAR-T cell recognition, we
incubated CD19-negative K562 cells with amph-P1to allow cell association, washed the cells, and
then incubated them with human CD19 CAR-T cells *in vitro* (**Fig. 2G**). As shown in **Fig. 2H**, co-
culture of CD19 CAR-T with K562 cells decorated by amph-P1 but not a control amph-peptide
(pepvIII, a minimal epitope from EGFRvIII) triggered robust IFN-γ secretion from the CAR-T cells.
Consistent with the need for the P1 to contain an intramolecular disulfide for CAR recognition,
CAR-T stimulation was lost when amph-P1-labeled cells were pre-treated with DTT (**Fig. 2H**). By
contrast, control EGFRvIII CAR-T cells had no response to amph-P1 decorated K562 cells

irrespective of the disulfide bond (**Fig. 2I**). These results confirmed that P1 is a functional mimotope with a cyclic conformation required for its specific recognition by FMC63.

To demonstrate that the yeast mimotope library is suitable for identifying mimotopes for any desired antibody, we repeated the library screen using an anti-mouse CD19 antibody (Clone 1D3) (**Extended Data Fig. 4A-D**) and an anti-human Anaplastic Lymphoma Kinase (ALK) antibody of interest for CAR-T cell treatment of lymphoma and some lung cancers (**Extended Data Fig. 4E-F**). These additional library screens successfully identified multiple binders for each antibody, highlighting the potential of yeast surface display to effectively identify antibody-specific mimotope peptides.

Affinity maturation of the CD19 mimotope

The affinity of a typical CAR toward its ligand is often low nanomolar[34], however, mimotope P1 only exhibited a low micromolar apparent affinity toward FMC63[IgG]. We therefore aimed to further evolve the mimotope to obtain variants with enhanced affinity. To this end, we generated a new mimotope library V2 bearing a shared motif derived from the two parental mimotopes, namely RXCPWXCXXX, with a diversity of $\sim 1.5 \times 10^8$ (**Fig. 3A**). A significant portion of the library V2 could be positively stained with 20 nM biotinylated FMC63[IgG] while the original P1 yeast clone showed no detectable binding at this concentration (**Fig. 3B**). Flow cytometry sorting of the top 1% of the positive yeast cells followed by Sanger sequencing yielded dozens of mimotope clones. Sequence analysis of these mimotopes revealed a consensus sequence with high-affinity binding potential to FMC63[IgG] (**Fig. 3C**). Several candidate mimotope clones resembling the consensus sequence were validated experimentally, and all of them exhibited stronger binding than the original clone P1 (**Fig. 3D**). We moved forward with mimotope F12 given its highest resemblance to the consensus sequence and highest binding affinity to FMC63[IgG] (apparent K_D = 15.6 nM). Alanine scanning across all AA positions in the peptide confirmed the determining role of R at position 1°, C at positions 3° and 7°, P at position 4°, and W at position 5° (**Fig. 3E**). Residues at the remaining AA positions contributed more modestly to the overall affinity. Interestingly, inserting an extra 1-2 alanines between position 6° and 7° completely disrupted F12 binding to FMC63 (**Fig. 3E**), likely through the disruption of the intra-mimotope disulfide bond. Given the labile nature of the disulfide bond in the mimotope, we sought to test if the disulfide bond can be replaced with a structurally similar yet non-reducible thioacetal bond to produce a more stable mimotope (**Extended Data Fig. 5A-B**). Replacing the disulfide bond with a thioacetal bond completely eliminated its binding to FMC63[IgG], perhaps because the thioacetal bond is 0.9 angstrom (Å) longer than the disulfide bond (**Extended Data Fig. 5C**).

Next, we sought to evaluate the contribution of the flanking sequence to mimotope binding with FMC63[IgG]. A 17 AA linker (GGGGS)$_3$AS was initially introduced between the 10-mer mimotope and Aga2p during yeast library construction to avoid steric hindrance. We chemically synthesized F12 mimotope variants with increasing length of the upstream linker and performed ELISA to verify FMC63[IgG] binding. Strikingly, the addition of even a minimal SAS tri-amino acid linker at the N-terminus of the peptide increased binding of F12 to FMC63[IgG] by $\sim$100-fold (**Fig. 3F**). Given the importance of flanking sequence on mimotope binding to FMC63[IgG], we created a third yeast library V3 with $\sim 1 \times 10^8$ diversity with 0-6 and 3-10 flanking residues extended from the N- and C-termini of F12, respectively (**Fig. 3G**), aiming to identify mimotope variants with even higher affinity. We performed kinetic sorting by competing off pre-bound biotinylated monovalent FMC63[scFv] using non-modified bivalent FMC63[IgG], and were able to identify yeast populations with prolonged retention of biotinylated FMC63[scFv], indicative of slower off-rates. Characterization of individual clones identified mimotope variants with further increased affinity compared to the parental F12 clone (**Fig. 3H**). Notably, clone F12-A1 exhibited a further $\sim$10-fold increased affinity toward FMC63[scFv] compared to F12. In the same competition assay as depicted in **Fig. 3G**, FMC63[scFv] remained bound on F12-A1 yeast surface over 24 hours in the presence of excess

303 soluble FMC63^{IgG}, but was competed off the parental F12 clone within half an hour (**Fig. 3I**). Thus,
304 these sequential affinity maturation screens effectively enabled identification of mimotopes with
305 much-improved affinity for the CAR antibody.
306
Optimized mimotopes bind to FMC63 without compromising its recognition of CD19

To gain a comprehensive view of how the CD19 mimotope interacts with FMC63, we leveraged a
recently solved crystal structure of FMC63^{scFv} in complex with CD19 (PDB:7URV)[30] and modeled
mimotope binding to FMC63^{scFv}. Both F12 and F12-A1 were modeled using Alphafold[35] and
Rosetta[36], and the FMC63^{scFv}-CD19, FMC63^{scFv}-F12, and FMC63^{scFv}-F12A1 binding interfaces
were analyzed using PDBePISA[37]. PDBePISA analysis recovered many of the previously defined
interactions between FMC63^{scFv} and CD19 through analysis of polarities and bond lengths **(Fig.**
4A), confirming the validity of PDBePISA in performing interaction interface analysis. Structural
modeling of F12 (**Fig. 4B**) or F12-A1 (**Fig. 4C**) interactions with FMC63^{scFv} showed that these
mimotopes engage FMC63^{scFv} at the same domain. The key residues on FMC63^{scFv} that are
essential for CD19 binding were found to be Y260, Y261, Y70, G263, W212, G262, S214, K69,
Y265, and G129 [30]. Of these key residues, both F12 and F12-A1 engage Y260 and S214 **(Fig.**
4D). However, only F12-A1 engages G263 (**Fig. 4D**), which likely explains its stronger binding to
FMC63^{scFv} than F12.
To experimentally assess potential overlaps in the binding interface of FMC63 to CD19 vs.
our mimotope peptides, we pre-blocked FMC63^{scFv}, FMC63^{IgG} or an alternative control anti-CD19
antibody HIB19^{IgG} with increasing concentrations of mimotopes F12 or F12-A1, and then added
the antibody to CD19$^+$ NALM6 cells (**Fig. 4E**). When FMC63^{scFv} was pre-blocked with F12 peptide,
binding to NALM6 cells was only compromised when the F12 concentration was 100 µM, and
NALM6 binding was nearly unaffected for FMC63^{IgG}. By contrast, F12-A1 abolished FMC63^{scFv} or
FMC63^{IgG} binding to CD19 when the mimotope concentrations were greater than 1 or 10 µM,
respectively, consistent with its higher affinity for FMC63. However, when FMC63 CAR-
expressing Jurkat T cells were pre-blocked with increasing concentrations of F12 or F12-A1 and
stimulated with NALM6 cells, CD69 induction on Jurkat cells was not affected (**Fig. 4F**). Thus,
despite some ability of the mimotopes to interfere with soluble FMC63 binding to CD19$^+$ cells at
high concentrations, FMC63 CAR-T cells retained their full potential of recognizing CD19$^+$
leukemic cells even in the presence of mimotopes.
334
Amph-mimotope vaccination triggers FMC63-mCAR-T expansion *in vivo*

To test if amph-mimotope vaccines could efficiently stimulate CD19 CAR-T cells in an
immunocompetent mouse model *in vivo*, we established a hybrid CAR by fusing the human
FMC63 scFv to murine CAR signaling domains, (FMC63-mCAR, **Fig. 5A**). First, we validated the
ability of the amph-mimotope variants to stimulate FMC63-mCAR T cells. K562 cells were pre-
labeled with amph-mimotope variants and co-cultured with FMC63-mCAR-T cells as in **Fig. 2G.**
Amph-mimotope-decorated target cells triggered IFN-γ production from the CAR-T cells, with the
stimulation efficiency positively correlating with FMC63-mimotope affinity and amph-F12-A1
providing the strongest CAR-T activation, comparable to stimulation by CD19$^+$ NALM6 cells (**Fig.**
5B). To determine if these amph-mimotopes could be effectively presented by DCs *in vivo*, we
vaccinated C57BL/6 mice with 10 nmol amph-F12-A1, a dose we previously found to be effective
for amph-peptide stimulation of CAR-T cells. We formulated the mimotope with or without cyclic-
di-GMP, a STING agonist added as an adjuvant[38]. Following vaccination, we isolated CD11b$^+$
myeloid cells and CD11c$^+$ DCs from the draining inguinal lymph nodes and analyzed the presence
of surface-displayed amph-peptides using flow cytometry by staining the cells with biotinylated
FMC63^{IgG} (**Extended Data Fig. 6A**). Interestingly, high levels of amph-mimotope were detected

on APCs for at least 24 hr, but only when amph-mimotope was co-administered with adjuvant
(**Fig. 5C**).
Next, we labeled FMC63-mCAR cells with cell trace violet (CTV) and transferred them into
C57BL/6 mice, followed by vaccination with amph-P1, amph-F12, amph-F12-A1 vaccine, or saline
as a control, followed by flow cytometry analysis to detect CAR-T cell proliferation (CTV dilution)
2 days later. This functional test of amph-mimoptope stimulation revealed a clear hierarchy of
mimotope activity *in vivo*, with amph-P1 eliciting no CAR-T cell division, amph-F12 eliciting limited
proliferation, and only amph-F12-A1 triggering proliferation of the entire CAR-T population (**Fig.
5D**). These data suggest there is an affinity threshold for mimotope vaccines to effectively
stimulate CAR-T cells *in vivo*.
Clinical evidence indicates that lack of antigen exposure soon after adoptive transfer may
compromise CAR-T engraftment[39]. To test if amph-mimotope vaccine boosting post adoptive cell
transfer could promote FMC63-mCAR-T cell expansion and engraftment in the absence of
leukemia, naïve CD45.2$^+$ C57BL/6 mice received sublethal lymphodepletion followed by
intravenous infusion of either CD45.1$^+$ FMC63-mCAR-T or control untransduced T cells (**Fig. 5E**).
Without additional treatment, peripheral CD45.1$^+$ CAR-T cells only occupied <5% of total CD8$^+$ T
cells by day 4 and decreased to <3% by day 14 (**Fig. 5E**). By contrast, amph-F12-A1 vaccine
boosting rapidly amplified CD45.1 CAR-T cells, reaching ~40% of total peripheral CD8$^+$ T cells
(~500 CAR-T cells per microliter of blood, **Fig. 5E, Extended Data Fig. 6B**) by day 14 after two
doses of vaccination, and then slowly contracting over the next few weeks.
371
Amph-mimotope-labeled DCs enhance human CD19 CAR-T expansion and leukemia
clearance in a humanized mouse model

Human CAR-T cells are routinely evaluated in immunodeficient NSG mice, but this animal model
is problematic for testing lymphatic-mediated amph-vaccine boosting, as these mice have
generally defective lymph node and lymphatic development due to their lack of native
lymphocytes[40]. To overcome this technical hurdle, we devised an alternate strategy to test amph-
vaccine boosting of human CAR-T cells: We previously showed that amph-ligand vaccine
molecules label lymph node-resident DCs, which are the key APC population mediating the amph-
vaccine stimulation[14]. To mimic this process in NSG mice in a human-relevant setting, we
differentiated human peripheral blood monocytes into DCs *in vitro* and matured them with
lipopolysaccharide (LPS) and IFN-γ (**Fig. 6A**). These matured human monocyte-derived DCs
(MoDCs) exhibited significantly increased expression of multiple co-stimulatory molecules,
including CD80/86, 41BBL, Ox40L, and ICOSL (**Fig. 6A**). Incubation of activated MoDCs with
amph-F12-A1 led to robust cell surface labeling with the mimotope (DC-mVax, **Fig. 6B**) and
efficiently stimulated human CD19 CAR-T cell activation *in vitro* (**Fig. 6C**). To determine if DC-
mVax could activate and expand human CD19 CAR-T cells *in vivo* in the absence of additional
antigen, we intravenously (i.v.) infused human CD19 CAR-T cells into non-leukemic NSG mice
with or without subsequent i.v. infusion of DC-mVax (**Fig. 6D-E**). Mice receiving a single dose of
DC-mVax 24 hours post CAR-T infusion exhibited significant expansion of CAR-T cells in the
peripheral blood compared to those receiving unmodified DCs, and circulating human CD19 CAR-
T cells remained detectable at day 30 (**Fig. 6E**). Notably, the extent of CAR-T cell expansion
markedly decreased when DC-mVax was infused 7 days post CAR-T cell transfer, indicating that
early antigen exposure plays an important role in promoting CAR-T expansion and engraftment
(**Fig. 6E**). Repeated dosing of DC-mVax on day 1 and day 7 days led to a further increase in CAR-
T cell expansion (**Fig. 6E**).
To determine if DC-mVax boosting could augment the anti-tumor activity of human CD19
CAR-T cells, we established leukemia in NSG mice by infusion of luciferase-expressing NALM6
tumor cells as previously described[41]. Suboptimal doses of CD19 CAR-T cells were administered
on day 4 followed 24 hours later by repeated i.v. infusion of either unmodified DCs or DC-mVax

(**Fig. 6F**). Despite the presence of high levels of target cells, CAR-T cells alone showed poor persistence/expansion compared to CAR-T cells boosted with DC-mVax (**Fig. 6G**). While CAR-T cells alone only slightly delayed tumor expansion and moderately prolonged survival, addition of the DC-mVax boosting enabled this dose of CAR-T cells to substantially slow tumor growth and 30% of the animals completed rejected their tumors (**Fig 6H-J**). Collectively, these results demonstrated that a suitably affinity-matured amph-mimotope could act as an effective vaccine to promote engraftment, expansion and reinvigoration of human T cells expressing the CAR currently used in all approved CD19-targeting CAR-T products, and enhance their anti-leukemic activity.

Discussion

Successful engraftment and persistence is critical for ensuring robust and long-term anti-leukemic activity by CAR-T cells[1,42]. Analysis of data from recent CAR-T cell clinical trials suggested that CD19-directed CAR-T cells initially expand more strongly if there is a high tumor burden, consistent with the idea of CAR-T expansion that is commensurate with the dose of antigen encountered. However, long-term persistence of CAR-T cells does not correlate with initial tumor burden, suggesting that this initial stimulus vanishes over time as leukemic cells are cleared from the body and/or that stimulation received from tumor cells or naïve B cells lacking appropriate costimulatory signals is insufficient to maintain the CAR-T cell population. We and others have recently demonstrated the ability of using a synthetic vaccine to boost CAR-T cells *in vivo* and achieved enhanced therapeutic outcomes[14,15,43]. Our strategy of linking a ligand for the CAR to a lymph node-targeting PEG-lipid amphiphile is simple, but requires a suitable peptide or protein ligand. The poor expression and folding of recombinant human CD19 ectodomain[29] makes use of the native target antigen impractical for this approach. Here we demonstrate that surrogate CAR ligands based on short peptides can be readily identified and affinity matured using directed evolution, enabling the generation of mimotope vaccines that efficiently boost CD19 CAR-T cells analogous to natural CD19 antigen exposure. Vaccine boosting significantly impacted the outcome of CD19 CAR-T cell therapy by increasing CAR-T cell expansion and engraftment, and we have shown previously, also augments CAR-T cell functionality. Importantly, the mimotope vaccine discovered here is applicable to all four FDA-approved FMC63-based CD19 CAR-T cell therapies.

In addition to the human CD19 CAR mimotope, mimotopes for anti-mouse CD19 and anti-human ALK CARs were also successfully identified using the same yeast mimotope library, indicating the broad applicability of this library for identifying mimotopes for any desired antibody. Experimental verification and structural modeling demonstrated that the CD19 mimotope discovered through the yeast library forms a cyclic conformation that is essential for its recognition by FMC63. Given that FMC63 recognizes a conformational epitope, it is possible that a mimotope also needs to form a desired secondary structure to achieve strong binding. Notably, the CD19 mimotope only partially occupies the natural epitope binding domain on FMC63. This partial occupancy appears to enable FMC63-based CD19 CAR-T cells to maintain their capacity to recognize CD19[+] leukemia cells even in the presence of the mimotope, which could be important for allowing the CAR-T cells to continue to sense tumor cells in lymph nodes during vaccine boosting. The transient surface presentation of amph-ligands also prevents CAR-T cells from receiving excessive and prolonged antigen exposure, thus reducing the likelihood of CAR-T cell exhaustion and severe cytokine release syndrome[14].

Despite expansion and efficacy of CD19 CAR-T cells following the first infusion, reinfusion of CAR-T cells into patients experiencing CD19[+] tumor relapse is often ineffective, with significantly lower CAR-T expansion compared to the first infusion[39,44–46]. While B cells retain the antigen for CAR-T, evidence suggests that B cells themselves may not be as effective as conventional DCs for restimulating CAR-T cells at least partially due to greater expression of co-

stimulatory molecules in DCs[26]. Our previous findings showed that costimulatory molecules
expressed by activated DCs are crucial for optimally expanding CAR-T cells with amph-ligands[14].
Therefore, one potential use of the mimotope vaccine is to support CAR-T cells during reinfusion.
Notably, even during the first infusion, CD19 CAR-T cells rapidly decline after a transient
expansion[9,47]. Despite the general correlation of initial CAR-T expansion with baseline tumor
burden, a long-term follow-up study in B-ALL found that a higher ratio of peak CAR-T-cell
expansion to tumor burden significantly correlated with event-free survival and overall survival[23].
Although this effective ratio more likely occurs in patients with lower tumor burden, administering
the mimotope vaccine upon CAR-T contraction during the first infusion could potentially
reinvigorate CAR-T cells and sustain CAR-T expansion to elevate this effective ratio in patients
with both low and high tumor burden to improve event-free survival.
One shortcoming of characterizing vaccines for human T cells is the lack of proper
humanized mouse models. The defective lymphatics, absent or disorganized lymph nodes, and
poor crosstalk of mouse APCs with human T cells is a barrier to evaluating vaccine boosting of
human CAR-T cells in NSG mice[40]. As an alternative, we evaluated vaccine boosting of mouse T
cells bearing a hybrid FMC63-mCAR in immunocompetent C57BL/6 mice via standard vaccine
administration routes, and boosting of human CD19 CAR-T cells using mimotope-decorated
human MoDCs in NSG mice via i.v. delivery. These two complementary approaches
demonstrated the feasibility of the mimotope vaccine in both the setting of natural trafficking and
distribution from lymphatics to lymph nodes as well as its capacity in stimulating human CAR-T
cells upon insertion into the surface of human DCs.
mRNA vaccines encoding the target tumor antigen can also be employed to stimulate
CAR-T cells[43]. Recently, an CLDN6-expressing mRNA vaccine was tested in a phase 1 clinical
trial in combination with CLDN6-directed CAR-T cells and showed promise in improving CAR-T
cell expansion in patients with solid tumors[48]. Importantly, CAR-T plus mRNA vaccination was well
tolerated, suggesting the overall safety profile associated with this combination therapy. However,
mRNA-LNPs differ from our amphiphile vaccines in their *in vivo* trafficking, biodistribution, and the
duration of antigen presentation on cell surface, therefore their differential effects on boosting
human CD19 CAR-T cells remain to be determined and is beyond the scope of this work.
In summary, we have designed and validated a general strategy to generate mimotope
ligands for any CAR T cell. Using this methodology, we demonstrate an amph-mimotope vaccine
that could be used clinically in tandem with all four FDA-approved CD19 CAR-T cell products.
The ability of amph-mimotope vaccines to efficiently stimulate human CD19 CAR-T cells *in vivo*
support its potential use as a robust and clinically-translatable approach for enhancing the
engraftment, long-term persistence, and anti-leukemic activity of current CD19 CAR-T cell
therapies.
487
488

ACKNOWLEDGMENTS
We thank the Koch Institute Swanson Biotechnology Center for technical support, specifically the flow
cytometry core facility. **Funding:** This work was supported by the NIH (award EB022433 to DJI), the
Marble Center for Nanomedicine (to DJI), and the Mark Foundation for Cancer Research (to DJI). L.M
was supported by an American Cancer Society postdoctoral fellowship, the Cell and Gene Therapy
Collaborative and the Junior faculty pilot program at CHOP, NIH New Innovators Award (DP2
AI164319-03), ITMAT at UPenn and NCATS (UL1TR001878). This work is also partially supported by
Cancer Center Support (core) Grant P30-CA14051 from the NCI to the Barbara K. Ostrom (1978)
Bioinformatics and Computing Core Facility of the Swanson Biotechnology Center. D.J.I. is an
investigator of the Howard Hughes Medical Institute. R.R. is supported by a National Science
Foundation Graduate Research Fellowship Program award. Parisa Yousefpour is supported by
an NRSA F32 fellowship (award AI164829) from the NIH. M.R. is supported by the NIH NCI P01
PCA214278C, the R01/37-CA262362-01A1, the Laffey McHugh Foundation, and the Berman and

Maguire Funds for Lymphoma Research at Penn. G.G is supported by the SITC-Mallinckrodt Pharmaceuticals Adverse Events in Cancer Immunotherapy Clinical Fellowship, and the Mario Luvini fellowship grant—Fondazione Ticinese per la Ricerca sul Cancro.

Data and materials availability: All data are available on request.

AUTHOR CONTRIBUTIONS

L.M., D.J.I. and K.D.W designed the studies. L.M. R.R. D.J.I. analyzed and interpreted the data and wrote the manuscript. L.M and N.M generated yeast mimotope libraries. L.M. performed the experiments. B.C and I.S assisted with mimotope identification. D.M.M. mined single cell RNA-sequencing data. L.M. and L.Z. carried out amphiphile-mimotope synthesis. R.R performed structural modeling. I.S. P.Y. and H.S. assisted with sample preparation. W.A. T.M.M. and A.C. assisted with animal experiments. S.K, R.M.M, S.A.G, L.P, G.G., S.J.S, N.F and M.R processed and provided the clinical data.

DECLARATION OF INTERESTS

L.M and D.J.I are inventors on patents filed related to the amphiphile-mimotope vaccine technology. G.G. served as a scientific consultant for viTToria Biotherapeutics. M.R. holds patents related to CD19 CAR-T cells, served as a consultant for NanoString, Bristol Myers Squibb,GlaxoSmithKline, Scaylite, Bayer, and AbClon, receives research funding from AbClon,NanoString, Oxford NanoImaging, viTToria biotherapeutics, CURIOX, and Beckman Coulter. M.R. is the scientific founder of viTToria Biotherapeutics

    FIGURE LEGENDS

Figure 1: CD19 CAR-T cell expansion and B cell aplasia in patients with initial high (≥40%)
or low tumor burden (<40%), and rationale for a DC-targeting CD19-CAR stimulating
vaccine. (**A**) Scatterplot of peripherally circulating CAR-T cells measured by quantitative
polymerase chain reaction (qPCR) over time during the first month after infusion. Solid lines show
trends generated by locally estimated scatterplot smoothing (loess). The median time to peak
CAR-T expansion was 10 days after infusion in both cohorts. (**B**) Maximum concentration and (**C**)
Area under the curve (AUC) of CAR-Transgene levels measured by qPCR, broken down by tumor
burden. ***, p<0.001, by Wilcoxon rank-sum tests. (**D**) Kaplan-Meier analysis of B cell aplasia for
all patients (black) or high (orange) or low (blue) tumor burden patients. B cell aplasia is defined
as the time to the emergence of ≥1% CD19-positive B cell in bone marrow aspirate or ≥3% B cell
by peripheral blood flow cytometry or CD19-positive relapse. Data were censored for patients
who had CD19-negative relapse, reinfusion for hematogones in the marrow, alternative therapy
including other CAR-T therapy or hematopoietic stem cell transplant, or non-relapse mortality. The
probability of continued B-cell aplasia at two years was similar in both cohorts (p = 0.76). At 6
months, it was 72% (95% CI 55-95) in high and 67% (95% CI 57-78) in low tumor burden patients.
(**E**) Box plot of costimulatory molecule module scores expressed by B cells (likely-healthy and
likely-malignant) and DC. Each point represents the average of all cells of that phenotype within
a patient. ****, p<0.0001, by a two-sided Wilcoxon rank-sum test.

Figure 2: Yeast surface display identifies CD19-CAR binding mimotopes. (**A**) Illustration of
CD19-CAR stimulation by natural CD19 or amph-mimotope. (B) Yeast surface display workflow
for identifying antibody-specific mimotopes. (**C**) Flow cytometry plots showing the successful
identification of yeast cell populations (P1 and P2) binding to 1 µM FMC63^{IgG} after a single round
of magnetic enrichment. (**D**) Structural modeling of mimotopes P1 and P2 binding to FMC63. (**E**)
Yeast cells bearing mimotope P1 was treated with increasing concentrations of DTT, and then
stained with 500 nM biotin-FMC63^{IgG} and an anti-HA tag antibody for analysis of binding by flow
cytometry. (**F**) ELISA analysis showing the impact of the disulfide bond on FMC63^{IgG} binding to
plate-bound P1 mimotope. (**G**) Scheme for amph-mimotope generation by click chemistry linkage
of azide-modified P1 to DBCO-PEG-DSPE, and schematic illustration of amph-P1 coating of
target cells. (**H-I**) CD19$^-$ K562 cells were labeled with control amph-pepvIII or amph-P1 molecules
in vitro followed with or without 1µM DTT treatment and subsequent co-culture with human CD19
CAR-T cells (**H**) or EGFRvIII CAR-T cells (**I**). IFN-γ ELISA was performed to monitor activation of
CAR-T cells. Error bars show mean ± s.d. with triplicate samples. **, p<0.01 by one-way ANOVA
with Turkey's post-test. A, B and G are created with BioRender.com.

Figure 3: Affinity maturation of CD19 mimotopes. (**A**) Schematic workflow for identifying
affinity-enhanced mimotope variants of P1 from a motif-fixed mimotope library V2. Created with
BioRender.com. (**B**) Flow cytometry plots of 20 nM FMC63IgG binding to yeast cells expressing
the V2 mimotope library. Yeast cells bearing mimotope P1 was included as a control. (**C**) Weblogo
showing the consensus mimotope sequence with high-affinity binding to FMC63^{IgG}. (**D**) Binding
of FMC63^{IgG} to select yeast clones assessed by flow cytometry. Amino acid sequences and
apparent binding affinity are shown for each yeast clone. (**E**) Yeast clones bearing F12 variants
with 1-2 alanine (Red) insertions or individual AA mutated to alanine (Red) were assessed for
FMC63^{IgG} binding by flow cytometry. (**F**) ELISA showing the impact of flanking residues on
mimotope^{F12} binding to FMC63^{IgG}. The apparent binding affinity was determined as in D. (**G**)
Schematic workflow for identifying affinity-enhanced mimotopes from mimotope library V3 using
kinetic sorting. Yeast library V3 were first stained with monovalent biotin-FMC63^{scFv} followed by
competition with non-biotinylated bivalent FMC63^{IgG}. Created with BioRender.com. (**H**) Binding of

FMC63^{scFv} to select yeast clones bearing various flanking sequences. The parental mimotope^{F12}
sequence was underlined. Apparent binding affinity was determined as in D. (**I**) Representative
histograms showing retention of prebound FMC63^{scFv} on yeast clone F12-A1 in the presence of
FMC63^{IgG}. Yeast cells were pre-stained with 50nM Biotin-FMC63^{scFv} followed by incubation with
1µM FMC63^{IgG}. Yeast cells were sampled at indicated times for PE-streptavidin staining as an
indicator for the retention of Biotin-FMC63^{scFv}.

Figure 4: Structural modeling of mimotopes complexed with FMC63. (A) Ribbon diagram of
FMC63^{scFv} (purple) in complex with human CD19 (green, PDB 7URV). Interactions involving key
residues of FMC63 are shown. Interacting sidechain atoms are shown and boxed. Interactions
involving backbone atoms are labeled without box. **(B-C)** Computational models generated using
Alphafold and Rosetta for FMC63^{scFv} bound to F12 (**B**) or F12-A1 (**C**), respectively. Interactions
involving previously mentioned key residues of FMC63 are labeled using same scheme as in A.
For F12-A1, the orientation of mimotope is shown to best exhibit how it engages FMC63. The
panel in the bottom left depicts the complex in the same orientation as F12-FMC63. **(D)** Interface
footprint of mimotopes on FMC63. On the left side, the CD19 interface is shown in green, the F12
interface shown in orange, and overlapping residues are shown in red. On the right side, the CD19
interface is shown in green, the F12A1 interface shown in blue, and overlapping are residues
shown in red. **(E)** Biotinylated FMC63^{scFv} was pre-incubated with various concentrations of F12
or F12-A1 prior to staining NALM6 cells and detection via PE-conjugated streptavidin. For
FMC63^{IgG} binding, NALM6 cells were stained with the same concentrations of mimotope and
FMC63^{IgG} with no prior incubation, and binding was detected using PE-conjugated anti-mouse-
IgG. A non-competing IgG clone HIB19 is shown as control. **(F)** FMC63 CAR$^+$ Jurkat T cells were
pre-incubated with F12 or F12-A1 at indicated concentrations, then co-cultured with NALM6 cells
at a 1:1 ratio for 16 hours followed by flow cytometry analysis of CD69 expression. Shown in E-F
are one representative results of three independent experiments.

Figure 5: Amph-mimotopes require a threshold affinity to stimulate CAR-T proliferation in
vivo. (**A**) Schematic of hybrid FMC63-mCAR design. (**B**) K562 cells were labeled with 100 nM of
amph-mimotope variants for 30 min, washed, and co-cultured with FMC63-mCAR-T cells at a 1:1
E:T ratio for 6 hours followed by ELISA measurement of IFN-γ secretion. Error bars show mean
± s.d. with triplicate samples. (**C**) C57BL/6 mice (*n*=3 animals/group) were vaccinated with 10
nmol amph-mimotope ± CDN adjuvant, and 24 or 48 hr later, mimotope uptake by macrophages
and cDCs was detected by staining with FMC63 followed by flow cytometry analysis. Shown are
representative histograms of FMC63 staining. (**D**) C57BL/6 mice (*n*=3 animals/group) were
lymphodepleted, adoptively transferred with 2x10^6 CTV-labeled CAR-T cells, then vaccinated 1
day later with 10 nmol amph-peptides + CDN adjuvant. Shown are representative histograms of
FMC63-mCAR-T cell proliferation in LNs 48 hours after vaccination. (**E**) C57BL/6 mice (*n*=5
animals/group) were lymphodepleted (L.D.), adoptively transferred with 10^6 FMC63 CAR-T cells,
and then vaccinated at indicated time points, and circulating FMC63-mCAR-T cells were
quantified in the blood by flow cytometry over time. Error bars show mean ± 95% CI. ***,
p<0.0001;**, p<0.01; n.s., not significant by one-way ANOVA with Turkey's post- test for B-D, two-way
ANOVA with Tukey's post-test for E.

Figure 6: Amph-mimotope vaccine-labeled DCs augment human CD19 CAR-T therapy
against leukemia. (**A**) Schematic of approach to generate "DC-mVax" and representative
histograms depicting LPS/IFNγ-induced upregulation of co-stimulatory markers on mature DCs.
(**B**) Activated MoDCs were incubated with 500 nM amph-F12-A1 or control amph-pepvIII for
30min, washed, and then stained with FMC63^{IgG}. Shown are representative histogram of

FMC63^{IgG} labeling. (**C**) Activated MoDCs were labeled with amph-F12-A1 or amph-pepvIII as in
(B) and then co-cultured human CD19 CAR-T cells at a 1:1 E:T ratio. Shown is IFN-γ measured
in the supernatant after 6 hr by ELISA. Error bars show mean ± s.d. with triplicate samples. (**D**)
Experimental setup and timeline for E. (**E**) Quantification of circulating human CD19 CAR-T cells
in the absence or presence of a single dose of DC-mVax at day 1 or day 7 or two weekly doses
of DC-mVax administered following timeline in D (n=5 animals/group). (**F**) Experimental setup and
timeline for G-J. (**G-J**) CAR-T cells were administered with or without DC-mVax as shown in
timeline (F) and tumor progression (H) was monitored using *in vivo* bioluminescent imaging (*n*=5
animals/group. Shown are enumeration of circulating human CD19 CAR-T cells on day 21 (**G**),
whole animal imaging at indicated times (**H**) and quantification of total photon counts over time
(**I**). Shown is one of two independent experiments. (**J**) NSG mice (*n*=10 animals/group) inoculated
with NALM6 tumor cells and treated with CAR-T cells ± DC-mVax as in F were followed for overall
survival. Shown are combined data from two independent experiments. Error bars show mean ±
95% CI. ***, p<0.0001; **, p<0.01 by Student's t-test for C and G, by two-way ANOVA with Tukey's
post-test for E and I, and by Log-rank (Mantel-Cox) test for J.

References

References

1. Cappell, K. M. & Kochenderfer, J. N. Long-term outcomes following CAR T cell therapy: what
we know so far. *Nat. Rev. Clin. Oncol.* **20**, 359–371 (2023).

2. Irvine, D. J., Maus, M. V., Mooney, D. J. & Wong, W. W. The future of engineered immune cell
therapies. *Science* **378**, 853–858 (2022).

3. Labanieh, L. & Mackall, C. L. CAR immune cells: design principles, resistance and the next
generation. *Nature* **614**, 635–648 (2022).

4. Bai, Z. *et al.* Single-cell antigen-specific landscape of CAR T infusion product identifies
determinants of CD19-positive relapse in patients with ALL. *Sci. Adv.* **8**, eabj2820 (2022).

5. Wang, X. & Rivière, I. Clinical manufacturing of CAR T cells: foundation of a promising therapy.
Mol. Ther. - Oncolytics **3**, 16015 (2016).

6. Neeser, A., Ramasubramanian, R., Wang, C. & Ma, L. Engineering enhanced chimeric antigen
receptor-T cell therapy for solid tumors. *Immuno-Oncol. Technol.* **19**, 100385 (2023).

7. Garfall, A. L. *et al.* T-cell phenotypes associated with effective CAR T-cell therapy in
postinduction vs relapsed multiple myeloma. *Blood Adv.* **3**, 2812–2815 (2019).

8. Turtle, C. J. *et al.* CD19 CAR–T cells of defined CD4+:CD8+ composition in adult B cell ALL
patients. *Journal of Clinical Investigation* **126**, 2123–2138 (2016).

9. Kadauke, S. *et al.* Risk-Adapted Preemptive Tocilizumab to Prevent Severe Cytokine Release
Syndrome After CTL019 for Pediatric B-Cell Acute Lymphoblastic Leukemia: A Prospective
Clinical Trial. *J. Clin. Oncol.* **39**, 920–930 (2021).

10. Seder, R. A., Darrah, P. A. & Roederer, M. T-cell quality in memory and protection:
implications for vaccine design. *Nat. Rev. Immunol.* **8**, 247–258 (2008).

11. Eyquem, J. *et al.* Targeting a CAR to the TRAC locus with CRISPR/Cas9 enhances tumour
rejection. *Nature* **543**, 113–117 (2017).

12. Rossig, C. *et al.* Vaccination to improve the persistence of CD19CAR gene-modified T cells
in relapsed pediatric acute lymphoblastic leukemia. **31**, 1087–1095 (2017).

13. Akahori, Y. *et al.* Antitumor activity of CAR-T cells targeting the intracellular oncoprotein WT1
can be enhanced by vaccination. *Blood* **132**, 1134–1145 (2018).

14. Ma, L. *et al.* Enhanced CAR–T cell activity against solid tumors by vaccine boosting through
the chimeric receptor. *Science* **365**, 162–168 (2019).

15. Ma, L. *et al.* Vaccine-boosted CAR T crosstalk with host immunity to reject tumors with antigen
heterogeneity. *Cell* **186**, 3148-3165.e20 (2023).

16. Boder, E. T. & Wittrup, K. D. Yeast surface display for screening combinatorial polypeptide
libraries. *Nat. Biotechnol.* **15**, 553–557 (1997).

17. Rafiq, S., Hackett, C. S. & Brentjens, R. J. Engineering strategies to overcome the current
roadblocks in CAR T cell therapy. *Nat Rev Clin Oncol* **17**, 147–167 (2020).

18. Maude, S. L. *et al.* Chimeric antigen receptor T cells for sustained remissions in leukemia.
New Engl J Medicine **371**, 1507–17 (2014).

19. Kadauke, S. *et al.* Risk-Adapted Preemptive Tocilizumab to Prevent Severe Cytokine Release
Syndrome After CTL019 for Pediatric B-Cell Acute Lymphoblastic Leukemia: A Prospective
Clinical Trial. *J Clin Oncol* JCO.20.02477 (2021) doi:10.1200/jco.20.02477.

20. Frey, N. V. *et al.* Optimizing Chimeric Antigen Receptor T-Cell Therapy for Adults With Acute
Lymphoblastic Leukemia. *J. Clin. Oncol. : Off. J. Am. Soc. Clin. Oncol.* **38**, 415–422 (2019).

21. J., S. S. *et al.* Chimeric Antigen Receptor T Cells in Refractory B-Cell Lymphomas. *N. Engl.*
J. Med. **377**, 2545–2554 (2017).

22. Maude, S. L. *et al.* Chimeric Antigen Receptor T Cells for Sustained Remissions in Leukemia.
New England Journal of Medicine **371**, 1507–1517 (2014).

23. Park, J. H. *et al.* Long-Term Follow-up of CD19 CAR Therapy in Acute Lymphoblastic
Leukemia. *N. Engl. J. Med.* **378**, 449–459 (2018).

24. Fischer, L. *et al.* Cellular dynamics following CAR T cell therapy are associated with response
and toxicity in relapsed/refractory myeloma. *Leukemia* 1–11 (2024) doi:10.1038/s41375-023-
02129-y.

25. Ho, W. Y., Cooke, M. P., Goodnow, C. C. & Davis, M. M. Resting and anergic B cells are
defective in CD28-dependent costimulation of naive CD4+ T cells. *J. Exp. Med.* **179**, 1539–1549
(1994).

26. Lim, T. S. *et al.* CD80 and CD86 Differentially Regulate Mechanical Interactions of T-Cells
with Antigen-Presenting Dendritic Cells and B-Cells. *PLoS ONE* **7**, e45185 (2012).

27. Witkowski, M. T. *et al.* Extensive Remodeling of the Immune Microenvironment in B Cell Acute
Lymphoblastic Leukemia. *Cancer Cell* **37**, 867-882.e12 (2020).

28. Klesmith, J. R., Wu, L., Lobb, R. R., Rennert, P. D. & Hackel, B. J. Fine Epitope Mapping of
the CD19 Extracellular Domain Promotes Design. *Biochemistry* **58**, 4869–4881 (2019).

29. Laurent, E. *et al.* Directed Evolution of Stabilized Monomeric CD19 for Monovalent CAR
Interaction Studies and Monitoring of CAR-T Cell Patients. *ACS Synth. Biol.* **10**, 1184–1198
(2021).

30. He, C. *et al.* CD19 CAR antigen engagement mechanisms and affinity tuning. *Sci. Immunol.*
8, eadf1426 (2023).

31. Li, N. *et al.* CAR T cells targeting tumor-associated exons of glypican 2 regress neuroblastoma
in mice. *Cell Rep. Med.* **2**, 100297 (2021).

32. Wellhausen, N. *et al.* Epitope base editing CD45 in hematopoietic cells enables universal
blood cancer immune therapy. *Sci. Transl. Med.* **15**, eadi1145 (2023).

33. Melnikov, S. *et al.* Molecular insights into protein synthesis with proline residues. *EMBO Rep.*
17, 1776–1784 (2016).

34. Foote, J. & Eisen, H. N. Breaking the affinity ceiling for antibodies and T cell receptors. *Proc.*
Natl. Acad. Sci. **97**, 10679–10681 (2000).

35. Jumper, J. *et al.* Highly accurate protein structure prediction with AlphaFold. *Nature* **596**, 583–
589 (2021).

36. Rohl, C. A., Strauss, C. E. M., Misura, K. M. S. & Baker, D. Protein Structure Prediction Using
Rosetta. *Methods Enzym.* **383**, 66–93 (2004).

37. Krissinel, E. & Henrick, K. Inference of Macromolecular Assemblies from Crystalline State. *J.*
Mol. Biol. **372**, 774–797 (2007).

38. Burdette, D. L. *et al.* STING is a direct innate immune sensor of cyclic-di-GMP. *Nature* **478**,
515–518 (2011).

39. Holland, E. M. *et al.* Efficacy of second CAR-T (CART2) infusion limited by poor CART
expansion and antigen modulation. *J. Immunother. Cancer* **10**, e004483 (2022).

40. Shultz, L. D., Brehm, M. A., Garcia-Martinez, J. V. & Greiner, D. L. Humanized mice for
immune system investigation: progress, promise and challenges. *Nat. Rev. Immunol.* **12**, 786–
798 (2012).

41. Gordon, K. S. *et al.* Screening for CD19-specific chimaeric antigen receptors with enhanced
signalling via a barcoded library of intracellular domains. *Nat Biomed Eng* **6**, 855–866 (2022).

42. Melenhorst, J. J. *et al.* Decade-long leukaemia remissions with persistence of CD4+ CAR T
cells. *Nature* **602**, 503–509 (2022).

43. Reinhard, K. *et al.* An RNA vaccine drives expansion and efficacy of claudin-CAR-T cells
against solid tumors. *Science* **367**, 446–453 (2020).

44. Gauthier, J. *et al.* Factors associated with outcomes after a second CD19-targeted CAR T-
cell infusion for refractory B-cell malignancies. *Blood* **137**, 323–335 (2021).

45. Myers, R. M. *et al.* Outcomes after Reinfusion of CD19-Specific Chimeric Antigen Receptor
(CAR)-Modified T Cells in Children and Young Adults with Relapsed/Refractory B-Cell Acute
Lymphoblastic Leukemia. *Blood* **138**, 474 (2021).

46. Wagner, D. L. *et al.* Immunogenicity of CAR T cells in cancer therapy. *Nat. Rev. Clin. Oncol.*
18, 379–393 (2021).

47. Faude, S. *et al.* Absolute lymphocyte count proliferation kinetics after CAR T-cell infusion
impact response and relapse. *Blood Adv.* **5**, 2128–2136 (2021).

48. Mackensen, A. *et al.* CLDN6-specific CAR-T cells plus amplifying RNA vaccine in relapsed or
refractory solid tumors: the phase 1 BNT211-01 trial. *Nat. Med.* **29**, 2844–2853 (2023).

776