1

Title:

Dissociable memory modulation mechanisms facilitate fear amnesia at different timescales

Yinmei Ni[1*], Ye Wang[1*], Zijian Zhu[2], Jingchu Hu[3], Daniela Schiller[4], Jian Li[1, 5]

[1]School of Psychological and Cognitive Sciences and Beijing Key Laboratory of Behavior and
Mental Health, Peking University, Beijing, China
[2]School of Psychology, Shaanxi Normal University, Xi'an, China
[3]Department of Anxiety Disorders, Shenzhen Kangning Hospital, Shenzhen Mental Health
Institute, Shenzhen, China.
[4]Departments of Psychiatry and Neuroscience, and Friedman Brain Institute, Icahn School of
Medicine at Mount Sinai, New York, NY, USA
[5]IDG/McGovern Institute for Brain Research, Peking University, Beijing, China

* Equal contribution
Correspondence: leekin@gmail.com

Keywords: fear memory, memory retrieval, reconsolidation, amnesia, cTBS, dlPFC

20
21
22
23

Abstract
Memory reactivation renders consolidated memory fragile and thereby opens the window for
memory updates, such as memory reconsolidation. However, whether memory retrieval facilitates
update mechanisms other than memory reconsolidation remains unclear. We tested this hypothesis
in three experiments with healthy human participants. First, we demonstrate that memory
retrieval-extinction protocol prevents the return of fear expression shortly after extinction training
and this short-term effect is memory reactivation dependent (Study 1, $N = 57$ adults). Furthermore,
across different timescales, the memory retrieval-extinction paradigm triggers distinct types of
fear amnesia in terms of cue-specificity and cognitive control dependence, suggesting that the
short-term fear amnesia might be caused by different mechanisms from the cue-specific amnesia
at a longer and separable timescale (Study 2, $N = 79$ adults). Finally, using continuous theta-burst
stimulation (Study 3, $N = 75$ adults), we directly manipulated brain activity in the dorsolateral
prefrontal cortex, and found that both memory reactivation and intact prefrontal cortex function
were necessary for the short-term fear amnesia after the retrieval-extinction protocol. The
differences in temporal scale, cue-specificity, and cognitive control ability dependence between
the short- and long-term amnesia suggest that memory retrieval and extinction training trigger
distinct underlying memory update mechanisms. These findings suggest the potential involvement
of coordinated memory modulation processes upon memory retrieval and may inform clinical
approaches for addressing persistent maladaptive memories.

Introduction

Memory retrieval provides a unique window through which consolidated memories can be modified or even neutralized. Studies of memory reconsolidation suggest that memories are rendered fragile each time they are retrieved *(Alberini, 2005; Dudai, 2006; Nader, Schafe, & LeDoux, 2000)*, possibly due to the disruption of the old memory trace. The reconsolidation of the new memory trace also requires *de novo* protein synthesis, which usually takes hours to complete. Pharmacological blockade of protein synthesis and behavioral interventions can both diminish the original fear memory expression in the long-term (24 hours later) memory test *(Lee, 2008; Lee et al., 2017; Schiller et al., 2013; Schiller et al., 2010)*, resulting in the cue-specific fear memory deficit *(Debiec et al., 2002; Lee, 2008; Nader, Schafe, & LeDoux, 2000)*. For example, during the reconsolidation window, retrieving a fear memory allows it to be updated through extinction training (i.e., the retrieval-extinction paradigm *(Agren, 2014; Agren et al., 2012; Lee, 2008; Lee et al., 2017; Schiller et al., 2013; Schiller et al., 2010)*, but also see *(Chalkia, Schroyens, et al., 2020; Chalkia, Van Oudenhove, & Beckers, 2020; Schiller, 2020)*.

However, the exact functional role of memory retrieval remains unclear. For example, while previous fear memory studies typically reported the reconsolidation effects emerging several hours later (the long-term effect) *(Kindt and Soeter, 2018; Lee, 2008; Speer et al., 2021)*, the temporal dynamics of the amnesia effect through behavioral manipulation such as retrieval-extinction is often overlooked in the fear memory literature *(Lee et al., 2017; Schiller et al., 2013; Schiller et al., 2010)*. It is possible that memory retrieval might facilitate memory modification at different temporal scales. This is in stark contrast to semantic and episodic memory studies where the behavioral manipulations such as the retrieval-induced forgetting (RIF) protocol or direct suppression result in short-term memory deficit within 30 minutes *(Anderson et al., 2000; Anderson et al., 1994)*. More recently, behavioral strategies originally developed in the declarative memory studies such as direct suppression had also been successfully applied to the fear memory paradigms in humans and prevented the return of fear shortly afterward (30mins), suggesting that the fear memory might be amenable to modulation and this effect can manifest at a more immediate timescale, to which the memory reconsolidation theory remain agnostic *(Wang et al., 2021)*.

Corresponding to the long-term behavioral manifestation, concurrent neural evidence supporting memory reconsolidation hypothesis emphasizes that synapse degradation and *de novo* protein synthesis are required for reconsolidation. These processes typically take several hours, if not longer, and are often aided by overnight sleep *(Kindt and Soeter, 2018)*. On the contrary, previous behavioral manipulations engendering the short-term amnesia on declarative memory, such as the think/no-think paradigm, hinges on the intact activities in brain areas such as dorsolateral prefrontal cortex (cognitive control) and its functional coupling with specific brain regions such as hippocampus (memory retrieval) *(Anderson and Green, 2001; Wimber et al., 2015)*. The declarative amnesia effect arises much earlier due to the more instant modulation of functional connectivity, rather than the slower processes of new protein synthesis in these brain regions *(Wang et al., 2021)*. However, it remains unclear whether experimental paradigms such as memory retrieval-extinction, a method developed to induce long-term prevention of fear expression based on the memory reconsolidation hypothesis, might also trigger short-term amnesia for the fear

memory.

To fully comprehend the temporal dynamics of the memory retrieval-extinction training effect, we
first carried out a simple fear memory study by pairing a neutral cue (conditioned stimulus, CS)
with the electric shock (unconditioned stimulus, US) to directly test whether the fear
retrieval-extinction paradigm would yield a short-term effect on fear expression. We then
performed another double cue CS-US fear memory experiment to study the temporal
characteristics of fear memory malleability to behavioral extinction at the short- (30mins),
medium- (6 hours) and long-term (24 hours) time scales.

We hypothesize that the labile state triggered by the memory retrieval may facilitate different
memory deficits following extinction training, and these deficits can be further disentangled
through the lens of temporal dynamics and cue-specificities. In theory, different cognitive
mechanisms underlying specific fear memory deficits, therefore, can be inferred based on the
difference between memory deficits. Specifically, memory reconsolidation effect, if exists, should
only be evident in the long-term (24h) memory test due to the requirement of new protein
synthesis and should be cue-dependent *(Monfils et al., 2009)*. In contrast, if more immediate
memory update mechanisms are involved, then the fear memory deficit should be observed
relatively early (30mins) after extinction training *(Anderson and Floresco, 2022; Anderson and*
Green, 2001). There may even exist a "limbo" state: when the immediate memory update effects
dissipate and the reconsolidation updating is yet to take effect (6h), the conditioned stimuli should
still elicit fear responses, regardless of the reminder (cue). Previous research in the short-term
declarative and fear memory deficits highlighted the role of cognitive control. Subjects with
stronger control over intrusive thoughts or stronger functional connectivity between brain regions
such as the dorsolateral prefrontal cortex (dlPFC) and hippocampus exhibited more pronounced
amnesia in the memory suppression tasks *(Anderson et al., 1994; Wells and Davies, 1994; Wimber*
et al., 2015). We reason that if the retrieval related short-term fear amnesia is also related to
cognitive control and the dlPFC activities, then the individual difference of such immediate fear
memory deficits should be associated with individual subject's control abilities over intrusive
thoughts *(Küpper et al., 2014)*, which can be measured via the Thought-Control Ability
Questionnaire (TCAQ) scale.

Indeed, through a series of experiments, we identified a short-term fear amnesia effect following
memory retrieval-extinction training, in addition to the long-term amnesia effect that appeared
much later. Finally, by applying the continuous theta-burst stimulation (cTBS) protocol over the
dlPFC, the brain region critical for cognitive control and goal maintenance, we showed that both
memory retrieval and intact dlPFC activity were necessary for the more immediate fear amnesia
effect.

Open Practices Statement
Neither of the studies reported in this article was preregistered. The data and analysis codes for all
studies are publicly accessible at https://osf.io/9agvk/.

Methods

Participants

All participants were students recruited from Peking University. They were right-handed with normal vision and had not participated in electric shock-related experiments before. None of the participants had current or history of psychiatric illness, nor any contraindication on the application of transcranial magnetic stimulation (TMS) *(Rossi et al., 2009; Rossini et al., 2015)*. All participants provided informed consent and were remunerated for their participation. This study was approved by the institutional review board of school of psychological and cognitive sciences at Peking University.

We first conducted a power analysis (G*Power) to determine the number of participants sufficient to detect a reliable effect *(Faul et al., 2009)*. Based on the reported effect size of the reinstated fear memory between the treatment and control groups on fear memories reported in the previous literature ($\eta^2 = 0.19, 0.216, 0.24$; N = 40, 55, 30 of sample size, respectively *(Kindt and Soeter, 2018; Kindt et al., 2009; Wang et al., 2021)*), 52 participants for both groups (study 1: reminder group and no-reminder group) were enough to detect a significant interaction effect ($\alpha = 0.05$, $\beta = 0.9$, 2 (groups) $\times$ 2 (phases) two-way ANOVA interaction effect). Similarly, a total of 72 participants for three groups (study 2: 30-min group, 6-h group and 24-h group) were required to detect a reliable effect ($\alpha = 0.05$, $\beta = 0.9$, 3 (groups) $\times$ 2 (phases) $\times$ 2 (CS+) three-way ANOVA interaction effect) *(Heo and Leon, 2010)*. For the rTMS study, based on the reported effect size of reinstated fear memory ($\eta^2 = 0.144, 0.19, 0.216, 0.24$; N =79, 40, 55, 30 total sample size, respectively *(Kindt and Soeter, 2018; Kindt et al., 2009; Wang et al., 2021)*), a total of 72 participants (study 3) were needed to detect a significant interaction effect ($\alpha = 0.05$, $\beta = 0.8$, 4 (groups) $\times$ 2 (phases) $\times$ 2 (CS+) three-way ANOVA interaction effect).

In study 1, we first recruited a total of 75 human subjects (41 females; mean age= 21.3, SD= 2.21). Eight participants discontinued after Day 1 due to their non-measurable spontaneous conditioned stimulus (CS) signal (the skin conductance response, SCR) towards different CSs (non-responders, SCR response to any CS was less than 0.02 μS, n = 0 & 8; reminder group and no-reminder group, respectively). Ten participants (7 females; mean age = 20.6, SD = 1.79) who were "non-learners" during fear acquisition (n = 3 & 1 in both groups; respectively) or extinction (n = 3 & 3, respectively) phase were also excluded from further analysis. The criteria for "non-learners" in the fear acquisition include smaller mean SCR of CS+ (CS that was partially paired with the electric shock) than that of the CS- (CS that was never paired with shock) in the trials in the latter half of acquisition, and a smaller mean SCR difference between CS+ and CS- in the latter than the first half trials of fear acquisition on day 1. Similarly, the criteria for "non-learners" in the fear extinction are larger mean SCRs of CS+ than the CS- responses in both the last trial and the latter-half trials of extinction and a larger mean SCR difference between CS+ and CS- in the latter than the first half trials during fear extinction on day 2 (see the exclusion criteria Table S1 in the Supplemental Material). Therefore, we had a total of 57 subjects for data analysis in study 1 (30 for the reminder group, 17 females, mean age = 21.4, SD = 2.37, and 27 for the no-reminder group, 15 females, mean age = 21.62, SD = 2.13). The group $\times$ gender interaction effect on participants' age was not statistically significant ($F_{1,49} = 0.137$, $P = 0.713$, $\eta^2 = 0.003$, two-way ANOVA), and gender ($\chi_1^2 = 0.007$, $P = 0.933$) and age ($t_{55} = -0.376$, $P = 0.708$) were not significantly different between two groups.

In study 2, we first recruited 119 participants (56 females; mean age= 20.85, SD= 2.59).
Thirty-seven participants discontinued after Day 1 for they were "non-responders" (n = 19, 5 and
13; 30-min group, 6-h group and 24-h group, respectively). Additionally, two participants (n =
1&1; 30-min group and 24-h group; respectively) failed to show the evidence of fear acquisition
and one participant (30-min group) failed to show the evidence of fear extinction on day 2 and
were also excluded from further analysis. Finally, we had a total of 79 subjects for data analysis in
study 2 (27 for the 30-min group, 15 females, mean age = 21.4, SD = 2.46; 26 for the 6-h group,
12 females, mean age = 20.54, SD = 1.74; and 26 for the 24-h group, 12 females, mean age =
20.96, SD = 2.990). The interaction effect of gender × group on participants' age was not
statistically significant ($F_{2,73} = 0.213$, $P = 0.809$, $\eta^2 = 0.006$, two-way ANOVA), and gender ($\chi_2^2 =$
0.021, $P = 0.990$) or age ($F_{2,76} = 0.661$, $P = 0.519$, $\eta^2 = 0.017$) was not significantly different
across three groups.

In study 3, we recruited a total of 90 participants (47 females; mean age = 21.01, SD = 2.44).
Fifteen participants (8 females; mean age = 21.2, SD = 1.22) were excluded from further analysis
since they failed to show SCR response to any stimulus (CS1+, CS2+ or CS-) (non-responders, n
= 3,4,4,4; R rdlPFC group, R vertex group, NR rdlPFC group and NR vertex group, respectively,
see Supplemental Table S1) *(de Voogd and Phelps, 2020; Dunsmoor et al., 2015; Dunsmoor et al.,*
2017; Hu et al., 2018; Raio et al., 2017; Schiller et al., 2013; Schiller et al., 2010; Schiller et al.,
2012). Our final sample included 75 participants for data analysis (19 for the Reminder right
dlPFC group, 10 females, mean age = 22.05, SD = 2.44; 18 for the Reminder vertex group, 9
females, mean age = 20.67, SD = 2.68; 18 for the No-reminder right dlPFC group, 10 females,
mean age = 20.28, SD = 2.85; and 20 for the No-reminder vertex group, 10 females, mean age =
20.85, SD = 2.43).

Conditioned stimuli and psychophysiological stimulation
For study 1, we employed two squares with different colors (yellow and blue) that served as CS+
and CS-. During fear acquisition, the CS+ was paired with a mild electrical shock (US) on a 37.5%
partial reinforcement scheme (6 CS+ paired with electric shock and 10 CS+ with no shock), and
the CS- was never paired with a shock (10 CS-). A pseudo-random CS delivery order was
generated for the fear acquisition, extinction and test phase for two groups with the rule that no
same trial-type (CS+ and CS-) repeated more than twice. Assignment of square colors to the CS
(CS+ and CS-) was counterbalanced across participants.

For study 2 & 3, three squares with different colors (yellow, red and blue) were used as CS1+,
CS2+ and CS-, respectively. During the fear acquisition phase, both CS1+ and CS2+ were paired
with electric shocks (US) on a 37.5% partial reinforcement scheme, and the CS- was not paired
with shocks (10 non-reinforced presentations of CS1+, CS2+ and CS- each, intermixed with an
extra of 6 CS1+ and 6 CS2+ that co-terminated with the shock). Again, a pseudo-random stimulus
order was generated for fear acquisition and extinction phases of three groups with the rule that no
same trial-type (CS1+, CS2+ and CS-) repeated more than twice. In the test phase, to exclude the
possibility that the difference between CS1+ and CS2+ was simply caused by the presentation

sequence of CS1+ and CS2+, half of the participants completed the test phase using a
pseudo-random stimuli sequence and the identities of CS1+ and CS2+ reversed in the other half of
the participants. For all three groups, the CS colors were counterbalanced across participants.
224
225   The US was a mild electrical shock delivered to the right wrist of participants via a DS-5 Isolated
226   Bipolar Constant Current Stimulator (Digitimer, Welwyn Garden City, UK). The US levels were
227   set by the participants themselves, starting with a low shock level (5v) and gradually increased and
228   settled to a level that they described as 'uncomfortable, but not painful' (with the maximum level
229   of 10v). The duration of all electric shocks was 200ms with 50 pulses per second.
230
SCR measurements were collected using two Ag-AgCI electrodes attached to the tips of the index
and middle fingers of each subject's left hand. All skin conductance data were recorded via the
Biopac® MP160 BioNomadix System and analyzed using the Acknowledgement 5.0 software.
The SCR level for each trial was calculated as the amplitude difference (in micro siemens) from
peak to trough during the 0.5 to 4.5s window after the CS (CS SCR) or US (US SCR) onset and
responses below $0.02\mu S$ were encoded as zero. The raw SCR scores were divided by each
subject's mean US responses and then square-root-transformed to normalize SCR across
individuals and groups *(Wang et al., 2021)*.
239
Continuous theta burst stimulation (cTBS), a specific form of repetitive transcranial magnetic
stimulation (rTMS), was applied with a Magstim Rapid 2 system (Magstim, Whitland, Wales,
UK). Before the experiments, we determined the stimulation intensity of the cTBS protocol that
was at 80% of the individual active motor threshold (AMT). Each transcranial magnetic
stimulation (TBS) burst consisted of three stimuli pulses at 50HZ, with each train being repeated
every 200ms (5Hz). In this cTBS protocol, a 40s train of uninterrupted TBS is given (600 pluses)
(Borgomaneri et al., 2020a; Su et al., 2022). During the stimulation, the cTBS was applied either
over the right dlPFC or the vertex, which was determined by standard F4 and Cz locations based
on international electroencephalogram 10-20 system measurements *(Censor et al., 2010)*.
249
The control ability over intrusive thoughts was measured by the 25-item Thought-Control Ability
Questionnaire (TCAQ) scale *(Wells and Davies, 1994)*. Participants were asked to rate on a
five-point Likert-type scale the extent to which they agreed with the statement from 1 (*completely*
disagree) to 5 (*completely agree*). At the end of the experiments, all participants completed the
TCAQ scale to assess their perceived control abilities over intrusive thoughts in daily life *(Küpper*
et al., 2014). The internal consistency (Cronbach's alpha) of TCAQ scale was 0.92, and the
reliability coefficient was satisfactory ($r = 0.88$).
257
258   To measure the extent to which fear returns after the presentation of unconditioned stimuli (US,
259   electric shock) in the test phase, we defined the fear recovery index as the SCR difference between
260   the first test trial and the last extinction trial for a specific CS for each subject. Similarly, in studies
261   2 and 3, differential fear recovery index was defined as the difference between fear recovery
262   indices of CS+ and CS- for both CS1+ and CS2+.
263

Experimental procedure

Study 1, 2 and 3 were carried out for two or three consecutive days with three phases. Before the experiments, all participants gave informed consent. During the experiments, subjects were required to stay relaxed and still, focus on the computer screen and pay attention to the relationship between color stimuli (CS) and the electric shock (US).

Day 1: Acquisition. For study 1, two groups of participants underwent a fear-conditioning paradigm (acquisition) where the CS+ was partially (37.5%) paired with the US and the CS- was never paired with the shock. For study 2 & 3, participants underwent fear conditioning using three color squares as the CSs (CS1+, CS2+ and CS-). The CSs were presented for 4s each with a 10-12s variable ITI in all the studies (1, 2 and 3).

Day 2: Retrieval-extinction phase. For study 1, the fear memory was reactivated before extinction in the reminder group. During reactivation, the CS+ was presented once without the delivery of the US. Followed by a 10-min break, the reminder group underwent extinction training with 10 CS+ and 11 CS- presentation non-reinforced. In the no-reminder group, the fear memory was not reactivated (no CS+ reminder). To match with the timeline of the reminder group, the no-reminder group subjects still needed to wait for 10-min after the beginning of the experiment on day 2 and then underwent extinction training with 11 CS+ and 11 CS+ non-reinforced presentation. During the 10-min break, all participants were asked to rest and stay still.

For study 2, only one of the CS+ (CS1+) was presented once without the US during reactivation in all three groups. After a 10-min break, all three groups underwent extinction training with 10 CS1+, 11CS2+ and 11CS- non-reinforced presentation.

For study 3, for the reminder groups (reminder-dlPFC and reminder-vertex groups), only one of the CS+ (CS1+) was presented once (without the US delivery) as the retrieval trial. No CS+ reminder was delivered for the no-reminder groups (no-reminder dlPFC and no-reminder vertex groups). Additionally, during the 10-min break (8 mins after the supposed CS+ reminder delivery), cTBS was applied either over the right dlPFC (dlPFC groups: PFC) or the vertex (vertex groups: VER), resulting in a 2 (reminder vs. no-reminder) x 2 (dlPFC vs. vertex) of 4 different groups (reminder-dlPFC: R-PFC, reminder-vertex: R-VER, no-reminder dlPFC: NR-PFC, no-reminder vertex: NR-VER).

Day 2 or Day 3: Reinstatement and test phases. For study 1, 30 minutes after the extinction training, both groups received four un-signaled electric shocks with 10s to 12s ITI (reinstatement). Eighteen seconds after the reinstatement phase, all subjects were presented with CS+ and CS- 10 times each without electric shocks and their SCRs were recorded (test phase).

For study 2, participants were assigned into three groups: the 30min group, the 6h group and the 24h group. All participants received four un-signaled electric shocks with 10s to 12s ITI (reinstatement) after their corresponding time delays (30min, 6h or 24h) between extinction and reinstatement. Eighteen seconds after the reinstatement phase, all subjects were tested via the presentation of CS1+, CS2+ and CS- 10 times each without electric shocks associated.

For study 3, the reinstatement and testing procedures were the same as in study 2 except that they
were conducted one hour after the extinction training.

Results
The short-term effect of the fear memory retrieval-extinction procedure
To test the memory-retrieval-induced fear memory deficits at different time scales in humans, we
designed two experiments to examine whether the fear memory retrieval-extinction procedure
would block the return of fear memory.

In the first study, we aimed to test whether there is a short-term amnesia effect following the fear
retrieval-extinction paradigm. We measured subjects' SCRs for the fear acquisition, extinction and
test phases to assess the associative fear memory and the recovery of fear memory. At each phase,
the differential fear response was calculated as the difference between SCRs to the CS+ and CS-.

To assess fear acquisition across groups (Figure 1B and C), we conducted a mixed two-way
ANOVA of group (reminder vs. no-reminder) x time (early vs. late part of the acquisition; first 5
and last 5 trials, correspondingly) on the differential fear SCR. Our results showed a significant
main effect of time (early vs. late; $F_{1,55} = 6.545$, $P = 0.013$, $\eta^2 = 0.106$), suggesting successful fear
acquisition in both groups. There was no main effect of group (reminder vs. no-reminder) or the
group x time interaction (group: $F_{1,55} = 0.057$, $P = 0.813$, $\eta^2 = 0.001$; interaction: $F_{1,55} = 0.066$, $P =$
0.798, $\eta^2 = 0.001$), indicating similar levels of fear acquisition between two groups. Post-hoc
t-tests confirmed that the fear responses to the CS+ were significantly higher than that of CS-
during the late part of acquisition phase in both groups (reminder group: $t_{29} = 6.642$, $P < 0.001$;
no-reminder group: $t_{26} = 8.522$, $P < 0.001$; Figure 1C). Importantly, the levels of acquisition were
equivalent in both groups (early acquisition: $t_{55} = -0.063$, $P = 0.950$; late acquisition: $t_{55} = -0.318$,
$P = 0.751$; Figure 1C).

To ensure equivalent and successful extinction across groups (Figure 1B and D), we performed
similar mixed two-way ANOVA (group x trial) on the differential SCR in the extinction phase.
Our results showed significant main effect of trial ($F_{10,550} = 2.825$, $P = 0.010$, $\eta^2 = 0.049$), but no
effect of group ($F_{1,55} = 0.063$, $P = 0.802$, $\eta^2 = 0.001$) or their interaction ($F_{10,550} = 1.77$, $P = 0.101$,
$\eta^2 = 0.031$). Follow-up t-tests further confirmed that in the last trial of extinction there was no
difference between fear responses to CS+ and CS- within group (reminder group: $t_{29} = -1.147$, $P =$
0.261; no-reminder group: $t_{26} = 0.261$, $P = 0.796$; Figure 1D). Importantly, the levels of extinction
did not show significant difference between two groups ($t_{55} = -1.074$, $P = 0.288$; Figure 1D).

Fear recovery was assessed using a two-way ANOVA with main effects of group (reminder vs.
no-reminder) and time (last trial of extinction vs. first trial of test) and it showed a significant time
× group interaction ($F_{1,55} = 4.087$, $P = 0.048$, $\eta^2 = 0.069$). We defined the fear recovery index as
the SCR difference between the first test trial and the last extinction trial for a specific CS. The
interaction effect was confirmed by the significant difference between the differential fear
recovery indexes between CS1+ and CS2+ in the reminder and no-reminder groups ($t_{55} = -2.022$,
$P = 0.048$; Figure 1E, also see Supplemental Material for the direct test of the test phase). Post hoc

t-tests showed that fear memories were resilient after regular extinction training, as demonstrated
by the significant difference between fear recovery indexes of the CS+ and CS- for the
no-reminder group ($t_{26} = 7.441$, $P < 0.001$; Bonferroni correction, Figure 1E), while subjects in the
reminder group showed no difference of fear recovery between CS+ and CS- ($t_{29} = 0.797$, $P =$
0.432, Figure 1E). Also, our results remain robust even with the "non-learners" included in the
analysis (Figure supplement 1 in the Supplemental Material).

Together, these results indicate that the retrieval-extinction procedure prevents the return of fear
expression in the short-term test, a phenomenon distinct from the hypothesized fear
reconsolidation effect at a much later time *(Schiller et al., 2013)*. In the current study, fear
recovery was tested 30 minutes after extinction training, whereas the effect of memory
reconsolidation was generally evident only several hours later and possibly with the help of sleep
(Kindt and Soeter, 2018; Speer et al., 2021), leaving open the possibility of a different cognitive
mechanism for the short-term fear amnesia related to the retrieval-extinction procedure.
Importantly, such a short-term effect is also retrieval dependent, suggesting the labile state of
memory may be necessary for the short-term memory update to take effect (Figure 1E).

Temporal and generalizing characteristics of the retrieval-extinction effect
Given the findings of short-term fear amnesia following the retrieval-extinction paradigm, we set
out to map the temporal dynamics of fear amnesia as well as its cue-specificities - whether the
amnesia triggered by a specific CS+ reminder (for example CS1+) generalizes to the fear memory
associated with the other CS+ (CS2+). We adopted a double-cue fear learning paradigm and
examined fear recovery in the short- (30 mins), medium- (6 hours) and long-term (24 hours) fear
memory tests in three separate groups of participants after the extinction training (Figure 2A).

Fear acquisition was assessed using a mixed three-way ANOVA with group (30min, 6h and 24h),
CS+ (CS1+ vs. CS2+, defined as the mean SCR differences between each CS+ and CS-) and trial
numbers (10 trials) as main factors. This analysis showed a significant main effect of trial ($F_{9,684} =$
12.782, $P < 0.001$, $\eta^2 = 0.144$), but no effect of group ($F_{2,76} = 2.121$, $P = 0.127$, $\eta^2 = 0.053$), CS+
($F_{1,76} = 0.576$, $P = 0.45$, $\eta^2 = 0.008$) or their interactions (all $Ps > 0.05$; Figure 2B). Furthermore, to
ensure equivalent fear acquisition across three groups, a mixed two-way ANOVA showed no
effect of group (30min, 6h and 24h; $F_{2,76} = 1.79$, $P = 0.174$, $\eta^2 = 0.045$), CS+ (CS1+ vs. CS2+;
$F_{1,76} = 1.337$, $P = 0.251$, $\eta^2 = 0.017$) or their interaction ($F_{2,76} = 0.959$, $P = 0.388$, $\eta^2 = 0.025$) on
the late phase of acquisition (last 5 trials). Post-hoc *t*-tests confirmed that the SCRs elicited by
CS1+ and CS2+ were significantly higher than that of CS- among three groups on the late phase
of acquisition (CS1+: $t_{26} = 6.806$, $P < 0.001$; CS2+: $t_{26} = 6.854$, $P < 0.001$ for the 30min group;
CS1+: $t_{25} = 9.822$, $P < 0.001$; CS2+: $t_{25} = 9.841$, $P < 0.001$ for the 6h group and CS1+: $t_{25} = 6.909$,
$P < 0.001$; CS2+: $t_{25} = 6.956$, $P < 0.001$ for the 24h group), indicating successful and similar fear
acquisition across all groups (Figure 2C).

We then set out to examine whether participants underwent successful extinction training on day 2.
The similar mixed three-way ANOVA showed a significant effect of trial ($F_{10,760} = 8.220$, $P <$
0.001, $\eta^2 = 0.098$), but no effect of group ($F_{2,76} = 1.822$, $P = 0.169$, $\eta^2 = 0.046$), CS+ ($F_{1,76} = 0.91$,
$P = 0.343$, $\eta^2 = 0.012$) or their interactions (all $Ps > 0.1$; Figure 2B). To test whether participants

across all groups achieved similar levels of extinction, a mixed two-way ANOVA showed no
effect of group (30min, 6h and 24h; $F_{2,76} = 0.059$, $P = 0.943$, $\eta^2 = 0.002$), CS+ (CS1+ vs. CS2+;
$F_{1,76} = 0.258$, $P = 0.613$, $\eta^2 = 0.003$) or their interaction ($F_{2,76} = 0.504$, $P = 0.606$, $\eta^2 = 0.013$) on
the last trial of extinction training. Post-hoc t-tests showed that there was no difference between
CS+ and CS- responses in all groups on the last trial of extinction (CS1+: $t_{26} = -0.397$, $P = 0.695$;
CS2+: $t_{26} = -0.505$, $P = 0.618$ for the 30min group; CS1+: $t_{25} = -0.054$, $P = 0.957$; CS2+: $t_{25} = $
0.140, $P = 0.890$ for the 6h group and CS1+: $t_{25} = 0.435$, $P = 0.668$; CS2+: $t_{25} = -0.317$, $P = 0.754$
for the 24h group; Figure 2D).
404
To assess the effects of the retrieval-extinction procedure as a function of the time delay between
fear extinction and test phases, we conducted a 3-way ANOVA with the within-subject factor CS+
(CS1+ vs. CS2+, defined by the mean SCR differences between each CS+ and CS-), time (last
trial of extinction vs. first trial of test) and between-subject factor group (30min, 6h and 24h) and
found a significant three-way interaction ($F_{2,76} = 6.376$, $P = 0.003$, $\eta^2 = 0.144$). To disentangle this
interaction, we further examined the effects in each group.
411
In the 30min group, there was no significant effect of time ($F_{1,26} = 0.208$, $P = 0.652$, $\eta^2 = 0.008$),
CS+ ($F_{1,26} = 0.888$, $P = 0.355$, $\eta^2 = 0.033$) or their interaction ($F_{1,26} = 1.039$, $P = 0.318$, $\eta^2 = 0.038$).
Post-hoc t-tests showed that the retrieval-extinction training diminished fear responses to both the
reminded CS+ (differential fear recovery index between CS1+ and CS-, $t_{26} = -0.787$, $P = 0.438$;
Figure 2C) and the non-reminded CS+ (differential fear recovery index between CS2+ and CS-, t_{26}
$= 0.088$, $P = 0.93$; Figure 2E), suggesting a cue-independent short-term amnesia effect. This result
indicates that the short-term amnesia effect observed in Study 2 is not reminder-cue specific and
can generalize to the non-reminded cues. In addition, there was no significant difference between
the fear recovery index associated with CS1+ and CS2+ ($t_{26} = -1.019$, $P = 0.318$; Figure 2E, also
see the supplemental material for direct test of the test phase SCRs).
422
In contrast to the short-term effect, there was a significant effect of time ($F_{1,25} = 9.654$, $P = 0.005$,
$\eta^2 = 0.279$), but no effect of CS+ ($F_{1,25} = 0.037$, $P = 0.85$, $\eta^2 = 0.001$) or their interaction ($F_{1,25} = $
0.028, $P = 0.869$, $\eta^2 = 0.001$) in the 6h (medium-term) group. Post-hoc t-tests showed that fear
memory recovered for the reminded CS+ (differential fear recovery index between CS1+ and CS-,
$t_{25} = 2.382$, $P = 0.025$), as well as the non-reminded CS+ (differential fear recovery index between
CS2+ and CS-, $t_{25} = 3.525$, $P = 0.002$) in the medium-term test of the retrieval-extinction
procedure effect, suggesting the failure to suppress the return of extinguished fear memory at such
a time scale (Figure 2E). There was also no significant difference measured by the fear recovery
index between CS1+ and CS2+ ($t_{25} = 0.166$, $P = 0.870$; Figure 2E).
432
Finally, in line with previous research on fear memory reconsolidation, significant SCR difference
of the fear reinstatement effect between the reminded CS+ and non-reminded CS+ (time × CS+
interaction: $F_{1,25} = 16.924$, $P < 0.001$, $\eta^2 = 0.404$) was detected in the 24h group via two-way
ANOVA. More specifically, the retrieval-extinction procedure only diminished fear responses to
the reminded CS+ (differential fear recovery index between CS1+ and CS-, $t_{25} = -0.25$, $P = 0.805$),
whereas the fear response to the non-reminded CS+ remained significant (differential fear
recovery index between CS2+ and CS-, $t_{25} = 3.269$, $P = 0.003$; Figure 2E). Post-hoc t-test showed

that the fear recovery index between CS1+ and CS2+ was significantly different (t_{25} = -4.114, $P <$
0.001; Figure 2E), aligning well with previous literature that suggested the cue-specificity of the
fear reconsolidation effect *(Lee et al., 2017; Monfils et al., 2009)*. Similarly, these results remain
robust even with the "non-learners" included in the analysis (Figure supplement 2 in the
Supplemental Material).

Altogether, in line with our hypothesis, the return of fear memory measured by the reinstatement
effect showed distinct patterns at different time scales. The retrieval-extinction procedure
diminished fear SCRs for both the reminded and non-reminded CS+, engendering
cue-independent amnesia in the short-term test (30min). On the other hand, at the long-term (24h)
fear memory test, fear responses were only diminished for the reminded CS+ (but not the
non-reminded CS+), consistent with previous literature on memory reconsolidation. Interestingly,
SCR measures of the fear memory re-appeared in the medium-term (6h) test and there was no
difference between fear responses elicited by the reminded and non-reminded CS+, suggesting
that the 6-hour timescale may be at a "limbo" stage where both the immediate and the delayed
(reconsolidation) memory update mechanisms are not at work.

Fear amnesia as a function of the thought-control ability
Finally, to assess whether the fear reinstatement effects at different time scale were related to
subjects' thought-control ability, we first ran a one-way ANOVA to confirm that subjects among
different groups did not differ in their thought-control abilities measured by the TCAQ scores
($F_{2,76}$ = 0.154, P = 0.857, η^2 = 0.004). We then ran separate linear regressions between the
differential fear recovery index (differential fear recovery index between CS+ and CS-) and
thought-control abilities (TCAQ scores) in the three groups. This analysis yielded significant
negative correlation between TCAQ score and fear recovery index for both CS+ (CS1+ and CS2+)
in the 30min group (Figure 3A, main effect of thought-control ability, t_{51} = -3.446, P = 0.003,
Bonferroni correction), indicating that subjects endowed with higher thought-control abilities were
less likely to experience the return of fear response. However, no significant correlation was
observed in the 6h or the 24h group (Figure 3B and C; Ps > 0.4). Post-hoc analysis of the 30min
group showed significant correlation between the fear recovery index of both CS1+ and CS2+
with subjects' thought-control abilities (Figure 3A, CS1+: r = -0.461, P = 0.015; CS2+: r = -0.413,
P = 0.032). No significant TCAQ and CS+ interactions were found in the three groups (Ps > 0.4).

It is worth noting that correlations between TCAQ scores and fear responses in all three groups
were not significant in the late phase of acquisition or the last trial of extinction (all Ps > 0.10),
indicating that the effect of thought-control ability was specifically reflected in the fear recovery
index.

In conclusion, thought-control abilities were associated with fear recovery indices only in the
short-term (30min) test for both the reminded and non-reminded CS+ but not at longer time scales
(6h and 24h). The 30min group results are consistent with previous research which suggested that
people with better capability to resist intrusive thoughts also performed better in motivated
forgetting in both declarative and associative memories *(Küpper et al., 2014; Wang et al., 2021)*.

dlPFC dependent short-term fear memory amnesia

In study 3 (Figure 4A), we examined whether intact activity of the right dlPFC (rdlPFC) was also

necessary for short-term fear amnesia, in addition to the memory retrieval cue (reminder). To test

this hypothesis, we set up a reminder CS+ (CS1+ vs. CS2+) x reminder (reminder vs. no-reminder)

x cTBS (rdlPFC vs. vertex) x trial number four-way ANOVA against the differential SCRs (CS+

minus CS-) for the acquisition and extinction phases. In the acquisition phase, as expected, there

were no significant main effects of reminder ($F_{1,71} = 1.291$, $P = 0.260$, $\eta^2 = 0.018$), cTBS ($F_{1,71} <$

0.001, $P = 0.990$, $\eta^2 < 0.001$), CS+ ($F_{1,71} = 1.927$, $P = 0.169$, $\eta^2 = 0.026$) or their interactions (all

$Ps > 0.1$; Figure 4B and C) but a significant main effect of trial number ($F_{9,639} = 8.054$, $P < 0.001$,

$\eta^2 = 0.102$), indicating successful learning of CS+ stimuli. Indeed, post-hoc tests confirmed that

the SCRs of CS+ was significantly higher than that of the CS- on the late phase (last 5 trials) of

acquisition in all four groups (CS1+: $t_{18} = 7.735$, $P < 0.001$; CS2+: $t_{18} = 8.546$, $P < 0.001$ for the

R-PFC group; CS1+: $t_{17} = 8.568$, $P < 0.001$; CS2+: $t_{17} = 8.478$, $P < 0.001$ for the R-VER group;

CS1+: $t_{17} = 6.032$, $P < 0.001$; CS2+: $t_{17} = 7.722$, $P < 0.001$ for the NR-PFC group and CS1+: $t_{19} =$

9.110, $P < 0.001$; CS2+: $t_{19} = 6.538$, $P < 0.001$ for the NR-VER group, Figure 4C) .

A similar mixed-effect four-way ANOVA was also performed on the SCRs in the extinction phase,

which showed significant effects of trial ($F_{10,710} = 8.567$, $P < 0.001$, $\eta^2 = 0.108$), cTBS ($F_{1,71} =$

6.269, $P = 0.015$, $\eta^2 = 0.081$) and cTBS $\times$ trial interaction ($F_{10,710} = 2.292$, $P = 0.030$, $\eta^2 = 0.031$),

but no effect of reminder ($F_{1,71} = 1.894$, $P = 0.173$, $\eta^2 = 0.026$), CS+ ($F_{1,71} = 0.398$, $P = 0.530$, $\eta^2 =$

0.006) or other interactions (all $Ps > 0.1$; Figure 4B). To test whether participants across all groups

achieved similar levels of extinction, a mixed-effect three-way ANOVA was performed on the last

trial of extinction training and there was no significant effect of reminder (reminder and

no-reminder; $F_{1,71} = 3.112$, $P = 0.082$, $\eta^2 = 0.042$), cTBS ($F_{1,71} = 0.929$, $P = 0.338$, $\eta^2 = 0.013$),

CS+ (CS1+ vs. CS2+; $F_{1,71} = 1.623$, $P = 0.207$, $\eta^2 = 0.022$) or their interactions (all $Ps > 0.1$;

Figure 4D). Furthermore, post-hoc t-tests showed that there was no difference between CS+ and

CS- responses in all groups on the last trial of extinction (CS1+: $t_{18} = 1.310$, $P = 0.207$; CS2+: $t_{18} =$

1.298, $P = 0.211$ for the R-PFC group; CS1+: $t_{17} = 0.839$, $P = 0.413$; CS2+: $t_{17} = 1.090$, $P = 0.291$

for the R-VER group; CS1+: $t_{17} = 0.380$, $P = 0.709$; CS2+: $t_{17} = 0.624$, $P = 0.541$ for the NR-PFC

group and CS1+: $t_{19} = -1.434$, $P = 0.168$; CS2+: $t_{19} = -0.535$, $P = 0.599$ for the NR-VER group;

Figure 4D).

To assess the effects of the right dlPFC on the memory retrieval related short-term amnesia, we

conducted a mixed-effect 3-way ANOVA with the within-subject factors CS+ (CS1+ vs. CS2+),

between-subject factors reminder (reminder vs. no-reminder) and cTBS (dlPFC vs. vertex) on fear

recovery index (the difference between the first test trial and the last extinction trial for CS1+ and

CS2+), which revealed a significant effect of reminder ($F_{1,71} = 8.920$, $P = 0.004$, $\eta^2 = 0.112$), cTBS

($F_{1,71} = 7.180$, $P = 0.009$, $\eta^2 = 0.092$) and reminder $\times$ cTBS interaction ($F_{1,71} = 10.613$, $P = 0.002$,

$\eta^2 = 0.130$), but no effect of CS+ ($F_{1,71} = 0.518$, $P = 0.474$, $\eta^2 = 0.007$) or other interactions (all $Ps >$

0.1; Figure 4C). Post-hoc t-tests showed that there were significant fear reinstatement responses in

the R-PFC group (CS1+: $t_{18} = 5.182$, $P < 0.001$; CS2+: $t_{18} = 5.258$, $P < 0.001$), the NR-PFC group

(CS1+: $t_{17} = 3.496$, $P = 0.003$; CS2+: $t_{17} = 4.628$, $P < 0.001$) and the NR-VER group (CS1+: $t_{19} =$

4.885, $P < 0.001$; CS2+: $t_{19} = 3.262$, $P = 0.004$), but not in the R-VER group (CS1+: $t_{17} = -1.640$,

$P = 0.119$; CS2+: $t_{17} = -0.744$, $P = 0.467$) (Figure 4E), suggesting that both memory retrieval and

intact dlPFC function are necessary for the short-term fear amnesia.

Finally, we went on to examine whether the fear reinstatement response was related to the
thought-control ability. A two-way ANOVA of reminder (reminder vs. no-reminder) x cTBS
(rdlPFC vs. vertex) confirmed that there were no significant effects of reminder ($F_{1,71} = 0.366$, $P =$
0.547, $\eta^2 = 0.005$), cTBS ($F_{1,71} = 0.606$, $P = 0.439$, $\eta^2 = 0.008$) or their interaction ($F_{1,71} = 1.216$, P
$= 0.274$, $\eta^2 = 0.017$) on the TCAQ scores. We then conducted simple linear regressions on
different groups (as in study 2) and found a significant correlation between the thought-control
ability and differential fear recovery index in the R-VER group (Figure 5B, $t_{33} = -3.283$, $P = 0.008$,
Bonferroni correction across 4 groups), consistent with the results of the short term (30min)
amnesia effect in study 2 (Figure 3A). Further post-hoc analysis of the individual CS+ showed
that the correlations between thought-control ability and fear recovery index were significant for
both CS+ (CS1+: $r = -0.557$, $P = 0.016$; CS2+: $r = -0.483$, $P = 0.042$). However, there was no
such correlation in the R-PFC group (Figure 5A, $t_{35} = 0.719$, $P = 0.477$), the NR-PFC group
(Figure 5C, $t_{33} = -1.334$, $P = 0.192$) or the NR-VER group (Figure 5D, $t_{37} = -0.025$, $P = 0.980$) and
the interaction effects of thought-control ability and CS+ (CS1+ and CS2+) were not significant in
all 4 groups ($Ps > 0.3$). Importantly, there was no significant correlation between thought-control
ability and SCR in the late phase of acquisition or the last trial of extinction in four groups (all $Ps >$
0.10), suggesting a specific relationship between thought-control ability and short-term fear
recovery.

Discussion
Our results provide direct evidence to support the hypothesis that the memory reactivation might
engage distinct cognitive mechanisms for fear memory modulation, which can be separated by
their temporal dynamics, cue specificity and the dependence of thought-control ability. In line
with previous research on fear memory reconsolidation *(Liu et al., 2014)*, our research replicated
the reported results that the retrieval-extinction paradigm blocked fear responses to the reminded
CS+ while leaving non-reminded CS+ response intact (cue-specificity) in the long-term test
(Figure 6A and B). However, thought-control ability was not related to the long-term fear amnesia
(Figure 6C). Therefore, findings of the short-term fear amnesia suggest that the reconsolidation
framework falls short to accommodate this more immediate effect (Figure 6A and B).

Research in both declarative and associative emotional memories suggest that memory retrieval is
not a simple act of obtaining a copy of stored information, but provides a critical time window
where the old memory is eligible to be integrated with new information *(Gershman, Schapiro, et*
al., 2013; Hupbach et al., 2007; Monfils et al., 2009). Research in the field of declarative memory
has shown that the motivated forgetting effect can appear early (10-30 minutes) after the memory
suppression practices such as the influential think/no-think (TNT) paradigm *(Anderson and Green,*
2001), suggesting that memory retrieval may also facilitate short-term memory update *(Benoit and*
Anderson, 2012; Engen and Anderson, 2018). Unlike the slow protein synthesis process involved
in memory reconsolidation, the motivated forgetting effect was accompanied by heightened
dorsolateral prefrontal cortex (dlPFC) activity, diminished hippocampal activation and altered
functional connectivity between these brain regions and hence more adaptive *(Anderson et al.,*
2004; Wimber et al., 2015).

Previous studies indicate that a suppression mechanism can be characterized by three distinct
features: first, the memory suppression effect tends to emerge early, usually 10-30 mins after
memory suppression practice and can be transient *(MacLeod and Macrae, 2001; Saunders and*
MacLeod, 2002); second, the memory suppression practice seems to directly act upon the
unwanted memory itself *(Levy and Anderson, 2002)*, such that the presentation of other cues
originally associated with the unwanted memory also fails in memory recall (cue-independence);
third, the magnitude of memory suppression effects is associated with individual difference in
control abilities over intrusive thoughts *(Küpper et al., 2014)*. The short-term fear amnesia effects
we identified are fundamentally different from memory reconsolidation effects and fit closely with
the characteristics of the memory suppression effect. Inspired by the similarities between our
results and suppression-induced declarative memory amnesia *(Gagnepain et al., 2017)*, we
speculate that the retrieval-extinction procedure might facilitate a spontaneous memory
suppression process and thus yield a short-term amnesia effect. Accordingly, the activated fear
memory induced by the retrieval cue would be subjected to an automatic fear memory suppression
through the extinction training *(Anderson and Floresco, 2022)*.

In our experiments, subjects were not explicitly instructed to suppress their fear expression, yet the
retrieval-extinction training significantly decreased short-term fear expression. These results are
consistent with the short-term amnesia induced with the more explicit suppression intervention
(Anderson et al., 1994; Kindt and Soeter, 2018; Speer et al., 2021; Wang et al., 2021; Wells and
Davies, 1994). It is worth noting that although consciously repelling unwanted memory is a
standard approach in memory suppression paradigm, it is possible that the engagement of the
suppression mechanism can be unconscious. For example, in the retrieval-induced forgetting (RIF)
paradigm, recall of a stored memory impairs the retention of related target memory and this
forgetting effect emerges as early as 20 minutes after the retrieval procedure, suggesting memory
suppression or inhibition can occur in a more spontaneous and automatic manner *(Imai et al.,*
2014). Moreover, subjects with trauma histories exhibited more suppression-induced forgetting for
both negative and neutral memories than those with little or no trauma *(Hulbert and Anderson,*
2018). Similarly, people with higher self-reported thought-control capabilities showed more severe
cue-independent memory recall deficit, suggesting that suppression mechanism is associated with
individual differences in spontaneous control abilities over intrusive thoughts *(Küpper et al.,*
2014). It has also been suggested that similar automatic mechanisms might be involved in organic
retrograde amnesia of traumatic childhood memories *(Schacter et al., 2012; Schacter et al., 1996)*.

Therefore, it is theoretically possible that similar brain mechanisms, such as the dynamic brain
activity and connectivity modulation, involved in memory suppression may be deployed
spontaneously and act as a general cognitive mechanism to cope with intrusive memories in daily
life. Indeed, recently researchers have proposed that the standard retrieval-extinction paradigm to
study fear memory might also engage automatic memory suppression *(Anderson and Floresco,*
2022). Consistent with this hypothesis, we showed that the retrieval cue was necessary to produce
the short-term amnesia effect (Figure 1) and critically, our results fitted well with the three key
characteristics of the active suppression mechanism: temporal dynamics, cue-specificity and the
dependence of thought-control ability (Figures 2 and 3).

To calibrate the temporal boundaries of the short and long-term effects, we also included a
separate group (medium-term group) where fear expression was tested 6 hours after the
retrieval-extinction training. Interestingly, fear returned at this time point (Figure 6A and B),
confirming the 6-hour lower time boundary for the hypothesized memory reconsolidation in
previous animal and human studies *(Agren, 2014; Agren et al., 2012; Debiec et al., 2002; Kindt*
and Soeter, 2018; Lee, 2008; Nader, Schafe, & Le Doux, 2000; Speer et al., 2021). However, the
return of fear expression in the medium-term group also indicates that the short-term fear amnesia
is transient, laying out the upper time boundary for the hypothesized fear memory suppression
effect. Although mixed results have been reported regarding the durability of suppression effects
in the declarative memory studies *(Meier et al., 2011; Storm et al., 2012)*, future research will be
needed to investigate whether the short-term effect we observed is specifically related to
associative memory or the spontaneous nature of suppression as in RIF (Figure 6C). Previous
research also showed that separate mnemonic processes can be involved after emotional memory
retrieval. For example, by varying the duration length and repetition number of memory cue
re-exposure, different levels of memory persistence (reconsolidation, extinction or a "limbo" state
in between) can be induced, suggesting a neural titration mechanism for memory de-stabilization
and re-stabilization *(Faliagkas et al., 2018; Merlo et al., 2014)*. These works identified important
"boundary conditions" of memory retrieval in affecting the retention of the maladaptive emotional
memories. In our study, however, we showed that even within a boundary condition previously
thought to elicit memory reconsolidation, mnemonic processes other than reconsolidation could
also be at work, and these processes jointly shape the persistence of fear memory.

Our results also corroborate with a recent study showing that blocking fear reconsolidation using
post-retrieval repetitive transcranial magnetic stimulation (rTMS) did not influence the short-term
fear memory expression *(Borgomaneri et al., 2020b)*. However, as we demonstrated in studies 2 &
3, memory retrieval and intact rdlPFC function were both necessary for the short-term fear
amnesia after the extinction training. Therefore, instead of "intact short-term fear expression by
the rTMS over the dlPFC", their results could also be interpreted such that the rTMS on the dlPFC
demolished the otherwise short-term fear amnesia induced by the memory retrieval cue. Indeed, a
series of neuroimaging studies have revealed that the prefrontal neural network including anterior
cingulate cortex, anterior ventrolateral prefrontal cortex and the dlPFC is directly engaged in
purging the unwanted declarative memory from consciousness *(Anderson et al., 2004; Benoit and*
Anderson, 2012; Levy and Anderson, 2002). Moreover, previous research has established a causal
relationship between dlPFC activity and the memory suppression mechanism underlying
retrieval-induced forgetting using transcranial direct current stimulation (tDCS) *(Penolazzi et al.,*
2014; Stramaccia et al., 2017; Valle et al., 2020). In our experiment (studies 2 and 3), when the
dlPFC function was intact, subjects' thought-control abilities were positively correlated with fear
amnesia in the test phase. Such correlations disappeared in the no-reminder groups (NR-VER and
NR-PFC) and the R-PFC group (Figures 3 and 5), establishing a causal link between the dlPFC
activity and short-term fear amnesia.

It should be noted that while our long-term amnesia results were consistent with the fear memory
reconsolidation literatures, there were also studies that failed to observe fear prevention *(Chalkia,*

Schroyens, et al., 2020; Chalkia, Van Oudenhove, & Beckers, 2020; Schroyens et al., 2023). Although the memory reconsolidation framework provides a viable explanation for the long-term amnesia, more evidence is required to validate the presence of reconsolidation, especially at the neurobiological level *(Elsey et al., 2018)*. While it is beyond the scope of the current study to discuss the discrepancies between these studies, one possibility to reconcile these results concerns the procedure for the retrieval-extinction training. It has been shown that the eligibility for old memory to be updated is contingent on whether the old memory and new observations can be inferred to have been generated by the same latent cause *(Gershman et al., 2017; Gershman and Niv, 2012)*. For example, prevention of the return of fear memory can be achieved through gradual extinction paradigm, which is thought to reduce the size of prediction errors to inhibit the formation of new latent causes *(Gershman, Jones, et al., 2013)*. Therefore, the effectiveness of the retrieval-extinction paradigm might depend on the reliability of such paradigm in inferring the same underlying latent cause. Furthermore, other studies highlighted the importance of memory storage *per se* and suggested that memory retention was encoded in the memory engram cell ensemble connectivity whereas the engram cell synaptic plasticity is crucial for memory retrieval *(Ryan et al., 2015; Tonegawa, Liu, et al., 2015; Tonegawa, Pignatelli, et al., 2015)*. It remains to be tested how the cue-independent short-term and cue-dependent long-term amnesia effects we observed could correspond to the engram cell synaptic plasticity and functional connectivity among engram cell ensembles (Figure 6). This is particularly important, since the cue-independent characteristic of the short-term amnesia suggest that either different memory cues fail to evoke engram cell activities, or the retrieval-extinction training transiently inhibits connectivity among engram cell ensembles. Finally, SCR is only one aspect of the fear expression, how the retrieval-extinction paradigm might affect subjects' other emotional (such as the startle response) and cognitive fear expressions such as reported fear expectancy needs to be tested in future studies since they do not always align with each other *(Kindt et al., 2009; Sevenster et al., 2012, 2013)*.

These results help shed light on the dynamics of memory modulation after memory activation and designing novel treatment of psychiatric disorders caused by excessive fear or anxiety *(Chen et al., 2021)*. Memory reconsolidation and suppression both have been studied thoroughly in separate fields. Specifically, aversive memories are usually associated with different retrieval cues, and interfering memory reconsolidation only blocks fear memory recall from the retrieval memory probe but not the others *(Debiec et al., 2002; Kindt and Soeter, 2018; Lee, 2008; Nader, Schafe, & Le Doux, 2000; Speer et al., 2021)*. On the other hand, memory suppression is cue-independent and directly suppresses the fear memory trace yet only effective with limited durability *(MacLeod and Macrae, 2001)*. Although future research is clearly needed to understand the brain mechanisms for the short-term amnesia and its potential connection to the memory suppression effect, especially in relevance to clinical populations such as posttraumatic stress disorder (PTSD) and phobia patients *(Homan et al., 2019; Stramaccia et al., 2021)*, our results provide a general framework to highlight the potential of memory retrieval to triggering different memory updating mechanisms with unique temporal dynamics.

Transparency: This work was supported by the National Science and Technology Innovation
2030 Major Program (2021ZD0203702), National Natural Science Foundation of China Grants
(32071090, 31871140) to J.L.

The authors have declared no conflict of interests.

Data availability

All the data of this study are available for download at https://osf.io/9agvk/.

Figure 1 | SCR responses to conditioned stimuli in two groups of study 1 indicating short-term
memory amnesia (**A**) Experimental design and timeline. (**B**) SCRs of fear conditioned stimuli
(CS+) and the control stimulus (CS-) across fear acquisition, extinction and test phases for two
groups (reminder and no-reminder). (**C**) Mean differential SCRs (CS+ minus CS-) in the
acquisition phase (trials in the latter half). (**D**) Differential SCRs (CS+ minus CS-) in the
extinction phase (last trial). (**E**) Differential fear recovery index between CS+ and CS- in the test
phase. ***$P < 0.001$, *$P < 0.05$. NS: Non-significant. Error bars represent standard errors.

Figure 2 | SCR responses to conditioned stimuli in three groups of study 2 indicating amnesia at
different timescales. (**A**) Experimental design and timeline. (**B**) SCRs of fear conditioned stimuli
CS1+ (reminder) and CS2+ (No-reminder), and the control stimulus (CS-) across the fear
acquisition, extinction and test phases for each group (30min, 6h and 24h groups). (**C**) Mean
differential SCRs (CS1+ minus CS- and CS2+ minus CS-) in the acquisition phase (trials in the
latter half). (**D**) Differential SCRs in the extinction phase (last trial). (**E**) Differential fear recovery
index between CS+ and CS- in the test phase. ***$P < 0.001$. **$P < 0.01$. *$P < 0.05$. NS:
Non-significant. Error bars represent standard errors.

Figure 3 | Fear recovery as a function of thought-control abilities. (**A**)Thought control ability was significantly correlated with fear recovery index in the 30min group ($P = 0.003$, Bonferroni correction), but not in the 6h (**B**) or 24h group (**C**, Ps > 0.7). The violin graphs indicate the distribution of fear recovery index across subjects (Figure 2E). $**P < 0.01$. NS: Non-significant.

738
Figure 4 | SCR responses to conditioned stimuli in the cTBS study (study 3) indicating that both
intact dlPFC and cue reminder are required for short-term memory amnesia. (**A**) Experimental
design and timeline. (**B**) SCRs of fear conditioned stimuli CS1+ (reminder) and CS2+
(No-reminder), and the control stimulus (CS-) across the fear acquisition, extinction and test
phases for each group (R-PFC, R-VER, NR-PFC and NR-VER groups). (**C**) Mean differential
SCRs (CS1+ minus CS- and CS2+ minus CS-) in the acquisition phase (trials in the latter half). (**D**)
Differential SCRs in the extinction phase (last trial). (**E**) Differential fear recovery index between

CS+ and CS- in the test phase. ***$P < 0.001$. *$P < 0.05$. NS: Non-significant. Error bars represent

standard errors.

Figure 5 | The correlation between differential fear recovery index and thought-control ability in
four cTBS groups. (**A**) There was no significant correlation between differential fear recovery
index and thought-control abilities in the R-PFC group (*P* > 0.4). (**B**) However, in the R-VER
group, the correlation between thought control ability scores and the differential fear recovery
index was significant (*P* = 0.008, Bonferroni correction), with high thought-control ability
participants showing less fear recovery for both CSs+. Such correlation was not observed in the
(**C**) NR-PFC or the (**D**) NR-VER group (*Ps* >0.4). ** *P* < 0.01, NS: Non-significant.
756

Figure 6 | Time courses of short- and long-term amnesia. (**A**) At the short interval (30min to 1h),

fear recovery of the reminded CS (CS1+) is inhibited (green). As time progresses (from 6h to 24h),

amnesia effect is likely due to the fear memory reconsolidation effect emerged later (orange).

Actual SCR data in black solid line. (**B**) Fear amnesia of the non-reminded CS (CS2+) is only

evident at the 30-min interval and such effect starts to decay as test interval increases (green).

However, long-term amnesia does not generalize to CS2+ (orange, cue-dependence). The

observed SCR data (black solid line) paralleled the prediction from both the short-term and the

long-term effects as the interval length increased (from 30min to 24h). (**C**) Schematics depicting

the effect of cue-reminder on fear memory retention. After both CS1+ (black) and CS2+ (grey)

successfully elicit fear responses in the acquisition phase, CS1+ is reminded (black) before both

CS+ go through the extinction training. The lack of both CS1+ and CS2+ fear responses in the

short-term memory (STM) test (30 min) might be explained by the dlPFC dependent direct

suppression effect (dotted circle of US representation). On the other hand, the cue-specific fear

amnesia effect in the long-term memory (LTM) test (24h) of CS1+ but not CS2+ could be

attributed to the reconsolidation effect specific to CS1+. * $P < 0.05$, Error bars represent standard

errors.

786
Reference

Agren, T. 2014. Human reconsolidation: a reactivation and update. *Brain Res Bull* **105**: 70-82.

https://doi.org/10.1016/j.brainresbull.2013.12.010

Agren, T., Engman, J., Frick, A., Björkstrand, J., Larsson, E. M., Furmark, T., & Fredrikson, M.

2012. Disruption of reconsolidation erases a fear memory trace in the human

amygdala. *Science* **337**: 1550-1552. https://doi.org/10.1126/science.1223006

Alberini, C. M. 2005. Mechanisms of memory stabilization: are consolidation and

reconsolidation similar or distinct processes? *Trends Neurosci* **28**: 51-56.

https://doi.org/10.1016/j.tins.2004.11.001

Anderson, M. C., Bjork, E. L., & Bjork, R. A. 2000. Retrieval-induced forgetting: evidence for a

recall-specific mechanism. *Psychon Bull Rev* **7**: 522-530.

https://doi.org/10.3758/bf03214366

Anderson, M. C., Bjork, R. A., & Bjork, E. L. 1994. Remembering can cause forgetting:

retrieval dynamics in long-term memory. *Journal of experimental psychology.*

Learning, memory, and cognition **20**: 1063-1087.

https://doi.org/10.1037//0278-7393.20.5.1063

Anderson, M. C., & Floresco, S. B. 2022. Prefrontal-hippocampal interactions supporting the

extinction of emotional memories: the retrieval stopping model.

Neuropsychopharmacology **47**: 180-195. https://doi.org/10.1038/s41386-021-01131-1

Anderson, M. C., & Green, C. 2001. Suppressing unwanted memories by executive control.

Nature **410**: 366-369. https://doi.org/10.1038/35066572

Anderson, M. C., Ochsner, K. N., Kuhl, B., Cooper, J., Robertson, E., Gabrieli, S. W., Glover,

G. H., & Gabrieli, J. D. 2004. Neural systems underlying the suppression of unwanted

memories. *Science* **303**: 232-235. https://doi.org/10.1126/science.1089504

Benoit, R. G., & Anderson, M. C. 2012. Opposing mechanisms support the voluntary forgetting

of   unwanted   memories.   *Neuron*   **76**:   450-460.

https://doi.org/10.1016/j.neuron.2012.07.025

Borgomaneri, S., Battaglia, S., Garofalo, S., Tortora, F., Avenanti, A., & di Pellegrino, G. 2020a.

State-Dependent TMS over Prefrontal Cortex Disrupts Fear-Memory Reconsolidation

and   Prevents   the   Return   of   Fear.   *Curr*   *Biol*   **30**:   3672-3679.e3674.

https://doi.org/10.1016/j.cub.2020.06.091

Borgomaneri, S., Battaglia, S., Garofalo, S., Tortora, F., Avenanti, A., & di Pellegrino, G. 2020b.

State-Dependent TMS over Prefrontal Cortex Disrupts Fear-Memory Reconsolidation

and   Prevents   the   Return   of   Fear.   *Curr*   *Biol*   **30**:   3672-3679   e3674.

https://doi.org/10.1016/j.cub.2020.06.091

Censor, N., Dimyan, M. A., & Cohen, L. G. 2010. Modification of existing human motor

memories is enabled by primary cortical processing during memory reactivation. *Curr*

Biol **20**: 1545-1549. https://doi.org/10.1016/j.cub.2010.07.047

Chalkia, A., Schroyens, N., Leng, L., Vanhasbroeck, N., Zenses, A.-K., Van Oudenhove, L., &

Beckers, T. 2020. No persistent attenuation of fear memories in humans: A registered

replication   of   the   reactivation-extinction   effect.   *Cortex*   **129**:   496-509.

https://doi.org/https://doi.org/10.1016/j.cortex.2020.04.017

Chalkia, A., Van Oudenhove, L., & Beckers, T. 2020. Preventing the return of fear in humans

using reconsolidation update mechanisms: A verification report of Schiller et al. (2010).

Cortex **129**: 510-525. https://doi.org/https://doi.org/10.1016/j.cortex.2020.03.031

Chen, W., Li, J., Zhang, X., Dong, Y., Shi, P., Luo, P., & Zheng, X. 2021. Retrieval-extinction

as a reconsolidation-based treatment for emotional disorders:Evidence from an

extinction retention test shortly after intervention. *Behav Res Ther* **139**: 103831.

https://doi.org/10.1016/j.brat.2021.103831

de Voogd, L. D., & Phelps, E. A. 2020. A cognitively demanding working-memory intervention

enhances extinction Research article. *Scientific report* **10**: 7020.

https://doi.org/10.1038/s41598-020-63811-0

Debiec, J., LeDoux, J. E., & Nader, K. 2002. Cellular and systems reconsolidation in the

hippocampus. *Neuron* **36**: 527-538. https://doi.org/10.1016/s0896-6273(02)01001-2

Dudai, Y. 2006. Reconsolidation: the advantage of being refocused. *Curr Opin Neurobiol* **16**:

174-178. https://doi.org/10.1016/j.conb.2006.03.010

Dunsmoor, J. E., Campese, V. D., Ceceli, A. O., LeDoux, J. E., & Phelps, E. A. 2015.

Novelty-facilitated extinction: providing a novel outcome in place of an expected threat

diminishes recovery of defensive responses. *Biological psychiatry* **78**: 203-209.

https://doi.org/10.1016/j.biopsych.2014.12.008

Dunsmoor, J. E., Otto, A. R., & Phelps, E. A. 2017. Stress promotes generalization of older but

not recent threat memories. *Proceedings of the National Academy of Sciences of the*

United States of America **114**: 9218-9223. https://doi.org/10.1073/pnas.1704428114

Elsey, J. W. B., Van Ast, V. A., & Kindt, M. 2018. Human memory reconsolidation: A guiding

framework and critical review of the evidence. *Psychological bulletin* **144**: 797-848.

https://doi.org/10.1037/bul0000152

Engen, H. G., & Anderson, M. C. 2018. Memory Control: A Fundamental Mechanism of

Emotion Regulation. *Trends in Cognitive Sciences* **22**: 982-995.

https://doi.org/10.1016/j.tics.2018.07.015

Faliagkas, L., Rao-Ruiz, P., & Kindt, M. 2018. Emotional memory expression is misleading:

delineating transitions between memory processes. *Current Opinion in Behavioral*

Sciences **19**: 116-122. https://doi.org/https://doi.org/10.1016/j.cobeha.2017.12.018

Faul, F., Erdfelder, E., Buchner, A., & Lang, A.-G. 2009. Statistical power analyses using

G*Power 3.1: Tests for correlation and regression analyses. *Behavior Research*

Methods **41**: 1149-1160. https://doi.org/10.3758/BRM.41.4.1149

Gagnepain, P., Hulbert, J., & Anderson, M. C. 2017. Parallel Regulation of Memory and

Emotion Supports the Suppression of Intrusive Memories. *The Journal of*

neuroscience **37**: 6423-6441. https://doi.org/10.1523/jneurosci.2732-16.2017

Gershman, S. J., Jones, C. E., Norman, K. A., Monfils, M. H., & Niv, Y. 2013. Gradual

extinction prevents the return of fear: implications for the discovery of state. *Front*

Behav Neurosci **7**: 164. https://doi.org/10.3389/fnbeh.2013.00164

Gershman, S. J., Monfils, M. H., Norman, K. A., & Niv, Y. 2017. The computational nature of

memory modification. *Elife* **6**. https://doi.org/10.7554/eLife.23763

Gershman, S. J., & Niv, Y. 2012. Exploring a latent cause theory of classical conditioning.

Learn Behav **40**: 255-268. https://doi.org/10.3758/s13420-012-0080-8

Gershman, S. J., Schapiro, A. C., Hupbach, A., & Norman, K. A. 2013. Neural context

reinstatement predicts memory misattribution. *The Journal of neuroscience* **33**:

8590-8595. https://doi.org/10.1523/jneurosci.0096-13.2013

Heo, M., & Leon, A. C. 2010. Sample sizes required to detect two-way and three-way

interactions involving slope differences in mixed-effects linear models. *J Biopharm*

Stat **20**: 787-802. https://doi.org/10.1080/10543401003618819

Homan, P., Levy, I., Feltham, E., Gordon, C., Hu, J., Li, J., Pietrzak, R. H., Southwick, S.,

Krystal, J. H., Harpaz-Rotem, I., & Schiller, D. 2019. Neural computations of threat in

the aftermath of combat trauma. *Nature neuroscience* **22**: 470-476.

https://doi.org/10.1038/s41593-018-0315-x

Hu, J., Wang, W., Homan, P., Wang, P., Zheng, X., & Schiller, D. 2018. Reminder duration

determines threat memory modification in humans. *Scientific report* **8**: 8848.

https://doi.org/10.1038/s41598-018-27252-0

Hulbert, J. C., & Anderson, M. C. 2018. What doesn't kill you makes you stronger:

Psychological trauma and its relationship to enhanced memory control. *Journal of*

experimental psychology. General **147**: 1931-1949.

https://doi.org/10.1037/xge0000461

Hupbach, A., Gomez, R., Hardt, O., & Nadel, L. 2007. Reconsolidation of episodic memories:

a subtle reminder triggers integration of new information. *Learn Mem* **14**: 47-53.

https://doi.org/10.1101/lm.365707

Imai, H., Kim, D., Sasaki, Y., & Watanabe, T. 2014. Reward eliminates retrieval-induced

forgetting. *Proceedings of the National Academy of Sciences of the United States of*

America **111**: 17326-17329. https://doi.org/10.1073/pnas.1404469111

Kindt, M., & Soeter, M. 2018. Pharmacologically induced amnesia for learned fear is time and

sleep dependent. *Nature communications* **9**: 1316.

https://doi.org/10.1038/s41467-018-03659-1

Kindt, M., Soeter, M., & Vervliet, B. 2009. Beyond extinction: erasing human fear responses

and preventing the return of fear. *Nature neuroscience* **12**: 256-258.

https://doi.org/10.1038/nn.2271

Küpper, C. S., Benoit, R. G., Dalgleish, T., & Anderson, M. C. 2014. Direct suppression as a

mechanism for controlling unpleasant memories in daily life. *Journal of experimental*

psychology. General **143**: 1443-1449. https://doi.org/10.1037/a0036518

Lee, J. L. 2008. Memory reconsolidation mediates the strengthening of memories by additional

learning. *Nature neuroscience* **11**: 1264-1266. https://doi.org/10.1038/nn.2205

Lee, J. L. C., Nader, K., & Schiller, D. 2017. An Update on Memory Reconsolidation Updating.

Trends in Cognitive Sciences **21**: 531-545. https://doi.org/10.1016/j.tics.2017.04.006

Levy, B. J., & Anderson, M. C. 2002. Inhibitory processes and the control of memory retrieval.

Trends in Cognitive Sciences **6**: 299-305.

https://doi.org/10.1016/s1364-6613(02)01923-x

Liu, J., Zhao, L., Xue, Y., Shi, J., Suo, L., Luo, Y., Chai, B., Yang, C., Fang, Q., Zhang, Y., Bao,

Y., Pickens, C. L., & Lu, L. 2014. An unconditioned stimulus retrieval extinction

procedure to prevent the return of fear memory. *Biological psychiatry* **76**: 895-901.

https://doi.org/10.1016/j.biopsych.2014.03.027

MacLeod, M. D., & Macrae, C. N. 2001. Gone but not forgotten: the transient nature of

retrieval-induced forgetting. *Psychological Science* **12**: 148-152.

https://doi.org/10.1111/1467-9280.00325

Meier, B., König, A., Parak, S., & Henke, K. 2011. Suppressed, but not forgotten. *Swiss*

Journal of Psychology / Schweizerische Zeitschrift für Psychologie / Revue Suisse de

Psychologie **70**: 5-11. https://doi.org/10.1024/1421-0185/a000033

Merlo, E., Milton, A. L., Goozée, Z. Y., Theobald, D. E., & Everitt, B. J. 2014. Reconsolidation

and extinction are dissociable and mutually exclusive processes: behavioral and

molecular evidence. *The Journal of neuroscience* **34**: 2422-2431.

https://doi.org/10.1523/jneurosci.4001-13.2014

Monfils, M. H., Cowansage, K. K., Klann, E., & LeDoux, J. E. 2009. Extinction-reconsolidation

boundaries: key to persistent attenuation of fear memories. *Science* **324**: 951-955.

https://doi.org/10.1126/science.1167975

Nader, K., Schafe, G. E., & Le Doux, J. E. 2000. Fear memories require protein synthesis in

the amygdala for reconsolidation after retrieval. *Nature* **406**: 722-726.

https://doi.org/10.1038/35021052

Nader, K., Schafe, G. E., & LeDoux, J. E. 2000. Fear memories require protein synthesis in the

amygdala for reconsolidation after retrieval. *Nature* **406**: 722-726.

https://doi.org/10.1038/35021052

Penolazzi, B., Stramaccia, D. F., Braga, M., Mondini, S., & Galfano, G. 2014. Human memory

retrieval and inhibitory control in the brain: beyond correlational evidence. *The Journal*

of neuroscience **34**: 6606-6610. https://doi.org/10.1523/jneurosci.0349-14.2014

Raio, C. M., Hartley, C. A., Orederu, T. A., Li, J., & Phelps, E. A. 2017. Stress attenuates the

flexible updating of aversive value. *Proceedings of the National Academy of Sciences*

of the United States of America **114**: 11241-11246.

https://doi.org/10.1073/pnas.1702565114

Rossi, S., Hallett, M., Rossini, P. M., & Pascual-Leone, A. 2009. Safety, ethical considerations,

and application guidelines for the use of transcranial magnetic stimulation in clinical

practice and research. *Clin Neurophysiol* **120**: 2008-2039.

https://doi.org/10.1016/j.clinph.2009.08.016

Rossini, P. M., Burke, D., Chen, R., Cohen, L. G., Daskalakis, Z., Di Iorio, R., Di Lazzaro, V.,

Ferreri, F., Fitzgerald, P. B., George, M. S., Hallett, M., Lefaucheur, J. P., Langguth, B.,

Matsumoto, H., Miniussi, C., Nitsche, M. A., Pascual-Leone, A., Paulus, W., Rossi,

S., . . . Ziemann, U. 2015. Non-invasive electrical and magnetic stimulation of the brain,

spinal cord, roots and peripheral nerves: Basic principles and procedures for routine

clinical and research application. An updated report from an I.F.C.N. Committee. *Clin*

Neurophysiol **126**: 1071-1107. https://doi.org/10.1016/j.clinph.2015.02.001

Ryan, T. J., Roy, D. S., Pignatelli, M., Arons, A., & Tonegawa, S. 2015. Memory. Engram cells

retain memory under retrograde amnesia. *Science* **348**: 1007-1013.

https://doi.org/10.1126/science.aaa5542

Saunders, J., & MacLeod, M. D. 2002. New evidence on the suggestibility of memory: the role

of retrieval-induced forgetting in misinformation effects. *J Exp Psychol Appl* **8**:

127-142.

Schacter, D. L., Addis, D. R., Hassabis, D., Martin, V. C., Spreng, R. N., & Szpunar, K. K. 2012.

The future of memory: remembering, imagining, and the brain. *Neuron* **76**: 677-694.

https://doi.org/10.1016/j.neuron.2012.11.001

Schacter, D. L., Koutstaal, W., & Norman, K. A. 1996. Can cognitive neuroscience illuminate

the nature of traumatic childhood memories? *Curr Opin Neurobiol* **6**: 207-214.

https://doi.org/10.1016/s0959-4388(96)80074-2

Schiller, D., Kanen, J. W., LeDoux, J. E., Monfils, M. H., & Phelps, E. A. 2013. Extinction

during reconsolidation of threat memory diminishes prefrontal cortex involvement.

Proceedings of the National Academy of Sciences of the United States of America **110**:

20040-20045. https://doi.org/10.1073/pnas.1320322110

Schiller, D., LeDoux, J. E., & Phelps, E. A. . 2020. Reply to Beckers, McIntosh and Chambers

on the verification of 'preventing the return of fear using retrieval-extinction in humans'.

https://doi.org/10.31234/osf.io/jn6uw

Schiller, D., Monfils, M. H., Raio, C. M., Johnson, D. C., Ledoux, J. E., & Phelps, E. A. 2010.

Preventing the return of fear in humans using reconsolidation update mechanisms.

Nature **463**: 49-53. https://doi.org/10.1038/nature08637

Schiller, D., Raio, C. M., & Phelps, E. A. 2012. Extinction training during the reconsolidation

window prevents recovery of fear. *J Vis Exp*: e3893. https://doi.org/10.3791/3893

Schroyens, N., Beckers, T., & Luyten, L. 2023. Appraising reconsolidation theory and its

empirical   validation.   *Psychonomic   Bulletin   &   Review*   **30**:   450-463.

https://doi.org/10.3758/s13423-022-02173-2

Sevenster, D., Beckers, T., & Kindt, M. 2012. Retrieval per se is not sufficient to trigger

reconsolidation   of   human   fear   memory.   *Neurobiol   Learn   Mem*   **97**:   338-345.

https://doi.org/10.1016/j.nlm.2012.01.009

Sevenster, D., Beckers, T., & Kindt, M. 2013. Prediction error governs pharmacologically

induced   amnesia   for   learned   fear.   *Science*   **339**:   830-833.

https://doi.org/10.1126/science.1231357

Speer, M. E., Ibrahim, S., Schiller, D., & Delgado, M. R. 2021. Finding positive meaning in

memories of negative events adaptively updates memory. *Nature communications* **12**:

6601. https://doi.org/10.1038/s41467-021-26906-4

Storm, B., Bjork, E., & Bjork, R. 2012. On the durability of retrieval-induced forgetting. *Journal*

of Cognitive Psychology **24**: 617-629. https://doi.org/10.1080/20445911.2012.674030

Stramaccia, D. F., Meyer, A. K., Rischer, K. M., Fawcett, J. M., & Benoit, R. G. 2021. Memory

suppression and its deficiency in psychological disorders: A focused meta-analysis.

Journal   of   experimental   psychology.   General **150**:   828-850.

https://doi.org/10.1037/xge0000971

Stramaccia, D. F., Penolazzi, B., Altoè, G., & Galfano, G. 2017. TDCS over the right inferior

frontal gyrus disrupts control of interference in memory: A retrieval-induced forgetting

study. *Neurobiol Learn Mem* **144**: 114-130. https://doi.org/10.1016/j.nlm.2017.07.005

Su, S., Deng, J., Yuan, K., Gong, Y., Zhang, Y., Li, H., Cao, K., Huang, X., Lin, X., Wu, P., Xue,

Y., Bao, Y., Shi, J., Shi, L., & Lu, L. 2022. Continuous theta-burst stimulation over the

right dorsolateral prefrontal cortex disrupts fear memory reconsolidation in humans.

iScience **25**: 103614. https://doi.org/https://doi.org/10.1016/j.isci.2021.103614

Tonegawa, S., Liu, X., Ramirez, S., & Redondo, R. 2015. Memory Engram Cells Have Come

of Age. *Neuron* **87**: 918-931. https://doi.org/10.1016/j.neuron.2015.08.002

Tonegawa, S., Pignatelli, M., Roy, D. S., & Ryan, T. J. 2015. Memory engram storage and

retrieval. *Curr Opin Neurobiol* **35**: 101-109. https://doi.org/10.1016/j.conb.2015.07.009

Valle, T. M., Bajo, M. T., & Gómez-Ariza, C. J. 2020. Cathodal transcranial direct current

stimulation over the right dorsolateral prefrontal cortex cancels out the cost of

selective retrieval on subsequent analogical reasoning. *Neuropsychologia* **141**:

107431. https://doi.org/10.1016/j.neuropsychologia.2020.107431

Wang, Y., Zhu, Z., Hu, J., Schiller, D., & Li, J. 2021. Active suppression prevents the return of

threat memory in humans. *Commun Biol* **4**: 609.

https://doi.org/10.1038/s42003-021-02120-2

Wells, A., & Davies, M. I. 1994. The Thought Control Questionnaire: a measure of individual

differences in the control of unwanted thoughts. *Behav Res Ther* **32**: 871-878.

https://doi.org/10.1016/0005-7967(94)90168-6

Wimber, M., Alink, A., Charest, I., Kriegeskorte, N., & Anderson, M. C. 2015. Retrieval induces

adaptive forgetting of competing memories via cortical pattern suppression. *Nature*

neuroscience **18**: 582-589. https://doi.org/10.1038/nn.3973

1018