Cure or Curse? Simulation Indicates that Microbes Proliferate under

Disinfection Measures in the Space Station

Shuaishuai Li[1,2,5], Xiaolei Liu[1,2], Letian Chen[1,2], Hong Liu[1,2,3,4], Dawei Hu[1,2,3,4,]*

[1] School of Biological Science and Medical Engineering, Beihang University, Beijing 100083,

China

[2] Institute of Environmental Biology and Life Support Technology, Beihang University, Beijing

100083, China

[3] International Joint Research Center of Aerospace Biotechnology and Medical Engineering,

Beijing 100083, China

[4] State Key Laboratory of Virtual Reality Technology and Systems, School of Computer Science

and Engineering, Beihang University, Beijing 100083, China

[5] Department of Biology, McGill University, Montréal, Québec, Canada H3A 1B1

* Corresponding author.

E-mail address: dwhu@163.com (D. Hu)

Postal address: 37 Xueyuan Road, Haidian District, Beijing, China

Email addresses of other authors: Shuaishuai Li: angku@outlook.com, Xiaolei Liu:

liuxiaolei1998@126.com, Letian Chen: clt407360571@outlook.com, Hong Liu:

lh64@buaa.edu.cn

Abstract

The long-term stay on the space station requirement poses a knotty issue about

the stable hardware and reliable guarantee of astronauts' health. Despite regular

cleaning with wipes, microbes still proliferate in the space station. This study distilled

three key factors—low-dose ionizing radiation, dilution and quaternary ammonium

compound from the scenario. Species abundance and differential metabolites were

applied to test their sole/combined influences on the three-simple-species microbial

community succession. Mechanisms were built in mathematical models to generate

the structural and behavior similarity. Results showed that LDIR, dilution and QAC,

solely or jointly, could contribute to microbial proliferation. The synergy of

disturbances might convert them from harmful factors to rewarding ones (e.g. the

transformation of QAC into nutrients under LDIR), leading to this "Ecological

Surprise". These results shed light on mechanisms driving microbial community

succession in the space station and highlight the need for tailored biocontrol strategies

in the specific environment.

Keywords: microbial community succession, disturbance, low-dose ionizing

radiation, dilution, quaternary ammonium compound(s), synergy.

Abbreviation:

LDIR: low-dose ionizing radiation; QAC: quaternary ammonium compound(s)

1. Introduction

Long-term residence on the space station needs stable hardware and guaranteed

health status of astronauts (Brooks et al., 2014; Rosado et al., 2010; Sun et al., 2016;

Yang et al., 2020), which is threatened by proliferative microbes on the inner surface

of the space station (Alekhova et al., 2005). When environmental factors are the same,

the species diversity on the space station is much higher than that on the ground (Lang

et al., 2017; Novikova et al., 2006), showing the paradox of the plankton (G. E.

Hutchinson, 1961). As astronauts and goods are main sources of microbes in this

closed environment (Lax et al., 2014; Singh et al., 2018), periodic cleaning is an

actual plan for it. When carrying out the cleaning task, astronauts use wipes

containing 0.1% quaternary ammonium compounds (QAC) (Singh et al., 2018) to

wipe down the internal surface of the cabin under the low-dose ionizing radiation

(LDIR) from the universe. However, the removal of microbes seems to be only

effective for a short period of time and it is not complete. Microbes always grow back

soon after and the community structure varies by time and location (Lang et al., 2017;

Singh et al., 2018).

Understanding community succession mechanisms is always one of core targets

of ecology, especially microbial ecology(Powell et al., 2015; Stegen et al., 2013; Zhou

& Ning, 2017). Disturbance, as a largely independent area of ecology, is the event that

influences populations, resources or the physical environment in community

succession (Pickett & White, 1985). Removal by wiping, translated into the

perspective of microbial community succession, is caused by abiotic environmental

disturbance. This operation results in microscopic mass non-specific or specific death,

macroscopic physical dilution and chemical disinfection events. Community

succession occurs when organisms are disturbed due to their sensitivity to the abiotic

environment (Baudoin et al., 2008; Dangles & Chauvet, 2003; Tolkkinen et al., 2013).

Although violent environmental degradation might lead to species extinction, some

species could be favored under that circumstance because of the interactions

necessary for survival in multi-species communities (Abreu et al., 2019, 2020). For

example, evenly increasing mortality favors the faster grower, reversing the winner

from the slower to faster one in competition (Hastings, 2013; Stewart & Levin, 1973).

Nevertheless, current oversimplified models might lose some key information about

the succession. Research to date has not yet determined how disturbances would work

in resource-explicit models and the influence of selective disturbances.

In this study, three different types of disturbances, LDIR, dilution and QAC,

were distilled from the scene of wiping the inner surface of the space station.

Experiments with the simple three-species microbial community were carried out to

demonstrate species absolute abundance and differential metabolites in the

community succession under different combinations and gradients of these

disturbances. Results showed that trends of community succession varied among

different treatments and all types of disturbances could cause the proliferation by the

relief of competition or property transformation of specific chemicals. Afterward,

some measures were proposed for microbe control in aerospace. Also, it is hoped that

this research will contribute to a deeper understanding of community succession

formations and mechanisms under different types of disturbances.

2. Materials and methods

2.1 Strains, culture and radiative environment simulation

Three laboratory-stored strains, *Bacillus subtilis* (CMCC(B) 63501), *Escherichia*

coli (CMCC(B) 44102) and *Pseudomonas aeruginosa* (CMCC(B) 10104) were

originally obtained from China Medical Culture Collection. These three species have

been reported many times from microbial surveys in the space station, spacecrafts as

well as related studies(Crabbé Aurélie et al., 2011; Singh et al., 2018; Urbaniak et al.,

2018; Yang et al., 2021; Zea et al., 2016) and it was quite easy to distinguish them by

genetic relationship.

To simulate the nutrient-deficient environment on the inner surface of the space

station, overnight cultures of above strains were inoculated in the 10ml modified

minimal R2A broth (Yang et al., 2021) with a total density of 1×10^{6}/ml calculated by

direct counting under microscopy (Z. Song et al., 2015). The composition of modified

minimal R2A was shown in Table S1. In the mixed culture, the initial ratio of three

species was 1:1:1.

103 The LDIR environmental simulator (Fig. S1-A) was a refitted artificial climate

104 chamber that simulated the LDIR environment in the space station. The modification

105 included three main parts: a rotating wheel for placing cultures, two X-ray tubes for

106 alternatively ionizing the air inside to plasma, and a protective door for isolating the

107 radiation. Two X-ray tubes (XCH201, Dandong Shenbo Electronics Co. Ltd, China)

108 were installed at the inside top of the chamber and run at 20kV, 0.2mA, alternately. To

109 ensure that each culture is exposed to the same radiation dose, six boxes were affixed

110 to the rotary wheel running at a constant angular velocity. These boxes contained

111 plates with LDIR-treated cultures, while control cultures without LDIR treatment

112 were placed outside the radiation-exposed area but in the same climate chamber. The

113 protective door was sealed to maintain a constant temperature of 30°C, and normal

114 pressure during operation. 3D geometric configuration of LDIR environmental

115 simulator was accurately established based on its internal dimensions (Fig. S1-B).

116 2.2 Periodic sampling and successive incubation

117 During a seven-day experiment cycle, 1ml samples were taken from cultures at

118 the same time on each day. Then the residual parts were differently treated for

119 successive incubation by distinct groups in Table S2. Each group contains three

120 replicates. In each 1ml sample, 700μl was centrifuged for untargeted metabolomics

121 test with its supernatant, and two portions of 90μl bacterial fluid were used to depict

122 species abundance curves.

123 The composition of QAC was shown in Table S3.

124 2.3 Species abundance curve depicted by qPCR

125 Two portions of 90μl bacterial fluid above were applied to depict curves for

126 *Bacillus subtilis* (Kirchhoff & Cypionka, 2017; Tut et al., 2021; Xie et al., 2016) by

127 direct qPCR (Sup. A) and *Escherichia coli* as well as *Pseudomonas aeruginosa* by

PMA-qPCR, separately. For the PMA-treated one, 10μl 500mmol/L PMAxx (Cat.

40069, Biotium Inc., Hayward, CA, USA) was added into the sample under dark

conditions for forming the 50mmol/L final concentration. Then the mixed solution

was incubated at 20□ for 40 min in the dark on a vortex with mild continuous shaking

(100rpm) followed by the change of tube. Subsequently, the tube was exposure to a

120W blue LED light at 4□ for 30 min (Sup. B).

DNA was subsequently extracted from the solution above (with PMA or not) by

direct boiling with glass beads (Li et al., 2023). 0.1g 0.1mm glass beads (Cat.

11079101, BioSpec Products, Inc., Bartlesville, OK, USA) were added into the tube

containing 100μl resuspending solution before the heating procedure. After

centrifugation, the supernatant is DNA solution.

Primers, their qPCR standard curves, their amplicons, the composition of qPCR

reaction buffer, and the qPCR amplification program were the same as our previous

study (Li et al., 2023). 2×SYBR Green PCRmix (Cat. SR1110, Solarbio Science &

Technology Co. Ltd., Beijing, China) was used to provide PCR Buffer, $MgCl_2$, dNTPs,

Hs Taq DNA Polymerase, SYBR Green □ and so on. The reaction was performed

using Gentier 96C real-time PCR system (Tianlong Science and Technology Co. Ltd.,

Xi'an, China). The qPCR amplification program included: pre-denaturation at 95°C

for 10 min with a heating rate of 6°C/s; then 40 cycles of denaturation at 95°C for 10

sec with a temperature change rate of 5°C/s, annealing at 60°C for 20 sec with a

cooling rate 5°C/s and elongation at 72°C for 30 sec with heating rate 6°C/s.

2.4 Untargeted metabolomics test

Metabolites were quantified using ultra-performance liquid

chromatography-triple quadrupole time-of-flight mass spectrometry (UPLC-MS)

(Waters Corp., Milford, MA, USA). Raw data obtained from UPLC-MS were

quantified in Progenesis QI (Waters Corp.). Significantly differential metabolites

analysis was performed using BMKCloud (www.biocloud.net).

2.5 Mechanical modeling and computer simulation

The kinetic model was established based on generalized Lotka-Volterra (gLV)

equations and hypothetical mechanisms of disturbances. Parameters implying

attributes in the community were embedded into the model. Afterward, a

corresponding simulation model with dde23 solver was established by S-Function of

MATLAB/Simulink (MathWorks, Inc., Natick, MA, USA) to confirm the proposed

hypothesis through behavior similarity between computer simulation and

experimental data.

3. Results

This study aimed to demonstrate the influence of three different types of

disturbances on microbial community succession. Experiments were taken to depict

the species absolute abundance and differential metabolites during the succession. The

following mechanical models and computer simulation were proposed to explain

mechanisms of phenomena in these experiments.

3.1 qPCR results

The overview of all conditions was shown in Fig. 1. As seen in this figure,

distinct succession could be obtained by different treatments. Two-way repeated

measured ANOVA was applied using R (4.0.2) to calculate p-values to test if curves

showed a significant difference between the specific experimental group and the

control group in the overall experimental period, where the interaction effect of

treatment and time was considered first. If the interaction effect was not significant,

the main effect of treatment would be applied, where the p-value was marked as "p_m".

As shown in Fig. S2, in the control group, the total biomass and the absolute

abundance of *E. coli* reached highest points on the sixth day, when the density for

both of them was about 1.5×10^{10}/ml.

When exposed to LDIR (Fig. 2), the microbial community thrived after a period

of "hibernation". Compared with the control group, the total biomass rose on 5-7th

days and still peaked on the 6^{th} day, approximately reaching 3.5×10^{10}/ml. The

majority was still *E. coli*. The abundance of *B. subtilis* showed the trend of damped

periodic oscillations.

Dilution with gradients from 10% to 0.1‰ per day had no obvious different

influence with each other, as shown in Fig. S3 & S4. Taking 1‰ as an example (Fig.

3), compared with the control group, the total biomass and the abundance of *E. coli*

decreased on 4-6th days, exhibiting the trend from oscillation to robustness. The

abundance of *B. subtilis* significantly fell ($p_m = 1 \times 10^{-5}$), while that of

P. aeruginosa rose dramatically ($p = 0.004$).

Imposing 100% and 10% stock solution of QAC could rapidly bring the total

biomass down to below the initial density, when only *P. aeruginosa* could survive

below the initial density under this condition (Fig. 1). However, 1% and 1‰ stock

solution of QAC did not have the similar impact (Fig. 4). When treated with 1% stock

solution of QAC, compared with the control group, the peak of total biomass moved

back to 4th day, having no significant difference with the peak of the former. *E. coli*

had a great contribution to the total biomass on the 1-4th days, followed by the sharp

drop and extinction on the 6th day. Meanwhile, the abundance of

P. aeruginosa increased from the 5th day and peaked on the 6th day. In the group of 1‰

treatment, the total biomass exceeded the control group on the 7th day before the lag

on the 4-6th days, presenting the trend of rise with fluctuations. The abundance of *E.*

coli fluctuated continuously, and that of *P. aeruginosa* increased steadily.

When diluted with a 1‰ ratio per day, the total biomass ($p_m = 4 \times 10^{-4}$) and

the abundance of *E. coli* ($p_m = 4 \times 10^{-4}$) significantly rose in the whole period of

the LDIR group, compared with the group without LDIR (Fig. S5).

When exposed under LDIR, compared with the group without QAC, the total

biomass of the 1% stock solution of QAC group increased on the 2-4th days but

decreased on the 5-7th days, in which the abundance of *E. coli* went extinct quickly on

the 5th day and that of *P. aeruginosa* peaked on 6th day (Fig. S6-A). The total biomass

of the 1‰ stock solution of QAC group rose in fluctuations, exceeding the group

without QAC on the 4th and 7th days but lagged on the 5-6th days, when the latter

group was experiencing proliferation (Fig. S6-B). The majority was still *E. coli* but

the magnitude of *P. aeruginosa* also improved.

When treated with 1% stock solution of QAC, the total biomass of the LDIR

group went up in the whole period compared with the group without LDIR except on

the 4th day, when the latter group peaked. The abundance of *E. coli* increased on 1-3rd

days, then dropped on 4th day but this species still survived later. The abundance of

P. aeruginosa rose during the whole period and still peaked on the 6th day (Fig. S7-A).

When treated with 1‰ stock solution of QAC, the total biomass ($p = 0.003$) and the

abundance of *E. coli* ($p = 0.02$) of the LDIR group dramatically rocketed with

growing gaps in the whole period compared with the group without LDIR (Fig. S7-B).

The abundance of *P. aeruginosa* shrank a little but the difference was not significant

($p > 0.05$).

When LDIR, 1‰ dilution and 1% stock solution of QAC were imposed,

compared with the control group (Fig. S8-A), *E. coli* went extinct quickly on the 2nd

day. However, the abundance of *P. aeruginosa* substantially increased ($p = 3 \times 10^{-6}$).

When LDIR, 1‰ dilution and 1‰ stock solution of QAC were imposed, compared

with the control group (Fig. S8-B), the total biomass increased on 1-4th and 7th days,

showing a trend of fluctuation. The abundance of *P. aeruginosa* markedly increased

($p = 3 \times 10^{-4}$).

3.2 Untargeted metabolomics test results

Samples involved in metabolite analysis were divided into 24 series according to

biological repeat, and the symbol corresponding to the experimental condition of each

group was shown in Table 1. The quality control, OPLS-DA as well as its validity test,

OPLS-DA permutation test and screening of metabolites were done before analysis

(Sup. C).

Compared with the control group, the concentration of p-coumaroyl agmatine in

the group exposed under LDIR between 4-6th days dropped significantly ($p <$0.05,

Series B&M, C&N, D&O. Fig. S9). On 1st and 4-6th days, most of metabolic

pathways were inhibited, including polysaccharide metabolism, amino acid

metabolism, vitamin metabolism, xenobiotic degradation and metabolism, as well as

antibiotic biosynthesis (A&L, B&M, C&N, D&O; Fig. S10).

Compared with the control group, in the group diluted with 1‰ ratio per day,

N-acetylmannosamine was enriched on the 4th day (B&F), which was markedly

correlated with the rise of the abundance of *P. aeruginosa* ($p <$0.01, Fig. S11).

Pathways inhibited dominated on 4th day, including nucleotide metabolism, sugar

degradation, antibiotic synthesis, and aminobenzoate degradation (Fig. S12).

When treated with 1% stock solution of QAC, N-acetylmannosamine was more

abundant on 4th day compared with the control group (B&G), which was also

correlated with the rise of the abundance of *E. coli* ($p <$0.05, Fig. S13). Meanwhile,

staurosporine and tetracycline biosynthesis were active. When treated with 1‰ stock

solution of QAC, echitovenine was rarer on the 7[th] day compared with the control

group (E&K), which was also correlated with the increase of the abundance of

P. aeruginosa ($p < 0.01$, Fig. S14). On the same day, aminobenzoate degradation as

well as biosynthesis of alkaloids (derived from histidine and purine) were inhibited

(Fig. S15).

When both LDIR and 1% stock solution of QAC were imposed, compared with

the group under 1% stock solution of QAC, alkaloid biosynthesis was inhibited (Fig.

S16), and some nitrogenous long-chain organic matters were enriched on 5-6[th] days

(H&S, I&T), including tetradecenoyl carnitine and R-palmitoyl-(2-methyl)

ethanolamide, ranging from $1 \times 10^{10} \sim 2 \times 10^{12}$ times; compared with the control

group, cytochrome P450 metabolism was active on 5[th] day (C&S). When LDIR, 1‰

dilution and 1% stock solution of QAC were imposed together, the concentration of

N-[(2R)-2-hydroxypropyl] hexadecanamide was 2×10^{12} times higher than the

control group on the 4[th] day (B&W). However, when the QAC concentration was

changed to 1‰ of the stock solution (B&V, B&X, E&W, J&V, K&W), the

phenomenon of enrichment of nitrogenous long-chain organic matters disappeared;

alternatively, cyanoamino acid metabolism, related to nitrogen metabolism (Ding et

al., 2022), was active. The concentration of albaflavenone sharply diminished

($p < 0.01$, Fig. S17) on 4[th] day in the group with both LDIR and 1‰ stock solution of

QAC, compared with the control group (B&V). Aminobenzoate degradation on the 4[th]

day and antibiotic synthesis on the 7[th] day (K&W. Fig. S18) were restrained in the

group with both LDIR and 1‰ stock solution of QAC, compared with 1‰ stock

solution of QAC group. Under LDIR, when comparing 1% and 1‰ stock solution of

QAC groups, eicosanoids metabolism was active in the higher concentration one on

the 4th day (R&V. Fig. S19).

3.3 Mechanical modeling and simulation results

3.3.1 State variables

The mechanical model was composed of six state variables in four types: *B.*

subtilis biomass (x_1), *E. coli* biomass (x_2), *P.aeruginosa b*iomass (x_3); amount of

substrate (S), amount of QAC (Q), and amount of QAC degradation products (P).

3.3.2 Rate equations

(1) growth rate

The growth rate without competition of species i could be expressed as:

$$v_{ai} = \mu_i \left(\frac{S}{K_s + S} \right) x_i \#(3.1)$$

285 where μ_i was the specific growth rate of species i, and K_s denotes

286 substrate-dependent half-saturation constant for species i.

287 (2) interspecific competition

288 When there was no joint treatment of LDIR and QAC, the interspecific

289 competition rate against species i could be written as follows:

$$v_{bi} = \sum_{j \neq i}^{3} \alpha_{ji} x_i x_j \#(3.2)$$

290 where α_{ji} denotes the competition inhibition coefficient from species j to

291 species i.

292 When the microbial community was treated with LDIR or QAC, the interspecific

293 competition rate against species i could be written as follows:

$$v_{ci} = \sum_{j \neq i}^{3} \alpha_{ji} e^{-(c_{ji\_R}R + c_{ji\_Q}Q)} x_i x_j \#(3.3)$$

Where I was a constant, representing the LDIR coefficient; e was the Euler

number; c_{ji_R} and c_{ji_Q} expressed the interspecies cross-feeding response coefficient

between species j and species i caused by LDIR and QAC, respectively.

(3) intraspecific competition

When there was no joint treatment of LDIR and QAC, the intraspecific

competition rate of species i could be written as follows:

$$v_{di} = \beta_i x_i^2 \#(3.4)$$

300   where β_i denoted the intraspecific competition inhibition coefficient of species

301 i.

302   When the microbial community was treated with LDIR or QAC, the intraspecific

303 competition rate of species i could be written as follows:

$$v_{fi} = \beta_i e^{-(f_{i\_R}R + f_{i\_Q}Q)} x_i^2 \#(3.5)$$

304   where, f_{i_R} and f_{i_Q} denoted the intraspecies cross-feeding response coefficient

305 of species i caused by LDIR and QAC, respectively.

306   (4) metabolism

307   The metabolism rate of species i could be expressed as:

$$v_{gi} = \delta_i x_i \#(3.6)$$

308   where δ_i was the metabolism coefficient of species i.

309   (5) QAC disinfection

$$v_{hi} = \varepsilon_i Q x_i \#(3.7)$$

310   where ε_i was the QAC disinfection rate on species i.

311   (6) QAC degradation

$$v_{mi} = \gamma R Q \#(3.8)$$

312   where γ was the first-order kinetic coefficient of QAC degradation rate.

313   (7) generation of QAC degradation products

$$v_{ni} = \omega v_{mi} \#(3.9)$$

314   where ω was the calibration coefficient.

315   (8) consumption of substrate

$$v_S = \sum_{i=1}^{3} \varphi_i \, v_{ai} \#(3.10)$$

316   where φ_i was the substrate consumption coefficient during the growth of

317   species i.

318   (9) transient influence of Dilution and QAC

319   a. only dilution

$$\begin{cases} S = S_0, t = 1,2,\dots,7d \\ \quad x_{i,t} = Dx_{i,t-\Delta t} \end{cases} \#(3.11)$$

320   where S_0 was the original substrate concentration of the R2A culture; $x_{i,t}$ was

321   the abundance of species i at t moment; Δt was a very short time interval; D was

322   the proportion of the culture just before dilution to the total volume.

323   b. only QAC

$$Q_t = Q_{t-\Delta t} + Q_{Add}, t = 1,2,\dots,7d \#(3.12)$$

324   where Q_{Add} denoted the QAC added.

325   c. both dilution and QAC

$$\begin{cases} \qquad S = S_0, t = 1,2,\dots,7d \\ Q_t = DQ_{t-\Delta t} + Q_{Add}, t = 1,2,\dots,7d \\ \qquad\qquad x_{i,t} = Dx_{i,t-\Delta t} \end{cases} \#(3.13)$$

3.3.3 The overall mechanical model

Based on Eq(3.1)~(3.13), the overall mechanical model, i.e. a state space model

with six first-order nonlinear kinetic equations, could be formulated in a top-down

manner as follows:

(1) When there was no joint treatment of LDIR and QAC, the population

dynamics equation of species i was:

$\frac{dx_i}{dt} = v_{ai} - v_{bi} - v_{di} - v_{gi} - v_{hi} = \mu_i \left(\frac{S}{K_s+S}\right) x_i - \sum_{j \neq i}^{3} \alpha_{ji} x_i x_j - \beta_i x_i^2 - \delta_i x_i - \varepsilon_i Q x_i \#(3.14)$

When there was joint treatment of LDIR and QAC, it was:

$$\frac{dx_i}{dt} = v_{ai} - v_{ci} - v_{fi} - v_{gi} - v_{hi} =$$

$$\mu_i \left(\frac{S}{K_s + S}\right) x_i - \sum_{j \neq i}^{3} \alpha_{ji} e^{-(c_{ji\_R}R + c_{ji\_Q}Q)} x_i x_j - \beta_i e^{-(f_{i\_R}R + f_{i\_Q}Q)} x_i^2 - \delta_i x_i - \varepsilon_i Q x_i \#(3.15)$$

334   (2) The QAC degradation dynamics equation was:

$$\frac{dQ}{dt} = -v_{mi} = -\gamma R Q \#(3.16)$$

335   (3) The generation of QAC degradation production dynamics equation was:

$$\frac{dP}{dt} = v_{ni} = \omega \gamma R Q \#(3.17)$$

336   (4) The substrate change dynamics equation was:

$$\frac{dS}{dt} = v_{ni} - v_S = \omega \gamma R Q - \sum_{i=1}^{3} \varphi_i \mu_i \left(\frac{S}{K_s + S}\right) x_i \#(3.18)$$

3.3.4 Computer simulation

The code of the mechanical model above was written by S-Function in

MatLab/Simulink(R2019b,   9.7.0.1190202).   Simulation   experiments   were

implemented for simulating the control group and treatment groups of LDIR, dilution,

QAC as well as their combinations, as shown in Fig. S20. The simulation results

showed that the mechanical model had considerable behavior similarity with

experiments, while the mathematical expression of the mechanical model matched the

growth process of microbes, indicating the structure similarity between them.

Therefore, the mechanical model was considered valid.

4. Discussion

In this study, LDIR, dilution and QAC were extracted from the scenario as

treatments, and the microbial community succession under these uniformed different

treatments was observed by measuring species abundance and metabolites.

Mechanical models were built to explain experiments, with which the computer

simulation showed structural and behavioral similarity. The overall mechanisms were

shown in Fig. 5.

At the later stage, LDIR improved the total biomass, every species abundance

and their peaks, when the Hydra Effect happened. This phenomenon was shown in

both with/without LDIR comparison and with/without LDIR comparison under

dilution (Fig. 2 & S5). P-coumaroylagmatine could thicken cell walls, limit bacterial

movement and improve plant defense against pathogens(Häusler et al., 2014; Macoy

et al., 2015), thus its decrease on the 4-6[th] days (B&M, C&N, D&O) indicated that the

physiological activity was enhanced under LDIR. The inhibited antibiotic synthesis

pathway (B&M) suggested that external pressure was reduced and interspecific

competition was diminished. Therefore, the Hydra Effect occurred under LDIR. This

effect was more significant as the main effect in the comparison of the group under

LDIR and 1‰ dilution and only 1‰ dilution (Fig. S5). Except in *P. aeruginosa*,

dilution reshuffled all other components everyday and undermined the interaction

effect of treatment and time, where the enhancement effect of LDIR emerged.

Dilution, by removing a certain part of biomass during serial transfers, was a

non-specific and highly controllable disturbance way, where the transient removal did

not depend on response traits in the microbial community (Gibbons et al., 2016).

However, under non-selective dilution, *P. aeruginosa* was selectively released from

competitive exclusion. Aminobenzoate was the signal of xenobiotic degradation (Patil

et al., 2020; Wang et al., 2020), thus its degradation inhibition on 4[th] day (B&F)

suggested that organisms tended not to consume energy storage material (Smitha et al.,

2017) and the competition in the community was mitigated. It could be inferred that

many niches would be released without distinction under dilution, when

P. aeruginosa could reproduce based on its merits beyond others' (Abreu et al., 2019)

and occupy these niches. This release happened only under dilution with all gradients;

when both dilution and LDIR were applied, *E.coli* took the majority again (Fig.

S21-A).

QAC stood for specific disturbances, causing non-random removal according to

the microbe's response traits(Banitz et al., 2020; Calderón et al., 2018; Jurburg et al.,

2017; Srivastava & Vellend, 2005). Its effect also counted on disturbance intensities

and frequencies. Different QAC concentrations had various influences on the

microbial community, from elimination to promotion. The Hydra Effect even

happened under 1% and 1‰ stock solution of QAC, where simple competition

mitigation could not explain. Polymers of N-acetylmannosamine played a protective

role in *E. coli* (Sellner et al., 2021), thus their respective increments with dramatical

correlation told the reason why the species abundance of *E. coli* peaked in advance on

the 4th day under 1% stock solution (B&G). Besides, the active biosynthetic pathways

of antibiotics and cytokinins on the same day corroborated that microbes combated

adversity by more cell division and the total biomass also peaked. Echitovenine was a

monoterpene alkaloid with antimicrobial activity (Mohammed et al., 2021), whose

attenuation could interpret the growth of *P. aeruginosa* on the 7th day under 1‰ stock

solution (E&K), given their notable correlation. QAC would periodically selectively

remove a portion of individuals in the community, leading the way of community

succession to the direction of dominance by species that can tolerate the environment,

which could be pronounced in comparisons across gradients (Fig. S21-B&C). With

low-concentration QAC treatment, the community had adapted to the environment

and utilized it. QAC, was a series of long-chain organics that could be used by

microbes, especially in microbial co-metabolism by *Pseudomonas* (Harris & Knowles,

1983; Leadbetter & Foster, 1960; Zhang et al., 2015), which was widely and

systemically utilized in treating wastewater (biological nitrogen removal). On the

oligotrophic internal surface of the space station, carbon sources were mostly wiped

down, when co-metabolism might allow microbes to survive with remaining nitrogen

sources. Meanwhile, the inhibition of aminobenzoate degradation pathway also

flanked the corresponding decrease of the consumption of energy storage in the

community.

The upsurge of some nitrogenous long-chain organic matters under both LDIR

and QAC, compared with only QAC or the control group, strongly implies that QAC

may undergo radiolysis under LDIR. Several studies (Baidak & LaVerne, 2010;

Dhiman & LaVerne, 2013; Nogami et al., 1996; Pillay, 1986) implied that radiolysis

would happen on substrates containing QAC, especially under water acceleration. 1%

stock solution of QAC is substantially transformed from the toxic bactericide to

organic nitrogenous long-chain nutrients, including tetradecenoyl carnitine,

R-palmitoyl-(2-methyl) ethanolamide and N-[(2R)-2-hydroxypropyl] hexadecanamide

with significant differences between Series B&W, H&S, I&T, separately. This might

accelerate the microbial co-metabolism process. Cytochrome P450-related enzymes

prefer long-chain fatty acids with 15-16 carbon atoms (Lewis & Wiseman, 2005), thus

its activation might be related to added long-chain QAC utilization. Inhibited alkaloid

biosynthesis (H&S, I&T) and ameliorated cytochrome P450 metabolism (C&S) also

shows that the community could adapt to and consume QAC under LDIR.

Nevertheless, when the QAC concentration is 1‰, significant differences are no

longer detected for organic nitrogenous long-chain nutrients (B&V, B&X, E&W, J&V,

K&W), suggesting that the community can entirely co-metabolize QAC in this

situation. The activation of cyanoamino acid metabolism (B&X), inhibition of

aminobenzoate degradation (J&V) as well as inhibition of antibiotic synthesis (K&W)

also reconfirm the speculation of utilization of exogenous QAC as the source of

nitrogen and less pressure of the community caused by LDIR. Comparing different

concentration of QAC under LDIR (V&R), the active eicosanoids metabolism in the

higher concentration one showed that microbes might transform QAC into long-chain

fatty acid substances for consuming.

The proliferative microbes on the inner surface of the space station (Alekhova et

al., 2005) have been threatening its normal operation, which is a serious problem and

could be shifted to the issue of the effect of disturbance on community succession.

Disturbance is acknowledged as a key factor of biodiversity via affecting

eco-processes (Banitz et al., 2020). However, mechanisms behind the observed

community diversity under disturbance need to be further clarified (Fox, 2013; Shea

et al., 2004). First, few quantitative studies on disturbances have been reported, while

more relative studies focus on the qualitative aspect — this is due to both the

difficulty in defining the attribute of frequency, intensity, as well as extent (Shade et

al., 2012), and the wide variety of their characteristics and ecological impacts

(Fraterrigo & Rusak, 2008). Second, few studies are explicitly based on the time

course of microbial community composition under disturbance; instead, many solely

studies are on the sensitivity of composition at a given moment (Allison & Martiny,

2008), which is attributed to the inertial thinking in community ecology and research

tools in microbiology. Our previous study noticed the benefit of sole LDIR for the

microbial community (Yang et al., 2021). Abreu et al. (Abreu et al., 2019, 2020)

expanded the effect of dilution on multispecies community composition, providing an

example for quantifying disturbances with the time course. Nonetheless, in their

research, when distinguishing disturbances by pulsed versus sustained, pulsed ones

like dilution are not applicable for being directly integrated into the typical LV

equation. Finally, studies with similar gradient-based methods on combined multiple

factors with synergy, are still at an early stage. Most studies still concentrate on a single factor, and only a few have noticed the combined influence of multiple factors (Banitz et al., 2020; Zeng et al., 2020). In contrast, a large number of phenomena in nature are caused by the complex interaction of the latter condition (Elmqvist et al., 2003; Martínez-Ramos et al., 2016), including "ecological surprises" such as drastic shifts in community compositions and biodiversity (Banitz et al., 2020; Filbee☐Dexter et al., 2017; Newman, 2019).

In this study, "ecological surprise" occurred when human expectations deviated from the observed community succession behavior (Doak et al., 2008; Filbee☐Dexter et al., 2017). And these unexpected dynamics arose from human management and disturbances (Doak et al., 2008). Dilution (wiping) and QAC were recognized as efficient measures to take control of microbes (Sattar et al., 2015; Zhang et al., 2015) and applied in the space station (Singh et al., 2018). However, given the results above, the synergy emerged between LDIR and QAC, thus low-concentration QAC was transformed from the bactericide to nitrogen-rich long-chain nutrients. Selectively released *P. aeruginosa* under suppression after dilution would take these nutrients to thrive, changing the community structure and dominant species. That could be a possible reason for the phenomenon in the space station, and also gave a potential mechanism for "ecological surprise" under synergistical disturbances. It reminded us that tailored biocontrol strategies ought to be considered for the specific environment. Sometimes, the control measures were taken for granted that they were "disturbances" affecting targets without the consideration of their own properties and changes. It cannot be neglected that general strategies might not function under certain conditions.

Given the current microbial prevention strategies have some disadvantages,

future improvements can be adopted. Concerning the low concentration of QAC,

except for co-metabolism and radiolysis of QAC mentioned above, pre-package of

wipes can also cause it by both reduced antimicrobial activity of QAC with a

long-term soaking (Bloß et al., 2010, 2010; Slopek et al., 2011) and the chemical

combination with the wipe fibre (Dharan et al., 1999; X. Song et al., 2019). Target

surface properties also influence the disinfection (X. Song et al., 2019). The more

organic contamination on (dirtier) the surface, the lower the disinfection effect

(Williams et al., 2007). Moreover, radiolysis could happen just after the spacecraft

entered the outer space. In terms of dilution, its rate was limited on the order of 1%

even using two wipes successively (Gold & Hitchins, 2013), which is not enough

given results in this study. Furthermore, "wiping" itself could lead to

cross-contamination of microorganisms in different locations by mechanical

stimulation (Ramm et al., 2015; Siani et al., 2011; Williams et al., 2007). To attenuate

these effects, several measures could be considered, such as augmenting local wiping

time, wiping with mechanical equipment to restraint power variation (Sattar et al.,

2015; Williams et al., 2007), special antibacterial treatment on the surface requiring

microbial prevention, applying different wipe materials harder to react with QAC,

adjusting the formulation of QAC for a lower degradation rate (Pillay, 1986; Zhang et

al., 2015), avoiding surface exposure to radiation after wiping.

Despite these promising results, questions still remain. Further work is needed to

develop reliable experimental and analytical methods (e.g. based on the

individual-based model) (Li et al., 2022) for testing community succession

considering the spatial factor. The liquid culture implemented in this study aborted it

because of current biotechnology limitations on detecting microbial individuals.

Additional research based on more methods in bioinformatics is needed to better

understand the successive changes in microbial behaviors under disturbances. This study focused on the level of community succession. Hence, there is abundant room for further progress in determining evolutionary characteristics changes of the species facing disturbances in long-term experiments.

5. Conclusion

In this study, the intricate relationship between disturbance and the microbial community succession in the space station was elucidated. The proliferation under LDIR, the selective species release under dilution and complex responses of the community to QAC concentration were observed in experiments and their similarity was tested in modeling. The synergy of combined disturbances, QAC and LDIR, leads to the transformation of them from harmful factors to supportive ones. Therefore, it is necessary to consider synergistic effects and to apply tailored strategies when addressing the microbial proliferation challenge as well as other similar circumstances.

Fundings

This work was supported by Funds for International Cooperation and Exchange of the National Natural Science Foundation of China (32261133528).

Acknowledgment

We sincerely acknowledge the assistance from Chuanmeng Cai, Dengbo Chen, and Yong Feng in the long and non-stop periodic experiment, and the assistance from Yong Feng for depicting Fig.S1-B.

Data Availability Statement

The data and code used to produce results of this study are openly available in Zenodo at https://doi.org/10.5281/zenodo.11094948.

Author Contributions

SL, HL and DH conceptualized and designed the study; SL, XL and LC performed

experiments; SL and DH built the model and did the computer simulation; SL wrote

the first draft of the manuscript; all authors contributed substantially to revisions.

References

Abreu, C. I., Andersen Woltz, V. L., Friedman, J., & Gore, J. (2020). Microbial communities display
alternative stable states in a fluctuating environment. *PLOS Computational Biology*, *16*(5),
e1007934. https://doi.org/10.1371/journal.pcbi.1007934

Abreu, C. I., Friedman, J., Andersen Woltz, V. L., & Gore, J. (2019). Mortality causes universal
changes in microbial community composition. *Nature Communications*, *10*(1), 2120.
https://doi.org/10.1038/s41467-019-09925-0

Agustí, G., Fittipaldi, M., & Codony, F. (2018). Optimization of a viability PCR method for the
detection of listeria monocytogenes in food samples. *Current Microbiology*, *75*(6), 779–785.
https://doi.org/10.1007/s00284-018-1448-6

Alekhova, T. A., Aleksandrova, A. A., Novozhilova, T. Y., Lysak, L. V., Zagustina, N. A., &
Bezborodov, A. M. (2005). Monitoring of microbial degraders in manned space stations.
Applied Biochemistry and Microbiology, *41*(4), 382–389.
https://doi.org/10.1007/s10438-005-0065-x

Allison, S. D., & Martiny, J. B. H. (2008). Resistance, resilience, and redundancy in microbial
communities. *Proceedings of the National Academy of Sciences*, *105*(Supplement 1),
11512–11519. https://doi.org/10.1073/pnas.0801925105

Baidak, A., & LaVerne, J. A. (2010). Radiation-induced decomposition of anion exchange resins.
Journal of Nuclear Materials, *407*(3), 211–219.
https://doi.org/10.1016/j.jnucmat.2010.10.025

Banihashemi, A., Dyke, M. I., & Huck, P. M. (2012). Long-amplicon propidium monoazide-PCR
enumeration assay to detect viable campylobacter and salmonella. *Journal of Applied*
Microbiology, *113*(4), 863–873. https://doi.org/10.1111/j.1365-2672.2012.05382.x

Banitz, T., Chatzinotas, A., & Worrich, A. (2020). Prospects for Integrating Disturbances,
Biodiversity and Ecosystem Functioning Using Microbial Systems. *Frontiers in Ecology and*
Evolution, *8*. https://www.frontiersin.org/articles/10.3389/fevo.2020.00021

Baudoin, J. M., Guérold, F., Felten, V., Chauvet, E., Wagner, P., & Rousselle, P. (2008). Elevated
aluminium concentration in acidified headwater streams lowers aquatic hyphomycete
diversity and impairs leaf-litter breakdown. *Microbial Ecology*, *56*(2), 260–269.
https://doi.org/10.1007/s00248-007-9344-9

Baymiev, An. Kh., Baymiev, Al. Kh., Kuluev, B. R., Shvets, K. Yu., Yamidanov, R. S., Matniyazov,
R. T., Chemeris, D. A., Zubov, V. V., Alekseev, Ya. I., Mavzyutov, A. R., Ivanenkov, Ya. A.,
& Chemeris, A. V. (2020). Modern approaches to differentiation of live and dead bacteria
using selective amplification of nucleic acids. *Microbiology*, *89*(1), 13–27.
https://doi.org/10.1134/S0026261720010038

Bloß, R., Meyer, S., & Kampf, G. (2010). Adsorption of active ingredients of surface disinfectants
depends on the type of fabric used for surface treatment. *Journal of Hospital Infection*, *75*(1),
56–61. https://doi.org/10.1016/j.jhin.2009.11.027

Brooks, B., Firek, B. A., Miller, C. S., Sharon, I., Thomas, B. C., Baker, R., Morowitz, M. J., & Banfield, J. F. (2014). Microbes in the neonatal intensive care unit resemble those found in the gut of premature infants. *Microbiome*, *2*(1), 1. https://doi.org/10.1186/2049-2618-2-1

Calderón, K., Philippot, L., Bizouard, F., Breuil, M.-C., Bru, D., & Spor, A. (2018). Compounded disturbance chronology modulates the resilience of soil microbial communities and N-cycle related functions. *Frontiers in Microbiology*, *9*, 2721. https://doi.org/10.3389/fmicb.2018.02721

Codony, F., Dinh-Thanh, M., & Agustí, G. (2020). Key factors for removing bias in viability PCR-based methods: A review. *Current Microbiology*, *77*(4), 682–687. https://doi.org/10.1007/s00284-019-01829-y

Crabbé Aurélie, Schurr Michael J., Monsieurs Pieter, Morici Lisa, Schurr Jill, Wilson James W., Ott C. Mark, Tsaprailis George, Pierson Duane L., Stefanyshyn-Piper Heidi, & Nickerson Cheryl A. (2011). Transcriptional and proteomic responses of pseudomonas aeruginosa PAO1 to spaceflight conditions involve hfq regulation and reveal a role for oxygen. *Applied and Environmental Microbiology*, *77*(4), 1221–1230. https://doi.org/10.1128/AEM.01582-10

Dangles, O., & Chauvet, E. (2003). Effects of stream acidification on fungal biomass in decaying beech leaves and leaf palatability. *Water Research*, *37*(3), 533–538. https://doi.org/10.1016/S0043-1354(02)00359-7

Dharan, S., Mourouga, P., Copin, P., Bessmer, G., Tschanz, B., & Pittet, D. (1999). Routine disinfection of patients' environmental surfaces. Myth or reality? *Journal of Hospital Infection*, *42*(2), 113–117. https://doi.org/10.1053/jhin.1999.0567

Dhiman, S. B., & LaVerne, J. A. (2013). Radiolysis of simple quaternary ammonium salt components of Amberlite resin. *Journal of Nuclear Materials*, *436*(1–3), 8–13. https://doi.org/10.1016/j.jnucmat.2013.01.294

Ding, C., Wang, S., Li, J., & Wang, Z. (2022). Transcriptomic analysis reveals the mechanism of host growth promotion by endophytic fungus of rumex gmelinii turcz. *Archives of Microbiology*, *204*(7), 1–13.

Doak, D. F., Estes, J. A., Halpern, B. S., Jacob, U., Lindberg, D. R., Lovvorn, J., Monson, D. H., Tinker, M. T., Williams, T. M., Wootton, J. T., & others. (2008). Understanding and predicting ecological dynamics: Are major surprises inevitable. *Ecology*, *89*(4), 952–961.

Elmqvist, T., Folke, C., Nyström, M., Peterson, G., Bengtsson, J., Walker, B., & Norberg, J. (2003). Response diversity, ecosystem change, and resilience. *Frontiers in Ecology and the Environment*, *1*(9), 488–494. https://doi.org/10.1890/1540-9295(2003)001[0488:RDECAR]2.0.CO;2

Emerson, J. B., Adams, R. I., Román, C. M. B., Brooks, B., Coil, D. A., Dahlhausen, K., Ganz, H. H., Hartmann, E. M., Hsu, T., Justice, N. B., Paulino-Lima, I. G., Luongo, J. C., Lymperopoulou, D. S., Gomez-Silvan, C., Rothschild-Mancinelli, B., Balk, M., Huttenhower, C., Nocker, A., Vaishampayan, P., & Rothschild, L. J. (2017). Schrödinger's microbes: Tools

for distinguishing the living from the dead in microbial ecosystems. *Microbiome*, *5*(1), 86.

https://doi.org/10.1186/s40168-017-0285-3

Filbee☐Dexter, K., Pittman, J., Haig, H. A., Alexander, S. M., Symons, C. C., & Burke, M. J. (2017).

Ecological surprise: Concept, synthesis, and social dimensions. *Ecosphere*, *8*(12), e02005.

https://doi.org/10.1002/ecs2.2005

Fittipaldi, M., Nocker, A., & Codony, F. (2012). Progress in understanding preferential detection of

live cells using viability dyes in combination with DNA amplification. *Journal of*

Microbiological Methods, *91*(2), 276–289. https://doi.org/10.1016/j.mimet.2012.08.007

Fleischmann, S., Robben, C., Alter, T., Rossmanith, P., & Mester, P. (2021). How to evaluate

non-growing cells—Current strategies for determining antimicrobial resistance of VBNC

bacteria. *Antibiotics*, *10*(2), 115. https://doi.org/10.3390/antibiotics10020115

Fox, J. W. (2013). The intermediate disturbance hypothesis is broadly defined, substantive issues

are key: A reply to sheil and burslem. *Trends in Ecology & Evolution*, *28*(10), 572–573.

https://doi.org/10.1016/j.tree.2013.07.008

Fraterrigo, J. M., & Rusak, J. A. (2008). Disturbance-driven changes in the variability of ecological

patterns and processes. *Ecology Letters*, *11*(7), 756–770.

https://doi.org/10.1111/j.1461-0248.2008.01191.x

G. E. Hutchinson. (1961). The paradox of the plankton. *The American Naturalist*, *95*(882),

137–145.

García-Fontana, C., Narváez-Reinaldo, J. J., Castillo, F., González-López, J., Luque, I., &

Manzanera, M. (2016). A new physiological role for the DNA molecule as a protector against

drying stress in desiccation-tolerant microorganisms. *Frontiers in Microbiology*, *7*.

https://doi.org/10.3389/fmicb.2016.02066

Gibbons, S. M., Scholz, M., Hutchison, A. L., Dinner, A. R., Gilbert, J. A., & Coleman, M. L.

(2016). Disturbance regimes predictably alter diversity in an ecologically complex bacterial

system. *mBio*, *7*(6). https://doi.org/10.1128/mBio.01372-16

Gold, K. M., & Hitchins, V. M. (2013). Cleaning assessment of disinfectant cleaning wipes on an

external surface of a medical device contaminated with artificial blood or *streptococcus*

pneumoniae. *American Journal of Infection Control*, *41*(10), 901–907.

https://doi.org/10.1016/j.ajic.2013.01.029

Harris, R., & Knowles, C. J. (1983). Isolation and growth of a pseudomonas species that utilizes

cyanide as a source of nitrogen. In *Microbiology* (Vol. 129, Issue 4, pp. 1005–1011).

Microbiology Society. https://doi.org/10.1099/00221287-129-4-1005

Hastings, A. (2013). *Population biology: Concepts and models*. Springer Science & Business

Media.

Häusler, R. E., Ludewig, F., & Krueger, S. (2014). Amino acids – a life between metabolism and

signaling. *Plant Science*, *229*, 225–237. https://doi.org/10.1016/j.plantsci.2014.09.011

Jurburg, S. D., Nunes, I., Brejnrod, A., Jacquiod, S., Priemé, A., Sørensen, S. J., Van Elsas, J. D., &

Salles, J. F. (2017). Legacy effects on the recovery of soil bacterial communities from
extreme temperature perturbation. *Frontiers in Microbiology*, *8*, 1832.
https://doi.org/10.3389/fmicb.2017.01832

Kirchhoff, C., & Cypionka, H. (2017). Propidium ion enters viable cells with high membrane
potential during live-dead staining. *Journal of Microbiological Methods*, *142*, 79–82.
https://doi.org/10.1016/j.mimet.2017.09.011

Kralik, P., Nocker, A., & Pavlik, I. (2010). Mycobacterium avium subsp. Paratuberculosis viability
determination using F57 quantitative PCR in combination with propidium monoazide
treatment. *International Journal of Food Microbiology*, *141*, S80–S86.
https://doi.org/10.1016/j.ijfoodmicro.2010.03.018

Lang, J. M., Coil, D. A., Neches, R. Y., Brown, W. E., Cavalier, D., Severance, M.,
Hampton-Marcell, J. T., Gilbert, J. A., & Eisen, J. A. (2017). A microbial survey of the
international space station (ISS). *PeerJ*, *5*, e4029. https://doi.org/10.7717/peerj.4029

Lax, S., Smith, D. P., Hampton-Marcell, J., Owens, S. M., Handley, K. M., Scott, N. M., Gibbons, S.
M., Larsen, P., Shogan, B. D., Weiss, S., Metcalf, J. L., Ursell, L. K., Vázquez-Baeza, Y., Van
Treuren, W., Hasan, N. A., Gibson, M. K., Colwell, R., Dantas, G., Knight, R., & Gilbert, J. A.
(2014). Longitudinal analysis of microbial interaction between humans and the indoor
environment. *Science*, *345*(6200), 1048–1052. https://doi.org/10.1126/science.1254529

Leadbetter, E. R., & Foster, J. W. (1960). Bacterial oxidation of gaseous alkanes. *Archiv Für*
Mikrobiologie, *35*(1), 92–104. https://doi.org/10.1007/BF00425597

Lewis, D. F. V., & Wiseman, A. (2005). A selective review of bacterial forms of cytochrome P450
enzymes. *Enzyme and Microbial Technology*, *36*(4), 377–384.
https://doi.org/10.1016/j.enzmictec.2004.07.018

Li, S., Liu, H., Liu, G., & Hu, D. (2022). Simulated artificial interventions maintain species
diversity of spatially structured microbial communities in closed ecosystem. *Acta*
Astronautica, *201*, 39–47. https://doi.org/10.1016/j.actaastro.2022.08.023

Li, S., Liu, X., Li, Z., Liu, H., & Hu, D. (2023). Combination of direct boiling and glass beads
increases the purity and accuracy of bacterial DNA extraction. *Biotechnology Journal*,
18(11), 2300135.

Loozen, G., Boon, N., Pauwels, M., Quirynen, M., & Teughels, W. (2011). Live/dead real-time
polymerase chain reaction to assess new therapies against dental plaque-related pathologies:
Vitality real-time PCR. *Molecular Oral Microbiology*, *26*(4), 253–261.
https://doi.org/10.1111/j.2041-1014.2011.00615.x

Macoy, D. M., Kim, W.-Y., Lee, S. Y., & Kim, M. G. (2015). Biotic stress related functions of
hydroxycinnamic acid amide in plants. *Journal of Plant Biology*, *58*(3), 156–163.
https://doi.org/10.1007/s12374-015-0104-y

Martin, B., Raurich, S., Garriga, M., & Aymerich, T. (2013). Effect of amplicon length in propidium
monoazide quantitative PCR for the enumeration of viable cells of salmonella in cooked ham.

Food Analytical Methods, *6*(2), 683–690. https://doi.org/10.1007/s12161-012-9460-0

Martínez-Ramos, M., Ortiz-Rodríguez, I. A., Piñero, D., Dirzo, R., & Sarukhán, J. (2016).

Anthropogenic disturbances jeopardize biodiversity conservation within tropical rainforest

reserves. *Proceedings of the National Academy of Sciences*, *113*(19), 5323–5328.

https://doi.org/10.1073/pnas.1602893113

Mohammed, A. E., Abdul-Hameed, Z. H., Alotaibi, M. O., Bawakid, N. O., Sobahi, T. R.,

Abdel-Lateff, A., & Alarif, W. M. (2021). Chemical diversity and bioactivities of

monoterpene indole alkaloids (MIAs) from six apocynaceae genera. *Molecules*, *26*(2), 488.

https://doi.org/10.3390/molecules26020488

Newman, E. A. (2019). Disturbance ecology in the anthropocene. *Frontiers in Ecology and*

Evolution, *7*, 147. https://doi.org/10.3389/fevo.2019.00147

Nocker, A., Sossa-Fernandez, P., Burr, M. D., & Camper, A. K. (2007). Use of propidium

monoazide for live/dead distinction in microbial ecology. *Applied and Environmental*

Microbiology, *73*(16), 5111–5117. https://doi.org/10.1128/AEM.02987-06

Nogami, M., Fujii, Y., & Sugo, T. (1996). Radiation resistance of pyridine type anion exchange

resins for spent fuel treatment. *Journal of Radioanalytical and Nuclear Chemistry Articles*,

203(1), 109–117. https://doi.org/10.1007/BF02060385

Novikova, N., De Boever, P., Poddubko, S., Deshevaya, E., Polikarpov, N., Rakova, N., Coninx, I.,

& Mergeay, M. (2006). Survey of environmental biocontamination on board the international

space station. *Research in Microbiology*, *157*(1), 5–12.

https://doi.org/10.1016/j.resmic.2005.07.010

Patil, S. M., Suryavanshi, M. V., Chandanshive, V. V., Kurade, M. B., Govindwar, S. P., & Jeon,

B.-H. (2020). Regeneration of textile wastewater deteriorated microbial diversity of soil

microcosm through bioaugmentation. *Chemical Engineering Journal*, *380*, 122533.

https://doi.org/10.1016/j.cej.2019.122533

Pickett, S., & White, P. (1985). *Ecology of natural disturbance and patch dynamics*.

Pillay, K. K. S. (1986). A review of the radiation stability of ion exchange materials. *Journal of*

Radioanalytical and Nuclear Chemistry Articles, *102*(1), 247–268.

https://doi.org/10.1007/BF02037966

Powell, J. R., Karunaratne, S., Campbell, C. D., Yao, H., Robinson, L., & Singh, B. K. (2015).

Deterministic processes vary during community assembly for ecologically dissimilar taxa.

Nature Communications, *6*(1), 8444. https://doi.org/10.1038/ncomms9444

Ramm, L., Siani, H., Wesgate, R., & Maillard, J.-Y. (2015). Pathogen transfer and high variability in

pathogen removal by detergent wipes. *American Journal of Infection Control*, *43*(7),

724–728. https://doi.org/10.1016/j.ajic.2015.03.024

Rosado, H., Doyle, M., Hinds, J., & Taylor, P. W. (2010). Low-shear modelled microgravity alters

expression of virulence determinants of staphylococcus aureus. *Acta Astronautica*, *66*(3),

408–413. https://doi.org/10.1016/j.actaastro.2009.06.007

Sattar, S. A., Bradley, C., Kibbee, R., Wesgate, R., Wilkinson, M. A. C., Sharpe, T., & Maillard, J.-Y.
(2015). Disinfectant wipes are appropriate to control microbial bioburden from surfaces: Use
of a new ASTM standard test protocol to demonstrate efficacy. *Journal of Hospital Infection*,
91(4), 319–325. https://doi.org/10.1016/j.jhin.2015.08.026

Sellner, B., Prakapaitė, R., van Berkum, M., Heinemann, M., Harms, A., & Jenal, U. (2021). A new
sugar for an old phage: A c-di-GMP-dependent polysaccharide pathway sensitizes
escherichia coli for bacteriophage infection. *mBio*, *12*(6), e03246-21.
https://doi.org/10.1128/mbio.03246-21

Shade, A., Peter, H., Allison, S. D., Baho, D. L., Berga, M., Bürgmann, H., Huber, D. H.,
Langenheder, S., Lennon, J. T., Martiny, J. B., & others. (2012). Fundamentals of microbial
community resistance and resilience. *Frontiers in Microbiology*, *3*, 417.

Shea, K., Roxburgh, S. H., & Rauschert, E. S. J. (2004). Moving from pattern to process:
Coexistence mechanisms under intermediate disturbance regimes. *Ecology Letters*, *7*(6),
491–508. https://doi.org/10.1111/j.1461-0248.2004.00600.x

Siani, H., Cooper, C., & Maillard, J.-Y. (2011). Efficacy of "sporicidal" wipes against Clostridium
difficile. *American Journal of Infection Control*, *39*(3), 212–218.
https://doi.org/10.1016/j.ajic.2011.01.006

Singh, N. K., Wood, J. M., Karouia, F., & Venkateswaran, K. (2018). Succession and persistence of
microbial communities and antimicrobial resistance genes associated with international
space station environmental surfaces. *Microbiome*, *6*(1), 204.
https://doi.org/10.1186/s40168-018-0585-2

Slopek, R., Condon, B., Sawhney, P., Reynolds, M., & Allen, C. (2011). Adsorption of
alkyl-dimethyl-benzyl-ammonium chloride on differently pretreated non-woven cotton
substrates. *Textile Research Journal*, *81*(15), 1617–1624.
https://doi.org/10.1177/0040517511407373

Smitha, M., Singh, S., & Singh, R. (2017). Microbial biotransformation: A process for chemical
alterations. *J Bacteriol Mycol Open Access*, *4*(2), 85.

Song, X., Vossebein, L., & Zille, A. (2019). Efficacy of disinfectant-impregnated wipes used for
surface disinfection in hospitals: A review. *Antimicrobial Resistance & Infection Control*,
8(1), 139. https://doi.org/10.1186/s13756-019-0595-2

Song, Z., Han, H., & Zhu, W. (2015). Morphological variation and recovery mechanism of residual
crude oil by biosurfactant from indigenous bacteria: Macro-and pore-scale experimental
investigations. *Journal of Microbiology and Biotechnology*, *25*(6), 918–929.

Srivastava, D. S., & Vellend, M. (2005). Biodiversity-ecosystem function research: Is it relevant to
conservation? *Annual Review of Ecology, Evolution, and Systematics*, *36*(1), 267–294.
https://doi.org/10.1146/annurev.ecolsys.36.102003.152636

Stegen, J. C., Lin, X., Fredrickson, J. K., Chen, X., Kennedy, D. W., Murray, C. J., Rockhold, M. L.,
& Konopka, A. (2013). Quantifying community assembly processes and identifying features

that impose them. *The ISME Journal*, *7*(11), 2069–2079. https://doi.org/10.1038/ismej.2013.93

Stewart, F. M., & Levin, B. R. (1973). Partitioning of resources and the outcome of interspecific competition: A model and some general considerations. *The American Naturalist*, *107*(954), 171–198. https://doi.org/10.1086/282825

Sun, Y., Hu, D., Li, L., Jing, Z., Wei, C., Zhang, L., Fu, Y., & Liu, H. (2016). Influence mechanism of low-dose ionizing radiation on escherichia coli DH5α population based on plasma theory and system dynamics simulation. *Journal of Environmental Radioactivity*, *151*, 185–192. https://doi.org/10.1016/j.jenvrad.2015.10.007

Tolkkinen, M., Mykrä, H., Markkola, A., Aisala, H., Vuori, K., Lumme, J., Pirttilä, A. M., & Muotka, T. (2013). Decomposer communities in human☐impacted streams: Species dominance rather than richness affects leaf decomposition. *Journal of Applied Ecology*, *50*(5), 1142–1151. https://doi.org/10.1111/1365-2664.12138

Tut, G., Magan, N., Brain, P., & Xu, X. (2021). Molecular assay development to monitor the kinetics of viable populations of two biocontrol agents, *bacillus subtilis* QST 713 and *gliocladium catenulatum* J1446, in the phyllosphere of lettuce leaves. *Biology*, *10*(3), 224. https://doi.org/10.3390/biology10030224

Urbaniak, C., Sielaff, A. C., Frey, K. G., Allen, J. E., Singh, N., Jaing, C., Wheeler, K., & Venkateswaran, K. (2018). Detection of antimicrobial resistance genes associated with the international space station environmental surfaces. *Scientific Reports*, *8*(1), 814. https://doi.org/10.1038/s41598-017-18506-4

Wang, Q., Liang, J., Zhang, S., Yoza, B. A., Li, Q. X., Zhan, Y., Ye, H., Zhao, P., & Chen, C. (2020). Characteristics of bacterial populations in an industrial scale petrochemical wastewater treatment plant: Composition, function and their association with environmental factors. *Environmental Research*, *189*, 109939. https://doi.org/10.1016/j.envres.2020.109939

Williams, G. J., Denyer, S. P., Hosein, I. K., Hill, D. W., & Maillard, J.-Y. (2007). The development of a new three-step protocol to determine the efficacy of disinfectant wipes on surfaces contaminated with *staphylococcus aureus*. *Journal of Hospital Infection*, *67*(4), 329–335. https://doi.org/10.1016/j.jhin.2007.08.012

Xie, X., Wang, S., Jiang, S. C., Bahnemann, J., & Hoffmann, M. R. (2016). Sunlight-activated propidium monoazide pretreatment for differentiation of viable and dead bacteria by quantitative real-time polymerase chain reaction. *Environmental Science & Technology Letters*, *3*(2), 57–61. https://doi.org/10.1021/acs.estlett.5b00348

Yang, X., Li, S., Song, G., Xu, X., & Hu, D. (2021). Microbial diversity formed and maintained through substrate feedback regulation and delayed responses induced by low-dose ionizing radiation. *Acta Astronautica*, *188*, 239–251. https://doi.org/10.1016/j.actaastro.2021.07.027

Yang, X., Xu, X., & Hu, D. (2020). Succession mechanism of microbial community with high species diversity in nutrient-deficient environments with low-dose ionizing radiation.

Ecological Modelling, *435*, 109270. https://doi.org/10.1016/j.ecolmodel.2020.109270

Zea, L., Prasad, N., Levy, S. E., Stodieck, L., Jones, A., Shrestha, S., & Klaus, D. (2016). A

molecular genetic basis explaining altered bacterial behavior in space. *PLOS ONE*, *11*(11),

e0164359. https://doi.org/10.1371/journal.pone.0164359

Zeng, X.-Y., Li, S.-W., Leng, Y., & Kang, X.-H. (2020). Structural and functional responses of

bacterial and fungal communities to multiple heavy metal exposure in arid loess. *Science of*

The Total Environment, *723*, 138081. https://doi.org/10.1016/j.scitotenv.2020.138081

Zhang, C., Cui, F., Zeng, G., Jiang, M., Yang, Z., Yu, Z., Zhu, M., & Shen, L. (2015). Quaternary

ammonium compounds (QACs): A review on occurrence, fate and toxicity in the

environment.   *Science   of   The   Total   Environment*,   *518–519*,   352–362.

https://doi.org/10.1016/j.scitotenv.2015.03.007

Zhou, J., & Ning, D. (2017). Stochastic community assembly: Does it matter in microbial ecology?

Microbiology and Molecular Biology Reviews, *81*(4), e00002-17.

Figures

Figure 1: The overview of the community succession under all treatments. The

x-axis represents the QAC concentration (e.g. 1 stands for 100%, 10^{-1} stands for 10%),

where the left side of axis includes groups without LDIR and the right side includes

groups with LDIR; y-axis represents the dilution ratio (e.g. 10^{-1} stands for 10%, 10^{-2}

stands for 1%). each position on the x-y plane corresponds to a different treatment.

z-axis denotes the number of days, and the seven spheres in each cluster mean seven

days of community succession for a treatment group. The radius of the spheres

indicates the total biomass in the community (compared to the maximum value and

taken as log ten). RGB color shows the percentage of different species (Blue, red and

green stand for *Bacillus subtilis*, *Escherichia coli* and *Pseudomonas aeruginosa*,

respectively). WO means "without". The 3D version is uploaded as a .fig file in the

supplementary materials.

Figure 2: The absolute abundance of three species (A-C) and the total biomass (D)

under LDIR (solid line) and in the control group (dotted line).

Figure 3: The absolute abundance of three species (A-C) and the total biomass (D)

under 1‰ dilution (solid line) and in the control group (dotted line).

Figure 4: The absolute abundance of *E. coli*, *P. aeruginosa* as well as the total

biomass under 1% QAC (A, solid line) and 1‰ QAC (B, solid line) and in the

control group (dotted line). *B. subtilis* has a very low abundance, so it is not shown

in the figure; the same as below.

Figure 5: The overall mechanisms of LDIR, dilution and QAC on microbial

community succession. LDIR activates all species' growth and reproduction. Dilution

allows specific species to be selectively released from suppression.

Low-concentration QAC undergoes radiolysis under LDIR and co-metabolism by

bacteria, thus changing from bactericide with harmful effects to nutrients. This figure

is created with BioRender.com.

Table

Table 1 The experimental conditions of distinct metabolomics test series

Series	LDIR	Dilution	QAC	Day
A	N	N	N	1
B	N	N	N	4
C	N	N	N	5
D	N	N	N	6
E	N	N	N	7
F	N	1‰	N	4
G	N	N	1%	4
H	N	N	1%	5
I	N	N	1%	6
J	N	N	1‰	4
K	N	N	1‰	7
L	Y	N	N	1
M	Y	N	N	4
N	Y	N	N	5
O	Y	N	N	6
P	Y	N	N	7
Q	Y	1‰	N	4
R	Y	N	1%	4
S	Y	N	1%	5
T	Y	N	1%	6
U	Y	1‰	1%	4

V	Y	N	1‰	4
W	Y	N	1‰	7
X	Y	1‰	1‰	4

Supplementary Materials

Figure S1: The LDIR environmental simulator (A) and its 3D geometric

configuration (B).

Figure S2: The absolute abundance of three species (A-C) and the total biomass

(D) in the control group.

Figure S3: The absolute abundance of three species (A-C) and the total biomass

(D) under different dilution ratios. The thinner the line, the more diluted the

community in the group is.

Figure S4: The relative abundance of three species under different dilution ratios.

A-D represent 10%, 1%, 1‰ and 0.1‰ dilution ratios, respectively.

Figure S5: The absolute abundance of three species (A-C) and the total biomass

(D) under LDIR and 1‰ dilution (solid line) as well as only 1‰ dilution (dotted

line).

Figure S6: The absolute abundance of *E. coli*, *P. aeruginosa* as well as the total

biomass under LDIR and 1% QAC (A, solid line) or LDIR and 1‰ QAC (B,

solid line) as well as only LDIR (dotted line).

Figure S7: The absolute abundance of *E. coli*, *P. aeruginosa* as well as the total

biomass under LDIR and 1% QAC (A, solid line) or LDIR and 1‰ QAC (B,

solid line) as well as only 1% QAC (A, dotted line) or 1‰ QAC (B, dotted line).

Figure S8: The absolute abundance of *E. coli*, *P. aeruginosa* as well as the total

biomass under LDIR, 1% dilution and 1% QAC (A, solid line) or LDIR, 1%

dilution and 1‰ QAC (B, solid line) as well as in the control group (dotted line).

Figure S9: the concentration of p-coumaroyl agmatine in the group exposed

under LDIR between the 4-6ᵗʰ days dropped significantly compared with the

control group. "**" meant (<).

Figure S10: the differential abundance scores of differential metabolisms

between the group under LDIR and the control group. A-D represent the 1, 4, 5,

6[th] day, respectively.

Figure S11: the enrichment of N-acetylmannosamine was markedly correlated

with the rise of the abundance of *P. aeruginosa* (<) on the 4[th] day of the

group diluted with 1‰ ratio per day, compared with the control group.

Metabolites with the top 10 highest concentrations in either group, or whose

were considered in the intergroup correlation analysis. "**" meant

(<), and "*" meant (<). The same is as below.

Figure S12: the differential abundance scores of differential metabolisms on 4th

day between the group diluted with 1‰ ratio per day and the control group.

Figure S13: the enrichment of N-acetylmannosamine was markedly correlated

with the rise of the abundance of *E. coli* (<) on the 4th day of the group

with 1% stock solution of QAC, compared with the control group.

Figure S14: the drop of echitovenine was markedly correlated with the increase

of the abundance of *P. aeruginosa* (<) on the 7th day of the group with 1‰

stock solution of QAC, compared with the control group.

Figure S15: the differential abundance scores of differential metabolisms on 7th

 day between the group with 1‰ stock solution of QAC and the control group.

 Figure S16: the differential abundance scores of differential metabolisms

 between the group with both LDIR and 1% stock solution of QAC and the group

891 **under 1% stock solution of QAC.** A and B represent the 5th and 6th day, separately.

892 **Figure S17: the concentration of albaflavenone sharply diminished on 4th day in**

893 **the group with both LDIR and 1‰ stock solution of QAC, compared with the**

894 **control group.**

Figure S18: the differential abundance scores of differential metabolisms between the group with both LDIR and 1‰ stock solution of QAC and the group under 1‰ stock solution of QAC. A and B represent the 4th and 7th day, separately.

Figure S19: the differential abundance scores of differential metabolisms on 4[th]

day between the groups with 1% stock solution of QAC and 1‰ stock solution of

QAC.

Figure S20: The simulation results from the model built in MatLab/Simulink.

A-H represent the control group, LDIR, low dilution rate, high dilution rate, low QAC

concentration, high QAC concentration, LDIR with low QAC concentration, all three

factors together, respectively.

Figure S21: The overview of the community succession under all treatments and

its slides. (A) The two planes in the yellow boxes showed that *E.coli* took the

majority when both dilution and LDIR were applied compared to only dilution. (B)

With higher gradients of QAC under 1%, *P. aeruginosa* gradually became the

dominant species with LDIR. (C) With higher gradients of QAC under 1%,

P. aeruginosa gradually became the dominant species without LDIR.

Figure S22: the actual and theoretical counts for 10× gradient dilution of *B.*

***subtilis* cells.**

Table S1 The composition of modified minimal R2A

Component	per 1000ml	Final concentration
NaCl	18.0g	1.8%
Peptone	0.5g	0.05%
Yeast Extract Powder	0.5g	0.05%
Casein Peptone	0.5g	0.05%
Sodium Pyruvate	0.3g	0.03%
ddH$_2$O	1000ml	

916 **Table S2 Conditions of Distinct Groups**

Group	LDIR	Dilution	QAC
1	N	N	N
2	N	10%	N
3	N	1%	N
4	N	1‰	N

5	N	0.1‰	N
6	N	N	100%
7	N	N	10%
8	N	N	1%
9	N	N	1‰
10	Y	N	N
11	Y	10%	N
12	Y	1%	N
13	Y	1‰	N
14	Y	0.1‰	N
15	Y	N	100%
16	Y	10%	100%
17	Y	1%	100%
18	Y	1‰	100%
19	Y	0.1‰	100%
20	Y	N	10%
21	Y	10%	10%
22	Y	1%	10%
23	Y	1‰	10%
24	Y	0.1‰	10%
25	Y	N	1%
26	Y	10%	1%
27	Y	1%	1%
28	Y	1‰	1%
29	Y	0.1‰	1%
30	Y	N	1‰
31	Y	10%	1‰
32	Y	1%	1‰

33	Y	1‰	1‰
34	Y	0.1‰	1‰

*"N" means this condition was not applicable in this group. The ratio of dilution

meant that part of the culture from t-th day was taken into the R2A solution for

gaining the 10ml new culture of $t + 1$-th day. No dilution meant that R2A solution

was added into the old culture to gain the 10ml new culture. The ratio of QAC meant

the stock solution of QAC was added into the culture to make that concentration of

the stock solution of QAC.

Table S3 The composition of QAC

Component	per 100ml	Final Concentration
Octyl Decyl Dimethyl Ammonium Chloride (Purity: 80%)	0.5g	0.04%
Dioctyl Dimethyl Ammonium Chloride (Purity: 80%)	0.25g	0.02%
Didecyl Dimethyl Ammonium Chloride (Purity: 95%)	0.21g	0.02%
Alkyl Dimethyl Benzyl Ammonium Chloride, (50% C14, 40% C12, 10% C16) (Purity: 80%)	0.66g	0.0532%
ddH$_2$O	98ml	-

924 **Table S4 Orthogonal Experiments for PMA-qPCR Conditions**

No	PMA Concentration (mmol/L)	PMA Incubation Time (min)	PMA Incubation Temperature (□)	Light Exposure Time (min)	Light Exposure Temperature (□)	Light Intensity (W)	Changing Tubes or not	$\Delta Ct(L)$	$\Delta Ct(D)$
1	50(B)	10	20(A)	30(B)	4(A)	120(B)	N(B)	-0.642	-12.604
2	25(A)	10(A)	37(B)	15(A)	20(B)	120	N	1.845	-8.208
3	25	10	20	15	4	60(A)	Y(A)	-0.712	-14.497
4	50	10	37	30	20	60	Y	0.468	-11.381
5	50	30(B)	37	15	4	60	N	1.614	-15.257
6	25	30	20	30	20	60	N	0.772	-9.731

No								ΔCt(L)	ΔCt(D)
7	50	30	20	15	20	120	Y	-0.592	-17.972
8	25	30	37	30	4	120	Y	0.17	-10.692
TL_A	2.075	0.959	-1.174*	2.155	0.43*	2.142	-0.666*		
TD_A	-43.128	-46.69	-54.804*	-38.779	-53.05*	-50.866	-54.542*		
TL_B	0.848 (p=0.092)	1.964	4.097	0.768 (p=0.059)	2.493	0.781 (p=0.063)	3.589		
TD_B	-57.214*	-53.652*	-45.538	-44.408*	-47.292	-49.476	-45.8		

*$\Delta Ct(L)$ and $\Delta Ct(D)$ represent the measured live/dead microbes abundance difference

between the control group and experimental group (with PMA), respectively. TL_A and

TD_A stand for the sum of the test results $\Delta Ct(L)$/ $\Delta Ct(D)$ for the column with level A,

respectively. TL_B and TD_B stand for the sum of the test results $\Delta Ct(L)$/ $\Delta Ct(D)$ for the

column with level B, respectively. "*" means $p > 0.05$.

Table S5 Gradient Experiments for PMA-qPCR Conditions

No	PMA Concentration (mmol/L)	PMA Incubation Time (min)	PMA Incubation Temperature (□)	Light Exposure Time (min)	Light Exposure Temperature (□)	Light Intensity (W)	Changing Tubes or not	ΔCt(L)	ΔCt(D)
1	50	40	4	30	4	120	Y	0.89	-10.421
2	50	30	20	30	4	120	Y	1.242	-16.583
3	50	10	20	30	4	120	Y	0.244	-13.626
4	50	5	4	30	4	120	Y	0.891	-11.641
5	50	10	4	30	4	120	Y	0.497	-13.451
6	50	30	4	30	4	120	Y	1.045	-18.622
7	50	5	20	30	4	120	Y	1.691	-14.067
8	50	40	20	30	4	120	Y	1.319	-19.609*

 Supplementary Material A

Several studies from different perspectives have shown a special phenomenon in

Bacillus subtilis (Kirchhoff & Cypionka, 2017; Tut et al., 2021; Xie et al., 2016),

which is, that some nucleic acid dyes like PMA could penetrate a large number of

living cells due to the excessively strong cell membrane potential. Therefore, at the

macro level, the rate of false-negative qPCR results would increase, and the threshold

for detection of living *Bacillus subtilis* cells through the PMA-qPCR method would

spike. In the pre-experiments of this study, *Bacillus subtilis* did not lead in the

community. Hence, daily dilution to 10% of the culture yesterday experiment was

performed, and the magnitude of counts of *Bacillus subtilis* was much higher than the

circumstance "No *Bacillus subtilis* survived, its DNA was diluted simultaneously", as

shown in Fig. S22. Therefore, direct qPCR was applicable for approximately counting

Bacillus subtilis.

Supplementary Material B

Eliminating the DNA of dead bacteria cells is the key aim of PMA in

PMA-qPCR. The standard PMA protocol might be suitable for natural environmental

samples, but not enough for high-abundance samples, like artificial microbial

communities used in this study. Therefore, it is important to optimize its conditions.

Related factors included PMA concentration, PMA incubation (with oscillation at

100rpm) time, PMA incubation temperature, light exposure time, light exposure

temperature, light intensity (expressed as the blue LED bulb power), whether

changing tubes between incubation and light exposure (Banihashemi et al., 2012;

Baymiev et al., 2020; Codony et al., 2020; Emerson et al., 2017; Fittipaldi et al., 2012;

Fleischmann et al., 2021; Loozen et al., 2011; Martin et al., 2013). To find the highest

DNA of dead bacteria cells elimination rate, orthogonal experiments (Table S4) were

implemented first on *P. aeruginosa* to screen the better condition of the two-level

factor (PMA was not applicable for *B. subtilis* as seen in Sup. A and the primer of *E.*

coli used for this study had a relatively high detection threshold). PMAxx was used as

the improved reagent. Conditions were selected in previous studies (Agustí et al.,

2018; Codony et al., 2020; Emerson et al., 2017; Fittipaldi et al., 2012; Fleischmann

et al., 2021; García-Fontana et al., 2016; Kralik et al., 2010; Nocker et al., 2007).

The larger the absolute value of $\Delta Ct(D)$, the higher the dead bacteria cells'

elimination rate. Based on no significant difference of $\Delta Ct(D)$, the closer $\Delta Ct(L)$ to

zero, the smaller the influence of PMA on live cells. From the orthogonal experiments,

these conditions were significantly better: the PMA concentration was 50 mmol/L, the

light exposure time was 30min, the light exposure temperature was 4□, and changing

tubes were applied. 120W light intensity has a p-value of 0.063 so it was also

selected.

Incubation time and temperature were tested another time in gradient

experiments on *P. aeruginosa* as Table S5. As seen in Table, when the PMA

incubation time was 40min and the PMA incubation temperature was 20□, the

absolute value of $\Delta Ct(D)$ was the largest with a significant difference. Therefore, the

optimal condition was: to incubate with 50mmol/L PMAxx (Cat. 40069, Biotium Inc.,

Hayward, CA, USA) 100rpm for 40min at 20□, change a new tube, and expose it

under a blue LED bulb still for 30min at 4□.

When applying this optimal condition to *E. coli*, the $\Delta Ct(D)=-18.268$. Compared

to the standard condition of PMA protocol, the detective elimination rate of dead *E.*

coli surged from 25.976% to 99.999%, and that of *P. aeruginosa* dropped from 25.097%

to 99.999%. The optimal condition can fulfill the need to measure high species

abundance in this study and it was used when treating samples for these two species'

abundance data.

Supplementary Material C

In this study, based on LC-QTOF, the qualitative and quantitative analysis of

untargeted metabolomics of 72 samples was carried out. A total of 12,079 peaks were

detected, of which 1,529 metabolites were annotated. The RSD ≤ 30% ones accounted

for more than 60%, so it could be assumed that QC samples have good reproducibility,

the instrument was stable, and the requirements of quality control were met.

In OPLS-DA (Orthogonal Projections to Latent Structures-Discriminant

Analysis), all results of the experimental and their corresponding control groups could

be completely separated, indicating significant overall changes in metabolites for each

condition. In the comparison of groups J and V, the Q2Y < 0.5, thus VIP could not be

considered as a screening condition for differential metabolites. Besides the above,

Q2Y of all other comparisons > 0.5, indicating that the OPLS-DA model was valid.

Permutation tests were performed to check the reliability of the OPLS-DA model. The

slopes of Q2Y were positive for all groups except for the comparison of J and V,

further supporting that the OPLS-DA model was significant. Most of the blue dots in

the scatterplots were located above the red dots, indicating good independence of the

OPLS-DA model.

In the following analysis, metabolites with the same trend on consecutive days,

$|log_2 FC| \geq 20$ (except for $SD = 0$ to remove the systematic error of the instrument)

or ranking top 10 were screened.