Cure or Curse? Simulation Indicates that Microbes Proliferate under Disinfection Measures in the Space Station

Shuaishuai Li[1,2,5], Xiaolei Liu[1,2], Letian Chen[1,2], Hong Liu[1,2,3,4], Dawei Hu[1,2,3,4,*]

[1] School of Biological Science and Medical Engineering, Beihang University, Beijing 100083, China

[2] Institute of Environmental Biology and Life Support Technology, Beihang University, Beijing 100083, China

[3] International Joint Research Center of Aerospace Biotechnology and Medical Engineering, Beijing 100083, China

[4] State Key Laboratory of Virtual Reality Technology and Systems, School of Computer Science and Engineering, Beihang University, Beijing 100083, China

[5] Department of Biology, McGill University, Montréal, Québec, Canada H3A 1B1

* Corresponding author.

E-mail address: dwhu@163.com (D. Hu)

Postal address: 37 Xueyuan Road, Haidian District, Beijing, China

Abstract

The long-term stay on the space station requirement poses a knotty issue about the stable hardware and reliable guarantee of astronauts' health. Despite regular cleaning with wipes, microbes still proliferate in the space station. This study distilled three key factors—low-dose ionizing radiation, dilution and quaternary ammonium compound from the scenario. Species abundance and differential metabolites were applied to test their sole/combined influences on the three-simple-species microbial community succession. Mechanisms were built in mathematical models to generate the structural and behavior similarity with the phenomena. The results showed that LDIR, dilution and QAC, solely or jointly, could contribute to microbial proliferation. The synergy of

disturbances might convert them from harmful factors to rewarding ones (e.g. the

transformation of QAC into nutrients under LDIR), leading to this "Ecological

Surprise". These results shed light on the mechanisms driving microbial community

succession in the space station and highlighted the need for tailored biocontrol

strategies in the specific environment.

Keywords: microbial community succession, disturbance, low-dose ionizing radiation,

dilution, quaternary ammonium compound(s).

Abbreviation:

LDIR: low-dose ionizing radiation; QAC: quaternary ammonium compound(s)

1. Introduction

Long-term residence on the space station needs stable hardware and guaranteed

health status of astronauts (Brooks et al., 2014; Rosado et al., 2010; Sun et al., 2016;

Yang et al., 2020), which is threatened by proliferative microbes on the inner surface of

the space station (Alekhova et al., 2005). When the environmental factors are the same,

the species diversity on the space station is much higher than that on the ground (Lang

et al., 2017; Novikova et al., 2006), showing the phenomenon of the paradox of the

plankton (G. E. Hutchinson, 1961; Yang et al., 2020). As astronauts and goods are the

main sources of microbes in this closed environment (Lax et al., 2014; Singh et al.,

2018), periodic cleaning is an actual plan for it. When carrying out the cleaning task,

astronauts use wipes containing 0.1% quaternary ammonium compounds (QAC) (Singh

et al., 2018) to wipe down the internal surface of the cabin under the low-dose ionizing

radiation (LDIR) from the universe. However, the removal of microbes seems to be

only effective for a short period of time and it is not complete. The microbes would

grow back after some time and the microbial community structure varies by time and

location (Lang et al., 2017; Singh et al., 2018).

Understanding community succession mechanisms is always one of the core

targets of ecology, especially microbial ecology(Powell et al., 2015; Stegen et al., 2013;

Zhou & Ning, 2017). Disturbance, as a largely independent area of ecology, is the event

that influences populations, resources or the physical environment in community

succession (Pickett & White, 1985). Microbial removal by wiping, translated into the

perspective of microbial community succession, is caused by abiotic environmental

disturbance. This operation results in microscopic mass non-specific or specific death,

macroscopic physical dilution and chemical disinfection events. Organisms are

sensitive to the abiotic environment and community succession occurs when they are

disturbed(Baudoin et al., 2008; Dangles & Chauvet, 2003; Tolkkinen et al., 2013).

Although violent environmental degradation might lead to species extinction, some

species could be favored under that circumstance because of the interactions necessary

for survival in multi-species communities (Abreu et al., 2019, 2020). For example,

evenly increasing mortality favors the faster grower, reversing the winner from the

slower to the faster one in the competition (Hastings, 2013; Stewart & Levin, 1973).

Nevertheless, current oversimplified models might lose some key information about the

succession. Research to date has not yet determined how disturbances would work in

resource-explicit models and the influence of selective disturbances.

In this study, three different types of disturbances, LDIR, dilution and QAC, were

distilled from the scene of wiping the inner surface of the space station. Experiments

with the simple three-species microbial community were carried out to demonstrate the

species absolute abundance and differential metabolites in the community succession

under different combinations and gradients of these disturbances. Results showed that

trends of community succession varied among different treatments and all types of

disturbances could cause the proliferation of the community by the relief of competition

or property transformation of specific chemicals. Afterward, some measures were

proposed for microbe control in aerospace. Also, it is hoped that this research will

contribute to a deeper understanding of community succession formations and

mechanisms under different types of disturbances.

2. Materials and methods

2.1 Strains, culture and radiative environment simulation

Three laboratory-stored strains, *Bacillus subtilis* (CMCC(B) 63501), *Escherichia*

coli (CMCC(B) 44102) and *Pseudomonas aeruginosa* (CMCC(B) 10104) were

originally obtained from CMCC (China Medical Culture Collection). These three

species were reported many times from microbial surveys in ISS, spacecrafts as well as

related studies(Crabbé Aurélie et al., 2011; Singh et al., 2018; Urbaniak et al., 2018;

Yang et al., 2021; Zea et al., 2016) and it was quite easy to distinguish them by genetic

relationship.

To simulate the nutrient-deficient environment on the inner surface of the space

station, the overnight cultures of the above strains were inoculated in the 10ml modified

minimal R2A broth (Yang et al., 2021) with a total density of 1×10^6/ml calculated by

direct counting under microscopy (Z. Song et al., 2015). The composition of modified

minimal R2A was shown in Table S1. In the mixed culture, the initial ratio of three

species was 1:1:1.

The LDIR environmental simulator (Fig. 1-A) was a refitted artificial climate

chamber that simulated the LDIR environment in the space station. The modification

included three main parts: a rotating wheel for placing the cultures, two X-ray tubes for

alternatively ionizing the air inside to plasma, and a protective door for isolating the

radiation. Two X-ray tubes (XCH201, Dandong Shenbo Electronics Co. Ltd, China)

were installed at the inside top of the chamber and run at 20kV, 0.2mA, alternately. To

ensure that each culture is exposed to the same radiation dose, six boxes are affixed to

the rotary wheel running at a constant angular velocity. These boxes contained plates

with LDIR-treated cultures, while the control cultures without LDIR treatment were

placed outside the radiation-exposed area but in the same climate chamber. The

protective door was sealed to maintain a constant temperature of 30°C, and normal

pressure during operation. The 3D geometric configuration of LDIR environmental

simulator was accurately established using SolidWorks based on its int ernal

dimensions, as shown in Fig. 1-B.

2.2 Periodic sampling and successive incubation

During a seven-day experiment cycle, 1ml samples were taken from cultures at the

same time on each day. Then the residual parts were differently treated for successive

incubation by distinct groups in Table 1. Each group contains three replicates. In each

1ml sample, 700μl was centrifuged for untargeted metabolomics test with its

supernatant, and two portions of 90μl bacterial fluid were used to depict species

abundance curves.

The composition of QAC was shown in Table 2.

2.3 Species abundance curve depicted by qPCR

Two portions of 90μl bacterial fluid above were applied to depict curves for

Bacillus subtilis (Kirchhoff & Cypionka, 2017; Tut et al., 2021; Xie et al., 2016) by

direct qPCR (Sup. A) and *Escherichia coli* as well as *Pseudomonas aeruginosa* by

PMA-qPCR, separately.

For the PMA-treated one, 10μl 500mmol/L PMAxx (Cat. 40069, Biotium Inc.,

Hayward, CA, USA) was added into the sample under dark conditions for forming the

50mmol/L final concentration. Then the mixed solution was incubated at 20°C for 40

min in the dark on a vortex with mild continuous shaking (100rpm) followed by the

change of tube. Subsequently, the tube was exposure to a 120W blue LED light at 4°C

for 30 min (Agustí et al., 2018; Codony et al., 2020; Emerson et al., 2017; Fittipaldi et

al., 2012; García-Fontana et al., 2016; Nocker et al., 2007) (Sup. B).

DNA was subsequently extracted from the solution above (with PMA or not) by

direct boiling with glass beads (Li et al., 2023). 0.1g 0.1mm glass beads (Cat. 11079101,

BioSpec Products, Inc., Bartlesville, OK, USA) were added into the tube containing

100μl resuspending solution before the heating procedure. After centrifugation, the

supernatant is DNA solution.

The primers, their qPCR standard curves, their amplicons and the composition of

qPCR reaction buffer were the same as our previous study (Li et al., 2023). 2×SYBR

Green PCRmix (Cat. SR1110, Solarbio Science & Technology Co. Ltd., Beijing, China)

was used to provide PCR Buffer, $MgCl_2$, dNTPs, Hs Taq DNA Polymerase, SYBR

Green I and so on. Primers and standards of their amplicons (Banihashemi et al., 2012;

Baymiev et al., 2020; Contreras et al., 2011; Emerson et al., 2017; Fittipaldi et al., 2012;

Loozen et al., 2011; Martin et al., 2013; Schnetzinger et al., 2013) were commercially

synthesized by Biomed Gene Technology Co., Ltd., Beijing, China. The reaction was

performed using Gentier 96C real-time PCR system (Tianlong Science and Technology

Co. Ltd., Xi'an, China). The qPCR amplification program included: pre-denaturation

at 95°C for 10 min with a heating rate of 6°C/s; then 40 cycles of denaturation at 95°C

for 10 sec with a temperature change rate of 5°C/s, annealing at 60°C for 20 sec with a

cooling rate 5°C/s and elongation at 72°C for 30 sec with heating rate 6°C/s.

2.4 Untargeted metabolomics test

Metabolites were quantified using ultra-performance liquid chromatography-triple

quadrupole time-of-flight mass spectrometry (UPLC-MS) (Waters Corp., Milford, MA,

USA). Raw data obtained from UPLC-MS were quantified in Progenesis QI (Waters

Corp.). Significantly differential metabolites analysis was performed using BMKCloud

(www.biocloud.net).

2.5 Mechanical modeling and computer simulation

The kinetic model of microbial community succession was established based on

generalized Lotka-Volterra (gLV) equations and hypothetical mechanisms of

disturbances. Parameters implying attributes in the community were embedded into the

model and generated through computer Monte-Carlo experiments. Afterward, a

corresponding simulation model with dde23 solver was established by S-Function of

MATLAB/Simulink (MathWorks, Inc., Natick, MA, USA) to confirm the proposed

hypothesis through behavior similarity between computer simulation and experimental

data.

3. Results

This study aimed to demonstrate the influence of three different types of

disturbances on microbial community succession. Experiments were taken to depict the

species absolute abundance and differential metabolites during the succession. The

following mechanical models and computer simulation were proposed to explain the

mechanisms of phenomena in these experiments.

3.1 qPCR results

The overview of all conditions was shown in Fig. 2. As seen in this figure, distinct

succession could be obtained by different treatments. Two-way repeated measured

ANOVA was applied using R (4.0.2) to calculate the p-values to test if the curves

showed a significant difference between the specific experimental group and the control

group in the overall experimental period, where the interaction effect of treatment and

time was considered first. If the interaction effect was not significant, the main effect

of treatment would be applied, where the p-value was marked as "p_m".

As shown in Fig. 3, in the control group, the total biomass and the absolute

abundance of *E. coli* reached the highest points on the sixth day, when the density for

both of them was about 1.5×10^{10}/ml.

When exposed to LDIR (Fig. 4), the microbial community thrived after a period

of "hibernation". Compared with the control group, the total biomass rose on 5-7th days

and still peaked on the 6th day, approximately reaching 3.5×10^{10}/ml. The majority

was still *E. coli*. The abundance of *B. subtilis* showed the trend of damped periodic

oscillations.

Dilution with gradients from 10% to 0.1‰ per day had no obvious different

influence on the microbial community succession, as shown in Fig. S1 & S2. Taking 1‰

as an example (Fig. 5), compared with the control group, the total biomass and the

abundance of *E. coli* decreased on 4-6th days, exhibiting the trend from oscillation to

robustness. The abundance of *B. subtilis* significantly fell ($p_m = 1 \times 10^{-5}$), while that

of *P. aeruginosa* rose dramatically ($p = 0.004$).

Imposing 100% and 10% stock solution of QAC could rapidly bring the total

biomass down to below the initial density, when only *P. aeruginosa* could survive

below the initial density under this condition (Fig. 2). However, 1% and 1‰ stock

solution of QAC did not have the similar impact (Fig. 6). When treated with 1% stock

solution of QAC, compared with the control group, the peak of total biomass moved

back to 4th day, having no significant difference with the peak of the former. *E. coli* had

a great contribution to the total biomass on the 1-4th days, followed by the sharp drop

and extinction on the 6th day. Meanwhile, the abundance of *P. aeruginosa* increased

from the 5th day and peaked on the 6th day. In the group of 1‰ treatment, the total

biomass exceeded the control group on the 7th day before the lag on the 4-6th days,

presenting the trend of rise with fluctuations. The abundance of *E. coli* fluctuated

continuously, and that of *P. aeruginosa* increased steadily.

When diluted with a 1‰ ratio per day, the total biomass ($p_m = 4 \times 10^{-4}$) and the

abundance of *E. coli* ($p_m = 4 \times 10^{-4}$) significantly rose in the whole period of the

LDIR group, compared with the group without LDIR (Fig. S3).

When exposed under LDIR, compared with the group without QAC, the total

biomass of the 1% stock solution of QAC group increased on the 2-4th days but

decreased on the 5-7th days, in which the abundance of *E. coli* went extinct quickly on

the 5th day and that of *P. aeruginosa* peaked on 6th day (Fig. S4-A). The total biomass

of the 1‰ stock solution of QAC group rose in fluctuations, exceeding the group

without QAC on the 4th and 7th days but lagged on the 5-6th days, when the latter group

was experiencing proliferation (Fig. S4-B). The majority was still *E. coli* but the

magnitude of *P. aeruginosa* also improved.

When treated with 1% stock solution of QAC, the total biomass of the LDIR group

went up in the whole period compared with the group without LDIR except on the 4th

day, when the latter group peaked. The abundance of *E. coli* increased on 1-3rd days,

then dropped on 4th day but this species still survived later. The abundance of

P. aeruginosa rose during the whole period and still peaked on the 6th day (Fig. S5-A).

When treated with 1‰ stock solution of QAC, the total biomass ($p = 0.003$) and the

abundance of *E. coli* ($p = 0.02$) of the LDIR group dramatically rocketed with growing

gaps in the whole period compared with the group without LDIR (Fig. S5-B). The

abundance of *P. aeruginosa* shrank a little but the difference was not significant ($p >$

0.05).

When LDIR, 1‰ dilution and 1% stock solution of QAC were imposed, compared

with the control group (Fig. S6-A), *E. coli* went extinct quickly on the 2^{nd} day. However,

the abundance of *P. aeruginosa* substantially increased ($p = 3 \times 10^{-6}$). When LDIR,

1‰ dilution and 1‰ stock solution of QAC were imposed, compared with the control

group (Fig. S6-B), the total biomass increased on 1-4^{th} and 7^{th} days, showing a trend of

fluctuation. The abundance of *P. aeruginosa* markedly increased ($p = 3 \times 10^{-4}$).

3.2 Untargeted metabolomics test results

The samples involved in metabolite analysis were divided into 24 series according

to biological repeat, and the symbol corresponding to the experimental condition of

each group was shown in Table 3. The quality control, OPLS-DA as well as its validity

test, OPLS-DA permutation test and screening of metabolites were done before analysis

(Sup. C).

Compared with the control group, the concentration of p-coumaroyl agmatine in

the group exposed under LDIR between 4-6^{th} days dropped significantly ($p < 0.05$,

Series B&M, C&N, D&O. Fig. S7). On 1^{st} and 4-6^{th} days, most of the metabolic

pathways were inhibited, including polysaccharide metabolism, amino acid metabolism,

vitamin metabolism, xenobiotic degradation and metabolism, as well as antibiotic

biosynthesis (Series A&L, B&M, C&N, D&O; Fig. S8).

Compared with the control group, in the group diluted with 1‰ ratio per day, N-

acetylmannosamine was enriched on the 4^{th} day (Series B&F), which was markedly

correlated with the rise of the abundance of *P. aeruginosa* ($p < 0.01$, Fig. S9). Pathways

inhibited dominated on 4^{th} day in the microbial community, including nucleotide

metabolism, sugar degradation, antibiotic synthesis, and aminobenzoate degradation

(Fig. S10).

When treated with 1% stock solution of QAC, N-acetylmannosamine was more

abundant on 4^{th} day compared with the control group (Series B&G), which was also

correlated with the rise of the abundance of *E. coli* ($p < 0.05$, Fig. S11). Meanwhile,

staurosporine and tetracycline biosynthesis were active. When treated with 1‰ stock

solution of QAC, echitovenine was rarer on the 7th day compared with the control group

(Series E&K), which was also correlated with the increase of the abundance of

P. aeruginosa ($p < 0.01$, Fig. S12). On the same day, aminobenzoate degradation as

well as biosynthesis of alkaloids (derived from histidine and purine) were inhibited (Fig.

S13).

When both LDIR and 1% stock solution of QAC were imposed, compared with

the group under 1% stock solution of QAC, alkaloid biosynthesis was inhibited (Fig.

S14), and some nitrogenous long-chain organic matters were enriched on 5-6th days

(Series H&S, I&T), including tetradecenoyl carnitine and R-palmitoyl-(2-methyl)

ethanolamide, ranging from $1 \times 10^{10} \sim 2 \times 10^{12}$ times; compared with the control

group, cytochrome P450 metabolism was active on 5th day (Series C&S). When LDIR,

1‰ dilution and 1% stock solution of QAC were imposed together, the concentration

of N-[(2R)-2-hydroxypropyl] hexadecanamide was 2×10^{12} times higher than the

control group on the 4th day (Series B&W). However, when the QAC concentration was

changed to 1‰ of the stock solution (Series B&V, B&X, E&W, J&V, K&W), the

phenomenon of enrichment of nitrogenous long-chain organic matters disappeared;

alternatively, cyanoamino acid metabolism, related to nitrogen metabolism (Ding et al.,

2022), was active. The concentration of albaflavenone sharply diminished ($p < 0.01$,

Fig. S15) on 4th day in the group with both LDIR and 1‰ stock solution of QAC,

compared with the control group (Series B&V). Aminobenzoate degradation on the 4th

day and antibiotic synthesis on the 7th day (Series K&W. Fig. S16) were restrained in

the group with both LDIR and 1‰ stock solution of QAC, compared with 1‰ stock

solution of QAC group. Under LDIR, when comparing 1% and 1‰ stock solution of

QAC groups, eicosanoids metabolism was active in the higher concentration one on the

4[th] day (Series R&V. Fig. S17).

3.3 Mechanical modeling and simulation results

3.3.1 State variables

The mechanical model was composed of six state variables in four types: *B.*

subtilis biomass (x_1), *E. coli* biomass (x_2), *P.aeruginosa b*iomass (x_3); amount of

substrate (*S*), amount of QAC (*Q*), and amount of QAC degradation products (*P*).

3.3.2 Rate equations

(1) growth rate

The growth rate without competition of species i could be expressed as:

288
$$v_{ai} = \mu_i \left(\frac{S}{K_s + S} \right) x_i \tag{3.1}$$

289　　　　where μ_i is the specific growth rate of species i, and K_s denotes substrate-

290　　dependent half-saturation constant for species i.

291　　　　(2) interspecific competition

292　　　　When there is no joint treatment of LDIR and QAC, the interspecific competition

293　　rate against species i could be written as follows:

294
$$v_{bi} = \sum_{j \neq i}^{3} \alpha_{ji} x_i x_j \tag{3.2}$$

295　　　　where α_{ji} denotes the competition inhibition coefficient from species j to

296　　species i.

297　　　　When the microbial community is treated with LDIR or QAC, the interspecific

298　　competition rate against species i could be written as follows:

299
$$v_{ci} = \sum_{j \neq i}^{3} \alpha_{ji} e^{-(c_{ji\_R}R + c_{ji\_Q}Q)} x_i x_j \tag{3.3}$$

300　　　　Where I is a constant, representing the LDIR coefficient; e is the Euler number;

301 c_{ji_R} and c_{ji_Q} express the interspecies cross-feeding response coefficient between

302 species j and species i caused by LDIR and QAC, respectively.

303 (3) intraspecific competition

304 When there is no joint treatment of LDIR and QAC, the intraspecific competition

305 rate of species i could be written as follows:

$$v_{di} = \beta_i x_i^2 \tag{3.4}$$

307 where β_i denotes the intraspecific competition inhibition coefficient of species i.

308 When the microbial community is treated with LDIR or QAC, the intraspecific

309 competition rate of species i could be written as follows:

$$v_{fi} = \beta_i e^{-(f_{i\_R}R + f_{i\_Q}Q)} x_i^2 \tag{3.5}$$

311 where, f_{i_R} and f_{i_Q} denote the intraspecies cross-feeding response coefficient

312 of species i caused by LDIR and QAC, respectively.

313 (4) metabolism

314 The metabolism rate of species i could be expressed as:

$$v_{gi} = \delta_i x_i \tag{3.6}$$

316 where δ_i is the metabolism coefficient of species i.

317 (5) QAC disinfection

$$v_{hi} = \varepsilon_i Q x_i \tag{3.7}$$

319 where ε_i is the QAC disinfection rate on species i.

320 (6) QAC degradation

$$v_{mi} = \gamma R Q \tag{3.8}$$

322 where γ is the first-order kinetic coefficient of QAC degradation rate.

323 (7) generation of QAC degradation products

$$v_{ni} = \omega v_{mi} \tag{3.9}$$

325 where ω is the calibration coefficient.

326   (8) consumption of substrate

$$v_S = \sum_{i=1}^{3} \varphi_i v_{ai} \tag{3.10}$$

328   where φ_i is the substrate consumption coefficient during the growth of species i.

329   (9) transient influence of Dilution and QAC

330   a. only dilution

$$\begin{cases} S = S_0, t = 1,2,\dots,7d \\ x_{i,t} = D x_{i,t-\Delta t} \end{cases} \tag{3.11}$$

332   where S_0 is the original substrate concentration of the R2A culture; $x_{i,t}$ is the

333   abundance of species i at t moment; Δt is a very short time interval; D is the

334   proportion of the culture just before dilution to the total volume.

335   b. only QAC

$$Q_t = Q_{t-\Delta t} + Q_{Add}, t = 1,2,\dots,7d \tag{3.12}$$

337   where Q_{Add} denotes the QAC added.

338   c. both dilution and QAC

$$\begin{cases} S = S_0, t = 1,2,\dots,7d \\ Q_t = D Q_{t-\Delta t} + Q_{Add}, t = 1,2,\dots,7d \\ x_{i,t} = D x_{i,t-\Delta t} \end{cases} \tag{3.13}$$

3.3.3 the overall mechanical model

Based on Eq(3.1)~(3.13), the overall mechanical model, i.e. a state space model

with six first-order nonlinear kinetic equations, could be formulated in a top-down

manner as follows:

(1) When there is no joint treatment of LDIR and QAC, the population dynamics

equation of species i is:

$\frac{dx_i}{dt} = v_{ai} - v_{bi} - v_{di} - v_{gi} - v_{hi} = \mu_i \left(\frac{S}{K_s+S} \right) x_i - \sum_{j\neq i}^{3} \alpha_{ji} x_i x_j - \beta_i x_i^2 - \delta_i x_i - \varepsilon_i Q x_i \tag{3.14}$

When there is joint treatment of LDIR and QAC, it is:

$$\frac{dx_i}{dt} = v_{ai} - v_{ci} - v_{fi} - v_{gi} - v_{hi} =$$

$$\mu_i \left(\frac{S}{K_s + S}\right) x_i - \sum_{j \neq i}^{3} \alpha_{ji} e^{-(c_{ji\_R}R + c_{ji\_Q}Q)} x_i x_j - \beta_i e^{-(f_{i\_R}R + f_{i\_Q}Q)} x_i^2 - \delta_i x_i - \varepsilon_i Q x_i \quad (3.15)$$

349   (2) The QAC degradation dynamics equation is:

$$\frac{dQ}{dt} = -v_{mi} = -\gamma R Q \qquad\qquad (3.16)$$

351   (3) The generation of QAC degradation production dynamics equation is:

$$\frac{dP}{dt} = v_{ni} = \omega \gamma R Q \qquad\qquad (3.17)$$

353   (4) The substrate change dynamics equation is:

$$\frac{dS}{dt} = v_{ni} - v_S = \omega \gamma R Q - \sum_{i=1}^{3} \varphi_i \mu_i \left(\frac{S}{K_s + S}\right) x_i \qquad (3.18)$$

3.3.4 computer simulation

The code of the mechanical model above (Sup. D) was written by S-Function in

MatLab/Simulink(R2019b, 9.7.0.1190202). The simulation experiments were

implemented for simulating the control group and treatment groups of LDIR, dilution,

QAC as well as their combinations, as shown in Fig. 7. The simulation results showed

that the mechanical model had considerable behavior similarity with experiments,

while the mathematical expression of the mechanical model matched the growth

process of microbes, indicating the structure similarity between them. Therefore, the

mechanical model was considered valid.

4. Discussion

In the present study, LDIR, dilution and QAC were extracted from the

environment on the inner surface of the space station as treatments, and the microbial

community succession under these uniformed different treatments was observed by

measuring species abundance and metabolites. Mechanical models were built to explain

the experiments, with which the computer simulation showed structural and behavioral

similarity. The overall mechanisms are shown in Fig. 8.

At the later stage, LDIR improved the total biomass, every species abundance and

their peaks, when the Hydra Effect happened. This phenomenon was shown in both

with/without LDIR comparison and with/without LDIR comparison under dilution (Fig.

4 & S3). The p-coumaroylagmatine could thicken cell walls, limit bacterial movement

and improve plant defense against pathogens(Häusler et al., 2014; Macoy et al., 2015),

thus its decrease on the 4-6th days (Series B&M, C&N, D&O) indicated that the

physiological activity of the microbial community was enhanced under LDIR. The

inhibited antibiotic synthesis pathway (Series B&M) suggested that external pressure

was reduced and interspecific competition was diminished. Therefore, the Hydra Effect

occurred under LDIR. This effect was more significant as the main effect in the

comparison of the group under LDIR and 1‰ dilution and only 1‰ dilution (Fig. S3).

Except in *P. aeruginosa*, dilution reshuffled all other components everyday and

undermined the interaction effect of treatment and time, where the enhancement effect

of LDIR emerged.

Dilution, by removing a certain part of biomass during serial transfers, was a non-

specific and highly controllable disturbance way, where the transient removal did not

depend on response traits in the microbial community (Gibbons et al., 2016). However,

under non-selective dilution, *P. aeruginosa* was selectively released from competitive

exclusion. Aminobenzoate was the signal of xenobiotic degradation (Patil et al., 2020;

Wang et al., 2020), thus its degradation inhibition on 4th day (Series B&F) suggested

that organisms tended not to consume energy storage material (Smitha et al., 2017) and

the competition in the community was mitigated. It could be inferred that many niches

would be released without distinction under dilution, when *P. aeruginosa* could

reproduce based on its merits beyond others' (Abreu et al., 2019) and occupy these

niches. This release happened only under dilution with all gradients; when both dilution

and LDIR were applied, *E.coli* took the majority again (Fig. S18-A).

QAC stood for specific disturbances, causing non-random removal according to

the microbe's response traits(Banitz et al., 2020; Calderón et al., 2018; Jurburg et al.,

2017; Srivastava & Vellend, 2005). Its effect also counted on the disturbance intensities

and frequencies. Different QAC concentrations had various influences on the microbial

community, from elimination to promotion. The Hydra Effect even happened under 1%

and 1‰ stock solution of QAC, where simple competition mitigation cannot explain.

Polymers of N-acetylmannosamine played a protective role in *E. coli* (Sellner et al.,

2021), thus their respective increments with dramatical correlation told the reason why

the species abundance of *E. coli* peaked in advance on the 4th day under 1% stock

solution (Series B&G). Besides, the active biosynthetic pathways of antibiotics and

cytokinins on the same day corroborated that microbes combated adversity by more

cell division and the total biomass also peaked. Echitovenine was a monoterpene

alkaloid with antimicrobial activity (Mohammed et al., 2021), whose attenuation could

interpret the growth of *P. aeruginosa* on the 7th day under 1‰ stock solution (Series

E&K), given their notable correlation. QAC would periodically selectively remove a

portion of individuals in the community, leading the way of community succession to

the direction of dominance by species that can tolerate the environment, which could

be pronounced in comparisons across gradients (Figure S18-B&C). With low-

concentration QAC treatment, the community had adapted to the environment and

utilized it. QAC, was a series of long-chain organics that could be used by microbes,

especially in microbial co-metabolism by *Pseudomonas* (Harris & Knowles, 1983;

Leadbetter & Foster, 1960; Zhang et al., 2015), which was widely and systemically

utilized in treating wastewater (biological nitrogen removal). On the oligotrophic

surface of the space station, the carbon sources were mostly wiped down, when co-

metabolism might allow microbes to survive with remaining nitrogen sources.

Meanwhile, the inhibition of aminobenzoate degradation pathway also flanked the

corresponding decrease of the consumption of energy storage in the community.

The upsurge of some nitrogenous long-chain organic matters under both LDIR and

QAC, compared with only QAC or the control group, strongly implies that QAC may

undergo radiolysis under LDIR. Several studies (Baidak & LaVerne, 2010; Dhiman &

LaVerne, 2013; Nogami et al., 1996; Pillay, 1986) implied that radiolysis would happen

on substrates containing QAC, especially under water acceleration. 1% stock solution

of QAC is substantially transformed from the toxic bactericide to organic nitrogenous

long-chain nutrients, including tetradecenoyl carnitine, R-palmitoyl-(2-methyl)

ethanolamide and N-[(2R)-2-hydroxypropyl] hexadecanamide with significant

differences between Series B&W, H&S, I&T. This might accelerate the microbial co-

metabolism process. Cytochrome P450-related enzymes prefer long-chain fatty acids

with 15-16 carbon atoms (Lewis & Wiseman, 2005), thus its activation might be related

to added long-chain QAC utilization. Inhibited alkaloid biosynthesis (Series H&S, I&T)

and ameliorated cytochrome P450 metabolism (Series C&S) also shows that the

community could adapt to and consume QAC under LDIR. Nevertheless, when the

QAC concentration is 1‰, significant differences are no longer detected for organic

nitrogenous long-chain nutrients (Series B&V, B&X, E&W, J&V, K&W), suggesting

that the community can entirely co-metabolize QAC in this situation. The activation of

cyanoamino acid metabolism (Series B&X), inhibition of aminobenzoate degradation

(Series J&V) as well as inhibition of antibiotic synthesis (Series K&W) also reconfirm

the speculation of utilization of exogenous QAC as the source of nitrogen and less

pressure of the community caused by LDIR. Comparing different concentration of

QAC under LDIR (Series V&R), the active eicosanoids metabolism in the higher

concentration one shows that the microbes might transform QAC into long-chain fatty

acid substances for consuming.

The proliferative microbes on the inner surface of the space station (Alekhova et

al., 2005) have been threatening its normal operation, which is a serious problem and

could be shifted to the issue of the effect of disturbance on community succession.

Disturbance is acknowledged as a key factor of biodiversity via affecting eco-processes

(Banitz et al., 2020). However, the mechanisms behind the observed community

diversity under disturbance need to be further clarified (Fox, 2013; Shea et al., 2004).

First, few quantitative studies on disturbances have been reported, while more relative

studies focus on the qualitative aspect — this is due to both the difficulty in defining

the attribute of frequency, intensity, as well as extent (Shade et al., 2012), and the wide

variety of their characteristics and ecological impacts (Fraterrigo & Rusak, 2008).

Second, few studies are explicitly based on the time course of microbial community

composition under disturbance; instead, many solely studies are on the sensitivity of

composition at a given moment (Allison & Martiny, 2008), which is attributed to the

inertial thinking in community ecology and research tools in microbiology. Our

previous study noticed the benefit of sole LDIR for the microbial community (Yang et

al., 2021). Abreu et al. (Abreu et al., 2019, 2020) expanded the effect of dilution on

multispecies community composition, providing an example for quantifying

disturbances with the time course. Nonetheless, in their research, when distinguishing

disturbances by pulsed versus sustained, pulsed ones like dilution are not applicable for

being directly integrated into the typical LV equation. Finally, studies with similar

gradient-based methods on combined multiple factors with synergy, are still at an early

stage. Most studies still concentrate on a single factor, and only a few have noticed the

combined influence of multiple factors (Banitz et al., 2020; Zeng et al., 2020). In

contrast, a large number of phenomena in nature are caused by the complex interaction

of the latter condition (Elmqvist et al., 2003; Martínez-Ramos et al., 2016), including

"ecological surprises" such as drastic shifts in community compositions and

biodiversity (Banitz et al., 2020; Filbee-Dexter et al., 2017; Newman, 2019).

In this study, "ecological surprise" occurred when human expectations deviated

from the observed community succession behavior (Doak et al., 2008; Filbee-Dexter et

al., 2017). And these unexpected dynamics arose from human management and

disturbances (Doak et al., 2008). Dilution (wiping) and QAC were recognized as

efficient measures to take control of microbes (Sattar et al., 2015; Zhang et al., 2015)

and applied in the space station (Singh et al., 2018). However, given the results above,

the synergy emerged between LDIR and QAC, thus low-concentration QAC was

transformed from the bactericide to nitrogen-rich long-chain nutrients. Selectively

released *P. aeruginosa* under suppression after dilution would take these nutrients to

thrive, changing the community structure and dominant species. That could be a

possible reason for the phenomenon in the space station, and also gave a potential

mechanism for "ecological surprise" under synergistical disturbances. It reminded us

that tailored biocontrol strategies ought to be considered for the specific environment.

Sometimes, the control measures were taken for granted that they were "disturbances"

affecting the targets without the consideration of their own properties and changes. It

cannot be neglected that general strategies might not function under certain conditions.

Given the current microbial prevention strategies have some disadvantages, future

improvements can be adopted. Concerning the low concentration of QAC, except for

co-metabolism and radiolysis of QAC mentioned above, the pre-package of wipes can

also cause it by both reduced antimicrobial activity of QAC with a long-term soaking

(Bloß et al., 2010, 2010; Slopek et al., 2011) and the chemical combination with the

wipe fibre (Dharan et al., 1999; X. Song et al., 2019). The target surface properties also

influence the disinfection (X. Song et al., 2019). More organic contamination on (dirtier)

the surface, lower the disinfection effect (Williams et al., 2007). Moreover, radiolysis

could happen just after the spacecraft entered the outer space. In terms of dilution, its

rate was limited on the order of 1% even using two wipes successively (Gold & Hitchins,

2013), which is not enough given the results in this study. Furthermore, "wiping" itself

could lead to cross-contamination of microorganisms in different locations by

mechanical stimulation (Ramm et al., 2015; Siani et al., 2011; Williams et al., 2007).

To attenuate these effects, several measures could be considered. Such as augmenting

local wiping time, wiping with mechanical equipment to restraint power variation

(Sattar et al., 2015; Williams et al., 2007), special antibacterial treatment on the surface

requiring microbial prevention, applying different wipe materials harder to react with

QAC, adjusting the formulation of QAC for a lower degradation rate (Pillay, 1986;

Zhang et al., 2015), avoiding surface exposure to radiation after wiping.

Despite these promising results, questions still remain. Further work is needed to

develop reliable experimental and analytical methods (e.g. based on the individual-

based model) (Li et al., 2022) for testing community succession considering the spatial

factor. The liquid culture implemented in this study aborted it because of current

biotechnology limitations on detecting microbial individuals. Additional research based

on more methods in bioinformatics is needed to better understand the successive

changes in microbial behaviors under disturbances. This study focused on the level of

community succession. Hence, there is abundant room for further progress in

determining the evolutionary characteristics changes of the species facing disturbances

in long-term experiments.

5. Conclusion

In this study, the intricate relationship between disturbance and the microbial community succession in the space station was elucidated. The proliferation under LDIR, the selective species release under dilution and complex responses of the community to QAC concentration were observed in experiments and their similarity was tested in modeling. The synergy of combined disturbances, QAC and LDIR, led to the transformation of them from harmful factors to supportive ones. Therefore, it is necessary to consider synergistic effects and to apply tailored strategies when addressing the microbial proliferation challenge as well as other similar circumstances.

Fundings

This study was financially supported by Funds for International Cooperation and Exchange of the National Natural Science Foundation of China (32261133528).

Acknowledgment

We sincerely acknowledge the assistance from Chuanmeng Cai, Dengbo Chen, and Yong Feng in the long and non-stop periodic experiment, and the assistance from Yong Feng for depicting Fig.1-B.

References

Abreu, C. I., Andersen Woltz, V. L., Friedman, J., & Gore, J. (2020). Microbial communities display
alternative stable states in a fluctuating environment. *PLOS Computational Biology*, *16*(5),
e1007934. https://doi.org/10.1371/journal.pcbi.1007934

Abreu, C. I., Friedman, J., Andersen Woltz, V. L., & Gore, J. (2019). Mortality causes universal
changes in microbial community composition. *Nature Communications*, *10*(1), 2120.
https://doi.org/10.1038/s41467-019-09925-0

Agustí, G., Fittipaldi, M., & Codony, F. (2018). Optimization of a Viability PCR Method for the
Detection of Listeria monocytogenes in Food Samples. *Current Microbiology*, *75*(6), 779–
785. https://doi.org/10.1007/s00284-018-1448-6

Alekhova, T. A., Aleksandrova, A. A., Novozhilova, T. Y., Lysak, L. V., Zagustina, N. A., &
Bezborodov, A. M. (2005). Monitoring of Microbial Degraders in Manned Space Stations.
Applied Biochemistry and Microbiology, *41*(4), 382–389. https://doi.org/10.1007/s10438-
005-0065-x

Allison, S. D., & Martiny, J. B. H. (2008). Resistance, resilience, and redundancy in microbial
communities. *Proceedings of the National Academy of Sciences*, *105*(Supplement 1), 11512–
11519. https://doi.org/10.1073/pnas.0801925105

Baidak, A., & LaVerne, J. A. (2010). Radiation-induced decomposition of anion exchange resins.
Journal   of   Nuclear   Materials,   *407*(3),   211–219.
https://doi.org/10.1016/j.jnucmat.2010.10.025

Banihashemi, A., Dyke, M. I., & Huck, P. M. (2012). Long-amplicon propidium monoazide-PCR
enumeration assay to detect viable Campylobacter and Salmonella. *Journal of Applied
Microbiology*, *113*(4), 863–873. https://doi.org/10.1111/j.1365-2672.2012.05382.x

Banitz, T., Chatzinotas, A., & Worrich, A. (2020). Prospects for Integrating Disturbances,
Biodiversity and Ecosystem Functioning Using Microbial Systems. *Frontiers in Ecology and
Evolution*, *8*. https://www.frontiersin.org/articles/10.3389/fevo.2020.00021

Baudoin, J. M., Guérold, F., Felten, V., Chauvet, E., Wagner, P., & Rousselle, P. (2008). Elevated
Aluminium Concentration in Acidified Headwater Streams Lowers Aquatic Hyphomycete
Diversity and Impairs Leaf-Litter Breakdown. *Microbial Ecology*, *56*(2), 260–269.
https://doi.org/10.1007/s00248-007-9344-9

Baymiev, An. Kh., Baymiev, Al. Kh., Kuluev, B. R., Shvets, K. Yu., Yamidanov, R. S., Matniyazov,
R. T., Chemeris, D. A., Zubov, V. V., Alekseev, Ya. I., Mavzyutov, A. R., Ivanenkov, Ya. A.,
& Chemeris, A. V. (2020). Modern Approaches to Differentiation of Live and Dead Bacteria
Using Selective Amplification of Nucleic Acids. *Microbiology*, *89*(1), 13–27.
https://doi.org/10.1134/S0026261720010038

Bloß, R., Meyer, S., & Kampf, G. (2010). Adsorption of active ingredients of surface disinfectants
depends on the type of fabric used for surface treatment. *Journal of Hospital Infection*, *75*(1),
56–61. https://doi.org/10.1016/j.jhin.2009.11.027

Brooks, B., Firek, B. A., Miller, C. S., Sharon, I., Thomas, B. C., Baker, R., Morowitz, M. J., &
Banfield, J. F. (2014). Microbes in the neonatal intensive care unit resemble those found in
the gut of premature infants. *Microbiome*, *2*(1), 1. https://doi.org/10.1186/2049-2618-2-1

Calderón, K., Philippot, L., Bizouard, F., Breuil, M.-C., Bru, D., & Spor, A. (2018). Compounded
Disturbance Chronology Modulates the Resilience of Soil Microbial Communities and N-
Cycle Related Functions. *Frontiers in Microbiology*, *9*, 2721.
https://doi.org/10.3389/fmicb.2018.02721

Codony, F., Dinh-Thanh, M., & Agustí, G. (2020). Key Factors for Removing Bias in Viability PCR-
Based Methods: A Review. *Current Microbiology*, *77*(4), 682–687.
https://doi.org/10.1007/s00284-019-01829-y

Contreras, P. J., Urrutia, H., Sossa, K., & Nocker, A. (2011). Effect of PCR amplicon length on
suppressing signals from membrane-compromised cells by propidium monoazide treatment.
Journal of Microbiological Methods, *87*(1), 89–95.
https://doi.org/10.1016/j.mimet.2011.07.016

Crabbé Aurélie, Schurr Michael J., Monsieurs Pieter, Morici Lisa, Schurr Jill, Wilson James W., Ott
C. Mark, Tsaprailis George, Pierson Duane L., Stefanyshyn-Piper Heidi, & Nickerson Cheryl
A. (2011). Transcriptional and Proteomic Responses of Pseudomonas aeruginosa PAO1 to
Spaceflight Conditions Involve Hfq Regulation and Reveal a Role for Oxygen. *Applied and*
Environmental Microbiology, *77*(4), 1221–1230. https://doi.org/10.1128/AEM.01582-10

Dangles, O., & Chauvet, E. (2003). Effects of stream acidification on fungal biomass in decaying
beech leaves and leaf palatability. *Water Research*, *37*(3), 533–538.
https://doi.org/10.1016/S0043-1354(02)00359-7

Dharan, S., Mourouga, P., Copin, P., Bessmer, G., Tschanz, B., & Pittet, D. (1999). Routine
disinfection of patients' environmental surfaces. Myth or reality? *Journal of Hospital*
Infection, *42*(2), 113–117. https://doi.org/10.1053/jhin.1999.0567

Dhiman, S. B., & LaVerne, J. A. (2013). Radiolysis of simple quaternary ammonium salt
components of Amberlite resin. *Journal of Nuclear Materials*, *436*(1–3), 8–13.
https://doi.org/10.1016/j.jnucmat.2013.01.294

Ding, C., Wang, S., Li, J., & Wang, Z. (2022). Transcriptomic analysis reveals the mechanism of
host growth promotion by endophytic fungus of Rumex gmelinii Turcz. *Archives of*
Microbiology, *204*(7), 1–13.

Doak, D. F., Estes, J. A., Halpern, B. S., Jacob, U., Lindberg, D. R., Lovvorn, J., Monson, D. H.,
Tinker, M. T., Williams, T. M., Wootton, J. T., & others. (2008). Understanding and predicting
ecological dynamics: Are major surprises inevitable. *Ecology*, *89*(4), 952–961.

Elmqvist, T., Folke, C., Nyström, M., Peterson, G., Bengtsson, J., Walker, B., & Norberg, J. (2003).
Response diversity, ecosystem change, and resilience. *Frontiers in Ecology and the*
Environment, *1*(9), 488–494. https://doi.org/10.1890/1540-
9295(2003)001[0488:RDECAR]2.0.CO;2

Emerson, J. B., Adams, R. I., Román, C. M. B., Brooks, B., Coil, D. A., Dahlhausen, K., Ganz, H.
H., Hartmann, E. M., Hsu, T., Justice, N. B., Paulino-Lima, I. G., Luongo, J. C.,
Lymperopoulou, D. S., Gomez-Silvan, C., Rothschild-Mancinelli, B., Balk, M., Huttenhower,
C., Nocker, A., Vaishampayan, P., & Rothschild, L. J. (2017). Schrödinger's microbes: Tools
for distinguishing the living from the dead in microbial ecosystems. *Microbiome*, *5*(1), 86.
https://doi.org/10.1186/s40168-017-0285-3

Filbee-Dexter, K., Pittman, J., Haig, H. A., Alexander, S. M., Symons, C. C., & Burke, M. J. (2017).
Ecological surprise: Concept, synthesis, and social dimensions. *Ecosphere*, *8*(12), e02005.
https://doi.org/10.1002/ecs2.2005

Fittipaldi, M., Nocker, A., & Codony, F. (2012). Progress in understanding preferential detection of
live cells using viability dyes in combination with DNA amplification. *Journal of*
Microbiological Methods, *91*(2), 276–289. https://doi.org/10.1016/j.mimet.2012.08.007

Fleischmann, S., Robben, C., Alter, T., Rossmanith, P., & Mester, P. (2021). How to Evaluate Non-
Growing Cells—Current Strategies for Determining Antimicrobial Resistance of VBNC
Bacteria. *Antibiotics*, *10*(2), 115. https://doi.org/10.3390/antibiotics10020115

Fox, J. W. (2013). The intermediate disturbance hypothesis is broadly defined, substantive issues
are key: A reply to Sheil and Burslem. *Trends in Ecology & Evolution*, *28*(10), 572–573.
https://doi.org/10.1016/j.tree.2013.07.008

Fraterrigo, J. M., & Rusak, J. A. (2008). Disturbance-driven changes in the variability of ecological
patterns and processes. *Ecology Letters*, *11*(7), 756–770. https://doi.org/10.1111/j.1461-
0248.2008.01191.x

G. E. Hutchinson. (1961). The Paradox of the Plankton. *The American Naturalist*, *95*(882), 137–
145.

García-Fontana, C., Narváez-Reinaldo, J. J., Castillo, F., González-López, J., Luque, I., &
Manzanera, M. (2016). A New Physiological Role for the DNA Molecule as a Protector
against Drying Stress in Desiccation-Tolerant Microorganisms. *Frontiers in Microbiology*, *7*.
https://doi.org/10.3389/fmicb.2016.02066

Gibbons, S. M., Scholz, M., Hutchison, A. L., Dinner, A. R., Gilbert, J. A., & Coleman, M. L. (2016).
Disturbance Regimes Predictably Alter Diversity in an Ecologically Complex Bacterial
System. *mBio*, *7*(6). https://doi.org/10.1128/mBio.01372-16

Gold, K. M., & Hitchins, V. M. (2013). Cleaning assessment of disinfectant cleaning wipes on an
external surface of a medical device contaminated with artificial blood or *Streptococcus*
pneumoniae. *American Journal of Infection Control*, *41*(10), 901–907.
https://doi.org/10.1016/j.ajic.2013.01.029

Harris, R., & Knowles, C. J. (1983). Isolation and Growth of a Pseudomonas Species that Utilizes
Cyanide as a Source of Nitrogen. In *Microbiology* (Vol. 129, Issue 4, pp. 1005–1011).
Microbiology Society. https://doi.org/10.1099/00221287-129-4-1005

Hastings, A. (2013). *Population biology: Concepts and models*. Springer Science & Business Media.

Häusler, R. E., Ludewig, F., & Krueger, S. (2014). Amino acids – A life between metabolism and signaling. *Plant Science*, *229*, 225–237. https://doi.org/10.1016/j.plantsci.2014.09.011

Jurburg, S. D., Nunes, I., Brejnrod, A., Jacquiod, S., Priemé, A., Sørensen, S. J., Van Elsas, J. D., & Salles, J. F. (2017). Legacy Effects on the Recovery of Soil Bacterial Communities from Extreme Temperature Perturbation. *Frontiers in Microbiology*, *8*, 1832. https://doi.org/10.3389/fmicb.2017.01832

Kirchhoff, C., & Cypionka, H. (2017). Propidium ion enters viable cells with high membrane potential during live-dead staining. *Journal of Microbiological Methods*, *142*, 79–82. https://doi.org/10.1016/j.mimet.2017.09.011

Kralik, P., Nocker, A., & Pavlik, I. (2010). Mycobacterium avium subsp. Paratuberculosis viability determination using F57 quantitative PCR in combination with propidium monoazide treatment. *International Journal of Food Microbiology*, *141*, S80–S86. https://doi.org/10.1016/j.ijfoodmicro.2010.03.018

Lang, J. M., Coil, D. A., Neches, R. Y., Brown, W. E., Cavalier, D., Severance, M., Hampton-Marcell, J. T., Gilbert, J. A., & Eisen, J. A. (2017). A microbial survey of the International Space Station (ISS). *PeerJ*, *5*, e4029. https://doi.org/10.7717/peerj.4029

Lax, S., Smith, D. P., Hampton-Marcell, J., Owens, S. M., Handley, K. M., Scott, N. M., Gibbons, S. M., Larsen, P., Shogan, B. D., Weiss, S., Metcalf, J. L., Ursell, L. K., Vázquez-Baeza, Y., Van Treuren, W., Hasan, N. A., Gibson, M. K., Colwell, R., Dantas, G., Knight, R., & Gilbert, J. A. (2014). Longitudinal analysis of microbial interaction between humans and the indoor environment. *Science*, *345*(6200), 1048–1052. https://doi.org/10.1126/science.1254529

Leadbetter, E. R., & Foster, J. W. (1960). Bacterial oxidation of gaseous alkanes. *Archiv Für Mikrobiologie*, *35*(1), 92–104. https://doi.org/10.1007/BF00425597

Lewis, D. F. V., & Wiseman, A. (2005). A selective review of bacterial forms of cytochrome P450 enzymes. *Enzyme and Microbial Technology*, *36*(4), 377–384. https://doi.org/10.1016/j.enzmictec.2004.07.018

Li, S., Liu, H., Liu, G., & Hu, D. (2022). Simulated artificial interventions maintain species diversity of spatially structured microbial communities in closed ecosystem. *Acta Astronautica*, *201*, 39–47. https://doi.org/10.1016/j.actaastro.2022.08.023

Li, S., Liu, X., Li, Z., Liu, H., & Hu, D. (2023). Combination of direct boiling and glass beads increases the purity and accuracy of bacterial DNA extraction. *Biotechnology Journal*, *18*(11), 2300135.

Loozen, G., Boon, N., Pauwels, M., Quirynen, M., & Teughels, W. (2011). Live/dead real-time polymerase chain reaction to assess new therapies against dental plaque-related pathologies: Vitality real-time PCR. *Molecular Oral Microbiology*, *26*(4), 253–261. https://doi.org/10.1111/j.2041-1014.2011.00615.x

Macoy, D. M., Kim, W.-Y., Lee, S. Y., & Kim, M. G. (2015). Biotic stress related functions of hydroxycinnamic acid amide in plants. *Journal of Plant Biology*, *58*(3), 156–163.

https://doi.org/10.1007/s12374-015-0104-y

Martin, B., Raurich, S., Garriga, M., & Aymerich, T. (2013). Effect of Amplicon Length in
Propidium Monoazide Quantitative PCR for the Enumeration of Viable Cells of Salmonella
in Cooked Ham. *Food Analytical Methods*, *6*(2), 683–690. https://doi.org/10.1007/s12161-
012-9460-0

Martínez-Ramos, M., Ortiz-Rodríguez, I. A., Piñero, D., Dirzo, R., & Sarukhán, J. (2016).
Anthropogenic disturbances jeopardize biodiversity conservation within tropical rainforest
reserves. *Proceedings of the National Academy of Sciences*, *113*(19), 5323–5328.
https://doi.org/10.1073/pnas.1602893113

Mohammed, A. E., Abdul-Hameed, Z. H., Alotaibi, M. O., Bawakid, N. O., Sobahi, T. R., Abdel-
Lateff, A., & Alarif, W. M. (2021). Chemical Diversity and Bioactivities of Monoterpene
Indole Alkaloids (MIAs) from Six Apocynaceae Genera. *Molecules*, *26*(2), 488.
https://doi.org/10.3390/molecules26020488

Newman, E. A. (2019). Disturbance Ecology in the Anthropocene. *Frontiers in Ecology and
Evolution*, *7*, 147. https://doi.org/10.3389/fevo.2019.00147

Nocker, A., Sossa, K. E., & Camper, A. K. (2007). Molecular monitoring of disinfection efficacy
using propidium monoazide in combination with quantitative PCR. *Journal of
Microbiological Methods*, *70*(2), 252–260. https://doi.org/10.1016/j.mimet.2007.04.014

Nogami, M., Fujii, Y., & Sugo, T. (1996). Radiation resistance of pyridine type anion exchange
resins for spent fuel treatment. *Journal of Radioanalytical and Nuclear Chemistry Articles*,
203(1), 109–117. https://doi.org/10.1007/BF02060385

Novikova, N., De Boever, P., Poddubko, S., Deshevaya, E., Polikarpov, N., Rakova, N., Coninx, I.,
& Mergeay, M. (2006). Survey of environmental biocontamination on board the International
Space Station. *Research in Microbiology*, *157*(1), 5–12.
https://doi.org/10.1016/j.resmic.2005.07.010

Patil, S. M., Suryavanshi, M. V., Chandanshive, V. V., Kurade, M. B., Govindwar, S. P., & Jeon, B.-
H. (2020). Regeneration of textile wastewater deteriorated microbial diversity of soil
microcosm through bioaugmentation. *Chemical Engineering Journal*, *380*, 122533.
https://doi.org/10.1016/j.cej.2019.122533

Pickett, S., & White, P. (1985). *Ecology of natural disturbance and patch dynamics*.

Pillay, K. K. S. (1986). A review of the radiation stability of ion exchange materials. *Journal of
Radioanalytical and Nuclear Chemistry Articles*, *102*(1), 247–268.
https://doi.org/10.1007/BF02037966

Powell, J. R., Karunaratne, S., Campbell, C. D., Yao, H., Robinson, L., & Singh, B. K. (2015).
Deterministic processes vary during community assembly for ecologically dissimilar taxa.
Nature Communications, *6*(1), 8444. https://doi.org/10.1038/ncomms9444

Ramm, L., Siani, H., Wesgate, R., & Maillard, J.-Y. (2015). Pathogen transfer and high variability
in pathogen removal by detergent wipes. *American Journal of Infection Control*, *43*(7), 724–

728. https://doi.org/10.1016/j.ajic.2015.03.024

Rosado, H., Doyle, M., Hinds, J., & Taylor, P. W. (2010). Low-shear modelled microgravity alters
expression of virulence determinants of Staphylococcus aureus. *Acta Astronautica*, *66*(3),
408–413. https://doi.org/10.1016/j.actaastro.2009.06.007

Sattar, S. A., Bradley, C., Kibbee, R., Wesgate, R., Wilkinson, M. A. C., Sharpe, T., & Maillard, J.-
Y. (2015). Disinfectant wipes are appropriate to control microbial bioburden from surfaces:
Use of a new ASTM standard test protocol to demonstrate efficacy. *Journal of Hospital*
Infection, *91*(4), 319–325. https://doi.org/10.1016/j.jhin.2015.08.026

Schnetzinger, F., Pan, Y., & Nocker, A. (2013). Use of propidium monoazide and increased amplicon
length reduce false-positive signals in quantitative PCR for bioburden analysis. *Applied*
Microbiology and Biotechnology, *97*(5), 2153–2162. https://doi.org/10.1007/s00253-013-
4711-6

Sellner, B., Prakapaitė, R., van Berkum, M., Heinemann, M., Harms, A., & Jenal, U. (2021). A New
Sugar for an Old Phage: A c-di-GMP-Dependent Polysaccharide Pathway Sensitizes
Escherichia coli for Bacteriophage Infection. *mBio*, *12*(6), e03246-21.
https://doi.org/10.1128/mbio.03246-21

Shade, A., Peter, H., Allison, S. D., Baho, D. L., Berga, M., Bürgmann, H., Huber, D. H.,
Langenheder, S., Lennon, J. T., Martiny, J. B., & others. (2012). Fundamentals of microbial
community resistance and resilience. *Frontiers in Microbiology*, *3*, 417.

Shea, K., Roxburgh, S. H., & Rauschert, E. S. J. (2004). Moving from pattern to process:
Coexistence mechanisms under intermediate disturbance regimes. *Ecology Letters*, *7*(6),
491–508. https://doi.org/10.1111/j.1461-0248.2004.00600.x

Siani, H., Cooper, C., & Maillard, J.-Y. (2011). Efficacy of "sporicidal" wipes against Clostridium
difficile. *American Journal of Infection Control*, *39*(3), 212–218.
https://doi.org/10.1016/j.ajic.2011.01.006

Singh, N. K., Wood, J. M., Karouia, F., & Venkateswaran, K. (2018). Succession and persistence of
microbial communities and antimicrobial resistance genes associated with International
Space Station environmental surfaces. *Microbiome*, *6*(1), 204.
https://doi.org/10.1186/s40168-018-0585-2

Slopek, R., Condon, B., Sawhney, P., Reynolds, M., & Allen, C. (2011). Adsorption of alkyl-
dimethyl-benzyl-ammonium chloride on differently pretreated non-woven cotton substrates.
Textile Research Journal, *81*(15), 1617–1624. https://doi.org/10.1177/0040517511407373

Smitha, M., Singh, S., & Singh, R. (2017). Microbial biotransformation: A process for chemical
alterations. *J Bacteriol Mycol Open Access*, *4*(2), 85.

Song, X., Vossebein, L., & Zille, A. (2019). Efficacy of disinfectant-impregnated wipes used for
surface disinfection in hospitals: A review. *Antimicrobial Resistance & Infection Control*,
8(1), 139. https://doi.org/10.1186/s13756-019-0595-2

Song, Z., Han, H., & Zhu, W. (2015). Morphological variation and recovery mechanism of residual

crude oil by biosurfactant from indigenous bacteria: Macro-and pore-scale experimental investigations. *Journal of Microbiology and Biotechnology*, *25*(6), 918–929.

Srivastava, D. S., & Vellend, M. (2005). Biodiversity-Ecosystem Function Research: Is It Relevant to Conservation? *Annual Review of Ecology, Evolution, and Systematics*, *36*(1), 267–294. https://doi.org/10.1146/annurev.ecolsys.36.102003.152636

Stegen, J. C., Lin, X., Fredrickson, J. K., Chen, X., Kennedy, D. W., Murray, C. J., Rockhold, M. L., & Konopka, A. (2013). Quantifying community assembly processes and identifying features that impose them. *The ISME Journal*, *7*(11), 2069–2079. https://doi.org/10.1038/ismej.2013.93

Stewart, F. M., & Levin, B. R. (1973). Partitioning of Resources and the Outcome of Interspecific Competition: A Model and Some General Considerations. *The American Naturalist*, *107*(954), 171–198. https://doi.org/10.1086/282825

Sun, Y., Hu, D., Li, L., Jing, Z., Wei, C., Zhang, L., Fu, Y., & Liu, H. (2016). Influence mechanism of low-dose ionizing radiation on Escherichia coli DH5α population based on plasma theory and system dynamics simulation. *Journal of Environmental Radioactivity*, *151*, 185–192. https://doi.org/10.1016/j.jenvrad.2015.10.007

Tolkkinen, M., Mykrä, H., Markkola, A., Aisala, H., Vuori, K., Lumme, J., Pirttilä, A. M., & Muotka, T. (2013). Decomposer communities in human-impacted streams: Species dominance rather than richness affects leaf decomposition. *Journal of Applied Ecology*, *50*(5), 1142–1151. https://doi.org/10.1111/1365-2664.12138

Tut, G., Magan, N., Brain, P., & Xu, X. (2021). Molecular Assay Development to Monitor the Kinetics of Viable Populations of Two Biocontrol Agents, *Bacillus subtilis* QST 713 and *Gliocladium catenulatum* J1446, in the Phyllosphere of Lettuce Leaves. *Biology*, *10*(3), 224. https://doi.org/10.3390/biology10030224

Urbaniak, C., Sielaff, A. C., Frey, K. G., Allen, J. E., Singh, N., Jaing, C., Wheeler, K., & Venkateswaran, K. (2018). Detection of antimicrobial resistance genes associated with the International Space Station environmental surfaces. *Scientific Reports*, *8*(1), 814. https://doi.org/10.1038/s41598-017-18506-4

Wang, Q., Liang, J., Zhang, S., Yoza, B. A., Li, Q. X., Zhan, Y., Ye, H., Zhao, P., & Chen, C. (2020). Characteristics of bacterial populations in an industrial scale petrochemical wastewater treatment plant: Composition, function and their association with environmental factors. *Environmental Research*, *189*, 109939. https://doi.org/10.1016/j.envres.2020.109939

Williams, G. J., Denyer, S. P., Hosein, I. K., Hill, D. W., & Maillard, J.-Y. (2007). The development of a new three-step protocol to determine the efficacy of disinfectant wipes on surfaces contaminated with *Staphylococcus aureus*. *Journal of Hospital Infection*, *67*(4), 329–335. https://doi.org/10.1016/j.jhin.2007.08.012

Xie, X., Wang, S., Jiang, S. C., Bahnemann, J., & Hoffmann, M. R. (2016). Sunlight-Activated Propidium Monoazide Pretreatment for Differentiation of Viable and Dead Bacteria by

Quantitative Real-Time Polymerase Chain Reaction. *Environmental Science & Technology*

Letters, *3*(2), 57–61. https://doi.org/10.1021/acs.estlett.5b00348

Yang, X., Li, S., Song, G., Xu, X., & Hu, D. (2021). Microbial diversity formed and maintained

through substrate feedback regulation and delayed responses induced by Low-Dose Ionizing

Radiation. *Acta Astronautica*, *188*, 239–251. https://doi.org/10.1016/j.actaastro.2021.07.027

Yang, X., Xu, X., & Hu, D. (2020). Succession mechanism of microbial community with high

species diversity in nutrient-deficient environments with low-dose ionizing radiation.

Ecological Modelling, *435*, 109270. https://doi.org/10.1016/j.ecolmodel.2020.109270

Zea, L., Prasad, N., Levy, S. E., Stodieck, L., Jones, A., Shrestha, S., & Klaus, D. (2016). A

Molecular Genetic Basis Explaining Altered Bacterial Behavior in Space. *PLOS ONE*, *11*(11),

e0164359. https://doi.org/10.1371/journal.pone.0164359

Zeng, X.-Y., Li, S.-W., Leng, Y., & Kang, X.-H. (2020). Structural and functional responses of

bacterial and fungal communities to multiple heavy metal exposure in arid loess. *Science of*

The Total Environment, *723*, 138081. https://doi.org/10.1016/j.scitotenv.2020.138081

Zhang, C., Cui, F., Zeng, G., Jiang, M., Yang, Z., Yu, Z., Zhu, M., & Shen, L. (2015). Quaternary

ammonium compounds (QACs): A review on occurrence, fate and toxicity in the

environment. *Science of The Total Environment*, *518–519*, 352–362.

https://doi.org/10.1016/j.scitotenv.2015.03.007

Zhou, J., & Ning, D. (2017). Stochastic community assembly: Does it matter in microbial ecology?

Microbiology and Molecular Biology Reviews, *81*(4), e00002-17.

Figure 1: The LDIR environmental simulator (A) and its 3D geometric

configuration (B).

Figure 2: The overview of the community succession under all treatments. The x-

axis represents the QAC concentration (e.g. 1 stands for 100%, 10^{-1} stands for 10%),

where the left side of axis includes groups without LDIR and the right side includes

groups with LDIR; y-axis represents the dilution ratio (e.g. 10^{-1} stands for 10%, 10^{-2}

stands for 1%). each position on the x-y plane corresponds to a different treatment. z-

axis denotes the number of days, and the seven spheres in each cluster mean seven days

of community succession for a treatment group. The radius of the spheres indicates the

total biomass in the community (compared to the maximum value and taken as log ten).

RGB color shows the percentage of different species (Blue, red and green stand for

Bacillus subtilis, *Escherichia coli* and *Pseudomonas aeruginosa*, respectively). WO

means "without". The 3D version is uploaded as a .fig file in the supplementary

materials.

Figure 3: The absolute abundance of three species (A-C) and the total biomass (D)

in the control group.

Figure 4: The absolute abundance of three species (A-C) and the total biomass (D)

under LDIR (solid line) and in the control group (dotted line).

Figure 5: The absolute abundance of three species (A-C) and the total biomass

(D) under 1‰ dilution (solid line) and in the control group (dotted line).

Figure 6: The absolute abundance of *E. coli*, *P. aeruginosa* as well as the total

biomass under 1% QAC (A, solid line) and 1‰ QAC (B, solid line) and in the

control group (dotted line). *B. subtilis* has a very low abundance, so it is not shown

in the figure; the same as below.

Figure 7: The simulation results from the model built in MatLab/Simulink. A-H

represent the control group, LDIR, low dilution rate, high dilution rate, low QAC

concentration, high QAC concentration, LDIR with low QAC concentration, all three

factors together, respectively.

Figure 8: The overall mechanisms of LDIR, dilution and QAC on microbial

community succession. LDIR activates all species' growth and reproduction. Dilution

allows specific species to be selectively released from suppression. Low-concentration

QAC undergoes radiolysis under LDIR and co-metabolism by bacteria, thus changing

from bactericide with harmful effects to nutrients. This figure is created with

BioRender.com.

Tables

862   **Table 1 Conditions of Distinct Groups**

Group	LDIR	Dilution	QAC
1	N	N	N
2	N	10%	N
3	N	1%	N
4	N	1‰	N
5	N	0.1‰	N
6	N	N	100%
7	N	N	10%
8	N	N	1%
9	N	N	1‰
10	Y	N	N
11	Y	10%	N
12	Y	1%	N
13	Y	1‰	N
14	Y	0.1‰	N
15	Y	N	100%
16	Y	10%	100%
17	Y	1%	100%
18	Y	1‰	100%
19	Y	0.1‰	100%
20	Y	N	10%
21	Y	10%	10%
22	Y	1%	10%

23	Y	1‰	10%
24	Y	0.1‰	10%
25	Y	N	1%
26	Y	10%	1%
27	Y	1%	1%
28	Y	1‰	1%
29	Y	0.1‰	1%
30	Y	N	1‰
31	Y	10%	1‰
32	Y	1%	1‰
33	Y	1‰	1‰
34	Y	0.1‰	1‰

"N" means this condition was not applicable in this group. The ratio of dilution meant

that part of the culture from t-th day was taken into the R2A solution for gaining the

10ml new culture of $t + 1$-th day. No dilution meant that R2A solution was added into

the old culture to gain the 10ml new culture. The ratio of QAC meant the stock solution

of QAC was added into the culture to make that concentration of the stock solution of

QAC.

Table 2 The composition of QAC

Component	per 100ml	Final Concentration
Octyl Decyl Dimethyl Ammonium Chloride (Purity: 80%)	0.5g	0.04%
Dioctyl Dimethyl Ammonium Chloride (Purity: 80%)	0.25g	0.02%
Didecyl Dimethyl Ammonium Chloride (Purity: 95%)	0.21g	0.02%
Alkyl Dimethyl Benzyl Ammonium Chloride, (50% C14, 40% C12, 10% C16) (Purity: 80%)	0.66g	0.0532%

	ddH$_2$O		98ml	-

 Table 3 The experimental conditions of distinct metabolomics test series

Series	LDIR	Dilution	QAC	Day
A	N	N	N	1
B	N	N	N	4
C	N	N	N	5
D	N	N	N	6
E	N	N	N	7
F	N	1‰	N	4
G	N	N	1%	4
H	N	N	1%	5
I	N	N	1%	6
J	N	N	1‰	4
K	N	N	1‰	7
L	Y	N	N	1
M	Y	N	N	4
N	Y	N	N	5
O	Y	N	N	6
P	Y	N	N	7
Q	Y	1‰	N	4
R	Y	N	1%	4
S	Y	N	1%	5
T	Y	N	1%	6
U	Y	1‰	1%	4
V	Y	N	1‰	4

W	Y	N	1‰	7
X	Y	1‰	1‰	4

Figure S1: The absolute abundance of three species (A-C) and the total biomass

(D) under different dilution ratios. The thinner the line, the more diluted the

community in the group is.

Figure S2: The relative abundance of three species under different dilution ratios.

A-D represent 10%, 1%, 1‰ and 0.1‰ dilution ratios, respectively.

Figure S3: The absolute abundance of three species (A-C) and the total biomass

(D) under LDIR and 1‰ dilution (solid line) as well as only 1‰ dilution (dotted

line).

Figure S4: The absolute abundance of *E. coli*, *P. aeruginosa* as well as the total

biomass under LDIR and 1% QAC (A, solid line) or LDIR and 1‰ QAC (B,

solid line) as well as only LDIR (dotted line).

Figure S5: The absolute abundance of *E. coli*, *P. aeruginosa* as well as the total biomass under LDIR and 1% QAC (A, solid line) or LDIR and 1‰ QAC (B, solid line) as well as only 1% QAC (A, dotted line) or 1‰ QAC (B, dotted line).

Figure S6: The absolute abundance of *E. coli*, *P. aeruginosa* as well as the total biomass under LDIR, 1% dilution and 1% QAC (A, solid line) or LDIR, 1% dilution and 1‰ QAC (B, solid line) as well as in the control group (dotted line).

Figure S7: the concentration of p-coumaroyl agmatine in the group exposed under LDIR between the 4-6ᵗʰ days dropped significantly compared with the control

Figure S8: the differential abundance scores of differential metabolisms between

the group under LDIR and the control group. A-D represent the 1, 4, 5, 6th day,

respectively.

Figure S9: the enrichment of N-acetylmannosamine was markedly correlated with

the rise of the abundance of *P. aeruginosa* ($p < 0.01$) on the 4th day of the group

diluted with 1‰ ratio per day, compared with the control group. Metabolites with

the top 10 highest concentrations in either group, or whose $|\log_2 \text{FC}| \geq 20$ were

 considered in the intergroup correlation analysis. "**" meant ($p < 0.01$), and "*" meant

($p < 0.05$). The same is as below.

Figure S10: the differential abundance scores of differential metabolisms on 4th

day between the group diluted with 1‰ ratio per day and the control group.

Figure S11: the enrichment of N-acetylmannosamine was markedly correlated

with the rise of the abundance of *E. coli* ($p < 0.05$) on the 4th day of the group with

1% stock solution of QAC, compared with the control group.

Figure S12: the drop of echitovenine was markedly correlated with the increase of

the abundance of *P. aeruginosa* ($p < 0.01$) on the 7th day of the group with 1‰

stock solution of QAC, compared with the control group.

Figure S13: the differential abundance scores of differential metabolisms on 7th

day between the group with 1‰ stock solution of QAC and the control group.

Figure S14: the differential abundance scores of differential metabolisms between the group with both LDIR and 1% stock solution of QAC and the group under 1% stock solution of QAC. A and B represent the 5th and 6th day, separately.

Figure S15: the concentration of albaflavenone sharply diminished on 4th day in the group with both LDIR and 1‰ stock solution of QAC, compared with the control group.

Figure S16: the differential abundance scores of differential metabolisms between the group with both LDIR and 1‰ stock solution of QAC and the group under 1‰ stock solution of QAC. A and B represent the 4th and 7th day, separately.

Figure S17: the differential abundance scores of differential metabolisms on 4th day between the groups with 1% stock solution of QAC and 1‰ stock solution of QAC.

Figure S18: The overview of the community succession under all treatments and

its slides. (A) The two planes in the yellow boxes showed that *E.coli* took the majority

when both dilution and LDIR were applied compared to only dilution. (B) With higher

gradients of QAC under 1%, *P. aeruginosa* gradually became the dominant species with

LDIR. (C) With higher gradients of QAC under 1%, *P. aeruginosa* gradually became

the dominant species without LDIR.

Figure S19: the actual and theoretical counts for 10× gradient dilution of *B.*

***subtilis* cells.**

935   **Table S1: The composition of modified minimal R2A**

Component	per 1000ml	Final concentration
NaCl	18.0g	1.8%
Peptone	0.5g	0.05%
Yeast Extract Powder	0.5g	0.05%
Casein Peptone	0.5g	0.05%
Sodium Pyruvate	0.3g	0.03%

936 **Table S2 Orthogonal Experiments for PMA-qPCR Conditions**

No	PMA Concentration (mmol/L)	PMA Incubation Time (min)	PMA Incubation Temperature (°C)	Light Exposure Time (min)	Light Exposure Temperature (°C)	Light Intensity (W)	Changing Tubes or not	ΔCt(L)	ΔCt(D)
1	50(B)	10	20(A)	30(B)	4(A)	120(B)	N(B)	-0.642	-12.604
2	25(A)	10(A)	37(B)	15(A)	20(B)	120	N	1.845	-8.208
3	25	10	20	15	4	60(A)	Y(A)	-0.712	-14.497
4	50	10	37	30	20	60	Y	0.468	-11.381
5	50	30(B)	37	15	4	60	N	1.614	-15.257
6	25	30	20	30	20	60	N	0.772	-9.731
7	50	30	20	15	20	120	Y	-0.592	-17.972
8	25	30	37	30	4	120	Y	0.17	-10.692
TL$_A$	2.075	0.959	-1.174*	2.155	0.43*	2.142	-0.666*		
TD$_A$	-43.128	-46.69	-54.804*	-38.779	-53.05*	-50.866	-54.542*		
TL$_B$	0.848 (p=0.092)	1.964	4.097	0.768 (p=0.059)	2.493	0.781 (p=0.063)	3.589		
TD$_B$	-57.214*	-53.652*	-45.538	-44.408*	-47.292	-49.476	-45.8		

ΔCt(L) and ΔCt(D) represent the measured live/dead microbes abundance difference

between the control group and experimental group (with PMA), respectively. TL$_A$ and

TD$_A$ stand for the sum of the test results ΔCt(L)/ ΔCt(D) for the column with level A,

respectively. TL$_B$ and TD$_B$ stand for the sum of the test results ΔCt(L)/ ΔCt(D) for the

column with level B, respectively. "*" means $p > 0.05$.

Table S3 Gradient Experiments for PMA-qPCR Conditions

No	PMA Concentration (mmol/L)	PMA Incubation Time	PMA Incubation Temperature	Light Exposure Time	Light Exposure Temperature	Light Intensity (W)	Changing Tubes or not	ΔCt(L)	ΔCt(D)

	(min)	(°C)	(min)	(°C)				
1	50	40	4	30	4	120	Y	0.89	-10.421
2	50	30	20	30	4	120	Y	1.242	-16.583
3	50	10	20	30	4	120	Y	0.244	-13.626
4	50	5	4	30	4	120	Y	0.891	-11.641
5	50	10	4	30	4	120	Y	0.497	-13.451
6	50	30	4	30	4	120	Y	1.045	-18.622
7	50	5	20	30	4	120	Y	1.691	-14.067
8	50	40	20	30	4	120	Y	1.319	-19.609*

Supplementary Material A

Several studies from different perspectives have shown a special phenomenon in *Bacillus subtilis* (Kirchhoff & Cypionka, 2017; Tut et al., 2021; Xie et al., 2016), which is, that some nucleic acid dyes like PMA could penetrate a large number of living cells due to the excessively strong cell membrane potential. Therefore, at the macro level, the rate of false-negative qPCR results would increase, and the threshold for detection of living *Bacillus subtilis* cells through the PMA-qPCR method would spike. In the pre-experiments of this study, *Bacillus subtilis* did not lead in the community. Hence, daily dilution to 10% of the culture yesterday experiment was performed, and the magnitude of counts of *Bacillus subtilis* was much higher than the circumstance "No *Bacillus subtilis* survived, its DNA was diluted simultaneously", as shown in Figure S19. Therefore, direct qPCR was applicable for approximately counting *Bacillus subtilis*.

Supplementary Material B

Eliminating the DNA of dead bacteria cells is the key aim of PMA in PMA-qPCR.

The standard PMA protocol might be suitable for natural environmental samples, but

not enough for high-abundance samples, like artificial microbial communities used in

this study. Therefore, it is important to optimize its conditions. Related factors included

PMA concentration, PMA incubation (with oscillation at 100rpm) time, PMA

incubation temperature, light exposure time, light exposure temperature, light intensity

(expressed as the blue LED bulb power), whether changing tubes between incubation

and light exposure (Banihashemi et al., 2012; Baymiev et al., 2020; Codony et al., 2020;

Emerson et al., 2017; Fittipaldi et al., 2012; Fleischmann et al., 2021; Loozen et al.,

2011; Martin et al., 2013). To find the highest DNA of dead bacteria cells elimination

rate, orthogonal experiments (Table S2) were implemented first on *P. aeruginosa* to

screen the better condition of the two-level factor (PMA was not applicable for *B.*

subtilis as seen in Sup. A and the primer of *E. coli* used for this study had a relatively

high detection threshold). Conditions were selected in previous studies (Agustí et al.,

2018; Codony et al., 2020; Emerson et al., 2017; Fittipaldi et al., 2012; Fleischmann et

al., 2021; Kralik et al., 2010).

The larger the absolute value of $\Delta Ct(D)$, the higher the dead bacteria cells'

elimination rate. Based on no significant difference of $\Delta Ct(D)$, the closer $\Delta Ct(L)$ to

zero, the smaller the influence of PMA on live cells. From the orthogonal experiments,

these conditions were significantly better: the PMA concentration was 50 mmol/L, the

light exposure time was 30min, the light exposure temperature was 4°C, and changing

tubes were applied. 120W light intensity has a p-value of 0.063 so it was also selected.

Incubation time and temperature were tested another time in gradient experiments

on *P. aeruginosa* as Table S3. As seen in Table, when the PMA incubation time was

40min and the PMA incubation temperature was 20°C, the absolute value of $\Delta Ct(D)$

was the largest with a significant difference. Therefore, the optimal condition was: to

incubate with 50mmol/L PMA 100rpm for 40min at 20°C, change a new tube, and

expose it under a blue LED bulb still for 30min at 4°C.

When applying this optimal condition to *E. coli*, the ΔCt(D)=-18.268. Compared

to the standard condition of PMA protocol, the detective elimination rate of dead *E. coli*

surged from 25.976% to 99.999%, and that of *P. aeruginosa* dropped from 25.097% to

99.999%. The optimal condition can fulfill the need to measure high species abundance

in this study.

Supplementary Material C

In this study, based on LC-QTOF, the qualitative and quantitative analysis of

untargeted metabolomics of 72 samples was carried out. A total of 12,079 peaks were

detected, of which 1,529 metabolites were annotated. The RSD ≤ 30% ones accounted

for more than 60%, so it could be assumed that QC samples have good reproducibility,

the instrument was stable, and the requirements of quality control were met.

In OPLS-DA (Orthogonal Projections to Latent Structures-Discriminant Analysis),

all results of the experimental and their corresponding control groups could be

completely separated, indicating significant overall changes in metabolites for each

condition. In the comparison of groups J and V, the Q2Y < 0.5, thus VIP could not be

considered as a screening condition for differential metabolites. Besides the above, Q2Y

of all other comparisons > 0.5, indicating that the OPLS-DA model was valid.

Permutation tests were performed to check the reliability of the OPLS-DA model. The

slopes of Q2Y were positive for all groups except for the comparison of J and V, further

supporting that the OPLS-DA model was significant. Most of the blue dots in the

scatterplots were located above the red dots, indicating good independence of the

OPLS-DA model.

1007  In the following analysis, metabolites with the same trend on consecutive days,

1008 $|log_2FC| \geq 20$ (except for $SD = 0$ to remove the systematic error of the instrument)

1009 or ranking top 10 were screened.

1010

1011 **Supplementary Material D**

1012 D1 S-Function

1013 function [sys,x0,str,ts] =lss_with2f(t,x,u,flag, X0,mu, Ks, Kf, beta, alpha, delta, phi,

1014 C_I, C_E,D_I,D_E,ep, h, gamma,I,c)

1015 % ===========================

1016 switch flag

1017  case 0

1018   [sys,x0,str,ts]=mdlInitializeSizes(X0);

1019  case 1

1020   sys=mdlDerivatives(t,x,u,mu, Ks, Kf, beta, alpha, delta, phi, C_I, C_E,

1021 D_I,D_E,ep, h, gamma,I,c);

1022  case 2

1023   sys=mdlUpdate(t,x,u);

1024  case 3

1025   sys=mdlOutputs(t,x,u);

1026  case 4

1027   sys=mdlGetTimeOfNextVarHit(t,x,u);

1028  case 9

1029   sys=mdlTerminate(t,x,u);

1030  otherwise

```matlab
1031          DAStudio.error('Simulink:blocks:unhandledFlag', num2str(flag));

1032    end

1033    % ========================================

1034    function [sys,x0,str,ts]=mdlInitializeSizes(X0)

1035    sizes = simsizes;

1036    sizes.NumContStates   = 6;

1037    sizes.NumDiscStates   = 0;

1038    sizes.NumOutputs      = 6;

1039    sizes.NumInputs       = 0;

1040    sizes.DirFeedthrough = 1;

1041    sizes.NumSampleTimes = 1;

1042    sys = simsizes(sizes);

1043    x0   = X0;

1044    str = [];

1045    ts   = [0 0];

1046    %=====================

1047    function sys=mdlDerivatives(t,x,u,mu, Ks, Kf, beta, alpha, delta, phi, C_I, C_E, D_I,

1048    D_E,ep, h, gamma,I,c)

1049    S=x(4);

1050    E=x(5);

1051    F=x(6);

1052    v1b=mu(1)*x(1)*(S/(Ks(1)+S));

1053    v2b=mu(2)*x(2)*(S/(Ks(2)+S));

1054    v3b=mu(3)*x(3)*(S/(Ks(3)+S));

1055    %================
```

```matlab
va1=beta(1)*exp(-C_I(1)*I-C_E(1)*E)*x(1)^2;

vr1=alpha(1,2)*exp(-D_I(1,2)*I-D_E(1,2)*E)*x(1)*x(2)+alpha(1,3)*exp(-
D_I(1,3)*I-D_E(1,3)*E)*x(1)*x(3)

vv1=delta(1)*x(1)+ep(1)*x(1)*E;

%================

va2=beta(2)*exp(-C_I(2)*I-C_E(2)*E)*x(2)^2;

vr2=alpha(2,1)*exp(-D_I(2,1)*I-D_E(2,1)*E)*x(2)*x(1)+alpha(2,3)*exp(-
D_I(2,3)*I-D_E(2,3)*E)*x(2)*x(3);

vv2=delta(2)*x(2)+ep(2)*x(2)*E;

%================

va3=beta(3)*exp(-C_I(3)*I-C_E(3)*E)*x(3)^2;

vr3=alpha(3,1)*exp(-D_I(3,1)*I-D_E(3,2)*E)*x(3)*x(1)+alpha(3,2)*exp(-
D_I(3,2)*I-D_E(3,2)*E)*x(3)*x(2);

vv3=delta(3)*x(3)+ep(3)*x(3)*E;

%================

ve=gamma*I*E;

%================

sys = [v1b-va1-vr1-vv1;...
    v2b-va2-vr2-vv2;...
    v3b-va3-vr3-vv3;...
   h*ve-phi(1)*v1b-phi(2)*v2b-phi(3)*v3b;...
    -ve;...
    h*ve];

%================

function sys=mdlUpdate(t,x,u)
```

```matlab
1081    sys = [];
1082    %===============
1083    function sys=mdlOutputs(t,x,u)
1084    sys = x(1:6);
1085    %===============
1086    function sys=mdlGetTimeOfNextVarHit(t,x,u)
1087    sys=[];
1088    %===============
1089    function sys=mdlTerminate(t,x,u)
1090    sys = [];
1091    %===============
1092
```

```
1093    D2 Main Function
1094    clear;clc;shg
1095    rng(5);
1096    set_param(gcs,'MaxStep','.08');
1097
1098    S0=50+2*rand;
1099    E0=0;
1100    F0=0;
1101    X0=[3.3333;3.3333;3.3333;S0;E0;F0];
1102    mu=[0.01;0.1;0.045];
1103    Ks=20+rand(3,1);
1104    Kf=0.05+0.005*rand(3,1);
1105    beta=[0.002;0.003;0.001];
1106    alpha=.003*rand(3);
1107    delta=[0.1;0.03;0.015];
1108    phi=rand(3,1);
1109    C_I=1*rand(3,1);
1110    C_E=1*rand(3,1);
1111    D_I=1*rand(3);
1112    D_E=1*rand(3);
1113    h=0.1;
1114    gamma=.1+.2*rand;
1115    I=0;
1116    c=rand(3,1);
1117    t=[];
1118    y=[];
1119    dilution=1;
1120    ep=[0.1;0.005;0.001];
1121    %      =========================
1122    for i=1:7
1123        lss=sim(gcs,[0,24]);
1124        t=[t;(lss.tout+24*(i-1)+0.08)];
1125        y=[y;lss.yout];
1126        X0(1:3)=dilution*y(end,(1:3));
```

```matlab
1127        if dilution<1
1128            X0(4)=S0;
1129        else
1130            X0(4)=y(end,4);
1131        end
1132        E0=1;
1133        X0(5)=dilution*y(end,5)+E0;
1134        X0(6)=dilution*y(end,6);
1135    end
1136    plot(t,y(:,1),'LineStyle','-','Color','#3366FF','linew',3);
1137    hold on
1138    plot(t,y(:,2),'LineStyle','-','Color','#c71585','linew',3);
1139    plot(t,y(:,3),'LineStyle','-','Color','#00a15c','linew',3);
1140    grid
1141    set(gca,'YLim',[0,inf],'XLim',[0,168],'XTickMode','manual','XTick',[0:24:168],'FontSi
1142    ze',30);
1143    legend('Biomass of the 1^s^t Species','Biomass of the 2^n^d Species','Biomass of the
1144    3^r^d Species','location','best','FontSize',30)
1145    xlabel('Time(h)','FontSize',25)
1146    ylabel('Biomass of One Species(refering to Abundance)','FontSize',25)
1147    hold off

1148
1149
```

1150   D3 MatLab/Simulink Block Diagrams

1151   Available on GitHub: https://github.com/angkuLC/codeforcureorcurse.

1152

1153