A Futile Cycle?

Tissue Homeostatic Trans-Membrane Water Co-Transport:

Kinetics, Thermodynamics, Metabolic Consequences.

Charles S. Springer, Jr.[a-e] Martin M. Pike,[a,c,e] Thomas M. Barbara[a]

[a]Advanced Imaging Research Center, [b] Department of Chemical Physiology and Biochemistry,
[c]Department of Biomedical Engineering, [d]Brenden-Colson Center for Pancreatic Care, [e]Knight Cancer Institute,
Oregon Health & Science University; Portland, Oregon

Corresponding author: C. S. Springer; springer@ohsu.edu
Advanced Imaging Research Center, L452
Oregon Health and Science University
3181 S. W. Sam Jackson Park Road
Portland, Oregon 97239

ABSTRACT:

The phenomenon of active trans-membrane water cycling (AWC) has emerged in little over a decade. Here, we consider H_2O transport across cell membranes from the origins of its study. Historically, trans-membrane water transport processes were classified into: A) compensating bidirectional fluxes ("*exchange*"), and B) unidirectional flux ("*net flow*") categories. Recent literature molecular structure determinations and molecular dynamic (MD) simulations indicate probably all the many different hydrophilic substrate membrane co-transporters have membrane-spanning hydrophilic pathways and co-transport water along with their substrates, and that they individually catalyze category A and/or B water flux processes, although usually not simultaneously. The AWC name signifies that, integrated over the all the cell's co-transporters, the rate of *homeostatic*, bidirectional trans-cytolemmal water exchange (category A) is synchronized with the metabolic rate of the crucial Na^+,K^+-ATPase (NKA) enzyme. A literature survey indicates the stoichiometric (category B) water/substrate ratios of individual co-transporters are often very large. The MD simulations also suggest how different co-transporter reactions can be *kinetically* coupled molecularly.

Is this (Na^+,K^+-ATPase rate-synchronized) cycling futile, or is it consequential? Conservatively representative literature metabolomic and proteinomic results enable comprehensive free energy analyses of the many transport reactions with known water stoichiometries. Free energy calculations, using literature intracellular pressure (P_i) values reveals there is an *outward* trans-membrane H_2O barochemical gradient of magnitude comparable to that of the well-known *inward* Na^+ electrochemical gradient. For most co-influxers, these gradients are finely balanced to maintain intracellular metabolite concentration values near their consuming enzyme Michaelis constants. The thermodynamic analyses include glucose, glutamate⁻, gamma-aminobutyric acid (GABA), and lactate⁻ transporters. 2%-4% P_i alterations can lead to disastrous concentration levels. For the neurotransmitters glutamate⁻ and GABA, very small astrocytic P_i changes can allow/disallow synaptic transmission. Unlike the Na^+ and K^+ electrochemical steady-states, the H_2O barochemical *steady-state* is in (or near) chemical *equilibrium*. The analyses show why the presence of aquaporins (AQPs) does not dissipate the trans-membrane pressure gradient. A feedback loop inherent in the opposing Na^+ electrochemical and H_2O barochemical gradients regulates AQP-catalyzed water flux as an integral AWC aspect. These results also require a re-consideration of the underlying nature of P_i. Active trans-membrane water cycling is not futile, but is inherent to the cell's "NKA system" - a new, fundamental aspect of biology.

SYNOPSIS:

Via intracellular pressure, membrane co-transported water influences thermodynamic control of cell metabolite maintenance.

INTRODUCTION:

One of the essential aspects of water in living tissue is its intra- extracellular compartmentalization. The kinetics

of cellular, homeostatic trans-membrane water transport provides the basis of a new non-invasive, high-resolution MRI

approach; metabolic activity diffusion imaging, MADI (1-3). It is providing novel views of cancer (4) and brain function

(5) metabolism. Here, we inquire into kinetic, thermodynamic, and metabolic consequences of water co-transport

phenomena in general.

Distinct Trans-membrane Water Transport Processes: Category B *unidirectional* "flux" processes driven

by trans-membrane osmotic gradients have been extensively investigated (6,7). These are characterized by cell volume

(V) changes [cell swelling/shrinking; edema; tissue hypertrophy]. Their kinetics are quantified with the "osmotic"

[flux, flow] water permeability coefficient, P_f.

However, the earliest studies of which we are aware also detected category A *bidirectional* trans-membrane

water "exchange" (8,9). In such processes, there is no V change – the systems are *homeostatic*. The kinetics are

quantified with the "diffusional" water permeability coefficient, P_d. Initial studies employed isotopic labeling ($^2HO^1H$),

requiring very rapid, careful solution mixing – to perturb the system from isotopic equilibrium. As tracer methods, such

approaches yield the $P_d \bullet a_e$ product: $a_e = \rho \langle A \rangle$, where ρ is the cell [number] density [cells/volume(tissue)] and $\langle A \rangle$ is

the tissue-averaged cell surface area (1). [In the older literature, a_e is given as S, the *total* cell surface area per

tissue ["ensemble"] volume.] P_d can be determined if a_e can be estimated. Early P_d values were compiled (10). We (11)

and others showed that, at steady-state, P_d confounds V and A with the homeostatic cellular water efflux rate constant,

k_{io}, $P_d = (V/A)k_{io}$. It is k_{io} that is a true measure of permeation probability (1). A nuclear magnetic resonance (NMR)

approach was introduced in 1972 (12), which we later termed the "shutter-speed" (SS-NMR) method (13). Major

SS-NMR advantages are it: **1.)** yields k_{io} un-confounded, and separated from $(1 + ((\rho \bullet V - 1)/f_w))$: f_w is the tissue water

volume fraction (1,14); **2.)** does not require rapid solution mixing; and **3.)** could, in principle, quantify the category B flux

and category A exchange processes simultaneously. An extracellular paramagnetic agent labels extracellular ^1H_2O

magnetization, not extracellular water molecules (15), and non-invasive radio frequency electromagnetic pulses perturb

the system magnetization from equilibrium. A compilation of early SS-NMR results was reported in (16). (Acronyms and

symbols are listed in the **Appendix A1**).

Molecular Aspects of Trans-membrane Water Transport: Over the years, many membrane-bound macromolecules

have been found to transport water between intra- and extracellular spaces. In 1988, Agre and co-workers reported

the selective water transport protein aquaporin (AQP) family (17). The impressive array of different AQP variants now

known has been reviewed (6). During the 1990's and 2000's, many other proteins were found to co-transport water

molecules, along with the metabolic substrates for which they are named (reviewed in (18,19)). **Figure 1** shows AQP and

Figure 1. An Inventory of water transporting membrane proteins. The water stoichiometric values are taken from (19), and should be thought of as means of shot-to-shot variations (see text).

the ten water co-transporters detailed by Zeuthen (19). Each of their reactions is reversible, but here they are classified

by their tendencies for influx or efflux under normal cellular conditions. These proteins all catalyze crucial processes,

and are profitably considered as enzymes.

There are two glucose influx families: those co-transporting Na^+ (SGLT), and those that do not (GLUT).

(Interestingly, though both families transport glucose, it has been recently noted (20) the SGLT family does not transport

2-deoxy-2-[18]F-fluoro-D-glucose (2-FDG), the [18]FDG-PET (positron emission tomography) glucose tracer. This can

present interpretative problems.) The excitatory amino acid transporter (EAAT1) and the GABA transporter (GAT1)

facilitate uptake of the principal excitatory and inhibitory neurotransmitters, glutamate[-] and GABA

[the gamma-aminobutyric acid zwitterion], respectively. (In the Fig. 1 EAAT1 influx reaction, note the H_3O^+ co-influx, and

the K^+ counter-*efflux* components (21).) The $Na^+,K^+,2Cl^-$ co-transporter (NKCC1) facilitates K^+ and Cl^- influx (22).

The K^+,Cl^- co-transporter (KCC4) provides an important pathway for K^+ and Cl^- efflux (22). The monocarboxylic

acid transporter (MCT1) extrudes the lactate[-] produced by cytoplasmic glycolytic-type metabolisms (HCO_3^- is

the analogous product of mitochondrial oxidative phosphorylation (23).

One is struck by the large water stoichiometries in most co-transporter cases. These values should

be considered averages. Surely, they fluctuate stochastically (shot-to-shot) with each enzyme cycle (see below).

Also, they likely vary with the cellular environment and biological condition. It is important to realize the Fig. 1

stoichiometries were determined in model systems *in vitro*, and by inducing net category B unidirectional fluxes (19).

Homeostatic exchange was not studied in this regard.

In 1989, Ye and Verkman presented an elaborate optical method to measure P_f and P_d for *simultaneous* water

efflux from and exchange in liposomes and in (adenosine tri-phosphate) ATP-free erythrocyte ghosts (24). Two very

important results from that study are the following. First, a given water transporter can be involved in both category B

flux and category A exchange processes, but not necessarily in the same proportions. Using $HgCl_2$ to inhibit AQP,

the authors found while AQP contributes 90% of ghost water efflux, it accounts for only 45% of ghost exchange flux.

Second, though P_f for ghost efflux is numerically four times larger than P_d for ghost exchange (in the same units),

the time-course for water efflux is 50 times longer. The exchange process is much, much faster. A confusing aspect is

that P_f and P_d are always reported with the same dimensions (length/time). However, the two permeability coefficients

are defined by rate laws of very different natures ((7); (10), pp. 44 ff), and are not at all the same. The P_f quantity

characterizes transporter efficiency only in the presence of an osmotic gradient, and with the system perturbed from

the steady-state. It gives no information on P_d.

Active Trans-Membrane Water Cycling (AWC): During most *in vitro* and *in vivo* studies, the tissue is in homeostasis:

there is no substantial cell swelling or shrinking. Only category A trans-membrane water bidirectional *exchange* obtains.

Evidence mounts for a homeostatic, metabolic AWC process in cells (reviewed (1,2,15,25)). This is a very fast water

molecule exchange, the kinetics of which are driven by, and synchronized to, those of the rate-limiting plasma

membrane Na$^+$,K$^+$-ATPase (NKA) "sodium pump." The AWC phenomenon was first detected using water proton (^{1}H$_2$O)

SS-NMR in 2011 (26). **Figure 2** is a cartoon illustrating the AWC process. For each NKA cycle, one intracellular ATP$_i$

molecule is hydrolyzed, three intracellular Na$_i^+$ ions expelled, and two extracellular K$_o^+$ ions imported. (First and second

approximations of Fig 2 have appeared (2,15).) However, generic transporters II and III, respectively, allow K$^+$ to re-exit

and Na$^+$ to re-enter exit the cell. Thus, NKA cycles extremely rapidly: depending on NKA expression, there can be

10^{10} (Na$^+$ + K$^+$) ions(cycled)/s/cell (15). These actions maintain the crucial trans-membrane ion concentration gradients,

[Na$_o^+$] > [Na$_i^+$], [K$_o^+$] < [K$_i^+$]; and the membrane electrical potential (in more negative than out, Figs. 1,2). Explicit

examples of II and III are KCC4 and SGLT, respectively. The NKA substrates intracellular ATP and Na$^+$ and extracellular K$^+$

(ATP$_i$, Na$_i^+$, K$_o^+$, respectively) are rendered in red in Fig. 2, as is the natural product ouabain, a specific extracellular NKA

inhibitor (ouabain$_o$).

Generic water co-transporters I and IV represent *secondary*, *active* obligate water symporters for water to exit

and enter cells, respectively. Most of the Fig. 1 enzymes fit into this classification: they share substrates (Na$_o^+$ and/or K$_i^+$)

with NKA, a *primary*, *active* transporter (hydrolyzes ATP directly). The first-order, unidirectional rate constants k$_{io}$ and k$_{oi}$

are those for *cellular*, *homeostatic* water efflux and influx, respectively. In the differential first-order chemical

trans-membrane water transport rate law, k$_{io}$ can be expressed as the sum of energetically active, k$_{io}$(a) and passive,

k$_{io}$(p), contributions that are further elaborated in **Equations (1)**, where: x is the overall, cellular AWC water

$$k_{io} = k_{io}(a) \, + \, k_{io}(p) = \, \left(\frac{x}{[H_2O_i]V}\right) \, {}^cMR_{NKA} + \left(\frac{A}{V}\right)P_W(p) \tag{1}$$

stoichiometric coefficient (H$_2$O molecules(cycled)/NKA(cycle)/s/cell), [H$_2$O$_i$] the intracellular water concentration,

^{c}MR$_{NKA}$ the cellular NKA metabolic rate (ATP(molecules hydrolyzed by NKA)/s/cell), and P$_W$(p) the non-metabolic, passive

diffusive water membrane permeability coefficient P$_d$(p) (1,2,15). The influx rate constant, k$_{oi}$, is ρ-dependent, and also

has active and passive components that need not be in the same proportions as for k$_{io}$. In healthy, living tissue, k$_{io}$(p)

and k$_{oi}$(p) seem negligible (2). There can be 10^{12} (H$_2$O molecules actively cycled)/s/cell (15). The value of x may be

of magnitude 10^6 (2).

The sodium pump is perhaps biology's most vital enzyme: it is found in all mammalian cells, and it's role

in the evolution of life is thought crucial (27). Since NKA homeostatic activity has never before been accessible *in vivo*,

medical MR imaging (MRI) applications of MADI are very promising (1,2,4,5). The water proton MR signal (^{1}H$_2$O) is by far

the largest from tissue (28). The AWC-based MADI approach has been deemed "a new paradigm" (3).

Figure 2. A cartoon of active trans-membrane water cycling (AWC). The rate constant for steady-state cellular water efflux is k_{io}; that for steady-state cellular water influx is k_{oi}. The Na$^+$,K$^+$-ATPase enzyme is indicated as NKA, while AQP, KCC4, and SGLT are defined in Figure 1. The actions of NKA substrates intracellular Na$^+$ and ATP (Na$_i^+$ and ATP$_i$) and extracellular K$^+$ (K$_o^+$) are indicated in red, as are the NKA inhibitor extracellular ouabain (ouabain$_o$) and an extracellular AQP inhibitor. Generic transporters I, II, III, and IV are exemplified in Figure 1: the actions of AQP, KCC4, and SGLT are shown here as specific examples. The quantity x is the AWC water stoichiometry integrated over the entire cell. Thus, for example, $^cMR_{H2O}(influx) = {^cMR_{H2O}}(efflux) = x^cMR_{NKA} = x(^cMR_{Na+}(influx))/3 = x(^cMR_{K+}(efflux))/2$, where ^cMR represents a cellular metabolic rate. First and second approximations of this cartoon have appeared in (15) and (2), respectively.

Water-selective AQP transporters can also be involved in AWC as long there is a thermodynamic driving force

for water flux (our calculations suggest their role in cellular homeostasis, see Discussion). AQP molecules themselves

comprise simple, *passive* channels (29,30).

The existence of AWC has been demonstrated by many different, deliberate manipulations of NKA kinetics with,

for example, ATP_i, K_o^+ and $oubain_o$ concentration alterations (Fig. 2, red), while monitoring k_{io}. More than

20 independent studies, of a range of tissues, have been itemized (2,25). AWC can be mapped *in vivo* with non-invasive

MADI, which estimates k_{io}, V, and ρ separately (1,2). In awake, resting human brain gray matter [GM], MADI mapping

of the cellular water *flux* $[H_2O_i]k_{io}V = {}^cMR_{AWC}$, an x^cMR_{NKA} estimate, correlates very well with the tissue metabolic rate

of glucose consumption, ${}^tMR_{glu}$ (glucose(consumed)/s/unit volume(tissue)) determined from quantitative ^{18}FDG-PET (2).

This is expected: overall ATP production (MR_{ATP}) and consumption remain balanced in the resting brain. The white

matter [WM] ${}^cMR_{AWC}/{}^tMR_{glu}$ ratio is larger than the GM ${}^cMR_{AWC}/{}^tMR_{glu}$ slope (2), likely indicating a more oxidative

metabolic mechanism.

The most efficient MR_{ATP} comes from mitochondrial oxidative phosphorylation (Fig. 2 of ref (15)), and MR_{AWC}

should be even more sensitive to this. Indeed, *in vitro* model brain tissue studies indicate $k_{io}V$ correlates strongly with

mitochondrial MR_{O2}(consumption) (31). This is shown in **Figure 3** in the context of MR_{NKA} stimulation *via* $[K_o^+]$ titration

(Fig. 2). The vertical axis measures the population-averaged $\langle k_{io}\rangle_n$ for organotypic, cultured [spiking] rat somatosensory

cortex superfused with a paramagnetic agent. This SS-NMR study also allowed the estimation that $\langle V\rangle_n$ was rather

constant during the titration (Fig. 3D of ref. (31)): thus $\langle k_{io}\rangle_n$ was proportional to $\langle k_{io}V\rangle_n = \langle {}^cMR_{AWC}\rangle_n$. The horizontal axis

measures $\langle {}^tMR_{O2}\rangle_n$, directly determined in completely independent $[K_o^+]$ titrations of isolated rat brain synaptosome

suspensions. It is the $[K_o^+]$ titrations that allow this correlation: these are shown as outer ordinate and abscissa scales.

They are non-linearly (Michaelis-Menten) related to the inner $\langle k_{io}\rangle_n$ and $\langle {}^tMR_{O2}\rangle_n$ scales, which are linear. The correlation

is excellent over most of the range. (It is interesting O_2 consumption continues in the synaptosome suspensions even

when $[K_o^+]$ is zero. This is not the case for k_{io} in the SS-NMR study of cultured cortex preparations.) The quantity $k_{io}V$ is

probably best thought of as a high-resolution *in vivo* measure of mitochondrial function.

In Figure 2, AWC participation of Fig. 1 enzymes is exemplified by SGLT (both a type III and a type IV transporter)

and by KCC4 (a type II and type I enzyme). These transporters share substrates with NKA; Na_o^+ for SGLT and K_i^+ for KCC4.

The SGLT and KCC4 reactions are thereby coupled with the NKA reaction. As individual proteins, I and IV respectively are

catalyzing *unidirectional* water effluxes and influxes. Working in concert with other transporters, and orchestrated

Figure 3. A correlation of k_{io} with ${}^tMR_{O2}$. Independent $[K_o^+]$ titrations of paramagnetic agent-superfused [SS-NMR] studies of organotypic, cultured [spiking] rat somatosensory cortex [ordinate] and direct studies of isolated rat brain synaptosome suspensions [abscissa] allow correlation of the population-averaged $\langle k_{io} \rangle_n$ and $\langle {}^tMR_{O2} \rangle_n$ quantities, respectively. The outer $[K_o^+]$ scales are non-linearly related to the linear, inner $\langle k_{io} \rangle_n$ and $\langle {}^tMR_{O2} \rangle_n$ scales due to their Michaelis-Menten relationships. The correlation of $\langle k_{io} \rangle_n$ and $\langle {}^tMR_{O2} \rangle_n$ is very strong. The SS-NMR studies also indicate the mean cell volume $\langle V \rangle_n$ is rather constant. Thus, $\langle k_{io}V \rangle_n = \langle {}^cMR_{AWC} \rangle_n$ (pL(H_2O_i)/s/cell) correlates with $\langle {}^tMR_{O2} \rangle_n$ (pmole(O_2)/s/mL(suspension)). (This is a combination of Figures 3C and 3E of reference (31), where details are provided. The points are synaptosome measurements: the dashed line is the Michaelis-Menten fitting (K_m = 4.5 mM) of the cortical culture measurements.)

226   by NKA activity however, they produce overall cellular homeostatic water *bidirectional* exchange. The cellular AWC

227   fluxes are given in **Equation (2)**. This is a "systems" aspect of the cell. Though none of the individual process metabolic

228
229

$$\frac{^cMR_{Na+}(effl)}{3} = \frac{^cMR_{K+}(infl)}{2} = \frac{^cMR_{H2O}(effl)}{x} = \ ^cMR_{NKA} = \frac{^cMR_{H2O}(infl)}{x} = \frac{^cMR_{K+}(effl)}{2} = \frac{^cMR_{Na+}(infl)}{3} \qquad (2)$$

rates (^cMR's) are equal to $^cMR_{NKA}$, they rise or fall with the latter. For example, $^cMR_{K+}(effl) = 2^cMR_{NKA}$ and $^cMR_{H2O}(effl) =$

$x^cMR_{NKA} = [H_2O_i]k_{io}V$. Thus, $^cMR_{H2O}(effl) = \ ^cMR_{H2O}(infl) = x^cMR_{NKA}$. NKA is the driver.

Molecular Dynamics (MD): *In silico* simulations of Fig. 1 transporter mechanisms using MD (29,30,32-37) can be very

informative. This is particularly so for studies of an SGLT enzyme: the bacterial transporter vSGLT forms a hydrophilic

membrane-spanning channel (32-36). Generally, the enzyme exhibits spasmodic (tens of ns) bursts, some featuring

net water unidirectional influx and some featuring bidirectional water *exchange* (33-35). That is, a given transporter can

indeed facilitate the net flux (category B) and the exchange (category A) processes at different times (32).

For a period after sodium ion release in the vSGLT sugar-bound state, water molecules pass inward and

outward, past the galactose molecule (a glucose stereoisomer) in the channel, in almost equal, small numbers

(~one molecule/ns) - homeostatic exchange. The sugar is bound to its site midway through the channel, while a Na$^+$

binding site is near the cytoplasmic mouth. In the ~100 ns after galactose is released, it proceeds through the channel

into the cytoplasm, and pushes the channel-filling water molecules before it <u>into</u> the cell - net influx - in a quantity

(33,35) generally consistent with the Fig. 1 SGLT water stoichiometries. Both such periods (exchange and net flux) are

only transient (34). However, both seem triggered by Na$^+$ release.

Since the common role of the Fig. 1 enzymes is to facilitate passage of hydrophilic substrates across

the hydrophobic bilayer membrane, it is likely they all have somewhat similar hydrophilic pathways. The structure

of the NKCC1 transporter channel has been determined (38). We surmise there are many more water co-transporting

membrane enzymes, possibly including Na$^+$,K$^+$-ATPase (NKA) itself, as yet unstudied for this aspect. The NKA channel

structure has been elucidated (39). Transient exchange periods simulated for AQP-1 (37) are very similar to those

for vSGLT (34). It is likely each of the Fig. 1 transporters can accomplish what AQPs accomplish, but in addition

the conduct of specific metabolites across the membrane. Water molecules will find their ways through any open

hydrophilic pathway ("pore").

The aforementioned vSGLT MD simulations also suggest possible molecular mechanisms for SGLT/NKA *kinetic*

coupling. As stated, large tranches of net water molecule influx (category B) generally occur only upon galactose release

from its site near the middle of the hydrophilic channel. But, the probability for this, and for exchange (category A), can

depend on whether a Na^+ ion is in a site: the Na^+ can block the sugar (and apparently *net* water) ingress into the cell

(35,36). More often than not, Na^+ release occurs before sugar release (32). At least 80 ns after the Na^+ release (35,36),

galactose release can occur – unless the Na^+ ion has re-bound or has been replaced by another Na^+ ion. While galactose

is still present, the duty cycle of the Na^+ site remaining empty depends on the intracellular sodium concentration [Na_i^+]:

this Na^+ can come from only inside the cell. With a normal $^cMR_{NKA}$, [Na_i^+] is small (see below). However, [Na_i^+] increases

as $^cMR_{NKA}$ decreases (40). Thus, the probability of the Na^+ site being re-occupied within 80 ns increases as $^cMR_{NKA}$

decreases. As a consequence, the sugar-release frequency decreases as $^cMR_{NKA}$ decreases. In this way, MR_{SGLT} and

MR_{NKA} are positively correlated kinetically. (This pair-wise SGLT/NKA consideration assumes all other Figs. 1,2 enzyme

activities remain unchanged.) Irrespective of the details, the sharing of any substrate with NKA should facilitate kinetic

coupling. The correlation mechanisms of other Fig. 1 water co-transporter activities with MR_{NKA} must bear some general

similarity to this. Pore openings and closings are metabolically controlled.

A Futile Cycle? When they were first realized, the Fig. 2 Na^+ and K^+ exchange processes may have been briefly viewed

as "futile cycles." This term is used to characterize a set of coupled biochemical reactions that consume ATP with

no apparent benefit (41). Obviously, it is generally thought biological evolution discards futile cycles as "energy

wasting." However, it was quickly understood the trans-membrane ion concentration gradients enjoyed by Na^+ and K^+

serve to partially store chemical (Gibbs) free energy released by NKA-catalyzed ATP hydrolysis, $\Delta G_{ATP} = -59$ kJ/mole

(42,43). As we will see below, $\Delta G_{Na}(infl)$ is typically near -14.5 kJ/mole, for Na^+ influx, and $\Delta G_K(effl)$ near 0 kJ/mole,

for K^+ efflux. Thus, though these cycles are *homeostatic*, they are not in *equilibrium*, but in *steady-state*. Energy is

required to maintain them. Subsequently, the free energy thus stored in the [Na^+] gradient is used to facilitate cellular

uptake of many crucial molecules (Fig. 1), and these processes help complete the Na^+ cycle. The free energy in the [K^+]

gradient is small only because of the cell's membrane potential, which the homeostatic K^+ efflux largely produces and

maintains (see below).

The "pump and leak mechanism" term used for water (19,27) implies waste. However, a question naturally

arises. Might the existence of active trans-membrane water cycling (Fig. 2) reveal an analogous metabolic energy

storage function? Is AWC an actual example of a futile cycle, or does it have some evolutionary advantage?

THEORY:

Reaction Free Energy Changes (ΔG): The issue of AWC futility is a chemical thermodynamic question. Thus, we must

consider ΔG for each permeant as it crosses the membrane, for example, into the cell; ΔG_{per}(infl), for *permeant* (per)

influx (infl). For transport reactions (as in Fig. 1), the standard free energy change (ΔG^0) is zero. There are potential

chemogenic, electrogenic, and barogenic ΔG contributions. These are, respectively, the three terms on the right-hand-

sides of **Equations (3)**; where i and o are, respectively, the inside and outside compartmental indices, [per] is

287
$$\Delta G_{per}(infl) = RTln\left\{\frac{[per_i]}{[per_o]}\right\} + Z_{per}FE_{m,oi} + RTln\left\{\frac{P_i}{P_o}\right\} = 2.58ln\left\{\frac{[per_i]}{[per_o]}\right\} + 0.0965Z_{per}E_{m,oi} + 2.58ln\left\{\frac{P_i}{P_o}\right\} \quad \left(\frac{kJ}{mole}\right) \quad (3)$$

the compartmental permeant concentration, Z_{per} the (signed) permeant particle electrical charge, F is the Faraday

constant (0.0965 kJ/(mV•mole) at physiological temperature, 310 K), $E_{m,oi}$ is the (signed) trans-membrane electrical

potential (in minus out) in mV, P_i/P_o is the intracellular/extracellular hydraulic pressure ratio (Figs. 1,2) ((44), p. 35), and

RT = 2.58 kJ/mole at 310 K. For H_2O as permeant, $Z_{H2O} = 0$, and the second term makes no contribution.

Equations (3) can be written alternatively with permeant *chemical potential* changes; $\Delta\mu_{per}$, where

$\mu_{per} \equiv (\partial G/\partial n_{per})_{T,P,n(\neq per)}$ ((44), p. 20). However, the ΔG notation is more inclusive, and more common in this field (42,43).

To obtain the free energy change for any Fig. 1 *reaction*, ΔG_{reac}(infl), one must sum over all permeants,

Equation (4); where s_{per} is the permeant stoichiometric coefficient (Fig. 1). It is obvious from Eqs. (3,4) the chemogenic

296
$$\Delta G_{reac}(infl) = \sum_{per} s_{per}\left[2.58ln\left\{\frac{[per_i]}{[per_o]}\right\} + 0.0965Z_{per}E_{m,oi} + 2.58ln\left\{\frac{P_i}{P_o}\right\}\right] \quad \left(\frac{kJ}{mole}\right) \quad (4)$$

terms require realistic compartmental permeant concentrations. Strictly, thermodynamic activity

(a = the concentration•activity coefficient product) values should be used. Below, and in **A2**, we justify the first-order

approximation that solute and solvent activity coefficients are each unity.

Concentration Scales: In this paper, a concentration is indicated with square brackets, *e.g.*, [permeant]. Sometimes,

the molarity scale, with unit M = moles(permeant)/L(compartment) (activity coefficient y_{per} ((45), pp. 26,27)) is used.

However, first-order evaluation of chemogenic terms in Eqs. (3) and (4) consider only the entropy of mixing.

This requires concentration scales with mole ratio- or mole fraction (X)-like quality. Molality, with unit

m = moles(permeant)/kg(water) (activity coefficient γ_{per} ((45), pp. 26,27)), is one such scale. Biological aqueous solutions

are sufficiently dilute in each solute that its m value is well-approximated by its M value; the *infinite dilution* assumption.

The use of an entropy-centric concentration scale here is tantamount to assuming solution "ideality;" there are

no specific solute/solute interactions. We will see below this is clearly not the case. The "volume molality" scale,

$_vm$ = moles(permeant)/L(solvent) ((44), p.37) will also be considered. (Using m or $_vm$ concentration values is tantamount

to assuming the small molecule metabolites may sample the entire compartmental space or only aqueous spaces,

respectively, (*i.e.*, the water-excluding macromolecules are more or less mobile or immobile (14).)   Since all

the different neutral and ionic solutes (thus, osmolytes plural) need be counted, we will further use the OsMolarity

(unit, OsM), or Osmolality (unit, Osm) concentration scales.

Since we also consider the solvent water as a permeant (Fig. 1), we must express its concentration.

For understanding water thermodynamic activity – its "*escaping tendency*" - the dimensionless X (or a related) scale is

best ((44), p. 56; (45), p.27; (46)).   Thus, for water, X_{H2O} = (moles(water))/(moles(osmolytes + water)), activity coefficient

f_{H2O} ((45), pp. 26,27)), is used.

One will further see milli- scales: mOsM, mOsm, and mOsX.

Chemogenic Contributions

Tissue Osmolyte Compartmental Concentrations.   For *in vivo* conditions, accurate determination of biological

compartmental solute concentrations is difficult, particularly for intracellular metabolites.   Recent metabolomic and

proteomic advances (for example, using calibrated liquid chromatography/mass spectrometry techniques) have given

realistic accountings.   From such a study of cultured murine kidney epithelial cells, 101 different intracellular small

molecule metabolites, and their concentrations, have been itemized (47).   We compile an inventory of intra- and extra-

cellular solute concentrations from a number of related studies (47-62) in **Table 1**.   The inorganic ion (Na$^+$, K$^+$, H$_3$O$^+$, Ca^{2+},

Mg^{2+}, Cl$^-$, HCO$_3^-$) concentrations are conventional.   The units are mOsM (thus, mOsm for osmolytes).   These should

be considered representative of interstitial and intracellular compartment osmolytes.   Of course, they are not the values

for any actual cell.   Concentrations surely differ from cell to cell, tissue to tissue, and in different biological conditions.

Since we are calculating general thermodynamic trends, the Table 1 values suffice.

We see the intracellular RNA, lipid assembly, and DNA mOsM concentrations are tiny.   Though these occupy

considerable intracellular volume (see below), because of their very large macromolecular masses they make negligible

osmolal contributions.   Since there are fewer (but non-zero: see below) extracellular macromolecules, we ignore their

osmolal contributions here.

Interestingly, we see the total osmolyte concentrations are ~0.3 OsM for the interstitial space and ~0.4 OsM

for the intracellular space.   It is widely assumed intracellular osmolarity is near 0.3 (52), and thus similar to that

in the extracellular space.   The implications of this trans-membrane osmolarity gradient are explored in the section

on intracellular pressure (P_i) below.

Table 1. Representative Osmolyte Concentrations (mOsM)[a-d]		
	interstitial	intracellular
Na^+	135.0	15.0
K^+	5.0	150.0
H_3O^+	0.00005	0.0001
Ca^{2+}	1.5	0.0001 [free]; 0.01 [total]
Mg^{2+}	0.5	0.5 [free]; 10 [total]
other cations	1.5	3.5
Cl^-	105.0	15.0
HCO_3^-	30.0	10.0
small molecule metabolite anions	3.0	135.0[a,b]
protein anions	7.5[a]	9.5[a,c,d]
small molecule metabolites		40.0[a,b]
RNA		0.5[d]
lipids		0.05[d]
DNA		0.000005[d]
total osmolytes	289	379
H_2O mole fraction (OsX_W)	0.995	0.993
H_2O millimolarity (mM)	45,000[e]	42,000[f]
f_W	$f_{W,o} = 45/55 = 0.82$	$f_{W,i} = 42/55 = 0.76$
osmotic stoichiometry (H_2O/osmolyte)	45/0.289 = ~156	42/0.379 = ~111
References (47-62). [a]ref. (48), [b]ref. (47), [c]ref. (49), [d]ref. (50), [e]ref. (62), [f]ref. (51).		

Tissue Water Compartmental Concentrations. Experimentally determining biological compartmental water contents is
even harder than for solutes (2). For many years, it has been stated, usually anecdotally, that cell water content is
70% (v/v) [or (w/w), assuming unit density, d] (53). Vinnakota and Bassingthwaighte presented a comprehensive review
and tabulation of rigorous, quantitative determinations of myocardial tissue compartmental densities, volumes, volume
fractions, and mass fractions (51). These derive from direct *ex vivo* mass analyses and *in vivo* tracer studies, and yield
the cell water content value 75 % (w/w). A more recent optical determination yields 85 % (v/v) for cultured mammalian
cell controls (54).

If we use the 75 % value (51) and take d = 1.1 g/mL, we obtain $[H_2O_i]$ = 42 M; *i.e.*, 42 moles(H_2O)/L(cell). This is
the most parsimonious interpretation of the data, and includes water in the cytoplasm, mitochondria, and sarcoplasmic
reticula. The magnitude is considerably reduced from the value for pure water, 55 M. The intracellular water volume
fraction, $f_{W,i}$, estimated from this is 42/55 = 0.76 (Table 1). This is in good agreement with quantitative MRI *in vivo*
measures (55). (Note: f_W is different from f_{H2O}.) Rand states, without attribution, $[H_2O_i]$ = 54 M (56).

From tracer ($^{15}OH_2$) studies (reviewed in (51)) and NMR (^1H_2O) studies (reviewed in (2)), essentially all
intracellular water is effectively "well-mixed," at homeostasis. There has been much consideration of "bound" water
in tissue (reviewed in (25,57-59)). There is no doubt water near macromolecule and membrane surfaces is
thermodynamically different from bulk water – it has at least lower entropy (see below) – but kinetically (tumbling- and
diffusion-wise) it remains in very rapid communication with all the water in the cell. The fraction of water molecules
in "buried" intra-macromolecular sites inside cells is miniscule. Though the latter have significant NMR consequences
(reviewed in (25)), even these water molecules have labile access to the cytoplasm (60).

Because of its small volume fraction (typically, 0.2) the extracellular compartmental water content
in parenchymal tissue has been even more difficult to determine. Vinnakota and Bassingthwaighte report myocardial
interstitial water content 92 % (w/w) and d = 1.0 g/mL, but this has a significantly greater uncertainty (51) and seems
excessively large. The water content of cartilage can be more directly determined, and this might be a good
extracellular model: it is an essentially acellular matrix of collagen fibrils (61). Lu and Mow imply a cartilage water
content value of 78 % to 85 % (61). Maroudas and co-workers carefully measured the value 81 % for unloaded human
femoral head cartilage (62). Again assuming unit density, this yields $[H_2O_o]$ = 45 M, also significantly reduced from 55 M,
though not as much as $[H_2O_i]$. The corresponding $f_{W,o}$ is 45/55 = 0.82 (Table 1).

In Table 1, we give intra- and extracellular "osmotic stoichiometry" values; the overall ratio of water molecules
to osmolyte particles in the compartment. We see the intra- and extracellular values are near 100 and 150 water
molecules per osmolyte, respectively. The fact that in Fig. 1 transporter osmotic stoichiometries for the reactants are
either above or below these values is consistent with the water stoichiometry being principally determined
by the hydrophilic channel volumes (35). The molecular action of any particular transporter has nothing to do with

the overall "osmotic stoichiometry." It is only the *net* water transported by all influxers and effluxers in the cell that

must be zero for homeostasis. However, even if values of 100 or 150 did obtain, this would be irrelevant

for the calculations below.

The large macromolecular volumes are the reason the 42 M intracellular H_2O concentration (Table 1) used

below is so much less than that of pure water (55 M). These large molecules exclude considerable water (Fig. 8 of (15)).

Obviously, there are also water-displacing extracellular macromolecules: 45 M (Table 1) is also smaller than 55 M.

However, the molar concentration scale is useful mainly for molecular counting (1 mole = 6 x 10^{23} molecules).

As justified above, we use the dimensionless mole fraction concentration scale for water (X_{H2O}). (This differs from

the tissue compartmental water mole fractions ("populations") p_i and p_o (important in SS-NMR).) For most practical

aqueous solutions, including those in biological compartments, however, X_{H2O} is always very nearly unity. "In dilute

solutions of electrolytes, the activities, and even the activity coefficients (f_{H2O}), of the *solvent* are little different

from unity" ((63), p.12). Very important consequences of this are considered below.

One kg of water is (1000/18.0154) 55.508 moles H_2O. Dividing this by (55.508 + the sum of osmolyte

concentrations) in each of the cytosolic and interstitial compartments yields corresponding water mole fractions

$OsX_{H2O,i}$ = 0.993 and $OsX_{H2O,o}$ = 0.995 (Table 1). $OsX_{H2O,i}$ is only 0.2% smaller than $OsX_{H2O,o}$, and each value is nearly that

for pure water. Given this situation, following Pitzer ((63), p.12), it is not unreasonable to assume equal activity

coefficients ($f_{H2O,i}$ = $f_{H2O,o}$ = 1) and thus use these mole fraction values in the Eqs. (3) and (4) terms for water as permeant.

Inserting {$OsX_{H2O,i}/OsX_{H2O,o}$} = 0.998 yields ΔG_{H2O}(infl) = **-** 0.00517 kJ/mole for the *chemogenic* contribution.

Therefore, unlike the other Table 1 species, water is almost (but not exactly) in trans-membrane chemical

equilibrium. (In this paper, "*chemical*" equilibrium is defined as: [per$_i$] = [per$_o$].) There is (only) ~5 J/mole favoring

cellular water influx. Strictly, this chemogenic ΔG_{H2O}(infl) value might be considered not significantly different from zero

(chemical equilibrium). However, analyses of surrogate solutions in **A2** give water activity values: $a_{H2O,i}$ = 0.994 and

$a_{H2O,o}$ = 0.995. These correspond to chemogenic ΔG_{H2O}(infl) = **-** 0.003 kJ/mole. Henceforth, we will use − 3 J/mole.

As we will see, this has no bearing on the important metabolic consequences detailed below. For co-transported water,

the barogenic contribution will be much greater than the chemogenic term.

It is probably unreasonable that chemogenic ΔG_{H2O}(infl) be positive. First-order osmotic considerations more

strongly favor water influx. However, the first-order is inadequate, see below.

Electrogenic Contributions

We make the common approximation that K^+ *efflux* permeabillity dominates over Na^+ and Cl^- permeabilities (in the P_f sense) in cellular homeostasis. Thus, we use a version of the Nernst **Equation (5)** ((44), p. 113) to calculate $E_{m,oi}$,

$$E_{m,oi} = -\left(\frac{RT}{FZ_{K+}}\right)\left\{ln\left(\frac{[K_i^+]}{[K_o^+]}\right)\right\} = \left(\frac{2.58}{0.0965}\right)\left\{ln\left(\frac{150}{5}\right)\right\} = -91\ mV \quad\quad (5)$$

where: RT = 2.58 kJ/mole and the Faraday constant F = 0.0965 kJ/mV•mole (both at 310 K) and insert the Table 1 $[K^+]$ values. Z must be the value for the potential-producing ion, Z_{K+} = 1. It is important to note the actual number of *excess* K_o^+ ions (and an equal number of uncompensated intracellular anions) is insufficient to significantly alter the Table 1 $[K^+]$ values. An excess of only 0.02 mM intracellular anionic charges (0.014% = ((100x0.02)/145); Table 1) produces $E_{m,oi} \approx$ - 100 mV for a small (1 fL) spherical cell ((53), p. 198.)

Barogenic Contributions

Intracellular pressure (P_i) can be counteracted with an externally applied (macroscopically uniform) mechanical pressure, in principle ((44), p.32), and elegantly transduced into an isotropic, hydraulic (mechanical) pressure with cellular insertion of a pressure-sensitive "osmotic" microelectrode (64,65). The trans-membrane pressure difference ($P_i - P_o$) force is oriented roughly vectorially radial to the cell surface. Experimentally, most mammalian cells have (P_i - P_o) values under 0.1 atm (27). However, an osmotic microelectrode study of perfused *ex vivo* murine lens (featuring relatively simple lens cells) found (P_i - P_o) values up to 0.5 atm (66). Most P_o values are near 1 atm: even "high" tumor interstitial P_o is only 1.02 atm. (67). For water, the standard state (f_{H2O} truly 1) pressure, P^0, is 1.0 atm.

RESULTS:

Free Energy Calculations

Permeant **Influx Free Energy Changes, Eq. (3).** In **Table 2**, we list many potential non-metabolite permeant (per)
species, and calculate chemogenic, electrogenic, and barogenic ΔG(infl) contributions (kJ/mole) for their influx.
For these calculations, we use the Table 1 water OsX_W and solute mOsm concentrations for the chemogenic terms.
We calculate the barogenic contributions for P_i/P_o ratios of 1.00 (blue) and 1.05 (red).

It is clear that, compared with those of the other permeants, the chemogenic and electrogenic contributions
for individual H_2O molecule transport are negligible. We see the slight chemogenic tendency for water molecule influx
when $P_i = P_o$ is reversed when $P_i \geq 1.05\ P_o$. In fact, for ΔG_{H2O}(infl) = - 0.003 kJ/mole, any $P_i > 1.001\ P_o$ would result
in water efflux if there were no other considerations.

Let us consider two other potential permeants, Na_o^+ and K_o^+ as examples. The electrogenic terms for their
influxes are identical, - 8.78 kJ/mole. Electrostatically, influx of each is favored. However, while the chemogenic Na_o^+
contribution (- 5.67 kJ/mole) also favors influx, the chemogenic K_o^+ term (8.78 kJ/mole) opposes influx. Thus,
the *electrochemical* potential gradient for Na_o^+ influx is very favorable, − 14.5 kJ/mole (- 5.67 − 8.78), but zero for K_o^+
influx (or K_i^+ efflux) when $P_i = P_o$. The barogenic contributions for Na_o^+ and K_o^+ are identical, but this has very small effect
for only Na_o^+. However, compared with Na_o^+, and indeed with all the other Table 2 permeants except K_o^+, the barogenic
contribution for H_2O is *proportionally* much greater than the chemo- and electrogenic terms. (For example, a P_i
of 1.05 atm causes a more than 40-fold increase in ΔG_{H2O}, but only a 1% change in ΔG_{Na}.) This has very important
consequences.

Transport *Reaction* **Free Energy Changes, Eq. (4).** We use Eq. (4) to calculate the total $\Delta G_{transporter}$(infl) for each
of the Fig. 1 transporter reactions - utilizing the stoichiometries seen there. The "metabolite" substrates (glucose,
glutamate⁻, GABA, and lactate⁻) are omitted because we study below how their transport is affected
by the thermodynamic contributions of the other "ancillary" (support) substrates.

Influx Reactions. Reactions that normally run in the *influx* direction (IV, III, or both, in Figs. 1,2) are presented in **Table 3a**
($E_{m,oi}$ is fixed at − 91 mV). Each is favorable (negative ΔG) when $P_i = P_o$ (blue). However, they each become less favorable
as P_i/P_o increases, and all become unfavorable (positive ΔG) when P_i has reached 1.05 P_o (red). This is due to P_i
opposition to water co-influx (the factors that are - 0.003 when $P_i = P_o$ become positive when P_i exceeds 1.001 P_o).
Because of the large water stoichiometries, the barogenic contribution can have a significant influence on secondary
active water co-transporter thermodynamics. This is due to essentially only the water co-transport. For the other
permeants, the blue/red values generally have very similar magnitudes.

Table 2. Free Energy Change for Influx, Eq. (3).				
potential permeant	ΔG_{per}(infl) (kJ/mole)			
	chemogenic	electrogenic	barogenic	sum
			$P_i/P_o =$ 1.00 ∥ 1.05	$P_i/P_o =$ 1.00 ∥ 1.05
H_2O_o	-0.003	0	0 ∥ 0.13	-0.003 ∥ 0.13
Na_o^+	-5.67	-8.78	0 ∥ 0.13	-14.5 ∥ -14.3
K_o^+	8.78	-8.78	0 ∥ 0.13	0 ∥ 0.13
$H_3O_o^+$	1.79	-8.78	0 ∥ 0.13	-6.99 ∥ -6.86
Ca_o^{2+}	-24.8	-17.6	0 ∥ 0.13	-42.4 ∥ -42.3
Cl_o^-	-5.02	8.76	0 ∥ 0.13	3.74 ∥ 3.87
$HCO_{3,o}^-$	-2.83	8.76	0 ∥ 0.13	5.93 ∥ 6.06
Concentrations in Table 1. $E_{m,oi}$ = -91 mV				

Table 3a. Transport Reaction Free Energy Changes, $\Delta G(infl)$ (kJ/mol); Eq. (4)   ($P_i/P_o = 1.00 \,||\, 1.05$)

influxer reactions [III, IV, or both, in Fig. 1] w/o metabolic substrates

enzyme	substrates and free energies				net
hSGLT1	glucose$_o$ +	2 Na$_o^+$ +	235 H$_2$O$_o$ $\rightarrow$		
		2(-14.5 || -14.3)	235(-0.003 || 0.13)		
		(-29 || -28.6) +	(-0.71 || 31) =		(-29.7 || 2.4)
rSGLT1	glucose$_o$ +	2 Na$_o^+$ +	380 H$_2$O$_o$ $\rightarrow$		
		2(-14.5 || -14.3)	380(-0.003 || 0.13)		
		(-29 || -28.6) +	(-1.1 || 49) =		(-30 || 20.4)
GLUT1	glucose$_o$ +		40 H$_2$O$_o$ $\rightarrow$		
			40(-0.003 || 0.13)		
			(-0.12 || 5.2) =		(-0.12 || 5.2)
GLUT2	glucose$_o$ +		100 H$_2$O$_o$ $\rightarrow$		
			100(-0.003 || 0.13)		
			(-0.30 || 13) =		(-0.30 || 13)
EAAT1	glutamate$_o$ +	2 Na$_o^+$ +	440 H$_2$O$_o$ +	- K$_i^+$ + H$_3$O$_o^+$ $\rightarrow$	
		2(-14.5 || -14.3)	440(-0.003 || 0.13)		
		(-29 || -26.8) +	(-1.3 || 57) =	(0 || -0.13) + (-6.99 || -6.86) =	(-37 || 23)
GAT1	GABA$_o$ +	2 Na$_o^+$ +	330 H$_2$O$_o$ +	Cl$_o$ $\rightarrow$	
		2(-14.5 || -14.3)	330(-0.003 || 0.13)		
		(-29 || -26.8) +	(-0.99 || 43) =	(3.74 || 3.87) =	(-26 || 20)
NKCC1	2Cl$_o$ +	Na$_o^+$ +	590 H$_2$O$_o$ +	K$_o^+$ $\rightarrow$	
	2(3.74 || 3.87)		590(-0.003 || 0.13)		
	(7.48 || 7.74)	(-14.5 || -14.3)	(-1.77 || 77)	(0 || 0.13)	(-8.8 || 71)

All stoichiometries from reference (19). ΔG_{per}, from Table 2. $E_{m,oi}$ = -91 mV

Table 3b, Transport Reaction Free Energy Changes, ΔG(effl) (kJ/mol), Efflux Version of Eq. (4) (P_i/P_o = 1.00 $\|$ 1.05)					
effluxer reactions (I, II, or both, in Fig. 1) w/o metabolic substrates					
KCC4	K_i^+ +	Cl_i^- +	500. H_2O_i $\rightarrow$		net
			500(0.003 $\|$ -0.13)		
	(0 $\|$ -0.13) +	(-3.74 $\|$ -3.87) +	(1.5 $\|$ -65) =		(-2.2 $\|$ -69)
MCT1	lactate$_i^-$ +	$H_3O_i^+$ +	500 H_2O_i $\rightarrow$		
			500(0.003 $\|$ -0.13)		
		(6.99 $\|$ 6.86) +	(1.5 $\|$ -65) =		(8.5 $\|$ -58)
NaDC1	dicarboxylate$_i^{2-}$ +		175 H_2O_i +	2 Na_i^+ $\rightarrow$	
			175(0.003 $\|$ -0.13)	2(14.5 $\|$ 14.3)	
			(0.53 $\|$ -23) +	(29 $\|$ 29) =	(30 $\|$ 6.0)
All stoichiometries from reference (19). -ΔG_{per}(infl) from Table 2. $E_{m,oi}$ = -91 mV					

<u>Glucose Uptake at Fixed [glucose$_i$].</u> For the rSGLT1 influx reaction as an example: ΔG = **- 30 kJ/mole** when $P_i = P_o$,
but 20.4 kJ/mole when $P_i = 1.05\ P_o$. ($P_i = 1.03\ P_o$ is sufficient to make influx unfavorable.) To illustrate the consequences
of this on glucose uptake, we fix the intracellular glucose concentration, [glc$_i$], at 2 µM. (The intracellular enzyme that
phosphorylates glucose, hexokinase, has a glucose$_o$ K_m value reported as 1.7 µM (20). A principle of central carbon
(glucose-related) metabolism is that any intracellular metabolite steady-state concentration should be near the K_m value
of the enzyme that consumes it (47).) Thus, we write Eq. (4) explicitly for the 2Na$^+$/glucose co-transporter **rSGLT1** influx
free energy change, ΔG$_{rSGLT1}$(infl), in **Equation (6)**. The first term represents the glucose barochemical contribution

$$\Delta G_{rSGLT1}(infl) = \left[2.58ln\left\{\frac{2}{[glc_o]}\right\} + 2.58ln\left\{\frac{P_i}{P_o}\right\}\right] + 2\left[-5.67 - 8.78 + 2.58ln\left\{\frac{P_i}{P_o}\right\}\right] + 380\left[-0.003 + 2.58ln\left\{\frac{P_i}{P_o}\right\}\right] \quad (6)$$

with [glc$_i$] fixed at 2 µM. The second and third terms are due to, respectively, the Na$^+$ electro- and barochemical and
the H$_2$O barochemical potential gradients. The former favors influx, while the latter opposes it.

If we fix $P_i = 1.05\ P_o$, and neglect the water co-transport, ΔG becomes positive (*i.e.*, influx ceases) only when
the extracellular glucose concentration [glc$_o$] decreases below 0.33 nM. On the other hand, if we keep $P_i = 1.05\ P_o$
but retain the water co-transport, the reaction becomes unfavorable when [glc$_o$] decreases below 2.4 mM –
three orders of magnitude *greater than* [glc$_i$]. The large electrochemical ΔG$_{Na+}$ favoring influx (- 28.6 kJ/mole) is
counteracted by the even larger barochemical ΔG$_{H2O}$(infl) favoring efflux (49 kJ/mole). A number of significant
consequences of this will be considered in the Discussion section.

To partially generalize these considerations, **Figure 4** shows a 3D plot of Eq. (6) for the rSGLT1 reaction. For this,
the [Na$_o^+$] and [Na$_i^+$] values were fixed at those in Table 1, [glc$_i$] at 2 µM, and E$_{m,oi}$ at − 91 mV. The vertical axis plots
the rSGLT1 influx reaction ΔG$_{rSGLT1}$(infl), while the (logarithmic) oblique axes increment [glc$_o$] and P_i/P_o. The free energy
surface is colored green when influx is possible (ΔG$_{rSGLT1}$(infl) < 0), and red when impossible (ΔG$_{rSGLT1}$(infl) > 0).
The intersection of the ΔG$_{rSGLT1}$(infl) surface with the horizontal ΔG$_{rSGLT1}$(infl) = 0 plane defines the trajectory
of the (P_i/P_o)-dependence of the minimum extracellular glucose concentration required for influx. While the exponential
nature of the (P_i/P_o)-dependence is hardly noticeable, that of [glc$_o$] is very evident.

To make these effects clearer, **Figure 5** shows the 2D plot of the Fig. 4 ΔG$_{rSGLT1}$(infl) = 0 plane. With H$_2$O
co-influx, increasing P_i/P_o strongly increases [glc$_o$]$_{min}$. For $P_i = 1.04\ P_o$, [glc$_o$]$_{min}$ is already ~10 µM (*greater than* [glc$_i$] here)
for rSGLT1. The location of the hexokinase glucose$_i$ K_m is shown as a horizontal dashed line. Of course, the position
of the surface depends also on the E$_{m,oi}$, [glc$_i$], and [Na$_i^+$] values, among other quantities. For example, making ΔG$_{Na}$(infl)
less negative (say, by increasing [Na$_i^+$]) would increase [glc$_o$]$_{min}$, all other factors being equal – a given P_i/P_o, for example.

Plausible water chemogenic ΔG$_{H2O}$(infl) valies (- 5 to 5 J/mole) are small compared with a typical barogenic
magnitude (2.58ln(P_i/P_o) = 2.58ln(1.03) = 76 J/mole), which favors efflux. Thus, it is relatively inconsequential whether

Figure 4. A 3D plot for the rSGLT1 reaction. The vertical axis measures the Gibbs free energy change (ΔG) for the influx direction shown, and calculated with Eq. (6). The logarithmic oblique axes plot the extracellular glucose concentration, [glucose$_o$], and the intracellular/extracellular hydraulic pressure ratio (P_i/P_o) over the experimentally measured range. For this calculation, the Table 1 concentrations and the Tables 2 and 3a free energy terms were used (chemogenic ΔG_{H2O}(infl) = - 3 J/mole). The intracellular glucose concentration, [glucose$_i$], the membrane potential, $E_{m,oi}$, and P_o were held fixed at 2 μM, - 91 mV, and 1 atm, respectively (T = 310 K). The surface is colored green when influx is thermodynamically possible and red when it is impossible. Thus, the intersection of the ΔG surface with the ΔG = 0 plane traces the trajectory of the (P_i/P_o)-dependence of the *minimum*, [glucose$_i$]$_{min}$, value. The value of the intracellular hexokinase K_m (1.7 μM) for glucose$_i$ is indicated.

Figure 5. The 2D plot of the Figure 4 ΔG$_{rSGLT1}$(influx) = 0 plane. The regions where the Na$^+$ electrochemical gradient dominate and the H$_2$O barochemical gradient dominate are indicated. The dependence is so strong that small percentage changes of P$_i$ cause very large changes of the minimum [glucose$_o$] required for glucose uptake. The hexokinase K$_m$ for glucose$_i$ (1.7 μM) is indicated with a horizontal line.

water is in chemical equilibrium ($OsX_{H2O,i}$ = $OsX_{H2O,o}$) or not. The larger steady-state water barochemical potential

gradient is mostly barogenic.

Since it is likely almost all cells have P_i 1.01 atm or greater, it seems that each of the Table 3a reactions will have

similar behaviors. The water barochemical contributions provide crucial counterbalances for other influx transporters

as well.

Though the GLUT family of glucose influxers lacks the large SGLT Na^+ electrochemical potential gradient favoring

influx, it also has a smaller H_2O barochemical force opposing influx when $P_i > P_o$: the GLUT H_2O stoichiometries are

smaller (Fig. 1). **Figure 6** contrasts the 2D plot of GLUT1 with that of rSGLT1 (from Fig. 5) and makes obvious

for the SGLT family the Na^+ electrochemical dominance at small P_i values and the H_2O barochemical dominance

at large P_i.

<u>Glucose Uptake at Fixed [glucose$_o$].</u> For an alternative perspective, we take the extracellular glucose

concentration, [glc$_o$], to be maintained at 5 mM (a large, but reasonable (68,69), value). With rSGLT1, if we fix P_i =

1.05 P_o and neglect water co-transport, ΔG remains negative until the intracellular glucose concentration, [glc$_i$], reaches

the absurdly large value of 300 M.   On the other hand, if we keep P_i = 1.05 P_o but retain the water co-transport,

the reaction becomes unfavorable when [glc$_i$] reaches only 4.1 µM. More generally for rSGLT1 with H_2O transport,

Figure 7 shows the (P_i/P_o)-dependence of [glc$_i$]$_{max}$, with [glc$_o$] fixed at 5 mM. The location of the hexokinase glucose$_i$ K_m

is shown as a horizontal dashed line. For such a plot, in the lower portion of the graph, the influx reaction is now

possible (green), while it is impossible (red) in the upper portion. The maximum value of [glc$_i$] is greatly suppressed

by increasing P_i/P_o.

For GLUT2 at P_i = 1.04 P_o and [glc$_o$] = 5 mM, [glc$_i$] could accumulate to 4.8 mM if there was no water co-influx,

but only 110 µM with H_2O co-influx.

<u>Neurotransmitter Uptake and Clearance.</u> There are considerable similarities of the rSGLT and excitatory amino

acid transporter (EAAT1) influx reactions − with the exceptions of the H_3O^+ co-influx and K_i^+ co-efflux of the latter.

Glutamate$^-$ influx is greatly aided by the Na^+ electrochemical gradient, but opposed by the H_2O barochemical gradient.

The analogous EAAT1 3D plot (not shown) is very similar to that for rSGLT (Fig. 4). However, the roles of these

two reactions are quite different. While the purpose of rSGLT is mainly to deliver glucose into cells, that of EAAT1 is

principally to clear glutamate$^-$ from synapses (*via* astrocytic uptake) after action potential transmission.

Now, besides $E_{m,oi}$ = **-** 91 mV as before, we fix *intracellular* [glutamate$_i^-$] at 1.4 mM, the glutamate$_i^-$ K_m

for glutamine synthetase (70), which converts astrocytic glutamate$_i^-$ to glutamine$_i$. **Figure 8** depicts the 2D

$ΔG_{EAAT1}$(infl) = 0 plane of the unshown 3D plot. The vertical axis measures log *extracellular* [glutamate$_o^-$], while

the horizontal axis plots P_i/P_o. The intersection of the $ΔG_{EAAT1}$(infl) surface with the $ΔG_{EAAT1}$(infl) = 0 plane traces

Figure 6. The (P_i/P_o)-dependences of [glucose$_o$]$_{min}$ for rSGLT1 (from Fig. 5) and for GLUT1. Equation (4) was used, with the intracellular glucose concentration, [glucose$_i$], the membrane potential, $E_{m,oi}$, and P_o held fixed at 2 µM, -91 mV, and 1 atm, respectively (T = 310 K). The regions of Na$^+$ electrochemical and H$_2$O barochemical dominance are very evident. Influx through the rSGLT1 transporter is much more sensitive to P_i/P_o than that (barely noticeable) through the GLUT1 transporter. This is due to the much greater water stoichiometry of the former (Fig. 1).

Figure 7. The 2D plot of the ΔG_{rSGLT1}(influx) = 0 plane from a 3D plot similar to Figure 4. Equation (6) was used and, in this case, the *extracellular* glucose concentration, [glucose$_o$], the membrane potential, $E_{m,oi}$, and P_o were held fixed at 5 mM, -91 mV, and 1 atm, respectively (T = 310 K). The surface is colored green when influx is thermodynamically possible and red when it is impossible. Thus, the intersection of the ΔG surface with the ΔG = 0 plane traces the trajectory of the (P_i/P_o)-dependence of the (in this case) *maximum* intracellular glucose$^-$ concentration, [glucose$_i$]$_{max}$, value allowing uptake. The K_m value of the cytoplasmic hexokinase for glucose$_i$ (1.7 μM) is indicated with a horizontal dashed line. It is clear when P_i is small, tremendous values of [glucose$_i$] are allowed, which would easily saturate hexokinase. However, this is not the case when P_i is only a few percentage points greater.

Figure 8.   The 2D plot of the ΔG_{EAAT1}(influx) = 0 plane from the 3D plot analogous to Figure 4.   The (in this case) astrocytic glutamate⁻ concentration, [glutamate$_i$⁻], the membrane potential, $E_{m,oi}$, and P_o were fixed at 1.4 mM (the K_m value for glutamine synthetase), -91 mV, and 1 atm, respectively (T = 310 K). The surface is colored green when influx is thermodynamically possible and red when it is impossible. Thus, the intersection of the ΔG surface with the ΔG = 0 plane traces the trajectory of the (P_i/P_o)-dependence of the *minimum* synaptic glutamate⁻ concentration, [glutamate$_o$⁻]$_{min}$, value required for astrocyte uptake. The EC$_{50}$ value of the iGlutR receptor for synaptic glutamate$_o$⁻ (2.3 μM) is indicated with a horizontal dashed line. It is clear glutamate$_o$⁻ is well-cleared from the synapse when P_i is small, and not when P_i is only a few percentage points greater. In the latter case, the receptors would be saturated, and synaptic transmission would be interrupted.

the trajectory of the logarithm of the *minimum* glutamate$_o^-$ concentration ([glutamate$_o^-$]$_{min}$) necessary to initiate influx,

as a function of P$_i$/P$_o$. Though strictly bi-exponential, the plot is almost a single exponential function (the exponential

nature of the P$_i$/P$_o$ axis is very weak). For a member of the ionotropic glutamate receptor (iGlutR) family, the position

of the L-glutamate$_o^-$ EC$_{50}$ value, 2.3 μM (71), is shown as a horizontal dashed line.

There is an influxer for which representative concentrations for all substrates are found in Table 2. K$_o^+$ influx *via*

NKCC1 is not affected by the concentration of a metabolite substrate.

Inclusion of the effects of E$_{m,oi}$ and ΔG$_{Na}$ variation would increase the dimensionalities of Figure 4-type plots.

Making ΔG$_{Na}$(infl) less negative, for example, would (non-linearly) shift the lines to the left in Figs. 5-8. Making E$_{m,oi}$ less

negative (by increasing [K$_o^+$] or by [K$_i^+$] reduction) should do the same. In this paper, however, we focus on the novel

(and large) effects of P$_i$ variation, consequent to water co-transport.

Efflux Reactions. The ΔG$_{transporter}$(effl) values for the normal *efflux* reactions (II, I, or both, in Figs. 1,2) are found

in **Table 3b** (again, E$_{m,oi}$ is fixed at − 91 mV). (Note: the calculations are for ΔG(effl), not ΔG(infl).) In contrast

to the Table 3a influx reactions, except for that of KCC4 the other efflux reactions are unfavorable at P$_i$/P$_o$ = 1.00 (blue).

But, they are made much more favorable as P$_i$/P$_o$ increases (red), because of water co-efflux.

There is an efflux reaction for which representative concentrations for all substrates are found in Table 2;

KCC4, the [K$^+$,Cl$^-$,500H$_2$O] effluxer. For this, we see the efflux favorability is greatly enhanced by water co-efflux, even

though it is slightly favorable even w/o H$_2$O co-transport. Thus, it will always operate in efflux mode. K$_i^+$ efflux *via* KCC4

is not limited by the concentration of a metabolite substrate.

We focus on the lactic acid exporter, MCT1. The reaction is unfavorable when P$_i$/P$_o$ = 1.00, ΔG$_{MCT1}$(effl) =

8.5 kJ/mole (blue), but very favorable, **-** 58 kJ/mole (red), when P$_i$/P$_o$ = 1.05. Again, this is due to H$_2$O co-efflux.

At P$_i$/P$_o$ = 1.05 atm, the [lac$_o^-$]/[lac$_i^-$] ratio could reach only two without water, but can attain 3 x 10^{10} with the 500 H$_2$O

molecules co-effluxed.

DISCUSSION:

Biochemical Roles for Water Co-Transport. The analyses above suggest co-transported water can exert significant

thermodynamic effects on its process *via* the cytoplasmic pressure (P_i). We elaborate some examples.

<u>Figure 1 Glucose Influxers</u>.

<u>Small [glucose$_o$]</u>. Commonly, the SGLT transporter family is thought of in terms of its potential for catalyzing

glucose influx against its concentration gradient, chemogenic ΔG_{glc}(infl) (72,73). However, Figs. 5 and 6 show this is true

only when the P_i value is relatively small. For rSGLT1, "uphill" glucose influx is impossible for any P_i above ~1.03 P_o.

A 1 % P_i increase to 1.04 atm requires a [glc$_o$] value of at least ~10 μM for influx. This is five times *larger than* the 2 μM

[glc$_i$] fixed for Figs. 5-6. Perhaps cells using exclusively SGLT transporters always have small P_i values. Maybe P_i

fluctuation serves to control glucose influx. The GLUT family of transporters, with its smaller H_2O stoichiometries

(Fig. 1), does not suffer this severe (P_i/P_o)-dependence (Fig. 6). For GLUT1, the H_2O/glucose transport ratio is much

smaller than for rSGLT1, and its [glc$_o$]$_{min}$ value is almost P_i-insensitive (Fig. 6). Perhaps cells with larger P_i values,

and in situations with smaller [glc$_o$] values, employ GLUT enzymes. On the other hand, if P_i values are small, cells with

SGLT transporters will take up glucose at smaller [glc$_i$] values than cells with GLUT transporters. When rSGLT1 P_i is small,

small [glc$_o$] can insure sufficient [glc$_i$]. Also, when P_i is small, besides carrying glucose uphill, SGLT transporters are

"pumping" water up a barochemical hill. They are maintaining the H_2O barochemical steady-state.

<u>Large [glucose$_o$]</u>. If there is a situation where extracellular glucose is maintained at a relatively large value

(*e.g.*, 5 mM), the role of P_i can be viewed differently. Setting $P_i = P_o$, or ignoring co-transported water, would allow

absurdly large [glc$_i$]$_{max}$ values for cells with SGLT enzymes (Fig. 7). This would surely amount to a sugar overload. Even

if that was not cytotoxic, it could cause cells in a tissue to partake differentially of any available glucose charge –

some cells initially reached by a bolus taking up more (or even all) sugar than others. This would mean a heterogeneous

cellular [glc$_i$] spatial distribution. However with H_2O co-transport, representative experimental P_i values (say, 4% larger

than P_o) suppress the [glc$_i$]$_{max}$ value to near the hexokinase K_m, 1.7 μM for glucose (Fig. 7). Cells with even larger P_i

values will take up less glucose than those with smaller P_i. For rSGLT1 when P_i/P_o = 1.05 and [glc$_o$] = 5 mM, [glc$_i$]$_{max}$ is

near 1 nM, insufficient for metabolism. With no H_2O co-transport, [glc$_i$]$_{max}$ would increase more than ten orders

of magnitude; many orders greater than the hexokinase K_m for glucose.

For a given extracellular glucose concentration, cells with exclusively GLUT transporters will take up more

glucose than cells with exclusively SGLT enzymes at larger P_i values. Perhaps GLUT transporters are found in cells with

larger P_i values also in environments where [glc$_o$] is large.

As shown in Fig. 1, both SGLT and GLUT transporters deliver glucose. It is interesting a switch from

GLUT1- to SGLT-mediated cellular glucose uptake during lung cancer progression has been reported (20).

More generally, GLUT1 expression (along with other genetic changes) is found to promote the cancer cell Warburg State

(74). However, among these families, only the GLUTs transport the [18]FDG-PET tracer, (2-fluorodeoxyglucose) 2-FDG (20).

This produces an interpretation problem for the metabolic rate of glucose uptake and consumption determined

by quantitative [18]FDG-PET.

Because of H_2O co-transport, cells seem never allowed much steady-state free glucose. The role of water

co-influx, common to all the glucose influxers [hSGLT1, rSGLT1, GLUT1, GLUT2], appears dominant in controlling [glc_i].

It seems water co-transport *via* an SGLT generally guarantees a [glc_i] near the hexokinase K_m for glucose. Glucose

is effectively metabolized immediately upon entering the cell. This protects cells from too much glucose, and tissues

from excessive cellular glucose uptake inequality. A large ([glc_o] - [glc_i]) difference makes [glc_i] hard to measure, even

with modern methods (47). Also, glc_i likely does not contribute to the glucoCEST NMR signal (75,76). On the other

hand, GLUT1 generally keeps [glc_i] near [glc_o] (Fig. 6). (GLUT2 will keep it somewhat smaller.)

<u>Figure 1 Neurotransmitter Influxers: astrocyte uptake / synaptic clearance</u>.

Similar considerations must also obtain for the Table 3a influx reactions of the principal excitatory and inhibitory

neurotransmitters glutamate$_o^-$ and GABA$_o$, respectively.   Once inside the astrocyte, these must be processed

immediately by their metabolizing enzymes. EAAT1 (Fig. 8) and GAT1 will not allow them to build-up. (Note: for EAAT1,

Z_{glt} = - 1, and there are $H_3O_o^+$ co-influx and K_i^+ co-*efflux* terms (21).)

However, an even more crucial aspect of the EAAT and GAT roles is the effective clearance of neurotransmitter

species from synaptic junctions in time to enable the next action potential. Figure 8 shows this for EAAT1 and

glutamate$_o^-$. It is clear the astrocyte P_i must be relatively small (< 1.02 atm) to ensure [L-glutamate$_o^-$] does not much

exceed 2.3 µM, its EC$_{50}$ value for the iGlutR receptor enzyme. At the same time, the astrocyte P_i must be large enough

(1.035 atm) to ensure astrocytosolic [L-glutamate$_i^-$] does not much exceed 1.4 mM, its K_d value for glutamine synthetase

(analogous to Fig. 7). The large H_2O stoichiometries for EAAT1 (and GAT1) transport (similar to that of rSGLT1, Fig. 1)

guarantee high sensitivity to astrocyte P_i.

The astrocytic uptake of synaptic glutamate$^-$ is particularly interesting. If astrocyte P_i/P_o is even ~1.03,

[glutamate$_o^-$] would have to reach ~100 µM (almost two orders of magnitude *over* 2.3 µM) to initiate uptake (Fig. 8).

This could be sufficient to saturate iGlutR, and quench further synaptic transmission. Perhaps this is a physicochemical

mechanism whereby astrocytes serve as "gatekeepers" for synaptic activity. Increases of P_i to only 3 or 4% above P_o

could allow receptor saturation, and the interruption of neuronal firing. Very small astrocyte P_i fluctuations could

enable or disable synaptic function. For excitatory glutamatergic synapses this means turning off or on neural *excitation*.

A glance at the Fig. 1 GAT1 influx stoichiometry indicates very similar considerations would obtain for GABAergic

synapses. In that case, neuronal *inhibition* could be halted by increased intra-astrocyte pressures – resulting in

increased *excitation*.

Astrocyte swelling, a sign of possible P_i change, has been implicated in the glymphatic processes occurring

during sleep (commentary (77)).   Also, it has been suggested some therapeutic interventions can disturb

the glutamate⁻/glutamine cycle leaving interstitial glutamine, which can act as a nutrient for brain cancer cells (78).

In non-neural cells, [glutamate$_i$⁻] values can become very large, 64 mM (47).   Perhaps this is because there is no

glutamine synthetase.

<u>The Figure 1 NKCC1 Influxer</u>.

The function of NKCC1 (reviewed in (22)) is to effect cellular K_o^+ and H_2O_o uptake against their respective

chemical and barochemical gradients. Because of the huge H_2O stoichiometry (Fig. 1), this process is predicted to be

strongly P_i-dependent.

<u>Figure 1 Effluxers.</u>

For the Table 3b effluxers, each of the reactions are made significantly more favorable by increased intracellular

pressure (only KCC4 is favorable when $P_i = P_o$, and NaDC1 requires $P_i/P_o > 1.05$). Again, this is due to the pressure effect

on the free energy caused by the large numbers of co-transported H_2O molecules.

We use the monocarboxylate transporter [MCT1] as an example. It's main role is ridding the cell of lactic acid

build-up from glycolytic-type metabolism. This is common in cancer cells (the Warburg Effect). Lactic acid can be

sufficiently cytotoxic that MCT1 inhibition has been considered as a cancer therapy (79). **Equation (7)** expresses

710
$$\Delta G_{MCT1}(effl) = \left[2.58ln\left\{\frac{[lac_o^-]}{[lac_i^-]}\right\} - 0.0965(-0.91) + 2.58ln\left\{\frac{P_o}{P_i}\right\}\right] + \left[2.58ln\left\{\frac{[H_3O_o^+]}{[H_3O_i^+]}\right\} + 0.0965(-0.91) + 2.58ln\left\{\frac{P_o}{P_i}\right\}\right] + 500\left[2.58ln\left\{\frac{OsX_{H2O,o}}{OsX_{H2O,i}}\right\} + 2.58ln\left\{\frac{P_o}{P_i}\right\}\right] \quad (7)$$

the free energy change for the lactic acid efflux reaction: the first, second, and third terms represent the lactate⁻, H_3O^+,

and H_2O contributions, respectively. Taking the Table 1 H_3O^+ and OsX_{H2O} concentrations, $E_{m,oi}$ = -0.91 mV, $\{P_o/P_i\}$ =

$\{1/1.05\}$ = 0.95, and letting $\Delta G_{MCT1}(effl)$ = 0, we solve for the maximum $\{[lac_o^-]/[lac_i^-]\}$ value that can be achieved.

We find $\{[lac_o^-]/[lac_i^-]\}$ = 2.6 x 10^{10} when 500 H_2O molecules are co-transported, compared with only 2 if there was

no H_2O co-transport. This ten order-of-magnitude increase insures essentially all lactic acid is expelled from the cell –

because of the barochemical water contribution. Once again, water co-transport is protecting the cells – even if they are

malignant. A consequence is the extracellular acidification common in tumors (79). This phenomenon is also found

for other cell types (*e.g.*, muscle tissue cramping).

We now scrutinize the nature of intracellular hydraulic pressure (P_i).

The Nature of Intracellular Pressure. It seems clear the crucial thermodynamic role played by co-transported water is

effectuated by the pressure difference across the cell membrane. The intracellular pressure (P_i) is commonly considered

"osmotic" in nature. That is, it is thought to result from the phenomenon of *water molecules* crossing the membrane.

The simple first-order picture (H_2O permeable; osmolytes (Na^+, K^+, Cl^-) impermeable in the $P_f(p)$ sense) invokes *selective*

water permeability, with influx and efflux becoming the same when ΔP ($\equiv P_i$ - P_o) equals RT times the trans-membrane

osmotic gradient (7): at T = 310 K, RT is 25.4 L•atm/mole. Strictly, osmotic effects are entropic in nature (the entropy

of mixing (27)).

The first-order expression for the trans-cytolemmal osmotic gradient ($\Delta\pi_{oi}$) in terms of osmolyte concentrations

is given in **Equation (8)**, where [osmolytes$_i$] and [osmolytes$_o$] are the intra- and extracellular concentrations,

729
$$\Delta\pi_{oi} = \pi_i - \pi_o = RT([osmolytes_i] - [osmolytes_o]) = 25.4([osmolytes_i] - [osmolytes_o]) \tag{8}$$

respectively. This equation assumes *ideal solutions* ((44), p 37; (80), p. 302 ff): *i.e.*, entropy only. In Eq. (8), the osmolyte

concentrations are given on the osmolarity scale. We use the representative values listed in Table 1. There, total

[osmolytes$_i$] = 0.379 OsM and total [osmolytes$_o$] = 0.289 OsM. (At such magnitudes, these are essentially identical

to their *molal* concentrations. We neglect the tiny mOsM intracellular RNA, lipid, and DNA concentrations.) Inserting

these into Eq. (8) yields $\Delta\pi_{oi}$ = 2.29 atm. With the Table 1 compartmental water contents, we can also estimate

volume molal ($_vm$ = OsM/f$_W$) osmolyte concentrations. We obtain: $_vm_i$ = 0.38/0.76 = 0.50 mole(osmolytes)/L(cell water),

and $_vm_o$ = 0.29/0.82 = 0.35 mole(osmolytes)/L(extracellular water).   Inserting these into Eq. (8) gives:

$\Delta\pi_{oi}$ = 25.4 ($_vm_i$ - $_vm_o$) = 25.4(0.50 – 0.35) = 3.8 atm, an even greater value.

If there were only first-order mixing entropy contributions, P_i would equal 2.3 or 3.8 atm at osmotic

"equilibrium" (really steady-state; assuming P_o = 1 atm). A molecular mechanism for selective water influx and efflux

could be free (*i.e.*, unregulated) bidirectional aquaporin-mediated transport, as has been suggested (81). However,

the largest experimental P_i value (1.5 atm (66)) is 1.5 times smaller than even 2.3 atm. The most pertinent P_i values

(1.02 to 1.05; Figs. 4-8) are more than 2.2 times smaller. A (P_i - P_o) of 1.03 would correspond to Δ[osmolytes] = 0.04 OsM

(1.03/25.4). This is 2.5 times smaller than the 0.1 OsM value we estimate in Table 1. Of course, the Table 1

Δ[osmolytes] value is not found for any real cell. However, it is a reasonably representative magnitude: it is very unlikely

to be less than 1.5 times any real value. If anything, 0.1 OsM may even be small (52). It is not clear any real cytoplasmic

lipid bilayer membrane could even survive ΔP of 2.3 atm (the hydraulic pressure *ca.* 38 ft below sea level) if the *internal*

pressure (P_i) is the greater, let alone 3.8 atm. Even if it could, a ΔP value of even 2.29 atm would yield a barogenic

Eq. (8) term: ΔG(infl) = 2.58ln(2.29) = 2.14 kJ/mole. This is so large, it would dominate the values in Tables 2 - 4, and

obliterate the realistic estimations of the last section.

Such first-order considerations carry the implicit assumption of solution *ideality*; no specific osmolyte/osmolyte

molecular interactions. Osmotic pressure is generally considered one of the "colligative" properties: the identity

of the osmolyte is irrelevant. This is why the mole fraction concentration scale, mOsX (or the related mOsm scale),

for solution components is the most appropriate for osmotic considerations ((44), p. 56; (45), p.27; (46)). But, since

the Table 1 osmolyte mOSM values are sufficiently small (in the absolute sense) that mOsm values can be used,

this suggests their activity coefficients ($\gamma_{osmolyte}$, or $y_{osmolyte}$) are near unity. However, such reasoning does

"fail to emphasize the departure from ideality indicated by the activity of the solute" ((45), p.29).

Furthermore, we see from Table 1 intra- and extracellular water mole fractions $OsX_{H2O,i}$ = 0.993 and $OsX_{H2O,o}$ =

0.995, respectively. $OsX_{H2O,i}$ is only 0.2% smaller than $OsX_{H2O,o}$, and each is nearly the value for pure water. Given this

situation, it is not unreasonable to also assume the water activity coefficients are equal, $f_{H2O,i}$ = $f_{H2O,o}$ (63). In **A.2**,

we present calculations of H_2O activities for surrogate interstitial and cytosolic solutions that support this contention.

Table 2 shows inserting the calculated values $a_{H2O,i}$ = 0.994 and $a_{H2O,o}$ = 0.995 into Eq. (3) or (4), leads to a water

chemogenic contribution to ΔG_{H2O}(infl) of only ~3 J/mole favoring cellular water influx. Also, we showed above

that chemogenic ΔG_{H2O}(infl) =- 0.003 kJ/mole is counteracted by a P_i/P_o ratio of only 1.001. This is only one tenth

of a conservatively small typical intracellular pressure, P_i/P_o = 1.01 (27,66). This adds to the suggestion the entropic

contribution to P_i is actually small, usually less than 10%.

All these considerations indicate the mixing entropic contribution (- $T\Delta S_{H2O}$(infl)) to water ΔG_{H2O}(infl) is small.

By definition, a thermodynamically non-ideal solution means there is an enthalpic (ΔH_{H2O}(infl)) contribution.

A representative P_i/P_o value of 1.05 corresponds to barochemical ΔG_{H2O}(infl) = - 0.13 kJ/mole. A P_i/P_o ratio of 1.001

implies ~90% ΔH_{H2O}(infl); *i.e.*, - 0.12 kJ/mole. The conclusion must be that enthalpic (ΔH_{H2O}(infl)) contributions dominate

P_i. The aqueous solutions inside (and, for that matter, outside) cells deviate greatly from ideality. There are highly

specific molecular interactions between the various solutes, and with water. It has long been known the "hydration"

of biological solutes is extremely important for many different cellular processes (57-59). If the solute-interacting water

entropy is smaller than in pure water, this could even give rise to a (not mixing) entropic driving force for *efflux*.

Whatever the actual case, the intracellular osmolytes do not seem to lower the water "escaping tendency" nearly as

much as the first-order osmotic pressure equation (8) would predict.

There is a long history of considering higher order contributions to Eq. (8) ((82), p. 210 ff; (83)). However, it is

probably not realistic to expect an environment as complex as the intracellular *milieu* to be well-modeled as

a homogeneous solution.

Recent reports suggest particular intracellular species contribute to regulating intracellular hydraulic pressure.

These include: the mechanosensitive transcriptional regulator YAP (Yes-associated protein) (84), tropomyosins 1.6 and

2.1 (Tpm 1.6 and Tpm 2.1) (85), and the capsaicin-activatable transient receptor potential vanilloid 1 (TRPV1) cation

channel, that interestingly, stimulates the Fig. 1 NKCC1 water co-influxer (86) as well as NKA. An especially intriguing

proposal is that intracellular macromolecular polyanion electrostatic interactions dominate P_i, while allowing

modulation by extracellular osmolality (52).

It may be often assumed tissue homeostasis means $\Delta\pi$ is zero, and is regulated to that point: *i.e.*, [osmolytes$_i$] =

[osmolytes$_o$] in Eq. (8): *i.e.*, "*isosmolality*." However, the word "*isotonal"* (6) suggests only equal pressures, and is

preferred. A 290 mOsm NaCl solution is often considered "*isotonic*." Bathing solutions hyper- or hypo-tonic relative

to this cause *in vitro* cells (19) and *ex vivo* tissue cells (87) to shrink or swell due to *net* water efflux or influx,

respectively. Bolus blood infusions of "*hypertonic*" solutions are used clinically to *transiently* open the blood-brain-

barrier by shrinking capillary endothelial cells – and thus deliver otherwise non-extravasating therapeutic drugs

to the cerebral parenchyma (88). In cell cultures and to some degree in perfused tissues, the investigator can specify

and control [osmolytes$_o$]. When such a system reaches homeostasis, a common presumption may be that [osmolytes$_i$] =

[osmolytes$_o$]. From this, it is usually assumed [osmolytes$_i$] $\approx$ 0.3 OsM, the Table 1 [osmolytes$_o$] value (52). However, our

estimated [osmolytes$_i$] $\approx$ 0.4 OsM (Table 1) indicates this may be rarely (never) true.

Likely further evidence of cytoplasmic non-ideality is that different tissues seem to require different bathing

solution osmolalities to achieve isotonicity. While ~320 mOsm suffices for the retina and olfactory bulb, almost

600 mOsm is required for cerebral cortex (87). Inhibition of NKA causes cortical cells to swell considerably; *i.e.*, a net

category B water influx (89).

Whatever the actual P_i physicochemical nature, the calculations in the last section remain valid because they

employ experimental P_i values.

Aquaporin Role in Cellular Homeostasis. The specific water-transporting membrane aquaporin (AQP) molecules are

found in almost all tissues (6,90,91). Like all the Fig. 1 transporters, they are capable of catalyzing category B

unidirectional (flux, P_f-type) or category A bidirectional (exchange, P_d-type) water transport. The famously large "single

channel" AQP4 water volume flux, 0.25 fL/s/AQP4 (8.3 H_2O molecules/ns/AQP4), was determined at $10^{o}C$ and with

a very large 4.4 atm osmotic gradient (92). Like all P_f measurements, this is derived from the *asymptotic* volume change

when transport is initiated away from osmotic "equilibrium." However, AQP efficiency is minimal when there is

no pressure gradient ($\Delta P = 0$) (37), *i.e.,* the P_d-type condition. This is consistent with molecular considerations (the water

single-file nature of the AQP channel structure (30) is not optimal for exchange). Thus, AQP's by themselves have been

thought of as intrinsically *passive* transporters. They can facilitate very large *net* water influxes or effluxes, but

only when the latter are driven by independent forces. Thus, though their involvement in non-homeostatic cell swelling

or shrinking has been widely investigated (6,90,91), their roles in homeostasis has been less clear. We are not aware

a "single channel" category A exchange magnitude has ever been experimentally determined.

The results presented here suggest there is thermodynamic (metabolic) energy stored in tissue water

compartmentalization itself, helping maintain the system in homeostasis. Thus, we have inserted AQPs explicitly into

the active trans-membrane water cycling (AWC) scheme, rate-limited by ^{c}MR$_{NKA}$ (Fig. 2). Since AWC is, by definition,

homeostatic, MR$_{H2O}$(infl) = MR$_{H2O}$(effl). However, it is unlikely AQP's contribute equally to water influx and efflux.

As stated above, if ΔP was actually osmotic in nature, unregulated dominant AQP activity would lead to ΔP values many

times those measured. Thus, it is likely MR$_{AQP}$(infl) ≠ MR$_{AQP}$(effl).

In cell suspensions (and *in vivo*) astroglial aquaporin AQP4 expression has been shown to contribute

to category A water exchange (81). Furthermore, studies with human C6 glioma (cancer) cell suspensions and TGN020

(a specific AQP4 inhibitor) are informative. The k_{io} rate constant ratios ($^{TGN020}k_{io}$)/($^{C6}k_{io}$) is 0.69, while ($^{ouabain}k_{io}$)/($^{C6}k_{io}$) is

0.65 (81). Inhibiting water exchange with extracellular TGN020 reduces k_{io} to the same extent as inhibiting it with

extracellular ouabain, the specific NKA inhibitor. This indicates AQP4 contributes to AWC. Cancer cells in suspension

may not be in the Warburg state (2,4).

The metabolite influxers (Fig. 1) can provide significant MR$_{H2O}$(infl) values. The cellular glucose consumption

rate, MR$_{glc}$(consump), has been determined to be 38 μM(glc)/s/cell for murine kidney epithelial cells (47) and for brain

tissue cells, assuming reasonable cell density and volume values (68). The glucose influx rate, MR$_{glc}$(infl) must be at least

MR$_{glc}$(consump). But GLUT1 and rSGLT1 have different stoichiometries (Fig. 1). Thus, MR$_{H2O}$(infl) would range from

1.8×10^9 to 17×10^9 H$_2$O/s/cell for 2pL cells with, respectively, exclusively GLUT1 or exclusively rSGLT1 transporters. Using

the ratio of the glutamate$^-$-to-glutamine conversion flux, MR$_{glu-gln}$(cycle) to MR$_{glc}$(consump) (93), we estimate MR$_{H2O}$(infl)

provided by the EAAT1 transporter to be 4×10^7 H$_2$O/s/cell for a 2 pL astrocyte. This would be in addition to MR$_{H2O}$(infl)

provided by a glucose influxer.

Some so-called "metabolic water" is generated within the cell after glucose is imported. However, there are

at most six H$_2$O molecules produced per glucose molecule metabolized, depending on the glycolysis and oxidative

phosphorylation proportions (94). This is small when compared with even the smallest Fig. 1 glucose influxer H$_2$O

stoichiometries.

Our most recent estimate of brain x, the MR$_{H2O}$(AWC)/MR$_{NKA}$ ratio, is $> 10^6$ (2). We opined "such large

stoichiometries suggest aquaporin participation in active trans-membrane water cycling." Given the large cellular water

influx rates, and the $P_i > P_o$ pressure gradient, it seems likely MR$_{AQP}$(effl) > MR$_{AQP}$(infl). That is, in homeostasis AQP is

likely mainly catalyzing water *efflux*. It is working in parallel with the KCC effluxer. This is why we indicate the inhibitor

blocking MR$_{AQP}$(effl) in Fig. 2.

Water should be considered a substrate for active trans-membrane water cycling. It has thermodynamic

consequences. If it turns out NKA itself has its own water stoichiometry, AQPs could be then considered as also

secondary active enzymes. If not, they are only tertiary active transporters – sharing water as a substrate
with only the secondary active water co-transporters.

An interesting analogy with potassium ion-specific channels (PCs) (reviewed in (29); (95)) might obtain. It is
thought the formally "passive" K_i^+ efflux through PCs dominates the production of $E_{m,oi}$. The idea is K_i^+ efflux driven
by the chemogenic term proceeds until the latter is balanced by the electrogenic term (Table 2), which actually requires
very few effluxed K^+ ions. After that, an "electrochemical" steady-state condition is reached, and *homeostatic* K_i^+ efflux
is synchronized with $^cMR_{NKA}$. If it is water transport that dominates the P_i value, perhaps there is an analogous H_2O_i
efflux driven by the barochemical gradient. This would tend to decrease P_i. However, diminished P_i would allow
the Na^+ electrochemical gradient a greater role. This would tend to increase the water co-influxes of the many
transporters that also bring in Na^+ (*e.g.*, NKCC- and SGLT-catalyzed influxes; Figs. 1-8), which would tend to increase P_i.
This "feedback loop" would then establish an NKA-maintained *steady-state*, and AQP-catalyzed water efflux is also
synchronized with $^cMR_{NKA}$. Thus, we have AQP playing a vital role in active trans-membrane water cycling (Fig. 2).

A trans-membrane pressure difference could also affect transporter *kinetics*. This is discussed in **A.3**.

CONCLUSION:

The Na$^+$,K$^+$-ATPase (NKA) enzyme function has long been known to maintain Na$^+$ and K$^+$ in trans-membrane

electrochemical *steady-states*, which are far from chemical *equilibria* ([Na$_i^+$] = [Na$_o^+$], [K$_i^+$] = [K$_o^+$]).   Here, we find

metabolic energy released by NKA-catalyzed ATP hydrolysis is also required to maintain (and is thus stored in)

in a trans-cytolemmal water barochemical steady-state.   This is an inherent part of NKA function.   An important

difference from the Na$^+$ and K$^+$ cases is the trans-membrane water distribution is in (or near) chemical equilibrium

([H$_2$O$_i$] $\approx$ [H$_2$O$_o$]).   Crucially, the barochemical H$_2$O steady-state is significant, and enzymatic water co-transport has very

important thermodynamic metabolic consequences.   Active trans-membrane water cycling does not represent

a futile cycle.

ACKNOWLEDGEMENTS:

The authors especially thank Professor Daniel Zuckerman for critical readings of manuscript drafts and very
insightful suggestions, and Dr. August George for accomplishing initial calculations. They also thank Professors
Joseph Ackerman, Valerie Anderson, James Balschi, James Bassingthwaighte, Cynthia Burrows, Deborah Burstein,
Kieran Clarke, Ira Cohen, Michael Grabe, Fahmeed Hyder, Joanne Ingwall, Christopher Kroenke, Kenneth Krohn,
Philip Kuchel, Xin Li, Daniel Liefwalker (for ref. (74)), Jeffrey Maki, Silvia Mangia, Richard Mathias, Anusha Mishra,
William Rooney, Douglas Rothman, Suzanne Scarlatta, William Skach, and Mark Woods, and Drs. Joseph Armstrong,
Mauro DiNuzzo, Bassam Haddad, Audrey O'Connor, and Gregory Wilson, and Messers. Eric Baetscher, Eric Baker,
Brendan Moloney, and Joshua Schlegel for stimulating discussions.

Support:

- OHSU Brenden-Colson Center for Pancreatic Care; Sheppard (PI), Springer (PI)
- NIH UL1TR02369; Ellison, Morris (PIs); awarded by the Oregon Clinical and Translation Institute, Biomedical Innovation Program; Pike (PI)

AUTHOR ROLES:

CSS conceived the approach, developed the theory, carried out the calculations, and drafted the manuscript.

MMP edited the manuscript drafts, with particular attention to the metabolic aspects.

TMB contributed calculations and edited the manuscript drafts, especially scrutinizing the theory.

CONFLICTS OF INTEREST:

CSS and **TMB** are co-inventors on U.S. patent 11,728,038, "Activity MRI" (issued 15 August, 2023), which describes the MADI approach.

REFERENCES:

1. C. S. Springer, E. M. Baker, X. Li, B. Moloney, G. J. Wilson, M. M. Pike, T. M. Barbara, W. D. Rooney, J. H. Maki,
Metabolic activity diffusion imaging [MADI]: I. Metabolic, cytometric modeling and simulations. *NMR Biomed.* **36,** e4781
(2023).

2. C. S. Springer, E. M. Baker, X. Li, B. Moloney, G. J. Wilson, V. C. Anderson, M. K. Sammi, M. M. Pike, M. G. Garzotto,
R. P. Kopp, F. V. Coakley, W. D. Rooney, J. H. Maki, Metabolic activity diffusion imaging [MADI]: II. Non-invasive,
high-resolution human brain imaging of sodium pump flux and cell metrics. *NMR Biomed.* **36,** e4782 (2023).

3. J. J. Neil, J. J. H. Ackerman, Metabolic activity diffusion imaging (MADI): A new paradigm. *NMR in Biomed.* **36,** e4841
(2023).

4. J. Schlegel, E. Baker, S. Holland, J. Stoller, W. Packwood, X. Li, R. Barajas, C. Springer, M. Pike, Metabolic activity
diffusion imaging [MADI] of rat brain glioma. *Proc. Int. Soc. Magn. Reson. Med.* **31,** 3924 (2023).

5. M. M. Pike, X. Li, E. Baetscher, T. M. Barbara, M. K. Sammi, A. A. Stevens, C. S. Springer, Does MADI detect temporal
brain metabolic activity changes?" *Proc. Int. Soc. Magn. Reson. Med.* **31,** 5177 (2023).

6. R. E. Day, P. Kitchen, D. S Owen, C. Bland, L. Marshall, A. C. Conner, R. M. Bill, M. T. Conner, Human aquaporins:
Regulators of transcellular water flow. *Biochim. Biophys. Acta* **1840,** 1492-1506 (2014).
[doi.org/10.1016/j.bbagen,2013,09,033]

7. A. S. Verkman, Water permeability measurements in living cells and complex tissues. *J. Membrane Biol.* **173,**
73-87 (2000).

8. E. Zeuthen, A sensitive 'Cartesian Diver' balance. *Nature* **159,** 440-441 (1947).

9. O. Kedem, A. Katchalsky, Thermodynamic analysis of the permeability of biological membranes to non-electrolytes.
Biochim. Biophys. Acta **27,** 229-246 (1958).

10. C. R. House, Water transport in cells and tissues. (Edward Arnold, London, 1974), p. 156.

11. S-T. Chen, C. S. Springer, Ionophore-catalyzed cation transport between phospholipid inverted micelles manifest
in DNMR. *Biophys. Chem.* **14,** 375-388 (1981).

12. T. Conlon, R. Outhred, Water diffusion permeability of erythrocytes using an NMR technique. *Biochim. Biophys. Acta*
288, 354-361 (1972).

13. C. S. Landis, X. Li, F. W. Telang, P. E. Molina, I. Pályka, G. Vétek, C. S. Springer, Equilibrium transcytolemmal water-
exchange kinetics in skeletal muscle *in vivo*. *Magn. Reson. Med.* **42,** 467-478 (1999).

14. X. Li, W. D. Rooney, C. S. Springer, A unified magnetic resonance imaging pharmacokinetic theory: Intravascular and
extracellular contrast reagents. *Magn. Reson. Med.* **54,** 1351-1359 (2005). [DOI 10.1002/mrm.20684]

15. C. S. Springer, Using ^1H_2O to measure and map sodium pump activity *in vivo*. *J. Magn. Reson.* **291,** 110-126 (2018).
[doi.org/10.1016/j/jmr.2018.02.018]

16. B. P. Hills, P. S. Belton, NMR studies of membrane transport. *Ann. Reports NMR Spectroscopy* **21,** 99-159 (1989).

17. B. M. Denker, B. L. Smith, F. P. Kuhajda, P. Agre, Identification, purification, and partial characterization of a novel
M_r 28,000 integral membrane protein from erythrocytes and renal tubules. *J. Biol. Chem.* **263,** 15634-15642 (1988).

18. N. MacAulay, Molecular mechanisms of brain water transport. *Nat. Rev. Neurosci.* **22,** 326-344 (2021).

19. T. Zeuthen, Water-transporting proteins. *J. Membrane Biol.* **234,** 57-73 (2010). [DOI 10.1007/s00232-009-9216-y]

20. J. R. Barrio, S. C. Huang, N. Satyamurthy, C. S. Scafoglio, A. S. Yu, A. Alavi, K. A. Krohn, Does 2-FDG-PET accurately
reflect quantitative *in vivo* glucose utilization. *J. Nucl. Med.* **61,** 931-937 (2020).

21. N. MacAulay, U. Gether, D. A. Klaerke, T. Zeuthen, Water transport by the human Na^+-coupled glutamate cotransporter
expressed in *Xenopus* oocytes. *J. Physiol.* **530,** 367-378 (2001).

22. K. Kaila, T. J. Price, J. A. Payne, M. Puskarjov, J. Voipio, Cation-chloride cotransporters in neuronal development, plasticity and disease. *Nature Rev. Neurosci.* **15,** 637-654 (2014). [doi:10.1038/nrn3819]

23. A. W. Autry, J. W. Gordon, H-Y. Chen, M. LaFontaine, R. Bok, M. Van Criekinge, J. B. Slater, L. Carvajal, J. E. Villanueva-Meyer, S. M. Chang, J. L. Clarke, J. M. Lupo, D. Xu, P. E. Z. Larson, D. B. Vigneron, Y. Li, Characterization of serial hyperpolarized ^{13}C metabolic imaging in patients with glioma. *NeuroImage: Clin.* **27,** 102323 (2020).

24. R. Ye, A. S. Verkman, Simultaneous optical measurement of osmotic and diffusional water permeability in cells and liposomes. *Biochem.* **28,** 824-829 (1989).

25. X. Li, S. Mangia, J-H. Lee, R. Bai, C. S. Springer, NMR shutter-speed elucidates apparent population inversion of ^{1}H$_2$O signals due to active transmembrane water cycling. *Magn. Reson. Med.* **82,** 411-424 (2019). [DOI: 10.1002/mrm.27725]

26. Y. Zhang, M. Poirier-Quinot, C. S. Springer, J. A. Balschi, Active trans-plasma membrane water cycling in yeast is revealed by NMR. *Biophys. J.* **101,** 2833-2842 (2011). [DOI:10.1016/j.bpj.2011.10.035]

27. Y. Li, K. Konstantopoulos, R. Zhao, Y. Mori, S. X. Sun, The importance of water and hydraulic pressure in cell dynamics. *J. Cell. Sci.* **133,** jcs240341 (2020).

28. C. S. Springer, Measurement of metal cation compartmentalization in tissue by high resolution metal cation NMR. *Ann. Rev. of Biophys. and Biophys. Chem.,* **16**, 375-399 (1987).

29. C. Öster, K. Kendriks, W. Kopec, V. Chevelkov, C. Shi, D. Michl, S. Lange, H. Sun, B. L. de Groot, A. Lange, The conduction pathway of potassium channels is water free under physiological conditions. *Sci. Adv.* **5,** eeaw6756 (2019).

30. B. L. de Groot, H. Grubmüller, Water permeation across biological membranes: Mechanism and dynamics of aquaporin-1 and GlpF. *Science* **294,** 2353-2357 (2001).

31. R. Bai, C. S. Springer, D. Plenz, P. J. Basser, Brain active trans-membrane water cycling measured by MR is associated with neuronal activity. *Magn. Reson. Med.* **81,** 1280-1295 (2019). [DOI:10.1002/mrm.27473]

32. J. L. Adelman, C. Ghezzi, P. Bisignano, D. D. F. Loo, S. Choe, J. Abramson, J. M. Rosenberg, E. M. Wright, M. Grabe, Stochastic steps in secondary active sugar transport. *Proc. Nat. Acad. Sci.* **113,** E3960-E3966 (2016).

33. J. L. Adelman, Y. Sheng, S. Choe, J. Abramson, E. M. Wright, J. M. Rosenberg, M. Grabe, Structural determinants of water permeation through the sodium-galactose transporter vSGLT. *Biophys. J.* **106,** 1280-1289 (2014).

34. J. Li, S. A. Shaikh, G. Enkavi, P-C. Wen, Z. Huang, E. Tajkhorshid, Transient formation of water-conducting states in membrane transporters. *Proc. Nat. Acad. Sci.* **110,** 7696-7701 (2013).

35. S. Choe, J. M. Rosenberg, J. Abramson, E. M. Wright, M. Grabe, Water permeation through the sodium-dependent galactose cotransporter vSGLT." *Biophys. J.* **99,** L56-L58 (2010).

36. A. Watanabe, S. Choe, V. Chaptal, J. M. Rosenberg, E. M. Wright, M. Grabe, J. Abramson, The mechanism of sodium and substrate release from the binding pocket of vSGLT. *Nature* **468,** 988-991 (2010).

37. F. Zhu, E. Tajkhorshid, K. Schulten, Theory and simulation of water permeation in aquaporin-1. *Biophys. J.* **86,** 50-57 (2004).

38. S. Zhang, J. Zhou, Y. Zhang, T. Liu, P. Friedel, W. Zhou, S. Somasekharan, K. Roy, L. Zhang, Y. Liu, X. Meng, H. Deng, W. Zeng, G. Li, B. Forbush, M. Yang, The structural basis of function and regulation of neuronal cotransporters NKCC1 and KCC2. *Comm. Biol.* **4,** 226 (2021).

39. H. Ogawa, T. Shinoda, F. Cornelius, C. Toyoshima, Crystal structure of the sodium-potassium pump (Na$^+$,K$^+$-ATPase) with bound potassium and sodium." *Proc. Nat. Acad. Sci.* **106,** 13742-13747 (2009).

40. M. M. Pike, J. C. Frazer, D. F. Dedrick, J. S. Ingwall, P. D. Allen, C. S. Springer, T. W. Smith, ^{23}Na and ^{39}K nuclear magnetic resonance studies of perfused rat hearts. *Biophys. J.* **48,** 159-173 (1985).

41. A. L. Lehninger, "*Principles of Biochemistry*." (Worth Pub., New York, 1982), p. 571.

42. D. M. Bers, W. H. Barry, S. Despa, Intracellular Na$^+$ regulation in cardiac myocytes. *Cardiovas. Res.* **57,** 897-912 (2003).

43. A. A. Kadir, B. J. Stubbs, C-R, Chong, H. Lee, M. Cole, C. Carr, D. Hauton, J McCullagh, R. D. Evans, K. Clarke, On the interdependence of ketone body oxidation, glycogen content, glycolysis, and energy metabolism in the heart. *J. Physiol.* **601.7,** 1207-1224 (2023). [doi:10,1113/JP284270]

44. A. G. Marshall, "*Biophysical Chemistry: Principles, Techniques, and Applications*" (Wiley, New York, 1978).

45. R. A. Robinson, R. H. Stokes, in "*Electrolyte Solutions*" (Dover, Mineola, ed. 2nd rev., 1970).

46. M. J. Blandamer, J. B. F. N. Engberts, P. T. Gleeson, J. C. R. Reis, Activity of water in aqueous systems: A frequently neglected property." *Chem. Soc. Rev.* **34,** 440-458 (2005).

47. J. O. Park, S. A. Rubin, Y-F. Xu, D. Amador-Noguez, J. Fan, T. Shlomi, J. D. Rabinowitz, Metabolite concentrations, fluxes, and free energies imply efficient enzyme usage. *Nat. Chem. Biol.* **12,** 482-489 (2016).

48. O. S. Andersen, "Cellular electrolyte metabolism," in *Enclycopedia of Metalloproteins*, R. H. Kretsinger, V. N. Uversky, A. Permiakov, Eds. (Springer, 2013), pp. 580-587.

49. R. Milo, What is the total number of protein molecules per cell volume? A call to rethink some published values. *Bioessays.* **35,** 1050-1055 (2013). [DOI 10.1002/bies.201300066]

50. F. C. Neidhardt, H. E. Umbarger, in *Escherichia coli and Salmonella: Cellular and Molecular Biology*, F. C. Neidhardt, Ed. (Am. Soc. Microbiol. Press, New York, ed. 2, 1996), vol. 1, chapt. 3.

51. K. C. Vinnakota, J. B. Bassingthwaighte, Myocardial density and composition: A basis for calculating intracellular metabolite concentrations. *Am. J. Physiol. Heart Circ. Physiol.* **286,** H1742-H1749 (2004). [10.1152/ajpheart.00478-2003]

52. H. Wennerström, M. Oliveberg, On the osmotic pressure of cells. *QRB Discovery.* **3,** e12 (2022).

53. R. Milo, R. Phillips, N. Orme, "*Cell Biology By the Numbers.*" (Garland Science, New York, 2016), pp. 70, 92, 93, 96, 106, 130, 198.

54. M. A. Model, E. Schonbrun, Optical determination of intracellular water in apoptotic cells. *J. Physiol.* **591.23,** 5843-5849 (2013). [DOI: 10.1113/jphysiol.2013.263228]

55. W. D. Rooney, G. Johnson, X. Li, E. R. Cohen, S-G. Kim, K. Ugurbil, C. S. Springer, Magnetic field and tissue dependencies of human brain longitudinal ^{1}H$_2$O relaxation *in vivo*. *Magn. Res. Med.* **57,** 308-318 (2007).

56. R. P. Rand, Probing the role of water in protein conformation and function. *Phil. Trans. R. Soc. Lond. B* **359,** 1277-1285 (2004). [DOI 10.1098/rstb.2004.1504]

57. P. Ball, Water is an active matrix of life for cell and molecular biology. *Proc. Nat. Acad. Sci.* **114:** 13327-13335 (2017). [doi/10.1073/pnas.1703781114]

58. M. Chaplin, Do we underestimate the importance of water in biology. *Nat. Rev. Mol. Cell Biol.* **7,** 862-866 (2006).

59. K. D. Garlid, The state of water in biological systems. *Int. Rev. Cytology* **192,** 281-302 (2000).

60. F. Persson, B. Halle, Transient access to the protein interior: Simulation versus NMR. *J. Am. Chem. Soc.* **135,** 8735-8748 (2013). [doi.org/10.1021/ja403405d]

61. X. L. Lu, V. C. Mow, Biomechanics of articular cartilage and determination of material properties. *Med. Sci. Sports Exerc.* **40,** 193-199 (2008).

62. E. Nimer, R. Schneiderman, A. Maroudas, Diffusion and partition of solutes in cartilage under static load. *Biophys. Chem.* **106,** 125-146 (2003). [doi:10.1016/S0301-4622(03)00157-1]

63. K. S. Pitzer, in "*Activity Coefficients in Electrolyte Solutions*" (CRC Press, Boca Raton, ed. 2, 1991).

64. J. Gao, X. Sun, L. C. Moore, T. W. White, P. R. Brink, R. T. Mathias, Lens intracellular pressure is generated by the circulation of sodium and modulated by gap junction coupling. *J. Gen. Physiol.* **137,** 507 520 (2011).

65. R. J. Petrie, H. Koo, Direct measurement of intracellular pressure. *Curr. Protoc. Cell Biol.* **63,** 12.9.1-12.9.9 (2015).

66. J. Gao, X. Sun, T. W. White, N. A, Delamere, R. T. Mathias, Feedback regulation of intracellular hydrostatic pressure in surface cells of the lens. *Biophys. J.* **109,** 1830-1839 (2015).

67. E. LoCastro, R. Paudyal, Y. Mazaheri, V. Hatzoglou, J. H. Oh, Y. Lu, A. S. Konar, K. vom Eigen, A. Ho, J. R. Ewing, N. Lee, J. O. Deasy, A. Shukla-Dave, Computational modeling of interstitial fluid pressure and velocity in head and neck cancer based on dynamic contrast-enhanced magnetic resonance imaging: Feasibility analysis. *Tomog.* **6,** 129-138 (2020).

68. M. DiNuzzo, G. A. Dienel, K. L. Behar, O. A. Petroff, H. Benveniste, F. Hyder, F. Giove, S. Michaeli, S. Mangia, S. Herculano-Houzel, D. L. Rothman, Neurovascular coupling is optimized to compensate for the increase in proton production from nonoxidative glycolysis and glycogenolysis during brain activation and maintain homeostasis of pH, pCO_2, and pO_2. J. Neurochem. **00,** 1-31 (2023). [DOI:10.1111/jnc.15839]

69. H. H. Chowdhury, Differences in cytosolic glucose dynamics in astrocytes and adipocytes measured by FRET-based nanosensors. *Biophys. Chem.* **261,** 106377 (2020).

70. I. Schomburg, A. Chang, S. Placzek, C. Sohnigen, M. Rother, M. Lang, C. Munaretto, S. Ulas, M. Stelzer, A. Grote, M. Scheer, D. Schomberg, BRENDA in 2013: Integrated reactions, kinetic data, enzyme function data, improved disease classification: New options and content in BRENDA. *Nucleic Acids Res.* **41,** D764-D772 (2013).

71. D. K. Patneau, M. L. Mayer, Structure-activity relationships for amino acid transmitter candidates acting at N-Methyl-D-aspartate and quisqualate Receptors. *J. Neurosci.* **10,** 2385-2399 (1990).

72. P. Bisignano, C. Ghezzi, H. Jo, N. F. Polizzi, T. Athoff, C. Kalyanaraman, R. Friemann, M. P. Jacobson, E. M. Wright, M. Grabe, Inhibitor binding mode and allosteric regulation of Na^+-glucose symporters. *Nature Comm.* **9,** 5245 (2018).

73. J. Jurcovicova, Glucose transport in brain – Effect of inflammation. *Endo. Reg.* **48,** 35-48 (2014). [doi:10.4149/endo_2014_01_35]

74. J-H. Lee, R. Liu, J. Li, Y. Wang, L. Tan, X-J. Li, X. Qian, C. Zhang, Y. Xia, D. Xu, W. Guo, Z. Ding, L. Du, Y. Zheng, Q. Chen, P. L. Lorenzi, G. B. Mills, T. Jiang, Z. Lu, EGFR-phosphorylated platelet isoform of phosphofructokinase 1 promotes PI3K activation. *Mol. Cell* **70,** 197-210 (2018).

75. X. Xu, A. A. Sehgal, N. N. Yadav, J. Laterra, L. Blair, J. Blakeley, A. Seidemo, J. M. Coughlin, M. G. Pomper, L. Knutsson, P. C. M. van Zijl, D-glucose weighted chemical exchange saturation transfer (glucoCEST)-based dynamic glucose enhanced (DGE) MRI at 3T: Early experience in healthy volunteers and brain tumor patients. *Magn. Reson. Med.* **84,** 247-262 (2020).

76. F. A. Nasrallah, G. Pagès, P. W. Kuchel, X. Golay, K-H. Chaung, Imaging brain deoxyglucose uptake and metabolism by GlucoCEST MRI. *J. Cerebr. Blood Flow & Metabol.* **33,** 1270-1278 (2013).

77. A. J. Smith, B-J. Jin, A. S. Verkman, Muddying the water in brain edema? *Trends Neurosci.* **38,** 331-332 (2015).

78. T. N. Seyfried, L. Shelton, G. Arismendi-Morillo, M. Kalamian, A. Elsakka, J. Maroon, P. Mukherjee, Provocative question: Should ketogenic metabolic therapy become the standard of care for glioblastoma? *Neurochem. Res.* **44,** 2392-2404 (2019).

79. K. Nath, D. S. Nelson, D. F. Heitjan, R. Zhou. D. B. Leeper, J. D. Glickson, Effects of hyperglycemia on lonidamine-induced acidification and de-energization of human melanoma xenografts and sensitization to melphalan. (2015). *NMR Biomed.* **28,** 395-403. [DOI:10.1002/nbm.3260]

80. H. B. Callen, "Thermodynamics and an introduction to thermostatistics." (Wiley, New York, ed. 2, 1985).

81. Y. Jia, S. Xu, G. Han, B. Wang, Z. Wang, C. Lan, P. Zhao, M. Gao, Y. Zhang, W. Jiang, B. Qiu, R. Liu, Y-C. Hsu, Y. Sun, C. Liu, Y. Liu, R. Bai, Transmembrane water-efflux rate measured by magnetic resonance imaging as a biomarker of the expression of aquaporin-4 in gliomas. *Nat. Biomed. Eng.* **7,** 236-252 (2023).[doi.org/10.1038/s41551-022-00960-9]

82. C. Tanford, "Physical Chemistry of Macromolecules." (Wiley, New York, 1961).

83. H. R. Naito, R. Okamoto, T. Sumi, K. Koga, Osmotic second virial coefficients for hydrophobic interactions as a function of solute size. *J. Chem. Phys.* **156,** 221104 (2022).

84. N. A. Perez-Gonzalez, N. D. Rochman, K. Yao, J. Tao, M-T. T. Le, S. Flanary, L. Sablich, B. Toler, E. Crentsil, F. Takaesu, B. Lambrus, J. Huang, V. Fu, P. Chengappa, T. M. Jones, A. J. Holland, S. An, D. Wirtz, R. J. Petrie, K-L. Guan, S. X. Sun, YAP and TAZ Regulate Cell Volume. *J. Cell Biol.* **218,** 3472-3488 (2019).

85. K. Sao, T. M. Jones, A. D. Doyle, D. Maity, G. Schevzov, Y. Chen, P. W. Gunning, R. J. Petrie, Myosin II governs intracellular pressure and traction by distinct tropomyosin-dependent mechanisms. *Mol. Biol. Cell* **30,** 1170-1181 (2019).

86. M. Shahidullah, A. Mandal, R. T. Mathias, J. Gao, D. Križaj, S. Redmon, N. A. Delamere, TRPV1 activation stimulates NKCC1 and increases hydrostatic pressure in the mouse lens. *Am. J. Physiol. Cell Physiol.* **318,** C969-C980 (2020).

87. M. Pallotto, P. V. Watkins, B. Fubara, J. H. Singer, K. L. Briggman, Extracellular space preservation aids the connectomic analysis of neural circuits. *eLife* **4,** e08206 (2015).

88. E. A. Neuwelt, Mechanisms of disease: The blood-brain barrier. *Neurosurgery* **54,** 131-142 (2004).

89. R. Bai, C. S. Springer, D. Plenz, P. J. Basser, Fast, Na^+/K^+ pump driven, steady-state transcytolemmal water exchange in neuronal tissue: A study of rat brain cortical cultures. *Magn. Reson. Med.* **79,** 3207-3217 (2018). [DOI:10.1002/mrm.26980]

90. R. Meli, C. Pirozzi, A. Pelagalli, New perspectives on the potential role of aquaporins (AQPs) in the physiology of inflamation. *Front. Physiol.* **9,** 101 (2018).

91. L. S. King, D. Kozono, P. Agre. From structure to disease: The evolving tale of aquaporin biology. *Nat. Rev. Mol. Cell Biol.* **5,** 687-698 (2004).

92. B. Yang, A. S. Verkman, Water and glycerol permeabilities of aquaporins 1-5 and MIP determined quantitatively by expression of epitope-tagged constructs in Xenopus oocytes. *J. Biol. Chem.* **272,** 16140-16146 (1997).

93. L. Hertz, The glutamate-glutamine (GABA) cycle: Importance of late postnatal development and potential reciprocal interactions between biosynthesis and degration. *Front. Endochrinol.* **4,** 59 (2013).

94. G. W. Koch, E. Schwartz, Isotopic labeling of metabolic water with $^{18}O_2$. *Rapid Comm. Mass Spectrom.* **37,** e9447 (2023).

95. Q. Kuang, P. Purhonen, H. Hebert, Structure of potassium channels. *Cell. Mol. Life Sci.* **72,** 3677-3693 (2015).

96. R. P. Rand, V. A. Parsegian, D. C. Rau, Intracellular osmotic action. *Cell. Mol. Life Sci.* **57,** 1018-1032 (2000).

97. H. Wiebe, J. Spooner, N. Boon, E. Deglint, E. Edwards, P. Dance, N. and Weinberg, Calculation of molecular volumes and volumes of activation using molecular dynamics simulations. *J. Phys. Chem. C* **116,** 2240-2245 (2012). [doi.org/10.1021/jp209088u]

98. C. D. Hubbard, R. van Eldik, Mechanistic information on some inorganic and bioinorganic reactions from volume profile analysis. *Inorg. Chim. Acta* **363,** 2357-2374 (2010). [doi: 10.1016/j.ica.2009.09.042]

99. A. Drljaca, C. D. Hubbard, R. van Eldik, T. Asano, M. V. Basilevsky, W. J. le Noble, Activation and reaction volumes in solution. 3. *Chem. Revs.* **98,** 2167-2289 (1998).

Appendix A.1. *Acronyms and Symbols.*

$\langle A \rangle$ tissue or voxel mean cell surface area
AQP aquaporin
AQP4 aquaporin 4
AQP4KO aquaporin 4 knock-out
ATP adenosine triphosphate
ATP_i intracellular ATP
AWC active trans-membrane water cycling
a_e $\rho \langle A \rangle$
a_{H2O} water thermodynamic activity
CEST chemical exchange saturation transfer
ΔG Gibbs free energy change
ΔG^0 Standard ΔG
ΔG_{YYY} ΔG for substrate, enzyme, reaction; YYY
$\Delta G_{YYY}(effl)$ ΔG for YYY efflux
$\Delta G_{YYY}(infl)$ ΔG for YYY influx
$\Delta H_{YYY}(infl)$ enthalpy change for YYY influx
$\Delta \pi_{oi}$ trans-cytolemmal osmotic gradient (in – out)
$\Delta S_{YYY}(infl)$ entropy change for YYY influx
$\Delta V^{\dagger\dagger}$ activation volume
d   density (mass/volume)
EAAT1 excitatory amino acid transporter 1
$E_{m,oi}$ trans-membrane electrical potential (in – out)
effl   an efflux process
FDG 2-deoxy-2-[18]Fluoro-D-glucose
f_{H2O} water activity coefficient on the mole fraction scale (X)
f_W tissue water volume fraction
$f_{W,i}$ intracellular f_W
$f_{W,o}$ extracellular f_W
ϕ osmotic coefficient
G   Gibbs free energy
GABA γ-amino butyric acid
GAT1 GABA transporter 1
GLUT1 glucose transporter 1
glc   glucose
gln   glutamine
glu   glutamate-
γ_{H2O} water activity coefficient on the molality scale (m)
^1H_2O water proton MR signal
H_2O_i intracellular water molecule
H_2O_o extracellular water molecule
$H_3O_i^+$ intracellular hydronium ion
$H_3O_o^+$ extracellular hydronium ion
iGlutR an ionotropic glutamate-receptor
infl an influx process
K_i^+ intracellular K^+
K_o^+ extracellular K^+
KCC4 potassium, chloride transporter 4
k   kinetic rate constant
k_{io} cellular water efflux k ($1/\tau_i$)
$k_{io}(a)$ active k_{io} contribution
$k_{io}(p)$ passive k_{io} contribution
$k_{oi}(p)$ passive cellular water influx k contribution
M   molarity concentration scale
MADI metabolic activity diffusion imaging
MCT1 mono-carboxylate transporter
MD molecular dynamics
MRI magnetic resonance imaging
MR_{YYY} YYY metabolic rate
$MR_{glu}(consump)$ MR of glucose consumption
$MR_{glu}(infl)$ MR of glucose influx
$MR_{Glu-gln}(cycle)$ MR of glutamate-glutamine cycling
MR_{O2} MR of O2 consumption
$^cMR_{YYY}$ cellular metabolic rate for YYY process = $^tMR_{YYY}/\rho$

$^tMR_{YYY}$ tissue metabolic rate for YYY process = ρ^cMR_{YYY}
MW_{H2O} molecular mass ("weight") of water
m   molality concentration scale
mOsM milli-osmolarity concentration scale
mOsm milli-osmolality concentration scale
mOsX milli-osmole fraction concentration scale
μ_{YYY} chemical potential of YYY; $(\partial G/\partial n_{YYY})_{T,P,n(\neq nYYY)}$
Na_i^+ intracellular Na^+
Na_o^+ extracellular Na^+
NKA Na^+,K^+-ATPase (sodium pump)
NKCC1 sodium, potassium, chloride transporter 1
NMR nuclear magnetic resonance
n number of moles
OsM osmolarity concentration scale
OsM_{YYY} YYY OsM
Osm osmolality concentration scale
Osm_{YYY} YYY Osm
OsX osmole fraction concentration scale
$OsX_{YYY,i}$ intracellular YYY OsX
$OsX_{YYY,o}$ extracellular YYY OsX
P   hydraulic (mechanical) pressure
P^0 standard state P
P_d diffusional permeability coefficient (exchange)
$P_d(p)$ passive P_d
P_{dYYY} YYY P_d
P_f flux (flow) permeability coefficient
P_{fYYY} YYY Pf
P_i intracellular P
P_o extracellular P
PC   potassium channel
PET positron emission tomography
p   tissue water mole fraction ("population")
p_i intracellular p
p_o extracellular p
per   permeant particle
R   ideal gas constant
ρ cell (number) density
S_{YYY} stoichiometric coefficient of YYY
SGLT sodium/glucose co-transporter
SS   trans-cytolemmal NMR shutter-speed
TGN020 specific AQP4 inhibitor
τ_i mean H_2O_i molecule lifetime ($1/k_{io}$)
$\langle V \rangle$ tissue or voxel mean cell volume
$_Vm$ volume molality concentration scale
$_Vm_i$ intracellular $_Vm$
$_Vm_o$ extracellular $_Vm$
v   tissue volume fraction
v_e extracellular v (ECV) (= $1 - v_i$)
v_i intracellular v [ρV]
X   mole fraction concentration scale
X_{YYY} YYY X
$X_{YYY,i}$ intracellular YYY X
$X_{YYY,o}$ extracellular YYY X
x   water cycling stoichiometry [H_2O/ATP]
x′   x for individual water co-transporter
[YYY] YYY concentration
[YYY_c] compartmental YYY concentration
[YYY_i] intracellular YYY concentration
[YYY_o] extracellular YYY concentration
[YYY_t] tissue YYY concentration = $v_i[YYY_i] = v_e[YYY_o]$
yL   yocto liter (= 10^{-24} L = 1 (nm)^3))
y_{H2O} water activity coefficient on the molarity scale (M)
Z_{YYY} signed electrical charge of YYY

A.2. Compartmental Water Thermodynamic Activities.

Solvent mole fraction values as large as $OsX_{H2O,o}$ = 0.995 and $OsX_{H2O,i}$ = 0.993 (Table 1) "fail to emphasize the departure from ideality indicated by the activity coefficient of the *solute*" ((45), p.29). However, here the *solvent* activity coefficient difference is also very small. Since the water activity a_{H2O} (dimensionless on this scale) is always so close to unity, it requires sophisticated apparatus for high-precision direct water vapor pressure determinations to measure (45). Using these, it has been found a_{H2O} is inversely, and exponentially, related to the solute osmolality, Osm_{solute}, *via* the empirical *osmotic coefficient*, φ, **Equation (A.2.1)** (19,45,46; (63), p.12). MW_{H2O} is the water molecular

$$a_{H2O} = e^{-\left(\varphi \bullet MW_{H2O} \bullet Osm_{solute}/1000\right)} = e^{-(\varphi \bullet 0.018054 \bullet Osm_{solute})} \qquad (A.2.1)$$

weight (mass), 18.0154 g/mole(H_2O). The dimensionless product ($MW_{H2O} \bullet Osm_{solute}/1000$) is the solute/solvent mole ratio (1000 is (g/kg)), and essentially the solute mole fraction, OsX_{solute}. The latter is also dimensionless, and since the exponent must be dimensionless, φ is also dimensionless. Equation (A.2.1) provides a way to evaluate a_{H2O}. (When Osm_{solute} is zero (pure water), a_{H2O} is exactly one.)

How might we more fruitfully evaluate Eq. (4) when per = H_2O? Solutions as complex as those in Table 1 have never been subjected to precise measurements such as those of the last paragraph. However, pure NaCl and KCl solutions have been so studied, at 298 K ((45), p.476). We could take as a surrogate interstitium 0.290 Osm NaCl, and a surrogate cytosol 0.375 Osm KCl. (A glance at the metabolomics study mentioned above (47) indicates a K^+glutamate$^-$ solution would be a better intracellular surrogate. However, that has never been studied.) Their respective osmotic coefficients are φ_o = 0.9249 and φ_i = 0.9141 ((45), p.476) and, *via* Eq. (A.2.1), give $a_{H2O,o}$ = 0.995 and $a_{H2O,i}$ = 0.994. Using these activities in Eq. (4) gives $\Delta G(infl)$ = **-** 0.003 kJ/mole for H_2O influx. This free energy change is again so small that correcting for temperature and actual solute content would make no difference. The conclusion is inescapable: there is a very small trans-membrane chemical potential difference (~3 J/mole) for water to enter the cell.

A.3. Intracellular Pressure and Kinetics (Reaction Activation Volumes).

Mechanical pressure can affect transporter free energies (as above), and macromolecular structures (96), but can also exert an effect on chemical reaction kinetics. This is characterized by the volume of activation, $\Delta V^{\dagger\dagger}$, the difference between the partial molar volume of the reaction transition state and the partial molar volume of the reactants (97-99). The pressure dependence of a rate constant, k, is expressed in **Equations (A.3.1) and (A.3.2)**,

$$\frac{\partial lnk}{\partial P} = -\frac{\Delta V^{\dagger\dagger}}{RT} = -\frac{\Delta V^{\dagger\dagger}}{25.4} \tag{A.3.1}$$

$$\frac{k_A}{k_B} = e^{\left[\left(\Delta V^{\dagger\dagger}/RT\right)(P_B - P_A)\right]} \tag{A.3.2}$$

where RT is 25.4 L•atm/mole at 310 K. If a process has $\Delta V^{\dagger\dagger}$ = 6 L/mole, a 10 atm pressure increase will cause its rate constant to decrease by 90%. Six L/mole is 10^{-23} L/molecule, or 10 yL/molecule (1 yocto liter = 1 yL = 10^{-24} L = 1 (nm)^3). The plasma membrane NKA is a macromolecular complex of mass 145 kDa (145,000 g/mole). Thus (assuming d = 1 g/mL), it has a partial molar volume of approximately 242 yL/molecule. A 6 yL $\Delta V^{\dagger\dagger}$ would represent only a 2.5 % volume change. For P_i = 1.5 atm (the maximum P_i reported), P_o = 1 atm, and $\Delta V^{\dagger\dagger}$ = 6 L/mole, Eq. (A.3.2) yields $k_{1.5\ atm}$ = 0.89 $k_{1\ atm}$, an 11% decrease.

It is easy to imagine $\Delta V^{\dagger\dagger}$ values of this magnitude, or greater. In a molecular dynamics study of a small 6.5 kDa protein, Persson and Halle report a relatively constrained, 2 kDa domain undergoes numerous spontaneous 3% conformational volume fluctuations in a 1000 μs period (60). Most membrane transporters are rather large macromolecular complexes. It is quite likely they undergo significant absolute volume changes in reaching their transition states. Thus, their enzyme activities could be quite sensitive to intracellular pressure development. Some would slow down, and some would speed up: $\Delta V^{\dagger\dagger}$ quantities can be negative (97-99). Furthermore, the activities of most membrane transporters are coupled to each other *via* common substrates or products (Fig. 1). Since the cell has evolved to have trans-membrane fluxes in homeostatic balance, it is likely a P_i change would disturb the latter. The consequent flux changes could serve an auto-regulatory function.