Human natural killer cells can activate NLRP1 and NLRP3 inflammasomes and drives pyroptosis

Antonio Astorga-Gamaza[1,2], Inés Muela-Zarzuela[1], Juan Miguel Suárez-Rivero[1], Juan López-Pérez[2,3],

Raquel de la Varga-Martínez[2,3], Auxiliadora Mazuecos[4] and Mario D. Cordero[1,5, &]

[1] Department of Molecular Biology and Biochemical Engineering, Universidad Pablo de Olavide, 41013 Seville, Spain.

[2] Instituto de Investigación e Innovación Biomédica de Cádiz, INiBICA, Hospital Universitario Puerta del Mar, Cádiz, Spain.

[3] Department of Immunology, Puerta del Mar Hospital, 11009 Cádiz, Spain.

[4] Department of Nephrology, Hospital Universitario Puerta del Mar, Cadiz, Spain.

[5] CIBER de Enfermedades Respiratorias (CIBERES), Instituto de Salud Carlos III, 28220 Madrid, Spain.

Running tittle: Natural killers can activate inflammasomes

[&]Corresponding author: Mario D. Cordero, PhD

Department of Molecular Biology and Biochemical Engineering, Universidad Pablo de Olavide, 41013 Seville, Spain. Email: mdcormor1@upo.es.

Summary

Innate immunity relies on inflammasomes as key components, defending the host against diverse harmful stimuli by orchestrating the release of pro-inflammatory cytokines and initiating pyroptotic cell death. While extensively studied in myeloid cells, the involvement of natural killer (NK) cells in inflammatory responses through inflammasome signaling remains underexplored. In this study, we elucidate the activation of the inflammasome sensors NLRP1 and NLRP3 in human primary NK cells upon treatment with nigericin and blockade of dipeptidyl peptidases (DPP) using Talabostat (Val-boroPro). Our findings demonstrate the induction of pyroptotic cell death in a subset of NK cells following these stimuli, characterized by the cleavage and activation of gasdermin D, a lytic pore-forming protein. Moreover, we observe the release of lactate dehydrogenase (LDH) and small amounts of interleukin-18 (IL-18). Notably, differential responses are noted between CD56[dim] and CD56[bright] NK cell subsets following pro-inflammatory stimulation.

Furthermore, analysis of samples from patients with renal dysfunction reveals sustained

inflammasome activation in NK cells, particularly NLRP1 and NLRP3, with a tendency towards a

more pro-inflammatory phenotype shortly post-kidney transplantation. These findings underscore

the significance of considering NK cells in the context of inflammation studies.

Keywords: Natural Killers, NLRP3-Inflammasome, NLRP1-Inflammasome, Kidney transplantation

Introduction

Inflammation is part of the body's defense response to harmful stimuli. However,

exacerbated inflammation is also associated with a range of diseases. In this regard, the activity of

natural killer (NK) cells has been shown to highly depend on the context, helping to reduce or even

promote inflammatory processes [1].

NK cells are innate lymphoid cells characterized by a huge diversity of phenotypes and

functions [2, 3], although primarily categorized into two main subpopulations based on the neural

cell adhesion molecule CD56 density. CD56^{dim} NK cells, known for their potent cytolytic activities,

and CD56^{bright} NK cells, distinguished by their cytokine secretion capabilities [4]. They play a pivotal

role in immune surveillance, effectively targeting malignant and infected cells to control

intracellular pathogens and shaping adaptive immune responses [5]. Equipped with a huge array

of cell surface receptors, NK cells sense stress signals in target cells, distinguishing between 'self'

and 'non-self' or 'aberrant' targets under normal physiological conditions [6]. Upon activation, NK

cells release cytotoxic granules containing perforin and granzyme. Also, they secrete interferon

(IFN)-γ and tumor necrosis factor (TNF), and immunoregulatory molecules such as interleukin (IL)-

10 and the growth factor GM-CSF, as well as chemokines including CCL3 (MIP-1α), CCL4 (MIP-1β),

CCL5 (RANTES), and CXCL8 (IL-8) [7]. Through these mechanisms, NK cells can drive inflammation

and modulate the activity of other immune cells.

A key part of the inflammatory process and the innate immune response is the assembly

of inflammasomes. These cytosolic multi-protein complexes regulate tissue inflammation in

response to multiple cellular stress patterns, including pathogen-associated molecular patterns

(PAMPs) such as bacterial lipopolysaccharide (LPS), or cellular damage-associated molecular

patterns (DAMPs), which include extracellular ATP and changes in intracellular ion concentrations.

In this regard, inflammasome biology has been mostly characterized in macrophages, and at least

partially studied in other cell types such as endothelial cells, fibroblasts, epithelial cells, and T cells

[8-11]. However, it is unknown if NK cells are equipped with functional inflammasome

components, and thus, this could represent an underexplored arm of the diverse immune

mechanisms mediated by these cells and an important source of inflammation.

The first steps in the triggering of inflammasomes comprise the detection of the damage

patterns via sensors of several families such as the NOD-like (NLR), Toll-like (TLR), AIM2-like (ALR),

or the caspase recruitment domain (CARD)-containing family of proteins, that include the

members NLRP1, NLRP3, NLRC4, AIM2, and CARD8, among others [12]. Following activation, some

sensors interact with adaptor proteins such as ASC (apoptosis-associated speck-like protein

containing a CARD), composing a pre-complex that then recruits inflammatory caspase-1. Next to

terminal assembly, the activation of caspase-1 in the inflammasome can lead to the proteolytic

cleavage and maturation of the IL-1 family cytokines IL-1β and IL-18 [13]. This pathway is known as

the 'canonical' activation. Another usual key event after the inflammasome activation is the

cleavage of gasdermins, a family of pore-forming proteins, into its N-terminal active domain which

induces a proinflammatory form of regulated cell death termed pyroptosis [14].

Here, we conducted a transcriptomic analysis of NK cells derived from various tissues,

revealing the expression of genes encoding several inflammasome components. Additionally, we

provide proteomic evidence demonstrating the expression of canonical NLRP3 and NLRP1

inflammasome components in both CD56^{dim} and CD56^{bright} NK cell subsets. More importantly, we

demonstrate the functionality of these inflammasomes in NK cells, as evidenced by IL-18 secretion

and induction of pyroptotic cell death. Interestingly, we observe distinct responses to pro-

inflammatory stimulis between the two NK cell subpopulations. Furthermore, we identify signs of

inflammasome activation in NK cells from patients with renal dysfunction, which persist after

kidney transplantation and despite immunosuppression. Our findings suggest that NK cells should

be considered in inflammation research and as potential therapeutic targets, as they might play an

important role in both pathogen defense and inflammatory disorders through inflammasome

signaling   and   pyroptosis.

RESULTS

Expression of inflammation-related genes in human NK cells from different tissues

To gain knowledge on the potential of human NK cells for shaping inflammatory responses and constituting functional inflammasomes, we conducted a transcriptomic analysis of NK cells from diverse tissues from a publicly available RNA-seq dataset [3], and assessed the expression of a list of genes related to inflammation. As described, donors were free of cancer, chronic diseases, and seronegative for hepatitis B, C, and HIV [3]. From a total of 2,686 genes, we found a significant expression of 1,619 in NK cells from blood, bone marrow, lymph nodes, lungs, and spleen. These genes appeared to be involved in different processes such as the regulation of the NF-κB activity, apoptosis, TNF signaling, cellular response to lipopolysaccharide, NOD-like and Toll-like receptor signaling, IL-6 and IL-1β production (**Figure 1A**). Next, we performed a detailed analysis of the expression of short-listed genes reported to play a role in activating inflammasome multiprotein complexes at different stages (**Figure 1B, C, D, E, and F**). Interestingly, we noticed the expression in NK cells of guanylate-binding proteins (GBP), which target intracellular pathogens and can mediate host defense via inflammasome induction [15, 16]. Moreover, we found expressed other receptors of DAMPs and PAMPs such as TLR-2-3 and 4, NLRP1, NLRP3, NLRP6, NLRC3, and NLRC4, the interferon-inducible protein AIM-2 and CARD. NK cells also expressed PYCARD, the gene coding for ASC protein, required to constitute canonical inflammasomes, and the serin proteases DPP8 and DPP9, whose pharmacological inhibition by Val-boroPro (Vbp) (Talabostat) is reported to activate both NLRP1 and CARD8 inflammasomes [17, 18]. Furthermore, NK cells from all analyzed tissues expressed downstream effectors in the inflammasome pathway such as caspase-1, IL-1β, IL-18, and gasdermin D genes (**Figure 1B, C, D, E, and F**). Moreover, we analyzed differential expression between the NK subpopulations CD56^{dim} and CD56^{bright} and found only a few significant differences that varied depending on the tissue location. This was the case of GBP5 and TLR3 in the blood (**Figure 4B**), and Caspase-1 in the lung (**Figure 4C**), more expressed in CD56^{dim} NK cells. Of note, NK cells exhibit an intrinsic, highly heterogeneous nature intra- and inter-individual [2]. Thus, larger datasets are likely needed to reveal more differences between different subpopulations.

Our results indicate that NK cells, both the two main subpopulations CD56^{dim} and CD56^{bright}, express the genetic elements needed to constitute functional inflammasomes.

Protein expression of NLRP1 and NLRP3 and the adaptor ASC in NK cells

Next, we aimed to assess the expression of the protein sensors NLRP1 and NLRP3, which have

been implicated in the regulation and pathogenesis of many conditions. We studied the

expression of the proteins NLRP1, NLRP3, and the adaptor ASC in NK cells by flow cytometry, using

blood-derived primary NK cells previously isolated by FACS (fluorescence-activated cell sorting)

(**Supplementary Figure 1**). The median purity of the cells was 90.7% and 94.8% for CD56^{dim} and

CD56^{bright} NK cells, respectively. A representative example of the staining can be found in **Figure**

2A. These proteins were widely expressed in NK cells, with slightly higher frequencies of ASC and

NLRP1 in the CD56^{bright} subpopulation (**Figure 2B and C**), but no differences in the percentage of

NLRP3 (**Figure 2D**) nor in the intensity of expression (mean fluorescence intensity, MFI) of any of

them (**Figure 2E, F, and G**). Therefore, NLRP3 and NLRP1 inflammasomes may play a role in NK

cell-mediated immunity if functional.

Activation of the NLRP3 and NLRP1 inflammasome complexes in NK cells triggered by specific
activators

As a next step, we directly tested the functionality of inflammasomes in NK cells. For that, we

treated NK cells previously isolated by FACS with known inductors of the NLRP3 and NLRP1-

inflammasomes. We included conditions of cells cultured with lipopolysaccharide (LPS) plus ATP

(LPS-ATP), nigericin (Nig), and the DPP8/DPP9 inhibitor Vbp for both CD56^{dim} and CD56^{bright} NK

subpopulations separately. Inflammasome activation was first assessed by measuring the content

of inflammatory cytokines in culture supernatants. Remarkably, we detected secretion of pro-

inflammatory IL-18 upon stimulation with nigericin of CD56^{dim} (**Figure 3A**), but not CD56^{bright} NK

cells (**Figure 3B**). The opposite pattern was obtained for Vbp, which triggered IL-18 production in

CD56^{bright} (**Figure 3C**) but not in CD56^{dim} cells (**Figure 3D**). No increment was detected after LPS-ATP

stimulation (**Supplementary Figures A and B**). IL-1β was not induced in general in these in vitro

assays, with the exemption of one donor that also showed higher levels of IL-6 (**Supplementary**

Figures C, D, and E). Besides, we measured other immune mediators by a multiplex Luminex

immunoassay. We failed to detect secretion of IL-10, GM-CSF, and the NK-associated cytotoxic and

regulatory molecules IFN-γ, IFN-α, or TNF-α in our assayed conditions, suggesting that in vitro

treatment with nigericin and Vbp does not induce NK unspecific cytotoxicity and inflammatory

response, and that an inflammasome specific response is triggered. Interestingly, ICAM-1, the
adhesion receptor and regulator of inflammation, seemed to increase its concentration after LPS-
ATP and nigericin treatment of CD56^{dim} cells (**Supplementary Figures F and G**). This suggests that
different pro-inflammatory stimuli affect differently distinct NK subpopulations.

Second, we assessed cell death by pyroptosis, another key process that follows inflammasome
activation. We detected by western blot the activation of the N-terminal fragment of Gasdermin D
after nigericin and Vbp treatment of NK cells (**Figure 3E**). Of note, this was successfully detected
only in CD56^{dim} NK cells. However, the release of LDH, which confirms cell death, was more
prominent in the CD56^{bright} subpopulation (**Figure 3F and G**), raising the possibility that more
aggressive death of CD56^{bright} NK cells may hinder the detection of active gasdermin D. Both pro-
inflammatory cytokine secretion and gasdermin-dependent pyroptosis rely on the previous
cleavage of inflammatory caspases into its active form. Importantly, we observed increased active
fragments of caspase-1 in NK cells stimulated with nigericin and Vbp (**Figure 3H**).

To obtain more clues about the activation of inflammasomes in NK cells, we assessed the
expression of proteins within control and treated cells by flow cytometry. The purity of the cells
employed in these assays was always higher than 90.0%. Moreover, any remaining cell
contaminants in the culture were discarded from our analysis using lineage markers for NK cells in
the flow panel. Treatment of NK cells, CD56^{dim} or CD56^{bright}, with LPS-ATP, nigericin, or Vbp, did not
alter the frequency of cells expressing NLRP3 (**Figures 3I and J**), NLRP1 (**Figures 3K and L**), or ASC
(**Figures 3M and N**). However, changes in the intensity of expression (MFI) were detected in the
case of nigericin exposure of CD56^{bright} NK cells, with a reduction in NLRP3 levels (**Figure 3O**). This
was not observed for the rest of the proteins or stimulus (**Supplementary Figures 3H, I, J, K, and
L**). This may indicate that nigericin and Vbp promote the reorganization of these molecules to
assemble the inflammasomes rather than drastically changing their expression levels. Also,
nigericin may promote more aggressive pyroptotic death of CD56^{bright} NK cells, explaining the
lower detected levels of NLRP3 and higher LDH release. Moreover, we studied the formation of
specks by flow cytometry as previously reported [19]. In the formation of inflammasomes, ASC
interacts with upstream sensors such as NLRP1 and NLRP3 that subsequently trigger its assembly
into a structure known as a speck composed of multimers of ASC dimers. The speck can be
detected by flow cytometry using the time of flight (TOF) signal of ASC, as it concurs with higher
pulse height to area (H:A) (**Figure 3P)**. We consistently observed signs of speck in CD56^{dim} NK cells

after nigericin stimulation (**Figure 3Q**). These results resembled that of IL-18 secretion, as
treatment with Vbp also induced speck in 5 out of 7 independent assays in CD56[bright] NK cells
(**Figure 3R**). Notably, we did not detect speck in LPS-ATP or Vbp treated CD56[dim] NK cells
(**Supplementary Figures 2N and O**), nor consistently in LPS-ATP and nigericin treated CD56[bright]
subpopulation (**Supplementary Figures 2P and Q**). Of note, we believe that pyroptosis of NK cells
activating the inflammasomes may be hindering/underestimating the quantification of specks in
these assays.

Overall, our results indicate that both NLRP3 and NLRP1-inflammasomes are functional in NK
cells, particularly under nigericin and Vbp treatment. CD56[dim] and CD56[bright] NK cells behave
differently under pro-inflammatory stimulation.

NK cells from patients with chronic kidney disease display inflammasome activation

Recent data support the hypothesis that imbalanced inflammation might be an important
driver of kidney disease that may persist even after organ transplantation. We wondered if NK
cells may contribute to the inflammatory milieu in patients with kidney dysfunction. We collected
PBMCs from patients with chronic kidney disease listed in **Supplementary Table 1**. Western
blotting studies showed high levels of NLRP1, active caspase-1, and gasdermin-D in these patients
before and during the first week after kidney transplantation (**Figure 4A and Supplementary**
Figure 3), confirming the activation of inflammasomes and pro-inflammatory cell death by
pyroptosis of circulating immune cells. We isolated by FACS the CD56[dim] and CD56[bright]
subpopulations from a fraction of these PBMC, but it was technically challenging to perform the
same western blot assays likely due to low cell numbers. Importantly, flow cytometry analyses (as
represented in **Supplementary Figures 4A and B**) confirmed that nearly 100.0 % of NK cells in
these patients expressed NLRP1 (**Figure 4B**), about 50.0 % expressed ASC (**Figure 4C**), and a
smaller fraction NLRP3, which increased three days after transplantation (**Figure 4D**). We also
detected a fraction of NK cells with signs of speck, being enriched three days post-transplantation
(**Figure 4E and Supplementary Figure 4C**). Speck levels in NK cells were much lower than those
detected in monocytes, as expected (**Supplementary Figure 4D**). Additional analyses distinguishing
between specking versus non-specking cells showed higher levels of NLRP1 and NLRP3 in speck[+]
cells (**Figures 4F, G, H and I**), suggesting the activation of both inflammasomes in NK cells from
patients with renal dysfunction.

Discussion

In this study, we try to shed light on the biology of inflammasomes in human NK cells, as dedicated studies are lacking. These multiprotein complexes are involved in innate defense and inflammatory responses to pathogens. Similarly, NK cells constitute a first line of defense against aberrant targets. However, whether inflammasomes encompass one of their multiple immune mechanisms of action and how it may impact health are unknown. Inflammasomes are composed of sensors, adaptors, and effectors components, and their activation triggers the release of inflammatory cytokines and pyroptosis [20]. Here, we show the expression of transcripts encoding several inflammasome-forming proteins in NK cells distributed across different tissues. Sensors of several inflammasomes, including the members of the NALP family NLRP3 and NLRP1 were found expressed. Consistent with a prior report [21], we found a high constitutive expression of NLRP3. We also noticed high levels of NLRP1 and the adaptor protein ASC in NK cells. More importantly, we prove here that the NLRP3 and NLRP1 inflammasomes can be assembled and are functional in these cells.

Several activators of the NLRP3 and NLRP1 inflammasomes have been identified, being reported potassium ion (K^+) efflux as one of the most potent in the case of NLRP3 [22], and recently also proven to activate NLRP1 [23], and inhibition of the serin dipeptidyl peptidases DPP8 and DPP9 for NLRP1 activation [17, 18]. Here, we used Nigericin, a toxin derived from *Streptomyces hygroscopicus* that facilitates H^+/K^+ anti-port across cell membranes and thereby causes potassium efflux [24], and Vbp, an inhibitor of the inhibitory interaction between DPP8/DPP9 and NLRP1 [25], to assess the functionality of both inflammasomes in NK cells. Our results revealed changes that are compatible with inflammasome activation, including secretion of IL-18 and more notable pyroptosis. Also seemed to indicate that the components of the NLRP3 and NLRP1 inflammasomes more likely undergo a reorganization rather than upregulation upon stimulation with nigericin or Vbp. Moreover, we found differences in the response of CD56[dim] and CD56[bright] NK cells to both compounds. This is not completely unexpected given the high plasticity and diversity in phenotypes and functions of NK cells [2, 3]. Nigericin induced the secretion of IL-

18 by CD56^{dim} NK cells while Vbp did the same with the CD56^{bright} subpopulation. Regarding the

pore-forming protein gasdermin D, capable of inducing pyroptosis, its active form was detected in

CD56^{dim} NK cells treated with nigericin and Vbp. LDH levels were also increased in the culture of

these cells, particularly in those treated with nigericin. Of note, although we were not able to

detect active gasdermin D in CD56^{bright} cells, a notable release of LDH by these cells was observed.

The release of LDH is considered a surrogate of cell pyroptosis as large proteins are usually filtered

out in the absence of cell death [26]. Thus, it may be plausible that CD56^{bright} NK cells undergo a

more aggressive pyroptosis upon *in vitro* activation of the NLRP3/NLRP1 inflammasomes,

hindering the detection of active gasdermin-D. Also, the low numbers of these cells in the blood,

which usually represent less than 10% of circulating NK cells [27], may suppose a technical

challenge for the detection of proteins. Different frequencies and proportions of NK cell

subpopulations can be found in tissues like the bone marrow, spleen, lung, liver, lymph nodes,

tonsils and intestines [3]. Thus, further studies with different tissue samples, NK subpopulations

and inflammasome-triggering agents or conditions could add valuable insight in the biology of

these lymphocytes and the inflammatory field.

To our knowledge, this is the first report clearly showing that NLRP3/NLRP1 inflammasome

activation and pyroptosis take place in human NK cells. Nevertheless, prior results in mice support

our results that canonical activation of inflammasomes occur in these immune cells, showing that

burn injury and radiation exposure induce the activation of caspase-1 in many cell subsets

including NK cells [28, 29]. Importantly, Fong et al also showed that the human pathogen group B

Streptococcus directly interacts with NK cells and suppress a pyroptotic-like response via the

interaction with siglec-7 [30]. In this regard, our results did not show a typical inflammasome

response after the LPS-ATP stimulation of NK cells. However, this could be explained by the low

surface expression of TLR4 in these cells [31]. Thus, we do not rule out the possibility that direct

bacterial infection might activate some of the inflammasomes in NK cells. In fact, our

transcriptomic analysis showed that many other inflammasome sensors are transcriptionally

expressed in these cells, including NLRP6, NLRC4 or AIM-2, although if they are functional remain

unknown and could be more suitable after direct infections. Thus, the results presented here

provides new insight in the biology of NK cells that might support a role of these lymphocytes in

different contexts via inflammasome activation. Elucidating when these responses take place and

whether are beneficial or harmful could help in the identification of new targets in inflammatory

disorders.

Inflammasome activity has been associated to chronic kidney disease and the rejection of
the graft in kidney-transplanted patients, particularly linked to AIM-2, NLRP3, and more recently
NLRP1 [32-36]. Kidney transplantation is considered the best therapy for patients with end-stage
kidney disease, being associated with better quality of life and patient survival compared with
dialysis. However, a fraction of kidney transplant recipients still faces short-term or long-term graft
rejection, in which inflammation seems to play a relevant role [37]. Here, we observed signs of
NLRP3 and NLRP1 inflammasome activation as well as activation of gasdermin-D in circulating
immune cells, including NK cells, in patients with end-stage kidney disease before and after
receiving a donor kidney. Previously, it has been shown that post-transplantation NK cells subsets
can change even at the peripheral level, with variations in number and phenotype [38]. Here, we
show the acquisition of a more proinflammatory phenotype in NK cells shortly post-
transplantation, characterized by higher levels of NLRP1, NLRP3 and ASC. This suggests that
different inflammatory milieus in vivo could induce the activity of inflammasomes not only in
classically studied subsets such as monocytes and macrophages, but also innate lymphoid cells
such as NK cells. Until now, two mechanisms are considered the main drivers for the graft
rejection in transplanted patients, the T-cell mediated rejection, and/or the antibody-mediated
rejection. In both, NK cells have been shown to cooperate, through maturation of dendritic cells
and activation of T cells and mediating a pathogenic ADCC response triggered by anti-HLA
antibodies in CD56^{dim} CD16-expressing NK cells [39]. Our results add evidence that NK cells might
also contribute to pathogenic inflammation in renal injured patients through pro-inflammatory
activity via inflammasome signaling. Although larger studies and longer follow-up periods are
needed, our results are tempting to speculate that inflammasome is a pathway by which NK cells
might influence renal disease and the graft outcome. Moreover, they might play a role in several
other inflammatory diseases such as ischemic stroke, sepsis or cancer, where persistent
inflammasome activity and activation of gasdermin D has been observed to cause undesired
effects including organelle dysfunction, cell lysis and persistent release of cytokines [14]. Also, it
will be relevant to determine other contexts where inducing inflammasome activity of NK cells
could be beneficial.

In conclusion, our findings underscore the importance of including NK cell subpopulations in future
studies investigating inflammasome functionality.

Acknowledgments

M.D.C is supported by PI21/01656 grant from Instituto de Salud Carlos III, Spain, EMC21_00033 EMERGIA program from Junta de Andalucia, AthenaDAO and PRF 2021-78 from Progeria Research Foundation. A.A.G was supported by the Spanish Secretariat of Science and Innovation post-doctoral contract Juan de la Cierva (FJC2021-047304-I). The funders had no role in study design, data collection and analysis, the decision to publish, or the preparation of the manuscript.

Author contributions

Conceptualization: A.A.G, M.D.C; Methodology: A.A.G, I.M.Z; Investigation: A.A.G, JM.S.R, M.D.C; Resources: A.A.G, J.L.P, R.V.M, M.D.C; Visualization: A.A.G, I.M.Z, M.D.C; Supervision: A.A.G, M.D.C; Funding acquisition: M.D.C; Writing – Original Draft: A.A.G; Writing –Review & Editing: A.A.G, M.D.C.

DECLARATION OF INTEREST

The authors declare no competing interest.

    FIGURE LEGENDS

    **Figure 1. RNA-Seq analysis of inflammatory transcriptional signatures in circulating and tissue
resident human NK cells.** Cells were isolated from donors without cancer or any other chronic
disease, and seronegative for hepatitis B, C, and HIV viral infections. **A)** Significance of the most
represented immune pathways according to the expression of a list of 2,686 genes related to
inflammation. Functional gene annotation was performed using DAVID. **B)** Heatmap of the
expression of short-listed genes key in the activation and triggering of effector functions of
inflammasomes in circulating NK cells. An analysis of the differential expression of these genes
between CD56^{bright}CD16^{-} and CD56^{dim}CD16^{+} NK cell subsets is also shown, and genes showing a
different pattern of expression between both NK subpopulations are marked with a red box. The
same is shown for **C)** NK cells in the lung, **D)** NK cells in the lymph nodes, **E)** NK cells in the bone
marrow, and **F)** NK cells in the spleen. Differentially expressed genes were identified using linear
models (Limma-Voom) and P values adjusted for multiple comparisons by applying the Benjamini-
Hochberg correction.

**Figure 2. Expression of the inflammasome sensors NLRP1 and NLRP3 and the adaptor protein
ASC in human NK cells.** Two subpopulations of NK cells: CD56^{dim}CD16^{+} and CD56^{bright}CD16$^{-/dim}$,
were isolated by cell sorting and interrogated for the expression of the mentioned proteins by
flow cytometry (N=8). **A)** Flow-cytometry plots showing a representative example for the
expression of NLRP1, NLRP3 and ASC in CD56^{dim} and CD56^{bright} NK cells, and the corresponding
Fluorescence Minus One (FMO) control for the proper gating. **B)** Frequency (%) of expression of
ASC, **C)** Frequency of NLRP1, **D)** Frequency of NLRP3, and Mean Fluorescence Intensity (MFI) signal
for each marker in **E)**, **F)** and **G)**. Median with range is represented. Statistical comparisons were
performed using the Wilcoxon matched-pairs signed-rank test.

Figure 3. **Activation of the NLRP1 and NLRP3 inflammasomes and induction of pyroptosis in
human NK cells ex vivo.** Isolated NK cells from blood samples were cultured in the presence of the
inflammasome activators LPS (1 µg/ml for 4 hours) and ATP (5 mM for 30 mins), nigericin (10 µM
for 30 mins) or Vbp (1 µM for 4 hours). After that, the concentration of cytokines of the IL-1 family
IL-18 and IL1β as well as lactate dehydrogenase (LDH) release were measured in the culture
supernatants. Changes in the cell expression of inflammasome sensors and downstream effectors
in the inflammatory cascade such as caspase-1 and gasdermin D were also measured by western
blot and flow cytometry. **A)** Concentration of IL-18 in cell cultures of untreated CD56^{dim} NK cells or
treated with nigericin (n=6). **B)** The same for CD56^{bright} NK cells. **C)** IL-18 concentration in cell
cultures of CD56^{bright} NK cells stimulated with Vbp (n=5). **D)** IL-18 levels after stimulation with Vbp
of CD56^{dim} NK cells. **E)** Western blot analysis of the full length and cleaved form of gasdermin D in
CD56^{dim} NK cells in basal conditions and after nigericin and Vbp exposure. Representative result
from n=5 independent experiments. **F)** LDH release in cell cultures of stimulated CD56^{dim} NK cells
(n=6), and **G)** stimulated CD56^{bright} NK cells (n=6). **H)** Western blot analysis of the precursor and
active forms of caspase-1 in CD56^{dim} NK cells in basal conditions and after nigericin and Vbp
exposure. Representative result from n=3 independent experiments. **I)** to **O)** represent the

frequency and MFI signal for the expression of NLRP1, NLRP3 and ASC in CD56^{dim} and CD56^{bright} NK
cells in non-stimulated and under pro-inflammatory stimuli with LPS-ATP, nigericin and Vbp
conditions (n=8). Graphs represent median with range. Statistical comparisons were performed
using the Wilcoxon matched-pairs signed-rank test. **P)** Representative flow cytometry plots of the
time-of-flight signal for ASC in NK cells, showing an increase in ASC area: height pulse, indicative of
specking, after nigericin in vitro stimulation. **Q)** Graph showing the frequency of CD56^{dim} NK cells
with signs of ASC specking after nigericin stimulation. **R)** The same for CD56^{bright} NK cells stimulated
with Vbp. Statistical comparisons were performed using the Wilcoxon matched-pairs signed-rank
test.

Figure 4. **Inflammasome activation in NK cells from patients with chronic kidney disease.** Blood
samples from patients with renal disease were collected at different time points, before and after
being subjected to kidney transplantation (n=7). Activation of the NLRP1 and NLRP3
inflammasomes in these patients was studied by western blotting and flow cytometry. **A)** Western
blot representative results in total peripheral blood mononuclear cells (PBMCs) from one renal-
injured patient (#FJP in **Supplementary Table 1**), showing the expression of NLRP1, NLRP3, pro-
active caspase-1, the active fragment of caspase-1, full-length gasdermin D, active gasdermin D,
and the control β-actin, pre-transplantation and 3 and 7 days after being transplanted. **B)**
Frequency of expression of NLRP1 in total CD56$^+$ NK cells. **C)** Frequency of ASC. **D)** Frequency of
NLRP3. **E)** Percentage of ASC specking in NK cells. Graphs represent median with range. Statistical
comparisons were performed using the Wilcoxon matched-pairs signed-rank test. Graphs in **F)** and
G) show the percentage and MFI of NLRP1 within speck$^-$ and speck$^+$ NK cells at the different time
points. The same for NLRP3 in **H)** and **I).** Graphs represent median with range. P values shown in
the graphs represent ANOVA Friedman test, and asterisks denote the multiple comparison Dunn's
test. *p<0.05; **p<0.01. Moreover, comparisons between the speck$^-$ and speck$^+$ NK cells in the
same time points were performed using the Wilcoxon matched-pairs signed-rank test.

MATERIALS AND METHODS

<u>Human Samples</u>

In this study, we used primary cells obtained from blood samples from healthy donors or people
with kidney disease and undergoing transplantation. Peripheral blood mononuclear cells (PBMCs)
were obtained from the Hospital Puerta del Mar in Cádiz, Spain. Study protocols were approved by
the corresponding Ethical Committees (Institutional Review Board numbers). All subjects recruited
for this study were adults who provided written informed consent. Information on renal disease
parameters from affected patients is summarized in Table S1. Gender and age are also indicated in
Table S1 unless not available. Since we utilized samples from donors of different genders in our
experiments, we can conclude that the results reported here apply to both men and women.
PBMCs were obtained by Ficoll-Paque density gradient centrifugation. PBMCs were cultured in

RPMI medium (Gibco) supplemented with 10% Fetal Bovine Serum (Gibco), 1% streptomycin-

penicillin (Thermo Fisher, Waltham, MA, USA, 11548876) (R10 medium), and maintained at 37ºC

in a 5% CO_2 incubator, when needed. NK cells were isolated from PBMCs by FACS (fluorescence-

activated cell sorting) (MELODY sorter) or using a commercial kit (MagniSort™ Human NK cell

Enrichment; eBioscience). The purity of the cells was always higher than 90%.

<u>Transcriptomic analysis of a publicly available RNAseq data</u>

We analyzed publicly available RNA-seq datasets. Data are available from Gene Expression

Omnibus (GEO) corresponding to the study with accession number GSE133383. This study

provides valuable transcriptomic data of two subpopulations of NK cells, $CD56^{dim}$ and $CD56^{bright}$, in

tissues difficult to obtain [3]. R version 4.1.2 was used to perform all bioinformatic analyses. Count

values were imported and processed using edgeR

[https://academic.oup.com/bioinformatics/article/26/1/139/182458?login=false]. Expression

values were normalized using the trimmed mean of M values (TMM) method and lowly-expressed

genes (<1 counts per million) were filtered out. Differentially expressed genes were identified

using linear models (Limma-Voom) [https://academic.oup.com/nar/article/43/7/e47/2414268],

and P values adjusted for multiple comparisons by applying the Benjamini-Hochberg correction.

Heatmaps were generated using the heatmap3 package [https://github.com/slzhao/heatmap3].

Glimma [https://github.com/Shians/Glimma] was used for interactive visualizations. Functional

gene annotation was performed using DAVID [https://david.ncifcrf.gov].

<u>Human NK cell isolation by FACS</u>

For Fluorescence Activated Cell Sorting (FACS), 80 million PBMCs were stained with LIVE/DEAD

AQUA viability (Invitrogen) for 20 minutes at RT. After washing with staining buffer (PBS 3% FBS),

cells were surface stained with anti-CD56-FITC (B159, Becton Dickinson), anti-CD3-PE-Cy7 (SK7,

Becton Dickinson), anti-CD16-BV786 (3G8, Becton Dickinson) and anti-CD4-APC (RPA-T4, Becton

Dickinson) antibodies for 20 minutes at RT. Cells were then washed with staining buffer and

immediately sorted using the MELODY Cell Sorter. We sorted the live populations $CD4^-CD3^-CD56^{dim}$

($CD56^{dim}$ NK cells) and $CD4^-CD3^-CD56^{bright}$ ($CD56^{bright}$ NK cells). The purity of the cells was >90% in all

cases.

<u>In vitro inflammasome activation assays</u>

For in vitro pro-inflammatory stimulation, NK cells were treated with 1μg/ml LPS (Sigma-Aldrich,

St. Louis, MI, USA, L4391-1MG) for 4 hours and 5 mM ATP (Santa Cruz Biotechnology, Santa Cruz,

CA, USA, sc-214507A) for 30 mins. NK cells were also stimulated with 1 μM Val-boroPro (Sigma-

Aldrich, St. Louis, MI, USA, 5314650001) for 4 hours or 10 μM nigericin for 30 mins. As the number

of isolated NK cells varied between donors, we maintained the condition of the cell culture,

stimulating the cells at a concentration of approximately 0,1M/ml, in 24-well plates for the

CD56^{dim} subpopulation and 96-well plates for the CD56^{bright} subpopulation. After the incubation,

we collected culture supernatants for LDH and cytokine measures, and cells for proteomics

analyses by flow cytometry or western blot.

<u>Flow cytometry</u>

For studying the protein expression, before and after pro-inflammatory stimulation, NK cells were

stained with LIVE/DEAD AQUA viability (Invitrogen) for 20 minutes at room temperature (RT).

After washing once with staining buffer (1X PBS 3% FBS), cells were surface stained with anti-

CD56-FITC (B159, Becton Dickinson), and anti-CD3-PE-Cy7 (SK7, Becton Dickinson) antibodies for

20 mins at RT. Next, we performed a washing step and cells were fixed and permeabilized with

Fixation/Permeabilization Solution (Becton Dickinson) for 20 minutes at 4ºC and then washed with

BD Perm/Wash buffer. After, cells were stained with rabbit anti-NLRP1 (A16212 ABclonal) for 20

mins at RT, washed, and detected by staining with an anti-rabbit secondary AF750 (ab175735,

Abcam) antibody for an additional 20 mins at RT. Another washing step was performed and

staining with anti-NLRP3-AF700 (768319, R&D Systems) and anti-ASC-PE (HASC-71, Biolegend) was

carried out for 20 minutes at RT. Finally, cells were washed with BD Perm/Wash.

We also studied the expression of the inflammasome components in NK cells and monocytes from

kidney-injured patients, before and after being subjected to kidney transplantation. PBMCs from

these patients were stained with LIVE/DEAD AQUA viability (Invitrogen), and then, with anti-CD56-

FITC (B159, Becton Dickinson), anti-CD3-PE-Cy7 (SK7, Becton Dickinson), anti-CD4-BV605 (RPA-T4,

Becton Dickinson), anti-HLA-DR-PE-Da594 (L243, Biolegend), and anti-CD14-APC-H7 (M5E2, Becton

Dickinson) antibodies. Cells were subsequently fixed and permeabilized with

Fixation/Permeabilization Solution (Becton Dickinson) and intracellularly stained with anti-NLRP1-

AF647 (vwr), anti-NLRP3-AF700 (768319, R&D Systems) and anti-ASC-PE (HASC-71, Biolegend).

Samples were acquired on a CELESTA flow cytometer, and data was analyzed using FlowJo V10

software. Gating was performed according to the different FMO controls.

Samples were acquired on a CELESTA flow cytometer, and data was analyzed using FlowJo V10
software. Gating was performed according to the different FMO controls.

Western blot assays

Western blotting was performed using standard methods. Total PMBCs or NK cell extracted
proteins were used for standard protein electrophoresis and Western blot assays. Gel
electrophoresis was performed using 4–20% Mini-PROTEAN® TGX Stain-Free™ Protein Gels, Biorad
at 200V in Tris-Glycine-SDS buffer for 40 mins. Protein transfer was made using a TurboTransfer
(Biorad, Hercules, CA, USA) at 25V for 7 mins. After transferring the proteins to 0.45 μM
nitrocellulose membranes (Biorad, Hercules, CA, USA), these were incubated for 1 hour in BSA 5%
in PBS-Tween20 0.05% and then overnight at 4ºC with primary antibodies at 1:1,000 dilution. Then
washed twice with PBS-Tween20 and incubated with the corresponding secondary antibody
coupled to horseradish peroxidase diluted 1:10,000 for 1h at RT. Protein loading was checked
using stain-free gel activation and tubulin protein amount. Stripping was not used.

The following primary antibodies were used: NLRP1 (ABclonal, A16212), NLRP3 (ABclonal, A5652),
Caspase-1 (Cell signaling, 3866S), Gasdermin D (Santacruz, sc-393581), GADPH (Cell signaling,
5174S). Anti-rabbit or anti-mouse IgG secondary antibody from Calbiochem was used.

Analysis of cytokine and immune factor secretion

Concentrations of the cytokines and molecules GM-CSF, IFN-γ, IL-1β, IL-6, IL-10, TNF-α, IL-18, IFN-
α, and ICAM-1 were quantified in 50 μL of supernatant from stimulated NK cells using a bead-
based multiplex immunoassay (ProcartaPlex; Invitrogen) according to the manufacturer's
recommendations. Measurements were performed using a Luminex Intelliflex instrument
(ThermoFisher Scientific) and analyzed using a standard curve for each cytokine.

LDH cytotoxicity assay

Cell death was measured by lactate dehydrogenase (LDH) release in the supernatant following the
manufacturer's instructions (abcam). In these experiments, cells were plated and stimulated in
RPMI 1640 without FBS to not interfere with the assay.

Quantification and statistical analysis

Statistical analyses were performed with Prism software, version 6.0 (GraphPad). A P value <0.05

was considered significant. The statistical details for the different experiments can be found in

each figure legend.

REFERENCES

1. Zitti, B. and Y.T. Bryceson, *Natural killer cells in inflammation and autoimmunity.* Cytokine
Growth Factor Rev, 2018. **42**: p. 37-46.
2. Horowitz, A., et al., *Genetic and environmental determinants of human NK cell diversity*
revealed by mass cytometry. Sci Transl Med, 2013. **5**(208): p. 208ra145.
3. Dogra, P., et al., *Tissue Determinants of Human NK Cell Development, Function, and*
Residence. Cell, 2020. **180**(4): p. 749-763 e13.
4. Moretta, L., *Dissecting CD56dim human NK cells.* Blood, 2010. **116**(19): p. 3689-91.
5. Vivier, E., et al., *Functions of natural killer cells.* Nat Immunol, 2008. **9**(5): p. 503-10.
6. Lanier, L.L., *Up on the tightrope: natural killer cell activation and inhibition.* Nat Immunol,
2008. **9**(5): p. 495-502.
7. Fauriat, C., et al., *Regulation of human NK-cell cytokine and chemokine production by*
target cell recognition. Blood, 2010. **115**(11): p. 2167-76.
8. Erlich, Z., et al., *Macrophages, rather than DCs, are responsible for inflammasome activity*
in the GM-CSF BMDC model. Nat Immunol, 2019. **20**(4): p. 397-406.
9. Migita, K., et al., *Serum amyloid A triggers the mosodium urate -mediated mature*
interleukin-1beta production from human synovial fibroblasts. Arthritis Res Ther, 2012.
14(3): p. R119.
10. Lei-Leston, A.C., A.G. Murphy, and K.J. Maloy, *Epithelial Cell Inflammasomes in Intestinal*
Immunity and Inflammation. Front Immunol, 2017. **8**: p. 1168.
11. Talley, S., D.J. Rademacher, and E.M. Campbell, *Inflammasome activation occurs in CD4(+)*
and CD8(+) T cells during graft-versus-host disease. Cell Death Dis, 2023. **14**(9): p. 632.
12. Zheng, D., T. Liwinski, and E. Elinav, *Inflammasome activation and regulation: toward a*
better understanding of complex mechanisms. Cell Discov, 2020. **6**: p. 36.
13. Swanson, K.V., M. Deng, and J.P. Ting, *The NLRP3 inflammasome: molecular activation and*
regulation to therapeutics. Nat Rev Immunol, 2019. **19**(8): p. 477-489.
14. Dai, Z., et al., *Gasdermin D-mediated pyroptosis: mechanisms, diseases, and inhibitors.*
Front Immunol, 2023. **14**: p. 1178662.
15. Costa Franco, M.M., et al., *Brucella abortus Triggers a cGAS-Independent STING Pathway*
To Induce Host Protection That Involves Guanylate-Binding Proteins and Inflammasome
Activation. J Immunol, 2018. **200**(2): p. 607-622.
16. Gomes, M.T.R., et al., *Guanylate-binding proteins at the crossroad of noncanonical*
inflammasome activation during bacterial infections. J Leukoc Biol, 2019. **106**(3): p. 553-
562.
17. Zhong, F.L., et al., *Human DPP9 represses NLRP1 inflammasome and protects against*
autoinflammatory diseases via both peptidase activity and FIIND domain binding. J Biol
Chem, 2018. **293**(49): p. 18864-18878.
18. Okondo, M.C., et al., *DPP8 and DPP9 inhibition induces pro-caspase-1-dependent*
monocyte and macrophage pyroptosis. Nat Chem Biol, 2017. **13**(1): p. 46-53.
19. Sester, D.P., et al., *A novel flow cytometric method to assess inflammasome formation.* J
Immunol, 2015. **194**(1): p. 455-62.

20.   Shi, J., et al., *Cleavage of GSDMD by inflammatory caspases determines pyroptotic cell
death.* Nature, 2015. **526**(7575): p. 660-5.
21.   Qiu, F., et al., *Activation of cytokine-producing and antitumor activities of natural killer
cells and macrophages by engagement of Toll-like and NOD-like receptors.* Innate Immun,
2011. **17**(4): p. 375-87.
22.   Broz, P. and V.M. Dixit, *Inflammasomes: mechanism of assembly, regulation and
signalling.* Nat Rev Immunol, 2016. **16**(7): p. 407-20.
23.   Rozario, P., et al., *Mechanistic basis for potassium efflux-driven activation of the human
NLRP1 inflammasome.* Proc Natl Acad Sci U S A, 2024. **121**(2): p. e2309579121.
24.   Henderson, P.J., *Ion transport by energy-conserving biological membranes.* Annu Rev
Microbiol, 1971. **25**: p. 393-428.
25.   Taabazuing, C.Y., A.R. Griswold, and D.A. Bachovchin, *The NLRP1 and CARD8
inflammasomes.* Immunol Rev, 2020. **297**(1): p. 13-25.
26.   Volchuk, A., et al., *Indirect regulation of HMGB1 release by gasdermin D.* Nat Commun,
2020. **11**(1): p. 4561.
27.   Michel, T., et al., *Human CD56bright NK Cells: An Update.* J Immunol, 2016. **196**(7): p.
2923-31.
28.   Osuka, A., et al., *A protective role for inflammasome activation following injury.* Shock,
2012. **37**(1): p. 47-55.
29.   Stoecklein, V.M., et al., *Radiation exposure induces inflammasome pathway activation in
immune cells.* J Immunol, 2015. **194**(3): p. 1178-89.
30.   Fong, J.J., et al., *Siglec-7 engagement by GBS beta-protein suppresses pyroptotic cell death
of natural killer cells.* Proc Natl Acad Sci U S A, 2018. **115**(41): p. 10410-10415.
31.   Souza-Fonseca-Guimaraes, F., et al., *Toll-like receptors expression and interferon-gamma
production by NK cells in human sepsis.* Crit Care, 2012. **16**(5): p. R206.
32.   Zheng, Z., et al., *NLRP3 associated with chronic kidney disease progression after
ischemia/reperfusion-induced acute kidney injury.* Cell Death Discov, 2021. **7**(1): p. 324.
33.   Granata, S., et al., *NLRP3 inflammasome activation in dialyzed chronic kidney disease
patients.* PLoS One, 2015. **10**(3): p. e0122272.
34.   Franchon Marques Tejada, N., et al., *AIM2 as a putative target in acute kidney graft
rejection.* Frontiers in Immunology, 2022. **13**.
35.   Dessing, M.C., et al., *Donor and recipient genetic variants in NLRP3 associate with early
acute rejection following kidney transplantation.* Sci Rep, 2016. **6**: p. 36315.
36.   Suarez-Rivero, J.M., et al., *Does NLRP1 inflammasome activation in immune cells in kidney
transplantation relate with donor organ age?* Antioxid Redox Signal, 2024. doi:
10.1089/ars.2024.0588
37.   Granata, S., et al., *Inflammasome pathway in kidney transplantation.* Front Med
(Lausanne), 2023. **10**: p. 1303110.
38.   Zhu, L., et al., *Changes of NK cell subsets with time post-transplant in peripheral blood of
renal transplant recipients.* Transpl Immunol, 2018. **49**: p. 59-71.
39.   Pontrelli, P., et al., *The Role of Natural Killer Cells in the Immune Response in Kidney
Transplantation.* Front Immunol, 2020. **11**: p. 1454.

Figure 1

Figure 2

Figure 3

Figure 4