Assessment of the level of pollution based on soil and *Tilia × europaea* leaves

Edina Simon[a,b]. Ismail Haziq Bin[a]. Bianka Sipos[a,b]. Vanda Éva Abriha-Molnár[a,b]. Dávid Abriha[c,] Dávid Tőzsér[a,d], Hernandez Rafael Boluda[e]. Pérez Luis Roca[e]

[a]Department of Ecology, University of Debrecen, Egyetem tér 1. H-4032 Debrecen, Hungary
[b]HUN-REN–UD Anthropocene Ecology Research Group, University of Debrecen, Egyetem tér 1. H-4032 Debrecen, Hungary
[c]Department of Physical Geography and Geoinformatics, University of Debrecen, Egyetem tér 1. H-4032 Debrecen, Hungary
[d]Circular Economy Analysis Center, Hungarian University of Agriculture and Life Sciences, Páter Károly u. 1, H□2100 Gödöllő, Hungary
[e]Facultat de Farmàcia, Dpt. Biologia Vegetal, Universitat de València

Corresponding author: E. Simon (edina.simon@gmail.com)

Abstract

Urban soil and plant leaves are widely used as ecological indicators to assess the effect of urbanization. This study aimed to evaluate the impact of urbanization based on the elemental concentration of soil and leaves of *Tilia × europaea* from urban, suburban, and rural areas along an urbanization gradient in Debrecen, Hungary. Using the ICP-OES technique, we measured the concentration of Ba, Cd, Co, Cr, Cu, Ni, Pb, Sr, and Zn and based on the measured concentration, bioaccumulation factor (BAF) was calculated. The highest concentration of all elements was found in soil samples from urban areas, with an increasing tendency along the urbanization gradient. A significant difference was found along the urbanization gradient only for Cr based on the plant leaves. A high BAF value was found for Cu and Sr. Our results suggest that urbanization has a remarkable effect on the elemental concentration of soil, which is a perfect ecological indicator. At the same time, we concluded that the *T. × europaea* was not sensitive enough to indicate the effect of urbanization.

Keywords: tree leaves, metals, bioindicators, Hungary

Introduction

Soil and air pollution by metals is one of the most critical environmental problems worldwide (Hu et al. 2013). Among anthropogenic activities, urbanization is one of the primary sources of polluted soil with heavy metals (Mirales et al. 2012). Urbanization is progressing rapidly between 2000 and 2030, and the urban population is anticipated to grow an average of 2.3% per year in the developing world (Chen 2007). The effects of urbanization on environmental pollution are different in countries and regions with various degrees of development (Son and Hu 2018). Earlier studies reported that urbanization directly and indirectly affects soil as an essential resource; accumulation of heavy metals can modify the quality of soil and agricultural products, and thus, they have adverse effects on humans, animals, and the ecosystem health (Nagajyoti et al. 2010, Ihenetu et al. 2024)

Trees play an essential role in urban vegetated areas (Ma et al. 2022, Wang and Zhang 2022). Plant leaves can accumulate air pollutants in high concentrations; air pollutant particles can also be deposited in the leaves of plants due to their stomatal openings and waxy cuticles (Bibi et al. 2023, Molnár et al. 2020, Simon et al. 2021). The genus Tilia is generally used as biomonitors of trace elements to assess the effect of urbanization and industrialization on terrestrial environments (Zeb et al. 2022, Andrianjara et al. 2024).

Therefore, *Tilia × europaea* was chosen for our study, a typically ornamental tree in European countries (Ţenche-Constantinescu et al. 2015). Our study aimed to investigate elemental concentration in soil and *Tilia × europaea* plant leaves along an urbanization gradient in Debrecen. Hungary. Our hypotheses are as follows: (i) there are significant differences in pollutant levels along the urbanization gradient (urban, suburban, and rural areas), (ii) the highest metal concentration in soil and leaves is in the urban area, while the lowest concentration is in the rural area, (iii) significant correlation is found between the soil and leaves metal concentrations.

Material and methods

Study area
Debrecen is the second largest city in Hungary, with a population of approximately 200.000 as of 2019. The area allocated for the city is 461 km^2, which results in a population density of 436/km^2. Along an urbanization gradient in Debrecen city, three different studied areas were chosen (Fig.1). The rural area was situated near Nagyerdő, a protected area for nature conservation. The suburban area is intermediate between the rural and urban areas regarding urban density. It is located adjacent to a road with apartments on one side, and the side is a forest. Lastly, the urban study area was at the center of the urban community to maximize the observable effect of urban pollution, and it was surrounded by housing and offices at the center of Debrecen. Soil samples (N = 15) were collected along an urbanization gradient representing rural, suburban, and urban areas. At each sampling area, five soil samples were collected. All samples were collected from 0–20 cm depth with a small hand spade. Samples were placed into poly bags and kept at -21 °C in the refrigerator.

Figure 1. Studied areas for Debrecen along an urbanization gradient. Notations: The red circle indicates the study areas

During the plant sampling, 10-15 leaves were collected from 3 individuals of the same tree species in the same location. Additionally, soil samples were collected within a 1-meter radius of the tree base to calculate the bioaccumulation later in this research. Trees showing any form of disease should be avoided, as this may alter the investigation results.

Sample preparation
The laboratory process started with a surface wash to remove any surface pollutants from the leaf samples. After washing, the leaf samples were put to dry in an oven set at 70 ^{o}C for 90 minutes. Soil samples were dried at room temperature, and after that stones, plant roots, and residues were removed with plastic tweezers. Samples were sieved in a 2-mm plastic sieve.

Then, the sample process was the same for soil and leaf samples. The samples process was published by Simon et al. (2014, 2016, 2021), Bibi et al. (2023a, b), and Molnár et al. (2020) (Fig. 2). For the elemental analysis, ICP-OES was used. Peach leaves (1547) and soil (SQC001) CRM were used, and the recoveries were within 10% of the certified values for the elements (Simon et al. 2013, 2014, 2016, 2021). The concentration of Ba, Cd, Co, Cr, Cu, Ni, Pb, Sr, and Zn was analyzed during the analysis.

Figure 2. Sample preparations and elemental analysis for soil and leaves

Bioaccumulation factor (BAF)

To study the accumulation of elements in leaves of *T. × europaea*, bioaccumulation factor (BAF) was used. BAF is the ratio of elemental concentration in leaves (C_{plant}) and elemental concentration in soil (C_{soil}) (Ivanciuc et al. 2006, van Gestel et al. 2011):

$$BAF = C_{plant} / C_{soil}$$

Statistical analyses

Statistical analyses were conducted with SPSS Statistics 20 (IBM) statistical software. The normality of the distribution was tested using the Shapiro–Wilk test. The homogeneity of variances was tested using Levene's test. The differences among samples were tested with the analysis of variance (ANOVA) for each variable. Tukey's test was performed for pairwise comparison between the groups. Canonical discriminant analysis (CDA) was used to reduce dimensions and to identify those variables that most efficiently discriminated the study area as the dependent variable (De Sá 2007). The correlation between the elemental concentration of soil and leaves was analyzed with the Pearson correlation coefficient (Zar 1996).

Results

There was a clear separation among the studied areas based on the elemental concentration of soil according to the CDA (Fig. 3). The canonical variables were significant ($p < 0.001$) in the first and second discriminant functions. The percentage of variance was 87.0 in the first, while 13.0 was in the second discriminant function. There was significantly higher elemental soil concentration in the urban area than in the rural area (Table 1).

Figure 3. Scatter plot of canonical discriminant analysis (CDA) based on the elemental concentration of soil among the studied areas

Table 1. Summary statistics of elemental concentration of soil (mg kg^{-1}, mean $\pm$ SE).

Elements	Urbanization gradient			F	p
	rural	suburban	urban		
Ba	14.2 ± 1.0	67.7 ± 16.2	97.1 ± 11.0	13.825	0.001
Cd	0.3 ± 0.1	0.8 ± 0.1	1.8 ± 0.2	55.623	<0.001
Co	1.3 ± 0.1	2.6 ± 0.1	6.4 ± 0.4	96.986	<0.001
Cr	3.1 ± 0.3	7.3 ± 0.7	28.0 ± 6.3	13.081	0.001
Cu	3.7 ± 0.3	13.8 ± 2.5	22.3 ± 2.7	19.031	<0.001
Ni	18.9 ± 4.5	28.6 ± 1.5	39.8 ± 4.2	8.229	0.006
Pb	5.5 ± 2.3	20.9 ±5.5	24.5 ± 3.9	5.971	0.016
Sr	5.5 ± 0.7	25.2 ± 6.8	32.6 ± 4.3	8.975	0.004
Zn	19.1 ± 1.2	61.5 ± 8.1	83.9 ± 13.7	12.735	0.001

Similar to the soil, separation was found among the studied areas based on the elemental concentration of *Tilia × europaea* according to the CDA (Fig. 4). The canonical variables were not significant ($p > 0.05$) in the first and second discriminant functions. The percentage of variance was 99.2 in the first, while 0.8 was in the second discriminant function. A significant difference was found only in the case of the Cr concentration of *Tilia × europaea*. Higher Cr concentration was found in the rural and suburban areas than in urban areas (Table 2).

Figure 4. Scatter plot of canonical discriminant analysis (CDA) based on the elemental concentration of *Tilia × europaea* leaves among the studied areas

Table 2. Summary statistics of elemental concentration of *Tilia × europaea* leaves (mg kg^{-1}, mean ± SE). The Cd, Co, Cr, and Pb concentrations were below the detection limit.

| Elements | Urbanization gradient | | | F | p |
	rural	suburban	urban		
Ba	12.9 ± 4.2	8.1 ± 0.6	5.6 ± 2.5	1.663	0.266
Cr	0.3 ± 0.1	0.3 ± 0.1	0.2 ± 0.1	7.426	0.024
Cu	7.7 ± 0.9	6.2 ± 0.6	4.9 ± 1.1	2.709	0.145
Ni	1.5 ± 0.5	0.6 ± 0.1	0.5 ± 0.1	3.844	0.084
Sr	28.8 ± 4.6	41. 9 ± 1.7	32.1 ± 5.8	2.456	0.166
Zn	18.5 ± 1.4	14.4 ± 0.4	17.5 ±0.9	4.85	0.056

In all studied areas, we did not find any significant correlation between the elemental concentration of soil and leaves (rural: r = 0.429, p = 0.397; suburban: r = 0.486, p = 0.329; urban: r = 0.371, p = 0.468). The BAF values were higher than 1 for Cu in the rural and for Sr in all studied areas (Fig. 5 and 6). The BAF values of Cu and Sr indicated that trees could accumulate these elements from soil indirectly or directly.

Figure 5. Value of bioaccumulation factor (BAF) for Cu (mean ± SE) along the urbanization gradient

Figure 6. Value of bioaccumulation factor (BAF) for Sr (mean ± SE) along the urbanization gradient based on the elemental concentration of soil and

Discussion

As hypothesized, we found differences based on the elemental concentration of soil along the urbanization gradient; significantly higher concentrations were found in urban areas than in rural ones. Thus, it was indicated that urbanization had a remarkable effect on the metal concentration of soil. Contrary to our hypothesis, the elemental concentration of *T.* × *europaea* did not differ significantly along the urbanization gradient, only the concentration of Cr, which is a typically anthropogenic element. At the same time, our results also indicated that *T.* × *europaea* was not sensitive enough to reveal the effect of urbanization.

Similar to our results, Hu et al. (2013) also demonstrated the effect of urbanization on the soil elemental concentration change, and they found the highest pollution level and potential ecological risk of surface soil polluted with metals in urban soil. Urbanization may cause an increasing risk of soil pollutants through waste disposal and acid deposition derived from urban air pollution (Chen 2007). Tešic et al. (2022) also found significantly higher Zn, Cu, Cd, Cr, and Ni in the soil of urban areas in Belgrade than in the rural. Simon et al. (2016) also reported that soil was a suitable bioindicator for monitoring the effects of urbanization and anthropogenic activities on terrestrial ecosystems because they found significant differences among areas based on the concentration of Ba, Cu, Mn, Sr, and Zn in soil. Simon et al. (2013) also confirmed that the elemental analysis of soil samples is one of the best ways to study the effects of urbanization. Bibi et al. (2023) also showed higher concentrations of Cu, Pb, Sr, and Zn were
in the urban and suburban areas than in the rural areas in Wien, Austria.

Our study by Molnár et al. (2020) demonstrated that *T.* × *europaea* is sensitive to air pollutants based on the air pollution tolerance index (APTI). Serbula et al. (2013) showed that pine needles responded more intensively to pollution than *Tilia*. Serbula et al. (2013) also indicated, based on the bioaccumulation and translocation index, that the high concentrations of Cu and Pb in aerial parts of pine and linden are mainly a consequence of airborne pollution, which explains that there was no correlation between the elemental concentration of soil and leaves in our study. Similarly to our finding, Țenche-Constantinescu et al. (2015) also demonstrated that *Tilia* sp. was very resistant to biotic and abiotic stress because they found a correlation only between traffic intensity and the concentration of Pb in leaves. The study of *Tilia* leaf showed a different capacity for accumulating potentially toxic elements in various locations by Mitrović et al. (2021) in urban areas with varying pollution levels. Jouraeva et al. (2002) studied the accumulation potential of PAHs and metals on the leaves of *Tilia×euchlora* and *Pyrus calleryana*, and they reported that leaves of *T.* ×*euchlora* accumulated higher concentrations of air pollutants than *P. calleryana*. Aničić et al. (2011) studied the metal concentration in the *Aesculus hippocastanum* leaves and *Tilia* spp. Similar to our finding, they also demonstrated that *Tilia* spp. was not so sensitive to assessing Pb and Cu atmospheric pollution (Aničić et al. 2011). Urošević et al. (2019) studied the metal concentration of different tree leaves, and they did not recommend *Tilia* sp. as a valuable bioindicator for air pollutants. In the case of plant species, the uptake of the elements may be accompanied by a species-specific pattern of physiological reaction (Drzewiecka et al. 2019).

Conclusion

Our study demonstrated differences in the elemental concentration of soil along the urbanization gradient. Slight difference was found in elemental concentrations of *T.* ×

europaea only in the case of Cr along the urbanization gradient. Based on the accumulation rate, *T.* × *europaea* could accumulate the Cu and Sr from soil directly or indirectly. Thus, we concluded that urbanization has a remarkable effect on the elemental concentration of soil, which is a perfect ecological indicator. At the same time, we concluded that the *T.* × *europaea* was not sensitive enough to indicate the effect of urbanization.

Acknowledgements
Research supported by 2021-1.2.4-TÉT-2021-00049 project and the University of Debrecen Program for Scientific Publication.

References
Andrianjara I, Cabassa C, Lata JC, Hansart A, Raynaud X, Renard M, Nold F, Genet G, Planchais S 2024. Characterization of stress indicators in *Tilia cordata* Mill. as early and long-term stress markers for water availability and trace element contamination in urban environments. Ecological Indicators 158, 111296. https://doi.org/10.1016/j.ecolind.2023.111296

Aničić M, Spasić T, Tomašević M, Rajšić S, Tasić M 2011. Trace elements accumulation and temporal trends in leaves of urban deciduous trees (*Aesculus hippocastanum* and *Tilia* spp.). Ecological Indicators 11, 824-830. https://doi.org/10.1016/j.ecolind.2010.10.009.

Aničić Urošević M, Jovanović G, Stević N, Deljanin I, Nikolić M, Tomašević M, Samson R 2019. Leaves of common urban tree species (*Aesculus hippocastanum*, *Acer platanoides*, *Betula pendula* and *Tilia cordata*) as a measure of particle and particle-bound pollution: a 4-year study. Air Quality, Atmosphere & Health 12, 1081–1090. https://doi.org/10.1007/s11869-019-00724-6

Bibi D, Tőzsér D, Sipos B, Molnár VÉ, Tóthmérész B, Simon E 2023b. Complex study of air pollution based on tree species in Vienna. Air Quality, Atmosphere & Health. https://doi.org/10.1007/s11869-023-01452-8

Bibi D, Tőzsér D, Sipos B, Tóthmérész B, Simon E 2023a. Heavy metal pollution of soil in Vienna, Austria. Water Air Soil Pollution 234, 232. https://doi.org/10.1007/s11270-023-06244-5

Chen J 2007. Rapid urbanization in China: A real challenge to soil protection and food security. Catena 69, 1-15. https://doi.org/10.1016/j.catena.2006.04.019

De Sá JPM 2007. Applied Statistics Using SPSS. STATISTICA. MATLAB and R. 2nd ed.; Springer: Berlin/Heidelberg. Germany. ISBN 978-3-540-71971-7.

Drzewiecka K, Piechalak A, Goliński P, Gąsecka M, Magdziak Z, Szostek M, Budzyńska S, Niedzielski P, Mlecze M 2019. Differences of *Acer platanoides* L. and *Tilia cordata* Mill. Response patterns/survival strategies during cultivation in extremely polluted mining sludge – A pot trial. Chemosphere 229, 589-601. https://doi.org/10.1016/j.chemosphere.2019.05.051

Drzewiecka K, Piechalak A, Goliński P, Gąsecka M, Magdziak Z, Szostek M, Budzyńska S, Niedzielski P, Mlecze M 2019. Differences of Acer platanoides L. and Tilia cordata Mill. Response patterns/survival strategies during cultivation in extremely polluted mining sludge – A pot trial. Chemosphere 229, 589-601. https://doi.org/10.1016/j.chemosphere.2019.05.051

Hu Y, Liu X, Bai J, Shih K, Zeng EY, Cheng H 2013.Assessing heavy metal pollution in the surface soils of a region that had undergone three decades of intense industrialization and urbanization. Environmental Science and Pollution Research 20, 6150–6159. https://doi.org/10.1007/s11356-013-1668-z

Ihenetu SC, Li G, Mo Y, Jacques KJ 2024. Impacts of microplastics and urbanization on soil health: An urgent concern for sustainable development. Green Analytical Chemistry 8, 100095. https://doi.org/10.1016/j.greeac.2024.100095

Ivanciuc T, Ivanciuc O, Klein DJ 2006. Modeling the bioconcentration factors andbioaccumulation factors of polychlorinated biphenyls with posetic quantitativesuper-structure/activity relationships (QSSAR). Molecular Diversity 10, 133–145.

Jouraeva VA, Johnson DL, Hassett JP, Nowak DJ 2002. Differences in accumulation of PAHs and metals on the leaves of Tilia×euchlora and Pyrus calleryana. Environmental Pollution 120, 331-338. https://doi.org/10.1016/S0269-7491(02)00121-5

Ma Z, Zhang P, Hu N, Wang G, Dong Y, Guo Y, Wang C, Fu Y, Ren Z 2022. Understanding the drivers of woody plant diversity in urban parks in a snow climate city of China. Journal of Forestry Research 34, 1021-1032. https://doi.org/10.1007/s11676-022-01535-9

Mireles F, Davila JI, Pinedo JL, Reyes E, Speakman RJ, Glascock MD 2012. Assessing urban soil pollution in the cities of Zacatecas and Guadalupe, Mexico by instrumental neutron activation analysis. Microchemical Journal 103, 158–164.https://doi.org/10.1016/j.microc.2012.02.00

Mitrović M, Blanusa T, Pavlović M, Pavlović D, Kostić O, Perović V, Jarić S, Pavlović P 2021. Using Fractionation Profile of Potentially Toxic Elements in Soils to Investigate Their Accumulation in Tilia sp. Leaves in Urban Areas with Different Pollution Levels. Sustainability 13, 9784. https://doi.org/10.3390/su13179784

Molnár VÉ, Simon E, Ninsawat S, Tóthmérész B, Szabó Sz 2020. Pollution assessment based on element concentration of tree leaves and topsoil in Ayutthaya Province, Thailand. International Journal of Environmental Research and Public Health 17, 5165.

Nagajyoti PC, Lee KD, Sreekanth TVM 2010. Heavy metals, occurrence and toxicity for plants: a review. Environmental Chemistry Letter 8, 199–216. https://doi.org/10.1007/s10311-010-0297-8

Serbula SM, Kalinovic TS, Ilic AA, Kalinovic JV, Steharnik MM 2013. Assessment of Airborne Heavy Metal Pollution Using Pinus spp. and Tilia spp. Aerosol and Air Quality Research 13, 563-573. https://doi.org/10.4209/aaqr.2012.06.0153.

Simon E, Baranyai E, Braun M, Cserháti C, Fábián I, Tóthmérész B 2014. Elemental concentrations in deposited dust on leaves along an urbanization gradient. Science of the Total Environment 490, 514–520. https://doi.org/10.1016/j.scitotenv.2014.05.028.

Simon E, Harangi S, Baranyai E, Braun M, Fábián I, Mizser Sz, Nagy L, Tóthmérész B 2016. Distribution of toxic elements between biotic and abiotic components of terrestrial ecosystem along an urbanization gradient: Soil. leaf litter and ground beetles. Ecological Indicators 60, 258–264. http://doi.org/10.1016/j.ecolind.2015.06.045.

Simon E, Molnár VÉ, Lajtos D, Bibi D, Tóthmérész B, Szabó S 2021. Usefulness of Tree Species as Urban Health Indicators. Plants 10, 2797. https://doi.org/10.3390/plants10122797.

Simon E, Vidic A, Braun M, Fábián I, Tóthmérész B 2013. Trace element concentrations in soils along urbanization gradients in the city of Vienna. Austria. Environmental Science and Pollution Research 20, 917–924. https://doi.org/10.1007/s11356-012-1091-x.

Song M, Hu C 2018.A coupling relationship between the eco-environment carrying capacity and new-type urbanization: a case study of the Wuhan metropolitan area in China. Sustainability 10, 4671. https://doi.org/10.3390/su10124671

Țenche-Constantinescu AM, Madoşa E, Chira D, Hernea C, Țenche-Constantinescu RV, Lalescu D, Borlea GF 2015. Tilia spp.- Urban Trees for future. Notulae Botanicae Horti Agrobotanici Cluj-Napoca. 43, 259-264. DOI:10.15835/nbha4319794

Tešić M, Stojanović N, Knežević M, Đunisijević-Bojović D, Petrović J, Pavlović P 2022. The Impact of the Degree of Urbanization on Spatial Distribution, Sources and Levels of Heavy Metals Pollution in Urban Soils—A Case Study of the City of Belgrade (Serbia). Sustainability 14, 13126. https://doi.org/10.3390/su142013126

Van Gestel CAM, Ortiz MD, Borgman E, Verweij RA 2011. The bioaccumulation of Molybdenum in the earthworm Eisenia andrei: influence of soil properties and ageing. Chemosphere 82, 1614–1619.

Wang S, Zhang H 2022. Tree composition and diversity in relation to urban park history in Hong Kong, China. Urban Forestry & Urban Greening 67, 127430. https://doi.org/10.1016/j.ufug.2021.127430

Zar IH 1996. Biostatistical Analysis. Prentice Hall, N.J.

Zeb J, Tahir H, Othman A, Habeebullah TM, Saygal A, Assaggaf HM, Ahmed OB, Sultan M, Mohiuddin S, Masood SS, Mirza AZ, Hajira B 2022. Geo-environmental approach to assess heavy metals around auto-body refinishing shops using biomonitors. Heliyon 8, e08809. https://doi.org/10.1016/j.heliyon.2022.e08809

rural   suburban   urban