1   Running title: (50 characters) Warm temperature inhibits *Rickettsiella*-induced CI

5   **Title:** Warm temperature inhibits cytoplasmic incompatibility induced by endosymbiotic
6   *Rickettsiella* in a spider host

8   Jordyn D. Proctor[1], Virginija Mackevicius-Dubickaja[2], Yuval Gottlieb[2], Jennifer A. White[1]*
9   [1] *S-225 Agricultural Science Center N, Department of Entomology, University of Kentucky,*
10   *Lexington, Kentucky.*
11   [2]*Koret School of Veterinary Medicine, The Robert H. Smith Faculty of Agriculture, Food and*
12   *Environment, The Hebrew University, Rehovot, Israel.*
13
14   *To whom correspondence should be addressed.
15   email: jenawhite@uky.edu

16
17

Data availability statement: All data associated with this manuscript are available in the
supplemental material, Supplemental Document 1 and Supplemental Document 2.

Funding statement: This material is based upon work supported by the National Science
Foundation under Grant No. 1953223 and the Binational, USA-Israel, Science Foundation (BSF)
under Grant No. 201697 to JAW and YG; and the National Institute of Food and Agriculture, U.S.
Department of Agriculture (Hatch No. 1020740).

Conflict of interest disclosure: The authors declare no conflicts of interest.

Author contributions: All authors designed the experiments; JDP and VMD conducted the
experiments; JAW conducted statistical analyses and generated figures; All authors contributed
to the manuscript and approved the final version.

ABSTRACT

Bacterial endosymbionts manipulate reproduction in arthropods to increase prevalence in the host
population. One such manipulation is cytoplasmic incompatibility (CI), wherein the bacteria sabotage
sperm in infected males to reduce hatch rate when mated with uninfected females, but zygotes are
"rescued" when that male mates with an infected female. In the spider *Mermessus fradeorum*
(Linyphiidae), *Rickettsiella* symbionts cause variable levels of CI. We hypothesized that temperature
affects CI strength and rescue in *M. fradeorum,* potentially mediated by bacterial titer. We reared
Rickettsiella-infected spiders in two temperature conditions (26°C vs 20°C) and tested CI induction in
males and CI rescue in females. In incompatible crosses between infected males and uninfected females,
hatch rate from warm males was doubled (Mean±S.E. = 0.687±0.052) relative to cool males
(0.348±0.046), indicating that CI induction is weaker in warm males. In rescue crosses between infected
females and infected males, female rearing temperature had a marginal effect on CI rescue, but hatch
rate remained high for both warm (0.960±0.023) and cool females (0.994±0.004). Bacterial titer as
measured by qPCR was lower in warm than cool spiders, particularly in females, suggesting that
bacterial titer may play a role in causing the temperature-mediated changes in CI.

Keywords

bacterial endosymbionts, bacterial titer, climate change, cytoplasmic incompatibility,
reproductive manipulation, symbiosis, temperature

INTRODUCTION

Maternally-inherited bacterial symbionts are common in arthropods, often manipulating host
reproduction to increase bacterial prevalence within the host population. Cytoplasmic
incompatibility (CI) is the most common and well-studied manipulation, sabotaging matings
between infected males and uninfected females, preventing the production of uninfected
offspring, and increasing the relative proportion of infected offspring in the population over
time (Shropshire et al. 2020). Numerous bacterial clades have been shown to cause CI,
including *Wolbachia*, *Cardinium*, *Spiroplasma*, and *Rickettsiella* (Hunter et al. 2003, Rosenwald
et al. 2020, Shropshire et al. 2020, Pollmann et al. 2022). The mechanistic underpinnings of CI
are only well characterized for *Wolbachia* (Shropshire et al. 2020), but genomic and phenotypic
studies with other symbionts suggest novel modes of action may be present (Poinsot et al.
2003, Penz et al. 2012, Pollmann et al. 2022). Consequently, caution should be used when
extrapolating inferences from *Wolbachia* CI studies to other bacterial symbionts.
59
There is growing evidence that symbiont-induced host phenotypes, such as CI, can be thermally
sensitive but potentially idiosyncratic (Corbin et al. 2017, Doremus et al. 2019, Shropshire et al.
2020, Chrostek et al. 2021, Corbin et al. 2021, Hague et al. 2022, Jones et al. 2023, Martins et al.
2023). The effect of temperature on *Rickettsiella*-induced CI has not previously been
investigated, but in a non-CI-inducing *Rickettsiella* strain in aphids, temperature affects
Rickettsiella titer, host phenotype, and spread in the host population (Gu et al. PNAS 2023).
More generally, several studies have suggested that thermal conditions might limit the
geographic distribution of symbionts, their phenotypes, and consequently even their host
species (Zhang et al 2019, Ross et al. 2020, Hague et al. 2022). Thus, under global climate

change, it becomes critical to understand thermal effects on arthropod symbionts, because
symbiont-induced phenotypes are likely to change and affect the spread and performance of
both pest and beneficial arthropod hosts.

Here, we test whether temperature affects the strength of *Rickettsiella*-induced CI and bacterial
titer in *Mermessus fradeorum* (Berland, 1932). This linyphiid spider is infected with several
maternally inherited symbionts, of which only *Rickettsiella* has been shown to induce CI (Curry
et al. 2015, Rosenwald et al. 2020). Using laboratory-reared colonies of spiders that were either
uninfected or infected only with *Rickettsiella* (Rosenwald et al. 2020), we reared spiders under
two temperature regimes (26°C or 20°C) and evaluated the ability of infected males to induce CI
and infected females to rescue CI as a function of temperature. We also evaluated bacterial
titer as a function of temperature because the strength of symbiont-induced phenotypes is
often mediated by bacterial quantity (Lopez-Madrigal and Duarte 2019).

EXPERIMENTAL PROCEDURES

Clutches of newly hatched *M. fradeorum* spiderlings infected with *Rickettsiella* were split into
individual 4cm diameter rearing cups with moistened plaster at the bottom for humidity control
(Rosenwald et al. 2020). Half of the spiderlings were randomly assigned to be reared in a 20˚C
environmental chamber and half to a 26˚C chamber. All were maintained on a diet of *Sinella*
curviseta collembola and *Drosophila melanogaster* fruit flies later on (Rosenwald et al. 2020).
Uninfected spiders were treated identically, except all were reared at 20˚C. After spiders
reached maturity, they were randomly assigned to mating crosses in either male CI induction or
female CI rescue experiments. In the CI induction experiment, *Rickettsiella* (R)-infected male
spiders were mated twice, first to uninfected females (the CI cross) and second to *R*-infected
females to verify the fertility of each male spider. Male spiders were maintained under
standard rearing conditions (fed twice a week with *D. melanogaster*) at their assigned
temperature between matings. In the female CI rescue experiment, all spiders were mated only
once. After matings were complete, male spiders were placed in 1.5mL microcentrifuge tubes
and stored at their rearing temperature without food for five to seven days to ensure that gut
contents had been emptied. Specimens were then preserved in 95% ethanol and stored at -
20˚C for further analysis. Female spiders were maintained under standard rearing conditions at
their assigned temperature until they deposited three eggmasses. Then, females were moved
to microcentrifuge tubes and underwent the same food deprivation and preservation sequence
as the males.

To evaluate the hatch rate, we dissected each eggmass after spiderlings emerged, and counted
the remaining unhatched eggs as in Rosenwald et al. (2020). Total hatched and unhatched eggs
per female were used for a series of logistical regression contrasts, using Williams' correction
for overdispersion (Arc v 1.06; Williams 1982). In the CI induction experiment, we first
compared both expected CI treatments, warm and cool R-infected males mated to uninfected
females, relative to control matings between uninfected males and uninfected females, to
ensure that CI was in evidence in the experiment. We then contrasted when these same males
were mated to cool-reared *R*-infected females, relative to uninfected males and *R*-infected

females. This contrast ensured that neither temperature nor *Rickettsiella* infection negatively
affected male fertility. Finally, we directly contrasted the hatch rates of the warm and cool CI
treatments to one another, to evaluate the effect of rearing temperature on CI induction. In the
CI rescue experiment, we first contrasted warm versus cool R-infected females mated to
uninfected males, to ensure rearing temperature didn't directly affect hatch rate. Next, we
contrasted cool R-infected females mated to cool R-infected males, relative to cool uninfected
females mated to cool R-infected males, to ensure that this group of males was capable of
inducing CI. Finally, we directly contrasted hatch rates of the warm and cool CI rescue
treatments to one another, to evaluate the effect of rearing temperature on CI rescue.

We evaluated symbiont titer via qPCR for a subset of 10-15 *R*-infected spiders from each sex
and temperature treatment. For the sake of consistency, all evaluated spiders originated from
the same experiment, the female CI rescue experiment. We placed specimens into 50% PBS
(EtOH: PBS) for 30 min, followed by 75% PBS (EtOH: PBS) for 30 min, and finally, kept in 100%
PBS overnight. We crushed these specimens in cryogenic vials containing solid-glass beads using
FastPrep® Tissue Homogenizer (MP Biomedicals), then extracted DNA using DNeasy Blood &
Tissue Kit (Qiagen) following the manufacturer's protocol. We designed short primers specific
to the *Rickettsiella recA* gene and *M. fradeorum 18S rRNA* gene using Geneious Prime Version
2021.2. Primer sequences, reaction mixtures, cycling conditions, and titer calculation
methodology are provided in Supplementary Document 1.

RESULTS

When we reared *R*-infected spiders at either 26°C or 20°C and then evaluated the strength of CI
induction (Figure 1A), we found that both warm and cool R-infected males caused reduced
hatch rates when mated to uninfected females, relative to control matings between uninfected
males and females (ΔDeviance = 58.95, d.f. = 2, P<0.001). When these same R-infected males
were mated with *R*-infected females, hatch rates were nearly perfect and did not differ from
control crosses between uninfected males and *R*-infected females (ΔDeviance = 2.87, d.f. = 2,
P=0.238), indicating that neither rearing temperature nor *Rickettsiella* infection intrinsically
decreased fertility of males. When compared directly, the hatch rate of incompatible crosses
between uninfected females and warm males was twice as high (0.687±0.052) as with cool
males (0.348±0.046; ΔDeviance = 21.30, d.f. = 1, P<0.001), indicating that warmer rearing
temperatures weakened CI induction.

Figure 1. Proportion hatch of eggs
from crosses between *Rickettsiella*-
infected *Mermessus fradeorum*
spiders that were individually reared
at 26°C (+) or 20°C (+), and
subsequently mated to either
infected (+) or uninfected (-) spiders
reared at 20°C as indicated on the x-
axes. Crosses to evaluate male CI
induction (Panel A) showed a
significantly higher hatch rate
(weaker CI) for offspring of warm (+)
than cool (+) males when mated to
uninfected (-) females. All control
crosses had high hatch rates. Crosses
to evaluate female CI rescue (Panel B)
showed marginally lower hatch rate
for warm (+) than cool (+) females
mated to infected (+) males, but high
hatch rates overall indicate
temperature had relatively little
effect on CI rescue. Incompatible
crosses between infected (+/+) males
and uninfected (-) females were
included as controls to aid
comparison across the two
experiments and validate that
infected males (+) used within this
experiment were capable of inducing
CI.
171

In the CI rescue experiment (Figure 1B), hatch rates were high and equivalent for warm and
cool R-infected females mated to uninfected males (ΔDeviance=0.20, d.f.=1, P=0.654),
indicating that rearing temperatures did not directly affect hatch rate. Cool R-infected females
also exhibited nearly perfect hatch rates when mated to cool R-infected males, much higher
than cool uninfected females mated to cool R-infected males (ΔDeviance=84.49, d.f.=1,
P<0.001). This control contrast establishes both that the R-infected males in this experiment
induced CI, and that cool R-infected females were capable of rescue. When we contrasted
rescue between the two female rearing temperatures, we found that maternal rearing
temperature had a marginally significant effect on hatch rate (ΔDeviance = 3.825, d.f. =1,
P=0.0505), suggesting that CI rescue ability can be weakened in at least some *R*-infected
females. Nevertheless, most females in these rescue crosses exhibited perfect hatch rates,
indicating that the CI rescue ability of *Rickettsiella* usually remains intact regardless of rearing
temperature.

When we evaluated bacterial titer in a subset of warm-reared versus cool-reared spiders
(Figure 2), we found an interactive effect between sex and temperature (Sex $F_{1,44}$ =0.329,
P=0.529, Temperature $F_{1,44}$ =6.026, P=0.018, Sex × Temp interaction $F_{1,44}$ =5.22, P=0.027). Titer
was lower in the warm treatments, but this lower titer was associated with females more than
males, even though temperature effects on symbiont phenotype were more evident in males
than females.

Figure 2. *Rickettsiella* titer in a
subset of warm (26°C) and cool
(20°C) spiders from the CI rescue
experiment. Log transformed titer
(normalized against spider 18S
quantity) was significantly affected
by the interaction between
temperature and sex, primarily
driven by lower titer in warm
females.

DISCUSSION

Warmer rearing temperatures strongly decreased CI induction by *Rickettsiella* in male *M.*
fradeorum spiders, but only mildly decreased CI rescue in female spiders. Thermal sensitivity for
Rickettsiella-induced phenotypes is consistent with results for a different strain of *Rickettsiella*
in an aphid host, which exhibits a different set of host phenotypic effects that don't include CI
(Gu et al. 2023). In the present study, CI strength overall was notably weaker than in prior
studies by our group (Rosenwald et al. 2020), which were conducted at room temperature, but
in a laboratory that was cool throughout the duration of the experiments (J. White, pers. obs.).
Highly variable hatch rates across all CI crosses (Fig 1A), suggest that other factors may
additionally be driving CI strength in this system (e.g., Shropshire et al. 2021). More generally,
bacterial symbionts often show reduced phenotypic penetrance at warmer temperatures
(Martins et al. 2023), although the phenomenon is not universal, and instances where
symbiont-induced phenotypes increase with warmer temperatures also exist (e.g., Corbin et al.
2021).

Warmer rearing temperature also decreased *Rickettsiella* titer, particularly in female spiders,
even though temperature effects on symbiont phenotype were more apparent in males than
females. This result is somewhat surprising because gravid females are usually expected to
carry higher loads of maternally transmitted symbionts than males; perhaps the use of post-
oviposition females in the present study depleted the bacterial loads of females. Associations
between temperature and symbiont titers have previously been demonstrated with other
endosymbionts (Breeuwer and Warren 1993, Lopez-Madrigal and Duarte 2019, Ross et al.
2020), and thus our results further support the potential links between temperature, titer, and
phenotypic outcome.

Environmental effects on symbiont phenotypic penetrance may have important implications in
both managed and natural ecosystems. *Wolbachia* is already in use in vector management in
several areas of the globe, with mixed results (Hoffmann et al. 2011, Dos Santos et al. 2022,
Hoffmann et al. 2024). Symbiont thermal tolerance is one factor that appears to limit the
success of this technique in some regions (Ross et al. 2020, 2023). As the use of symbionts to
control pests and vectors comes under consideration in a range of other systems (e.g., Gong et
al. 2020, Mateos et al. 2020) it becomes critical to characterize the environmental constraints
that might influence efficacy. Even in less-managed systems, the thermal optima of symbionts
might play a critical role in determining the distribution of pests and the ability of natural
enemies to control them (Zhang et al. 2019, Hague et al. 2022). Our study contributes to a
growing body of literature that suggests environmental contingency may be the rule for
interactions between hosts and reproductive manipulators, not the exception.

ACKNOWLEDGEMENTS
We thank L. Rosenwald, K. Butler, and E. Williams for assistance in the lab, and M. Doremus for
helpful comments on a previous version of this manuscript.

REFERENCES

Breeuwer, J.A. and Werren, J.H. (1993). Cytoplasmic incompatibility and bacterial density in *Nasonia vitripennis*. Genetics, 135, 565-574.

Chrostek, E., Martins, N., Marialva, M.S. and Teixeira, L. (2021). *Wolbachia*-conferred antiviral protection is determined by developmental temperature. mBio, 12, 10-1128.

Corbin, C., Heyworth, E.R., Ferrari, J., and Hurst, G.D.D. (2017). Heritable symbionts in a world of varying temperature. Heredity, 118, 10-20.

Corbin C., J. E. J., Chrostek E., Fenton A., and Hurst G.D.D. (2021). Thermal sensitivity of the *Spiroplasma-Drosophila hydei* protective symbiosis: The best of climes, the worst of climes. Molecular Ecology, 30, 1336-1344.

Curry, M. M., Paliulis, L. V., Welch, K. D., Harwood, J. D., and White, J. A. (2015). Multiple endosymbiont infections and reproductive manipulations in a linyphiid spider population. Heredity, 115, 146-152.

Doremus M.R., K. S. E., and Hunter M.S. (2019). Exposure to opposing temperature extremes causes comparable effects on *Cardinium* density but contrasting effects on *Cardinium*-induced cytoplasmic incompatibility. PLoS Pathogens, 15, 1008022.

Dos Santos, G.R., Durovni, B., Saraceni, V., Riback, T.I.S., Pinto, S.B., Anders, K.L., Moreira, L.A. and Salje, H. (2022). Estimating the effect of the wMel release programme on the incidence of dengue and chikungunya in Rio de Janeiro, Brazil: a spatiotemporal modelling study. The Lancet Infectious Diseases, 22, 1587-1595.

Gong, J.T., Li, Y., Li, T.P., Liang, Y., Hu, L., Zhang, D., Zhou, C.Y., Yang, C., Zhang, X., Zha, S.S. and Duan, X.Z. (2020). Stable introduction of plant-virus-inhibiting *Wolbachia* into planthoppers for rice protection. Current Biology, 30, 4837-4845.

Gu, S., Ross, P.A., Gill, A., Yang, Q., Ansermin, E., Sharma, S., Soleimannjad, S., Sharma, K., Callahan, A., Brown, C., Umina, P.A., Kristensen, T.N., and Hoffmann. A.A. (2023). A rapidly spreading deleterious aphid endosymbiont that uses horizontal as well as vertical transmission. PNAS, 120, 2217278120.Hague, M.T., Shropshire, J.D., Caldwell, C.N., Statz, J.P., Stanek, K.A., Conner, W.R. and Cooper, B.S. (2022). Temperature effects on cellular host-microbe interactions explain continent-wide endosymbiont prevalence. Current Biology, 32, 878-888.

Hoffmann, A. A., Montgomery, B.L., Popovici, J., Iturbe-Ormaetxe, I., Johnson, P.H., Muzzi, F., Greenfield, M., Durkan, M., Leong, Y.S., Dong, Y., Cook, H., Axford, J., Callahan, A.G., Kenny, N., Omodei, C., McGraw, E.A., Ryan, P.A., Ritchie, S.A., Turelli, M., and O'Neill, S.L. (2011). Successful establishment of *Wolbachia* in *Aedes* populations to suppress dengue transmission. Nature, 476, 454-457. Hoffmann, A.A., Ahmad, N.W., Keong, W.M., Ling, C.Y., Ahmad, N.A., Golding, N., Tierney, N., Jelip, J., Putit, P.W., Mokhtar, N. and Sandhu, S.S. (2024). Introduction

of *Aedes aegypti* mosquitoes carrying *wAlbB Wolbachia* sharply decreases dengue incidence in
disease hotspots. iScience, 27.

Hunter, M. S., Perlman S.J., and Kelly S.E. (2003). A bacterial symbiont in the Bacteroidetes
induces cytoplasmic incompatibility in the parasitoid wasp *Encarsia pergandiella*. The Royal
Society, Proceedings B, 270, 2185-2190.

Jones, J. E., Hurst, G.D.D. (2023). History matters: thermal environment before, but not during
wasp attack determines the efficiency of symbiont-mediated protection. Molecular Ecology, 32,
3340-3351.

López-Madrigal, S. and Duarte, E.H. (2019). Titer regulation in arthropod-*Wolbachia*
symbioses. FEMS microbiology letters, 366, 232.Martins, M., César, C.S. and Cogni, R. (2023).
The effects of temperature on prevalence of facultative insect heritable symbionts across
spatial and seasonal scales. Frontiers in Microbiology, 14, 1321341.

Mateos, M., Martinez Montoya, H., Lanzavecchia, S.B., Conte, C., Guillén, K., Morán-Aceves,
B.M., Toledo, J., Liedo, P., Asimakis, E.D., Doudoumis, V. and Kyritsis, G.A. (2020). *Wolbachia*
pipientis associated with tephritid fruit fly pests: from basic research to applications. Frontiers
in Microbiology, 11, 521402.

Penz, T., Schmitz-Esser, S., Kelly, S.E., Cass, B.N., Müller, A., Woyke, T., Malfatti, S.A., Hunter,
M.S., and Horn, M. (2012). Comparative genomics Suggests an independent origin of
cytoplasmic incompatibility in *Cardinium hertigii*. PLoS Genetics, 8, 1003012.

Poinsot, D., Charlat, S., and Merçot, H. (2003). On the mechanism of *Wolbachia*-induced
cytoplasmic incompatibility: confronting the models with facts. BioEssays 25, 259-265.

Pollmann M., M., L.D., Krimmer, P.D., Hasselmann, M., Perlman, S.J., Ballinger, M.J., Steidle,
J.L.M., and Gottlieb, Y. (2022). Highly transmissible cytoplasmic incompatibility by the
extracellular insect symbiont *Spiroplasma*. iScience, 25,104335.

Rosenwald L.C., Sitvarin, M. I. and White, J. A. (2020). Endosymbiotic *Rickettsiella* causes
cytoplasmic incompatibility in a spider host. The Royal Society, Proceedings B, 278, 20201107

Ross, P.A., Axford, J.K., Yang, Q., Staunton, K.M., Ritchie, S.A., Richardson, K.M. and Hoffmann,
A.A. (2020). Heatwaves cause fluctuations in *wMel Wolbachia* densities and frequencies in
Aedes aegypti. PLoS neglected tropical diseases, 14, e0007958. Ross, P.A., Elfekih, S., Collier, S.,
Klein, M.J., Lee, S.S., Dunn, M., Jackson, S., Zhang, Y., Axford, J.K., Gu, X. and Home, J.L. (2023).
Developing *Wolbachia*-based disease interventions for an extreme environment. PLoS
Pathogens, 19, e1011117.

Shropshire J.D., L. B., and Bordenstein S.R. (2020). Symbiont-mediated cytoplasmic
incompatibility: What have we learned in 50 years? eLife, 9, e61989.

Shropshire, J.D., Hamant, E. and Cooper, B.S., 2021. Male age and *Wolbachia* dynamics:
investigating how fast and why bacterial densities and cytoplasmic incompatibility strengths
vary. *MBio*, *12*, e02998-21.

Williams, D.A. (1982). Extra-binomial variation in logistic linear models. Journal of the Royal
Statistical Society: Series C (Applied Statistics), 31, 144-148.

Zhang, B., Leonard, S.P., Li, Y. and Moran, N.A. (2019). Obligate bacterial endosymbionts limit
thermal tolerance of insect host species. Proceedings of the National Academy of Sciences, 116,
24712-24718.