1 # Induction by caterpillars of stored and emitted volatiles in terpene
2 # chemotypes from populations of wild cotton (*Gossypium hirsutum*)
3

<u>Marine Mamin</u>[1], Mary V Clancy[1], Galien Flückiger[1], Teresa Quijano-Medina[2], Biiniza Pérez-Niño[2], Luis
Abdala-Roberts[2], Ted C.J. Turlings[1,3*], Carlos Bustos-Segura[1,4*]

[1] University of Neuchâtel, Institute of Biology, Fundamental and Applied Research in Chemical Ecology,
Neuchâtel, Switzerland

[2] Departamento de Ecología Tropical, Campus de Ciencias Biológicas y Agropecuarias, Universidad Autó-
noma de Yucatán, Mérida, México

[3] State Key Laboratory of Crop Stress Adaptation and Improvement, State Key Laboratory of Cotton Bio-
breeding and Integrated Utilization, School of Life Sciences, Henan University, Kaifeng, 475004, China

[4] Present address : Institute of Ecology and Environmental Sciences-Paris, INRAE, Sorbonne Université,
CNRS, IRD, Université de Paris, UPEC, Route de St Cyr, 78026 Versailles, France
14
15 *Correspondence: ted.turlings@unine.ch; carlos.bustos@inrae.fr

Abstract

Background

Upland cotton (*Gossypium hirsutum*) plants constitutively store volatile terpenes in their leaves, which are steadily emitted at low levels. Herbivory leads to a greater release of these stored volatiles. Additionally, damaged plants increase the accumulation of volatile terpenes in their leaves and begin to emit other terpenes and additional compounds. This has been well characterised for cultivated *G. hirsutum*, but little is known about volatile production in response to herbivory in wild populations. We investigated how damage by the beet armyworm (*Spodoptera exigua*) affects leaf-stored and emitted volatiles in wild *G. hirsutum* plants. These plants were grown in a greenhouse using seeds collected from populations found along the Yucatan coast, from where this cotton species originates. We assessed whether the differences in leaf terpene profiles between two known chemotypes persisted upon herbivory, in leaves and in headspace emissions, and whether these chemotypes also differed in the production and release of herbivory-induced volatiles. In addition to chemotypic variation, we further investigated intraspecific variation in the volatile response to herbivory among genotypes, populations, and four geographic regions.

Results

The difference between the two chemotypes persisted after herbivory in the stored volatile profile of induced leaves, as well in the emissions from damaged plants. Therefore, wild cotton chemotypes may differ in their airborne interactions with their environment. The specific terpenes distinguishing these chemotypes showed a weak inducibility, raising questions about their functions. Herbivory triggered changes in stored and emitted volatiles similar to what is known for cultivated varieties of *G. hirsutum.* However, we report for the first time on the emission of volatile aldoximes by cotton plants, which were only detected in the headspace upon herbivory, and displayed chemotypic and interpopulation variation. Intraspecific variation was also observed in the induced emissions of nitriles and certain terpenes. Moreover, chemotypes differed in their induction of (*E*)-β-ocimene stored in the leaves.

Conclusions

This comprehensive insight into herbivore-induced volatiles of wild cotton reveals variation in production and emission among populations. A full understanding of their ecological role may help in the development of future pest-management strategies for cotton crops.

Keywords

47 *Gossypium hirsutum*; wild cotton; herbivore-induced plant volatiles (HIPVs); intraspecific variation; chemo-

48 type; terpenes; aldoximes

Background

Plants emit volatiles that can play various defensive roles against insect herbivores. Volatile plant metab-

olites can function as toxins, repel insect feeding and oviposition, and attract natural enemies of their

attackers [1,2]. Volatiles also mediate signaling within plants [3] and between neighboring plants, increas-

ing their resistance against herbivores [4–6]. Unharmed plants can produce and emit certain volatiles con-

stitutively, a phenomenon particularly prevalent in plant species that accumulate volatile terpenes in spe-

cialized structures. These stored terpenes are continuously emitted at low levels by intact plants and are

released in much larger quantities when the storage organs are physically damaged, for example by chew-

ing herbivores [7–9]. In addition, when plants detect an insect attack, they may actively synthesize so-

called herbivore-induced plant volatiles (HIPVs), which can include compounds that are produced consti-

tutively but synthesized in greater amounts upon damage, or compounds whose synthesis is triggered

exclusively in response to herbivory [2,10].

Several wild plant species have been shown to exhibit intraspecific variation in their chemical phenotypes.

This chemodiversity often has a genetic basis and can modulate the interactions between plants and their

environment [11–13]. Intraspecific variation in volatile production has been described both for constitu-

tive and herbivore-induced compounds [14,15]. Various species that constitutively produce and store vol-

atile terpenes notably present discrete chemical polymorphisms in their terpene blend (chemotypes here-

after) [e.g. 14–18]. Importantly, this variation has been associated with differences in resistance to abiotic

stress [21,22], allelopathy [23], defense against pathogens [24] and herbivores [24–28], as well as airborne

signaling between plants when stored terpenes are released by damage [6,29]. Concerning HIPVs, varia-

tion has been reported for various classes of compounds across several wild species, such as: *Nicotiana*

attenuata [30,31], *Datura wrightii* [32], *Solanum carolinense* [33,34], *Arabidopsis thaliana* [35,36], *Ascle-*

pias syriaca [37], *Quercus robur* [38] and *Vicia sepium* [39]. Variation in HIPVs can be linked to differences

in synthetic pathways [40,41], but also to differences in the endogenous signaling cascades that regulate

the plants' response to herbivory [31]. In addition, differences in the attraction of herbivores [34,38] and

their natural enemies [34,36] have been found to be associated with variation in volatile induction.

Upland cotton (*Gossypium hirsutum* L. Malvaceae) is cultivated worldwide as a major source of natural

fiber, and suffers from high levels of herbivory from multiple insect pests [42]. Increasing our understand-

ing of the natural defense mechanisms of cotton could aid in the advancement of sustainable pest control

strategies and the development of more resistant cultivars [2,43–45]. Cotton plants constitutively store

volatile mono- and sesquiterpenes in specialized glands present in their leaf tissues [42,46]. Herbivory on

older leaves induces an increase in the concentrations of these terpenes in young undamaged leaves

[47,48]. These leaf-stored terpenes are steadily emitted at low levels from intact leaves, however large

quantities are immediately released following gland rupture, for instance caused by chewing insects [49–

52]. Therefore, volatile emissions instantly resulting from insect damage primarily comprise locally re-

leased leaf mono- and sesquiterpenes, together with green leaf volatiles (GLVs; aldehydes and alcohols),

that are linked to tissue disruption [49,50]. Herbivory also induces a delayed emission of mono-, sesqui-

and homoterpenes, which are systematically emitted (i.e. also by undamaged parts) following their in-

duced synthesis, and concurrently also triggers the emissions of other newly synthesized volatiles, such as

the aromatic indole and GLV esters [49–55]. When active physical damaging ceases, the emission of con-

stitutive leaf-stored terpenes drops rapidly, while induced compounds continue to be emitted [49]. The

distinction between constitutive and truly inducible terpenes is not so clear cut, and some studies report

systemic emissions of terpenes constitutively stored in leaves [54,55]. Collectively, volatile emissions of

insect-damaged cotton plants appear to serve as direct and indirect defenses [reviewed in 42], and are

involved in plant-plant interactions .

Domestication often leads to a reduction in defenses against herbivory in crop plants [59,60]. Modern

cultivars of *G. hirsutum* exhibit a depletion of genetic diversity [61] and also appear to be less resistant to

herbivory compared to wild-growing plants [62–65]. Therefore, exploring useful defensive traits in wild

populations of *G. hirsutum* should be considered. While volatile production in cultivated varieties has been

well characterized, less is known about the volatiles produced by wild *G. hirsutum* plants in response to

herbivory, and studies conducted so far mostly relate to naturalized varieties [62,63,66]. *G. hirsutum* was

domesticated at least 4,000 years ago in the Yucatán Peninsula (southeast Mexico) [61], where wild pop-

ulations still grow naturally along its northern coastline [67,68].

Recently, we described two heritable constitutive leaf-terpene chemotypes in wild cotton plants from the

Yucatan coast: one chemotype containing either none or low amounts of γ-terpinene (and other co-ex-

pressed monoterpenes), and the other chemotype exhibiting distinctively higher abundances of these

compounds in their leaves [69]. It is worthwhile to note that production of γ-terpinene has been only

described in wild/naturalized cotton and from crosses between cultivated and wild cotton [62,63,66]. If γ-
terpinene variation in leaves influences the composition of the volatile emissions of wild cotton plants, it
could impact the plants' interactions with organisms that use these volatiles as a cue, such as herbivores
looking for a host plant, their natural enemies, or neighboring plants.

The present study investigated the effect of damage by caterpillars (beet armyworm, *Spodoptera exigua*
Hübner; Lepidoptera; Noctuidae) on both the volatile terpenes stored in leaves and the volatile emissions
of plants from wild populations of *G. hirsutum*. Plants were grown in a greenhouse from seeds collected
from sixteen populations forming four distinct geographic zones distributed along the Yucatan coastline.
We assessed the consistency of the two leaf-terpene chemotypes, previously described as constitutive, in
the stored volatile profile of induced leaves, as well in the emissions of both undamaged and damaged
plants. In addition, we evaluated the variations in the induction by herbivory of other stored and released
volatiles between the two chemotypes, and among three nested levels of intraspecific variation: geno-
types, populations, and the geographic regions grouping the populations. As the prevalence of the two
chemotypes along the coast follows a geographic gradient [69], chemotypic variation is interrelated with
variation among populations.

The results presented here shed new light on the production of volatiles by wild *G. hirsutum* plants in
response to caterpillar herbivory, and uncover sources of variation in volatile induction, which may be
genetically based, thereby opening new avenues to study and exploit cotton HIPVs as a defense mecha-
nism against insect pests.

Methods

1 Plants and insects

Wild *Gossypium hirsutum* seeds were collected in 2019 and 2020 on the northern coast of the Yucatán
Peninsula (Mexico) and grown as described in [69]. From July to September, plants were grown in individ-
ual pots filled with 250 ml of soil (Profi Substrat, Einheitserde, Germany), randomly arranged within a sin-
gle greenhouse located in Neuchâtel. Seeds were sampled from several mother plants from 16 populations
in four regions along the coast (Table S1 and Fig. 1). As cotton cross-pollinates, seeds from the same
mother are either full or half-siblings, of unknown paternal source. Offspring from the same mother plant
is referred to here as genotype. Plants were grown in a greenhouse for approximately four weeks. Even-
tually, we obtained plants from one to six mother plants for each population, representing around 35 to

60 individuals per population, except for a few populations with particularly low germination rates (Table S1). To induce volatile production in the cotton plants, we used larvae of the moth *Spodoptera exigua* (Hübner; Lepidoptera; Noctuidae), a cosmopolitan generalist pest that also feeds on cotton. Eggs were supplied by Entocare (Wageningen, The Netherlands) and reared in the laboratory on wheat-germ based artificial diet (Frontier Scientific Services, Newark, USA).

2 Experimental process and sample collection

Plants were used when the fourth leaf was fully expanded but still developing. A total of 322 plants were subjected to herbivory and 353 were left undamaged, in six experimental batches. The numbers of replicates per region and population are described in Table S1. Second to early third-instar *S. exigua* larvae fed on the first and second leaves for two days, with a single larva enclosed on each leaf in clip cages (4.52 cm^2, BioQuip product, USA). Undamaged and damaged plants were kept in two adjacent rooms with similar configurations and conditions (L14:D10, T=28°C), to prevent any signaling effect of HIPVs. The treatments were alternated between rooms after each batch. After two days of herbivory, we collected the volatile emissions of a subset of the plants. A total of 45 undamaged and 48 damaged individuals were used in four batches for headspace collection, chosen to represent the various populations (Table S1). Afterwards, the fourth leaf of all plants was harvested, photographed, frozen in liquid nitrogen, and stored at -80 °C until processed. Photographs of all leaves were taken to quantify the total leaf area consumed by *S. exigua*, as well as to determine the area (size) of the fourth leaves and the total leaf area (including cotyledons and secondary leaves) of the plants used for the headspace collections. For this purpose, we used the Fiji distribution of ImageJ software [70].

3 Headspace volatile collection and GC-MS analyses

S. exigua larvae were removed before headspace sampling, to avoid the release of volatiles due to physical tissue disruption by active feeding. The caterpillars' frass was cleared away from the leaves. Pots were covered with aluminum foil to limit potential emissions from the soil. Headspace volatiles were collected in a push-pull air system of custom-made glass bottles (VTQ SA, Switzerland). Charcoal-purified and humidified air was pushed at 1 L min^{-1} and pulled at 0.6 L min^{-1} through trapping filters (25 mg of 80/100 mesh Hayesep-Q adsorbent, Sigma, Switzerland) for two hours. Filters were eluted with 100 µL of dichloromethane, with the addition of 10 µL of internal standard solution (nonyl-acetate, 20 ng µL^{-1}). Samples were

163 stored at -80 °C before being run on a gas chromatograph (GC Agilent 7890B) coupled to a mass spectrom-

164 eter (MSD Agilent 5977B): 1.5 µL were autoinjected in splitless mode onto an Agilent HP-5MS column (30

165 m × 250 µm × 0.25 µm). The temperature was kept at 40 °C for 3.5 min, then ramped to 100 °C at 8 °C min^{-1}

166 and to 230 °C at 5 °C min^{-1}. Helium was used as the carrier gas, with a stable flow of 0.9 mL min^{-1}.

4 Hexane leaf extractions and GC-MS analyses

Hexane leaf extractions and subsequent analyses were performed as described in Clancy et al. (2023).

Briefly, 100 mg of finely ground frozen leaf material was extracted with 1 mL of hexane and kept at 4 °C

for 24 hours. 800 µL was recovered and 8 µL of internal standard solution (nonyl acetate, 2000 ng µL^{-1})

was added. The samples were directly injected into a gas chromatograph (GC Agilent 8890A) coupled to a

mass spectrometer (MSD Agilent 5977B): 1.5 µL were injected in splitless mode onto an Agilent HP-5MS

column (30 m × 250 µm × 0.25 µm). Temperature started at 40°C for 0 min, ramped to 130 °C at 15 °C min^{-1},

then to 190 °C at 60 °C min^{-1}, and to 290 °C at 30 °C min^{-1}, then held for 0.5 min. Helium was used as

carrier gas, with a stable flow of 0.9 mL min^{-1}.

5 GC-MS data processing

PyMS (O'Callaghan et al., 2012) was used to integrate and align peaks from the total ion chromatograms

of all samples (https://github.com/bustossc/PyMS-routine/). Analytes were quantified using relative re-

sponse factors obtained from a series of external calibration curves of commercial analytical standards in

comparison to nonyl acetate (Table S2 and S3). For compounds lacking available response factors, we em-

ployed the relative response factor associated with the closest standard in retention time and similar

chemical structure, while normalizing by molecular masses (Kreuzwieser et al. 2014). When possible, ana-

lytes were identified based on comparison with commercial standards. Aldoximes were identified with

synthesized standards kindly provided by Dr. Tobias Köllner. Compounds with no standard available were

identified based on mass spectra comparison with NIST 2.3 mass spectral library, an in-house library, and

Kovats retention indices [71] (Table S2 and S3). Components retained in the analyses were present in at

least 10% of the samples. For the hexane extracts, values were normalized by the fresh weights of the

leaves (units: ng mg^{-1} FW). For headspace samples, values were normalized by hours of collection and the

total leaf area of the plants (units: ng h^{-1} cm^{-2}).

Statistical analyses

Statistical analyses were carried out with R 4.3.2 [72]. A significance level of α = 0.05 was used for all analyses. Plots were built with "ggplot2" [73] (v. 3.4.4).

1 Multivariate analyses of volatiles composition: pRDAs and correlation analyses

Multivariate analyses were always performed on the proportions of compounds relative to the sum of all analytes of interest. To deal with the left-censored and compositional nature of the data, zeroes were imputed by multiplicative simple replacement with multRepl() from "zCompositions" [74] (v. 1.5.0.1), using the smallest non-zero value of each compound as detection limits and a delta of 0.65. We subsequently employed a centered log-ratio (clr) transformation with clr() from "compositions"[75] (v. 2.0.8). Data were finally scaled and centered using scale() from the R base package. To evaluate correlations among compounds, hierarchical clustering was performed on the correlation-based (Pearson) dissimilarity matrix with Ward's minimum variance method (Ward.D2) using pvclust() from "pvclust" [76] (v. 2.2.0). Approximately Unbiased (AU) p-values were computed by multiscale bootstrap resampling with 10,000 replications. Correlations heatmaps were plotted with "ComplexHeatmap" [77] (v. 2.16.0). Partial redundancy analyses (pRDAs) were conducted with rda() from "vegan"[78] (v. 2.6.4). Models' structures are described in the following paragraphs. We then performed a type II permutation test with MVA.anova() from "RVAide-Memoire"[79] (v. 0.9.83.7). Pairwise comparisons were performed with pairwise.factorfit() from the same package.

2 Analyses of single compounds: GLMMs

Univariate analyses of concentrations of compounds in leaves or emission rates were performed with generalized linear mixed models (GLMMs) using glmmTMB [80](v. 1.1.9) with Gamma or Tweedie exponential families of distribution, depending on models' fit. Models' fits were checked with the package "DHARMa"[81] (v. 0.4.6). Effects of the explanatory variables were tested with Wald chi-squared tests (type-II) using Anova() from "car" [82] (v. 3.1.2). We plotted estimated marginal means and their 95% confidence intervals (CI), extracted from the models with ggemmeans() from "ggeffects" [83] (v. 1.3.4). When plotted, data points represented observed values. In all models, we controlled for the effects of plants' "size" (leaf extracts: fourth leaf area; headspace emissions: total leaf area) and of experimental batches (leaf extracts: n = 6; headspace emissions: n = 4) by including them as fixed effects (Table S2 and S3). γ-Terpinene and coexpressed compounds were analyzed within the producing chemotype only, as the other chemotype either does not produce them, or only in very low amounts. Population and genotype

220 were included as random intercepts in all models, except for those relying solely on the headspace emis-
221 sions of damaged plants, where 86% of the data consisted of individual data points per genotype (Table
222 S2 and S3). When necessary, pairwise comparisons were performed with emmeans() from "emmeans",
223 with multiple test adjustment (FDR)[84] (v. 1.10.0).

3 Volatile terpenes in leaves: herbivory and variation in induction

To assess how herbivory affected the leaf composition in volatile terpenes for the two chemotypes, a pRDA
was performed on the relative abundances of all leaf mono- and sesquiterpenes. Herbivory, chemotype
and their interaction were used as explanatory variables while leaf area was partialled out. Then, we ana-
lyzed the effect of herbivory and damage intensity (total leaf area consumed by caterpillars) on the total
concentrations of the leaf volatile terpenes and a subset of compounds using GLMMs (Table S2: model
structures A and E). We tested whether there was variation in the induction (i.e. increase in the leaf con-
centrations; group-by-herbivory interaction) of volatile terpenes between the two chemotypes and among
nested levels of intraspecific variation: 88 genotypes, 16 populations and four geographic regions. Varia-
tion in the response of genotypes and populations to herbivory was modelled with random slopes and
their significance was tested by comparing nested models with a random slope component to a model
without (Restricted Maximum Likelihood method) with Likelihood ratio tests. Chemotype was included in
the fixed effects of these models (Table S2: model structures B and C). The random slope component in-
cluded a term for the variance in slopes and one for the covariance between slopes and intercepts unless
the model did not converge. In that case, the covariance term was removed. Variation in response to her-
bivory among region and chemotypes was modelled as interaction between these groups and herbivory,
set as fixed effects). Because of their geographic distribution, chemotypes were partially nested in regions.
Therefore, to avoid collinearity, they were always analyzed separately (Table S2: model structures A and
D).

4 Headspace volatile emissions: herbivory and variation in induction

To examine whether the intraspecific variation observed in the leaf-stored terpenes by Clancy et al. (2023)
was reflected in the emissions, we performed Spearman correlation tests, separately for undamaged and
damaged plants, on the ratio of concentrations of γ-terpinene and α-pinene (the dominant leaf-stored
monoterpene) in leaves and in headspace emissions. To visualize the composition of the emitted blends
of undamaged and damaged plants of both chemotypes, we plotted the median proportions with
"treemapify" [85] (v. 2.5.6). To further understand how herbivory and chemotypes affected the composi-
tion of headspace emissions a pRDA was conducted on the relative abundances of all emitted volatiles

together. Herbivory, chemotype and their interaction were used as explanatory variables while the total leaf area was partialled out. Then, we analyzed the effect of herbivory and damage intensity (total leaf area consumed by caterpillars) on the total emission rates and on the emissions of same terpenes analyzed for the leaves (Table S3: model structures A and C).

Variation in the volatile emissions induced by caterpillar herbivory was evaluated between chemotypes, as well as two levels of intraspecific variation: among the four regions and 16 populations. Genotypes were not assessed because of the lack of replication at this level. To evaluate variation in the volatile response to herbivory rather than pre-existing differences in compounds already emitted by intact plants, we selected only highly inducible volatiles, defined from the pRDA assessing the effect of herbivory on the headspace emissions. We first evaluated whether there was variation in the total emissions of these highly inducible compounds (Table S3, model structures C, D and E). Variation among population was modelled through random intercepts (Table S3: model structures E) and their significance was tested by comparing nested models with random intercept component to models without (Restricted Maximum Likelihood method) with Likelihood ratio tests. Chemotypes and regions were analyzed as fixed effects, and separately, as mentioned above. To unveil potential variation in the composition of the induced emissions, pRDAs were performed on the proportions of highly inducible volatiles, selected from the transformed and scaled matrix of the relative abundances of all compounds emitted by damaged plants, separately for chemotype and region. Chemotype/region, the leaf surface damaged and their interaction were used as explanatory variables while the total leaf area was partialled out. We then further tested whether chemotypes and regions varied significantly in their emission rates of the different candidate compounds highlighted by the pRDAs with GLMMs, as explained for the total induced emissions (Table S3, model structure C and D). Compounds with significant variations were also assessed for variation among populations (Table 4, model structure E). For inducible compounds differing between chemotypes or regions but also emitted by undamaged plants, the interaction effect between chemotype/region and herbivory was tested (Table S3, model structure A and B). A significant interaction implied that chemotypes/regions differed in their response to herbivory, not solely by pre-existing disparities.

Results

1 Leaf-stored volatiles

We extracted volatile content from the leaves of undamaged (n=353) and damaged (n=322) wild cotton plants (Fig. 1). A total of 15 monoterpenes, 18 sesquiterpenes and three GLVs were quantified in the leaf extracts (Table S4). Subsequent analyses focus on terpenes.

1.1 Monoterpene chemotypes

Chemotypes were assigned to each plant based on the leaf volatile profile. First, undamaged plants were chemotyped based on the contribution of six strongly correlated compounds relative to the total mono-terpene concentration (γ-terpinene, limonene, α-thujene, terpinolene, p-cymene, and α-terpinene, in or-der of abundance; *sensu* [69]; Fig. S1, Table S4). Plants containing higher proportions (> 4%) of these com-pounds were assigned to chemotype A; plants with less than 4% were assigned to chemotype B (Fig. 2). The six variable monoterpenes remained highly correlated upon the two days of herbivory (Fig. S2) and damaged plants could be separated in two chemotypes in the same way as undamaged plants (Fig. 2).

1.2 Effect of herbivory on the leaf-stored volatile terpene profiles

Following herbivory during two days on the first and second leaves of four-leaf plants, the total leaf con-centration of mono- and sesquiterpenes increased by 9% on average in the undamaged fourth leaves (Fig. 3A, $\chi^2_{(1)}$ = 12.407, p < 0.001). The same compounds were detected in the leaf terpene profiles of undam-aged and damaged plants (Table S4), but a partial redundancy analysis (pRDA) revealed that herbivory significantly affected their relative abundances (Fig. 3B, $F_{(1,670)}$ = 6.414, p = 0.001). This change in compo-sition upon herbivory appeared to be mainly linked to the increase in the proportion of the monoterpene (*E*)-β-ocimene, followed by several sesquiterpenes, notably δ-cadinene and α-copaene (Fig. 3B). The in-duction of (*E*)-β-ocimene depended on chemotype (Fig. 3A; chemotype x herbivory: $\chi^2_{(1)}$ = 7.694, p = 0.005), with a 64% increase in chemotype A plants (contrast of EMMs: p < 0.001) and a 108% increase in chemo-type B plants (contrast of EMMs: p < 0.001). Leaf content of the sesquiterpene δ-cadinene increased fol-lowing herbivory by 27% (Fig. 3A, $\chi^2_{(1)}$ = 27.201, p < 0.001). (*E*)-β-caryophyllene, the most abundant ses-quiterpene in leaves (Table S4), increased by 11% (Fig. 2A: $\chi^2_{(1)}$ = 14.254, p < 0.001). In contrast, levels of α-pinene, the most abundant monoterpene in leaves (Table S4), did not change upon herbivory (Fig. 3A, $\chi^2_{(1)}$ = 0.437, p = 0.509). This was also the case for γ-terpinene, the main monoterpene characteristic for chemotype A (Fig. 3A, $\chi^2_{(1)}$ = 0.766, p = 0.381). The total accumulation of (*E*)-β-ocimene increased with the

306   amount of leaf tissue consumed by *S. exigua,* but this effect of damage intensity was not significant for the

307   other terpenes tested (Fig. S3: (*E*)-β-ocimene: $\chi^2_{(1)}$ = 5.619, p = 0.018).

1.3 Variation in the induction of leaf-stored volatiles

We investigated whether there was intraspecific variation in the volatile response to herbivory for the

total leaf terpene content and for the aforementioned compounds. As volatile leaf terpenes are constitu-

tively produced, we assessed the variation in the increase of leaf concentrations induced by herbivory, i.e.

a group-by-herbivory interaction, groups being either chemotypes, genotypes, populations or regions (Ta-

ble S5). As mentioned above, induction of (*E*)-β-ocimene varied between the two chemotypes (Fig. 3A and

Table S5; chemotype x herbivory: $\chi^2_{(1)}$ = 7.694, p = 0.005), with a stronger induction observed in chemotype

B plants. Chemotype A plants exhibited constitutively higher amounts of (*E*)-β-ocimene in their leaves than

chemotype B plants (Fig. S4, $\chi^2_{(1)}$ = 8.247, p = 0.004); this difference was no longer significant following

herbivory (Fig. S4, $\chi^2_{(1)}$ = 0.793, p = 0.373). We did not find any significant variation in induction of any

tested compounds among genotypes, populations, and regions (Table S5).

2 Headspace volatile emissions

The volatile emissions of undamaged and damaged wild cotton plants were collected and characterized in

a subset of the plants that were afterwards used for leaf extraction (Fig. 1; 45 undamaged and 48 damaged

plants). We quantified a total of 54 compounds in the headspace (Table S6). Terpenes represented the

richest group, with 13 monoterpenes, 13 sesquiterpenes, and the homoterpenes DMNT and TMTT identi-

fied. Additionally, headspace emissions contained many fatty acids derivatives, which comprised GLV al-

dehydes, alcohols, and esters, and (*Z*)-jasmone (Table S6). Also, we identified three aliphatic aldoximes

and several aromatic compounds including two nitriles. Most of the mono- and sesquiterpenes detected

in the headspace were also found in the leaf extracts, as well as a GLV aldehyde (Fig. 4A).

2.1 Monoterpene chemotypes

We assessed whether the leaf-terpene chemotypes could also be distinguished from their headspace emis-

sions, both for undamaged and damaged plants. Of the six monoterpenes that were used to define chemo-

types A and B using leaf extracts, the following compounds were also detected in the headspace emissions

(in order of abundance): γ-terpinene, limonene, p-cymene, and α-thujene (Table S6). For both undamaged

and damaged plants, the ratio of γ-terpinene to α-pinene in headspace emissions was correlated to their

ratios in the leaves (Fig. 5) and volatile emissions of the two chemotypes were clearly distinguishable by

the prevalence of γ-terpinene, p-cymene, and α-thujene (Fig. 4B and 6A), which were highly correlated

(Fig. S5 and S6).

2.2 Effect of herbivory on the emitted volatile profile

The volatile blend emitted by undamaged plants was mainly composed of mono- and sesquiterpenes that

were also found in leaf extracts, with (E)-β-caryophyllene and α-pinene dominating the median profile of

both chemotype A and B plants (Fig. 4B). Herbivory caused the total volatile emissions to increase 8.7-fold

(Fig. 6B, $\chi^2_{(1)}$ = 146.449, p < 0.001) and triggered qualitative and quantitative changes in their composition

(Fig. 4, 6A and S7). Damaged plants released several compounds that were absent from the headspace of

intact plants, specifically aliphatic aldoximes, benzyl nitrile and another unidentified nitrile, the aromatics

indole and methyl anthranilate, butyric and hexenoic acid esters of GLVs, (Z)-jasmone, the acyclic sesquit-

erpenes (E)-β-farnesene and (E,E)-α-farnesene, and monoterpene (E)-myroxide (Fig. 4). Other compounds

such as the acyclic monoterpenes (E)-β-ocimene, linalool and (Z)-linalool oxide, the homoterpenes DMNT

and TMTT, and the GLV ester (Z)-3-hexenyl acetate were emitted in low amounts by some of the undam-

aged plants, but their prevalence greatly increased following herbivory (Fig. 4B, 6A and S7). Indole, DMNT,

(Z)-3-hexenyl acetate, and (E)-β-ocimene were major constituents of the volatile blend from damaged

plants of both chemotypes (Fig. 4B). The emission rates of the mono- and sesquiterpenes that were also

found in the leaf extracts generally exhibited small increases upon herbivory (Fig. 6B). However, due to

greater increases of other compounds, their relative contribution to the emissions of damaged plants de-

creased (Fig. 4B and 6A). This was not the case of (E)-β-ocimene, which exhibited a distinctly high increase

in response to herbivory compared to the other terpenes found in both headspace and leaves. Irrespective

of chemotype ($\chi^2_{(1)}$ = 1.901, p = 0.168), the amounts emitted of (E)-β-ocimene rose 26-fold in response to

herbivory (Fig. 6B, $\chi^2_{(1)}$ = 139.057, p < 0.001). In contrast, (E)-β-caryophyllene and α-pinene only increased

2.6-fold (Fig. 6B, $\chi^2_{(1)}$ = 33.800, p < 0.001) and 3.2-fold (Fig. 6B, $\chi^2_{(1)}$ = 56.859, p < 0.001) respectively.

Emissions of δ-cadinene showed an intermediate increase of 11-fold (Fig. 6B, $\chi^2_{(1)}$ = 62.593, p < 0.001). A

particularly low response was observed for γ-terpinene, with only a marginal increase in the headspace of

1.7-fold following herbivory (Fig. 6B, χ2(1) = 3.868, p = 0.049).Emissions of γ-terpinene by damaged plants

were independent of damage level (Fig 6C, $\chi^2_{(1)}$ = 1.299, p = 0.254), as with p-cymene, α-thujene, and lim-

onene (Fig. S8). The emissions of all other terpenes detected in both headspace and leaf extracts were

positively correlated with damage intensity (Fig. S8).

2.3 Variation in the induction of volatile emissions

We investigated whether there were differences in the volatile emissions induced by *S. exigua* between chemotypes, as well as among regions and populations. Genotypes were not assessed because of the low level of replication. Our analyses focused on highly inducible volatiles. These were defined as the compounds with a negative score on the first axis, which separated damaged and undamaged plants, of the pRDA in Figure 5A (see Fig. S8 for a biplot with all compounds displayed). Total emissions of these highly inducible compounds did not differ between chemotypes (Fig. S9, $\chi^2_{(2)}$ = 0.003, p = 0.954), nor among regions (Fig. S9, $\chi^2_{(3)}$ = 2.893, p = 0.408) or populations (Fig. S9, $\chi^2_{(1)}$ = 1.344, p = 0.246). To further investigate differences between chemotypes and among regions, a new set of pRDAs were conducted on the abundances of these highly inducible compounds relative to the whole volatile blend emitted by damaged plants. These analyses revealed differences between chemotypes (Fig. S10, $F_{(1,42)}$ = 1.189, p = 0.009) and among regions (Fig. S11, $F_{(3,38)}$ = 1.904, p = 0.005) in the composition of their HIPV emissions. Of the compounds highlighted by the pRDAs, the terpenes (*E,E*)-α-farnesene, (*E*)-myroxide, the three aldoximes and the two nitriles showed statistically significant differences in emission rates between chemotypes and/or regions (Table S7). These compounds were all only detected in the headspace of damaged plants. We also observed variations in the emissions of TMTT and δ-cadinene (Fig. S10 and S11); these volatiles were also found in undamaged plants but in lower amounts. The absence of significant interactions between region/chemotype and herbivory suggests that the observed differences might not be specifically linked to a distinct response to herbivory (Fig. S10 and S11). Emission rates of (*E,E*)-α-farnesene and (*E*)-myroxide varied among regions, as well as among populations, but not between chemotypes (Table S7, Fig. 7 and Fig. S9). The three aldoximes, which were strongly correlated (Fig. S6), varied both among regions and between chemotypes. Plants from Celestún (far west) and Sisal (west) produced higher quantities of aldoximes compared to plants from regions further east (Table S7, Fig. 6 and Fig. S9). Chemotype B plants released higher amounts of these compounds than chemotype A plants (Table S7, Fig. 6 and Fig. S9). Thus, variation could stem both from the chemotype and the geographic region, which are partially nested. However, even when considering the effect of chemotype in the models, significant variation was observed at the population level for the two methylbutyraldoximes (Table S7). Benzyl nitrile showed the same trend in variation as the aldoximes, while the opposite was observed for the unknown nitrile (Table S7, Fig. 6). For these latter two compounds, however, no evidence of variation among populations was observed (Table S7).

Discussion

1 Consistency of the leaf-terpenes chemotypes

The two previously reported constitutive leaf-terpene chemotypes [69] were also distinct in their leaf volatile terpene profile after herbivory. They were also distinctive in their headspace emissions, both from undamaged and damaged plants. The fact that the variation in the leaf content of γ-terpinene and its correlated compounds was mirrored in the emissions suggests that, in both cases, the primary source of emission for these volatiles are the leaf storage pools, and that the chemotypes of wild cotton plants could differ in their airborne interactions with their environment. Volatile terpenes likely released from stored pools have been found to repel adult moths and piercing-sucking pests of cultivated cotton [86,87], and appear to function as foraging cues for some species of parasitoid wasps [87–90]. Additional research is needed to evaluate how the two chemotypes may impact wild cotton's interactions with its insect herbivores and their natural enemies. Our results suggest that the difference between the chemotypes may be of lesser importance in this context, as γ-terpinene exhibited a notably low inducibility in comparison to other leaf monoterpenes. Measurements at additional time points and molecular studies would be required to confirm how the synthesis of γ-terpinene and correlated compounds respond to *S. exigua* herbivory. Moreover, as induction of volatiles in cotton can be specific to the species of an attacking herbivore [91–94], we recommend that future work investigates the response of wild cotton chemotypes to other herbivore species.

Increased production of monoterpenes has also been reported in several plant species in response to abiotic stresses such as drought and excessive temperatures, against which they may offer protective properties [7,95–98]. Interestingly, some plant species specifically increase γ-terpinene concentrations in their tissues under drought conditions [99,100], and it has been demonstrated that γ-terpinene has a particularly high antioxidant capacity [101,102]. Previous work on Yucatanese wild cotton documented that populations in higher-temperature environments tend to exhibit a greater density of glands (albeit non-significantly), while herbivory pressure and gland density are not correlated among populations [103]. These findings suggests that abiotic conditions could potentially be significant factors driving variation in gland density and their chemical content. Considering the particularly harsh environments in which wild cotton grows [68], it would be interesting to investigate the relationship between γ-terpinene content and tolerance to extreme environmental conditions.

2 Effects of herbivory on the leaf-stored and emitted volatile profiles

Two days of herbivory on the first and second leaves led to a rapid accumulation of several volatile terpenes in the undamaged fourth leaf of the same plants. The leaf terpenes differed in their inducibility, which resulted in compositional changes in leaf volatile profiles triggered by herbivory. A similar variation in inducibility among compounds has previously been described for domesticated varieties, with (*E*)-β-ocimene being the most responsive to damage, and with sesquiterpenes exhibiting a stronger response than monoterpenes [47,48]. The higher inducibility of (*E*)-β-ocimene may allow for the increased synthesis of heliocides H1 and H4, the most inducible gland-stored, non-volatile defensive terpenoid aldehydes produced by cotton, for which (*E)*-β-ocimene is a precursor, together with hemigossypolone [47,104,105].

The constitutive volatile emissions of wild cotton consisted mostly of terpenes that also accumulated in the leaves: several cyclic terpenes, along with the linear terpenes β-myrcene and (*E*)-β-ocimene. This parallels previous descriptions of cultivated varieties, where these compounds are known to be stored in glands and constitutively released in small amounts [46,49,52]. For most of these terpenes, we found a small increase in their emission rates following herbivory. As there was no active damage occurring on the plant during sampling, this increase could reflect passive release from storage pools in induced undamaged leaves, although we cannot exclude some leakage from very fresh wounds, nor can a release from sources other than storage pools be fully ruled out [9,106]. The latter is already known for (*E*)-β-ocimene in cultivated cotton, which is not only accumulated in the leaves but is also *de novo* synthesized and systemically emitted upon induction [52]. Consistent with this previously observed phenomenon, we found a particularly strong induction of (*E*)-β-ocimene emissions in comparison to the other terpenes detected in both leaves and headspace. Together with (*E*)-β-ocimene, the major compounds induced by *S. exigua* herbivory in wild cotton plants included indole, (*Z*)-3-hexenyl acetate, and several terpenes that were distinct from those found in stored pools, namely the linear terpenoids linalool, (*E*)-β-farnesene, (*E,E*)-α-farnesene, DMNT, and TMTT. This aligns with previous findings in cultivated cotton, showing that these compounds are newly synthesized following damage [49–53,55].

 We identified several compounds that, to the best of our knowledge, have not yet been reported for cotton, including aliphatic aldoximes, nitriles, and the aromatic compound 2-phenylethanol. A nitrile compound previously described in wild cotton from Yucatán [58] likely corresponds to the nitrile we could not conclusively identify. Aldoximes are nitrogenous compounds synthesized from amino acids by cytochrome P450 monooxygenases of the CYP79 family, found across all angiosperms, and are precursors to nitriles [107,108]. Aldoximes and nitriles may be precursors to volatile alcohols, such as 2-phenylethanol

[108,109]. Various plant species emit volatile aliphatic aldoximes in response to herbivory or jasmonic acid

treatment [108]. In *Populus trichocarpa,* herbivore-induced CYP79 enzymes produce (*E/Z*)-2-methylbutyr-

aldoximes from L-isoleucine in response to caterpillar herbivory [107], and CYP71 enzymes convert aldox-

imes into nitriles, including benzyl nitrile, from the semi-volatile compound phenylacetaldoxime [109].

Volatile aldoximes and nitriles could be involved in cotton indirect defenses; in *Populus nigra*, methylbutyr-

aldoximes and benzyl nitrile are central to the recruitment of parasitoid wasps, despite being only minor

components of the full induced volatile blend [110]. Benzyl nitrile is involved in direct defense mechanisms

of *Populus trichocarpa and P. nigra* [109,111]. Aldoximes and nitriles might be emitted only in small

amounts by cultivated *G. hirsutum*, and therefore more difficult to detect with classic sampling and anal-

yses [112]. Our results could imply that wild cotton plants produce more aldoximes and nitriles than culti-

vated varieties. This might be of interest for future studies as it could offer an opportunity to enhance the

resistance of cultivated cotton varieties to herbivorous pests.

466   3 Variation in the induction of leaf-stored and emitted volatiles

In terms of leaf volatile content, the two chemotypes varied in their inducibility (magnitude of the increase

in leaf concentrations) of the monoterpene (*E*)-β-ocimene, whereas there was no difference between

chemotypes in the induction of (*E*)-β-ocimene in the headspace emissions. This is likely due to the ealier

mentioned possibility that most of its induced emissions may derive from new synthesis rather than stored

pools [52]. This variation in the inducibility of (E)-β-ocimene in leaves could have implications for wild

cotton plants' airborne interactions, particularly in scenarios involving subsequent leaf damage, as it would

lead to the release of the induced stored pools [51,53]. As for headspace HIPV emissions, we observed

considerable intraspecific variation at the population and regional levels in the emissions of the terpenes

(*E,E*)-α-farnesene and (*E*)-myroxide, as well as in the emissions of aldoximes. Nitriles were also found to

differ, to a lesser extent, among regions. Quantities of aldoximes and nitriles were also observed to vary

between the two leaf-terpene chemotypes. As the prevalence of the two chemotypes was partially nested

within region, the source of variation in emissions of these compounds cannot be conclusively ascertained.

Having said this, the similarity between damaged plants from Chicxulub (high proportion of chemotype B)

and Coloradas (dominance of chemotype A) in the production of both classes of compounds suggests that

this variation is more likely linked to interpopulation differences than to the chemotypes themselves. In

contrast, chemotype-specific responses to caterpillar herbivory have been described in another species

(*Tanacetum vulgare*) for both leaf-stored and emitted volatiles [113]. Despite being confined to coastal

habitats, Yucatanese wild cotton is subjected to substantial spatial variability in herbivore pressure and

abiotic conditions [103,114–116], which could be associated with the origin of the observed variation in HIPVs. Further work is required to understand the drivers of these differences. Given the multiple proposed functions of HIPVs in plant defense against insect herbivores [2], the differences in HIPVs that we observed could have consequences for wild cotton's resistance to herbivores. Genotypes producing higher amounts of certain HIPVs that enhance their direct or indirect defenses as well as those of neighboring plants [57] could be a valuable genetic resource in the development of more resistant cotton cultivars. The leaf volatiles may also enhance tolerance to abiotic stresses such as extreme temperatures and drought [7], which is worthy of further exploration.

Conclusions

In this study, we present a comprehensive description of the quantitative and qualitative changes triggered by caterpillar herbivory in the leaf-stored volatile terpenes and compounds emitted by wild *G. hirsutum* plants from the Yucatán Peninsula. We show that two previously described monoterpene chemotypes can also be distinguished based on their headspace emissions, which is true for both undamaged and caterpillar-damaged plants. Furthermore, we provide a first report of aldoximes being identified as cotton HIPVs. Wild cotton volatile responses to herbivory were found to vary among chemotype, geographic location and population. Future research should examine the consequences of this variation for the plant's resistance to herbivory, or other biotic and abiotic stresses.

Ethics approval and consent to participate

Not applicable

Consent for publication

Not applicable

Availability of data and materials

The datasets generated during the current study, as well as the R code for the analyses are available in the Zenodo repository, https://doi.org/10.5281/zenodo.10927508. The procedure for spectra analyses of GC-MS data is available at https://github.com/bustossc/PyMS-routine.

Competing interests

The authors declare that they have no competing interests.

Funding

This work was supported by the Swiss National Science Foundation [grant 315230_185319]. Permit information for sampling of plant material in Yucatán: SEMARNAT- SCPA/DGVS/00585/20 SEMARNAT SCPA/DGVS/2974/20.

Authors' contributions

T.C.J.T. and L.A-R., C.B-S., M.V.C. and M.M. designed the research. L.A-R., T.Q-M. and B.P-N. collected the cotton seeds. C.B-S., M.V.C., M.M. and G.F. performed the experiment and processed the samples. M.M. and C.B-S. processed and analyzed the data. M.M. wrote the first draft of the manuscript with contributions by all authors.

Acknowledgments

The authors thank Thomas Degen (UniNE) for the illustrations of cotton and for helping to create figure 1, Karla Ku Durán (UADY) for laboratory support, Dominic Davis-Foster (Staffordshire U) for maintaining the PyMassSpec repository, Tobias Köllner (Max Planck Institute for Chemical Ecology) for the aldoximes standards, and Fanny Deiss, Sara Leite Dias, Jonathan Interian, and Miguel Cárdenas for help with sample collection.

References

[1]   Unsicker SB, Kunert G, Gershenzon J. Protective perfumes: the role of vegetative volatiles in plant defense against herbivores. Curr Opin Plant Biol 2009;12:479–85. https://doi.org/10.1016/j.pbi.2009.04.001.
[2]   Turlings TC, Erb M. Tritrophic interactions mediated by herbivore-induced plant volatiles: mechanisms, ecological relevance, and application potential. Annu Rev Entomol 2018;63:433–52. https://doi.org/10.1146/annurev-ento-020117-043507.
[3]   Rodriguez-Saona CR, Frost CJ. New evidence for a multi-functional role of herbivore-induced plant volatiles in defense against herbivores. Plant Signal Behav 2010;5:58–60. https://doi.org/10.4161/psb.5.1.10160.
[4]   Karban R, Yang LH, Edwards KF. Volatile communication between plants that affects herbivory: A meta-analysis. Ecol Lett 2014;17:44–52. https://doi.org/10.1111/ele.12205.
[5]   Erb M, Veyrat N, Robert CAM, Xu H, Frey M, Ton J, et al. Indole is an essential herbivore-induced volatile priming signal in maize. Nat Commun 2015;6:6273. https://doi.org/10.1038/ncomms7273.
[6]   Grof-Tisza P, Kruizenga N, Tervahauta AI, Blande JD. Volatile-mediated induced and passively acquired resistance in sagebrush (*Artemisia tridentata*). J Chem Ecol 2022;48:730–45. https://doi.org/10.1007/s10886-022-01378-y.

[7] Loreto F, Schnitzler J-P. Abiotic stresses and induced BVOCs. Trends Plant Sci
2010;15:154–66. https://doi.org/10.1016/j.tplants.2009.12.006.
[8] Lange BM. The evolution of plant secretory structures and emergence of terpenoid chemi-
cal diversity. Annu Rev Plant Biol 2015;66:139–59. https://doi.org/10.1146/annurev-ar-
plant-043014-114639.
[9] Bustos-Segura C, Foley WJ. Foliar terpene chemotypes and herbivory determine variation
in plant volatile emissions. J Chem Ecol 2018;44:51–61. https://doi.org/10.1007/s10886-
017-0919-8.
[10] Dicke M, Baldwin IT. The evolutionary context for herbivore-induced plant volatiles: be-
yond the 'cry for help.' Trends Plant Sci 2010;15:167–75.
https://doi.org/10.1016/j.tplants.2009.12.002.
[11] Moore BD, Andrew RL, Külheim C, Foley WJ. Explaining intraspecific diversity in plant
secondary metabolites in an ecological context. New Phytol 2014;201:733–50.
https://doi.org/10.1111/nph.12526.
[12] Kessler A, Kalske A. Plant Secondary Metabolite Diversity and Species Interactions. Annu
Rev Ecol Evol Syst 2018;49:115–38. https://doi.org/10.1146/annurev-ecolsys-110617-
062406.
[13] Müller C, Junker RR. Chemical phenotype as important and dynamic niche dimension of
plants. New Phytol 2022;234:1168–74. https://doi.org/10.1111/nph.18075.
[14] Hare JD. Ecological role of volatiles produced by plants in response to damage by herbivo-
rous insects. Annu Rev Entomol 2011;56:161–80. https://doi.org/10.1146/annurev-ento-
120709-144753.
[15] Niederbacher B, Winkler JB, Schnitzler JP. Volatile organic compounds as non-invasive
markers for plant phenotyping. J Exp Bot 2015;66:5403–16.
https://doi.org/10.1093/jxb/erv219.
[16] Keefover-Ring K, Thompson JD, Linhart YB. Beyond six scents: defining a seventh *Thy-*
mus vulgaris chemotype new to southern France by ethanol extraction. Flavour Fragr J
2009;24:117–22. https://doi.org/10.1002/ffj.1921.
[17] Taft S, Najar A, Godbout J, Bousquet J, Erbilgin N. Variations in foliar monoterpenes
across the range of jack pine reveal three widespread chemotypes: implications to host ex-
pansion of invasive mountain pine beetle. Front Plant Sci 2015;6:342.
https://doi.org/10.3389/fpls.2015.00342.
[18] Clancy MV, Zytynska SE, Senft M, Weisser WW, Schnitzler J-P. Chemotypic variation in
terpenes emitted from storage pools influences early aphid colonisation on tansy. Sci Rep
2016;6:38087. https://doi.org/10.1038/srep38087.
[19] Bustos-Segura C, Dillon S, Keszei A, Foley WJ, Külheim C. Intraspecific diversity of ter-
penes of *Eucalyptus camaldulensis* (Myrtaceae) at a continental scale. Aust J Bot
2017;65:257–69. https://doi.org/10.1071/BT16183.
[20] Grof-Tisza P, Karban R, Rasheed MU, Saunier A, Blande JD. Risk of herbivory negatively
correlates with the diversity of volatile emissions involved in plant communication. Proc R
Soc B Biol Sci 2021;288:20211790. https://doi.org/10.1098/rspb.2021.1790.
[21] Amiot J, Salmon Y, Collin C, Thompson JD. Differential resistance to freezing and spatial
distribution in a chemically polymorphic plant *Thymus vulgaris*. Ecol Lett 2005;8:370–7.
https://doi.org/10.1111/j.1461-0248.2005.00728.x.
[22] Kleine S, Müller C. Drought stress and leaf herbivory affect root terpenoid concentrations
and growth of *Tanacetum vulgare*. J Chem Ecol 2014;40:1115–25.
https://doi.org/10.1007/s10886-014-0505-2.

[23] Linhart YB, Gauthier P, Keefover-Ring K, Thompson JD. Variable phytotoxic effects of *Thymus vulgaris* (Lamiaceae) terpenes on associated species. Int J Plant Sci 2015;176:20–30. https://doi.org/10.1086/678772.

[24] Bustos-Segura C, Külheim C, Foley W. Effects of terpene chemotypes of *Melaleuca alternifolia* on two specialist leaf beetles and susceptibility to myrtle rust. J Chem Ecol 2015;41:937–47. https://doi.org/10.1007/s10886-015-0628-0.

[25] Linhart YB, Thompson JD. Thyme is of the essence: biochemical polymorphism and multi-species deterrence. Evol Ecol Res 1999;1:151–71.

[26] Kleine S, Müller C. Intraspecific plant chemical diversity and its relation to herbivory. Oecologia 2011;166:175–86. https://doi.org/10.1007/s00442-010-1827-6.

[27] Karban R, Grof-Tisza P, Blande JD. Chemotypic variation in volatiles and herbivory for sagebrush. J Chem Ecol 2016;42:829–40.

[28] Senft M, Clancy MV, Weisser WW, Schnitzler J, Zytynska SE. Additive effects of plant chemotype, mutualistic ants and predators on aphid performance and survival. Funct Ecol 2019;33:139–51. https://doi.org/10.1111/1365-2435.13227.

[29] Karban R, Wetzel WC, Shiojiri K, Ishizaki S, Ramirez SR, Blande JD. Deciphering the language of plant communication: volatile chemotypes of sagebrush. New Phytol 2014;204:380–5. https://doi.org/10.1111/nph.12887.

[30] Halitschke R, Keßler A, Kahl J, Lorenz A, Baldwin IT. Ecophysiological comparison of direct and indirect defenses in *Nicotiana attenuata*. Oecologia 2000;124:408–17. https://doi.org/10.1007/s004420000389.

[31] Schuman MC, Heinzel N, Gaquerel E, Svatos A, Baldwin IT. Polymorphism in jasmonate signaling partially accounts for the variety of volatiles produced by *Nicotiana attenuata* plants in a native population. New Phytol 2009;183:1134–48. https://doi.org/10.1111/j.1469-8137.2009.02894.x.

[32] Hare JD. Variation in herbivore and methyl jasmonate-induced volatiles among genetic lines of *Datura wrightii*. J Chem Ecol 2007;33:2028–43. https://doi.org/10.1007/s10886-007-9375-1.

[33] Delphia CM, Rohr JR, Stephenson AG, De Moraes CM, Mescher MC. Effects of Genetic Variation and Inbreeding on Volatile Production in a Field Population of Horsenettle. Int J Plant Sci 2009;170:12–20. https://doi.org/10.1086/593039.

[34] Kariyat RR, Mauck KE, De Moraes CM, Stephenson AG, Mescher MC. Inbreeding alters volatile signalling phenotypes and influences tri-trophic interactions in horsenettle (*Solanum carolinense* L.): Inbreeding impacts plant volatiles and insect attraction. Ecol Lett 2012;15:301–9. https://doi.org/10.1111/j.1461-0248.2011.01738.x.

[35] Huang M, Abel C, Sohrabi R, Petri J, Haupt I, Cosimano J, et al. Variation of herbivore-induced volatile terpenes among Arabidopsis ecotypes depends on allelic differences and subcellular targeting of two terpene synthases, TPS02 and TPS03. Plant Physiol 2010;153:1293–310. https://doi.org/10.1104/pp.110.154864.

[36] Snoeren TAL, Kappers IF, Broekgaarden C, Mumm R, Dicke M, Bouwmeester HJ. Natural variation in herbivore-induced volatiles in Arabidopsis thaliana. J Exp Bot 2010;61:3041–56. https://doi.org/10.1093/jxb/erq127.

[37] Wason EL, Agrawal AA, Hunter MD. A genetically-based latitudinal cline in the emission of herbivore-induced plant volatile oganic compounds. J Chem Ecol 2013;39:1101–11. https://doi.org/10.1007/s10886-013-0309-9.

[38] Ghirardo A, Heller W, Fladung M, Schnitzler J-P, Schroeder H. Function of defensive vol-
atiles in pedunculate oak (*Quercus robur*) is tricked by the moth *Tortrix viridana*: Detri-
mental effect of HIPV emissions in oaks. Plant Cell Environ 2012;35:2192–207.
https://doi.org/10.1111/j.1365-3040.2012.02545.x.
[39] Rasmann S, Buri A, Gallot-Lavallée M, Joaquim J, Purcell J, Pellissier L. Differential allo-
cation and deployment of direct and indirect defences by *Vicia sepium* along elevation gra-
dients. J Ecol 2014;102:930–8. https://doi.org/10.1111/1365-2745.12253.
[40] He J, Fandino RA, Halitschke R, Luck K, Köllner TG, Murdock MH, et al. An unbiased
approach elucidates variation in (*S*)-(+)-linalool, a context-specific mediator of a tri-
trophic interaction in wild tobacco. Proc Natl Acad Sci 2019;116:14651–60.
https://doi.org/10.1073/pnas.1818585116.
[41] Xu S, Kreitzer C, McGale E, Lackus ND, Guo H, Köllner TG, et al. Allelic differences of
clustered terpene synthases contribute to correlated intraspecific variation of floral and her-
bivory-induced volatiles in a wild tobacco. New Phytol 2020;228:1083–96.
https://doi.org/10.1111/nph.16739.
[42] Hagenbucher S, Olson DM, Ruberson JR, Wäckers FL, Romeis J. Resistance mechanisms
against arthropod herbivores in cotton and their interactions with natural enemies. Crit Rev
Plant Sci 2013;32:458–82. https://doi.org/10.1080/07352689.2013.809293.
[43] Mitchell C, Brennan RM, Graham J, Karley AJ. Plant defense against herbivorous pests:
exploiting resistance and tolerance traits for sustainable crop protection. Front Plant Sci
2016;7:1132. https://doi.org/10.3389/fpls.2016.01132.
[44] Pickett JA, Khan ZR. Plant volatile-mediated signalling and its application in agriculture:
successes and challenges. New Phytol 2016;212:856–70.
https://doi.org/10.1111/nph.14274.
[45] Villamar-Torres R, Mehdi Jazayeri S, Liuba-Delfini G, García Cruzaty LC, Viot C-R. Vol-
atile organic compounds: plant natural defense mechanisms against herbivorous arthropods
and an opportunity for plant breeding of cotton. Sci Agropecu 2018;9:287–97.
https://doi.org/10.17268/sci.agropecu.2018.01.14.
[46] Elzen GW, Williams HJ, Bell AA, Stipanovic RD, Vinson SB. Quantification of volatile
terpenes of glanded and glandless *Gossypium hirsutum* L. cultivars and lines by gas chro-
matography. J Agric Food Chem 1985;33:1079–82. https://doi.org/10.1021/jf00066a015.
[47] Opitz S, Kunert G, Gershenzon J. Increased terpenoid accumulation in cotton (*Gossypium
hirsutum*) foliage is a general wound response. J Chem Ecol 2008;34:508–22.
https://doi.org/10.1007/s10886-008-9453-z.
[48] Mamin M, Vallat A, Turlings TCJ. Cotton plants as ideal models for teaching and research
on inducible direct plant defenses. Front Ecol Evol 2023;11.
https://doi.org/10.3389/fevo.2023.1119472.
[49] Loughrin JH, Manukian A, Heath RR, Turlings TC, Tumlinson JH. Diurnal cycle of emis-
sion of induced volatile terpenoids by herbivore-injured cotton plant. Proc Natl Acad Sci
1994;91:11836–40. https://doi.org/10.1073/pnas.91.25.11836.
[50] McCall PJ, Turlings TCJ, Loughrin J, Proveaux AT, Tumlinson JH. Herbivore-induced
volatile emissions from cotton (*Gossypium hirsutum L.*) seedlings. J Chem Ecol
1994;20:3039–50. https://doi.org/10.1007/BF02033709.
[51] Röse USR, Manukian A, Heath RR, Tumlinson JH. Volatile semiochemicals released from
undamaged cotton leaves (A systemic response of living plants to caterpillar damage).
Plant Physiol 1996;111:487–95. https://doi.org/10.1104/pp.111.2.487.

[52]   Paré PW, Tumlinson JH. De novo biosynthesis of volatiles induced by insect herbivory in
cotton plants 1997;114:7. https://doi.org/10.1016/S0031-9422(97)00442-1.

[53]   McAuslane HJ, Alborn HT. Systemic induction of allelochemicals in glanded and gland-
less isogenic cotton by *Spodoptera exigua* feeding. J Chem Ecol 1998;24:399–416.
https://doi.org/10.1023/A:1022596812049.

[54]   Röse USR, Tumlinson JH. Systemic induction of volatile release in cotton: How specific is
the signal to herbivory? Planta 2005;222:327–35. https://doi.org/10.1007/s00425-005-
1528-2.

[55]   Arce CM, Besomi G, Glauser G, Turlings TC. Caterpillar-induced volatile emissions in
cotton: The relative importance of damage and insect-derived factors. Front Plant Sci
2021:1503. https://doi.org/10.3389/fpls.2021.709858.

[56]   Zakir A, Sadek MM, Bengtsson M, Hansson BS, Witzgall P, Anderson P. Herbivore-in-
duced plant volatiles provide associational resistance against an ovipositing herbivore. J
Ecol 2013;101:410–7. https://doi.org/10.1111/1365-2745.12041.

[57]   Llandres AL, Almohamad R, Brévault T, Renou A, Téréta I, Jean J, et al. Plant training for
induced defense against insect pests: a promising tool for integrated pest management in
cotton: Cotton training for induced defense against pests. Pest Manag Sci 2018;74:2004–
12. https://doi.org/10.1002/ps.5039.

[58]   Briones-May Y, Quijano-Medina T, Pérez-Niño B, Benrey B, Turlings TCJ, Bustos-Se-
gura C, et al. Soil salinization disrupts plant–plant signaling effects on extra-floral nectar
induction in wild cotton. Oecologia 2023; 202:313-23. https://doi.org/10.1007/s00442-023-
05395-w.

[59]   Fernandez AR, Sáez A, Quintero C, Gleiser G, Aizen MA. Intentional and unintentional
selection during plant domestication: herbivore damage, plant defensive traits and nutri-
tional quality of fruit and seed crops. New Phytol 2021;231:1586–98.
https://doi.org/10.1111/nph.17452.

[60]   Chen YH, Gols R, Benrey B. Crop domestication and its impact on naturally selected
trophic interactions. Annu Rev Entomol 2015;60:35–58.

[61]   Yuan D, Grover CE, Hu G, Pan M, Miller ER, Conover JL, et al. Parallel and intertwining
threads of domestication in allopolyploid cotton. Adv Sci 2021;8:2003634.
https://doi.org/10.1002/advs.202003634.

[62]   Loughrin JH, Manukian A, Heath RR, Tumlinson JH. Volatiles emitted by different cotton
varieties damaged by feeding beet armyworm larvae. J Chem Ecol 1995;21:1217–27.
https://doi.org/10.1007/BF02228321.

[63]   Hagenbucher S, Birgersson G, Chattington S, Anderson P. Domestication influences
choice behavior and performance of a generalist herbivore. Perspect Plant Ecol Evol Syst
2016;23:63–72. https://doi.org/10.6084/m9.figshare. 3467747.

[64]   Shaver T, Lukefahr M. A bioassay technique for detecting resistance of cotton strains to
tobacco budworms. J Econ Entomol 1971;64:1274–7.

[65]   de Carvalho LP, Lukefhar MJ, Farias FJC, de Macedo Vieira R, Moreira J de AN, da Costa
JN. Seleção de algodoeiro com resistência ao bicudo. Pesqui Agropecuária Bras
1996;31:195–9.

[66]   Magalhães DM, Borges M, Laumann RA, Caulfield JC, Birkett MA, Blassioli-Moraes
MC. Inefficient weapon—the role of plant secondary metabolites in cotton defence against
the boll weevil. Planta 2020;252:94. https://doi.org/10.1007/s00425-020-03497-w.

[67]   Wegier A, Piñeyro-Nelson A, Alarcón J, Gálvez-Mariscal A, Álvarez-Buylla E, Piñero D.
Recent long-distance transgene flow into wild populations conforms to historical patterns

of gene flow in cotton (*Gossypium hirsutum*) at its centre of origin. Mol Ecol
2011;20:4182–94. https://doi.org/10.1111/j.1365-294X.2011.05258.x.

[68]   Coppens d'Eeckenbrugge G, Lacape J-M. Distribution and Differentiation of Wild, Feral,
and Cultivated Populations of Perennial Upland Cotton *(Gossypium hirsutum* L.) in Meso-
america and the Caribbean. PLoS ONE 2014;9:e107458. https://doi.org/10.1371/jour-
nal.pone.0107458.

[69]   Clancy MV, Mamin M, Flückiger G, Quijano-Medina T, Pérez-Niño B, Abdala-Roberts L,
et al. Terpene chemotypes in *Gossypium hirsutum* (wild cotton) from the Yucatan Penin-
sula, Mexico. Phytochemistry 2023;205:113454. https://doi.org/10.1016/j.phyto-
chem.2022.113454.

[70]   Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, et al. Fiji: an
open-source platform for biological-image analysis. Nat Methods 2012;9:676–82.
https://doi.org/10.1038/nmeth.2019.

[71]   Van Den Dool H, Kratz PD. A generalization of the retention index system including lin-
ear temperature programmed gas-liquid partition chromatography. J Chromatogr 1963.

[72]   R Core Team. R: A Language and Environment for Statistical Computing. Vienna, Aus-
tria: R Foundation for Statistical Computing; 2022.

[73]   Wickham H. ggplot2: Elegant Graphics for Data Analysis. Springer-Verlag New York;
2016.

[74]   Palarea-Albaladejo J, Martin-Fernandez JA. zCompositions – R package for multivariate
imputation of left-censored data under a compositional approach. Chemom Intell Lab Syst
2015;143:85–96.

[75]   Boogaart KG van den, Tolosana-Delgado R, Bren M. compositions: Compositional Data
Analysis. 2023.

[76]   Suzuki R, Terada Y, Shimodaira H. pvclust: Hierarchical Clustering with P-Values via
Multiscale Bootstrap Resampling. 2019.

[77]   Gu Z. Complex heatmap visualization. iMeta 2022;1:e43. https://doi.org/10.1002/imt2.43.

[78]   Oksanen J, Simpson GL, Blanchet FG, Kindt R, Legendre P, Minchin PR, et al. vegan:
Community Ecology Package. 2022.

[79]   Hervé M. RVAideMemoire: Testing and Plotting Procedures for Biostatistics. 2022.

[80]   Brooks ME, Kristensen K, Benthem KJ van, Magnusson A, Berg CW, Nielsen A, et al.
Glmmtmb balances speed and flexibility among packages for zero-inflated generalized lin-
ear mixed modeling. R J 2017;9:378–400. https://doi.org/10.3929/ethz-b-000240890.

[81]   Hartig F. DHARMa: Residual Diagnostics for Hierarchical (Multi-Level / Mixed) Regres-
sion Models. 2022.

[82]   Fox J, Weisberg S. An R Companion to Applied Regression. Third. Thousand Oaks CA:
Sage; 2019.

[83]   Lüdecke D. Ggeffects: tidy data frames of marginal effects from regression models. J Open
Source Softw 2018;3:772. https://doi.org/10.21105/joss.00772.

[84]   Lenth RV. emmeans: Estimated Marginal Means, aka Least-Squares Means. 2023.

[85]   Wilkins D. treemapify: Draw Treemaps in'ggplot2'. R package version 2.5. 2 2021.

[86]   Jenkins JN, Maxwell FG, Lafever HN. The comparative preference of insects for glanded
and glandless cottons. J Econ Entomol 1966;59:352–6.
https://doi.org/10.1093/jee/59.2.352.

[87]   Zhang L, Lu G, Huang X, Guo H, Su X, Han L, et al. Overexpression of the caryophyllene
synthase gene *GhTPS1* in cotton negatively affects multiple pests while attracting parasi-
toids. Pest Manag Sci 2020;76:1722–30. https://doi.org/10.1002/ps.5695.

[88] Elzen G, Williams H, Vinson S. Isolation and identification of cotton synomones mediat-
ing searching behavior by parasitoid *Campoletis sonorensis*. J Chem Ecol 1984;10:1251–
64. https://doi.org/10.1007/BF00988552.
[89] Elzen G, Williams H, Vinson S. Wind tunnel flight responses by hymenopterous parasitoid
Campoletis sonorensis to cotton cultivars and lines. Entomol Exp Appl 1986;42:285–9.
https://doi.org/10.1111/j.1570-7458.1986.tb01034.x.
[90] McCall PJ, Turlings TCJ, Lewis WJ, Tumlinson JH. Role of plant volatiles in host location
by the specialist parasitoid *Microplitis croceipes* cresson (Braconidae: Hymenoptera). J In-
sect Behav 1993;6:625–39. https://doi.org/10.1007/BF01048128.
[91] Rodriguez-Saona C, Crafts-Brandner SJ. Volatile emissions triggered by multiple herbi-
vore damage: beet armyworm and whitefly feeding on cotton plants. J Chem Ecol
2003;29:2539–50. https://doi.org/10.1023/a:1026314102866.
[92] Ngumbi E, Chen L, Fadamiro HY. Comparative GC-EAD responses of a specialist (*Mi-
croplitis croceipes*) and a generalist (*Cotesia marginiventris*) parasitoid to cotton volatiles
induced by two caterpillar species. J Chem Ecol 2009;35:1009–20.
https://doi.org/10.1007/s10886-009-9700-y.
[93] Harris CM, Ruberson JR, Meagher R, Tumlinson JH. Host suitability affects odor associa-
tion in *Cotesia marginiventris*: Implications in generalist parasitoid host-finding. J Chem
Ecol 2012;38:340–7. https://doi.org/10.1007/s10886-012-0095-9.
[94] Huang X, Kou J, Jing W, Han X, Liu D, Ghasemzadeh S, et al. Transcriptomic and metab-
olomic reprogramming in cotton after *Apolygus lucorum* feeding implicated in enhancing
recruitment of the parasitoid *Peristenus spretus*. J Pest Sci 2022;95:249–62.
https://doi.org/10.1007/s10340-021-01369-0.
[95] Holopainen JK. Can forest trees compensate for stress-generated growth losses by induced
production of volatile compounds? Tree Physiol 2011;31:1356–77.
https://doi.org/10.1093/treephys/tpr111.
[96] Selmar D, Kleinwächter M. Stress enhances the synthesis of secondary plant products: The
impact of stress-related over-reduction on the accumulation of natural products. Plant Cell
Physiol 2013;54:817–26. https://doi.org/10.1093/pcp/pct054.
[97] Tian Z, Luo Q, Li Y, Zuo Z. Terpinene and β-pinene acting as signaling molecules to im-
prove *Cinnamomum camphora* thermotolerance. Ind Crops Prod 2020;154:112641.
https://doi.org/10.1016/j.indcrop.2020.112641.
[98] Bertamini M, Faralli M, Varotto C, Grando MS, Cappellin L. Leaf monoterpene emission
limits photosynthetic downregulation under heat stress in field-grown grapevine. Plants
2021;10:181. https://doi.org/10.3390/plants10010181.
[99] Alavi-Samani SM, Kachouei MA, Pirbalouti AG. Growth, yield, chemical composition,
and antioxidant activity of essential oils from two thyme species under foliar application of
jasmonic acid and water deficit conditions. Hortic Environ Biotechnol 2015;56:411–20.
https://doi.org/10.1007/s13580-015-0117-y.
[100] Razavizadeh R, Komatsu S. Changes in essential oil and physiological parameters of callus
and seedlings of *Carum copticum* L. under in vitro drought stress. J Food Meas Charact
2018;12:1581–92. https://doi.org/10.1007/s11694-018-9773-9.
[101] Ruberto G, Baratta MT. Antioxidant activity of selected essential oil components in two
lipid model systems. Food Chem 2000;69:167–74. https://doi.org/10.1016/S0308-
8146(99)00247-2.

[102] De Sousa DP, Damasceno ROS, Amorati R, Elshabrawy HA, De Castro RD, Bezerra DP,
et al. Essential Oils: Chemistry and Pharmacological Activities. Biomolecules
2023;13:1144. https://doi.org/10.3390/biom13071144.
[103] Abdala-Roberts L, Quijano-Medina T, Moreira X, Vázquez-González C, Parra-Tabla V,
Berny Mier Y Terán JC, et al. Bottom-up control of geographic variation in insect her-
bivory on wild cotton (*Gossypium hirsutum*) by plant defenses and climate. Am J Bot
2019;106:1059–67. https://doi.org/10.1002/ajb2.1330.
[104] Stipanovic RD, Bell AA, O'Brien DH, Lukefahr MJ. Heliocide H1. A new insecticidal
C25 terpenoid from cotton (*Gossypium hirsutum*). J Agric Food Chem 1978;26:115–8.
[105] McAuslane HJ, Alborn HT, Toth JP. Systemic induction of terpenoid aldehydes in cotton
pigment glands by feeding of larval *Spodoptera exigua*. J Chem Ecol 1997;23:2861–79.
https://doi.org/10.1023/A:1022575313325.
[106] Gershenzon J, McConkey ME, Croteau RB. Regulation of monoterpene accumulation in
leaves of peppermint. Plant Physiol 2000;122:205–14.
https://doi.org/10.1104/pp.122.1.205.
[107] Irmisch S, Clavijo McCormick A, Boeckler GA, Schmidt A, Reichelt M, Schneider B, et
al. Two herbivore-induced cytochrome P450 enzymes CYP79D6 and CYP79D7 catalyze
the formation of volatile aldoximes involved in poplar defense. Plant Cell 2013;25:4737–
54. https://doi.org/10.1105/tpc.113.118265.
[108] Sørensen M, Neilson EHJ, Møller BL. Oximes: Unrecognized chameleons in general and
specialized plant metabolism. Mol Plant 2018;11:95–117.
https://doi.org/10.1016/j.molp.2017.12.014.
[109] Irmisch S, Clavijo McCormick A, Günther J, Schmidt A, Boeckler GA, Gershenzon J, et
al. Herbivore-induced poplar cytochrome P450 enzymes of the CYP71 family convert ald-
oximes to nitriles which repel a generalist caterpillar. Plant J 2014;80:1095–107.
https://doi.org/10.1111/tpj.12711.
[110] Clavijo McCormick A, Irmisch S, Reinecke A, Boeckler GA, Veit D, Reichelt M, et al.
Herbivore-induced volatile emission in black poplar: regulation and role in attracting her-
bivore enemies. Plant Cell Environ 2014;37:1909–23. https://doi.org/10.1111/pce.12287.
[111] McCormick AC, Reinecke A, Gershenzon J, Unsicker SB. Feeding experience affects the
behavioral response of polyphagous gypsy moth caterpillars to herbivore-induced poplar
volatiles. J Chem Ecol 2016;42:382–93. https://doi.org/10.1007/s10886-016-0698-7.
[112] Clavijo McCormick A, Gershenzon J, Unsicker SB. Little peaks with big effects: establish-
ing the role of minor plant volatiles in plant-insect interactions: Minor plant volatiles. Plant
Cell Environ 2014;37:1836–44. https://doi.org/10.1111/pce.12357.
[113] Clancy MV, Haberer G, Jud W, Niederbacher B, Niederbacher S, Senft M, et al. Under
fire-simultaneous volatilome and transcriptome analysis unravels fine-scale responses of
tansy chemotypes to dual herbivore attack. BMC Plant Biol 2020;20:551.
https://doi.org/10.1186/s12870-020-02745-1.
[114] Munguía-Rosas MA, Angulo DF, Arceo-Gómez G, Parra-Tabla V. Variation in leaf traits
across a precipitation gradient in coastal sand dunes in Yucatan Peninsula. J Arid Environ
2019;162:10–7. https://doi.org/10.1016/j.jaridenv.2018.12.001.
[115] Quijano-Medina T, Covelo F, Moreira X, Abdala-Roberts L. Compensation to simulated
insect leaf herbivory in wild cotton (*Gossypium hirsutum*): responses to multiple levels of
damage and associated traits. Plant Biol 2019;21:805–12.
https://doi.org/10.1111/plb.13002.

[116] Quijano-Medina T, Turlings TCJ, Sosenski P, Grandi L, Cervera JC, Moreira X, et al. Ef-
fects of soil salinity on the expression of direct and indirect defences in wild cotton *Gossy-*
pium hirsutum. J Ecol 2021;109:354–68. https://doi.org/10.1111/1365-2745.13483.
871

Figures

Figure 1. Illustration of the experimental process. Seeds were collected from 16 populations (red points)
in four regions along the northern coast of the Yucatán Peninsula. Plants were grown in a greenhouse until
they developed the fourth leaf, at which time half of the plants were damaged by *Spodoptera exigua* larvae
on their first and second leaves for two days and the other half was kept intact. Emitted volatiles were
collected by headspace sampling and stored leaf volatiles by hexane extraction of the undamaged fourth
leaves.

Figure 2. Chemotypic variation associated with γ-terpinene in leaf volatile profiles of wild cotton plants.
Proportion of γ-terpinene, limonene, α-thujene, terpinolene, p-cymene, and α-terpinene («γ-terpinene
group») among monoterpenes stored in wild cotton leaves of undamaged and damaged wild cotton
plants.

Figure 3. Effect of herbivory on stored leaf mono- and sesquiterpenes for both wild cotton chemotypes. Volatile compounds were extracted from the fourth leaves of 353 undamaged wild cotton plants and 322 plants damaged by *Spodoptera exigua* for two days on their first and second leaves. **A:** Effect of herbivory on the leaf concentrations of a subset of terpenes. Estimated marginal means and 95% CI from GLMMs. γ-Terpinene was analyzed within chemotype A while all other compounds were analyzed for both chemotypes together, with chemotype set as fixed factor. **B**: Biplot with symmetric scaling of a pRDA on the relative abundances (clr-transformed) of leaf mono- and sesquiterpenes. It elucidates the relationship between volatile composition and herbivory, chemotype and their interaction, while accounting for the effect of leaf area. The proximity of the points indicates the similarity in volatile composition. The table provides a summary of the permutation test results, with the percentage of variation explained by each variable (% Var.), the F-statistic (F) and the p-value (P).

Figure 4. Relative composition of the headspace volatiles emitted by undamaged and damaged wild cotton plants. **A:** Compounds detected both in the plants' leaves extracts and headspace, or in their headspace only. RI = Retention indices. The retention index for the unknown aldoxime could not be calculated because it eluted before the smallest alkane. † indicates compounds specific to damaged plants (detected in less than 10% of undamaged plants, but in between 43% and 100% of damaged plants). * indicates compounds whose identity was confirmed with standards. Distinction between the two 2-methylbutyrald-oximes isomers is tentative. FA = fatty acid derivatives; MT = monoterpenes; SQT= sesquiterpenes; Ox. = aldoximes; Ar. = aromatics; HT = homoterpenes. **B:** Composition of medians emitted volatile profiles of undamaged and damaged plants of both chemotypes. Area of the rectangles is proportional to the contribution of the compound to the profile. Numbers refer to A.

914 **Figure 5. Effect of γ-terpinene variation in leaves on headspace emissions:** correlation (Spearman) be-
915 tween the ratio of γ-terpinene and α-pinene concentrations in leaves (x-axis) and in headspace emissions
916 (y-axis).

Figure 6. Effect of herbivory on the volatiles emitted by the two wild cotton chemotypes. Headspace volatiles were sampled from 45 undamaged wild cotton plants and 48 plants damaged by *Spodoptera exigua* for two days on their first and second leaves. **A:** Biplot with symmetric scaling of a pRDA on the relative abundances (clr-transformed) of all compounds quantified in the headspace samples. Only compounds accounting for more than 10% of inertia are plotted (full biplot in Fig. S7). The plot elucidates the relationship between volatile composition and herbivory, chemotype and their interaction, while accounting for the effect of the total leaf area of the plants. The proximity of the points indicates the similarity in volatile composition. The table provides a summary of the permutation test results, with the percentage of variation explained by each variable (% Var.), the F-statistic (F) and the p-value (P). B: Effect of herbivory on emission rates of all emitted volatiles and of a subset of emitted terpenes found also in the leaf extracts. Estimated marginal means and 95% CI from GLMMs are plotted. γ-Terpinene was analyzed within chemotype A while all other compounds were analyzed for both chemotypes together, with

chemotype set as fixed factor. C: relationship between damage intensity (leaf area damaged) and the
emission rates of a subset of terpenes. Left: multiplicative increases with the leaf area damaged (back-
transformed coefficient estimates from GLMMs and their 95% CIs). The dotted line indicates a coefficient
of 1, where quantities emitted remain constant with damaged area. Right: adjusted predictions and 95%
CI from GLMMs with observed values as data points. γ-Terpinene was analyzed within chemotype A
while all other compounds were analyzed for both chemotypes together, with chemotype set as fixed
factor.

Figure 7. Variation in emitted HIPVs among geographic regions and between wild cotton chemotypes. Headspace volatile emissions were sampled from 48 wild cotton plant damaged by *Spodoptera exigua* for two days. Regions are displayed from west to east. Prevalence of chemotypes is nested with regions, as indicated by the color of data points (grey points indicate chemotype A; blue points indicate chemotype B), therefore they were analyzed separately. Estimated marginal means and 95% CI from GLMMs with observed values as data points. Damage intensity was controlled for as covariate. Letters indicate significant pairwise comparisons after multiple test adjustment (FDR). For regions, % values indicate the difference between the region with the lowest and highest estimated mean emission rates.