Induction by caterpillars of stored and emitted volatiles among populations of wild cotton (*Gossypium hirsutum*)

Short title: Plant volatiles of wild cotton chemotypes

Marine Mamin[1], Mary V Clancy[1], Galien Flückiger[1], Teresa Quijano-Medina[2], Biiniza Pérez-Niño[2], Luis Abdala-Roberts[2], Ted C.J. Turlings[1*], Carlos Bustos-Segura[1,3*]

[1] University of Neuchâtel, Institute of Biology, Fundamental and Applied Research in Chemical Ecology, Neuchâtel, Switzerland

[2] Departamento de Ecología Tropical, Campus de Ciencias Biológicas y Agropecuarias, Universidad Autónoma de Yucatán, Mérida, México

[3] Sorbonne Université, Institut National de Recherche pour l'Agriculture, l'Alimentation et l'Environnement, CNRS, Institut de Recherche pour le Développement, Université Paris-Est-Créteil-Val-de-Marne, Université Paris Cité, Institut d'Ecologie et des Sciences de l'Environnement de Paris, Versailles 78026, France

*Correspondence: ted.turlings@unine.ch; carlos.bustos@inrae.fr

ABSTRACT

Upland cotton (*Gossypium hirsutum*) plants constitutively store volatile terpenes in their leaves, which are steadily emitted at low levels. Herbivory leads to a greater release of these stored volatiles. Additionally, damaged plants increase terpene accumulation and emit other terpenes and additional volatiles. This has been well characterised for cultivated cotton, but little is known about volatile emissions in wild cotton. We investigated how herbivory by the beet armyworm (*Spodoptera exigua*) affects both leaf-stored and emitted volatiles in wild cotton plants from populations found along the Yucatan coast and assessed different scales of intraspecific variation in induced volatiles. Furthermore, we examined how leaf-terpene chemotypes differ in volatile emissions from intact and damaged plants. Herbivory triggered changes in stored and emitted volatiles in a manner similar to what is known for cultivated cotton. However, we report for the first time the emission of volatile aldoximes by cotton plants, which were only detected in the headspace upon herbivory, and displayed interpopulation variation. Intraspecific variation was also observed in the induced emissions of nitriles and some terpenes. In addition, chemotypes differed in their induction of the (*E*)-β-ocimene in leaves. Distinctions between terpene chemotypes were reflected in the emissions from both intact and damaged plants, suggesting differences in how dominant terpenes of each chemotype might affect airborne interactions. However, these variable terpenes showed weak inducibility, raising questions about their functions. This comprehensive insight into herbivore-induced volatiles in wild cotton may help in the development of future pest-management strategies for cotton crops.

Keywords: *Gossypium hirsutum*; wild cotton; herbivore-induced plant volatiles (HIPVs); intraspecific variation; chemotype; terpenes; aldoximes

INTRODUCTION

Plants emit volatiles that can play various defensive roles against insect herbivores. Volatile plant metabolites can function as toxins, repel insect feeding and oviposition, and attract natural enemies of their attackers (Unsicker et al., 2009; Turlings and Erb, 2018). Volatiles also mediate signaling within plants (Rodriguez-Saona and Frost, 2010) and may be passively used as protection, or actively perceived by neighboring plants (Karban et al., 2014b; Erb et al., 2015; Grof-Tisza et al., 2022). Un-harmed plants can produce and emit certain volatiles constitutively, a phenomenon particularly prevalent in plant species that accumulate volatile terpenes in specialized structures. These stored terpenes are continuously emitted at low levels by intact plants and are released in larger quantities when the storage organs are physically damaged, for example by chewing herbi-vores (Loreto and Schnitzler, 2010; Lange, 2015; Bustos-Segura and Foley, 2018). Additionally, when plants detect an insect attack, they may actively synthesize so-called herbivore-induced plant volatiles (HIPVs), which can include compounds that are produced constitutively but synthesized in greater amounts upon damage, or compounds whose synthesis is triggered exclusively in response to herbivory (Dicke and Baldwin, 2010; Turlings and Erb, 2018).

Several wild plant species have been shown to exhibit intraspecific variation in their chemical phenotypes. This chemodiversity often has a genetic basis and can modulate the interactions between plants and their environment (Moore et al., 2014; Kessler and Kalske, 2018; Müller and Junker, 2022). Various species that constitutively produce volatile terpenes present discrete chemical polymorphisms (chemotypes hereafter) (e.g. Keefover-Ring et al., 2009; Taft et al., 2015; Clancy et al., 2016; Bustos-Segura et al., 2017; Grof-Tisza et al., 2021). Importantly, this variation has been associated with differences in resistance to abiotic stress (Amiot et al., 2005; Kleine and Müller, 2014), allelopathy (Linhart et al., 2015), defense against pathogens (Bustos-Segura et al., 2015) and herbivores (Linhart and Thompson, 1999; Kleine and Müller, 2011; Bustos-Segura et al., 2015; Karban et al., 2016; Senft et al., 2019), as well as airborne signaling between plants when stored terpenes are released by damage (Karban et al., 2014a; Grof-Tisza et al., 2022). This variation is likely linked to genetic polymorphisms of the synthetiz-ing enzymes: for instance, variability in a terpene synthase (TPS) gene cluster underlies the constitutive chemotypes of *Mela-leuca alternifolia* (Voelker et al., 2023).

Intraspecific variation in herbivory-induced volatiles has also been described (Hare, 2011). Differences in the induction of volatile emissions have been reported across several wild species, such as: *Nicotiana attenuata* (Halitschke et al., 2000; Schu-man et al., 2009), *Datura wrightii*, (Hare, 2007), *Solanum carolinense* (Delphia et al., 2009; Kariyat et al., 2012), *Arabidopsis thaliana* (Huang et al., 2010), *Asclepias syriaca* (Wason et al., 2013), *Quercus robur* (Ghirardo et al., 2012) and *Vicia sepium* (Rasmann et al., 2014). As with constitutive volatiles, variation in induced volatiles can be linked to differences in synthetic pathways. In *N. attenuata* for instance, differences in the emission rates of the herbivore-induced terpenes linalool, (*E*)-β-ocimene and (*E*)-α-bergamotene are associated with allelic variations of *TPS* genes (He et al., 2019; Xu et al., 2020). However, fluctuations in (*E*)-α-bergamotene are also correlated with differences in the accumulation of the phytohormone jasmonic acid (JA) (Schuman et al., 2009); showing that variation in volatile induction can also be associated to differences in the en-dogenous signaling cascades that regulate the plants' response to herbivory.

Upland cotton (*Gossypium hirsutum* L. Malvaceae) is cultivated worldwide as a major source of natural fiber and suffers from high levels of herbivory from multiple insect pests (Hagenbucher et al., 2013). Increasing our understanding of the natural defense mechanisms of cotton could aid in the advancement of sustainable pest control strategies and the development of more resistant cultivars (Mitchell et al., 2016; Pickett and Khan, 2016; Turlings and Erb, 2018; Villamar-Torres et al., 2018). Cotton plants constitutively store volatile mono- and sesquiterpenes in specialized glands present in their leaf tissues (Elzen et al., 1985; Hagenbucher et al., 2013). Herbivory on older leaves induces an increase in the concentrations of these terpenes in young undamaged leaves (Opitz et al., 2008; Mamin et al., 2023). These leaf-stored terpenes are steadily emitted at low levels from intact leaves, however large quantities are immediately released following gland rupture, as caused by chewing insects (Loughrin et al., 1994; McCall et al., 1994; Röse et al., 1996; Paré and Tumlinson, 1997). Therefore, the volatile emissions instantly resulting from insect damage primarily comprise locally released leaf mono- and sesquiterpenes, together with green leaf volatiles (GLVs; aldehydes and alcohols), that are linked to tissue disruption (Loughrin et al., 1994; McCall et al., 1994). Herbivory also induces a delayed emission of mono-, sesqui- and homoterpenes, which are systematically emitted (i.e. also by undamaged parts) following *de novo* synthesis, and concurrently also triggers the emissions of other newly synthesized volatiles, such as the aromatic indole and GLV esters (Loughrin et al., 1994; McCall et al., 1994; Röse et al., 1996; Paré and Tumlinson, 1997; McAuslane and Alborn, 1998; Röse and Tumlinson, 2005; Arce et al., 2021). Thus, volatile emissions from cultivated cotton plants sustaining caterpillar herbivory mainly comprise terpenes released from constitutively stored pools at damaged sites, alongside compounds whose emissions are truly induced, with the latter coming from biosynthesis activated in response to herbivory. When active physical damaging ceases, the emission of constitutive leaf-stored terpenes drops rapidly, while induced compounds continue to be emitted (Loughrin et al., 1994). The distinction between constitutive and truly inducible terpenes is not so clear cut with some studies reporting systemic emissions of terpenes described as being constitutively stored in leaves (Röse and Tumlinson, 2005; Arce et al., 2021). Collectively, volatile emissions of insect-damaged cotton plants appear to serve as direct and indirect defenses (reviewed in Hagenbucher et al., 2013), and are involved in plant-plant interactions (Zakir et al., 2013; Llandres et al., 2018; Briones-May et al., 2023).

Given the depletion of genetic diversity observed in modern cultivars of *G. hirsutum* (Yuan et al., 2021), exploring useful defensive traits in wild cotton should be considered. While volatile production in cultivated cotton has been well characterized, less is known about the volatiles produced by wild cotton (but see Loughrin et al., 1995; Hagenbucher et al., 2016; Quijano-Medina et al., 2021; Briones-May et al., 2023). *G. hirsutum* was domesticated at least 4,000 years ago in the Yucatán Peninsula (southeast Mexico) (Yuan et al., 2021), where wild populations still grow naturally along its northern coastline (Wegier et al., 2011; Coppens d'Eeckenbrugge and Lacape, 2014). Recently, we described two heritable constitutive leaf-terpene chemotypes in wild cotton plants from the Yucatan coast: one chemotype containing none or low amounts of γ-terpinene (and other co-expressed monoterpenes), and the other exhibiting distinctively higher abundances of these compounds in their leaves (Clancy et al., 2023). It is worthwhile to note that production of γ-terpinene has been only described in wild/naturalized cotton (Loughrin et al., 1995; Hagenbucher et al., 2016). If γ-terpinene variation in leaves influences the composition of the volatile emissions of intact and/or herbivore-damaged wild cotton plants, it could impact the plants' interactions with organisms that use these volatiles as a cue, such as herbivores looking for a host plant, their natural enemies, or neighboring plants.

The present study investigated the volatile emissions of wild cotton plants, and the induction by caterpillar herbivory of foliar terpenes as well as volatiles emitted to the headspace. We determined the consistency of leaf-terpene chemotypes in the emissions of intact and damaged plants. Furthermore, we evaluated differences in the induction of stored and emitted volatiles between the two chemotypes and at different scales of intraspecific variation. To this end, we analyzed the stored volatiles in leaves and headspace volatile emissions of both undamaged plants and plants that were continuously damaged for two days by the beet armyworm (*Spodoptera exigua* (Hübner; Lepidoptera; Noctuidae)). Plants were grown in a common garden environment from seeds collected from sixteen wild cotton populations forming four distinct geographic zones distributed along the Yucatan coastline. Prevalence of the two chemotypes along the coast was previously found to follow a geographic gradient, with increasing relative abundance of the γ-terpinene chemotype from west to east (Clancy et al., 2023). We examined the variation in the increase of volatiles upon herbivory (i.e. group-by-herbivory interaction) and the variation in production rates of volatiles only produced by damaged plants (i.e. direct effect of the group of interest) between chemotypes and across the three nested levels: genotype, population and the geographic region grouping the populations. The results presented here shed new light on the production of volatiles by wild cotton plants in response to caterpillar herbivory, and uncover sources of variation, which may be genetically based, thereby opening new avenues to study and exploit cotton HIPVs as a defense mechanism against insect pests.

RESULTS

1 VOLATILE TERPENES IN LEAVES

We analyzed volatile content from the leaves of undamaged (n=353) and damaged (n=322) wild cotton plants with using hexane extraction (Fig. 1). A total of 15 monoterpenes and 18 sesquiterpenes were quantified in the leaf extracts (Table S2).

1.1 TERPENE CHEMOTYPES IN LEAVES

Undamaged and damaged plants were chemotyped based on the contribution of six strongly correlated compounds relative to the total monoterpene concentration (γ-terpinene, limonene, α-thujene, terpinolene, p-cymene, and α-terpinene, in order of abundance; *sensu* Clancy et al. 2023; Fig. S1 and S2 and Table S2). Plants containing higher proportions (> 4%) of these compounds were assigned to chemotype A; plants with less than 4% were assigned to chemotype B (Fig. S3).

1.2 EFFECT OF HERBIVORY ON THE ACCUMULATION OF VOLATILE TERPENES IN LEAVES

Following herbivory on the first and second leaves of four-leaf plants, the total leaf concentration of mono- and sesquiterpenes increased by 9% on average in the undamaged fourth leaves (Fig. 2A, $\chi^2_{(1)}$ = 12.407, p < 0.001). The same compounds were detected in the leaf terpene profiles of undamaged and damaged plants (Table S2), but a partial redundancy analysis (pRDA) revealed that herbivory affected their relative abundances (Fig. 2B, $F_{(1,670)}$ = 6.414, p = 0.001). This was mainly due to the increase in the proportion of the monoterpene (*E*)-β-ocimene upon herbivory, followed by several sesquiterpenes, notably δ-cadinene, α-copaene (Fig. 2B). The induction of (*E*)-β-ocimene depended on chemotype (Fig. 2A; chemotype x herbivory: $\chi^2_{(1)}$ = 7.694, p = 0.006), with a 64% increase in chemotype A plants (contrast of EMMs: p < 0.001) and a 108% increase in chemotype

B plants (contrast of EMMs: p < 0.001). Leaf content of the sesquiterpenes δ-cadinene and (*E*)-β-caryophyllene increased

following herbivory, by 27% (Fig. 2A, $\chi^2_{(1)}$ = 27.201, p < 0.001) and 11% (Fig. 2A: $\chi^2_{(1)}$ = 14.254, p < 0.001) respectively. In

contrast, levels of α-pinene, the most abundant monoterpene in leaves (Table S2), did not change upon herbivory (Fig. 2A,

$\chi^2_{(1)}$ = 0.437, p = 0.509). This was also the case for γ-terpinene, the main monoterpene characteristic for chemotype A (Fig. 2A,

$\chi^2_{(1)}$ = 0.766, p = 0.381). The total accumulation of (*E*)-β-ocimene increased with the amount of leaf tissue consumed by *S.*

exigua, however this effect of damage intensity was not significant for the other terpenes tested (Fig. S4: (*E*)-β-ocimene: $\chi^2_{(1)}$

= 5.619, p = 0.018).

1.2 VARIATION IN THE INDUCTION OF VOLATILE TERPENES IN LEAVES

We tested variation in induction (i.e. variation in the increase in leaf concentrations in response to herbivory; group-by-her-

bivory interaction) for the total leaf terpene content and for the aforementioned compounds between the two chemotypes

and among genotypes, populations and the four geographic regions (Table 1). As mentioned above, induction of (*E*)-β-oci-

mene in wild cotton leaves varied between the two chemotypes (Fig. 2A and Table 1; chemotype x herbivory: $\chi^2_{(1)}$ = 7.694, p

= 0.005), with a stronger induction observed in chemotype B plants. Chemotype A plants exhibited constitutively higher

amounts of (*E*)-β-ocimene in their leaves than chemotype B plants (Fig. S5, $\chi^2_{(1)}$ = 8.247, p = 0.004); this difference was no

longer significant following herbivory (Fig. S5, $\chi^2_{(1)}$ = 0.793, p = 0.373). We did not find any significant variation in induction of

any tested compounds among genotypes, populations, and regions (Table 1).

2 HEADSPACE VOLATILE EMISSIONS

The volatile emissions of undamaged and damaged wild cotton plants were collected and characterized in a subset of the

plants that were afterwards used for leaf extraction (Fig. 1; 45 undamaged and 48 damaged plants). We quantified a total of

54 compounds in the headspace (Fig. 3A and Table S3). Terpenes comprised the most diverse group, with 13 monoterpenes,

13 sesquiterpenes, and the homoterpenes DMNT and TMTT identified (Fig. 3A). Additionally, headspace emissions contained

many fatty acids derivatives, which comprised GLV aldehydes, alcohols, and esters, and (*Z*)-jasmone (Fig. 3A). Also, we iden-

tified three aliphatic aldoximes and several aromatic compounds including two nitriles. Most of the mono- and sesquiterpenes

detected in the headspace were also found in the leaf extracts, as well as a GLV aldehyde (Fig. 3A).

2.1 TERPENE CHEMOTYPES IN VOLATILE EMISSIONS

159 Of the six monoterpenes that were used to define chemotypes A and B using leaf extracts, the following compounds were

160 also detected in the headspace emissions (in order of abundance): γ-terpinene, limonene, p-cymene, and α-thujene. For both

161 undamaged and damaged plants, volatile emissions of the two chemotypes were clearly distinguishable by the prevalence of

162 γ-terpinene, p-cymene, and α-thujene (Fig. 3B and 4A), which were highly correlated (Fig. S6 and S7). The ratio of γ-terpinene

163 to α-pinene emissions was highly correlated to their ratios in the leaves (Fig. 5).

2.2 EFFECT OF HERBIVORY ON VOLATILE EMISSIONS

The volatile blend emitted by undamaged plants was mainly composed of mono- and sesquiterpenes that were also found in leaf extracts, with (*E*)-β-caryophyllene and α-pinene dominating the median profile of both chemotype A and B plants (Fig. 3B). Herbivory caused the total volatile emissions to increase 8.7-fold (Fig. 4B, $\chi^2_{(1)}$ = 146.449, p < 0.001) and triggered qualitative and quantitative changes in their composition (Fig. 3, 4A and S8). Damaged plants released several compounds that were absent from the headspace of intact plants, specifically aliphatic aldoximes, benzyl nitrile and another unidentified nitrile, the aromatics indole and methyl anthranilate, butyric and hexenoic acid esters of GLVs, (*Z*)-jasmone, the acyclic sesquiterpenes (*E*)-β-farnesene and (*E,E*)-α-farnesene, and monoterpene (*E*)-myroxide (Fig. 3). Other compounds such as the acyclic monoterpenes (*E*)-β-ocimene, linalool and (*Z*)-linalool oxide, the homoterpenes DMNT and TMTT, and the GLV ester (*Z*)-3-hexenyl acetate were emitted in low amounts by some of the undamaged plants, but their prevalence greatly increased following herbivory (Fig. 3B, 4A and S8). Indole, DMNT, (*Z*)-3-hexenyl acetate, and (*E*)-β-ocimene were major constituents of the volatile blend from damaged plants of both chemotypes (Fig. 3B). The emission rates of the mono- and sesquiterpenes that were also found in the leaf extracts generally exhibited small increases upon herbivory (Fig. 4B). However, due to greater increases of other compounds, their relative contribution to the emissions of damaged plants decreased (Fig. 3B and 4A). This was not the case of (*E*)-β-ocimene, which exhibited a distinctly high increase in response to herbivory compared to the other terpenes found in both headspace and leaves. Irrespective of chemotype ($\chi^2_{(1)}$ = 1.901, p = 0.168), the amounts emitted of (*E*)-β-ocimene rose 26-fold in response to herbivory (Fig. 4B, $\chi^2_{(1)}$ = 139.057, p < 0.001). In contrast, (*E*)-β-caryophyllene and α-pinene only increased 2.6-fold (Fig. 4B, $\chi^2_{(1)}$ = 33.800, p < 0.001) and 3.2-fold (Fig. 4B, $\chi^2_{(1)}$ = 56.859, p < 0.001) respectively. Emissions of δ-cadinene showed an intermediate increase of 11-fold (Fig. 4B, $\chi^2_{(1)}$ = 62.593, p < 0.001). A particularly low response was observed for γ-terpinene, with only a marginal increase in the headspace of 1.7-fold following herbivory (Fig. 4B, $\chi^2_{(1)}$ = 3.868, p = 0.049). Emissions of γ-terpinene were independent of damage level (Fig 4C, $\chi^2_{(1)}$ = 1.299, p = 0.254), similar to p-cymene, α-thujene, and limonene (Fig. S9). The emissions of all other terpenes detected in both headspace and leaf extracts were positively correlated with damage intensity (Fig. S9).

2.3 VARIATION IN HERBIVORY-INDUCED VOLATILE EMISSIONS

We investigated whether there were differences in the volatile emissions induced by *S. exigua* between chemotypes, as well as among regions and populations. Our analyses focused on highly inducible volatiles in order to evaluate variation in response to herbivory. The highly inducible volatiles were defined as the compounds with a negative score on the first axis of the pRDA in Figure 4A. Total emissions of these highly inducible compounds did not differ between chemotypes (Fig. S10, $\chi^2_{(2)}$ = 0.003, p = 0.954), nor among regions (Fig. S10, $\chi^2_{(3)}$ = 2.893, p = 0.408) or populations ($\chi^2_{(1)}$ = 1.344, p = 0.246). However, the pRDA revealed differences between chemotypes (Fig. S11, $F_{(1,42)}$ = 1.189, p = 0.009) and among regions (Fig. S12, $F_{(3,38)}$ = 1.904, p = 0.005) in the composition of their HIPV emissions. Of the compounds highlighted by the pRDAs, the terpenes (*E,E*)-α-farnesene, (*E*)-myroxide, the three aldoximes and the two nitriles showed statistically significant differences in the emission rates (Table 2). These compounds were all only detected in the headspace of damaged plants. We also observed variations in the emissions of TMTT and δ-cadinene (Fig. S11 and S12); these volatiles were also found in undamaged plants but in lower amounts. The absence of significant interactions between region/chemotype and herbivory suggests that the observed differences might not be specifically linked to a distinct response to herbivory (Fig. S11 and S12). Emission rates of (*E,E*)-α-farnesene

and (*E*)-myroxide varied among regions, as well as among populations, but not between chemotypes (Table 2, Fig. 6 and Fig. S10). The three aldoximes, which were strongly correlated (Fig. S7), varied both among regions and between chemotypes. Plants from Celestún (far west) and Sisal (west) produced higher quantities of aldoximes compared to plants from regions further east (Table 2, Fig. 6 and Fig. S10). Chemotype B plants released higher amounts of these compounds than chemotype A plants (Table 2, Fig. 6 and Fig. S10). Thus, variation could stem both from the chemotype and the geographic region, which are partially nested. However, even when taking into account the effect of chemotype, significant variation was observed at the population level for the two methylbutyraldoximes (Table 2). Benzyl nitrile showed the same trend in variation as the aldoximes, while the opposite was observed for the unknown nitrile (Table 2, Fig. 6). For these latter two compounds, however, no evidence of variation among population was observed (Table 2).

DISCUSSION

In this study, we investigated the effect of caterpillar herbivory on quantitative and qualitative changes in leaf-stored and emitted volatiles of wild cotton plants from 16 population along the northern coast of the Yucatán Peninsula. Moreover, we show that two previously described chemotypes found within these populations could also be distinguished based on their headspace emissions, which was true for both undamaged and caterpillar-damaged plants. Responses of wild cotton plants to herbivory were found to differ between chemotypes in the induction of (*E*)-β-ocimene in leaves; for the emission of several HIPV chemical classes there were differences between chemotypes and among regions and/or populations.

1 INDUCTION OF VOLATILE MONO- AND SESQUITERPENES IN LEAVES

Two days of herbivory on the first and second leaves led to a rapid accumulation of several volatile terpenes in the undamaged fourth leaf of the same plants. The leaf terpenes differed in their inducibility, which resulted in compositional changes in leaf volatile profiles triggered by herbivory. A similar variation in inducibility among compounds has previously been described for domesticated varieties, with (*E*)-β-ocimene being the most responsive to damage, and with sesquiterpenes exhibiting a stronger response than monoterpenes (Opitz et al., 2008; Mamin et al., 2023). The higher inducibility of (*E*)-β-ocimene may allow for the increased synthesis of heliocides H1 and H4, the most inducible gland-stored, non-volatile defensive terpenoid aldehydes produced by cotton, for which (*E*)-β-ocimene is a precursor, together with hemigossypolone (Stipanovic et al., 1978; McAuslane et al., 1997; Opitz et al., 2008).

2 INDUCTION OF VOLATILE EMISSIONS

The constitutive volatile emissions of wild cotton consisted mostly of terpenes that were also found accumulated in the leaves: several cyclic terpenes, along with the linear terpenes β-myrcene and (*E*)-β-ocimene. This parallels previous descriptions of cultivated varieties, where these compounds are known to be stored in glands and constitutively released in small amounts (Elzen et al., 1985; Loughrin et al., 1994; Paré and Tumlinson, 1997). For most of these terpenes, we found a small increase in their emission rates following herbivory. As there was no active damage occurring on the plant during sampling, this increase could reflect passive release from storage pools in induced undamaged leaves, although we cannot exclude some leakage

from very fresh wounds, nor can a release from sources other than storage pools be fully ruled out (Gershenzon et al., 2000;

Bustos-Segura and Foley, 2018). The latter is already known for (*E*)-β-ocimene in cultivated cotton, which is not only accumu-

lated in the leaves but is also *de novo* synthesized and systemically emitted upon induction (Paré and Tumlinson, 1997). Con-

sistent with this previously observed phenomenon, we found a particularly strong induction of (*E*)-β-ocimene emissions in

comparison to the other terpenes detected in both leaves and headspace. Together with (*E*)-β-ocimene, the major compounds

induced by *S. exigua* herbivory in wild cotton plants included indole, (*Z*)-3-hexenyl acetate, and several terpenes that were

distinct from those found in stored pools, namely the linear terpenoids linalool, (*E*)-β-farnesene, (*E,E*)-α-farnesene, DMNT,

and TMTT. This aligns with previous findings in cultivated cotton, showing that these compounds are newly synthesized fol-

lowing damage (Loughrin et al., 1994; McCall et al., 1994; Röse et al., 1996; Paré and Tumlinson, 1997; McAuslane and Alborn,

1998; Arce et al., 2021).

We identified several compounds that to the best of our knowledge have not yet been reported for cotton, including aliphatic

aldoximes, nitriles, and the aromatic compound 2-phenylethanol. A nitrile compound previously described in wild cotton from

Yucatán (Briones-May et al., 2023) likely corresponds to the nitrile we could not conclusively identify. Aldoximes are nitroge-

nous compounds synthesized from amino acids by cytochrome P450 monooxygenases of the CYP79 family, found across all

angiosperms, and are precursors to nitriles (Irmisch et al., 2013; Sørensen et al., 2018). Aldoximes and nitriles may be precur-

sors to volatile alcohols, such as 2-phenylethanol (Irmisch et al., 2014; Sørensen et al., 2018). Various plant species, including

several Fabaceae*,* Populus, and Erythroxylum species, *Solanum melongena*, *Zea mays, Cucumis sativus*, and *Gerbera jamesonii*

emit volatile aliphatic aldoximes in response to herbivory or jasmonic acid treatment (Sørensen et al., 2018). In *Populus tricho-*

carpa, herbivore-induced CYP79 enzymes produce (*E/Z*)-2-methylbutyraldoximes from L-isoleucine in response to caterpillar

herbivory (Irmisch et al., 2013), and CYP71 enzymes convert aldoximes into nitriles, including benzyl nitrile, from the semi-

volatile compound phenylacetaldoxime (Irmisch et al., 2014). Volatile aldoximes and nitriles could be involved in cotton indi-

rect defenses; in *Populus nigra*, methylbutyraldoximes and benzyl nitrile are central to the recruitment of parasitoid wasps,

despite being only minor components of the full induced volatile blend (Clavijo McCormick et al., 2014b). Benzyl nitrile is also

involved in direct defense mechanisms of *Populus trichocarpa and P. nigra* (Irmisch et al., 2014; McCormick et al., 2016).

Aldoximes and nitriles might be emitted in small amounts by cultivated *G. hirsutum*, and therefore more difficult to detect

with classic sampling and analyses (Clavijo McCormick et al., 2014a). Our results seem to imply wild cotton plants produce

more aldoximes and nitriles than cultivated varieties. This might be of interest for future studies as it could offer an oppor-

tunity to enhance the resistance of cultivated cotton varieties to herbivorous pests.

3 CHEMOTYPES AND THEIR VOLATILE TERPENE EMISSIONS

The two previously reported leaf-terpene chemotypes (Clancy et al., 2023) were also distinct in their headspace emissions, as

the variation in the leaf content of γ-terpinene and its correlated compounds was mirrored in the emissions from both un-

damaged and damaged plants. This suggests that, in both cases, the primary source of emission for these compounds are the

leaf storage pools, and that the chemotypes of wild cotton plants could differ in their airborne interactions with their environ-

ment. Volatile terpenes likely released from stored pools have been found to repel adult moths and piercing-sucking pests of

cultivated cotton (Jenkins et al., 1966; Zhang et al., 2020), and appear to function as foraging cues for some species of parasi-

toid wasps (Elzen et al., 1984, 1986; McCall et al., 1993; Zhang et al., 2020). Additional research is needed to evaluate how the

two chemotypes may impact wild cotton's interactions with its insect herbivores and their natural enemies. Our results sug-

gest that their role in this context may be of lesser importance, as γ-terpinene exhibited a notably low inducibility in compar-

ison to other monoterpenes. Measurements at additional time points would be required to confirm this, but it appears that

the TPSs associated with γ-terpinene synthesis are not responsive to caterpillar herbivory, suggesting that this compound is

not involved in defense against *S. exigua*. As induction of volatiles in cotton can be specific to the species of an attacking

herbivore (Rodriguez-Saona and Crafts-Brandner, 2003; Ngumbi et al., 2009; Harris et al., 2012; Huang et al., 2022), we rec-

ommend that future work investigates the response of wild cotton chemotypes to other herbivore species.

Increased production of monoterpenes has also been reported in several plant species in response to abiotic stresses such as

drought and excessive temperatures, against which they may offer protective effects (Loreto and Schnitzler, 2010; Holopainen,

2011; Selmar and Kleinwächter, 2013; Tian et al., 2020; Bertamini et al., 2021). Interestingly, some plant species specifically

increase γ-terpinene concentrations in their tissues under drought conditions (Alavi-Samani et al., 2015; Razavizadeh and

Komatsu, 2018), and it has been demonstrated that γ-terpinene has a particularly high antioxidant capacity (Ruberto and

Baratta, 2000; De Sousa et al., 2023). Previous work on Yucatanese wild cotton documented that populations in higher-tem-

perature environments tend to exhibit a greater density of glands (albeit non-significantly), while herbivory pressure and gland

density are not correlated among populations (Abdala-Roberts et al., 2019). This finding suggests that abiotic conditions could

potentially be significant factors driving variation in gland density and their chemical content. Considering the particularly

harsh environments in which wild cotton grows, it would be interesting to investigate the relationship between γ-terpinene

content and tolerance to extreme environmental conditions.

286 4 VARIATION IN THE INDUCTION OF LEAF-STORED VOLATILES AND IN HIPV EMISSIONS

In terms of leaf volatile content, the two chemotypes varied in their inducibility (magnitude of the increase in leaf concentra-

tions) of the monoterpene (*E*)-β-ocimene, whereas there was no difference between chemotypes in the induction of (*E*)-β-

ocimene in the headspace emissions. This is likely due to the possibility, as mentioned earlier, that most of its induced emis-

sions may derive from *de novo* synthesis rather than stored pools (Paré and Tumlinson, 1997). Nevertheless, this variation in

the inducibility of (*E*)-β-ocimene in leaves could have implications for wild cotton plants' airborne interactions, particularly in

scenarios involving subsequent leaf damage, as it would lead to the release of the induced stored pools (Röse et al., 1996;

McAuslane and Alborn, 1998). As for headspace HIPV emissions, we observed considerable intraspecific variation at the pop-

ulation and region levels in the emissions of the terpenes (*E,E*)-α-farnesene and (*E*)-myroxide, as well as in the emissions of

aldoximes. Nitriles were also found to differ, to a lesser extent, among regions. Aldoximes and nitriles were also observed to

vary between the two leaf-terpene chemotypes. As the prevalence of the two chemotypes was partially nested within region,

the source of variation in emissions of these compounds cannot be conclusively ascertained. Having said this, the similarity

between damaged plants from Chicxulub (high proportion of chemotype B) and Coloradas (dominance of chemotype A) in the

production of both classes of compounds suggests that this variation is more likely linked to interpopulation differences than

to the chemotypes themselves. In contrast, chemotype-specific responses to caterpillar herbivory have been described in

another species (*Tanacetum vulgare*) for both leaf-stored and emitted volatiles (Clancy et al., 2020). Despite being confined

to coastal habitats, Yucatanese wild cotton is subjected to substantial spatial variability in herbivore pressure and abiotic

conditions (Abdala-Roberts et al., 2019; Munguía-Rosas et al., 2019; Quijano-Medina et al., 2019, 2021), which could be asso-

ciated with the origin of the observed variation in HIPVs. Further work is required to understand the drivers of these differ-

ences. Given the multiple proposed functions of HIPVs in plant defense against insect herbivores (Turlings and Erb, 2018), the

differences in HIPVs that we observed could have consequences for wild cotton's resistance to biotic and abiotic stresses,

which is worthy of further exploration. Genotypes producing higher amounts of various HIPVs that improve their direct or

indirect defenses or those of neighboring plants could be a valuable genetic resource in the development of more resistant

cotton cultivars.

MATERIALS AND METHODS

1 PLANTS AND INSECTS

Wild *Gossypium hirsutum* seeds were collected in 2019 and 2020 on the northern coast of the Yucatán Peninsula (Mexico) and

grown as described in Clancy et al. (2023). Seeds were sampled from several mother plants from 16 populations in four regions

along the coast (Table S1 and Fig. 1). As cotton cross-pollinates, seeds from the same mother are either full or half-siblings, of

unknown paternal source. Offspring from the same mother plant is referred to here as genotype. Plants were grown in a

greenhouse for around four weeks. Eventually, we obtained plants from one to six mother plants for each population, repre-

senting around 35 to 60 individuals per population, except for a few populations with particularly low germination rates (Table

S1). To induce volatile production in the cotton plants we used larvae of the moth *Spodoptera exigua* (Hübner; Lepidoptera;

Noctuidae), a cosmopolitan generalist pest that feeds on cotton. Eggs were supplied by Entocare (Wageningen, The Nether-

lands) and reared in the laboratory on wheat germ-based artificial diet (Frontier Scientific Services, Newark, USA).

2 EXPERIMENTAL PROCESS AND SAMPLE COLLECTION

Plants were used when the fourth leaf was fully expanded but still developing. A total of 322 plants were subjected to her-

bivory and 353 were left undamaged, in six experimental batches. The numbers of replicates per region and population are

described in Table S1. Second to early third-instar *S. exigua* larvae fed on the first and second leaves for two days, with a single

larva enclosed on each leaf in clip cages (4.52 cm^2, BioQuip product, USA). Undamaged and damaged plants were kept in two

adjacent rooms with similar configurations and conditions (L14:D10, T=28°C), to prevent any signaling effect of HIPVs. The

treatments were alternated between rooms after each batch. After two days of herbivory, we collected the volatile emissions

of a subset of the plants. A total of 45 undamaged and 48 damaged individuals were used in four batches for headspace

collection, chosen to represent the various populations (Table S1). Afterwards, the fourth leaf of all plants was harvested,

photographed, frozen in liquid nitrogen, and stored at -80 °C until processed. Photographs of all leaves were taken to quantify

the total leaf area consumed by *S. exigua*, as well as to determine the area (size) of the fourth leaves and the total leaf area

(including cotyledons and secondary leaves) of the plants used for the headspace collections. For this purpose, we used the

Fiji distribution of ImageJ software (Schindelin et al., 2012).

3 HEADSPACE VOLATILE COLLECTION AND GC-MS ANALYSES

S. exigua larvae were removed before headspace sampling, to avoid the release of volatiles due to physical tissue disruption

by active feeding. The caterpillars' frass was removed from the leaves. Pots were covered with aluminum foil to limit potential

emissions from the soil. Headspace volatiles were collected in a push-pull air system of custom-made glass bottles (VTQ SA,

Switzerland). Charcoal-purified and humidified air was pushed at 1 L min^{-1} and pulled at 0.6 L min^{-1} through trapping filters (25

mg of 80/100 mesh Hayesep-Q adsorbent, Sigma, Switzerland) for two hours. Filters were eluted with 100 µL of dichloro-

methane, with the addition of 10 µL of internal standard solution (nonyl-acetate, 20 ng $µL^{-1}$). Samples were stored at -80 °C

before being run on a gas chromatograph (GC Agilent 7890B) coupled to a mass spectrometer (MSD Agilent 5977B): 1.5 µL

were autoinjected in splitless mode onto an Agilent HP-5MS column (30 m × 250 µm × 0.25 µm). The temperature was kept

at 40 °C for 3.5 min, then ramped to 100 °C at 8 °C min^{-1} and to 230 °C at 5 °C min^{-1}. Helium was used as the carrier gas, with

a stable flow of 0.9 mL min^{-1}.

4 HEXANE LEAF EXTRACTIONS AND GC-MS ANALYSES

Hexane leaf extractions and subsequent analyzes were performed as described in Clancy et al. (2023). Briefly, 100 mg of

grounded frozen leaf material was extracted with 1 mL of hexane and kept at 4 °C for 24 hours. We recovered 800 µL and

added 8 µL of internal standard solution (nonyl acetate, 2000 ng $µL^{-1}$). The samples were directly injected into a gas chromato-

graph (GC Agilent 8890A) coupled to a mass spectrometer (MSD Agilent 5977B): 1.5 µL were injected in splitless mode onto

an Agilent HP-5MS column (30 m × 250 µm × 0.25 µm). Temperature started at 40°C for 0 min, ramped to 130 °C at 15 °C min^{-}

^{1}, then to 190 °C at 60 °C min^{-1}, and to 290 °C at 30 °C min^{-1}, then held for 0.5 min. Helium was used as carrier gas, with a stable

flow of 0.9 mL $min^{-1.}$

5 GC-MS DATA PROCESSING

PyMS (O'Callaghan et al., 2012) was used to integrate and align peaks from the total ion chromatograms of all samples

(https://github.com/bustossc/PyMS-routine/). Analytes were quantified using relative response factors obtained from a series

of external calibration curves of commercial analytical standards in comparison to nonyl acetate (Table S2 and S3). For com-

pounds lacking available response factors, we employed the relative response factor associated with the closest standard in

retention time and similar chemical structure, while normalizing by molecular masses (Kreuzwieser et al. 2014). When possi-

ble, analytes were identified based on comparison with commercial standards. Aldoximes were identified with synthesized

standards kindly provided by Prof. Tobias Köllner. Compounds with no standard available were identified based on mass

spectra comparison with NIST 2.3 mass spectral library, an in-house library, and Kovats retention indices (Van den Dool and

Kratz, 1963) (Table S2 and S3). Components retained in the analyzes were present in at least 10% of the samples. For the

hexane extracts, values were normalized by the fresh weights of the leaves (units: ng mg^{-1} FW). For headspace samples, values

were normalized by hours of collection and the total leaf area of the plants (units: ng h^{-1} cm^{-2}).

STATISTICAL ANALYZES

366 Statistical analyzes were carried out with R 4.2.3 (R Core Team, 2022). A significance level of $\alpha = 0.05$ was used for all analyzes.

367 Plots were built with "ggplot2" (v. 3.3.6, Wickham, 2016).

1 MULTIVARIATE ANALYZES OF VOLATILES COMPOSITION: PRDAS AND CLUSTERING

Multivariate analyzes were always performed on the proportions of compounds relative to the sum of all analytes of interest.

To deal with the left-censored and compositional nature of the data, zeroes were imputed by multiplicative simple replace-

ment with multRepl() from "zCompositions" (v. 1.4.0.1, Palarea-Albaladejo and Martin-Fernandez, 2015), using the smallest

non-zero value of each compound as detection limits and a delta of 0.65. We subsequently employed a centered log-ratio

(clr) transformation with clr() from "compositions" (v. 2.0.5, Boogaart et al., 2023). Data were finally scaled and centered with

scale() from the R base package. To evaluate correlations among compounds, hierarchical clustering was performed on the

correlation-based (Pearson) dissimilarity matrix with Ward's minimum variance method (Ward.D2) with pvclust() from

"pvclust" (v. 2.2.0, Suzuki et al., 2019). Approximately Unbiased p-values were computed by multiscale bootstrap resampling

with 10,000 replications. Correlations heatmaps were plotted with "ComplexHeatmap" (v. 2.16.0, Gu. Z., 2022). Partial redun-

dancy analyzes (pRDAs) were conducted with rda() from "vegan" (v. 2.6.4 Oksanen et al., 2022). Models' structures are de-

scribed in the following paragraphs. We then performed a type II permutation test with MVA.anova() from "RVAideMemoire"

(v. 0.9.81.2, Hervé, 2022). Pairwise comparisons were performed with pairwise.factorfit() from the same package.

2 ANALYZES OF SINGLE COMPOUNDS: GLMMS

Univariate analyzes of concentrations of compounds in leaves or emission rates were performed with generalized linear mixed

models (GLMMs) using glmmTMB (v. 1.1.5, Brooks et al., 2017) with Gamma or Tweedie exponential families of distribution,

depending on models' fit. Models' fits were checked with the package "DHARMa" (v. 0.4.6, Hartig, 2022). Effects of the ex-

planatory variables were tested with Wald chi-squared tests (type-II) using Anova() from "car" (v. 3.1.1, Fox and Weisberg,

2019). We plotted estimated marginal means and their 95% confidence intervals (CI), extracted from the models with ggem-

means() from "ggeffects" (v. 1.2.0, Lüdecke, 2018). When plotted, data points represented observed values. In all models, we

controlled for the effects of plants "size" (leaf extracts: fourth leaf area; headspace emissions: total leaf area) and of experi-

mental batches (leaf extracts: n = 6; headspace emissions: n = 4) by including them as fixed effects (Table 3 and 4). γ-Terpinene

was analyzed within the producing chemotype only, as the other chemotype does not produce it, or only in very low amounts.

Population and genotype were included as random intercepts in all models, except for those relying solely on the headspace

emissions of damaged plants, where 86% of the data consisted of individual data points per genotype (Table 3 and 4). When

necessary, pairwise comparisons were performed with emmeans() from "emmeans", with multiple test adjustment (FDR). (v.
1.8.7, Lenth, 2023).

3 VOLATILE TERPENES IN LEAVES: HERBIVORY AND VARIATION IN INDUCTION

To assess how herbivory affected the leaf composition in volatile terpenes for the two chemotypes, a pRDA was performed
on the relative abundances of all leaf mono- and sesquiterpenes. Herbivory, chemotype and their interaction were used as
explanatory variables while leaf area was partialled out. Then, we analyzed the effect of herbivory and damage intensity (total
leaf area consumed by caterpillars) on the total concentrations of the leaf volatile terpenes and a subset of compounds using
GLMMs (Table 3: model structures A and E). We tested whether there was variation in the induction (i.e. increase in the leaf
concentrations; group-by-herbivory interaction) of volatile terpenes between the two chemotypes and among nested levels
of intraspecific variation: 88 genotypes, 16 populations and four geographic regions. Variation in the response of genotypes
and populations to herbivory was modelled with random slopes and their significance was tested by comparing nested models
with a random slope component to a model without (Restricted Maximum Likelihood method) with Likelihood ratio tests.
Chemotype was included in the fixed effects of these models (Table 3: model structures B and C). The random slope compo-
nent included a term for the variance in slopes and one for the covariance between slopes and intercepts unless the model
did not converge. In that case, the covariance term was removed. Variation in response to herbivory among region and chemo-
types was modelled as interaction between these groups and herbivory, set as fixed effects). Because of their geographic
distribution, chemotypes were partially nested in regions. Therefore, to avoid collinearity, they were always analyzed sepa-
rately (Table 3: model structures A and D).

4 HEADSPACE VOLATILE EMISSIONS: HERBIVORY AND VARIATION IN INDUCED COMPOUNDS

To examine whether the intraspecific variation observed for the leaf-stored terpenes by Clancy et al. (2023) was translated to
the emissions, we performed Spearman correlation tests, separately for undamaged and damaged plants, on the ratio of
concentrations of γ-terpinene and α-pinene (the dominant leaf-stored monoterpene) in leaves and in headspace emissions.
To visualize the composition of the emitted blends of undamaged and damaged plants of both chemotypes, we plotted the
median proportions with "treemapify" (v. 2.5.5, Wilkins 2021). To further understand how herbivory and chemotypes affected
the composition of headspace emissions a pRDA was conducted on the relative abundances of all emitted volatiles together.
Herbivory, chemotype and their interaction were used as explanatory variables while the total leaf area was partialled out.
Then, we analyzed the effect of herbivory and damage intensity (total leaf area consumed by caterpillars) on the total emission
rates and on the emissions of same terpenes analyzed for the leaves (Table 4: model structures A and C).

Variation in the volatile emissions induced by caterpillar herbivory was evaluated between chemotypes, as well as two levels
of intraspecific variation: among the four regions and 16 populations. Genotypes were not assessed because of the lack of
replication at this level. To evaluate variation in the volatile response to herbivory rather than pre-existing differences in com-
pounds already emitted by intact plants, we selected only highly inducible volatiles, defined from the pRDA assessing the
effect of herbivory on the headspace emissions. We first evaluated whether there was variation in the total emissions of these
highly inducible compounds (Table 4, model structures C, D and E). Variation among population was modelled through random

intercepts (Table 4: model structures E) and their significance was tested by comparing nested models with random intercept component to models without (Restricted Maximum Likelihood method) with Likelihood ratio tests. Chemotypes and regions were analyzed as fixed effects, and separately, as mentioned above. To unveil potential variation in the composition of the induced emissions, pRDAs were performed on the proportions of highly inducible volatiles, selected from the transformed and scaled matrix of the relative abundances of all compounds emitted by damaged plants, separately for chemotype and region. Chemotype/region, the leaf surface damaged and their interaction were used as explanatory variables while the total leaf area was partialled out. We then further tested whether chemotypes and regions varied significantly in their emission rates of the different candidate compounds highlighted by the pRDAs with GLMMs, as explained for the total induced emissions (Table 4, model structure C and D). Compounds with significant variations were also assessed for variation among populations (Table 4, model structure E). For inducible compounds differing between chemotypes or regions but also emitted by undamaged plants, the interaction effect between chemotype/region and herbivory was tested (Table 4, model structure A and B). A significant interaction implied that chemotypes/regions differed in their response to herbivory, not solely by pre-existing disparities.

FUNDING INFORMATION

This work was supported by the Swiss National Science Foundation [grant 315230_185319]. Permit information for sampling of plant material in Yucatán: SEMARNAT- SCPA/DGVS/00585/20 SEMARNAT SCPA/DGVS/2974/20.

ACKNOWLEDGMENTS

The authors thank Thomas Degen (UniNE) for the illustrations of cotton and for helping to build figure 1, Karla Ku Durán (UADY) laboratory support, Dominic Davis-Foster (Staffordshire U) for maintaining the PyMassSpec repository, Tobias Köllner (Max Planck Institute for Chemical Ecology) for the aldoximes standards, and Fanny Deiss, Sara Leite Dias, Jonathan Interian, and Miguel Cárdenas for help with sample collection.

AUTHOR CONTRIBUTIONS

T.C.J.T. and L.A-R., C.B-S., M.V.C. and M.M. designed the research. L.A-R., T.Q-M. and B.P.-N. collected the cotton seeds. C.B-S., M.V.C., M.M. and G.F. performed the experiment and processed the samples. M.M. and C.B-S. processed and analyzed the data. M.M. wrote the first draft of the manuscript with contributions by all authors.

DECLARATION OF COMPETING INTEREST

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

DATA AVAILABILITY

Data and code can be found at: https://doi.org/10.5281/zenodo.10927508

REFERENCES

Abdala-Roberts, L., Quijano-Medina, T., Moreira, X., Vázquez-González, C., Parra-Tabla, V., Berny Mier Y Terán, J. C., et al. (2019). Bottom-up control of geographic variation in insect herbivory on wild cotton (*Gossypium hirsutum*) by plant defenses and climate. *Am. J. Bot.* 106, 1059–1067. doi: 10.1002/ajb2.1330

Alavi-Samani, S. M., Kachouei, M. A., and Pirbalouti, A. G. (2015). Growth, yield, chemical composition, and antioxidant activity of essential oils from two thyme species under foliar application of jasmonic acid and water deficit conditions. *Hortic. Environ. Biotechnol.* 56, 411–420. doi: 10.1007/s13580-015-0117-y

Amiot, J., Salmon, Y., Collin, C., and Thompson, J. D. (2005). Differential resistance to freezing and spatial distribution in a chemically polymorphic plant *Thymus vulgaris*. *Ecol. Lett.* 8, 370–377. doi: 10.1111/j.1461-0248.2005.00728.x

Arce, C. M., Besomi, G., Glauser, G., and Turlings, T. C. (2021). Caterpillar-induced volatile emissions in cotton: The relative importance of damage and insect-derived factors. *Front. Plant Sci.*, 1503. doi: 10.3389/fpls.2021.709858

Bertamini, M., Faralli, M., Varotto, C., Grando, M. S., and Cappellin, L. (2021). Leaf monoterpene emission limits photosynthetic downregulation under heat stress in field-grown grapevine. *Plants* 10, 181. doi: 10.3390/plants10010181

Boogaart, K. G. van den, Tolosana-Delgado, R., and Bren, M. (2023). *compositions: Compositional Data Analysis*. Available at: https://CRAN.R-project.org/package=compositions

Briones-May, Y., Quijano-Medina, T., Pérez-Niño, B., Benrey, B., Turlings, T. C. J., Bustos-Segura, C., et al. (2023). Soil salinization disrupts plant–plant signaling effects on extra-floral nectar induction in wild cotton. *Oecologia*. doi: 10.1007/s00442-023-05395-w

Bustos-Segura, C., Dillon, S., Keszei, A., Foley, W. J., and Külheim, C. (2017). Intraspecific diversity of terpenes of *Eucalyptus camaldulensis* (Myrtaceae) at a continental scale. *Aust. J. Bot.* 65, 257–269. doi: 10.1071/BT16183

Bustos-Segura, C., and Foley, W. J. (2018). Foliar terpene chemotypes and herbivory determine variation in plant volatile emissions. *J. Chem. Ecol.* 44, 51–61. doi: 10.1007/s10886-017-0919-8

Bustos-Segura, C., Külheim, C., and Foley, W. (2015). Effects of terpene chemotypes of *Melaleuca alternifolia* on two specialist leaf beetles and susceptibility to myrtle rust. *J. Chem. Ecol.* 41, 937–947. doi: 10.1007/s10886-015-0628-0

Clancy, M. V., Haberer, G., Jud, W., Niederbacher, B., Niederbacher, S., Senft, M., et al. (2020). Under fire-simultaneous volatilome and transcriptome analysis unravels fine-scale responses of tansy chemotypes to dual herbivore attack. *BMC Plant Biol.* 20, 551. doi: 10.1186/s12870-020-02745-1

Clancy, M. V., Mamin, M., Flückiger, G., Quijano-Medina, T., Pérez-Niño, B., Abdala-Roberts, L., et al. (2023). Terpene chemotypes in *Gossypium hirsutum* (wild cotton) from the Yucatan Peninsula, Mexico. *Phytochemistry* 205, 113454. doi: 10.1016/j.phytochem.2022.113454

Clancy, M. V., Zytynska, S. E., Senft, M., Weisser, W. W., and Schnitzler, J.-P. (2016). Chemotypic variation in terpenes emitted from storage pools influences early aphid colonisation on tansy. *Sci. Rep.* 6, 38087. doi: 10.1038/srep38087

Clavijo McCormick, A., Gershenzon, J., and Unsicker, S. B. (2014a). Little peaks with big effects: establishing the role of minor
plant volatiles in plant-insect interactions: Minor plant volatiles. *Plant Cell Environ.* 37, 1836–1844. doi:
10.1111/pce.12357

Clavijo McCormick, A., Irmisch, S., Reinecke, A., Boeckler, G. A., Veit, D., Reichelt, M., et al. (2014b). Herbivore-induced volatile
emission in black poplar: regulation and role in attracting herbivore enemies. *Plant Cell Environ.* 37, 1909–1923. doi:
10.1111/pce.12287

Coppens d'Eeckenbrugge, G., and Lacape, J.-M. (2014). Distribution and Differentiation of Wild, Feral, and Cultivated Popula-
tions of Perennial Upland Cotton *(Gossypium hirsutum* L.) in Mesoamerica and the Caribbean. *PLoS ONE* 9, e107458.
doi: 10.1371/journal.pone.0107458

De Sousa, D. P., Damasceno, R. O. S., Amorati, R., Elshabrawy, H. A., De Castro, R. D., Bezerra, D. P., et al. (2023). Essential Oils:
Chemistry and Pharmacological Activities. *Biomolecules* 13, 1144. doi: 10.3390/biom13071144

Delphia, C. M., Rohr, J. R., Stephenson, A. G., De Moraes, C. M., and Mescher, M. C. (2009). Effects of Genetic Variation and
Inbreeding on Volatile Production in a Field Population of Horsenettle. *Int. J. Plant Sci.* 170, 12–20. doi:
10.1086/593039

Dicke, M., and Baldwin, I. T. (2010). The evolutionary context for herbivore-induced plant volatiles: beyond the 'cry for help.'
Trends Plant Sci. 15, 167–175. doi: 10.1016/j.tplants.2009.12.002

Elzen, G. W., Williams, H. J., Bell, A. A., Stipanovic, R. D., and Vinson, S. B. (1985). Quantification of volatile terpenes of glanded
and glandless *Gossypium hirsutum* L. cultivars and lines by gas chromatography. *J. Agric. Food Chem.* 33, 1079–1082.
doi: 10.1021/jf00066a015

Elzen, G., Williams, H., and Vinson, S. (1984). Isolation and identification of cotton synomones mediating searching behavior
by parasitoid *Campoletis sonorensis*. *J. Chem. Ecol.* 10, 1251–1264. doi: 10.1007/BF00988552

Elzen, G., Williams, H., and Vinson, S. (1986). Wind tunnel flight responses by hymenopterous parasitoid *Campoletis sonorensis*
to cotton cultivars and lines. *Entomol. Exp. Appl.* 42, 285–289. doi: 10.1111/j.1570-7458.1986.tb01034.x

Erb, M., Veyrat, N., Robert, C. A. M., Xu, H., Frey, M., Ton, J., et al. (2015). Indole is an essential herbivore-induced volatile
priming signal in maize. *Nat. Commun.* 6, 6273. doi: 10.1038/ncomms7273

Gershenzon, J., McConkey, M. E., and Croteau, R. B. (2000). Regulation of monoterpene accumulation in leaves of peppermint.
Plant Physiol. 122, 205–214. doi: 10.1104/pp.122.1.205

Ghirardo, A., Heller, W., Fladung, M., Schnitzler, J.-P., and Schroeder, H. (2012). Function of defensive volatiles in pedunculate
oak (*Quercus robur*) is tricked by the moth *Tortrix viridana*: Detrimental effect of HIPV emissions in oaks. *Plant Cell
Environ.* 35, 2192–2207. doi: 10.1111/j.1365-3040.2012.02545.x

Grof-Tisza, P., Karban, R., Rasheed, M. U., Saunier, A., and Blande, J. D. (2021). Risk of herbivory negatively correlates with the
diversity of volatile emissions involved in plant communication. *Proc. R. Soc. B Biol. Sci.* 288, 20211790. doi:
10.1098/rspb.2021.1790

Grof-Tisza, P., Kruizenga, N., Tervahauta, A. I., and Blande, J. D. (2022). Volatile-mediated induced and passively acquired
resistance in sagebrush (*Artemisia tridentata*). *J. Chem. Ecol.* 48, 730–745. doi: 10.1007/s10886-022-01378-y

Hagenbucher, S., Birgersson, G., Chattington, S., and Anderson, P. (2016). Domestication influences choice behavior and per-
formance of a generalist herbivore. *Perspect. Plant Ecol. Evol. Syst.* 23, 63–72. doi: 10.6084/m9.figshare. 3467747

Hagenbucher, S., Olson, D. M., Ruberson, J. R., Wäckers, F. L., and Romeis, J. (2013). Resistance mechanisms against arthropod
herbivores in cotton and their interactions with natural enemies. *Crit. Rev. Plant Sci.* 32, 458–482. doi:
10.1080/07352689.2013.809293

Halitschke, R., Keßler, A., Kahl, J., Lorenz, A., and Baldwin, I. T. (2000). Ecophysiological comparison of direct and indirect
defenses in *Nicotiana attenuata*. *Oecologia* 124, 408–417. doi: 10.1007/s004420000389

Hare, J. D. (2007). Variation in herbivore and methyl jasmonate-induced volatiles among genetic lines of *Datura wrightii*. *J.*
Chem. Ecol. 33, 2028–2043. doi: 10.1007/s10886-007-9375-1

Hare, J. D. (2011). Ecological role of volatiles produced by plants in response to damage by herbivorous insects. *Annu. Rev.*
Entomol. 56, 161–180. doi: 10.1146/annurev-ento-120709-144753

Harris, C. M., Ruberson, J. R., Meagher, R., and Tumlinson, J. H. (2012). Host suitability affects odor association in *Cotesia*
marginiventris: Implications in generalist parasitoid host-finding. *J. Chem. Ecol.* 38, 340–347. doi: 10.1007/s10886-
012-0095-9

He, J., Fandino, R. A., Halitschke, R., Luck, K., Köllner, T. G., Murdock, M. H., et al. (2019). An unbiased approach elucidates
variation in (*S*)-(+)-linalool, a context-specific mediator of a tri-trophic interaction in wild tobacco. *Proc. Natl. Acad.*
Sci. 116, 14651–14660. doi: 10.1073/pnas.1818585116

Hervé, M. (2022). *RVAideMemoire: Testing and Plotting Procedures for Biostatistics*. Available at: https://CRAN.R-pro-
ject.org/package=RVAideMemoire

Holopainen, J. K. (2011). Can forest trees compensate for stress-generated growth losses by induced production of volatile
compounds? *Tree Physiol.* 31, 1356–1377. doi: 10.1093/treephys/tpr111

Huang, M., Abel, C., Sohrabi, R., Petri, J., Haupt, I., Cosimano, J., et al. (2010). Variation of herbivore-induced volatile terpenes
among Arabidopsis ecotypes depends on allelic differences and subcellular targeting of two terpene synthases, TPS02
and TPS03. *Plant Physiol.* 153, 1293–1310. doi: 10.1104/pp.110.154864

Huang, X., Kou, J., Jing, W., Han, X., Liu, D., Ghasemzadeh, S., et al. (2022). Transcriptomic and metabolomic reprogramming
in cotton after *Apolygus lucorum* feeding implicated in enhancing recruitment of the parasitoid *Peristenus spretus*. *J.*
Pest Sci. 95, 249–262. doi: 10.1007/s10340-021-01369-0

Irmisch, S., Clavijo McCormick, A., Boeckler, G. A., Schmidt, A., Reichelt, M., Schneider, B., et al. (2013). Two herbivore-induced
cytochrome P450 enzymes CYP79D6 and CYP79D7 catalyze the formation of volatile aldoximes involved in poplar
defense. *Plant Cell* 25, 4737–4754. doi: 10.1105/tpc.113.118265

Irmisch, S., Clavijo McCormick, A., Günther, J., Schmidt, A., Boeckler, G. A., Gershenzon, J., et al. (2014). Herbivore-induced
poplar cytochrome P450 enzymes of the CYP71 family convert aldoximes to nitriles which repel a generalist caterpil-
lar. *Plant J.* 80, 1095–1107. doi: https://doi.org/10.1111/tpj.12711

Jenkins, J. N., Maxwell, F. G., and Lafever, H. N. (1966). The comparative preference of insects for glanded and glandless cot-
tons. *J. Econ. Entomol.* 59, 352–356. doi: 10.1093/jee/59.2.352

Karban, R., Grof-Tisza, P., and Blande, J. D. (2016). Chemotypic variation in volatiles and herbivory for sagebrush. *J. Chem. Ecol.*
42, 829–840.

Karban, R., Wetzel, W. C., Shiojiri, K., Ishizaki, S., Ramirez, S. R., and Blande, J. D. (2014a). Deciphering the language of plant
communication: volatile chemotypes of sagebrush. *New Phytol.* 204, 380–385. doi: 10.1111/nph.12887

Karban, R., Yang, L. H., and Edwards, K. F. (2014b). Volatile communication between plants that affects herbivory: A meta-analysis. *Ecol. Lett.* 17, 44–52. doi: 10.1111/ele.12205

Kariyat, R. R., Mauck, K. E., De Moraes, C. M., Stephenson, A. G., and Mescher, M. C. (2012). Inbreeding alters volatile signalling phenotypes and influences tri-trophic interactions in horsenettle (*Solanum carolinense* L.): Inbreeding impacts plant volatiles and insect attraction. *Ecol. Lett.* 15, 301–309. doi: 10.1111/j.1461-0248.2011.01738.x

Keefover-Ring, K., Thompson, J. D., and Linhart, Y. B. (2009). Beyond six scents: defining a seventh *Thymus vulgaris* chemotype new to southern France by ethanol extraction. *Flavour Fragr. J.* 24, 117–122. doi: 10.1002/ffj.1921

Kessler, A., and Kalske, A. (2018). Plant Secondary Metabolite Diversity and Species Interactions. *Annu. Rev. Ecol. Evol. Syst.* 49, 115–138. doi: 10.1146/annurev-ecolsys-110617-062406

Kleine, S., and Müller, C. (2011). Intraspecific plant chemical diversity and its relation to herbivory. *Oecologia* 166, 175–186. doi: 10.1007/s00442-010-1827-6

Kleine, S., and Müller, C. (2014). Drought stress and leaf herbivory affect root terpenoid concentrations and growth of *Tanace-tum vulgare*. *J. Chem. Ecol.* 40, 1115–1125. doi: 10.1007/s10886-014-0505-2

Lange, B. M. (2015). The evolution of plant secretory structures and emergence of terpenoid chemical diversity. *Annu. Rev. Plant Biol.* 66, 139–159. doi: 10.1146/annurev-arplant-043014-114639

Lenth, R. V. (2023). *emmeans: Estimated Marginal Means, aka Least-Squares Means*. Available at: https://CRAN.R-project.org/package=emmeans

Linhart, Y. B., Gauthier, P., Keefover-Ring, K., and Thompson, J. D. (2015). Variable phytotoxic effects of *Thymus vulgaris* (Lamiaceae) terpenes on associated species. *Int. J. Plant Sci.* 176, 20–30. doi: 10.1086/678772

Linhart, Y. B., and Thompson, J. D. (1999). Thyme is of the essence: biochemical polymorphism and multi-species deterrence. *Evol. Ecol. Res.* 1, 151–171.

Llandres, A. L., Almohamad, R., Brévault, T., Renou, A., Téréta, I., Jean, J., et al. (2018). Plant training for induced defense against insect pests: a promising tool for integrated pest management in cotton: Cotton training for induced defense against pests. *Pest Manag. Sci.* 74, 2004–2012. doi: 10.1002/ps.5039

Loreto, F., and Schnitzler, J.-P. (2010). Abiotic stresses and induced BVOCs. *Trends Plant Sci.* 15, 154–166. doi: 10.1016/j.tplants.2009.12.006

Loughrin, J. H., Manukian, A., Heath, R. R., and Tumlinson, J. H. (1995). Volatiles emitted by different cotton varieties damaged by feeding beet armyworm larvae. *J. Chem. Ecol.* 21, 1217–1227. doi: 10.1007/BF02228321

Loughrin, J. H., Manukian, A., Heath, R. R., Turlings, T. C., and Tumlinson, J. H. (1994). Diurnal cycle of emission of induced volatile terpenoids by herbivore-injured cotton plant. *Proc. Natl. Acad. Sci.* 91, 11836–11840. doi: 10.1073/pnas.91.25.11836

Mamin, M., Vallat, A., and Turlings, T. C. J. (2023). Cotton plants as ideal models for teaching and research on inducible direct plant defenses. *Front. Ecol. Evol.* 11. doi: 10.3389/fevo.2023.1119472

McAuslane, H. J., and Alborn, H. T. (1998). Systemic induction of allelochemicals in glanded and glandless isogenic cotton by *Spodoptera exigua* feeding. *J. Chem. Ecol.* 24, 399–416. doi: 10.1023/A:1022596812049

McAuslane, H. J., Alborn, H. T., and Toth, J. P. (1997). Systemic induction of terpenoid aldehydes in cotton pigment glands by feeding of larval *Spodoptera exigua*. *J. Chem. Ecol.* 23, 2861–2879. doi: 10.1023/A:1022575313325

McCall, P. J., Turlings, T. C. J., Lewis, W. J., and Tumlinson, J. H. (1993). Role of plant volatiles in host location by the specialist
parasitoid *Microplitis croceipes* cresson (Braconidae: Hymenoptera). *J. Insect Behav.* 6, 625–639. doi:
10.1007/BF01048128

McCall, P. J., Turlings, T. C. J., Loughrin, J., Proveaux, A. T., and Tumlinson, J. H. (1994). Herbivore-induced volatile emissions
from cotton (*Gossypium hirsutum L.*) seedlings. *J. Chem. Ecol.* 20, 3039–3050. doi: 10.1007/BF02033709

McCormick, A. C., Reinecke, A., Gershenzon, J., and Unsicker, S. B. (2016). Feeding experience affects the behavioral response
of polyphagous gypsy moth caterpillars to herbivore-induced poplar volatiles. *J. Chem. Ecol.* 42, 382–393. doi:
10.1007/s10886-016-0698-7

Mitchell, C., Brennan, R. M., Graham, J., and Karley, A. J. (2016). Plant defense against herbivorous pests: exploiting resistance
and tolerance traits for sustainable crop protection. *Front. Plant Sci.* 7, 1132. doi: 10.3389/fpls.2016.01132

Moore, B. D., Andrew, R. L., Külheim, C., and Foley, W. J. (2014). Explaining intraspecific diversity in plant secondary metabo-
lites in an ecological context. *New Phytol.* 201, 733–750. doi: 10.1111/nph.12526

Müller, C., and Junker, R. R. (2022). Chemical phenotype as important and dynamic niche dimension of plants. *New Phytol.*
234, 1168–1174. doi: 10.1111/nph.18075

Munguía-Rosas, M. A., Angulo, D. F., Arceo-Gómez, G., and Parra-Tabla, V. (2019). Variation in leaf traits across a precipitation
gradient in coastal sand dunes in Yucatan Peninsula. *J. Arid Environ.* 162, 10–17. doi: 10.1016/j.jaridenv.2018.12.001

Ngumbi, E., Chen, L., and Fadamiro, H. Y. (2009). Comparative GC-EAD responses of a specialist (*Microplitis croceipes*) and a
generalist (*Cotesia marginiventris*) parasitoid to cotton volatiles induced by two caterpillar species. *J. Chem. Ecol.* 35,
1009–1020. doi: 10.1007/s10886-009-9700-y

Oksanen, J., Simpson, G. L., Blanchet, F. G., Kindt, R., Legendre, P., Minchin, P. R., et al. (2022). *vegan: Community Ecology*
Package. Available at: https://CRAN.R-project.org/package=vegan

Opitz, S., Kunert, G., and Gershenzon, J. (2008). Increased terpenoid accumulation in cotton (*Gossypium hirsutum*) foliage is a
general wound response. *J. Chem. Ecol.* 34, 508–522. doi: 10.1007/s10886-008-9453-z

Palarea-Albaladejo, J., and Martin-Fernandez, J. A. (2015). zCompositions – R package for multivariate imputation of left-cen-
sored data under a compositional approach. *Chemom. Intell. Lab. Syst.* 143, 85–96.

Paré, P. W., and Tumlinson, J. H. (1997). De novo biosynthesis of volatiles induced by insect herbivory in cotton plants. 114, 7.
doi: 10.1016/S0031-9422(97)00442-1

Pickett, J. A., and Khan, Z. R. (2016). Plant volatile-mediated signalling and its application in agriculture: successes and chal-
lenges. *New Phytol.* 212, 856–870. doi: 10.1111/nph.14274

Quijano-Medina, T., Covelo, F., Moreira, X., and Abdala-Roberts, L. (2019). Compensation to simulated insect leaf herbivory in
wild cotton (*Gossypium hirsutum*): responses to multiple levels of damage and associated traits. *Plant Biol.* 21, 805–
812. doi: https://doi.org/10.1111/plb.13002

Quijano-Medina, T., Turlings, T. C. J., Sosenski, P., Grandi, L., Cervera, J. C., Moreira, X., et al. (2021). Effects of soil salinity on
the expression of direct and indirect defences in wild cotton *Gossypium hirsutum*. *J. Ecol.* 109, 354–368. doi:
https://doi.org/10.1111/1365-2745.13483

R Core Team (2022). *R: A Language and Environment for Statistical Computing*. Vienna, Austria: R Foundation for Statistical
Computing. Available at: https://www.R-project.org/

Rasmann, S., Buri, A., Gallot-Lavallée, M., Joaquim, J., Purcell, J., and Pellissier, L. (2014). Differential allocation and deployment
of direct and indirect defences by *Vicia sepium* along elevation gradients. *J. Ecol.* 102, 930–938. doi: 10.1111/1365-
2745.12253

Razavizadeh, R., and Komatsu, S. (2018). Changes in essential oil and physiological parameters of callus and seedlings of *Carum*
copticum L. under in vitro drought stress. *J. Food Meas. Charact.* 12, 1581–1592. doi: 10.1007/s11694-018-9773-9

Rodriguez-Saona, C., and Crafts-Brandner, S. J. (2003). Volatile emissions triggered by multiple herbivore damage: beet
armyworm and whitefly feeding on cotton plants. *J. Chem. Ecol.* doi: 10.1023/a:1026314102866

Rodriguez-Saona, C. R., and Frost, C. J. (2010). New evidence for a multi-functional role of herbivore-induced plant volatiles in
defense against herbivores. *Plant Signal. Behav.* 5, 58–60. doi: 10.4161/psb.5.1.10160

Röse, U. S. R., Manukian, A., Heath, R. R., and Tumlinson, J. H. (1996). Volatile semiochemicals released from undamaged
cotton leaves (A systemic response of living plants to caterpillar damage). *Plant Physiol.* 111, 487–495. doi:
10.1104/pp.111.2.487

Röse, U. S. R., and Tumlinson, J. H. (2005). Systemic induction of volatile release in cotton: How specific is the signal to her-
bivory? *Planta* 222, 327–335. doi: 10.1007/s00425-005-1528-2

Ruberto, G., and Baratta, M. T. (2000). Antioxidant activity of selected essential oil components in two lipid model systems.
Food Chem. 69, 167–174. doi: 10.1016/S0308-8146(99)00247-2

Schuman, M. C., Heinzel, N., Gaquerel, E., Svatos, A., and Baldwin, I. T. (2009). Polymorphism in jasmonate signaling partially
accounts for the variety of volatiles produced by *Nicotiana attenuata* plants in a native population. *New Phytol.* 183,
1134–1148. doi: 10.1111/j.1469-8137.2009.02894.x

Selmar, D., and Kleinwächter, M. (2013). Stress enhances the synthesis of secondary plant products: The impact of stress-
related over-reduction on the accumulation of natural products. *Plant Cell Physiol.* 54, 817–826. doi:
10.1093/pcp/pct054

Senft, M., Clancy, M. V., Weisser, W. W., Schnitzler, J., and Zytynska, S. E. (2019). Additive effects of plant chemotype, mutu-
alistic ants and predators on aphid performance and survival. *Funct. Ecol.* 33, 139–151. doi: 10.1111/1365-
2435.13227

Sørensen, M., Neilson, E. H. J., and Møller, B. L. (2018). Oximes: Unrecognized chameleons in general and specialized plant
metabolism. *Mol. Plant* 11, 95–117. doi: 10.1016/j.molp.2017.12.014

Stipanovic, R. D., Bell, A. A., O'Brien, D. H., and Lukefahr, M. J. (1978). Heliocide H1. A new insecticidal C25 terpenoid from
cotton (*Gossypium hirsutum*). *J. Agric. Food Chem.* 26, 115–118.

Taft, S., Najar, A., Godbout, J., Bousquet, J., and Erbilgin, N. (2015). Variations in foliar monoterpenes across the range of jack
pine reveal three widespread chemotypes: implications to host expansion of invasive mountain pine beetle. *Front.*
Plant Sci. 6. doi: 10.3389/fpls.2015.00342

Tian, Z., Luo, Q., Li, Y., and Zuo, Z. (2020). Terpinene and β-pinene acting as signaling molecules to improve *Cinnamomum*
camphora thermotolerance. *Ind. Crops Prod.* 154, 112641. doi: 10.1016/j.indcrop.2020.112641

Turlings, T. C., and Erb, M. (2018). Tritrophic interactions mediated by herbivore-induced plant volatiles: mechanisms, ecolog-
ical relevance, and application potential. *Annu. Rev. Entomol.* 63, 433–452. doi: 10.1146/annurev-ento-020117-
043507

Unsicker, S. B., Kunert, G., and Gershenzon, J. (2009). Protective perfumes: the role of vegetative volatiles in plant defense
against herbivores. *Curr. Opin. Plant Biol.* 12, 479–485. doi: 10.1016/j.pbi.2009.04.001

Villamar-Torres, R., Mehdi Jazayeri, S., Liuba-Delfini, G., García Cruzaty, L. C., and Viot, C.-R. (2018). Volatile organic com-
pounds: plant natural defense mechanisms against herbivorous arthropods and an opportunity for plant breeding of
cotton. *Sci. Agropecu.* 9, 287–297. doi: 10.17268/sci.agropecu.2018.01.14

Voelker, J., Mauleon, R., and Shepherd, M. (2023). The terpene synthase genes of Melaleuca alternifolia (tea tree) and com-
parative gene family analysis among Myrtaceae essential oil crops. *Plant Syst. Evol.* 309, 13. doi: 10.1007/s00606-
023-01847-1

Wason, E. L., Agrawal, A. A., and Hunter, M. D. (2013). A genetically-based latitudinal cline in the emission of herbivore-induced
plant volatile oganic compounds. *J. Chem. Ecol.* 39, 1101–1111. doi: 10.1007/s10886-013-0309-9

Wegier, A., Piñeyro-Nelson, A., Alarcón, J., Gálvez-Mariscal, A., Álvarez-Buylla, E., and Piñero, D. (2011). Recent long-distance
transgene flow into wild populations conforms to historical patterns of gene flow in cotton (*Gossypium hirsutum*) at
its centre of origin. *Mol. Ecol.* 20, 4182–4194. doi: 10.1111/j.1365-294X.2011.05258.x

Wu, J., Hettenhausen, C., Schuman, M. C., and Baldwin, I. T. (2008). A comparison of two *Nicotiana attenuata* accessions
reveals large differences in signaling induced by oral secretions of the specialist herbivore *Manduca sexta*. *Plant Phys-
iol.* 146, 927–939. doi: 10.1104/pp.107.114785

Xu, S., Kreitzer, C., McGale, E., Lackus, N. D., Guo, H., Köllner, T. G., et al. (2020). Allelic differences of clustered terpene syn-
thases contribute to correlated intraspecific variation of floral and herbivory-induced volatiles in a wild tobacco. *New
Phytol.* 228, 1083–1096. doi: 10.1111/nph.16739

Yuan, D., Grover, C. E., Hu, G., Pan, M., Miller, E. R., Conover, J. L., et al. (2021). Parallel and intertwining threads of domesti-
cation in allopolyploid cotton. *Adv. Sci.* 8, 2003634. doi: 10.1002/advs.202003634

Zakir, A., Sadek, M. M., Bengtsson, M., Hansson, B. S., Witzgall, P., and Anderson, P. (2013). Herbivore-induced plant volatiles
provide associational resistance against an ovipositing herbivore. *J. Ecol.* 101, 410–417. doi: 10.1111/1365-
2745.12041

Zhang, L., Lu, G., Huang, X., Guo, H., Su, X., Han, L., et al. (2020). Overexpression of the caryophyllene synthase gene *GhTPS1*
in cotton negatively affects multiple pests while attracting parasitoids. *Pest Manag. Sci.* 76, 1722–1730. doi:
10.1002/ps.5695

Table 1: Summary of the GLMMs analyzing the variation in induction (group-by-herbivory interaction) of volatile terpenes concentration in wild cotton leaves among genotypes and populations (random slopes; Likelihood ratio tests on nested models), as well as among geographic regions and between chemotypes (fixed effects; Wald chi-squared tests). Because they were partially nested, the effects of chemotype and region were analysed in separate models. Chemotype was included as fixed effect when testing variation among populations and genotypes. Chi-squared (χ^2) values are shown, and statistically significant (p < 0.05) effects are indicated in bold. Df = Degrees of freedom of χ^2 tests. The random slope component includes a term for the variance in slopes and one for the covariance between slopes and intercepts, unless the model could not converge, and the covariance term was excluded (italicized values).

Leaf volatiles	Genotype x herbivory (χ^2 value, df = 2)	Population x herbivory (χ^2 value, df = 2)	Region x herbivory (χ^2 value, df = 1)	Chemotype x herbivory (χ^2 value, df = 1)
Total terpenes	0.35	1.22	2.88	0.19
(E)-β-Ocimene	5.23	0.71	2.02	**7.69**
δ-Cadinene	*0.00*	0.88	0.94	0.29
β-Caryophyllene	0.02	0.48	4.00	0.02
α-Pinene	*0.00*	1.42	3.25	0.01
γ-Terpinene	0.04	0.59	1.35	-

Table 2: Summary of the GLMMs analyzing the variation in HIPV emission rates from damaged wild cotton plants among populations (random intercept; Likelihood ratio tests on nested models), as well as among geographic regions and between chemotypes (fixed effects; Wald chi-squared tests). Because they were partially nested, the effects of chemotype and region were analysed in separate models. Chemotype was included as fixed effect when testing variation among populations. Damage intensity was controlled for in all models. Chi-squared (χ^2) values are shown, and statistically significant (p < 0.05) effects are indicated in bold. Df = Degrees of freedom. Ce: Celestún, Si: Sisal, Ch: Chixculub, Co: Coloradas; A: chemotype A. B: chemotype B.

HIPV emitted	Population (χ^2 value, df = 1)	Region (χ^2 value, df = 3)	Effect	Chemotype (χ^2 value, df = 1)	Effect	Figure
Total HIPVs	1.34	2.89	-	0.00	-	Fig. S10
(E,E)-α-Farnesene	**4.23**	**25.45**	Ch < Co ≤ Ce ≤ Si	0.72	-	Fig. 6
(E)-Myroxide	**21.88**	**11.44**	Ch ≤ Co ≤ Ce ≤ Si	0.04	-	Fig. S10
(E)-Methylbutyraldoxime	**19.25**	**62.02**	Co ≤ Ch < Ce ≤ Si	**7.24**	B > A	Fig. 6
(Z)-Methylbutyraldoxime	**20.86**	**71.29**	Co ≤ Ch < Ce ≤ Si	3.50	-	Fig. S10
Unknown oxime	1.89	**25.79**	Co ≤ Ch < Ce ≤ Si	**5.07**	B > A	Fig. 6
Benzyl nitrile	0.11	**8.83**	Ch ≤ Co ≤ Si ≤ Ce	**4.71**	B > A	Fig. 6
Unknown nitrile	1.55	**8.03**	Ce ≤ Si ≤ Ch ≤ Co	**7.10**	A > B	Fig. 6

Table 3: structures of the GLMMs used to analyze concentrations of volatile terpenes in wild cotton leaves. Chemo = chemotype, Area = fourth leaf area, Damage = damage intensity (total leaf area consumed by *S. exigua*). Full fixed effects are shown, interaction terms were removed if not significantly contributing to the model.

All plants			
	Effects tested	**Fixed effects**	**Random effects (intercepts and slopes)**
A	Herbivory / Chemotype / Chemotype-by-herbivory interaction	Herbivory + Chemo + Chemo:Herbivory + Area + Batch + Area:Batch	Population + Genotype
B	Genotype-by-herbivory interaction	Herbivory + Chemo + Chemo:Herbivory + Area + Batch + Area:Batch	Population + Herbivory:Genotype
C	Population-by-herbivory interaction	Herbivory + Chemo + Chemo:Herbivory + Area + Batch + Area:Batch	Herbivory: Population + Genotype
D	Region-by-herbivory interaction	Herbivory + Region + Region:Herbivory + Area + Batch + Area:Batch	Population + Genotype
Damaged plants			
	Effects tested	**Fixed effects**	**Random effects (intercepts and slopes)**
E	Damage intensity	Damage + Chemo + Chemo:Damage + Area + Batch + Area: Batch	Population + Genotype

Table 4: structures of the GLMMs used to analyze emission rates of volatiles by wild cotton plants. Chemo = chemotype, Area = fourth leaf area, Damage = damage intensity (total leaf area consumed by *S. exigua*). Full models are shown, interaction terms were removed if not significantly contributing to the model.

All plants			
	Effects tested	**Fixed effects**	**Random effects (intercepts)**
A	Herbivory / Chemotype-by-Herbivory int.	Herbivory + Chemo + Chemo:Herbivory + Area + Batch + Area:Batch	Population + Genotype
B	Region-by-Herbivory int.	Herbivory + Region + Region:Herbivory + Area + Batch + Area:Batch	Population + Genotype
Damaged plants			
	Effects tested	**Fixed effects**	**Random effects (intercepts)**
C	Chemotype / Damage intensity	Chemo + Damage + Chemo:Damage + Area + Batch + Area:Batch	Population
D	Region	Region + Damage + Region:Damage + Area + Batch + Area:Batch	Population
E	Population	Chemo + Damage + Chemo:Damage + Area + Batch + Area:Batch	Population

Figure 1. Illustration of the experimental process. Seeds were collected from 16 populations (red points) in four regions along the northern coast of the Yucatán Peninsula. Plants were grown in a common greenhouse until they developed the fourth leaf, at which time half of the plants were damaged by *Spodoptera exigua* larvae on their first and second leaves for two days and the other half was kept intact. Emitted volatiles were collected by headspace sampling and stored leaf volatiles by hexane extraction of the undamaged fourth leaves.

Figure 2. Effect of herbivory on stored leaf mono- and sesquiterpenes for both wild cotton chemotypes. The volatile content
was extracted from the fourth leaves of 353 undamaged wild cotton plants and 322 plants damaged by *Spodoptera exigua* for
two days on their first and second leaves. **A:** Effect of herbivory on the leaf concentrations of a subset of terpenes. Estimated
marginal means and 95% CI from GLMMs. γ-Terpinene was analyzed within chemotype A while all other compounds were
analyzed for both chemotypes together, with chemotype set as fixed factor. **B**: Biplot with symmetric scaling of a pRDA on the
relative abundances (clr-transformed) of leaf mono- and sesquiterpenes.

Figure 3: Composition of the headspace volatiles emitted by undamaged wild cotton plants and plants damaged for two days by *Spodoptera exigua*. A: Compounds detected both in the plants' leaves extracts and headspace, or in their headspace only. RI = Retention indices. The retention index for the unknown aldoxime could not be calculated because it eluted before the smallest alkane. † indicates compounds specific to damaged plants (detected in less than 10% of undamaged plants, but in between 43% and 100% of damaged plants). * indicates compounds whose identity was confirmed with standards. Distinction between the two 2-methylbutyraldoximes isomers is tentative. FA = fatty acid derivatives*;* MT = monoterpenes; SQT= sesquiterpenes; Ox. = aldoximes; Ar. = aromatics; HT = homoterpenes. **B:** Composition of medians emitted volatile profiles of undamaged and damaged plants of both chemotypes. Area of the rectangles is proportional to the contribution of the compound to the profile. Numbers refer to A.

Figure 4. Effect of herbivory on the volatiles emitted by the two wild cotton chemotypes. Headspace volatiles were sampled
from 45 undamaged wild cotton plants and 48 plants damaged by *Spodoptera exigua* for two days on their first and second
leaves. **A:** Biplot with symmetric scaling of a pRDA on the relative abundances (clr-transformed) of all compounds quantified
in the headspace samples. Only compounds accounting for more than 10% of inertia are plotted (full biplot in Fig. S8). **B**:
Effect of herbivory on emission rates of all emitted volatiles and of a subset of emitted terpenes found also in the leaf extracts.
Estimated marginal means and 95% CI from GLMMs are plotted. γ-Terpenene was analyzed within chemotype A while all other
compounds were analyzed for both chemotypes together, with chemotype set as fixed factor. **C:** relationship between damage
intensity (leaf area damaged) and the emission rates of a subset of terpenes. Left: multiplicative increases with the leaf area
damaged (back-transformed coefficient estimates from GLMMs and their 95% CIs). The dotted line indicates a coefficient of
1, where quantities emitted remain constant with damaged area. Right: adjusted predictions and 95% CI from GLMMs with
observed values as data points. γ-Terpenene was analyzed within chemotype A while all other compounds were analyzed for
both chemotypes together, with chemotype set as fixed factor.

Figure 5. Influence of γ-terpinene variation in leaves on headspace emissions: correlation (Spearman) between the ratio of γ-terpinene and α-pinene concentrations in leaves (x-axis) and in headspace emissions (y-axis).

Figure 6. Variation in emissions of herbivory-induced volatiles among the four regions grouping wild cotton populations along the Yucatan coast and between the two terpene-chemotypes. Headspace volatile emissions were sampled from 48 wild cotton plant damaged by *Spodoptera exigua* for two days. Regions are displayed from west to east. Prevalence of chemotypes is nested with regions, as indicated by the color of data points (grey points indicate chemotype A; blue points indicate chemotype B), therefore they were analyzed separately. Estimated marginal means and 95% CI from GLMMs with observed values as data points. Damage intensity was controlled for as covariate. Letters indicate significant pairwise comparisons after multiple test adjustment (FDR). For regions, % values indicate the difference between the region with the lowest and highest estimated mean emission rates.

Parsed Citations

Abdala-Roberts, L., Quijano-Medina, T., Moreira, X., Vázquez-González, C., Parra-Tabla, V., Berny Mier Y Terán, J. C., et al. (2019). Bottom-up control of geographic variation in insect herbivory on wild cotton (Gossypium hirsutum) by plant defenses and climate. Am. J. Bot. 106, 1059–1067. doi: 10.1002/ajb2.1330

Google Scholar: Author Only Title Only Author and Title

Alavi-Samani, S. M., Kachouei, M. A., and Pirbalouti, A. G. (2015). Growth, yield, chemical composition, and antioxidant activity of essential oils from two thyme species under foliar application of jasmonic acid and water deficit conditions. Hortic. Environ. Biotechnol. 56, 411–420. doi: 10.1007/s13580-015-0117-y

Google Scholar: Author Only Title Only Author and Title

Amiot, J., Salmon, Y., Collin, C., and Thompson, J. D. (2005). Differential resistance to freezing and spatial distribution in a chemically polymorphic plant Thymus vulgaris. Ecol. Lett. 8, 370–377. doi: 10.1111/j.1461-0248.2005.00728.x

Google Scholar: Author Only Title Only Author and Title

Arce, C. M., Besomi, G., Glauser, G., and Turlings, T. C. (2021). Caterpillar-induced volatile emissions in cotton: The relative importance of damage and insect-derived factors. Front. Plant Sci., 1503. doi: 10.3389/fpls.2021.709858

Google Scholar: Author Only Title Only Author and Title

Bertamini, M., Faralli, M., Varotto, C., Grando, M. S., and Cappellin, L. (2021). Leaf monoterpene emission limits photosynthetic downregulation under heat stress in field-grown grapevine. Plants 10, 181. doi: 10.3390/plants10010181

Google Scholar: Author Only Title Only Author and Title

Boogaart, K. G. van den, Tolosana-Delgado, R., and Bren, M. (2023). compositions: Compositional Data Analysis. Available at: https://CRAN.R-project.org/package=compositions

Google Scholar: Author Only Title Only Author and Title

Briones-May, Y., Quijano-Medina, T., Pérez-Niño, B., Benrey, B., Turlings, T. C. J., Bustos-Segura, C., et al. (2023). Soil salinization disrupts plant–plant signaling effects on extra-floral nectar induction in wild cotton. Oecologia. doi: 10.1007/s00442-023-05395-w

Google Scholar: Author Only Title Only Author and Title

Bustos-Segura, C., Dillon, S., Keszei, A, Foley, W. J., and Külheim, C. (2017). Intraspecific diversity of terpenes of Eucalyptus camaldulensis (Myrtaceae) at a continental scale. Aust. J. Bot. 65, 257–269. doi: 10.1071/BT16183

Google Scholar: Author Only Title Only Author and Title

Bustos-Segura, C., and Foley, W. J. (2018). Foliar terpene chemotypes and herbivory determine variation in plant volatile emissions. J. Chem. Ecol. 44, 51–61. doi: 10.1007/s10886-017-0919-8

Google Scholar: Author Only Title Only Author and Title

Bustos-Segura, C., Külheim, C., and Foley, W. (2015). Effects of terpene chemotypes of Melaleuca alternifolia on two specialist leaf beetles and susceptibility to myrtle rust. J. Chem. Ecol. 41, 937–947. doi: 10.1007/s10886-015-0628-0

Google Scholar: Author Only Title Only Author and Title

Clancy, M. V., Haberer, G., Jud, W., Niederbacher, B., Niederbacher, S., Senft, M., et al. (2020). Under fire-simultaneous volatilome and transcriptome analysis unravels fine-scale responses of tansy chemotypes to dual herbivore attack. BMC Plant Biol. 20, 551. doi: 10.1186/s12870-020-02745-1

Google Scholar: Author Only Title Only Author and Title

Clancy, M. V., Mamin, M., Flückiger, G., Quijano-Medina, T., Pérez-Niño, B., Abdala-Roberts, L., et al. (2023). Terpene chemotypes in Gossypium hirsutum (wild cotton) from the Yucatan Peninsula, Mexico. Phytochemistry 205, 113454. doi: 10.1016/j.phytochem.2022.113454

Google Scholar: Author Only Title Only Author and Title

Clancy, M. V., Zytynska, S. E., Senft, M., Weisser, W. W., and Schnitzler, J.-P. (2016). Chemotypic variation in terpenes emitted from storage pools influences early aphid colonisation on tansy. Sci. Rep. 6, 38087. doi: 10.1038/srep38087

Google Scholar: Author Only Title Only Author and Title

Clavijo McCormick, A, Gershenzon, J., and Unsicker, S. B. (2014a). Little peaks with big effects: establishing the role of minor plant volatiles in plant-insect interactions: Minor plant volatiles. Plant Cell Environ. 37, 1836–1844. doi: 10.1111/pce.12357

Google Scholar: Author Only Title Only Author and Title

Clavijo McCormick, A, Irmisch, S., Reinecke, A, Boeckler, G. A, Veit, D., Reichelt, M., et al. (2014b). Herbivore-induced volatile emission in black poplar: regulation and role in attracting herbivore enemies. Plant Cell Environ. 37, 1909–1923. doi: 10.1111/pce.12287

Google Scholar: Author Only Title Only Author and Title

Coppens d'Eeckenbrugge, G., and Lacape, J.-M. (2014). Distribution and Differentiation of Wild, Feral, and Cultivated Populations of Perennial Upland Cotton (Gossypium hirsutum L.) in Mesoamerica and the Caribbean. PLoS ONE 9, e107458. doi:

10.1371/journal.pone.0107458

Google Scholar: Author Only Title Only Author and Title

De Sousa, D. P., Damasceno, R. O. S., Amorati, R., Elshabrawy, H. A, De Castro, R. D., Bezerra, D. P., et al. (2023). Essential Oils: Chemistry and Pharmacological Activities. Biomolecules 13, 1144. doi: 10.3390/biom13071144

Google Scholar: Author Only Title Only Author and Title

Delphia, C. M., Rohr, J. R., Stephenson, A. G., De Moraes, C. M., and Mescher, M. C. (2009). Effects of Genetic Variation and Inbreeding on Volatile Production in a Field Population of Horsenettle. Int. J. Plant Sci. 170, 12–20. doi: 10.1086/593039

Google Scholar: Author Only Title Only Author and Title

Dicke, M., and Baldwin, I. T. (2010). The evolutionary context for herbivore-induced plant volatiles: beyond the 'cry for help.' Trends Plant Sci. 15, 167–175. doi: 10.1016/j.tplants.2009.12.002

Google Scholar: Author Only Title Only Author and Title

Elzen, G. W., Williams, H. J., Bell, A A, Stipanovic, R. D., and Vinson, S. B. (1985). Quantification of volatile terpenes of glanded and glandless Gossypium hirsutum L. cultivars and lines by gas chromatography. J. Agric. Food Chem. 33, 1079–1082. doi: 10.1021/jf00066a015

Google Scholar: Author Only Title Only Author and Title

Elzen, G., Williams, H., and Vinson, S. (1984). Isolation and identification of cotton synomones mediating searching behavior by parasitoid Campoletis sonorensis. J. Chem. Ecol. 10, 1251–1264. doi: 10.1007/BF00988552

Google Scholar: Author Only Title Only Author and Title

Elzen, G., Williams, H., and Vinson, S. (1986). Wind tunnel flight responses by hymenopterous parasitoid Campoletis sonorensis to cotton cultivars and lines. Entomol. Exp. Appl. 42, 285–289. doi: 10.1111/j.1570-7458.1986.tb01034.x

Google Scholar: Author Only Title Only Author and Title

Erb, M., Veyrat, N., Robert, C. A M., Xu, H., Frey, M., Ton, J., et al. (2015). Indole is an essential herbivore-induced volatile priming signal in maize. Nat. Commun. 6, 6273. doi: 10.1038/ncomms7273

Google Scholar: Author Only Title Only Author and Title

Gershenzon, J., McConkey, M. E., and Croteau, R. B. (2000). Regulation of monoterpene accumulation in leaves of peppermint. Plant Physiol. 122, 205–214. doi: 10.1104/pp.122.1.205

Google Scholar: Author Only Title Only Author and Title

Ghirardo, A, Heller, W., Fladung, M., Schnitzler, J.-P., and Schroeder, H. (2012). Function of defensive volatiles in pedunculate oak (Quercus robur) is tricked by the moth Tortrix viridana: Detrimental effect of HIPV emissions in oaks. Plant Cell Environ. 35, 2192–2207. doi: 10.1111/j.1365-3040.2012.02545.x

Google Scholar: Author Only Title Only Author and Title

Grof-Tisza, P., Karban, R., Rasheed, M. U., Saunier, A, and Blande, J. D. (2021). Risk of herbivory negatively correlates with the diversity of volatile emissions involved in plant communication. Proc. R. Soc. B Biol. Sci. 288, 20211790. doi: 10.1098/rspb.2021.1790

Google Scholar: Author Only Title Only Author and Title

Grof-Tisza, P., Kruizenga, N., Tervahauta, A. I., and Blande, J. D. (2022). Volatile-mediated induced and passively acquired resistance in sagebrush (Artemisia tridentata). J. Chem. Ecol. 48, 730–745. doi: 10.1007/s10886-022-01378-y

Google Scholar: Author Only Title Only Author and Title

Hagenbucher, S., Birgersson, G., Chattington, S., and Anderson, P. (2016). Domestication influences choice behavior and performance of a generalist herbivore. Perspect. Plant Ecol. Evol. Syst. 23, 63–72. doi: 10.6084/m9.figshare. 3467747

Google Scholar: Author Only Title Only Author and Title

Hagenbucher, S., Olson, D. M., Ruberson, J. R., Wäckers, F. L., and Romeis, J. (2013). Resistance mechanisms against arthropod herbivores in cotton and their interactions with natural enemies. Crit. Rev. Plant Sci. 32, 458–482. doi: 10.1080/07352689.2013.809293

Google Scholar: Author Only Title Only Author and Title

Halitschke, R., Keßler, A, Kahl, J., Lorenz, A, and Baldwin, I. T. (2000). Ecophysiological comparison of direct and indirect defenses in Nicotiana attenuata. Oecologia 124, 408–417. doi: 10.1007/s004420000389

Google Scholar: Author Only Title Only Author and Title

Hare, J. D. (2007). Variation in herbivore and methyl jasmonate-induced volatiles among genetic lines of Datura wrightii. J. Chem. Ecol. 33, 2028–2043. doi: 10.1007/s10886-007-9375-1

Google Scholar: Author Only Title Only Author and Title

Hare, J. D. (2011). Ecological role of volatiles produced by plants in response to damage by herbivorous insects. Annu. Rev. Entomol. 56, 161–180. doi: 10.1146/annurev-ento-120709-144753

Google Scholar: Author Only Title Only Author and Title

Harris, C. M., Ruberson, J. R., Meagher, R., and Tumlinson, J. H. (2012). Host suitability affects odor association in Cotesia marginiventris: Implications in generalist parasitoid host-finding. J. Chem. Ecol. 38, 340–347. doi: 10.1007/s10886-012-0095-9

Google Scholar: Author Only Title Only Author and Title

He, J., Fandino, R. A, Halitschke, R., Luck, K., Köllner, T. G., Murdock, M. H., et al. (2019). An unbiased approach elucidates variation in (S)-(+)-linalool, a context-specific mediator of a tri-trophic interaction in wild tobacco. Proc. Natl. Acad. Sci. 116, 14651–14660. doi: 10.1073/pnas.1818585116

Google Scholar: Author Only Title Only Author and Title

Hervé, M. (2022). RVAideMemoire: Testing and Plotting Procedures for Biostatistics. Available at: https://CRAN.R-project.org/package=RVAideMemoire

Google Scholar: Author Only Title Only Author and Title

Holopainen, J. K. (2011). Can forest trees compensate for stress-generated growth losses by induced production of volatile compounds? Tree Physiol. 31, 1356–1377. doi: 10.1093/treephys/tpr111

Google Scholar: Author Only Title Only Author and Title

Huang, M., Abel, C., Sohrabi, R., Petri, J., Haupt, I., Cosimano, J., et al. (2010). Variation of herbivore-induced volatile terpenes among Arabidopsis ecotypes depends on allelic differences and subcellular targeting of two terpene synthases, TPS02 and TPS03. Plant Physiol. 153, 1293–1310. doi: 10.1104/pp.110.154864

Google Scholar: Author Only Title Only Author and Title

Huang, X., Kou, J., Jing, W., Han, X., Liu, D., Ghasemzadeh, S., et al. (2022). Transcriptomic and metabolomic reprogramming in cotton after Apolygus lucorum feeding implicated in enhancing recruitment of the parasitoid Peristenus spretus. J. Pest Sci. 95, 249–262. doi: 10.1007/s10340-021-01369-0

Google Scholar: Author Only Title Only Author and Title

Irmisch, S., Clavijo McCormick, A, Boeckler, G. A, Schmidt, A, Reichelt, M., Schneider, B., et al. (2013). Two herbivore-induced cytochrome P450 enzymes CYP79D6 and CYP79D7 catalyze the formation of volatile aldoximes involved in poplar defense. Plant Cell 25, 4737–4754. doi: 10.1105/tpc.113.118265

Google Scholar: Author Only Title Only Author and Title

Irmisch, S., Clavijo McCormick, A, Günther, J., Schmidt, A, Boeckler, G. A, Gershenzon, J., et al. (2014). Herbivore-induced poplar cytochrome P450 enzymes of the CYP71 family convert aldoximes to nitriles which repel a generalist caterpillar. Plant J. 80, 1095–1107. doi: https://doi.org/10.1111/tpj.12711

Google Scholar: Author Only Title Only Author and Title

Jenkins, J. N., Maxwell, F. G., and Lafever, H. N. (1966). The comparative preference of insects for glanded and glandless cottons. J. Econ. Entomol. 59, 352–356. doi: 10.1093/jee/59.2.352

Google Scholar: Author Only Title Only Author and Title

Karban, R., Grof-Tisza, P., and Blande, J. D. (2016). Chemotypic variation in volatiles and herbivory for sagebrush. J. Chem. Ecol. 42, 829–840.

Google Scholar: Author Only Title Only Author and Title

Karban, R., Wetzel, W. C., Shiojiri, K., Ishizaki, S., Ramirez, S. R., and Blande, J. D. (2014a). Deciphering the language of plant communication: volatile chemotypes of sagebrush. New Phytol. 204, 380–385. doi: 10.1111/nph.12887

Google Scholar: Author Only Title Only Author and Title

Karban, R., Yang, L. H., and Edwards, K. F. (2014b). Volatile communication between plants that affects herbivory: A meta-analysis. Ecol. Lett. 17, 44–52. doi: 10.1111/ele.12205

Google Scholar: Author Only Title Only Author and Title

Kariyat, R. R., Mauck, K. E., De Moraes, C. M., Stephenson, A. G., and Mescher, M. C. (2012). Inbreeding alters volatile signalling phenotypes and influences tri-trophic interactions in horsenettle (Solanum carolinense L.): Inbreeding impacts plant volatiles and insect attraction. Ecol. Lett. 15, 301–309. doi: 10.1111/j.1461-0248.2011.01738.x

Google Scholar: Author Only Title Only Author and Title

Keefover-Ring, K., Thompson, J. D., and Linhart, Y. B. (2009). Beyond six scents: defining a seventh Thymus vulgaris chemotype new to southern France by ethanol extraction. Flavour Fragr. J. 24, 117–122. doi: 10.1002/ffj.1921

Google Scholar: Author Only Title Only Author and Title

Kessler, A., and Kalske, A. (2018). Plant Secondary Metabolite Diversity and Species Interactions. Annu. Rev. Ecol. Evol. Syst. 49, 115–138. doi: 10.1146/annurev-ecolsys-110617-062406

Google Scholar: Author Only Title Only Author and Title

Kleine, S., and Müller, C. (2011). Intraspecific plant chemical diversity and its relation to herbivory. Oecologia 166, 175–186. doi: 10.1007/s00442-010-1827-6

Google Scholar: Author Only Title Only Author and Title

Kleine, S., and Müller, C. (2014). Drought stress and leaf herbivory affect root terpenoid concentrations and growth of Tanacetum vulgare. J. Chem. Ecol. 40, 1115–1125. doi: 10.1007/s10886-014-0505-2

Google Scholar: Author Only Title Only Author and Title

Lange, B. M. (2015). The evolution of plant secretory structures and emergence of terpenoid chemical diversity. Annu. Rev. Plant Biol. 66, 139–159. doi: 10.1146/annurev-arplant-043014-114639

Google Scholar: Author Only Title Only Author and Title

Lenth, R. V. (2023). emmeans: Estimated Marginal Means, aka Least-Squares Means. Available at: https://CRAN.R-project.org/package=emmeans

Google Scholar: Author Only Title Only Author and Title

Linhart, Y. B., Gauthier, P., Keefover-Ring, K., and Thompson, J. D. (2015). Variable phytotoxic effects of Thymus vulgaris (Lamiaceae) terpenes on associated species. Int. J. Plant Sci. 176, 20–30. doi: 10.1086/678772

Google Scholar: Author Only Title Only Author and Title

Linhart, Y. B., and Thompson, J. D. (1999). Thyme is of the essence: biochemical polymorphism and multi-species deterrence. Evol. Ecol. Res. 1, 151–171.

Google Scholar: Author Only Title Only Author and Title

Llandres, A. L., Almohamad, R., Brévault, T., Renou, A., Téréta, I., Jean, J., et al. (2018). Plant training for induced defense against insect pests: a promising tool for integrated pest management in cotton: Cotton training for induced defense against pests. Pest Manag. Sci. 74, 2004–2012. doi: 10.1002/ps.5039

Google Scholar: Author Only Title Only Author and Title

Loreto, F., and Schnitzler, J.-P. (2010). Abiotic stresses and induced BVOCs. Trends Plant Sci. 15, 154–166. doi: 10.1016/j.tplants.2009.12.006

Google Scholar: Author Only Title Only Author and Title

Loughrin, J. H., Manukian, A, Heath, R. R., and Tumlinson, J. H. (1995). Volatiles emitted by different cotton varieties damaged by feeding beet armyworm larvae. J. Chem. Ecol. 21, 1217–1227. doi: 10.1007/BF02228321

Google Scholar: Author Only Title Only Author and Title

Loughrin, J. H., Manukian, A, Heath, R. R., Turlings, T. C., and Tumlinson, J. H. (1994). Diurnal cycle of emission of induced volatile terpenoids by herbivore-injured cotton plant. Proc. Natl. Acad. Sci. 91, 11836–11840. doi: 10.1073/pnas.91.25.11836

Google Scholar: Author Only Title Only Author and Title

Mamin, M., Vallat, A, and Turlings, T. C. J. (2023). Cotton plants as ideal models for teaching and research on inducible direct plant defenses. Front. Ecol. Evol. 11. doi: 10.3389/fevo.2023.1119472

Google Scholar: Author Only Title Only Author and Title

McAuslane, H. J., and Alborn, H. T. (1998). Systemic induction of allelochemicals in glanded and glandless isogenic cotton by Spodoptera exigua feeding. J. Chem. Ecol. 24, 399–416. doi: 10.1023/A:1022596812049

Google Scholar: Author Only Title Only Author and Title

McAuslane, H. J., Alborn, H. T., and Toth, J. P. (1997). Systemic induction of terpenoid aldehydes in cotton pigment glands by feeding of larval Spodoptera exigua. J. Chem. Ecol. 23, 2861–2879. doi: 10.1023/A:1022575313325

Google Scholar: Author Only Title Only Author and Title

McCall, P. J., Turlings, T. C. J., Lewis, W. J., and Tumlinson, J. H. (1993). Role of plant volatiles in host location by the specialist parasitoid Microplitis croceipes cresson (Braconidae: Hymenoptera). J. Insect Behav. 6, 625–639. doi: 10.1007/BF01048128

Google Scholar: Author Only Title Only Author and Title

McCall, P. J., Turlings, T. C. J., Loughrin, J., Proveaux, A. T., and Tumlinson, J. H. (1994). Herbivore-induced volatile emissions from cotton (Gossypium hirsutum L.) seedlings. J. Chem. Ecol. 20, 3039–3050. doi: 10.1007/BF02033709

Google Scholar: Author Only Title Only Author and Title

McCormick, A. C., Reinecke, A, Gershenzon, J., and Unsicker, S. B. (2016). Feeding experience affects the behavioral response of polyphagous gypsy moth caterpillars to herbivore-induced poplar volatiles. J. Chem. Ecol. 42, 382–393. doi: 10.1007/s10886-016-0698-7

Google Scholar: Author Only Title Only Author and Title

Mitchell, C., Brennan, R. M., Graham, J., and Karley, A. J. (2016). Plant defense against herbivorous pests: exploiting resistance and tolerance traits for sustainable crop protection. Front. Plant Sci. 7, 1132. doi: 10.3389/fpls.2016.01132

Google Scholar: Author Only Title Only Author and Title

Moore, B. D., Andrew, R. L., Külheim, C., and Foley, W. J. (2014). Explaining intraspecific diversity in plant secondary metabolites in an ecological context. New Phytol. 201, 733–750. doi: 10.1111/nph.12526

Google Scholar: Author Only Title Only Author and Title

Müller, C., and Junker, R. R. (2022). Chemical phenotype as important and dynamic niche dimension of plants. New Phytol. 234, 1168–1174. doi: 10.1111/nph.18075

Google Scholar: Author Only Title Only Author and Title

Munguía-Rosas, M. A., Angulo, D. F., Arceo-Gómez, G., and Parra-Tabla, V. (2019). Variation in leaf traits across a precipitation gradient in coastal sand dunes in Yucatan Peninsula. J. Arid Environ. 162, 10–17. doi: 10.1016/j.jaridenv.2018.12.001

Google Scholar: Author Only Title Only Author and Title

Ngumbi, E., Chen, L., and Fadamiro, H. Y. (2009). Comparative GC-EAD responses of a specialist (Microplitis croceipes) and a generalist (Cotesia marginiventris) parasitoid to cotton volatiles induced by two caterpillar species. J. Chem. Ecol. 35, 1009–1020. doi: 10.1007/s10886-009-9700-y

Google Scholar: Author Only Title Only Author and Title

Oksanen, J., Simpson, G. L., Blanchet, F. G., Kindt, R., Legendre, P., Minchin, P. R., et al. (2022). vegan: Community Ecology Package. Available at: https://CRAN.R-project.org/package=vegan

Google Scholar: Author Only Title Only Author and Title

Opitz, S., Kunert, G., and Gershenzon, J. (2008). Increased terpenoid accumulation in cotton (Gossypium hirsutum) foliage is a general wound response. J. Chem. Ecol. 34, 508–522. doi: 10.1007/s10886-008-9453-z

Google Scholar: Author Only Title Only Author and Title

Palarea-Albaladejo, J., and Martin-Fernandez, J. A. (2015). zCompositions – R package for multivariate imputation of left-censored data under a compositional approach. Chemom. Intell. Lab. Syst. 143, 85–96.

Google Scholar: Author Only Title Only Author and Title

Paré, P. W., and Tumlinson, J. H. (1997). De novo biosynthesis of volatiles induced by insect herbivory in cotton plants. 114, 7. doi: 10.1016/S0031-9422(97)00442-1

Google Scholar: Author Only Title Only Author and Title

Pickett, J. A., and Khan, Z. R. (2016). Plant volatile-mediated signalling and its application in agriculture: successes and challenges. New Phytol. 212, 856–870. doi: 10.1111/nph.14274

Google Scholar: Author Only Title Only Author and Title

Quijano-Medina, T., Covelo, F., Moreira, X., and Abdala-Roberts, L. (2019). Compensation to simulated insect leaf herbivory in wild cotton (Gossypium hirsutum): responses to multiple levels of damage and associated traits. Plant Biol. 21, 805–812. doi: https://doi.org/10.1111/plb.13002

Google Scholar: Author Only Title Only Author and Title

Quijano-Medina, T., Turlings, T. C. J., Sosenski, P., Grandi, L., Cervera, J. C., Moreira, X., et al. (2021). Effects of soil salinity on the expression of direct and indirect defences in wild cotton Gossypium hirsutum. J. Ecol. 109, 354–368. doi: https://doi.org/10.1111/1365-2745.13483

Google Scholar: Author Only Title Only Author and Title

R Core Team (2022). R: A Language and Environment for Statistical Computing. Vienna, Austria: R Foundation for Statistical Computing. Available at: https://www.R-project.org/

Google Scholar: Author Only Title Only Author and Title

Rasmann, S., Buri, A., Gallot-Lavallée, M., Joaquim, J., Purcell, J., and Pellissier, L. (2014). Differential allocation and deployment of direct and indirect defences by Vicia sepium along elevation gradients. J. Ecol. 102, 930–938. doi: 10.1111/1365-2745.12253

Google Scholar: Author Only Title Only Author and Title

Razavizadeh, R., and Komatsu, S. (2018). Changes in essential oil and physiological parameters of callus and seedlings of Carum copticum L. under in vitro drought stress. J. Food Meas. Charact. 12, 1581–1592. doi: 10.1007/s11694-018-9773-9

Google Scholar: Author Only Title Only Author and Title

Rodriguez-Saona, C., and Crafts-Brandner, S. J. (2003). Volatile emissions triggered by multiple herbivore damage: beet armyworm and whitefly feeding on cotton plants. J. Chem. Ecol. doi: 10.1023/a:1026314102866

Google Scholar: Author Only Title Only Author and Title

Rodriguez-Saona, C. R., and Frost, C. J. (2010). New evidence for a multi-functional role of herbivore-induced plant volatiles in defense against herbivores. Plant Signal. Behav. 5, 58–60. doi: 10.4161/psb.5.1.10160

Google Scholar: Author Only Title Only Author and Title

Röse, U. S. R., Manukian, A, Heath, R. R., and Tumlinson, J. H. (1996). Volatile semiochemicals released from undamaged cotton leaves (A systemic response of living plants to caterpillar damage). Plant Physiol. 111, 487–495. doi: 10.1104/pp.111.2.487

Google Scholar: Author Only Title Only Author and Title

Röse, U. S. R., and Tumlinson, J. H. (2005). Systemic induction of volatile release in cotton: How specific is the signal to herbivory? Planta 222, 327–335. doi: 10.1007/s00425-005-1528-2

Google Scholar: Author Only Title Only Author and Title

Ruberto, G., and Baratta, M. T. (2000). Antioxidant activity of selected essential oil components in two lipid model systems. Food Chem. 69, 167–174. doi: 10.1016/S0308-8146(99)00247-2

Google Scholar: Author Only Title Only Author and Title

Schuman, M. C., Heinzel, N., Gaquerel, E., Svatos, A., and Baldwin, I. T. (2009). Polymorphism in jasmonate signaling partially accounts for the variety of volatiles produced by Nicotiana attenuata plants in a native population. New Phytol. 183, 1134–1148. doi: 10.1111/j.1469-8137.2009.02894.x

Google Scholar: Author Only Title Only Author and Title

Selmar, D., and Kleinwächter, M. (2013). Stress enhances the synthesis of secondary plant products: The impact of stress-related over-reduction on the accumulation of natural products. Plant Cell Physiol. 54, 817–826. doi: 10.1093/pcp/pct054

Google Scholar: Author Only Title Only Author and Title

Senft, M., Clancy, M. V., Weisser, W. W., Schnitzler, J., and Zytynska, S. E. (2019). Additive effects of plant chemotype, mutualistic ants and predators on aphid performance and survival. Funct. Ecol. 33, 139–151. doi: 10.1111/1365-2435.13227

Google Scholar: Author Only Title Only Author and Title

Sørensen, M., Neilson, E. H. J., and Møller, B. L. (2018). Oximes: Unrecognized chameleons in general and specialized plant metabolism. Mol. Plant 11, 95–117. doi: 10.1016/j.molp.2017.12.014

Google Scholar: Author Only Title Only Author and Title

Stipanovic, R. D., Bell, A. A, O'Brien, D. H., and Lukefahr, M. J. (1978). Heliocide H1. A new insecticidal C25 terpenoid from cotton (Gossypium hirsutum). J. Agric. Food Chem. 26, 115–118.

Google Scholar: Author Only Title Only Author and Title

Taft, S., Najar, A, Godbout, J., Bousquet, J., and Erbilgin, N. (2015). Variations in foliar monoterpenes across the range of jack pine reveal three widespread chemotypes: implications to host expansion of invasive mountain pine beetle. Front. Plant Sci. 6. doi: 10.3389/fpls.2015.00342

Google Scholar: Author Only Title Only Author and Title

Tian, Z, Luo, Q., Li, Y., and Zuo, Z (2020). Terpinene and β-pinene acting as signaling molecules to improve Cinnamomum camphora thermotolerance. Ind. Crops Prod. 154, 112641. doi: 10.1016/j.indcrop.2020.112641

Google Scholar: Author Only Title Only Author and Title

Turlings, T. C., and Erb, M. (2018). Tritrophic interactions mediated by herbivore-induced plant volatiles: mechanisms, ecological relevance, and application potential. Annu. Rev. Entomol. 63, 433–452. doi: 10.1146/annurev-ento-020117-043507

Google Scholar: Author Only Title Only Author and Title

Unsicker, S. B., Kunert, G., and Gershenzon, J. (2009). Protective perfumes: the role of vegetative volatiles in plant defense against herbivores. Curr. Opin. Plant Biol. 12, 479–485. doi: 10.1016/j.pbi.2009.04.001

Google Scholar: Author Only Title Only Author and Title

Villamar-Torres, R., Mehdi Jazayeri, S., Liuba-Delfini, G., García Cruzaty, L. C., and Viot, C.-R. (2018). Volatile organic compounds: plant natural defense mechanisms against herbivorous arthropods and an opportunity for plant breeding of cotton. Sci. Agropecu. 9, 287–297. doi: 10.17268/sci.agropecu.2018.01.14

Google Scholar: Author Only Title Only Author and Title

Voelker, J., Mauleon, R., and Shepherd, M. (2023). The terpene synthase genes of Melaleuca alternifolia (tea tree) and comparative gene family analysis among Myrtaceae essential oil crops. Plant Syst. Evol. 309, 13. doi: 10.1007/s00606-023-01847-1

Google Scholar: Author Only Title Only Author and Title

Wason, E. L., Agrawal, A. A, and Hunter, M. D. (2013). A genetically-based latitudinal cline in the emission of herbivore-induced plant volatile oganic compounds. J. Chem. Ecol. 39, 1101–1111. doi: 10.1007/s10886-013-0309-9

Google Scholar: Author Only Title Only Author and Title

Wegier, A, Piñeyro-Nelson, A, Alarcón, J., Gálvez-Mariscal, A, Álvarez-Buylla, E., and Piñero, D. (2011). Recent long-distance transgene flow into wild populations conforms to historical patterns of gene flow in cotton (Gossypium hirsutum) at its centre of origin. Mol. Ecol. 20, 4182–4194. doi: 10.1111/j.1365-294X.2011.05258.x

Google Scholar: Author Only Title Only Author and Title

Wu, J., Hettenhausen, C., Schuman, M. C., and Baldwin, I. T. (2008). A comparison of two Nicotiana attenuata accessions reveals large differences in signaling induced by oral secretions of the specialist herbivore Manduca sexta. Plant Physiol. 146, 927–939. doi: 10.1104/pp.107.114785

Google Scholar: Author Only Title Only Author and Title

Xu, S., Kreitzer, C., McGale, E., Lackus, N. D., Guo, H., Köllner, T. G., et al. (2020). Allelic differences of clustered terpene synthases contribute to correlated intraspecific variation of floral and herbivory-induced volatiles in a wild tobacco. New Phytol. 228, 1083–1096. doi: 10.1111/nph.16739

Google Scholar: Author Only Title Only Author and Title

Yuan, D., Grover, C. E., Hu, G., Pan, M., Miller, E. R., Conover, J. L., et al. (2021). Parallel and intertwining threads of domestication in allopolyploid cotton. Adv. Sci. 8, 2003634. doi: 10.1002/advs.202003634

Google Scholar: Author Only Title Only Author and Title

Zakir, A, Sadek, M. M., Bengtsson, M., Hansson, B. S., Witzgall, P., and Anderson, P. (2013). Herbivore-induced plant volatiles provide associational resistance against an ovipositing herbivore. J. Ecol. 101, 410–417. doi: 10.1111/1365-2745.12041

Google Scholar: Author Only Title Only Author and Title

Zhang, L., Lu, G., Huang, X., Guo, H., Su, X., Han, L., et al. (2020). Overexpression of the caryophyllene synthase gene GhTPS1 in cotton negatively affects multiple pests while attracting parasitoids. Pest Manag. Sci. 76, 1722–1730. doi: 10.1002/ps.5695

Google Scholar: Author Only Title Only Author and Title