Spirolactone, an unprecedented antifungal β-lactone spiroketal macrolide from *Streptomyces iranensis*

Zhijie Yang,[a] Yijun Qiao,[a] Emil Strøbech,[a] Jens Preben Morth,[a] Grit Walther,[b,c] Tue Sparholt Jørgensen,[d] Gundela Peschel,[e] Miriam A. Rosenbaum,[e] Viola Previtali,[f] Mads Hartvig Clausen,[f] Marie Vestergaard Lukassen,[a] Charlotte H. Gotfredsen,[g] Oliver Kurzai,[b,c] Tilmann Weber,[d] Ling Ding[a,]*

[a] Department of Biotechnology and Biomedicine, Technical University of Denmark, Søltofts Plads 221, 2800, Kgs Lyngby, Denmark.

[b] German National Reference Center for Invasive Fungal Infections, Leibniz Institute for Natural Product Research and Infection Biology – Hans Knöll Institute, Beutenbergstraße 11A 07745, Jena, Germany.

[c] Institute for Hygiene and Microbiology, University of Würzburg, Würzburg, Germany.

[d] The Novo Nordisk Foundation Center for Biosustainability, Technical University of Denmark, Søltofts Plads 220, 2800, Kgs Lyngby, Denmark.

[e] BioPilot Plant, Leibniz Institute for Natural Product Research and Infection Biology – Hans Knöll Institute, Beutenbergstraße 11A, 07745, Jena, Germany.

[f] Center for Nanomedicine and Theranostics, Department of Chemistry, Technical University of Denmark, Kemitorvet 207, 2800, Kgs. Lyngby, Denmark

[g] Department of Chemistry, Technical University of Denmark, Kemitorvet 207, 2800, Kgs. Lyngby, Denmark

*Corresponding author: Ling Ding lidi@dtu.dk

Abstract

Fungal infections pose a great threat to public health. There are only four classes of antifungals that have limitations due to high toxicity, drug-drug interactions, and emerging drug-resistance. *Streptomyces* spp. represent an important source of antimicrobial substances, notably including the antifungal agent amphotericin B. The rapamycin-producer *Streptomyces iranensis* displayed strong antifungal activities against *Aspergillus*. Revisiting its genome revealed several intriguing biosynthetic gene clusters, including one unparalleled Type I polyketide synthase, which codes for uncharacterized metabolites. The identification of a novel macrolide spirolactone (**1**) and its biosynthetic gene cluster was facilitated through CRISPR-based gene editing, HR-ESI-MS analysis, followed by fermentation and purification processes. Their structures and absolute configurations were confirmed by NMR, MS and X-ray crystallography. Spirolactone harbors an undescribed carbon skeleton with 13 chiral centers, featuring a rare *β*-lactone moiety, a [6,6]-spiroketal ring, and an unprecedented 7-oxo-octylmalonyl-CoA extender unit incorporated by a potential novel *Crotonyl*-CoA carboxylase/reductase. Spirolactone displayed profound antifungal effects against numerous fungal pathogens, e.g. the genus *Talaromyces* and several sections of *Aspergillus* including clinically relevant species such as *Aspergillus niger* and *A. tubingensis* (section Nigri), *A. terreus* (section Terrei) and the azol-resistant *A. calidoustus* (section Usti). Proteomics analysis revealed spirolactone potentially disrupted the integrity of fungal cell walls and induced the expression of stress-response proteins in *A. niger*. Spirolactone represents a new class of potential drug candidate to combat fungal infections.

Introduction

About 1.2 billion people worldwide are estimated to suffer from a fungal disease.[1] Invasive fungal diseases are particularly common among immunocompromised populations, such as patients with HIV/AIDS, chronic lung diseases, prior tuberculosis, cancer, diabetes, and patients receiving immunosuppressant treatment and invasive medical procedures.[2] Invasive fungal diseases are associated with unacceptably high mortality rates. The recent report suggested an annual incidence of 6.5 million invasive fungal infections and 3.8 million deaths.[3] Among those infections, *Aspergillus* has emerged as one of the most common causes

of infectious death in severely immunocompromised patients, with mortality rates up to 40% to 50% in patients with acute leukemia and recipients of hematopoietic stem cell transplantation.[1] Furthermore, new risk cohorts are emerging, including patients with severe respiratory virus infection such as influenza or COVID-19.[4] Currently, only four classes of systemic antifungal medicines (azoles, echinocandins, pyrimidines, and polyenes) are used in clinical practice, and only a limited number of candidates in the clinical development pipeline.[5-8] While existing antifungal medications are effective in numerous instances, challenges persist due to side effects and complications arising from drug-drug interactions.[9] Moreover, the emergence of resistance has rendered some existing medications ineffective, and this is partly driven by the inappropriate use of antifungals.[10] For example, agricultural use of azoles has been linked to rising rates of azole-resistant *Aspergillus fumigatus* infections, with resistance rates of 15-20% reported in parts of Europe and exceeding 80% in environmental samples from Asia.[10-12]

Streptomyces are producers of bioactive small molecules exhibiting a broad range of structural and functional diversity called secondary or specialised metabolites (SMs). Secondary metabolites are of great relevance to human health, offering pharmaceutical properties such as antibacterial, antifungal, anticancer, and immunosuppressive activities. Remarkably, all new drug approvals from 1981-2019 over 60% of the 1394 small molecule drugs were either secondary metabolites (SMs) or derivatives thereof.[13-15] Nearly two-thirds of antibiotics approved for clinical use originate from *Streptomyces*, and of Polyketides represent a large and diverse group of *Streptomyces*-derived SMs, and many of them possess valuable antifungal properties, among which nystatin and amphotericin B are two active polyenes used clinically as first-line drugs to treat fungal infections. The first example is the antifungal drug nystatin isolated from *Streptomyces noursei* in 1950 by Hazen and Brown.[16] Amphotericin B, isolated from *Streptomyces nodosus* in 1955, remains a crucial component in the therapeutic arsenal for combating invasive fungal diseases and leishmaniasis.[17] Recent studies revealed that the mode of action of polyenes antibiotics involves acting as sterol "sponges" and extracting ergosterol from the fungal membrane, causing cell death, in addition to the classical model of pore formation in the fungal cell membrane.[18-20] Amphotericin B has been used to treat mucormycosis, aspergillosis, blastomycosis,

candidiasis, coccidioidomycosis, and cryptococcosis. However, adverse effects for amphotericin B treatment are common, with nephrotoxicity being the most serious.[21,22] There has been a continued interest in finding new antifungal metabolites, such as turonicin A, hygrobafilomycin, iseolides, cyphomycin, azalomycins, niphimycins, and resistomycin.[23-29] However, most exhibit non-selectivity and display toxicity to human cells. Thus, it is of clinical importance to develop effective selective antifungal drug candidates, with a new mode-of-action and low cytotoxicity.

Given the significant number of biosynthetic gene clusters (BGCs) present in *Streptomyces*, a plethora of their metabolites remains undiscovered.[30] Access to modern analytical chemistry techniques in combination with genetics and bioinformatics tools has enabled us to revisit the "talented" microbes discovered decades ago.[31] *Streptomyces iranensis*, initially identified as a rapamycin producer, was isolated from soil in Isfahan City, Iran.[32] Subsequent investigations revealed its great capability to produce elaiophylin, azalomycins, nigericin, and other new metabolites yet to be discovered.[33] In our recent research, we found that *S. iranensis* produced pteridic acids H and F, which aid plants in coping with abiotic stress.[33] During our pipeline search for antifungal polyketides, *S. iranensis* exhibited profound antifungal activity in co-cultivation with *Aspergillus tubingensis* (*Supplementary information* Fig. 1), suggesting the production of antifungal metabolites. Genome mining of *S. iranensis* indicated the presence of an uncharacterized type-I modular polyketides BGC (*spl*) featuring sixteen modules, capturing our particular interest. The product of this BGC namely spirolactone was carried out through metabolomic analysis, large-scale fermentation and isolation. BGC (*spl*) was confirmed by CRISPR-based gene mutation. The structures and absolute configurations of spirolactone was elucidated using nuclear magnetic resonance (NMR), mass spectrometry (MS), and X-ray crystallography. The striking structural features of spirolactone include a spiroketal ring, a rare β-lactone moiety, and a distinctive polyketide synthetase (PKS) extender unit. Spirolactone exhibits potent antifungal activity, potentially due to disrupting fungal cell wall integrity, as suggested by label-free quantitative proteomics analysis. This study provides a paradigm for mining novel molecules from extensive natural microorganisms, with the potential to lead to the discovery of innovative antifungal agents to combat antimicrobial resistance.

Results and Discussion

Genome mining of *S. iranensis*. Due to the incompleteness of previously reported genomic data, we performed whole-genome resequencing of *S. iranensis* using the combination of the Oxford Nanopore Technologies MinION and Illumina MiSeq system. The high-resolution, full-genome sequencing of *S. iranensis* reveals a linear chromosome spanning 12,213,033 nucleotides, featuring inverted terminal repeats comprising 156,145 nucleotides. The BGC annotation of the acquired *S. iranensis* genome was carried out using antiSMASH version 7.0,[34] which resulted in the identification of a cryptic BGC (*spl* gene cluster) situated at the terminal region of the linear chromosome (Fig. 1 and *Supplementary information* Fig. 2). The gene cluster family (GCF) annotation of *spl* gene cluster showed it belongs to GCF_00315 and a total of 11 hits were detected (distance ≤ 900.0) in the BiG-FAM database (*Supplementary information* Tab. 2).[35] The distinctive biosynthetic gene architecture of *spl* gene cluster suggests its novelty as a modular Type-I PKS BGC, setting it apart from others (*Supplementary information* Fig. 3). This finding spurred our interest in conducting a thorough investigation of this individual BGC.

Fig. 1. Organization of the spirolactone biosynthetic gene cluster from *S. iranensis*. ORFs are shown as arrows indicating the size and direction of transcription.

To locate the products of the potential *spl* gene cluster, we were elucidating the products combining both genome editing and metabolomics. Initially, we constructed PKS knock-out mutant (Δ*splA*) and crotonyl-CoA carboxylase/reductase-inactivated mutant (Δ*splH*) using the CRISPR-Cas9 method and CRISPR base editing,[36] respectively (**Fig. 2**). Both mutants nearly abolished the antifungal activity against multiple *Aspergillus* strains. Comparative

metabolic profiling using LC-MS and LC-UV led to the identification of the potential bioactive principle as a major compound **1** with *m/z* 881.4656 ([M + Na]⁺) and other derivatives, which were specifically abolished in both mutants (**Fig. 2**). The complementation of pGM1190-based *splH* in mutant *S. iranensis*/Δ*splH* led to the restoration of the production of **1** (**Fig. 2**).

Fig. 2. Conformation of the polyketide biosynthetic gene cluster through inactivation of *splA* (**a**) and inactivation of *splH* (**b**) in *S. iranensis*. Extracted Ion Chromatogram (*m/z* 881.4656 [M + Na]$^+$) showing the abolishment of production of spirolatone in the mutants (*ΔsplA* and *ΔsplH*) compared to the wild strain.

Characterization of spirolactone. To obtain the products of the *spl* BGC, a 200 L fermentation was carried out followed by a downstream processing using the open-column chromatography on Amberchrom 161c resin, silica gel, Sephadex LH-20, and High-Performance Liquid Chromatography (HPLC) to yield pure spirolactone (**1**). The structure was elucidated by HR-ECI-MS, NMR spectroscopy, and X-ray crystallography.

Spirolactone (**1**) was obtained as a white solid and a molecular formula of $C_{47}H_{70}O_{14}$ by a sodium adduct ion (calcd *m/z* 881.4658 [M + Na]$^+$) was determined by HR-ESI-MS. It was assigned to the molecular formula on HR-ESI-MS combined with the NMR data. ^{1}H NMR spectrum revealed signals for six methyl groups H3-39 (δ 2.15), H3-41 (δ 0.94), H3-43 (δ 0.98), H3-44 (δ 0.93), H3-45 (δ 2.10), and OCH$_3$-8 (δ 3.37). Additionally, four olefinic protons were observed for H-2 (δ 6.19), H-3 (δ 7.91), H-4 (δ 6.47) and H-28 (δ 6.48). The coupling constant between H-2 and H-3 (*J* 15.4 Hz) confirmed a trans-orientation of a double bond. Besides the presence of several oxygen-bearing methines (H-6, H-8, H-10, H-14, H-16, and H-22), various other aliphatic proton signals were also observed (*Supplementary information* Fig. 5).

The ^{13}C NMR spectrum showed signals for several carbonyl groups, one carboxylic acid C-1 (δ 168.6), three keto- groups C-12 (δ 207.3), C-27 (δ 201.3) and C-38 (δ 209.6) and two ester groups C-40 (δ 161.8) and C-46 (δ 174.9), with C-40 from the β-lactone ring. Furthermore, signals were observed for six olefinic methines C-2 (δ 128.5), C-3 (δ 137.8), C-4 (δ 129.1), C-5 (δ 143.2), C-28 (δ 123.0), and C-29 (δ 156.4). Finally, signals for the methyl groups C-39 (δ 29.9), C-41 (δ 8.4), C-42 (δ 9.2), C-43 (δ 14.3), C-44 (δ 19.5), C-45 (δ 17.4) and the oxygenated methyl group 8-OMe (δ 59.6) connected to C-8 were also observed (*Supplementary information* Fig. 6). The chemical shift of C-18 (δ 101.0) indicated the presence of an acetal moiety. The ^{1}H and ^{13}C NMR data of **1** is shown in *Supplementary information*, Tab. 4.

COSY correlations further established six fragments including one saturated aliphatic chain. HMBC correlations (**Fig. 3**) further established a *β*-lactone moiety as HMBC correlations

between H-6 and C-40 were observed. A further connection of this β-lactone to a polyene moiety was established by HMBC correlations between H-4 and C-6, as well as H-4 and C-40. A macrolactone bridge between H-10 and C-46 was established by HMBC correlation between them. The key HMBC correlation between H-13 and C-18 indicated the presence of an oxane ring. Moreover, H-16 and Me-43 showed HMBC correlations with C-18. Considering the double bond equivalence and the number of oxygen atoms inferred from HR-ESI-MS, a spiroketal structure was proposed. Finally, a planar structure with a novel carbon skeleton was proposed and named spirolactone. The complex structure exhibited no similarities to reported natural products. Meanwhile, the dispersion of the thirteen chiral centers made it challenging to elucidate the absolute configuration. Interestingly, the compound could form crystals from a dichloromethane/methanol solution. One crystal was subjected to X-ray crystallography analysis, which successfully elucidated the configurations (**Fig. 3**).

Fig. 3. Structure elucidation of spirolactone (**1**). a: Chemical structure of **1** and selected ^{1}H-^{1}H COSY and HMBC correlations; b: ORTEP drawing of compound **1**.

Proposed biosynthesis of spirolactone. The proposed *spl* gene cluster spanned 87.99 kb with 12 open reading frames (Fig. 1), including five core polyketide synthases (*splA*, *splC*, *splD*, *splE*, and *splF*), a crotonyl-CoA carboxylase/reductase (*splH*), an *O*-methyltransferase (*splG*), three cytochrome P450 (*splB*, *splK* and *splL*), and two hypothetical enzymes (*splI* and *splJ*) (*Supplementary information* Tab. 3). The core PKS genes transcribed in the same direction encoding genes within the *spl* BGC consist of one loading module and fifteen extender modules, which are sequentially extended to form a linear polyketide chain by catalysis of ketosynthase (KS) domain, acyltransferase (AT) domain, and acyl carrier protein (ACP), with additional ketoreductase (KR), dehydratase (DH), and enoyl reductase (ER) domains. The proposed biosynthesis is shown in **Fig. 4**. Based on multiple sequence alignment, most KS domains keep the conserved catalytic sites comprised of a cysteine (TA**C**SSS motif) as well as two histidines (EA**H**GTG and KSNIG**H**T motifs).[37] Only the reactive cysteine in KS$_1$ was replaced by glutamine (**Fig. 4**), which was observed in the platensimycin/FabF (the type II FAS homology of KS) complex.[38] The conserved fingerprint

residues for extender unit selectivity, the GHSIG and HAFH motifs are present in the nine AT domains (AT_1, AT_2, AT_5, AT_7, AT_8, AT_9, AT_{11}, AT_{12}, and AT_{14}) that are specifically binding to malonyl-CoA while GHSQG and YASH motifs are present in the six AT domains (AT_3, AT_4, AT_6, AT_{10}, AT_{13}, and AT_{15}) that are specifically binding to (*2S*)-methylmalonyl-CoA (*Supplementary information* Tab. 5 and Fig. 14).[37] AT_{16} from module 15, which was thought to be accountable for loading distinct extender units, exhibits an unusual IASH motif. KR domains have been previously classified into A1-, A2-, B1-, and B2-types, which reduce ketones to their L-hydroxy or D-hydroxy counterparts.[39] We determined that the KR_3, KR_4, and KR_{15} appear to belong to the B1-type: all have the fingerprint LDD motif but the absence of a P residue in the catalytic region. KR_5 and KR_{12}, with the characteristic W residue but no LDD motif and H residue, were classified as A1-type KRs (*Supplementary information* Fig. 15). The other KR domains couldn't be classified according to the sequence, but they should be functional according to the structure of spirolactone. Similarly, combined with structural information and conserved site analysis, the five DH domains (DH_6, DH_7, DH_8, DH_9 and DH_{10}) were considered inactive during spirolactone biosynthesis (*Supplementary information* Fig. 16). ER_{11} is classified as L-type due to the presence of a tyrosine residue that donates a proton to the enol intermediate in L-type ER, whereas ER_{13} and ER_{14} belong to D-type because it lacks tyrosine residue (*Supplementary information* Fig. 17). The final release and cyclization of the linear product was probably accomplished by TE domain in the last module, which contains an α/β-hydrolase catalytic core and loop regions that form a substrate-binding lid.[40]

Fig. 4. Proposed biosynthesis pathway of spirolactone (**1**)

In the *spl* BGC, there are three cytochromes P450 enzymes (SplB, SplK, and SplL) with conserved motifs of this family (*Supplementary information* Fig. 18).[41] The cytochrome P_{450} enzyme BonL from *Burkholderia gladioli* was previously identified to confer the C-22 carboxyl group in the biosynthesis of bongkrekic acid via sequential six-electron-oxidation.[42] In addition, there are some cytochromes P_{450} enzymes that catalyze the carboxyl group formation in microbial SMs, such as XiaM, PimG, AmphN, NysN, and FscP.[43-45] To identify the cytochromes P450s responsible for carboxylation, we built the Hidden Markov model (HMM) based on these sequences and the results showed that SplL exhibited the best match with these known P450s with an E-value of 4e-117. In addition, bioinformatic analysis of XiaM indicated that the P450s with carboxylation function harbor highly conserved segment AGHET, which was also presented in SplK (*Supplementary information* Fig. 18). Therefore, SplK and SplL, were considered putative candidates for catalyzing the two-step oxidation to form carboxyl groups at positions C-1 and C-5 during spirolactone biosynthesis. SplB, homologous cytochrome P450 monooxygenase CftA in clifednamide biosynthesis gene cluster with 45.6% similarity,[46] was likely to accomplish the oxidation at C-27 in spirolactone. We constructed mutant strains of *S. iranensis* inactivating the SplB, SplK and SplL enzymes (denoted as Δ*splB*, Δ*splK*, and Δ*splL*) (*Supplementary information* Fig. 19).

Despite our efforts, no expected intermediates were observed in either mutant, likely due to their low yield.

Currently, there are only few enzymatic mechanisms corresponding to β-lactone ring formation that have been reported,[47] including: (1) the intramolecular cyclization from seven-membered ring, catalyzed by cyclase VibC in vibralactone biosynthesis;[48] (2) the tandem aldol-lactonization bicyclization reaction to generate the γ-lactam-β-lactone structure, catalyzed by standalone ketosynthase SalC in salinosporamide A biosynthesis;[49] (3) the β-lactone formation during the intramolecular attack of the β-hydroxyl group onto the thioester carbonyl, catalyzed by the C-terminal TE domain of ObiF in obafluorin biosynthesis or esterase GloD in globilactone A biosynthesis;[50,51] (4) the conversion of β-hydroxyl to β-lactone, catalyzed by β-lactone synthase OleC in olefin biosynthesis.[52,53] We proposed that the beta-lactone formation in spirolactone involves a two-step oxidation to form a carboxylic acid, followed by dehydration. It is likely that either the cytochrome P450 enzyme SplK or SplL is involved in the process, and we also identified several genes with unknown functions, such as *splJ* and *splI*. SplJ was annotated as a member of UbiD family decarboxylases, but it exhibits very distinguished divergences from other proteins in this family based on sequence similarity network (*Supplementary information* Fig. 20). In addition, NCBI Blastp results showed that proteins homologous to SplI have not been reported to have any clear biological functions. Inactivation of both genes abolished the production of spirolactone (Supplementary information Fig. 19).

Phylogenetic analysis indicated that SplH is a separate branch and clusters with other crotonyl-CoA reductase/carboxylases (CCRs) that catalyze long-chain precursors like butylmalonyl-CoA and hexylmalonyl-CoA.[54,55] The previous study showed that the large residue Phe380 in the CCR from *S. coelicolor* may constrain the potential pocket to accept long-chain substrates, which were also found in most ethylmalonyl-CoA specific crotonyl-CoA reductase/carboxylases, while its corresponding residue in SplH is smaller cysteine (*Supplementary information* Fig. 21).[56] Hence, we proposed that SplH catalyzes the conversion leading to the formation of 7-oxo-octylmalonyl-CoA in spirolactone, which is likely synthesized from a precursor that has previously been isolated from other *Streptomyces* species.[57] SplG, an O-methyltransferase homologous to AveD in avermectin biosynthesis

(40% identity and 55% similarity),[58] was proposed to catalyze the methylation of a hydroxyl group in C-8 of spirolactone.

Further investigation on the key enzymes involved in the biosynthesis of spirolactone will be carried out.

Antifungal and cytotoxic assays of spirolactone. Microbial β-lactone natural products are chemically diverse and have been employed in antimicrobial, antiviral, anticancer, and antiobesity therapeutics.[47] In the initial antifungal assay against *A. niger*, spirolactone exhibited the strongest antifungal effect. *In vitro* antifungal susceptibility of spirolactone against 34 fungal species represented by 38 clinical isolates was further tested in comparison to approved antifungals using a microdilution assay according to the EUCAST protocol. Spirolactone demonstrated antifungal effect against several sections of the genus *Aspergillus* and the genus *Talaromyces* (Tab. 1, *Supplementary information* Tab. 6). The lowest MICs for spirolactone were found in the *Aspergillus* section *Terrei* with intrinsically high MICs for amphotericin B and in section *Usti* with intrinsically high MICs for azoles. In the *Aspergillus* sections *Flavi*, *Nidulantes*, and *Nigri* as well as in the genus *Talaromyces* a distinctly changed morphology of the hyphae was found for all species but low MICs were observed only in some species. Only a weak effect was detected for the two *Fusarium* species studied. *Aspergillus* species belonging to section *Fumigati*, *Candida* spp., *Scedosporium prolificans*, *Purpureocillium lilacinum*, *Trichoderma longibrachiatum* and Mucorales were not affected by spirolactone (*Supplementary information* Tab. 6). Spirolactone did not exhibited cytotoxicity against the human HL-60 (acute promyelocytic leukemia) cell line (IC$_{50}$ >10 µg/mL).

Tab. 1 Antifungal activities of spirolactone (**1**) in comparison with amphotericin B, itraconazole and voriconazole, The values for MIC (MEC) are in mg/L.

Species	Section	amphotericin B	itraconazole	voriconazole M	spirolactone
A. brasiliensis	Nigri	0.5	2	2	0.5 (0.25)
A. luchuensis	Nigri	0.25	n.a.	1	0.5 (0.25)
A. citrinoterreus	Terrei	4	0.5	0.5	0.5 (0.25)
A. neoafricanus	Terrei	2	0.5	0.5	0.5 (0.25)
A. terreus	Terrei	8	0.5	1	0.125 (0.125)
A. calidoustus	Usti	16	8	4	0.5 (0.25)
A. ustus	Usti	1	>8	>8	0.125 (0.06)
T. marneffei		0.06	≤0.016	≤0.016	0.125 (0.125)
T. purpureogenes		1	>8	8	0.5 (0.25)

MIC: minimum inhibitory concentration.

Studies of the mechanism of action of spirolactone. To get a clue of how spirolactone functions and how it impacts *Aspergillus*, we carried out a label-free quantitative proteomics analysis of model organism *A. niger* ATCC 1015. We initially cultured *A. niger* in PDB liquid medium for 16 hours followed by the 1-hour and 4-hour treatment with spirolactone. The proteomics analysis showed that a total of 4,447 proteins (40.4% of the total encoded proteins) were detected, and 3,992 proteins (89.8% of the total detected proteins) didn't show significant differences between the treatment (1-hour or 4-hour treatment with spirolactone) and the control (without spirolactone treatment). Compared with the control, the abundance of 134 proteins significantly increased and 60 proteins significantly decreased after 1-hour spirolactone treatment. In comparison, the abundance of 273 proteins significantly increased and 113 proteins significantly decreased after 4-hour spirolactone treatment (Fig. 5). The Cluster of Orthologous Groups (COGs) functional classification of these differential proteins was performed by eggNOG-mapper v2.[59] Except for (S) function unknown, these proteins were mainly clustered into a few biological processes including (Q) Secondary structure, (C) Energy production and conversion, (E) Amino acid metabolism and transport, and (O) Post-translational modifications, protein turnover, chaperone functions (*Supplementary information* Fig. 22). The Gene Ontology (GO) annotation of these proteins was mainly involved in the metabolic process, cellular process, and single-organism process during the biological process (*Supplementary information* Fig. 23). Cellular component analysis showed that these proteins were mainly localized intracellularly, especially in organelles. Meanwhile,

most of these proteins participated in catalytic activity and binding in the category of molecular function. The Kyoto Encyclopedia of Genes and Genomes (KEGG) pathway enrichment analysis showed that these proteins mainly participate in amino acids metabolism and purine metabolism (*Supplementary information* Fig. 24).

Proteomic changes in fungi induced by known antifungal agents with well-described targets were subsequently studied. The abundance of Erg13 (EHA22977.1, involved in ergosterol biosynthesis) and manganese-superoxide dismutase (EHA27012.1, involved in oxidative stress response), which was regarded as polyene antibiotic (such as amphotericin B) responsive protein,[60] was not significantly altered in the presence of spirolactone. Allylamine antibiotics (naftifine and terbinafine) inhibit squalene epoxidase, causing the accumulation of squalene, and thereby damaging the intracellular membranes of fungi.[61] However, spirolactone is not belonging to the allylamine family and didn't induce any changes in the squalene epoxidase (EHA21184.1). Azoles are synthetic antifungal agents of non-natural origin that inhibit cytochrome P450 sterol-14α-demethylase (CYP51) to block the ergosterol biosynthesis in the fungi.[62] Similarly, spirolactone lacked the azoles substructure and had no impact on the abundance of CYP51 (EHA19435.1). Natural echinocandins comprise cyclic hexapeptide and lipid residues, like caspofungin, anidulafungin, and micafungin, preventing the biosynthesis of glucans of the fungal cell wall through non-competitive inhibition of 1,3-β-D-glucan synthase.[63] Previous proteomics analyses had proven that the level of chitinase ChiA1 in *A. fumigatus* was significantly decreased in response to caspofungin.[64] It may represent a common effect of self-resistance development due to caspofungin increased chitin content via induction of chitin synthases during cell wall remodeling.[65] We did not observe the abundance of 1,3-β-D-glucan synthase (EHA18547.1) and ChiA1-like chitinase (EHA28582.1) altered significantly after 1-hour and 4-hour treatment of spirolactone in comparison to the control group.

Fig. 5 Volcano plot of the abundance of protein after spirolactone treatment based on proteomics analysis.

The fungal cell wall is an essential structure that maintains cell shape and protects fungi against environmental stress and play roles in growth, invading ecological niches, and counteracting the host immune response.[66] Fungal cell walls are composed mainly of glucans, chitin, and glycoproteins, synthesized by glycosyltransferases, glycoside hydrolases, and transglycosylases.[67] Congo Red Hypersensitivity (CRH) family transglycosylases are usually highly expressed in multiple stages during the conidial germination of fungi and may be involved in cell wall synthesis and stability. The volcano plot of differential proteins indicated that a CRH family transglycosylase (EHA23738.1) was significantly inhibited, suggesting spirolactone might interfere with the formation of the fungal cell wall. Trehalose-6-phosphate synthase/phosphatase is another important enzyme for cell wall integrity and fungal virulence in various *Aspergillus* species.[68] The relative abundance of trehalose 6-phosphate synthase (EHA27700.1) was also significantly decreased after spirolactone treatment. The above data indicated that the mode of action of spirolactone could be through disruption of fungal cell wall biosynthesis and stability (Fig. 5). Besides, we found that the two enzymes Erg24 (EHA26587.1) and Erg27 (EHA20711.1), which catalyze the biosynthesis of 4,4-dimethylcholesta-8,14,24-trienol to fecosterol,[69] were remarkably upregulated after spirolactone treatment (Fig. 5). Drug efflux and resistance in fungi could be mediated by ATP-binding cassette (ABC) transporters, such as CDR1 was induced to high expression in *A. fumigatus* AF293 in the presence of azole antibiotics.[70] The expression of an

ABC transporter (EHA25684.1) was activated after spirolactone treatment. We observed the abundance of several putative spirolactone-responsive proteins were significantly altered. For example, the relative abundance of several proteins including inositol hexaphosphate kinase KCS1 (EHA27881.1), ubiquitin-protein ligase ASI3 (EHA19328.1), mitochondrial phosphate carrier protein PIC2 (EHA22842.1), pyridoxamine 5'-phosphate oxidase (EHA19800.1) and glucohydrolase (EHA21384.1) was significantly increased, but oxygen-dependent FAD-linked oxidoreductase (EHA22577.1) and dienelactone hydrolase (EHA27638.1) was significantly decreased in spirolactone-treated samples. Interestingly, an amount of unknown function proteins was found to increase or decrease in abundance affected by spirolactone and their functions still need to be further explored.

Conclusion

Herein, we report the discovery of a novel class of antifungal agent spirolactone, featuring an unprecedented carbon skeleton. Spirolactone exhibits selective antifungal activities against several pathogenic *Aspergillus* species, some of which are intrinsically resistant to amphotericin B or azoles, as well as against *Talaromyces* species. We confirmed the BGC of spirolactone through a variety of in silico and *in vivo* analyses. Proteomics data revealed that treatment with spirolactone leads to the inhibition of transglycosylase and trehalose-6-phosphate synthase/phosphatase, suggesting that it may target fungal cell wall biosynthesis.

Methods and Materials

Strains, plasmids, and culture conditions. All strains and plasmids used in this study were summarized in *Supplementary information* Tab. 7. All primers used in this study were summarized in *Supplementary information* Tab. 8. All constructed *E. coli* strains were growing on lysogeny broth (LB) liquid or agar medium at 37 °C. Wild-type *Streptomyces* strains and its mutants were cultivated on Mannitol Soya Flour (MS) agar medium (20.0 g mannitol, 20.0 g soya flour, 20.0 g agar, 1.0 L tap water, pH=7.0-7.5) at 30 □. The small-scale fermentation of *Streptomyces* strains was using 50 mL MS liquid medium in 250 mL flask, shaking at 200 rpm, 30 □ for 7 days. The large-scale fermentation of *Streptomyces* strains were carried out in medium 2 (CaCl$_2$·2H$_2$O, 3.0 g; citric acid/Fe III, 1.0 g;

MnSO$_4$·H$_2$O, 0.2 g; ZnCl$_2$, 0.1 g; CuSO$_4$·5H$_2$O, 0.025 g; Na$_2$B$_4$O$_7$·10H$_2$O, 0.02 g; Na$_2$MoO$_4$·2H$_2$O, 0.01 g; and oatmeal, 20.0 g, in 1.0 L distilled water), at 200 L scale in a 300 L fermentation vessel, for 6 days with aeration of 25-50 L min^{-1}, stirring at 200 rpm, 28 °C, with a pH range of 5.4-6.4. Antibiotics such as apramycin (50 mg mL^{-1}), kanamycin (50 mg mL^{-1}) or chloramphenicol (25 mg mL^{-1}) were appropriately used for resistance selection.

Genomic DNA extraction, sequencing, and assembly. The *S. iranensis* culture was growing in 50 mL sterilized liquid ISP2 medium (yeast extract 4.0 g; malt extract 10.0 g; and dextrose 4.0 g in 1.0 L distilled water, pH = 7.2) in 250 mL flask at 30 °C and 160 rpm for 5 days to generate sufficient biomass. The genomic DNA of *S. iranensis* was isolated using the QIAGEN Genomic-tip 20/G kit using a modified protocol and grinding with a mortar and pestle submerged in liquid nitrogen.[71] The procedure of genomic DNA sequencing was combining Oxford Nanopore Technologies MinION and Illumina MiSeq systems. To generate the nanopore data, both the SQK-RBK004 and the SKQ-RBK110-96 rapid barcoding kits were used for construction of three libraries, which were sequenced on two separate 9.4.1 flowcells. The *de novo* assembly was performed using Flye (v2.9-b1768) with the parameters --nano-hq --t 12. Since the genome could not be fully resolved using existing assemblers, the Terminal inverted repeat edge from the unambiguous assembly repeat graph were manually attached to both ends of the chromosome edge and the orientations were verified by mapping of the reads on the new assembly and manual inspection. This approach was later suggested by the Flye assembler Misha. The Unicycler (v0.4.8) polishing module was used to polish the nanopore assembly with illumina data.

Genetic manipulation. To verify the BGC of spirolactone and its individual biosynthetic genes, the classical CRISPR-Cas9 method, and advanced CRISPR-cBEST base editing toolkit were used to construct gene-inactive mutants.[36] The function-specific plasmids (Supplementary information Tab. 7) were constructed according to the respective protocol followed by introducing into wild-type *S. iranensis* HM 35 by conjugation with donor strain ET12567/pUZ8002 on SFM solid medium (soya flour 20.0 g; mannitol 20.0 g; and bacteria agar 20.0 g, in 1.0 L distilled water, pH = 7.2) according to modified protocol. The

exconjugants after resistance screening (50 µg mL^{-1} apramycin and 25 µg mL^{-1} nalidixic acid) were further verified by DNA extraction, PCR reaction, and Sanger sequencing.

DNA polymerases (Q5$^®$ High-Fidelity 2X Master Mix with Standard Buffer and OneTaq$^®$ 2X Master Mix with Standard Buffer) and restriction enzymes (NcoI, EcoRI) were purchased from New England Biolabs. PCR amplifications and restriction enzyme digestions were carried out on Bio-Rad's thermal cyclers according to the manufacturer's instructions. Plasmid DNA extraction was performed using NucleoSpin Plasmid EasyPure Kit (Macherey-Nagel, Germany). DNA purification was conducted on 1% Tris-acetate-EDTA (TAE) agarose gel followed by using NucleoSpin Gel and PCR Clean-up Kits (Macherey-Nagel, Germany). One Shot™ Mach1™ T1 Phage-Resistant Chemically Competent *E. coli* from Invitrogen™ was used for transformation. All oligonucleotides were ordered from Integrated DNA Technologies (Supplementary information Tab. 8) and Sanger sequencing was offered by Eurofins Genomics (Luxembourg). All solvents used for chromatography and HR-MS were purchased from VWR Chemicals with LC-MS grade, while for metabolites extraction, the solvents were of HPLC grade.

HR-ESI-MS analysis. HR-ESI-MS analysis was carried out via ultra-high-performance liquid chromatography-diode array detection–quadrupole time-of-flight mass spectrometry (UHPLC-DAD-QTOFMS) depending on an Agilent Infinity 1290 UHPLC system equipped with a diode array detector. Separation was achieved on a 250 × 2.1 mm i.d., 2.7 µm, Poroshell 120 Phenyl Hexyl column (Agilent Technologies) held at 60 °C. The sample, 1 µL, was eluted at a flow rate of 0.35 mL min^{-1} using a linear gradient from 10% acetonitrile in Milli-Q water buffered with 20 mM formic acid increasing to 100% in 15 min, staying there for 2 min before returning to 10% in 0.1 min. Starting conditions were held for 3 min before the following run. MS detection was performed on an Agilent 6545 QTOF MS equipped with Agilent Dual Jet Stream ESI with a drying gas temperature of 160 °C, a gas flow of 13 L min^{-1}, sheath gas temperature of 300 °C and flow of 16 L min^{-1}. Capillary voltage was set to 4000 V and nozzle voltage to 500 V in positive mode. All data were processed using Agilent MassHunter Qualitative Analysis software (Agilent Technologies, USA).

Large-scale fermentation and isolation. The culture broth from large-scale fermentation (described above) was filtered and loaded onto an Amberchrom 161c resin LC column (200 × 20 cm, 6 L). Elution with a linear gradient of H_2O-MeOH (from 30% to 100% v/v, flow rate 0.5 L min^{-1}, in 58 min) afforded seven fractions (Fr.A-Fr.G). Fr.G was first fractionated by silica gel chromatography with a CH_2Cl_2-CH_3OH gradient to yield 16 fractions, Fr.1-Fr.16. Fr.7 was further separated by a Sephadex LH-20 (MeOH) column, and twelve sub-fractions were obtained. The sub-fractions were separated by semipreparative HPLC RP-C18 (MeCN-H_2O as gradient) to afford compound **1**.

Spirolactone (**1**): white solid; $[\alpha]_D^{25}$ 15 (0.1 mg/mL, CH_3OH), UV/vis (CH_3CN/H_2O) λ_{max} 230, 270 nm; ECD λ_{ext} ($\Delta\varepsilon$) (CH_3OH) 235 (–9.78), 267 (7.02), 301 (–3.31) nm; IR (ATR) v_{max} 2932, 2748, 2704, 1810, 1713, 1686, 1619, 1457, 1382, 1173, 1134, 1085, 1057, 1013, 991, 947 cm^{-1}; (-)-HR-ESI-MS m/z 881.4659 [M + Na]$^+$ (calcd for $C_{47}H_{70}O_{14}$, 881.4658).

NMR spectroscopy. NMR spectra were recorded on 800 MHz Bruker Avance III spectrometer equipped with a TCI CryoProbe using standard pulse sequences. Measurements were carried out using ^1H and ^{13}C NMR, ^1H-^{13}C heteronuclear single quantum coherence (HSQC), ^1H-^{13}C heteronuclear multiple bond correlation (HMBC), ^1H-^1H correlation spectroscopy (COSY), ^1H-^1H nuclear overhauser effect spectroscopy (NOESY). Chemical shifts (δ) were reported in parts per million (ppm) and referenced to the solvent signals. Data are described as follows: chemical shift, multiplicity (br = broad, s = singlet, d = doublet, t = triplet, dd = doublet of doublet, m = multiplet and ov = overlapped) and coupling constants (in hertz). All NMR data were processed using MestReNova 14.0.

Crystal Data for spirolactone. The suitable crystal was selected and mounted in a nylon loop directly from the ethanol suspension and frozen in liquid nitrogen on a Synchrotron diffractometer. X-ray data collection of **1** was performed on an Agilent Supernova Diffractometer using CuKα radiation, and the crystal was kept at 100 K during data collection. Using Olex2,[72] the structure was solved with the XT[73] structure solution program using Intrinsic Phasing and refined with the SHELXL[74] refinement package using Least Squares minimization. Crystal data for $C_{47}H_{70}O_{14}$ (M =859.03 g/mol): monoclinic, space

group P2$_1$ (no. 4), a = 10.118(7) Å, b = 16.799(7) Å, c = 14.081(5) Å, β = 103.385(12)°, V = 2328(2) Å^3, Z = 2, T = 100 K, μ (synchrotron) = 0.089 mm^{-1}, $Dcalc$ = 1.225 g/cm^3, 17409 reflections measured (3.836° $\leq$ 2θ $\leq$ 45.2°), 5762 unique (R_{int} = 0.0646, R_{sigma} = 0.0641) which were used in all calculations. The final R_1 was 0.0646 (I > 2σ(I)) and wR_2 was 0.1523 (all data).

Sample preparation for proteomics analysis. Approximately 1.0 x 10^7 conidia mL^{-1} of *A. niger* ATCC 1015 strain were used to inoculate 100 mL (three biological replicates) of liquid cultures (PDB) in 1 L flask that were incubated in a reciprocal shaker at 28 ☐ in 180 rpm for 16 hours. Afterwards, samples in the treatment group were treated with 0.2 µg (concentration of 2 ng/mL) of spirolactone for 1-hour and 4-hour. Mycelia were then harvested by filtering, washed thoroughly with sterile water and quickly frozen in liquid nitrogen. Pellets were lysed in 100 µL lysis buffer (6 M Guanidium Hydrochloride, 10 mM TCEP, 40 mM CAA, 50 mM HEPES, pH 8.5) by boiling the samples at 95 ☐ for 5 mins, sonicating on high for 5 x 60 sec on/30sec off using the bioruptor pico sonication water bath (Diagenode) and finally disrupting twice with the tissuelyser (QIAGEN) going from 3 to 30 Hz in 1 min. Lysed samples were centrifuged at 18,000 g for 10 mins, and supernatants were transferred to clean LoBind Eppendorf tubes. Protein concentration was determined by BCA rapid gold (Thermo) and 10 µg of protein was taken forward for digestion. Samples were diluted 1:3 with digestion buffer (10% acetonitrile in 50mM HEPES pH 8.5) and incubated with 1:100 enzyme to protein ratio of LysC (MS Grade, Wako) at 37 ☐ for 4 hours. Samples were further diluted to a final 1:10 with more digestion buffer and digested with 1:100 trypsin for 18 hours at 37 ☐. After digestion, samples were acidified with TFA and desalted using the SOLAµ^{TM} SPE plate (HRP, Thermo).[75] Between each application, the solvents were spun through by centrifugation at 1,500 rpm. For each sample, the filters were activated with 200 µL of 100% Methanol, then 200 µL of 80% Acetonitrile, 0.1% formic acid. The filters were subsequently equilibrated 2x with 200 µL of 1% TFA, 3% acetonitrile, after which the sample was loaded. After washing the tips twice with 200 µL of 0.1% formic acid, the peptides were eluted into clean 0.5 mL Eppendorf tubes using 40% acetonitrile, 0.1% formic

acid. The eluted peptides were concentrated in an Eppendorf Speedvac. Samples were reconstituted in 12 µL A* buffer with iRT peptides (Biognosys).

MS analysis for proteomics analysis. Peptides were loaded onto a 2cm C18 trap column (ThermoFisher 164946), connected in-line to a 15cm C18 reverse-phase analytical column (Thermo EasySpray ES904) using 100% Buffer A (0.1% Formic acid in water) at 750 bar, using the Thermo EasyLC 1200 HPLC system, and the column oven operating at 30°C. Peptides were eluted over a 70 minutes gradient ranging from 10% to 60% of Buffer B (80% acetonitrile, 0.1% formic acid) at 250 µL/min, and the Orbitrap Exploris instrument (Thermo Fisher Scientific) was run in DIA mode with FAIMS ProTM Interface (ThermoFisher Scientific) with CV of -45 V. Full MS spectra were collected at a resolution of 120,000, with an AGC target of 300% or maximum injection time set to 'auto' and a scan range of 400–1000 m/z. The MS2 spectra were obtained in DIA mode in the orbitrap operating at a resolution of 60.000, with an AGC target 1000% or maximum injection time set to 'auto', a normalised HCD collision energy of 32. The isolation window was set to 6 m/z with a 1 m/z overlap and window placement on. Each DIA experiment covered a range of 200 m/z resulting in three DIA experiments (400-600 m/z, 600-800 m/z and 800-1000 *m/z*). Between the DIA experiments a full MS scan is performed. MS performance was verified for consistency by running complex cell lysate quality control standards, and chromatography was monitored to check for reproducibility.

Data process for proteomics analysis. The raw files were analyzed using Spectronaut[TM] (version 17.4) spectra were matched against the reviewed *A. niger* ATCC 1015 NCBI database. Dynamic modifications were set as Oxidation (M) and Acetyl on protein N-termini. Cysteine carbamidomethyl was set as a static modification. All results were filtered to a 1% FDR, and protein quantitation done on the MS1 level. Only proteotypic peptides were used for quantification and protein groups were inferred by IDPicker.

Antifungal susceptibility testing. In vitro antifungal susceptibility of spirolactone against 34 fungal species represented by 38 clinical isolates was tested in comparison to approved

antifungals amphotericin B (AMB; European Pharmacopoeia, Strasbourg, France); itraconazole (ITZ), and voriconazole (VCZ; Pfizer Inc., Peapack, NJ, USA) by using the broth microdilution technique following the European Committee on Antimicrobial Susceptibility Testing (EUCAST) standard methodology for yeasts or filamentous fungi respectively, in the main parts. In contrast to the EUCAST protocol, microdilution plates were prepared by 2-fold serial dilutions of the antifungal agents. Filamentous fungi were grown on malt extract agar (MEA) for 2 to 7 days at 35 °C and yeasts were cultivated on yeast extract peptone dextrose agar (YPD) for 24 hours. Spore or yeast cell suspensions were counted with a hemocytometer. Minimum inhibitory concentrations (MIC) endpoints of filamentous fungi were defined as 100% reduction in growth and were determined visually using a mirror after 48 h of incubation at 35 °C. Microdilution plates of yeasts were read with a microdilution plate reader (Infinite® M Nano plus, Tecan) and MIC endpoints were defined as the lowest drug concentration giving inhibition of growth of $\geq$ 50% of that of the drug-free control. Since the mode of action of spirolactone is unknown, the minimum effective concentration (MEC) was determined as well by reading the microplates with the aid of an inverted microscope (Eclipse Ts2, Nikon). *A. fumigatus* ATCC 204305 and *Candida parapsilosis* ATCC 22019 were used as reference strains.[76,77]

Cytotoxic activities of spirolactone. Cytotoxicity against the human cell line HL-60 was evaluated using Alamar Blue (Thermos Scientific, Kansas, USA). The assay (Hamid et al. 2004) was performed in 96 well plates (Costar 3595, Corning, New York, USA), with an assay volume of 200 µL. The software Prism 5.03 was used for data analysis (GraphPad Software, USA).[78]

Data availability. The genome sequence of *S. iranensis* HM35 has been deposited in the NCBI under BioProject accession number PRJNA1026072 and BioSample accession number SAMN37733152. The genome sequence can be found under Genbank accession number CP136563. The crystal data of spirolactone was deposited in CCDC database (https://www.ccdc.cam.ac.uk/) with accession number 2178562. The metabolomics data of wild-type and mutant strains has been deposited in the MassIVE repository

(https://massive.ucsd.edu/) under identifier MSV 000094394. The mass spectrometry for proteomics analysis have been deposited in the ProteomeXchange Consortium (https://www.proteomexchange.org/) with data set identifier MassIVE MSV 000094235. The NMR data were deposited at NP-MRD, NP0332733.

Acknowledgements

L.D. acknowledges funding from DTU Skylab, Novo Nordisk Foundation (Grant NNF23OC0082881) and Innovation Fund Denmark (Grant 3141-00013A). M.H.C. acknowledges funding for the DTU Screening Core from the Novo Nordisk Foundation (Grant NNF19OC0055818) and the Carlsberg Foundation (Grant CF19-0072). Y.Q. acknowledges funding from Novo Nordisk Foundation (NNF22OC0079928). The NMR Center DTU and the Villum Foundation are acknowledged for access to the 800 MHz spectrometer. The metabolomic data was generated at DTU Metabolomics Core facilities with help of Dr. A. Andersen. Thanks to Prof. Mohamed Shaaban for repurification of spirolactone and recording the ECD spectrum. Thanks to Prof. Jacob Blæsbjerg Hoof for the *A. niger* strain and advice.

Competing interests

The authors declare that there are no competing interests. Two patent applications were filed with the numbers of PCT/EP2023/067953 and EP24150176.6.

Additional Information

Supplementary information: The online version contains supplementary material available at xxx.

Correspondence and requests for materials should be addressed to Ling Ding.

References

1. Brown, G. D., et al. Hidden killers: human fungal infections. *Sci. Transl. Med.* **4**, 165rv13 (2012).

2. Bongomin, F., Gago, S., Oladele, R. O. & Denning, D. W. Global and multi-national prevalence of fungal diseases-estimate precision. *J. Fungi.* **3**, 57 (2017).

3. Denning, D. W. Global incidence and mortality of severe fungal disease. *Lancet Infect. Dis.* S1473-3099, 00692-8 (2024).

4. Sarden, N., et al. A B1a-natural IgG-neutrophil axis is impaired in viral- and steroid-associated aspergillosis. *Sci. Transl. Med.* **14**, eabq6682 (2022).

5. Osherov, N. & Kontoyiannis, D. P. The anti-*Aspergillus* drug pipeline: Is the glass half full or empty? *Med. Mycol.* **55**, 118-124 (2017).

6. Hoenigl, M., et al. The antifungal pipeline: fosmanogepix, ibrexafungerp, olorofim, opelconazole, and rezafungin. *Drugs* **81**, 1703-1729 (2021).

7. Perfect, J. R. The antifungal pipeline: a reality check. *Nat. Rev. Drug Discovery* **16**, 603-616 (2017).

8. Denning, D. W. & Bromley, M. J. Infectious Disease. How to bolster the antifungal pipeline. *Science* **347**, 1414-1416 (2015).

9. Gubbins, P. O. & Amsden, J. R. Drug-drug interactions of antifungal agents and implications for patient care. *Expert Opin. Pharmacother.* **6**, 2231-2243 (2005).

10. Rhodes, J., et al. Population genomics confirms acquisition of drug-resistant *Aspergillus fumigatus* infection by humans from the environment. *Nat. Microbiol.* **7**, 663-674 (2022).

11. Vermeulen, E., Lagrou, K. & Verweij, P. E. Azole resistance in *Aspergillus fumigatus*: a growing public health concern. *Curr. Opin. Infect. Dis.* **26**, 493-500 (2013).

12. Zhou, D., et al. Extensive genetic diversity and widespread azole resistance in greenhouse populations of *Aspergillus fumigatus* in Yunnan, China. *mSphere* **6**, e00066-21 (2021).

13. Newman, D. J. & Cragg, G. M. Natural products as sources of new drugs over the nearly four decades from 01/1981 to 09/2019. *J. Nat. Prod.* **83**, 770-803 (2020).

14. Jiang, T., Pu, H., Duan, Y., Yan, X. & Huang, Y. New natural products of *Streptomyces* sourced from deep-sea, desert, volcanic, and polar regions from 2009 to 2020. *Chin. J. Org. Chem.* **41**, 1804-1820 (2021).

15. Alam, K., et al. *Streptomyces*: The biofactory of secondary metabolites. *Front. Microbiol.* **13**, 968053 (2022).

16. Hazen, E. L. & Brown, R. Fungicidin, an antibiotic produced by a soil Actinomycete. *Proc. Soc. Exp. Biol. Med.* **76**, 93-97 (1951).

17. Oura, M., Sternberg, T. H. & Wright, E. T. A new antifungal antibiotic, amphotericin B. *Antibiot. Annu.* **3**, 566-573 (1955).

18. Anderson, T. M., et al. Amphotericin forms an extramembranous and fungicidal sterol sponge. *Nat. Chem. Biol.* **10**, 400-406 (2014).

19. Lewandowska, A., et al. Fungicidal amphotericin B sponges are assemblies of staggered asymmetric homodimers encasing large void volumes. *Nat. Struct. Mol. Biol.* **28**, 972-981 (2021).

20. Gray, K. C., et al. Amphotericin primarily kills yeast by simply binding ergosterol. *Proc. Natl. Acad. Sci. USA* **109**, 2234-2239 (2012).

21. Hamill, R. J. Amphotericin B formulations: a comparative review of efficacy and toxicity. *Drugs* **73**, 919-934 (2013).

22. Fanos, V. & Cataldi, L. Amphotericin B-induced nephrotoxicity: A review. *J. Chemother.* **12**, 463-470 (2000).

23. Chen, R., et al. Turonicin A, an antifungal linear polyene polyketide from an Australian *Streptomyces* sp. *J. Nat. Prod.* **86**, 2054-2058 (2023).

24. Tchize Ndejouong, B., et al. Hygrobafilomycin, a cytotoxic and antifungal macrolide bearing a unique monoalkylmaleic anhydride moiety, from *Streptomyces varsoviensis*. *J. Antibiot.* **63**, 359-363 (2010).

25. Zhang, Z. W., Zhou, T., Harunari, E., Oku, N. & Igarashi, Y. Iseolides A-C, antifungal macrolides from a coral-derived actinomycete of the genus *Streptomyces*. *J. Antibiot.* **73**, 534-541 (2020).

26. Chevrette, M. G., et al. The antimicrobial potential of *Streptomyces* from insect microbiomes. *Nat. Commun.* **10**, 516 (2019).

27. Arai, M. Isolation of three main components. F3, F4 and F5, from azalomycin F-complex. *J. Antibiot.* **23**, 107-112 (1970).

28. Chen, Y., et al. Discovery of niphimycin C from *Streptomyces yongxingensis* sp. nov. as a promising agrochemical fungicide for controlling banana *Fusarium* Wilt by destroying the mitochondrial structure and function. *J. Agric. Food Chem.* **70**, 12874-12795 (2022).

29. Zhang, Y., et al. Antifungal activities of metabolites produced by a termite-associated *Streptomyces canus* BYB02. *J. Agric. Food Chem.* **61**, 1521-1524 (2013).

30. Nguyen, C. T., Dhakal, D., Pham, V. T. T., Nguyen, H. T. & Sohng, J. K. Recent advances in strategies for activation and discovery/characterization of cryptic biosynthetic gene clusters in *Streptomyces*. *Microorganisms* **8**, 616 (2020).

31. Ziemert, N., Alanjary, M. & Weber, T. The evolution of genome mining in microbes - a review. *Nat. Prod. Rep.* **33**, 988-1005 (2016).

32. Hamedi, J., et al. *Streptomyces iranensis* sp nov., isolated from soil. *Int. J. Syst. Evol. Microbiol.* **60**, 1504-1509 (2010).

33. Yang, Z., et al. *Streptomyces* alleviate abiotic stress in plant by producing pteridic acids. *Nat. Commun.* **14**, 7398 (2023).

34. Blin, K., et al. antiSMASH 7.0: new and improved predictions for detection, regulation, chemical structures and visualisation. *Nucleic Acids Res.* **51**, W46-W50 (2023).

35. Kautsar, S. A., et al. BiG-FAM: the biosynthetic gene cluster families database. *Nucleic Acids Res.* **49**, D490-D497 (2021).

36. Tong, Y., et al. CRISPR-Cas9, CRISPRi and CRISPR-BEST-mediated genetic manipulation in streptomycetes. *Nat. Protoc.* **15**, 2470-2502 (2020).

37. Keatinge-Clay, A. T. The structures of type I polyketide synthases. *Nat. Prod. Rep.* **29**, 1050-1073 (2012).

38. Wang, J., et al. Platensimycin is a selective FabF inhibitor with potent antibiotic properties. *Nature* **441**, 358-361 (2006).

39. Keatinge-Clay, A. T. A tylosin ketoreductase reveals how chirality is determined in polyketides. *Chem. Biol.* **14**, 898-908 (2007).

40. Gehret, J. J., et al. Terminal alkene formation by the thioesterase of curacin a biosynthesis structure of a decarboxylating thioesterase. *J. Biol. Chem.* **286**, 14445-14454 (2011).

41. Rudolf, J. D., Chang, C. Y., Ma, M. & Shen, B. Cytochromes P450 for natural product biosynthesis in *Streptomyces*: sequence, structure, and function. *Nat. Prod. Rep.* **34**, 1141-1172 (2017).

42. Moebius, N., et al. Biosynthesis of the respiratory toxin bongkrekic acid in the pathogenic bacterium *Burkholderia gladioli*. *Chem. Biol.* **19**, 1164-1174 (2012).

43. Zhang, Q., et al. Carboxyl formation from methyl via triple hydroxylations by XiaM in xiamycin A biosynthesis. *Org. Lett.* **14**, 6142-6145 (2012).

44. Aparicio, J. F., Caffrey, P., Gil, J. A. & Zotchev, S. B. Polyene antibiotic biosynthesis gene clusters. *Appl. Microbiol. Biotechnol.* **61**, 179-188 (2003).

45. Chen, S., et al. Tailoring the P450 monooxygenase gene for FR-008/Candicidin biosynthesis. *Appl. Environ. Microbiol.* **75**, 1778-1781 (2009).

46. Yang, J., Qi, Y., Blodgett, J. & Wencewicz, T. Multifunctional P450 monooxygenase CftA diversifies the clifednamide pool through tandem C-H bond activations. *J. Nat. Prod.* **85**, 47-55 (2022).

47. Robinson, S. L., Christenson, J. K. & Wackett, L. P. Biosynthesis and chemical diversity of β-lactone natural products. *Nat. Prod. Rep.* **36**, 458-475 (2019).

48. Feng, K. N., et al. A hydrolase-catalyzed cyclization forms the fused bicyclic β-Lactone in vibralactone. *Angew. Chem. Int. Ed.* **59**, 7209-7213 (2020).

49. Bauman, K. D., et al. Enzymatic assembly of the salinosporamide γ-lactam-β-lactone anticancer warhead. *Nat. Chem. Biol.* **18**, 538-546 (2022).

50. Schaffer, J. E., Reck, M. R., Prasad, N. K. & Wencewicz, T. A. β-Lactone formation during product release from a nonribosomal peptide synthetase. *Nat. Chem. Biol.* **13**, 737-744 (2017).

51. Xu, Z. F., et al. Discovery and biosynthetic pathway analysis of cyclopentane-β-lactone globilactone A. *Nat. Synth.* **3**, 99-110 (2024).

52. Christenson, J. K., et al. β-Lactone synthetase found in the olefin biosynthesis pathway. *Biochemistry* **56**, 348-351 (2017).

53. Robinson, S. L., et al. Mechanism of a standalone β-lactone synthetase: new continuous assay for a widespread ANL superfamily enzyme. *ChemBioChem* **20**, 1701-1711 (2019).

54. Chan, Y. A., Podevels, A. M., Kevany, B. M. & Thomas, M. G. Biosynthesis of polyketide synthase extender units. *Nat. Prod. Rep.* **26**, 90-114 (2009).

55. Wilson, M. C. & Moore, B. S. Beyond ethylmalonyl-CoA: The functional role of crotonyl-CoA carboxylase/reductase homologs in expanding polyketide diversity. *Nat. Prod. Rep.* **29**, 72-86 (2012).

56. Quade, N., Huo, L. J., Rachid, S., Heinz, D. W. & Muller, R. Unusual carbon fixation gives rise to diverse polyketide extender units. *Nat. Chem. Biol.* **8**, 117-124 (2012).

57. Omura, S., Imai, H., Takeshima, H., & Nakagawa, A. Structure of a new antimicrobial unsaturated fatty acid from *Sm. kitasatoenis* NU-23-1. *Chem. Pharm. Bull.* 24(12), 3139-3143 (1976).

58. Ikeda, H. & Omura, S. Avermectin biosynthesis. *Chem. Rev.* **97**, 2591-2610 (1997).

59. Cantalapiedra, C. P., Hernández-Plaza, A., Letunic, I., Bork, P. & Huerta-Cepas, J. eggNOG-mapper v2: Functional annotation, orthology assignments, and domain prediction at the metagenomic scale. *Mol. Biol. Evol.* **38**, 5825-5829 (2021).

60. Gautam, P., et al. Proteomic and transcriptomic analysis of *Aspergillus fumigatus* on exposure to amphotericin B. *Antimicrob. Agents Chemother.* **52**, 4220-4227 (2008).

61. Ghannoum, M., et al. *In vitro* antifungal activity of naftifine hydrochloride against dermatophytes. *Antimicrob. Agents Chemother.* **57**, 4369-4372 (2013).

62. Hitchcock, C. A. Cytochrome-P-450-dependent 14α-sterol demethylase of *Candida albicans* and its interaction with azole antifungals. *Biochem. Soc. Trans.* **19**, 782-787 (1991).

63. Sawistowska-Schröder, E. T., Kerridge, D. & Perry, H. Echinocandin inhibition of 1,3-β-D-glucan synthase from *Candida albicans*. *FEBS Lett.* **173**, 134-138 (1984).

64. Cagas, S. E., Jain, M. R., Li, H. & Perlin, D. S. Profiling the *Aspergillus fumigatus* proteome in response to caspofungin. *Antimicrob. Agents Chemother.* **55**, 146-154 (2011).

65. Walker, L. A., Gow, N. A. R. & Munro, C. A. Elevated chitin content reduces the susceptibility of *Candida* species to caspofungin. *Antimicrob. Agents Chemother.* **57**, 146-154 (2013).

66. Garcia-Rubio, R., de Oliveira, H. C., Rivera, J. & Trevijano-Contador, N. The fungal cell wall: *Candida*, *Cryptococcus*, and *Aspergillus* species. *Front. Microbiol.* **10**, 2993 (2020).

67. Fang, W., et al. Mechanisms of redundancy and specificity of the *Aspergillus fumigatus* Crh transglycosylases. *Nat. Commun.* **10**, 1669 (2019).

68. Puttikamonkul, S., et al. Trehalose 6-phosphate phosphatase is required for cell wall integrity and fungal virulence but not trehalose biosynthesis in the human fungal pathogen *Aspergillus fumigatus*. *Mol. Microbiol.* **77**, 891-911 (2010).

69. Dhingra, S. & Cramer, R. A. Regulation of sterol biosynthesis in the human fungal pathogen *Aspergillus fumigatus*: opportunities for therapeutic development. *Front. Microbiol.* **8**, 92 (2017).

70. Fraczek, M. G., et al. The *cdr1B* efflux transporter is associated with non-*cyp51a*-mediated itraconazole resistance in *Aspergillus fumigatus*. *J. Antimicrob. Chemother.* **68**, 1486-1496 (2013).

71. Alvarez-Arevalo, M., et al. Extraction and Oxford Nanopore sequencing of genomic DNA from filamentous Actinobacteria. *STAR protoc.* **4**, 101955 (2023).

72. Dolomanov, O. V., et al. *OLEX2*: a complete structure solution, refinement and analysis program. *J. Appl. Crystallogr.* **42**, 339-341 (2009).

73. Sheldrick, G. M. Crystal structure refinement with SHELXL. *Acta Crystallogr C Struct. Chem.* **71**, 3-8 (2015).

74. Sheldrick, G. M. SHELXT - integrated space-group and crystal-structure determination. *Acta Crystallogr. A Found. Adv.* **71**, 3-8 (2015).

75. Rappsilber, J., Mann, M. & Ishihama, Y. Protocol for micro-purification, enrichment, pre-fractionation and storage of peptides for proteomics using StageTips. *Nat. Protoc.* **2**, 1896-1906 (2007).

76. Arendrup, M. C., Cuenca-Estrella, M., Lass-Flörl, C., Hope, W. & EUCAST-AFST. EUCAST technical note on the EUCAST definitive document EDef 7.2: Method for the determination of broth dilution minimum inhibitory concentrations of antifungal agents for yeasts EDef 7.2 (EUCAST-AFST). *Clin. Microbiol. Infect.* **18**, E246-247 (2012).

77. Subcommittee on Antifungal Susceptibility Testing of the ESCMID European Committee for Antimicrobial Susceptibility Testing. EUCAST Technical Note on the method for the determination of broth dilution minimum inhibitory concentrations of antifungal agents for conidia forming moulds. *Clin. Microbiol. Infect.* **14**, 982-984 (2008).

78. Lima, T. S., et al. Molecular profiling of docetaxel-resistant prostate cancer cells identifies multiple mechanisms of therapeutic resistance. *Cancers* **13**, 1290 (2021).