Transcriptome-wide identification of 5-methylcytosine by deaminase and reader protein-assisted sequencing

Jiale Zhou[1][#], Ding Zhao[1][#], Jinze Li[2][#], Deqiang Kong[1], Xiangrui Li[2], Renquan Zhang[2],

Yuru Liang[2], Xun Gao[2], Yuqiang Qian[2], Di Wang[2], Jiahui Chen[2], Liangxue Lai [1,3,4*],

Yang Han[2*], Zhanjun Li [2*]

# These authors contributed equally: Jiale Zhou, Ding Zhao, Jinze Li.

[1]Key Laboratory of Zoonosis Research, Ministry of Education, College of Veterinary

Medicine, Jilin University, Changchun, 130062, China

[2]Laboratory Animal Center, College of Animal Science, Jilin University, Changchun,

130062, China

[3]Guangzhou Regenerative Medicine and Health Guang Dong Laboratory

(GRMH-GDL), Guangzhou, 510530, China

[4]Institute for Stem Cell and Regeneration, Chinese Academy of Sciences, Beijing,

510005, China

* To whom correspondence should be addressed:

Tel: (86)431-87836176; Fax: (86)431-87980131.

*e-mail:

hanyang8584@jlu.edu.cn (Yang Han); lizj_1998@jlu.edu.cn (Zhanjun Li);

lai_liangxue@gibh.ac.cn (Liangxue lai).

Abstract

5-Methylcytosine (m5C) is one of the major post-transcriptional modifications in mRNA and is highly involved in the pathogenesis of various diseases. However, the capacity of existing assays for accurately and comprehensively transcriptome-wide m5C mapping still needs improvement. Here, we develop a detection method named DRAM (deaminase and reader protein assisted RNA methylation analysis), in which deaminases (APOBEC1 and TadA-8e) are fused with m5C reader proteins (ALYREF and YBX1) to identify the m5C sites through deamination events neighboring the methylation sites. This antibody-free and bisulfite-free approach provides transcriptome-wide editing regions which are highly overlapped with the publicly available BS-seq datasets and allows for a more stable and comprehensive identification of the m5C loci. In addition, DRAM system even supports ultra-low input RNA (10ng) and monitors the dynamic accumulation of cellular m5C. We anticipate that the DRAM system could pave the way for uncovering further biological functions of m5C modifications.

Keywords: RNA, 5-methylcytosine, deaminase, m5C reader

Introduction

Epigenetics refers to stable inheritance without changing the basic sequence of DNA, involving various forms such as DNA methylation, histone modification and RNA modification. In recent years, RNA sequencing technology has boosted research on RNA epigenetics. More than 170 RNA modifications have been identified, mainly including m^6A, m^5C, m^1A, m^7G and others [1,2]. Notably, RNA m^5C methylation represents a crucial post-transcriptional modification observed across different RNA types, such as tRNA, mRNA, rRNA, vault RNA, microRNA, long non-coding RNA and enhancer RNA[3-8]. Numerous studies have revealed multiple molecular functions of m^5C in numerous key stages of RNA metabolism, such as mRNA stability, translation, and nuclear export[5,9-13]. The dynamic alterations of m^5C play integral roles in many physiological and pathological processes, such as early embryonic development[14], neurodevelopmental disorder[15,16] and multifarious tumorigenesis and migration[17-20]. Moreover, this modification significantly contributes to the regulation of gene expression[5,9-13,17]. Therefore, the detection of m^5C sites appears to be essential for understanding their underlying effects on cellular function and disease states.

With the recent advances in sequencing techniques, several high-throughput assays have been developed for qualitative or quantitative analysis of m^5C. To date, bisulfite-sequencing (BS-seq) has been proven to be the gold standard method for RNA m^5C methylation analysis[5,21,22]. This approach chemically deaminates unmethylated cytosine to uracil, while keeping methylated cytosine unchanged. The m^5C methylation sites can be identified by subsequent library construction and sequencing. However, bisulfite treatment of BS-seq is extremely detrimental to RNA, thus resulting in unstable detection of m^5C in low abundance RNA or highly structured RNA, which directly affects the confidence of results[23,24]. Another major type of global m^5C analysis depends on antibody-assisted immunoprecipitation of m^5C methylated RNAs, such as m^5C-RIP-seq[25-27], AZA-IP-seq[28] or miCLIP-seq[7]. These methods are unable to recognize methylation on mRNAs with low abundance and secondary structure. Moreover, these methods are highly dependent on antibody specificity, which usually leads to unspecific binding of RNA and a low amount of m^5C-modified regions. Moreover, TAWO-seq, originally developed for the identification of hm^5C, is also capable of m^5C analysis, but it highly depends on the oxidation efficiency of perovskite, which usually causes false positives and unstable conversion [29,30]. Furthermore, the emerging third-generation sequencing, such as Nanopore-seq, can directly map m^5C by tracking the characteristic changes of bases, but it still faces challenges of a high error rate[31-33]. These together largely hamper its wide application on transcriptome profiling of m^5C (Supplementary Table 1). Hence, there is an urgent need for a simple, efficient, sensitive, and antibody-independent method for global m^5C detection.

The RNA-binding protein ALYREF is the initially recognized nuclear m^5C reader that binds directly to m^5C sites in mRNA and plays key roles in promoting mRNA nuclear export or tumor progression[5]. Another well-known m^5C reader, YBX1, can also recognize m^5C-modified mRNA through its cold-shock domain and participates in a variety of RNA-dependent events such as mRNA packaging, mRNA stabilization and translational regulation[9,18]. RNA affinity chromatography and mass spectrometry analyses using biotin-labelled oligonucleotides with or without m^5C were performed in previous reports, which indicated that ALYREF and YBX1 had a more prominent binding ability to m^5C-modified oligonucleotides[5,18]. YBX1 can preferentially

recognize mRNAs with m[5]C modifications via key amino acids W65-N70 (WFNVRN)[18], while
K171 is essential for the specific binding of ALYREF to m[5]C sites [5]. Previous studies have shown
that mutations in key amino acids responsible for recognising m[5]C binding in ALYREF and YBX1
lead to a significant reduction in their binding levels to m[5]C-containing oligonucleotides[5,18].
Nucleic acid deaminases, primarily categorized as cytosine deaminases and adenine deaminases,
are zinc-dependent enzymes which facilitate the deamination of cytosine or adenine within DNA
or RNA substrates[34]. APOBEC1, an evolutionarily conserved family member of APOBEC
proteins, can specifically catalyze the deamination of cytosine in single-stranded RNA (ssRNA) or
DNA (ssDNA) to uracil[35-37]. TadA8e is an adenine deaminase optimized through re-engineering of
TadA and it induces conversion of adenine to inosine (eventually read as guanine by transcriptases)
in ssRNA or ssDNA[38,39]. APOBEC1 and TadA8e, with their prominent deamination efficiency,
have been employed for the development of precise and efficient base editors such as CBE and
ABE8e, which find widespread application in studies related to genome editing[37,38].

Here we aim to establish a deaminase and m[5]C reader-assisted RNA methylation sequencing
approach (DRAM-seq), which identifies the m[5]C sites through reader-mediated recognitions and
deaminase-mediated point mutations neighboring the m[5]C methylation sites. This bisulfite-free
and antibody-free method is anticipated to provide more comprehensive and cost-effective
transcriptome-wide detection of m[5]C methylation, which may better assist on exploring its further
regulatory mechanisms.

Results

Development of DRAM system for m[5]C detection

Our sequencing platform is inspired by the concept of the m[6]A DART-seq assay, in which C
near the m[6]A site is converted into U without affecting sequences near non-m6A sites[40]. Therefore,
we hypothesized that, by utilizing the targeted binding of m[5]C readers, deaminase can be recruited
to achieve deamination of cytosine or adenine in the vicinity of the m[5]C sites on single-stranded
RNA, thereby facilitating the detection of the m[5]C site. This approach was named DRAM
(deaminase and m[5]C reader-assisted RNA methylation sequencing). As RNA-binding proteins,
ALYREF and YBX1 also could bind to RNAs without m[5]C modification[5,18]. To exclude the
false-positive detection of DRAM due to the non-m[5]C specific binding of ALYREF and YBX1,
knockout of W65-N70 (WFNVRN) amino acids in YBX1and K171A mutation in ALYREF were
introduced seperately, resulting in the DRAM[mut] system (Fig. S1A-S1D). DRAM[mut] exhibits
substantially reduced binding to m[5]C in RNA pulldown assays (Fig. S1D-S1F). To confirm the
recognition of m[5]C site by DRAM system, DRAM, DRAM[mut] and Deaminase system were
transfected into the human HEK293T cells, respectively. Finally, we considered the presence of
m[5]C modification in the vicinity only if the deamination changes produced under DRAM
induction were significantly different from those produced under DRAM[mut] or Deaminase
induction (Fig. 1A).

Previous studies have indicated that there is no uniform intrinsic signature motif sequence
that can characterize all m[5]C sites[5,26,41,42]. To comprehensively detect the m[5]C loci, the readers of
m[5]C (ALYREF and YBX1) were separately fused to the C-terminus of the deaminases
(APOBEC1 and TadA-8e), namely DRAM-ABE and DRAM-CBE system (Fig.1B).

DRAM detection system is assayed in an m^{5}C-dependent form

To confirm the recognition of m^{5}C site by DRAM system, DRAM, DRAM^{mut} and Deaminase were transfected into the human HEK293T cells, respectively. To evaluate candidate DRAM constructs within a cellular environment, we performed fluorescence microscopy to analyze the expression of DRAM. The results showed that DRAM-ABE and DRAM-CBE were properly expressed in HEK293T cells (Fig. S2A-S2B). In addition, flow cytometry displayed ~60% of cells were GFP-positive (Fig. S2C). Two previously reported m^{5}C sites in RPSA and AP5Z1 were selected for the analysis [5,21], and their methylation status was verified by bisulfite sequencing PCR. The deep sequencing results showed that the m□C fraction of RPSA and SZRD1 was 75.5% and 27.25%, respectively (Fig.2A and B). Sanger sequencing following RT-PCR was then performed to determine the editing of neighbouring m^{5}C sites by DRAM system in these two mRNA. Notably, adenine close to the m^{5}C site in RPSA mRNA was mutated into guanine, resulting in an A-to-G editing rate of 14.7% by DRAM-ABE, whereas this was rarely observed with TadA-8e or DRAM^{mut}-ABE (Fig.2C). DRAM-CBE induced C to U editing in the vicinity of the m^{5}C site in AP5Z1 mRNA, with 13.6% C-to-U editing, while this effect was significantly reduced with APOBEC1 or DRAM^{mut}-CBE (Fig.2D). Subsequently, in order to investigate whether the DRAM system can detect other types of RNA, such as tRNA, 28S rRNA, or others, we performed PCR amplification of the flanking sequences of the m^{5}C sites 3782 and 4447 on 28S rRNA and several m^{5}C sites on tRNA, such as the m^{5}C48 and m^{5}C49 sites of tRNA^{Val}, the m^{5}C48 and m^{5}C49 sites of tRNA^{Asp}, and the m^{5}C48 site of tRNA^{Lys}. But Sanger sequencing showed that there was no valid A-to-G/C-to-U mutation detected, which is most likely due to the fact that ALYTEF and YBX1 are mainly responsible for the mRNA m^{5}C binding proteins, and thus the DRAM system is more suitable for the mRNA m^{5}C detection (Fig. S3). Taken together, the fusion of m^{5}C reader and deaminase can effectively and selectively deaminate cytosine/adenine in the vicinity of the mRNA m^{5}C sites.

NSUN2[43] and NSUN6[44], two family members of NOL1/NSUN protein, were both identified as m^{5}C methyltransferase of mRNA[45]. To verify that the detection of DRAM occurs in the presence of m^{5}C, we individually depleted NSUN2 and NSUN6 in HEK293T cells and performed DRAM transfection. The knockout efficiency has been confirmed by western blotting (Fig.2E, 2F and Fig. S4A,4B). It has been previously demonstrated that m^{5}C methylation of AP5Z1 and RPSA is catalyzed by NSUN2 and NSUN6, respectively[21,46]. In line with this, sanger sequencing following RT-PCT showed a significant reduction in C-to-U or A-to-G mutations near the m^{5}C sites in methyltransferase-deficient cells compared with WT cells (Fig. 2G and H). Overall, these findings suggest that the DRAM detection system is assayed in an m^{5}C-dependent form.

DRAM enables transcriptome-wide analysis of m^{5}C methylation

Subsequently, we performed RNA-seq analysis after DRAM transfection by detecting C-to-U/A-to-G editing events to accomplish transcriptome-wide detection of m^{5}C (Fig.3A). To serve as positive controls, two previously published BS-seq datasets were also integrated[5,21]. Mutations were detected near the m^{5}C site in RPSA as A-to-G by DRAM-ABE (Fig.3B), and DRAM-CBE detected the presence of C-to-U mutations near the AP5Z1 m^{5}C site (Fig.3C). However, the DRAM^{mut} and Deaminase systems induced few effective mutations close to these sites. Examination of multiple reported high-confidence RNA m^{5}C sites showed that DRAM-seq

editing events were also enriched in the vicinity of the BS-seq sites (Fig.3B, 3C and Fig. S5).

DRAM-seq analysis further confirmed that DRAM was detected in an m^{5}C-dependent
manner and that mutations in AP5Z1 and RPSA mRNA were reduced in methyltransferase
knockout cells compared to wild-type cells (Fig. 3D, 3E). Moreover, the knockout cells exhibited
overall rare DRAM-seq editing events close to m^{5}C sites in other mRNAs (Fig. S6). Unfortunately
however due to the inability of the DRAM detection system to achieve m^{5}C detection with single
base resolution, we did not find sequence preferences for the m^{5}C sites responsible for catalysis by
each of the two methyltransferases NSUN2 and NSUN6(Fig. S4D).

A comparison of three biological replicates from each experimental group revealed a strong
reproducibility of A-to-G/C-to-U mutations in HEK293T cells expressing DRAM-ABE and
DRAM-CBE (Fig. S7). Moreover, the integration and analysis of the experimental data revealed a
high degree of overlap between the three biological replicates (Fig. S4C). And a recent study by
Wang et al. showed that ALYREF deletion affects the expression of 94 mRNAs[47], and only 55.32%
of these ALYREF-regulated mRNAs can be detected by the DRAM system (Fig. S4E). These
findings suggest that DRAM selectively targets specific RNAs for editing, exhibiting a high
degree of consistency across samples.

To obtain information on a set of high-confidence DRAM-seq data, we filtered the list of
sites transfected with deaminase alone and screened the sequencing results with methyltransferase
depleted, pooled editing events occurring in at least 10% of reads across multiple samples to
obtain a set of high-confidence editing sites (Fig. 3F and Supplementary Table 2), and integrated
genes with editing sites occurring in DRAM-ABE and DRAM-CBE (Fig. 3F and Supplementary
Table 3).

Previous studies have indicated that m^{5}C sites are predominantly distributed in the coding
sequences (CDS) and notably enriched near the initiation codon[5,25,26,48-50]. To further delineate the
characteristics of the DRAM-seq data, we compared the distribution of DRAM-seq editing sites
within the gene structure, specifically examining their occurrences in the 5'untranslated region
(5'UTR), 3' untranslated region (3'UTR), CDS and Intergenic/Intron region. Our analysis
revealed that DRAM-seq editing events in cells expressing DRAM-ABE and DRAM-CBE were
primarily located in the CDS and 3'UTR, indicating a non-random distribution of m^{5}C (Fig.3G,
Fig. S8A and 8B). Moreover, plotting the distribution of DRAM-seq editing sites in mRNA
segments (5'UTR, CDS, and 3'UTR) highlighted a significant enrichment in the CDS (Fig.3H). In
contrast, cells expressing the deaminase exhibited a distinct distribution pattern of editing sites,
characterized by a prevalence throughout the 3'UTR (Fig.3H). This finding reaffirms that the
specific editing pattern observed in DRAM-seq across the transcriptome depends on its capacity to
bind m^{5}C.

Comparative analysis of the DRAM-seq editing sites with the previously published BS-seq
m^{5}C sites indicated that the likelihood of editing was notably higher in closer proximity to the
m^{5}C sites (Fig.3I). Furthermore, the editing window of DRAM exhibited enrichment
approximately 20bp before and after the m^{5}C site (Fig.3I). Investigation into the sequences
surrounding the editing window revealed that AC motifs were the most significantly enriched in
DRAM-CBE, whereas (U/C) A motifs were most notably enriched in DRAM-ABE. In contrast,
the APOBEC1 and TadA-8e samples displayed no significantly enriched motifs, with mutations
being more randomly orientated (Fig.3J, 3K).

DRAM-seq provides stable and comprehensive identification of m[5]C loci

Subsequently, we then evaluated the ability of DRAM-seq to detect m[5]C across the entire transcriptome and compared its performance to that of the previously reported BS-seq. Although both previous studies employed bisulfite treatment, the resulting data obtained significant discrepancies due to variations in their treatment and analysis methodologies. Our results indicated that DRAM-seq identified the presence of m[5]C modifications covering 79.6% of the genes detected by Yang et al.[5] and 91.9% of the genes detected by Zhang et al.[21] (Fig.4A and C). Remarkably, certain pivotal regulators with diverse biological functions, such as ATG16L1(coordinats autophagy pathway)[51] and ARHGEF25 (plays an important role in actin cytoskeleton reorganisation)[52], were identified by Zhang et al. and DRAM-seq, but not by Yang et al. (Fig.4B). Conversely, FANCD2 (Maintains chromosome stability)[53] and RPL15(components of the large ribosomal subunit)[54,55], were discovered by Yang et al. and DRAM-seq, but not by Zhang et al. (Fig.4D). Hence, DRAM-seq appears to offer a more stable and comprehensive identification of the m[5]C loci.

To provide functional insights into m[5]C RNA-modified genes in HEK293T cells, we conducted Gene Ontology (GO) and Kyoto Encyclopedia of Genes and Genomes (KEGG) analyses. These results highlighted the involvement of these genes in the regulation of diverse key biological processes, such as cell division, cell cycle, mRNA splicing, protein processing in the endoplasmic reticulum, nucleocytoplasmic transport, translation, DNA repair and others (Fig.4E, 4F, Fig. S8C and S8D).

DRAM monitors m[5]C dynamics in cellular RNA

Next, we aim to determine whether changes in m[5]C can be detected in real time by DRAM. Previous studies have established that oxidative stress induced by 200 μM sodium arsenite (NaAsO$_2$) treatment inhibited the expression of NSUN2 and NSUN6, resulting in a gradual decrease in m[5]C methylation[56,57]. As cells exposed to NaAsO$_2$ typically exhibited slowed cell cycle progression and NaAsO$_2$-induced apoptosis, we performed a low concentration screen[58,59]. After gradient treatment of NaAsO$_2$ (50μM, 100μM, 150μM, and 200μM, separately) in cells, 200μM group was identified as the optimal concentration, leading to highly significant decreases of NSUN2 and NSUN6 expression (Fig.5A). Subsequently, HEK293T cells expressing DRAM with NaAsO$_2$ for 3 hours revealed a decreasing trend in the mRNA expression of NSUN2 and NSUN6 (Fig.5B). Bisulfite sequencing PCR also verified a decreasing trend in m[5]C levels in selected mRNAs after NaAsO$_2$ treatment (Fig. 5C and D).

Next, we performed DRAM system assays, and Sanger sequencing results showed that induction of the DRAM system after NaAsO$_2$ treatment resulted in a highly significant reduction of A-to-G/C-to-U mutations on both RPSA and AP5Z1 mRNAs compared to cells without NaAsO$_2$ treatment (Fig.5E and F). Taken together, with our previous evidence that the methyltransferase-deleted cells exhibited overall rare DRAM editing events close to m[5]C sites, we suggest that the DRAM could effectively monitor the accumulation of m[5]C in individual RNAs in response to the alteration of cellular conditions.

DRAM enables low-input m[5]C profiling

A significant challenge in m[5]C detection lies in the specificity of antibodies and the substantial amount of input RNA required for sequencing. RNA is susceptible to degradation

during denaturation, sodium bisulfite treatment and desulfurization steps in the BS-seq assay[60].
Immunoprecipitation-based m[5]C assays and LC-MS/MS also impose high demand for sample
input[7,25,61]. Several experiments have highlighted the requirement of 100-500 ng of RNA for
m[5]C-RIP-seq, while BS-seq necessitates an even more demanding 750-1000 ng of RNA[21,25,62]. To
assess the detection limits of DRAM-Sanger, we attempted to amplify two representative
m[5]C-containing sites in the RPSA and AP5Z1 transcripts from diluted RNA samples.

Remarkably, we successfully generated PCR products of these two mRNAs from cDNAs
corresponding to 250 ng, 50 ng, and 10 ng of total RNA. Quantitative analysis by Sanger
sequencing demonstrated nearly identical Sanger traces across these dilutions (Fig.5G and H).
This finding underscores that the specificity of DRAM editing depended on its ability to bind m[5]C,
and DRAM is proficient in low-input m[5]C analyses. Furthermore, cell viability was determined by
CCK8 assay on HEK293T cells transfected with DRAM (Fig.5I). Importantly, there was no
significant difference in the relative proliferative capacity of the cells compared to untransfected
cells (NC), indicating that DRAM expression did not adversely affect cell viability (Fig.5J).

Transfection of the DRAM system in cells results in the transient overexpression of fusion
proteins. To investigate how varying expression levels of these proteins influence A-to-G and
C-to-U editing within the same m[5]C region, we conducted a gradient transfection using plasmid
concentrations of 1500 ng, 750 ng and 300 ng. This approach allowed us to progressively reduce
the expression levels of the fusion proteins (Fig. S9A and S9B). Sanger sequencing revealed that
the editing efficiency of A-to-G and C-to-U within the m[5]C region significantly decreased as
fusion protein expression diminished (Fig. S9C and S9D). These findings suggest that the
transfection efficiency of the DRAM system is concentration-dependent and that the ratio of
editing efficiency to transfection efficiency may assist in the quantitative analysis of m[5]C using
the DRAM system.

279
280

Discussion

In recent years, m[5]C methylation modifications have received increasing attention, with
multiple reports detailing the distribution of RNA m[5]C methylation modifications across various
species and tissues, elucidating their characteristics. Despite the relatively low abundance of m[5]C,
its highly dynamic changes hold significant implications for the regulation of physiological and
pathological processes[5,21,44]. However, due to the limitations of sequencing methods and the
variability of data processing, there remains ample room for progress in the study of m[5]C
detection methods.

In this study, we developed a site-specific, depth-sequencing-free m[5]C detection method
using DRAM-Sanger. This workflow relies on conventional molecular biology assays such as
RT-PCR and Sanger sequencing, eliminating the need for specialized techniques and thereby
simplifying the process of m[5]C detection.

DRAM-seq introduces a novel strategy for transcriptome-wide m[5]C detection, overcoming
inherent limitations in existing methods. Notably, DRAM-seq covered around 80% of the
high-confidence m[5]C-modified genes detected by BS-seq and identified more potential m[5]C sites.
This can be attributed to the avoidance of bisulfite treatment by DRAM-seq, preventing RNA
damage and ensuring a more comprehensive representation of RNA samples. This feature also

likely contributes to the observed stability of DRAM-seq in comparison to BS-seq. Additionally,
DRAM-seq is not limited by antibody specificity and is resistant to chemical-induced damage.

A prominent challenge in existing m^5C profiling methods is their reliance on substantial
amounts of input RNA samples. In contrast, DRAM operates through the deamination activity of
deamination activity of deaminase, preserving RNA integrity and preventing degradation. The
notable advantage of DRAM lies in its capacity for low-input m^5C detection. Our analysis
demonstrates that DRAM requires as low as 10ng of total RNA for m^5C detection. While DRAM
is currently well-suited for detecting m^5C on a transcriptome-wide scale, the potential for future
applications involving third-generation sequencing could extend its utility to individual mRNAs,
particularly m^5C heterogeneity on mRNA splicing variants. In addition, the DRAM system
depends on the specific recognition of m^5C modifications on ssRNA by the reader protein,
theoretically avoiding the false-positive effects of 5-hydroxymethylation modifications in other
assays, such as BS-seq[21-23]. This potential feature could enhance the accuracy of the DRAM assay,
albeit it still requires careful validation.

In our study, m^5C detection was performed following the transient transfection of the DRAM
detection system into mammalian cells, which might result in a lower mutation rate at the
corresponding site. Therefore, employing lentiviral-mediated transfection into cell lines of interest
could potentially enhance the efficiency of m^5C detection. In the present study, the editing
efficiency of the DRAM system, which is believed to detect dynamic changes in m^5C, was
significantly reduced following sodium arsenite treatment. However, sodium arsenite can
adversely affect all proteins containing thiol groups, impacting not only NSUN proteins but
potentially also the function of reader proteins and deaminases. As a result, this validation cannot
entirely rule out the possibility that the observed decrease in editing frequency is due to the
inactivation of fusion proteins by sodium arsenite. The optimization of this validation may require
the development of more specific m^5C inhibitors in the future. Further elucidation of the key
amino acids directing ALYREF and YBX1's binding to m^5C methylation sites should enable more
accurate and sensitive m^5C detection by DRAM-seq. Due to the lack of a fixed base composition
for characterizing all m^5C modification sites, DRAM has an apparent limitation in achieving
single-base resolution for detecting m^5C. However, our present study proved that the measuring
resolution of DRAM is around 40nt, which facilitates higher precision than that of m^5C-RIP-seq
(~100nt). In the future, with more in-depth analyses of m^5C reader structures and the identification
of new potential m^5C readers, we expect to achieve more precise m^5C localization and more
comprehensive m^5C modification detection. Moreover, the substitution of deaminases, such as
A3A and A3G (the family members of APOBEC), could also potentially enhance the efficiency of
the DRAM detection [63-65].

Following the purification of DRAM fusion protein, achieving *in vitro* detection of RNA m^5C
methylation could broaden the applicability to a more diverse range of samples. Another potential
application for DRAM-seq could be the expression of drug-inducible DRAM systems *in vivo*
using various animal models for m^5C analysis. These will together provide novel insights into
m^5C modifications for biological and clinical research.

Conclusions

In summary, we developed a novel deaminase and reader protein-assisted RNA m^5C
methylation approach that detects the m^5C region by deaminating As or Cs in close proximity to

the m^{5}C sites, which does not rely on antibodies or bisulfite, thus leading to unprecedently

comprehensive transcriptome-wide RNA m^{5}C methylation profiling. We anticipated that this

system could pave the way for uncovering further biological functions of m^{5}C modifications and

facilitate the development of therapeutic interventions for associated diseases.

Materials and methods

Plasmid construction

ALYREF and YBX1 expression plasmids were purchased from MIAOLING BIOLOGY

(http://www.miaolingbio.com/), and the ALYREF and YBX1 fractions were then amplified using

specific primer. The ALYREF and YBX1 portions were amplified using pCMV-APOBEC1-YTH

(Addgene plasmid no. 131636; https://www.addgene.org/131636/) and ABE8e (Addgene plasmid

no. 138489; https://www.addgene.org/ 138489/) to amplify the deaminase portion and the essential

plasmid construct proxies, and finally the fragments were recombined by the ClonExpress Ultra

One Step Cloning Kit to complete the plasmid vector construction. Both DRAM^{mut}-ABE and

DRAM^{mut}-CBE related vectors were obtained by introducing the corresponding key amino acid

mutations using Fast Site-Directed Mutagenesis Kit (TIANGEN Biotech). The primer sequences

used are listed in Supplementary Table 4.

Cell culture and plasmid transfection

HEK293T cell line (ATCC) was cultured in Dulbecco's Modified Eagle Medium (DMEM)

supplemented with 10% fetal bovine serum (CLARK BIOSCIENCE) and 1% penicillin (100

U/ml)-streptomycin (100μg/ml). The cells were seeded in 12-well plates and transfected using

Hieff Trans™ Liposomal Transfection Reagent (Yeasen).

NSUN2-depleted cell lines were generated by cloning NSUN2-targeting single guide RNA

sequences into the pSpCas9(BB)–2A-Puro (PX459) V2.0 plasmid (Addgene plasmid no. 62988;

http://n2t.net/addgene:62988). Plasmids were then transfected into HEK293T cells and Puromycin

(Meilunbio) was added at a final concentration of 3 μg/ml to enrich the positively transfected cells

24 h after transfection. After 72 h, the cells were collected and used for genotyping by Sanger

sequencing. NSUN6-depleted cell lines were generated in the same way. The primers used for

genotyping and single guide RNA sequences are listed in Supplemental Table 4.

Cell viability measurements

HEK293T cells were transfected with DRAM plasmid and cultured at 37°C for 24 h.

Subsequently, 1000 cells were seeded in 96-well plates. After waiting for the cells to attach to the

wall, the cell activity was detected by Cell Counting Kit-8 (Meilunbio). Cell Counting Kit-8

contains WST-8, which in the presence of the electronically coupled reagent 1-Methoxy PMS can

be reduced by mitochondrial dehydrogenase to the orange-colored metazan product Formazan, the

absorbance of which is measured at 450 nm to analyze cellular activity.

Western blotting

For protein blotting, samples were lysed in RIPA Lysis Buffer (Meilunbio) with

Phenylmethanesulfonyl fluoride (PMSF) and the BCA protein assay kit (Beyotime Biotechnology)

was used to Protein concentration was measured. Total protein extracts were separated by

SDS-PAGE on a 10% gel and then transferred to 0.22 nm polyvinylidene fluoride membranes

(Boster). Subsequently, the proteins were probed with specific antibodies after the blot was

blocked with 5% non-fat milk (Boster). Images were quantified using ImageJ software and all data
are expressed as mean ± SEM.

The following antibodies and concentrations were used: NSUN2 Polyclonal antibody
(Proteintech; Cat No.20854-1-AP; 1:7500), NSUN6 Polyclonal antibody (Proteintech; Cat No.
17240-1-AP; 1:2000), RabbitAnti-GAPDH antibody (Bioss; bs-41373R; 1:2000), Alpha Tubulin
Polyclonal antibody (Proteintech; Cat No. 11224-1-AP; 1:2000), HRP-labeled Goat Anti-Rabbit
IgG(H+L) (Beyotime Biotechnology; A0208; 1:2000).

cDNA synthesis and Sanger sequencing

Total cellular RNA was extracted with TRIzol reagent (TIANGEN Biotech) and cDNA was
synthesized using PrimeScript™ II 1st Strand cDNA Synthesis Kit (Takara Bio) according to the

manufacturer's recommendations. PCR was then performed using 2×Taq PCR MasterMix II

(TIANGEN Biotech) and primers flanking m^5C target sites, and the purified PCR products were
directly sequenced by Sanger sequencing. The Sanger sequencing results were analyzed using
EditR 1.0.10 to calculate the mutation frequency[66]. The primers used in this study are shown in
Supplemental Table 4.

Real-time quantitative PCR

cDNA was synthesized using FastKing RT kit (with gDNase) (TIANGEN Biotech) according
to the manufacturer's recommendations. RT-qPCR assay was performed using SuperReal PreMix
Plus (SYBR Green) (TIANGEN Biotech). GAPDH was used as an endogenous control, and the
expression levels were normalized to the control and calculated by the $2^{-\Delta\Delta Ct}$ formula. All samples
were analyzed in triplicate and each mRNA quantification represents the average of at least three
measurements. All data are expressed as mean ± SEM. The primers used in this study are shown in
Supplemental Table 4.

Protein structure modelling

Protein structure simulations were performed using the SWISS-MODEL online website
(https://swissmodel.expasy.org/interactive)[67]. The SWISS-MODEL database is able to provide
up-to-date annotated 3D protein models, which are generated from automated homology
modelling of related model organisms and experimental structural information for all sequences in
UniProtKB, with reliable structural information, and subsequently protein structure observations
were performed using PyMOL[68].

Bisulfite sequencing PCR

We referenced bisulfite sequencing PCR, an assay established by Matthias Schaefer et al. We
chemically deaminated cytosine in RNA using the EZ RNA methylation kit (50) (ZYMO
RESEARCH) and then quantified m^5C methylation levels based on PCR amplification of cDNA
combined with deep sequencing[23].

RNA Conversion Reagent was premixed with prepared RNA samples, and the RNA was
denatured at 70°C for 5 minutes, followed by a reaction period of 45 minutes at 54°C. Finally, the
purified RNA samples were recovered after desulfurization by RNA Desulphonation Buffe. cDNA
was synthesized using PrimeScript™ II 1st Strand cDNA Synthesis Kit (Takara Bio) according to
the manufacturer's recommendations. PCR was then performed using 2× EpiArt HS Taq Master
Mix (Dye Plus) (Vazyme) and m^5C target site-specific Bisulfite Primer (primer sequences were
designed at https://zymoresearch.eu/pages/bisulfite-primer- seeker), the products were purified by
TIANgel Midi Purification Kit (TIANGEN Biotech), and the connectors for second-generation

sequencing were attached at both ends of the products for sequencing. Finally, deep sequencing
was performed by HiTOM analysis to detect the methylation level (The number of reads >1000 in
deep sequencing) [69]. The primers used in this study are shown in Supplemental Table 4.

Library construction and next-generation sequencing

1μg of total cellular RNA was used for sequencing library generation by NEBNext Ultra
RNA Library Prep Kit for Illumina (NEB, USA, Catalog #: E7530L) following manufacturer's
recommendations and index codes were added to attribute sequences to each sample. Briefly,
mRNA was purified from total RNA using poly-T oligo-attached magnetic beads. Fragmentation
was carried out using divalent cations under elevated temperature in NEB Next First Strand
Synthesis Reaction Buffer(5X). First-strand cDNA was synthesized using random hexamer primer
and M-MuLV Reverse Transcriptase (RNase H). Second-strand cDNA synthesis was subsequently
performed using DNA Polymerase I and RNase H. Remaining overhangs was converted into blunt
ends via exonuclease/polymerase activities. After adenylation of 3' ends of DNA fragments, NEB
Next Adaptor with hairpin loop structure was ligated to prepare for hybridization. To select cDNA
fragments of preferentially 370~420 bp in length, the library fragments were purified with
AMPure XP system (Beverly, USA). Then 3 μL USER Enzyme (NEB, USA) was used with size
selected, adaptor-ligated cDNA at 37°C for 15 min followed by 5 min at 95 °C before PCR. Then
PCR was performed with Phusion High-Fidelity DNA polymerase, Universal PCR primers and
Index (X) Primer. At last, PCR products were purified (AMPure XP system) and library quality
was assessed on the Agilent 5400 system(Agilent, USA)and quantified by QPCR (library
concentration ≥ 1.5 nM). The qualified libraries were pooled and sequenced on Illumina
platforms with PE150 strategy in Novogene Bioinformatics Technology Co., Ltd (Beijing, China),
according to effective library concentration and data amount required.

DRAM-seq analysis and calling of edited sites

The raw fastq sequencing data were cleaned by trimming the adapter sequences using Fastp
(v0.23.1) and were aligned to the human genome (hg19) using STAR (v2.7.7) in paired-end mode.
The aligned BAM files were sorted and PCR duplicates were removed using Samtools (v1.12).
The cite calling pf DRAM-seq was performed using Bullseye, a previously customized pipeline to
look for C-to-U or A-to-G edited sites throughout the transcriptome[40]. Briefly, the sorted and
deduplicated BAM files were initially parsed by parseBAM.pl script.

Then, Find_edit_site.pl script was employed to find C-to-U or A-to-G editing events by
DRAM-seq with at least 10 reads of coverage, an edit ratio of 5%-95%, an edit ratio at least
1.5-fold higher than NSUN2 or NSUN6-knockout samples, and at least 2 editing events at a given
site. Sites that were only found in one replicate of each DRAM protein variant were removed.
Editing events appeared in cells expressing merely APOBEC1 or TadA8e were also removed. For
high confidence filtering, we further adjusted the Find_edit_site.pl parameters to the edit ratio of
10%-60%, an edit ratio of control samples at least 2-fold higher than NSUN2 or NSUN6-knockout
samples, and at least 4 editing events at a given site.

Metagene and motif analyses

Metagene analysis was performed using hg19 annotations according to previously reported
tool, MetaplotR[70]. For motif analysis, the 20bp flanking sequence of each DRAM-seq editing site
was extracted by Bedtools (v2.30.0)[71]. The motif logos were then plotted by WebLogo (v3.7.12)[72].

Replicates analysis

Independent biological replicates of DRAM-ABE or DRAM-CBE in DRAM-seq analysis
were separately compared by computing the Pearson correlation coefficient between the number
of C-to-U mutations per mRNA between any two replicate experiments.

GO and KEGG analysis

GO and KEGG analysis of DRAM-seq edited mRNAs was performed using the DAVID
bioinformatic database [73]. GO terms with a P value of less than 0.05 were considered statistically
significant.

RNA pulldown assays

RNAs containing C or m^5C residues and proteins expressing DRAM or DRAM^{mut} were
prepared in advance, processed using the Pierce™ Magnetic RNA-Protein Pull-Down Kit
(Thermo) and finally analysed by Western blotting.

Statistical analysis

All data are expressed as mean ± S.E.M of three independent determinations. Data were
analyzed through a two-tailed t-test. A probability of $P < 0.05$ was considered statistically
significant; □, $P < 0.05$, □□, $P < 0.01$, *, $P < 0.05$, **, $P < 0.01$, ***, $P < 0.001$ and ****, $P <$
0.0001 denote the significance thresholds; ns denotes not significant.

Data and Materials Availability

487   The data supporting the findings of this study are available within the article and its
488   Supplementary Information. Other data and reagents are available from the corresponding authors
489   upon reasonable request.

AUTHOR CONTRIBUTIONS

Conceptualization: JZ, YH, LL, ZL

Methodology: JZ, DZ, JL

Investigation: JZ, DZ, JL, DK, XL, RZ, YL

Visualization: XG, YQ, DW, JC

Supervision: DK, XL, RZ, YL, XG, YQ, DW, JC, YH

Funding acquisition: YH, LL, ZL

Data curation: JZ, YH

Writing—original draft: JZ, DZ, JL

Writing—review & editing: JZ, YH, LL, ZL

Competing Interests

503   All other authors declare they have no competing interests.

ACKNOWLEDGEMENTS

506   We thank Yuning Song, Yuanyuan Xu and Tingting Sui for critical feedbacks on the work and

507 manuscript.

FUNDING

509 This work was supported by the National Natural Science Foundation of China
510 (Nos.32200466).

REFERENCES

1   Wiener, D. & Schwartz, S. The epitranscriptome beyond m(6)A. *Nat Rev Genet* **22**, 119-131
(2021). https://doi.org:10.1038/s41576-020-00295-8
2   Li, S. & Mason, C. E. The pivotal regulatory landscape of RNA modifications. *Annu Rev*
Genomics   Hum   Genet   **15**,   127-150   (2014).
https://doi.org:10.1146/annurev-genom-090413-025405
3   Van Haute, L. *et al.* NSUN2 introduces 5-methylcytosines in mammalian mitochondrial
tRNAs. *Nucleic Acids Res* **47**, 8720-8733 (2019). https://doi.org:10.1093/nar/gkz559
4   Blaze, J. *et al.* Neuronal Nsun2 deficiency produces tRNA epitranscriptomic alterations and
proteomic shifts impacting synaptic signaling and behavior. *Nat Commun* **12**, 4913 (2021).
https://doi.org:10.1038/s41467-021-24969-x
5   Yang, X. *et al.* 5-methylcytosine promotes mRNA export - NSUN2 as the methyltransferase
and   ALYREF   as   an   m(5)C   reader.   *Cell   Res*   **27**,   606-625   (2017).
https://doi.org:10.1038/cr.2017.55
6   Sharma, S., Yang, J., Watzinger, P., Kötter, P. & Entian, K. D. Yeast Nop2 and Rcm1 methylate
C2870 and C2278 of the 25S rRNA, respectively. *Nucleic Acids Res* **41**, 9062-9076 (2013).
https://doi.org:10.1093/nar/gkt679
7   Hussain, S. *et al.* NSun2-mediated cytosine-5 methylation of vault noncoding RNA
determines its processing into regulatory small RNAs. *Cell Rep* **4**, 255-261 (2013).
https://doi.org:10.1016/j.celrep.2013.06.029
8   Aguilo, F. *et al.* Deposition of 5-Methylcytosine on Enhancer RNAs Enables the Coactivator
Function   of   PGC-1α.   *Cell   Rep*   **14**,   479-492   (2016).
https://doi.org:10.1016/j.celrep.2015.12.043
9   Yang, Y. *et al.* RNA 5-Methylcytosine Facilitates the Maternal-to-Zygotic Transition by
Preventing   Maternal   mRNA   Decay.   *Mol   Cell*   **75**,   1188-1202.e1111   (2019).
https://doi.org:10.1016/j.molcel.2019.06.033
10   García-Vílchez, R., Sevilla, A. & Blanco, S. Post-transcriptional regulation by cytosine-5
methylation of RNA. *Biochim Biophys Acta Gene Regul Mech* **1862**, 240-252 (2019).
https://doi.org:10.1016/j.bbagrm.2018.12.003
11   Trixl, L. & Lusser, A. The dynamic RNA modification 5-methylcytosine and its emerging role
as   an   epitranscriptomic   mark.   *Wiley   Interdiscip   Rev   RNA*   **10**,   e1510   (2019).
https://doi.org:10.1002/wrna.1510
12   Boo, S. H. & Kim, Y. K. The emerging role of RNA modifications in the regulation of mRNA
stability. *Exp Mol Med* **52**, 400-408 (2020). https://doi.org:10.1038/s12276-020-0407-z
13   Courtney, D. G. *et al.* Epitranscriptomic Addition of m(5)C to HIV-1 Transcripts Regulates
Viral   Gene   Expression.   *Cell   Host   Microbe*   **26**,   217-227.e216   (2019).

548 https://doi.org/10.1016/j.chom.2019.07.005

549 14 Liu, J. *et al.* Developmental mRNA m(5)C landscape and regulatory innovations of massive
550 m(5)C modification of maternal mRNAs in animals. *Nat Commun* **13**, 2484 (2022).
551 https://doi.org/10.1038/s41467-022-30210-0

552 15 Chen, P., Zhang, T., Yuan, Z., Shen, B. & Chen, L. Expression of the RNA methyltransferase
553 Nsun5 is essential for developing cerebral cortex. *Mol Brain* **12**, 74 (2019).
554 https://doi.org/10.1186/s13041-019-0496-6

555 16 Flores, J. V. *et al.* Cytosine-5 RNA Methylation Regulates Neural Stem Cell Differentiation
556 and Motility. *Stem Cell Reports* **8**, 112-124 (2017).
557 https://doi.org/10.1016/j.stemcr.2016.11.014

558 17 Chen, S. Y. *et al.* RNA bisulfite sequencing reveals NSUN2-mediated suppression of epithelial
559 differentiation in pancreatic cancer. *Oncogene* **41**, 3162-3176 (2022).
560 https://doi.org/10.1038/s41388-022-02325-7

561 18 Chen, X. *et al.* 5-methylcytosine promotes pathogenesis of bladder cancer through stabilizing
562 mRNAs. *Nat Cell Biol* **21**, 978-990 (2019). https://doi.org/10.1038/s41556-019-0361-y

563 19 Yang, R. *et al.* The RNA methyltransferase NSUN6 suppresses pancreatic cancer development
564 by regulating cell proliferation. *EBioMedicine* **63**, 103195 (2021).
565 https://doi.org/10.1016/j.ebiom.2020.103195

566 20 Li, Y. *et al.* Long noncoding RNA DIAPH2-AS1 promotes neural invasion of gastric cancer
567 via stabilizing NSUN2 to enhance the m5C modification of NTN1. *Cell Death Dis* **14**, 260
568 (2023). https://doi.org:10.1038/s41419-023-05781-5

569 21 Huang, T., Chen, W., Liu, J., Gu, N. & Zhang, R. Genome-wide identification of mRNA
570 5-methylcytosine in mammals. *Nat Struct Mol Biol* **26**, 380-388 (2019).
571 https://doi.org/10.1038/s41594-019-0218-x

572 22 Trixl, L., Rieder, D., Amort, T. & Lusser, A. Bisulfite Sequencing of RNA for
573 Transcriptome-Wide Detection of 5-Methylcytosine. *Methods Mol Biol* **1870**, 1-21 (2019).
574 https://doi.org/10.1007/978-1-4939-8808-2_1

575 23 Schaefer, M., Pollex, T., Hanna, K. & Lyko, F. RNA cytosine methylation analysis by bisulfite
576 sequencing. *Nucleic Acids Res* **37**, e12 (2009). https://doi.org:10.1093/nar/gkn954

577 24 Amort, T. & Lusser, A. Detection of 5-Methylcytosine in Specific Poly(A) RNAs by Bisulfite
578 Sequencing. *Methods Mol Biol* **1562**, 107-121 (2017).
579 https://doi.org/10.1007/978-1-4939-6807-7_8

580 25 Cui, X. *et al.* 5-Methylcytosine RNA Methylation in Arabidopsis Thaliana. *Mol Plant* **10**,
581 1387-1399 (2017). https://doi.org/10.1016/j.molp.2017.09.013

582 26 Edelheit, S., Schwartz, S., Mumbach, M. R., Wurtzel, O. & Sorek, R. Transcriptome-wide
583 mapping of 5-methylcytidine RNA modifications in bacteria, archaea, and yeast reveals m5C
584 within archaeal mRNAs. *PLoS Genet* **9**, e1003602 (2013).
585 https://doi.org/10.1371/journal.pgen.1003602

586 27 Saplaoura, E., Perrera, V., Colot, V. & Kragler, F. Methylated RNA Immunoprecipitation
587 Assay to Study m5C Modification in Arabidopsis. *J Vis Exp* (2020).
588 https://doi.org/10.3791/61231

589 28 Khoddami, V. & Cairns, B. R. Identification of direct targets and modified bases of RNA
590 cytosine methyltransferases. *Nat Biotechnol* **31**, 458-464 (2013).
591 https://doi.org/10.1038/nbt.2566

29 Yuan, F. *et al.* Bisulfite-free and base-resolution analysis of 5-methylcytidine and
5-hydroxymethylcytidine in RNA with peroxotungstate. *Chem Commun (Camb)* **55**,
2328-2331 (2019). https://doi.org/10.1039/c9cc00274j

30 Hayashi, G. *et al.* Base-Resolution Analysis of 5-Hydroxymethylcytosine by One-Pot
Bisulfite-Free Chemical Conversion with Peroxotungstate. *J Am Chem Soc* **138**, 14178-14181
(2016). https://doi.org/10.1021/jacs.6b06428

31 Harel, N., Meir, M., Gophna, U. & Stern, A. Direct sequencing of RNA with MinION
Nanopore: detecting mutations based on associations. *Nucleic Acids Res* **47**, e148 (2019).
https://doi.org/10.1093/nar/gkz907

32 Chang, J. J. *et al.* Transcriptional and epi-transcriptional dynamics of SARS-CoV-2 during
cellular infection. *Cell Rep* **35**, 109108 (2021). https://doi.org/10.1016/j.celrep.2021.109108

33 Wan, Y. K., Hendra, C., Pratanwanich, P. N. & Göke, J. Beyond sequencing: machine learning
algorithms extract biology hidden in Nanopore signal data. *Trends Genet* **38**, 246-257 (2022).
https://doi.org/10.1016/j.tig.2021.09.001

34 Budzko, L., Hoffa-Sobiech, K., Jackowiak, P. & Figlerowicz, M. Engineered deaminases as a
key component of DNA and RNA editing tools. *Mol Ther Nucleic Acids* **34**, 102062 (2023).
https://doi.org/10.1016/j.omtn.2023.102062

35 Rosenberg, B. R., Hamilton, C. E., Mwangi, M. M., Dewell, S. & Papavasiliou, F. N.
Transcriptome-wide sequencing reveals numerous APOBEC1 mRNA-editing targets in
transcript 3' UTRs. *Nat Struct Mol Biol* **18**, 230-236 (2011).
https://doi.org/10.1038/nsmb.1975

36 Qian, X. *et al.* Low expression of the apolipoprotein B mRNA-editing transgene in mice
reduces LDL levels but does not cause liver dysplasia or tumors. *Arterioscler Thromb Vasc*
Biol **18**, 1013-1020 (1998). https://doi.org/10.1161/01.atv.18.6.1013

37 Siriwardena, S. U., Chen, K. & Bhagwat, A. S. Functions and Malfunctions of Mammalian
DNA-Cytosine Deaminases. *Chem Rev* **116**, 12688-12710 (2016).
https://doi.org/10.1021/acs.chemrev.6b00296

38 Richter, M. F. *et al.* Phage-assisted evolution of an adenine base editor with improved Cas
domain compatibility and activity. *Nat Biotechnol* **38**, 883-891 (2020).
https://doi.org/10.1038/s41587-020-0453-z

39 Gaudelli, N. M. *et al.* Directed evolution of adenine base editors with increased activity and
therapeutic application. *Nat Biotechnol* **38**, 892-900 (2020).
https://doi.org/10.1038/s41587-020-0491-6

40 Meyer, K. D. DART-seq: an antibody-free method for global m(6)A detection. *Nat Methods*
16, 1275-1280 (2019). https://doi.org/10.1038/s41592-019-0570-0

41 Sun, Z. *et al.* Effects of NSUN2 deficiency on the mRNA 5-methylcytosine modification and
gene expression profile in HEK293 cells. *Epigenomics* **11**, 439-453 (2019).
https://doi.org/10.2217/epi-2018-0169

42 Zhang, Q., Zheng, Q., Yu, X., He, Y. & Guo, W. Overview of distinct 5-methylcytosine
profiles of messenger RNA in human hepatocellular carcinoma and paired adjacent non-tumor
tissues. *J Transl Med* **18**, 245 (2020). https://doi.org/10.1186/s12967-020-02417-6

43 Zhang, Q. *et al.* The role of RNA m(5)C modification in cancer metastasis. *Int J Biol Sci* **17**,
3369-3380 (2021). https://doi.org/10.7150/ijbs.61439

44 Liu, J. *et al.* Sequence- and structure-selective mRNA m(5)C methylation by NSUN6 in

animals. *Natl Sci Rev* **8**, nwaa273 (2021). https://doi.org:10.1093/nsr/nwaa273

45   Schumann, U. *et al.* Multiple links between 5-methylcytosine content of mRNA and
translation. *BMC Biol* **18**, 40 (2020). https://doi.org:10.1186/s12915-020-00769-5

46   Fang, L. *et al.* CIGAR-seq, a CRISPR/Cas-based method for unbiased screening of novel
mRNA modification regulators. *Mol Syst Biol* **16**, e10025 (2020).
https://doi.org:10.15252/msb.202010025

47   Wang, N. *et al.* m(5)C-dependent cross-regulation between nuclear reader ALYREF and writer
NSUN2 promotes urothelial bladder cancer malignancy through facilitating RABL6/TK1
mRNAs splicing and stabilization. *Cell Death Dis* **14**, 139 (2023).
https://doi.org:10.1038/s41419-023-05661-y

48   Tang, Y. *et al.* OsNSUN2-Mediated 5-Methylcytosine mRNA Modification Enhances Rice
Adaptation to High Temperature. *Dev Cell* **53**, 272-286.e277 (2020).
https://doi.org:10.1016/j.devcel.2020.03.009

49   Squires, J. E. *et al.* Widespread occurrence of 5-methylcytosine in human coding and
non-coding RNA. *Nucleic Acids Res* **40**, 5023-5033 (2012).
https://doi.org:10.1093/nar/gks144

50   Zhang, P., Wang, Y. & Gu, X. RNA 5-Methylcytosine Controls Plant Development and
Environmental Adaptation. *Trends Plant Sci* **25**, 954-958 (2020).
https://doi.org:10.1016/j.tplants.2020.07.004

51   Fletcher, K. *et al.* The WD40 domain of ATG16L1 is required for its non-canonical role in
lipidation of LC3 at single membranes. *EMBO J* **37** (2018).
https://doi.org:10.15252/embj.201797840

52   Guo, X. *et al.* A Rac/Cdc42-specific exchange factor, GEFT, induces cell proliferation,
transformation, and migration. *J Biol Chem* **278**, 13207-13215 (2003).
https://doi.org:10.1074/jbc.M208896200

53   Naim, V. & Rosselli, F. The FANC pathway and BLM collaborate during mitosis to prevent
micro-nucleation and chromosome abnormalities. *Nat Cell Biol* **11**, 761-768 (2009).
https://doi.org:10.1038/ncb1883

54   Anger, A. M. *et al.* Structures of the human and Drosophila 80S ribosome. *Nature* **497**, 80-85
(2013). https://doi.org:10.1038/nature12104

55   Liang, X. *et al.* Structural snapshots of human pre-60S ribosomal particles before and after
nuclear export. *Nat Commun* **11**, 3542 (2020). https://doi.org:10.1038/s41467-020-17237-x

56   Gkatza, N. A. *et al.* Cytosine-5 RNA methylation links protein synthesis to cell metabolism.
PLoS Biol **17**, e3000297 (2019). https://doi.org:10.1371/journal.pbio.3000297

57   Ye, F. *et al.* SIRT1/PGC-1α is involved in arsenic-induced male reproductive damage through
mitochondrial dysfunction, which is blocked by the antioxidative effect of zinc. *Environ*
Pollut **320**, 121084 (2023). https://doi.org:10.1016/j.envpol.2023.121084

58   McCollum, G., Keng, P. C., States, J. C. & McCabe, M. J., Jr. Arsenite delays progression
through each cell cycle phase and induces apoptosis following G2/M arrest in U937 myeloid
leukemia cells. *J Pharmacol Exp Ther* **313**, 877-887 (2005).
https://doi.org:10.1124/jpet.104.080713

59   Ran, S. *et al.* NaAsO(2) decreases GSH synthesis by inhibiting GCLC and induces apoptosis
through Hela cell mitochondrial damage, mediating the activation of the NF-κB/miR-21
signaling pathway. *Ecotoxicol Environ Saf* **234**, 113380 (2022).

680   https://doi.org/10.1016/j.ecoenv.2022.113380

681   60   Schaefer, M. RNA 5-Methylcytosine Analysis by Bisulfite Sequencing. *Methods Enzymol* **560**,
682   297-329 (2015). https://doi.org/10.1016/bs.mie.2015.03.007

683   61   Bourgeois, G. *et al.* Eukaryotic rRNA Modification by Yeast
684   5-Methylcytosine-Methyltransferases and Human Proliferation-Associated Antigen p120.
685   *PLoS ONE* **10**, e0133321 (2015). https://doi.org:10.1371/journal.pone.0133321

686   62   Gu, X. & Liang, Z. Transcriptome-Wide Mapping 5-Methylcytosine by m(5)C RNA
687   Immunoprecipitation Followed by Deep Sequencing in Plant. *Methods Mol Biol* **1933**,
688   389-394 (2019). https://doi.org:10.1007/978-1-4939-9045-0_24

689   63   Kim, K., Shi, A. B., Kelley, K. & Chen, X. S. Unraveling the Enzyme-Substrate Properties for
690   APOBEC3A-Mediated RNA Editing. *J Mol Biol* **435**, 168198 (2023).
691   https://doi.org/10.1016/j.jmb.2023.168198

692   64   Barka, A. *et al.* The Base-Editing Enzyme APOBEC3A Catalyzes Cytosine Deamination in
693   RNA with Low Proficiency and High Selectivity. *ACS Chem Biol* **17**, 629-636 (2022).
694   https://doi.org/10.1021/acschembio.1c00919

695   65   Pan, Y. *et al.* Nanoscale Characterization of Interaction of APOBEC3G with RNA.
696   *Biochemistry* **56**, 1473-1481 (2017). https://doi.org/10.1021/acs.biochem.6b01189

697   66   Kluesner, M. G. *et al.* EditR: A Method to Quantify Base Editing from Sanger Sequencing.
698   *Crispr j* **1**, 239-250 (2018). https://doi.org:10.1089/crispr.2018.0014

699   67   Bienert, S. *et al.* The SWISS-MODEL Repository-new features and functionality. *Nucleic*
700   *Acids Res* **45**, D313-d319 (2017). https://doi.org:10.1093/nar/gkw1132

701   68   Delano, W. L. The PyMol Molecular Graphics System. *Proteins Structure Function and*
702   *Bioinformatics* **30**, 442-454 (2002).

703   69   Liu, Q. *et al.* Hi-TOM: a platform for high-throughput tracking of mutations induced by
704   CRISPR/Cas systems. *Sci China Life Sci* **62**, 1-7 (2019).
705   https://doi.org/10.1007/s11427-018-9402-9

706   70   Olarerin-George, A. O. & Jaffrey, S. R. MetaPlotR: a Perl/R pipeline for plotting metagenes of
707   nucleotide modifications and other transcriptomic sites. *Bioinformatics* **33**, 1563-1564 (2017).
708   https://doi.org/10.1093/bioinformatics/btx002

709   71   Quinlan, A. R. & Hall, I. M. BEDTools: a flexible suite of utilities for comparing genomic
710   features. *Bioinformatics* **26**, 841-842 (2010). https://doi.org/10.1093/bioinformatics/btq033

711   72   Crooks, G. E., Hon, G., Chandonia, J. M. & Brenner, S. E. WebLogo: a sequence logo
712   generator. *Genome Res* **14**, 1188-1190 (2004). https://doi.org/10.1101/gr.849004

713   73   Sherman, B. T. *et al.* DAVID: a web server for functional enrichment analysis and functional
714   annotation of gene lists (2021 update). *Nucleic Acids Res* **50**, W216-w221 (2022).
715   https://doi.org/10.1093/nar/gkac194

FIGURES LEGENDS

Fig.1: **Development of DRAM system for m^{5}C detection.**

(A) Schematic diagram of the DRAM assay. DRAM, DRAM^{mut} and Deaminase system were transfected into HEK293T cells separately. After DRAM transfection, the deaminase was directed by m^{5}C reader to the vicinity of the m^{5}C site and induce C-to-U/A-to-G mutations, whereas transfection of the DRAM^{mut} or Deaminase system failed to effectively induce similar mutations due to the absence of the m^{5}C-recognition-binding domain.

(B) The overall design of DRAM, DRAM^{mut} and Deaminase system.

Fig.2: DRAM detection system was assayed in an m⁵C -dependent form.

(**A, B**) Two m⁵C sites from RPSA (**A**) and AP5Z1 (**B**) mRNA detected by deep sequencing of bisulfite sequencing PCR in HEK293T cells. The m⁵C sites are highlighted by red color. The m□C fraction of RPSA and AP5Z1 were 75.5% and 27.25% (The number of reads is greater than 1000).

(**C, D**) Sanger sequencing following RT-PCR verified two m⁵C sites from RPSA (**C**) and AP5Z1 (**D**) mRNAs in DRAM-transfected HEK293T cells, respectively. HEK293T cells only expressing DRAM^mut or Deaminase were served as negative controls. The left panel illustrates the location of DRAM induced mutation sites, which is highlighted in red asterisk. The right panel shows the corresponding quantification of sanger sequencing.

(**E, F**) The knockout efficiency of NSUN2 (**E**) and NSUN6 (**F**) in HEK293T cell lines verified by Western blotting. The protein level of α-Tubulin and GAPDH were served as loading controls, separately.

741   **(G, H)** DRAM induced mutations close to m^{5}C sites in AP5Z1 (**G**) and RPSA (**H**) mRNAs after
742   NSUN2 and NSUN6 knockout in HEK293T cells. The left panel illustrates the location of DRAM
743   induced mutation sites, which is highlighted in red asterisk. The right panel shows the
744   corresponding quantification of sanger sequencing.
745

Fig.3: DRAM enables transcriptome-wide analysis of m⁵C methylation.

(A) Schematic of the DRAM-seq method.

(B, C) Integrative genomics viewer (IGV) browser traces of DRAM-seq data expressing the

indicated constructs in RPSA (**B**, left panel), TARBP2 (**B**, right panel), AP5Z1(**C**, left panel), and

TRAF7 (**C**, left panel) mRNAs. C-to-U or A-to-G mutations found in at least 10% of reads are

indicated by coloring. The previously published RNA BS-seq datasets from two individual studies

were displayed as panel "Yang et al." and "Zhang et al.". (n(DRAM)=3 independent samples,

n(Deaminase)=2 independent samples, and n(DRAM^mut)=1 independent sample.)

(D, E) Integrative genomics viewer (IGV) browser traces of DRAM-seq data in wildtype and

methyltranferases knockout cells in AP5Z1 (**D**) and RPSA (**E**) mRNAs. C-to-U or A-to-G

mutations were found in at least 10% of reads are indicated by coloring. The previously published

RNA BS-seq datasets from two individual studies were displayed as panel "Yang et al." and

"Zhang et al.". n=3 independent samples.

(F) Screening process for DRAM-seq assays and principles for screening high-confidence genes.

(G) The pie chart shows the distribution of editing sites in different transcript region in cells

expressing DRAM (n=3 independent samples).

(H) The density map showing the distribution of editing events across the mRNA transcripts

detected by DRAM-seq.

(I) The frequency plot shows the distribution of the distances of edit events in DRAM-seq relative

to the m⁵C sites from the published BS-seq datasets. The position of each m⁵C site of BS-seq is

determined as 0, and the relative distance of each site to the nearest edit event in DRAM-seq is

calculated and plotted. The plots are presented separately based on the cutoff of upstream and

downstream 3000bp (above) and 80bp (below) windows.

(J, K) Motif analysis discovered within the ±20nt region around the C-to-U or A-to-G editing site

in cells expressing DRAM-CBE **(J)**, APOBEC1**(J)**, DRAM-ABE **(K)** and TadA-8e **(K)**.

Fig.4: DRAM monitors m^{5}C dynamics in cellular RNA.

(**A**) Venn diagram showing the overlap between DRAM-seq and Yang et al.'s edited genes.

(**C**) Venn diagram showing the overlap between DRAM-seq and Zhang et al.'s edited genes.

(**B**, **D**) Integrative genomics viewer (IGV) browser traces of DRAM-seq data expressing the

indicated constructs in the ATG16L1(**B**), ARHGEF25(**B**), FANCD2(**D**), and RPL15(**D**) mRNAs.

C-to-U/A-to-G mutations found in at least 10% of reads are indicated by coloring, and the m^{5}C

site found by BS-seq is also labelled.

(**E**) Genes with DRAM-seq editing events were analyzed for KEGG bioprocess enrichment.

(**F**) GO biological processes enrichment analysis of genes with DRAM-seq editing events.

Statistical analyses were performed using the DAVID tool.

Fig.5: DRAM monitors m^{5}C dynamics in cellular RNA.

(**A**) Relative expression levels of NSUN2 and NSUN6 mRNA in HEK293T cells treated with 50

μM, 100 μM, 150μM and 200μM NaAsO$_2$.

(**B**) Relative expression levels of NSUN2 and NSUN6 mRNA in DRAM-expressing HEK293T

cells treated with 200μM NaAsO$_2$ for 3 hours.

(**C, D**) Bisulfite sequencing PCR showing the m^{5}C levels in RPSA and AP5Z1 mRNAs after 3 h

of control and NaAsO$_2$ treatment in cells.

(**E, F**) The m^{5}C adjacent mutations from RPSA (**C**) and AP5Z1 (**D**) mRNA in DRAM-expressing

HEK293T cells treated with NaAsO$_2$. The left panel indicates the sanger sequencing results

following RT-PCR, while the corresponding quantifications of DRAM-induced mutations are

shown on the right panel.

(**G, H**) DRAM analysis of RPSA (**G**) and AP5Z1 (**H**) mRNAs with 250 ng, 50 ng, and 10 ng of

input RNA. Representative Sanger sequencing plots are shown on the left panel, with mutation

sites marked with asterisks. The mutation rates are quantified on the right panel.

(**I**) Flowchart illustrating Cell viability analysis by CCK8 reagent after DRAM transfection in

HEK293T cells.

(**J**) Quantitative comparison of the relative proliferative capacity of DRAM-expressing and

untransfected cells.

803