Transcriptome-wide identification of 5-methylcytosine by deaminase and reader protein-assisted sequencing

Jiale Zhou[1][#], Ding Zhao[1][#], Jinze Li[2][#], Deqiang Kong[1], Xiangrui Li[2], Renquan Zhang[2],

Yuru Liang[2], Xun Gao[2], Yuqiang Qian[2], Di Wang[2], Jiahui Chen[2], Liangxue Lai [1,3,4*],

Yang Han[2*], Zhanjun Li [2*]

# These authors contributed equally: Jiale Zhou, Ding Zhao, Jinze Li.

[1]Key Laboratory of Zoonosis Research, Ministry of Education, College of Veterinary

Medicine, Jilin University, Changchun, 130062, China

[2]Laboratory Animal Center, College of Animal Science, Jilin University, Changchun,

130062, China

[3]Guangzhou Regenerative Medicine and Health Guang Dong Laboratory

(GRMH-GDL), Guangzhou, 510530, China

[4]Institute for Stem Cell and Regeneration, Chinese Academy of Sciences, Beijing,

510005, China

* To whom correspondence should be addressed:

Tel: (86)431-87836176; Fax: (86)431-87980131.

*e-mail:

hanyang8584@jlu.edu.cn (Yang Han); lizj_1998@jlu.edu.cn (Zhanjun Li);

lai_liangxue@gibh.ac.cn (Liangxue lai).

Abstract

5-Methylcytosine (m^{5}C) is one of the major post-transcriptional modifications in mRNA and is highly involved in the pathogenesis of various diseases. However, the capacity of existing assays for accurately and comprehensively transcriptome-wide m^{5}C mapping still needs improvement. Here, we develop a detection method named DRAM (deaminase and reader protein assisted RNA methylation analysis), in which deaminases (APOBEC1 and TadA-8e) are fused with m^{5}C reader proteins (ALYREF and YBX1) to identify the m^{5}C sites through deamination events neighboring the methylation sites. This antibody-free and bisulfite-free approach provides transcriptome-wide editing regions which are highly overlapped with the publicly available BS-seq datasets and allows for a more stable and comprehensive identification of the m^{5}C loci. In addition, DRAM system even supports ultra-low input RNA (10ng). We anticipate that the DRAM system could pave the way for uncovering further biological functions of m^{5}C modifications.

Keywords: RNA, 5-methylcytosine, deaminase, m^{5}C reader

Introduction

Epigenetics refers to stable inheritance without changing the basic sequence of DNA, involving various forms such as DNA methylation, histone modification and RNA modification. In recent years, RNA sequencing technology has boosted research on RNA epigenetics. More than 170 RNA modifications have been identified, mainly including m^6A, m^5C, m^1A, m^7G and others [1,2]. Notably, RNA m^5C methylation represents a crucial post-transcriptional modification observed across different RNA types, such as tRNA, mRNA, rRNA, vault RNA, microRNA, long non-coding RNA and enhancer RNA [3-8]. Numerous studies have revealed multiple molecular functions of m^5C in numerous key stages of RNA metabolism, such as mRNA stability, translation, and nuclear export [5,9-13]. The dynamic alterations of m^5C play integral roles in many physiological and pathological processes, such as early embryonic development [14], neurodevelopmental disorder [15,16] and multifarious tumorigenesis and migration [17-20]. Moreover, this modification significantly contributes to the regulation of gene expression [5,9-13,17]. Therefore, the detection of m^5C sites appears to be essential for understanding their underlying effects on cellular function and disease states.

With the recent advances in sequencing techniques, several high-throughput assays have been developed for qualitative or quantitative analysis of m^5C. To date, bisulfite-sequencing (BS-seq) has been proven to be the gold standard method for RNA m^5C methylation analysis [5,21,22]. This approach chemically deaminates unmethylated cytosine to uracil, while keeping methylated cytosine unchanged. The m^5C methylation sites can be identified by subsequent library construction and sequencing. However, bisulfite treatment of BS-seq is extremely detrimental to RNA, thus resulting in unstable detection of m^5C in low abundance RNA or highly structured RNA, which directly affects the confidence of results [23,24]. Another major type of global m^5C analysis depends on antibody-assisted immunoprecipitation of m^5C methylated RNAs, such as m^5C-RIP-seq [25-27], AZA-IP-seq [28] or miCLIP-seq [7]. These methods are unable to recognize methylation on mRNAs with low abundance and secondary structure. Moreover, these methods are highly dependent on antibody specificity, which usually leads to unspecific binding of RNA and a low amount of m^5C-modified regions. Moreover, TAWO-seq, originally developed for the identification of hm^5C, is also capable of m^5C analysis, but it highly depends on the oxidation efficiency of perovskite, which usually causes false positives and unstable conversion [29,30]. Furthermore, the emerging third-generation sequencing, such as Nanopore-seq, can directly map m^5C by tracking the characteristic changes of bases, but it still faces challenges of a high error rate [31-33]. These together largely hamper its wide application on transcriptome profiling of m^5C (Supplementary Table 1). Hence, there is an urgent need for a simple, efficient, sensitive, and antibody-independent method for global m^5C detection.

The RNA-binding protein ALYREF is the initially recognized nuclear m^5C reader that binds directly to m^5C sites in mRNA and plays key roles in promoting mRNA nuclear export or tumor progression [5]. Another well-known m^5C reader, YBX1, can also recognize m^5C-modified mRNA through its cold-shock domain and participates in a variety of RNA-dependent events such as mRNA packaging, mRNA stabilization and translational regulation [9,18]. RNA affinity chromatography and mass spectrometry analyses using biotin-labelled oligonucleotides with or without m^5C were performed in previous reports, which indicated that ALYREF and YBX1 had a more prominent binding ability to m^5C-modified oligonucleotides [5,18]. YBX1 can preferentially

recognize mRNAs with m[5]C modifications via key amino acids W65-N70 (WFNVRN)[18], while

K171 is essential for the specific binding of ALYREF to m[5]C sites [5]. Previous studies have shown

that mutations in key amino acids responsible for recognising m[5]C binding in ALYREF and YBX1

lead to a significant reduction in their binding levels to m[5]C-containing oligonucleotides[5,18].

Nucleic acid deaminases, primarily categorized as cytosine deaminases and adenine deaminases,

are zinc-dependent enzymes which facilitate the deamination of cytosine or adenine within DNA

or RNA substrates[34]. APOBEC1, an evolutionarily conserved family member of APOBEC

proteins, can specifically catalyze the deamination of cytosine in single-stranded RNA (ssRNA) or

DNA (ssDNA) to uracil[35-37]. TadA8e is an adenine deaminase optimized through re-engineering of

TadA and it induces conversion of adenine to inosine (eventually read as guanine by transcriptases)

in ssRNA or ssDNA[38,39]. APOBEC1 and TadA8e, with their prominent deamination efficiency,

have been employed for the development of precise and efficient base editors such as CBE and

ABE8e, which find widespread application in studies related to genome editing[37,38].

Here we aim to establish a deaminase and m[5]C reader-assisted RNA methylation sequencing

approach (DRAM-seq), which identifies the m[5]C sites through reader-mediated recognitions and

deaminase-mediated point mutations neighboring the m[5]C methylation sites. This bisulfite-free

and antibody-free method is anticipated to provide more comprehensive and cost-effective

transcriptome-wide detection of m[5]C methylation, which may better assist on exploring its further

regulatory mechanisms.

Results

Development of DRAM system for m[5]C detection

Our sequencing platform is inspired by the concept of the m[6]A DART-seq assay, in which C

near the m[6]A site is converted into U without affecting sequences near non-m6A sites[40]. Therefore,

we hypothesized that, by utilizing the targeted binding of m[5]C readers, deaminase can be recruited

to achieve deamination of cytosine or adenine in the vicinity of the m[5]C sites on single-stranded

RNA, thereby facilitating the detection of the m[5]C site. This approach was named DRAM

(deaminase and m[5]C reader-assisted RNA methylation sequencing). As RNA-binding proteins,

ALYREF and YBX1 also could bind to RNAs without m[5]C modification[5,18]. To exclude the

false-positive detection of DRAM due to the non-m[5]C specific binding of ALYREF and YBX1,

knockout of W65-N70 (WFNVRN) amino acids in YBX1and K171A mutation in ALYREF were

introduced seperately, resulting in the DRAM[mut] system (Fig. S1A-S1D). Subsequently, we

verified the affinity ability of YBX1 and ALYREF for m[5]C-modified RNAs by RNA pull-down

experiments. Consistent with previous reports[5,18], those two m[5]C readers preferentially bound

RNAs containing m[5]C modifications. Furthermore, mutating key amino acids involved in their

interaction with m[5]C significantly reduced their binding ability, indicating that ALYREF and

YBX1 exhibit specificity for m[5]C-methylated mRNAs. (Fig. S1E-S1H). To confirm the

recognition of m[5]C site by DRAM system, DRAM, DRAM[mut] and Deaminase system were

transfected into the human HEK293T cells, respectively. Finally, we considered the presence of

m[5]C modification in the vicinity only if the deamination changes produced under DRAM

induction were significantly different from those produced under DRAM[mut] or Deaminase

induction (Fig. 1A).

Previous studies have indicated that there is no uniform intrinsic signature motif sequence
that can characterize all m5C sites[5,26,41,42]. To comprehensively detect the m5C loci, the readers of
m5C (ALYREF and YBX1) were separately fused to the C-terminus of the deaminases
(APOBEC1 and TadA-8e), namely DRAM-ABE and DRAM-CBE system (Fig.1B).
126

DRAM detection system is assayed in an m5C-dependent form

To confirm the recognition of m5C site by DRAM system, DRAM, DRAM^mut and Deaminase
were transfected into the human HEK293T cells, respectively. To evaluate candidate DRAM
constructs within a cellular environment, we performed fluorescence microscopy to analyze the
expression of DRAM. The results showed that DRAM-ABE and DRAM-CBE were properly
expressed in HEK293T cells (Fig. S2A-S2B). In addition, flow cytometry displayed ~60% of
cells were GFP-positive (Fig. S2C). Two previously reported m5C sites in RPSA and AP5Z1 were
selected for the analysis [5,21], and their methylation status was verified by bisulfite sequencing PCR.
The deep sequencing results showed that the m□C fraction of RPSA and SZRD1 was 75.5% and
27.25%, respectively (Fig.2A and B). Sanger sequencing following RT-PCR was then performed
to determine the editing of neighbouring m5C sites by DRAM system in these two mRNA.
Notably, adenine close to the m5C site in RPSA mRNA was mutated into guanine, resulting in an
A-to-G editing rate of 14.7% by DRAM-ABE, whereas this was rarely observed with TadA-8e or
DRAM^mut-ABE (Fig.2C). DRAM-CBE induced C to U editing in the vicinity of the m5C site in
AP5Z1 mRNA, with 13.6% C-to-U editing, while this effect was significantly reduced with
APOBEC1 or DRAM^mut-CBE (Fig.2D). Subsequently, in order to investigate whether the DRAM
system can detect other types of RNA, such as tRNA, 28S rRNA, or others, we performed PCR
amplification of the flanking sequences of the m5C sites 3782 and 4447 on 28S rRNA and several
m5C sites on tRNA, such as the m5C48 and m5C49 sites of tRNA^Val, the m5C48 and m5C49 sites
of tRNA^Asp, and the m5C48 site of tRNA^Lys. But Sanger sequencing showed that there was no
valid A-to-G/C-to-U mutation detected, which is most likely due to the fact that ALYTEF and
YBX1 are mainly responsible for the mRNA m5C binding proteins, and thus the DRAM system is
more suitable for the mRNA m5C detection (Fig. S3). Taken together, the fusion of m5C reader
and deaminase can effectively and selectively deaminate cytosine/adenine in the vicinity of the
mRNA m5C sites.
NSUN2[43] and NSUN6[44], two family members of NOL1/NSUN protein, were both identified
as m5C methyltransferase of mRNA[45]. To verify that the detection of DRAM occurs in the
presence of m5C, we individually depleted NSUN2 and NSUN6 in HEK293T cells and performed
DRAM transfection. The knockout efficiency has been confirmed by western blotting (Fig.2E, 2F
and Fig. S4A,4B). It has been previously demonstrated that m5C methylation of AP5Z1 and RPSA
is catalyzed by NSUN2 and NSUN6, respectively[21,46]. In line with this, sanger sequencing
following RT-PCT showed a significant reduction in C-to-U or A-to-G mutations near the m5C
sites in methyltransferase-deficient cells compared with WT cells (Fig. 2G and H). Overall, these
findings suggest that the DRAM detection system is assayed in an m5C-dependent form.
161

DRAM enables transcriptome-wide analysis of m5C methylation

Subsequently, we performed RNA-seq analysis after DRAM transfection by detecting
C-to-U/A-to-G editing events to accomplish transcriptome-wide detection of m5C (Fig.3A). To
serve as positive controls, two previously published BS-seq datasets were also integrated[5,21].

Mutations were detected near the m^{5}C site in RPSA as A-to-G by DRAM-ABE (Fig.3B), and
DRAM-CBE detected the presence of C-to-U mutations near the AP5Z1 m^{5}C site (Fig.3C).
However, the DRAM^{mut} and Deaminase systems induced few effective mutations close to these
sites. Examination of multiple reported high-confidence RNA m^{5}C sites showed that DRAM-seq
editing events were also enriched in the vicinity of the BS-seq sites (Fig.3B, 3C and Fig. S5).
DRAM-seq analysis further confirmed that DRAM was detected in an m^{5}C-dependent
manner and that mutations in AP5Z1 and RPSA mRNA were reduced in methyltransferase
knockout cells compared to wild-type cells (Fig. 3D, 3E). Moreover, the knockout cells exhibited
overall rare DRAM-seq editing events close to m^{5}C sites in other mRNAs (Fig. S6). Unfortunately
however due to the inability of the DRAM detection system to achieve m^{5}C detection with single
base resolution, we did not find sequence preferences for the m^{5}C sites responsible for catalysis by
each of the two methyltransferases NSUN2 and NSUN6(Fig. S4D).
A comparison of three biological replicates from each experimental group revealed a strong
reproducibility of A-to-G/C-to-U mutations in HEK293T cells expressing DRAM-ABE and
DRAM-CBE (Fig. S7). Moreover, the integration and analysis of the experimental data revealed a
high degree of overlap between the three biological replicates (Fig. S4C). And a recent study by
Wang et al. showed that ALYREF deletion affects the expression of 94 mRNAs[47], and only 55.32%
of these ALYREF-regulated mRNAs can be detected by the DRAM system (Fig. S4E). These
findings suggest that DRAM selectively targets specific RNAs for editing, exhibiting a high
degree of consistency across samples.
To obtain information on a set of high-confidence DRAM-seq data, we filtered the list of
sites transfected with deaminase alone and screened the sequencing results with methyltransferase
depleted, pooled editing events occurring in at least 10% of reads across multiple samples to
obtain a set of high-confidence editing sites (Fig. 3F and Supplementary Table 2), and integrated
genes with editing sites occurring in DRAM-ABE and DRAM-CBE (Fig. 3F and Supplementary
Table 3).
Previous studies have indicated that m^{5}C sites are predominantly distributed in the coding
sequences (CDS) and notably enriched near the initiation codon[5,25,26,48-50]. To further delineate the
characteristics of the DRAM-seq data, we compared the distribution of DRAM-seq editing sites
within the gene structure, specifically examining their occurrences in the 5'untranslated region
(5'UTR), 3' untranslated region (3'UTR), CDS and Intergenic/Intron region. Our analysis
revealed that DRAM-seq editing events in cells expressing DRAM-ABE and DRAM-CBE were
primarily located in the CDS and 3'UTR, indicating a non-random distribution of m^{5}C (Fig.3G,
Fig. S8A and 8B). Moreover, plotting the distribution of DRAM-seq editing sites in mRNA
segments (5'UTR, CDS, and 3'UTR) highlighted a significant enrichment in the CDS (Fig.3H). In
contrast, cells expressing the deaminase exhibited a distinct distribution pattern of editing sites,
characterized by a prevalence throughout the 3'UTR (Fig.3H). This finding reaffirms that the
specific editing pattern observed in DRAM-seq across the transcriptome depends on its capacity to
bind m^{5}C.
Comparative analysis of the DRAM-seq editing sites with the previously published BS-seq
m^{5}C sites indicated that the likelihood of editing was notably higher in closer proximity to the
m^{5}C sites (Fig.3I). Furthermore, the editing window of DRAM exhibited enrichment
approximately 20bp before and after the m^{5}C site (Fig.3I). Investigation into the sequences
surrounding the editing window revealed that AC motifs were the most significantly enriched in

DRAM-CBE, whereas (U/C) A motifs were most notably enriched in DRAM-ABE. In contrast,

the APOBEC1 and TadA-8e samples displayed no significantly enriched motifs, with mutations

being more randomly orientated (Fig.3J, 3K).

DRAM-seq provides stable and comprehensive identification of m^{5}C loci

Subsequently, we then evaluated the ability of DRAM-seq to detect m^{5}C across the entire

transcriptome and compared its performance to that of the previously reported BS-seq. Although

both previous studies employed bisulfite treatment, the resulting data obtained significant

discrepancies due to variations in their treatment and analysis methodologies. We first complied

the overall distribution of mutant regions identified by DRAM-seq, presenting both the mutant

sites detected by the DRAM system and those reported in previous studies[5,18] across each

chromosome (Fig.4A). Our results indicated that DRAM-seq identified the presence of m^{5}C

modifications covering 79.6% of the genes detected by Yang et al.[5] and 91.9% of the genes

detected by Zhang et al.[21] (Fig.4B and D). Remarkably, certain pivotal regulators with diverse

biological functions, such as ATG16L1(coordinats autophagy pathway)[51] and ARHGEF25 (plays

an important role in actin cytoskeleton reorganisation)[52], were identified by Zhang et al. and

DRAM-seq, but not by Yang et al. (Fig.4C). Conversely, FANCD2 (Maintains chromosome

stability)[53] and RPL15(components of the large ribosomal subunit)[54,55], were discovered by Yang

et al. and DRAM-seq, but not by Zhang et al. (Fig.4E). Hence, DRAM-seq appears to offer a more

stable and comprehensive identification of the m^{5}C loci.

To provide functional insights into m^{5}C RNA-modified genes in HEK293T cells, we

conducted Gene Ontology (GO) and Kyoto Encyclopedia of Genes and Genomes (KEGG)

analyses. These results highlighted the involvement of these genes in the regulation of diverse key

biological processes, such as cell division, cell cycle, mRNA splicing, protein processing in the

endoplasmic reticulum, nucleocytoplasmic transport, translation, DNA repair and others (Fig.4F,

4G, Fig. S8C and S8D).

DRAM enables low-input m^{5}C profiling

A significant challenge in m^{5}C detection lies in the specificity of antibodies and the

substantial amount of input RNA required for sequencing. RNA is susceptible to degradation

during denaturation, sodium bisulfite treatment and desulfurization steps in the BS-seq assay[56].

Immunoprecipitation-based m^{5}C assays and LC-MS/MS also impose high demand for sample

input[7,25,57]. Several experiments have highlighted the requirement of 100-500 ng of RNA for

m^{5}C-RIP-seq, while BS-seq necessitates an even more demanding 750-1000 ng of RNA[21,25,58]. To

assess the detection limits of DRAM-Sanger, we attempted to amplify two representative

m^{5}C-containing sites in the RPSA and AP5Z1 transcripts from diluted RNA samples.

Remarkably, we successfully generated PCR products of these two mRNAs from cDNAs

corresponding to 250 ng, 50 ng, and 10 ng of total RNA. Quantitative analysis by Sanger

sequencing demonstrated nearly identical Sanger traces across these dilutions (Fig.5A and B). This

finding underscores that the specificity of DRAM editing depended on its ability to bind m^{5}C, and

DRAM is proficient in low-input m^{5}C analyses. Furthermore, cell viability was determined by

CCK8 assay on HEK293T cells transfected with DRAM (Fig.5C). Importantly, there was no

significant difference in the relative proliferative capacity of the cells compared to untransfected

cells (NC), indicating that DRAM expression did not adversely affect cell viability (Fig.5D).

Transfection of the DRAM system in cells results in the transient overexpression of fusion
proteins. To investigate how varying expression levels of these proteins influence A-to-G and
C-to-U editing within the same m^{5}C region, we conducted a gradient transfection using plasmid
concentrations of 1500 ng, 1000 ng and 500 ng. This approach allowed us to progressively reduce
the expression levels of the fusion proteins (Fig. 5E and 5F). Sanger sequencing revealed that the
editing efficiency of A-to-G and C-to-U within the m^{5}C region significantly decreased as fusion
protein expression diminished (Fig. 5G and 5H). These findings suggest that the transfection
efficiency of the DRAM system is concentration-dependent and that the ratio of editing efficiency
to transfection efficiency may assist in the quantitative analysis of m^{5}C using the DRAM system.
263
264

Discussion

In recent years, m^{5}C methylation modifications have received increasing attention, with
multiple reports detailing the distribution of RNA m^{5}C methylation modifications across various
species and tissues, elucidating their characteristics. Despite the relatively low abundance of m^{5}C,
its highly dynamic changes hold significant implications for the regulation of physiological and
pathological processes[5,21,44]. However, due to the limitations of sequencing methods and the
variability of data processing, there remains ample room for progress in the study of m^{5}C
detection methods.
In this study, we developed a site-specific, depth-sequencing-free m^{5}C detection method
using DRAM-Sanger. This workflow relies on conventional molecular biology assays such as
RT-PCR and Sanger sequencing, eliminating the need for specialized techniques and thereby
simplifying the process of m^{5}C detection.
DRAM-seq introduces a novel strategy for transcriptome-wide m^{5}C detection, overcoming
inherent limitations in existing methods. Notably, DRAM-seq covered around 80% of the
high-confidence m^{5}C-modified genes detected by BS-seq and identified more potential m^{5}C sites.
This can be attributed to the avoidance of bisulfite treatment by DRAM-seq, preventing RNA
damage and ensuring a more comprehensive representation of RNA samples. This feature also
likely contributes to the observed stability of DRAM-seq in comparison to BS-seq. Additionally,
DRAM-seq is not limited by antibody specificity and is resistant to chemical-induced damage.
A prominent challenge in existing m^{5}C profiling methods is their reliance on substantial
amounts of input RNA samples. In contrast, DRAM operates through the deamination activity of
deamination activity of deaminase, preserving RNA integrity and preventing degradation. The
notable advantage of DRAM lies in its capacity for low-input m^{5}C detection. Our analysis
demonstrates that DRAM requires as low as 10ng of total RNA for m^{5}C detection. While DRAM
is currently well-suited for detecting m^{5}C on a transcriptome-wide scale, the potential for future
applications involving third-generation sequencing could extend its utility to individual mRNAs,
particularly m^{5}C heterogeneity on mRNA splicing variants. In addition, the DRAM system
depends on the specific recognition of m^{5}C modifications on ssRNA by the reader protein,
theoretically avoiding the false-positive effects of 5-hydroxymethylation modifications in other
assays, such as BS-seq[21-23]. This potential feature could enhance the accuracy of the DRAM assay,
albeit it still requires careful validation.
In our study, m^{5}C detection was performed following the transient transfection of the DRAM

detection system into mammalian cells, which might result in a lower mutation rate at the corresponding site. Therefore, employing lentiviral-mediated transfection into cell lines of interest could potentially enhance the efficiency of m5C detection. Our results confirm that YBX1 and ALYREF exhibit specificity as m5C readers, binding preferentially to RNAs with m5C modifications, thereby validating the reliability of the DRAM detection system. However, mutations in the key amino acids responsible for m5C binding reduced their affinity while retaining some binding capacity. Further elucidation of the key amino acids directing ALYREF and YBX1's binding to m5C methylation sites should enable more accurate and sensitive m5C detection by DRAM-seq. Due to the lack of a fixed base composition for characterizing all m5C modification sites, DRAM has an apparent limitation in achieving single-base resolution for detecting m5C. However, our present study proved that the measuring resolution of DRAM is around 40nt, which facilitates higher precision than that of m5C-RIP-seq (~100nt). In the future, with more in-depth analyses of m5C reader structures and the identification of new potential m5C readers, we expect to achieve more precise m5C localization and more comprehensive m5C modification detection. Moreover, the substitution of deaminases, such as A3A and A3G (the family members of APOBEC), could also potentially enhance the efficiency of the DRAM detection [59-61].

Although the m5C assay can be performed using the DRAM system alone, comparing it with the DRAM^mut and deaminase controls could enhance the accuracy of m5C detection in specific regions. Given that the expression of DRAM fusion proteins significantly influences m5C detection, it is advisable to transfect the same batch of cells during the assay to ensure consistent transfection efficiency across experimental groups and thus can better standardize the detection.

One future direction of endeavour is the purification of DRAM fusion proteins to facilitate *in vitro* detection of RNA m5C methylation, which could extend the scope of DRAM-seq to diverse sample types. Another potential application for DRAM-seq could be the expression of drug-inducible DRAM systems *in vivo* using various animal models for m5C analysis. These will together provide novel insights into m5C modifications for biological and clinical research.

Conclusions

In summary, we developed a novel deaminase and reader protein-assisted RNA m5C methylation approach that detects the m5C region by deaminating As or Cs in close proximity to the m5C sites, which does not rely on antibodies or bisulfite, thus leading to unprecedently comprehensive transcriptome-wide RNA m5C methylation profiling. We anticipated that this system could pave the way for uncovering further biological functions of m5C modifications and facilitate the development of therapeutic interventions for associated diseases.

Materials and methods

Plasmid construction

ALYREF and YBX1 expression plasmids were purchased from MIAOLING BIOLOGY (http://www.miaolingbio.com/), and the ALYREF and YBX1 fractions were then amplified using specific primer. The ALYREF and YBX1 portions were amplified using pCMV-APOBEC1-YTH (Addgene plasmid no. 131636; https://www.addgene.org/131636/) and ABE8e (Addgene plasmid

no. 138489; https://www.addgene.org/ 138489/) to amplify the deaminase portion and the essential

plasmid construct proxies, and finally the fragments were recombined by the ClonExpress Ultra

One Step Cloning Kit to complete the plasmid vector construction. Both DRAM[mut]-ABE and

DRAM[mut]-CBE related vectors were obtained by introducing the corresponding key amino acid

mutations using Fast Site-Directed Mutagenesis Kit (TIANGEN Biotech). The primer sequences

used are listed in Supplementary Table 4.

Cell culture and plasmid transfection

HEK293T cell line (ATCC) was cultured in Dulbecco's Modified Eagle Medium (DMEM)

supplemented with 10% fetal bovine serum (CLARK BIOSCIENCE) and 1% penicillin (100

U/ml)-streptomycin (100μg/ml). The cells were seeded in 12-well plates and transfected using

Hieff Trans™ Liposomal Transfection Reagent (Yeasen).

NSUN2-depleted cell lines were generated by cloning NSUN2-targeting single guide RNA

sequences into the pSpCas9(BB)–2A-Puro (PX459) V2.0 plasmid (Addgene plasmid no. 62988;

http://n2t.net/addgene:62988). Plasmids were then transfected into HEK293T cells and Puromycin

(Meilunbio) was added at a final concentration of 3 μg/ml to enrich the positively transfected cells

24 h after transfection. After 72 h, the cells were collected and used for genotyping by Sanger

sequencing. NSUN6-depleted cell lines were generated in the same way. The primers used for

genotyping and single guide RNA sequences are listed in Supplemental Table 4.

Cell viability measurements

HEK293T cells were transfected with DRAM plasmid and cultured at 37°C for 24 h.

Subsequently, 1000 cells were seeded in 96-well plates. After waiting for the cells to attach to the

wall, the cell activity was detected by Cell Counting Kit-8 (Meilunbio). Cell Counting Kit-8

contains WST-8, which in the presence of the electronically coupled reagent 1-Methoxy PMS can

be reduced by mitochondrial dehydrogenase to the orange-colored metazan product Formazan, the

absorbance of which is measured at 450 nm to analyze cellular activity.

Western blotting

For protein blotting, samples were lysed in RIPA Lysis Buffer (Meilunbio) with

Phenylmethanesulfonyl fluoride (PMSF) and the BCA protein assay kit (Beyotime Biotechnology)

was used to Protein concentration was measured. Total protein extracts were separated by

SDS-PAGE on a 10% gel and then transferred to 0.22 nm polyvinylidene fluoride membranes

(Boster). Subsequently, the proteins were probed with specific antibodies after the blot was

blocked with 5% non-fat milk (Boster). Images were quantified using ImageJ software and all data

are expressed as mean ± SEM.

The following antibodies and concentrations were used: NSUN2 Polyclonal antibody

(Proteintech; Cat No.20854-1-AP; 1:7500), NSUN6 Polyclonal antibody (Proteintech; Cat No.

17240-1-AP; 1:2000), RabbitAnti-GAPDH antibody (Bioss; bs-41373R; 1:2000), Alpha Tubulin

Polyclonal antibody (Proteintech; Cat No. 11224-1-AP; 1:2000), HRP-labeled Goat Anti-Rabbit

IgG(H+L) (Beyotime Biotechnology; A0208; 1:2000).

cDNA synthesis and Sanger sequencing

Total cellular RNA was extracted with TRIzol reagent (TIANGEN Biotech) and cDNA was

synthesized using PrimeScript™ II 1st Strand cDNA Synthesis Kit (Takara Bio) according to the

manufacturer's recommendations. PCR was then performed using 2×Taq PCR MasterMix II

(TIANGEN Biotech) and primers flanking m[5]C target sites, and the purified PCR products were

directly sequenced by Sanger sequencing. The Sanger sequencing results were analyzed using

EditR 1.0.10 to calculate the mutation frequency[62]. The primers used in this study are shown in

Supplemental Table 4.

Real-time quantitative PCR

cDNA was synthesized using FastKing RT kit (with gDNase) (TIANGEN Biotech) according

to the manufacturer's recommendations. RT-qPCR assay was performed using SuperReal PreMix

Plus (SYBR Green) (TIANGEN Biotech). GAPDH was used as an endogenous control, and the

expression levels were normalized to the control and calculated by the $2^{-\Delta\Delta Ct}$ formula. All samples

were analyzed in triplicate and each mRNA quantification represents the average of at least three

measurements. All data are expressed as mean ± SEM. The primers used in this study are shown in

Supplemental Table 4.

Protein structure modelling

Protein structure simulations were performed using the SWISS-MODEL online website

(https://swissmodel.expasy.org/interactive)[63]. The SWISS-MODEL database is able to provide

up-to-date annotated 3D protein models, which are generated from automated homology

modelling of related model organisms and experimental structural information for all sequences in

UniProtKB, with reliable structural information, and subsequently protein structure observations

were performed using PyMOL[64].

Bisulfite sequencing PCR

We referenced bisulfite sequencing PCR, an assay established by Matthias Schaefer et al. We

chemically deaminated cytosine in RNA using the EZ RNA methylation kit (50) (ZYMO

RESEARCH) and then quantified m^5C methylation levels based on PCR amplification of cDNA

combined with deep sequencing[23].

RNA Conversion Reagent was premixed with prepared RNA samples, and the RNA was

denatured at 70°C for 5 minutes, followed by a reaction period of 45 minutes at 54°C. Finally, the

purified RNA samples were recovered after desulfurization by RNA Desulphonation Buffe. cDNA

was synthesized using PrimeScript™ II 1st Strand cDNA Synthesis Kit (Takara Bio) according to

the manufacturer's recommendations. PCR was then performed using 2× EpiArt HS Taq Master

Mix (Dye Plus) (Vazyme) and m^5C target site-specific Bisulfite Primer (primer sequences were

designed at https://zymoresearch.eu/pages/bisulfite-primer- seeker), the products were purified by

TIANgel Midi Purification Kit (TIANGEN Biotech), and the connectors for second-generation

sequencing were attached at both ends of the products for sequencing. Finally, deep sequencing

was performed by HiTOM analysis to detect the methylation level (The number of reads >1000 in

deep sequencing) [65]. The primers used in this study are shown in Supplemental Table 4.

Library construction and next-generation sequencing

1μg of total cellular RNA was used for sequencing library generation by NEBNext Ultra

RNA Library Prep Kit for Illumina (NEB, USA, Catalog #: E7530L) following manufacturer's

recommendations and index codes were added to attribute sequences to each sample. Briefly,

mRNA was purified from total RNA using poly-T oligo-attached magnetic beads. Fragmentation

was carried out using divalent cations under elevated temperature in NEB Next First Strand

Synthesis Reaction Buffer(5X). First-strand cDNA was synthesized using random hexamer primer

and M-MuLV Reverse Transcriptase (RNase H). Second-strand cDNA synthesis was subsequently

performed using DNA Polymerase I and RNase H. Remaining overhangs was converted into blunt

ends via exonuclease/polymerase activities. After adenylation of 3' ends of DNA fragments, NEB

Next Adaptor with hairpin loop structure was ligated to prepare for hybridization. To select cDNA
fragments of preferentially 370~420 bp in length, the library fragments were purified with
AMPure XP system (Beverly, USA). Then 3 μL USER Enzyme (NEB, USA) was used with size
selected, adaptor-ligated cDNA at 37°C for 15 min followed by 5 min at 95 °C before PCR. Then
PCR was performed with Phusion High-Fidelity DNA polymerase, Universal PCR primers and
Index (X) Primer. At last, PCR products were purified (AMPure XP system) and library quality
was assessed on the Agilent 5400 system(Agilent, USA)and quantified by QPCR (library

concentration ≥ 1.5 nM). The qualified libraries were pooled and sequenced on Illumina

platforms with PE150 strategy in Novogene Bioinformatics Technology Co., Ltd (Beijing, China),
according to effective library concentration and data amount required.

DRAM-seq analysis and calling of edited sites
The raw fastq sequencing data were cleaned by trimming the adapter sequences using Fastp
(v0.23.1) and were aligned to the human genome (hg19) using STAR (v2.7.7) in paired-end mode.
The aligned BAM files were sorted and PCR duplicates were removed using Samtools (v1.12).
The cite calling pf DRAM-seq was performed using Bullseye, a previously customized pipeline to
look for C-to-U or A-to-G edited sites throughout the transcriptome[40]. Briefly, the sorted and
deduplicated BAM files were initially parsed by parseBAM.pl script.

Then, Find_edit_site.pl script was employed to find C-to-U or A-to-G editing events by
DRAM-seq with at least 10 reads of coverage, an edit ratio of 5%-95%, an edit ratio at least
1.5-fold higher than NSUN2 or NSUN6-knockout samples, and at least 2 editing events at a given
site. Sites that were only found in one replicate of each DRAM protein variant were removed.
Editing events appeared in cells expressing merely APOBEC1 or TadA8e were also removed. For
high confidence filtering, we further adjusted the Find_edit_site.pl parameters to the edit ratio of
10%-60%, an edit ratio of control samples at least 2-fold higher than NSUN2 or NSUN6-knockout
samples, and at least 4 editing events at a given site.

Metagene and motif analyses
Metagene analysis was performed using hg19 annotations according to previously reported
tool, MetaplotR[66]. For motif analysis, the 20bp flanking sequence of each DRAM-seq editing site
was extracted by Bedtools (v2.30.0)[67]. The motif logos were then plotted by WebLogo (v3.7.12)[68].

Replicates analysis
Independent biological replicates of DRAM-ABE or DRAM-CBE in DRAM-seq analysis
were separately compared by computing the Pearson correlation coefficient between the number
of C-to-U mutations per mRNA between any two replicate experiments.

GO and KEGG analysis
GO and KEGG analysis of DRAM-seq edited mRNAs was performed using the DAVID
bioinformatic database [69]. GO terms with a P value of less than 0.05 were considered statistically
significant.

RNA pulldown assay
RNAs containing C (5′-biotin-GAGGUAUGAACUGUAAGUU-3′,
5′-biotin-GAAAGGAGAUCGCCAUUAUCC-3′) or m^{5}C residues
(5′-biotin-GAGGUAUGAA/i5Me-rC/UGUAAGUU-3′,
5′-biotin-GAAAGGAGAU/i5Me-rC/GCCAUUAUCC-3′) were prepared in advance. Protein
lysates were isolated from HEK293T cells transfected with YBX1, YBX1^{mut}, ALYREF or

ALYREF^{mut} for 24 h using lysis buffer. RNA pull-down assays were performed with the
Pierce™ Magnetic RNA-Protein Pull-Down Kit (Thermo) following the manufacturer's
instructions, and the results were finally analyzed by Western blotting.

Statistical analysis

All data are expressed as mean ± S.E.M of three independent determinations. Data were
analyzed through a two-tailed t-test. A probability of $P < 0.05$ was considered statistically
significant; □, $P < 0.05$, □□, $P < 0.01$, *, $P < 0.05$, **, $P < 0.01$, ***, $P < 0.001$ and ****, $P <$
0.0001 denote the significance thresholds; ns denotes not significant.

Data and Materials Availability

478 The data supporting the findings of this study are available within the article and its
479 Supplementary Information. Other data and reagents are available from the corresponding authors
480 upon reasonable request.

AUTHOR CONTRIBUTIONS

Conceptualization: JZ, YH, LL, ZL
Methodology: JZ, DZ, JL
Investigation: JZ, DZ, JL, DK, XL, RZ, YL
Visualization: XG, YQ, DW, JC
Supervision: DK, XL, RZ, YL, XG, YQ, DW, JC, YH
Funding acquisition: YH, LL, ZL
Data curation: JZ, YH
Writing—original draft: JZ, DZ, JL
Writing—review & editing: JZ, YH, LL, ZL
492

Competing Interests

494 All other authors declare they have no competing interests.
495

ACKNOWLEDGEMENTS

497 We thank Yuning Song, Yuanyuan Xu and Tingting Sui for critical feedback on the work and
498 manuscript.

FUNDING

500 This work was supported by the National Key Research and Development Program of China
501 (2023YFF1000204) and the National Natural Science Foundation of China (Nos.32200466).

REFERENCES

Wiener, D. & Schwartz, S. The epitranscriptome beyond m(6)A. *Nat Rev Genet* **22**, 119-131 (2021). https://doi.org/10.1038/s41576-020-00295-8

Li, S. & Mason, C. E. The pivotal regulatory landscape of RNA modifications. *Annu Rev Genomics Hum Genet* **15**, 127-150 (2014). https://doi.org/10.1146/annurev-genom-090413-025405

Van Haute, L. *et al.* NSUN2 introduces 5-methylcytosines in mammalian mitochondrial tRNAs. *Nucleic Acids Res* **47**, 8720-8733 (2019). https://doi.org/10.1093/nar/gkz559

Blaze, J. *et al.* Neuronal Nsun2 deficiency produces tRNA epitranscriptomic alterations and proteomic shifts impacting synaptic signaling and behavior. *Nat Commun* **12**, 4913 (2021). https://doi.org/10.1038/s41467-021-24969-x

Yang, X. *et al.* 5-methylcytosine promotes mRNA export - NSUN2 as the methyltransferase and ALYREF as an m(5)C reader. *Cell Res* **27**, 606-625 (2017). https://doi.org/10.1038/cr.2017.55

Sharma, S., Yang, J., Watzinger, P., Kötter, P. & Entian, K. D. Yeast Nop2 and Rcm1 methylate C2870 and C2278 of the 25S rRNA, respectively. *Nucleic Acids Res* **41**, 9062-9076 (2013). https://doi.org/10.1093/nar/gkt679

Hussain, S. *et al.* NSun2-mediated cytosine-5 methylation of vault noncoding RNA determines its processing into regulatory small RNAs. *Cell Rep* **4**, 255-261 (2013). https://doi.org/10.1016/j.celrep.2013.06.029

Aguilo, F. *et al.* Deposition of 5-Methylcytosine on Enhancer RNAs Enables the Coactivator Function of PGC-1α. *Cell Rep* **14**, 479-492 (2016). https://doi.org/10.1016/j.celrep.2015.12.043

Yang, Y. *et al.* RNA 5-Methylcytosine Facilitates the Maternal-to-Zygotic Transition by Preventing Maternal mRNA Decay. *Mol Cell* **75**, 1188-1202.e1111 (2019). https://doi.org/10.1016/j.molcel.2019.06.033

García-Vílchez, R., Sevilla, A. & Blanco, S. Post-transcriptional regulation by cytosine-5 methylation of RNA. *Biochim Biophys Acta Gene Regul Mech* **1862**, 240-252 (2019). https://doi.org/10.1016/j.bbagrm.2018.12.003

Trixl, L. & Lusser, A. The dynamic RNA modification 5-methylcytosine and its emerging role as an epitranscriptomic mark. *Wiley Interdiscip Rev RNA* **10**, e1510 (2019). https://doi.org/10.1002/wrna.1510

Boo, S. H. & Kim, Y. K. The emerging role of RNA modifications in the regulation of mRNA stability. *Exp Mol Med* **52**, 400-408 (2020). https://doi.org/10.1038/s12276-020-0407-z

Courtney, D. G. *et al.* Epitranscriptomic Addition of m(5)C to HIV-1 Transcripts Regulates Viral Gene Expression. *Cell Host Microbe* **26**, 217-227.e216 (2019). https://doi.org/10.1016/j.chom.2019.07.005

Liu, J. *et al.* Developmental mRNA m(5)C landscape and regulatory innovations of massive m(5)C modification of maternal mRNAs in animals. *Nat Commun* **13**, 2484 (2022). https://doi.org/10.1038/s41467-022-30210-0

Chen, P., Zhang, T., Yuan, Z., Shen, B. & Chen, L. Expression of the RNA methyltransferase Nsun5 is essential for developing cerebral cortex. *Mol Brain* **12**, 74 (2019). https://doi.org/10.1186/s13041-019-0496-6

16   Flores, J. V. *et al.* Cytosine-5 RNA Methylation Regulates Neural Stem Cell Differentiation
and Motility. *Stem Cell Reports* **8**, 112-124 (2017).
https://doi.org:10.1016/j.stemcr.2016.11.014

17   Chen, S. Y. *et al.* RNA bisulfite sequencing reveals NSUN2-mediated suppression of epithelial
differentiation in pancreatic cancer. *Oncogene* **41**, 3162-3176 (2022).
https://doi.org:10.1038/s41388-022-02325-7

18   Chen, X. *et al.* 5-methylcytosine promotes pathogenesis of bladder cancer through stabilizing
mRNAs. *Nat Cell Biol* **21**, 978-990 (2019). https://doi.org:10.1038/s41556-019-0361-y

19   Yang, R. *et al.* The RNA methyltransferase NSUN6 suppresses pancreatic cancer development
by regulating cell proliferation. *EBioMedicine* **63**, 103195 (2021).
https://doi.org:10.1016/j.ebiom.2020.103195

20   Li, Y. *et al.* Long noncoding RNA DIAPH2-AS1 promotes neural invasion of gastric cancer
via stabilizing NSUN2 to enhance the m5C modification of NTN1. *Cell Death Dis* **14**, 260
(2023). https://doi.org:10.1038/s41419-023-05781-5

21   Huang, T., Chen, W., Liu, J., Gu, N. & Zhang, R. Genome-wide identification of mRNA
5-methylcytosine in mammals. *Nat Struct Mol Biol* **26**, 380-388 (2019).
https://doi.org:10.1038/s41594-019-0218-x

22   Trixl, L., Rieder, D., Amort, T. & Lusser, A. Bisulfite Sequencing of RNA for
Transcriptome-Wide Detection of 5-Methylcytosine. *Methods Mol Biol* **1870**, 1-21 (2019).
https://doi.org:10.1007/978-1-4939-8808-2_1

23   Schaefer, M., Pollex, T., Hanna, K. & Lyko, F. RNA cytosine methylation analysis by bisulfite
sequencing. *Nucleic Acids Res* **37**, e12 (2009). https://doi.org:10.1093/nar/gkn954

24   Amort, T. & Lusser, A. Detection of 5-Methylcytosine in Specific Poly(A) RNAs by Bisulfite
Sequencing. *Methods Mol Biol* **1562**, 107-121 (2017).
https://doi.org:10.1007/978-1-4939-6807-7_8

25   Cui, X. *et al.* 5-Methylcytosine RNA Methylation in Arabidopsis Thaliana. *Mol Plant* **10**,
1387-1399 (2017). https://doi.org:10.1016/j.molp.2017.09.013

26   Edelheit, S., Schwartz, S., Mumbach, M. R., Wurtzel, O. & Sorek, R. Transcriptome-wide
mapping of 5-methylcytidine RNA modifications in bacteria, archaea, and yeast reveals m5C
within archaeal mRNAs. *PLoS Genet* **9**, e1003602 (2013).
https://doi.org:10.1371/journal.pgen.1003602

27   Saplaoura, E., Perrera, V., Colot, V. & Kragler, F. Methylated RNA Immunoprecipitation
Assay to Study m5C Modification in Arabidopsis. *J Vis Exp* (2020).
https://doi.org:10.3791/61231

28   Khoddami, V. & Cairns, B. R. Identification of direct targets and modified bases of RNA
cytosine methyltransferases. *Nat Biotechnol* **31**, 458-464 (2013).
https://doi.org:10.1038/nbt.2566

29   Yuan, F. *et al.* Bisulfite-free and base-resolution analysis of 5-methylcytidine and
5-hydroxymethylcytidine in RNA with peroxotungstate. *Chem Commun (Camb)* **55**,
2328-2331 (2019). https://doi.org:10.1039/c9cc00274j

30   Hayashi, G. *et al.* Base-Resolution Analysis of 5-Hydroxymethylcytosine by One-Pot
Bisulfite-Free Chemical Conversion with Peroxotungstate. *J Am Chem Soc* **138**, 14178-14181
(2016). https://doi.org:10.1021/jacs.6b06428

31   Harel, N., Meir, M., Gophna, U. & Stern, A. Direct sequencing of RNA with MinION

Nanopore: detecting mutations based on associations. *Nucleic Acids Res* **47**, e148 (2019).
https://doi.org/10.1093/nar/gkz907

32   Chang, J. J. *et al.* Transcriptional and epi-transcriptional dynamics of SARS-CoV-2 during
cellular infection. *Cell Rep* **35**, 109108 (2021). https://doi.org:10.1016/j.celrep.2021.109108

33   Wan, Y. K., Hendra, C., Pratanwanich, P. N. & Göke, J. Beyond sequencing: machine learning
algorithms extract biology hidden in Nanopore signal data. *Trends Genet* **38**, 246-257 (2022).
https://doi.org:10.1016/j.tig.2021.09.001

34   Budzko, L., Hoffa-Sobiech, K., Jackowiak, P. & Figlerowicz, M. Engineered deaminases as a
key component of DNA and RNA editing tools. *Mol Ther Nucleic Acids* **34**, 102062 (2023).
https://doi.org:10.1016/j.omtn.2023.102062

35   Rosenberg, B. R., Hamilton, C. E., Mwangi, M. M., Dewell, S. & Papavasiliou, F. N.
Transcriptome-wide sequencing reveals numerous APOBEC1 mRNA-editing targets in
transcript 3' UTRs. *Nat Struct Mol Biol* **18**, 230-236 (2011).
https://doi.org:10.1038/nsmb.1975

36   Qian, X. *et al.* Low expression of the apolipoprotein B mRNA-editing transgene in mice
reduces LDL levels but does not cause liver dysplasia or tumors. *Arterioscler Thromb Vasc*
Biol **18**, 1013-1020 (1998). https://doi.org:10.1161/01.atv.18.6.1013

37   Siriwardena, S. U., Chen, K. & Bhagwat, A. S. Functions and Malfunctions of Mammalian
DNA-Cytosine Deaminases. *Chem Rev* **116**, 12688-12710 (2016).
https://doi.org:10.1021/acs.chemrev.6b00296

38   Richter, M. F. *et al.* Phage-assisted evolution of an adenine base editor with improved Cas
domain compatibility and activity. *Nat Biotechnol* **38**, 883-891 (2020).
https://doi.org:10.1038/s41587-020-0453-z

39   Gaudelli, N. M. *et al.* Directed evolution of adenine base editors with increased activity and
therapeutic application. *Nat Biotechnol* **38**, 892-900 (2020).
https://doi.org:10.1038/s41587-020-0491-6

40   Meyer, K. D. DART-seq: an antibody-free method for global m(6)A detection. *Nat Methods*
16, 1275-1280 (2019). https://doi.org:10.1038/s41592-019-0570-0

41   Sun, Z. *et al.* Effects of NSUN2 deficiency on the mRNA 5-methylcytosine modification and
gene expression profile in HEK293 cells. *Epigenomics* **11**, 439-453 (2019).
https://doi.org:10.2217/epi-2018-0169

42   Zhang, Q., Zheng, Q., Yu, X., He, Y. & Guo, W. Overview of distinct 5-methylcytosine
profiles of messenger RNA in human hepatocellular carcinoma and paired adjacent non-tumor
tissues. *J Transl Med* **18**, 245 (2020). https://doi.org:10.1186/s12967-020-02417-6

43   Zhang, Q. *et al.* The role of RNA m(5)C modification in cancer metastasis. *Int J Biol Sci* **17**,
3369-3380 (2021). https://doi.org:10.7150/ijbs.61439

44   Liu, J. *et al.* Sequence- and structure-selective mRNA m(5)C methylation by NSUN6 in
animals. *Natl Sci Rev* **8**, nwaa273 (2021). https://doi.org:10.1093/nsr/nwaa273

45   Schumann, U. *et al.* Multiple links between 5-methylcytosine content of mRNA and
translation. *BMC Biol* **18**, 40 (2020). https://doi.org:10.1186/s12915-020-00769-5

46   Fang, L. *et al.* CIGAR-seq, a CRISPR/Cas-based method for unbiased screening of novel
mRNA modification regulators. *Mol Syst Biol* **16**, e10025 (2020).
https://doi.org:10.15252/msb.202010025

47   Wang, N. *et al.* m(5)C-dependent cross-regulation between nuclear reader ALYREF and writer

NSUN2 promotes urothelial bladder cancer malignancy through facilitating RABL6/TK1
mRNAs splicing and stabilization. *Cell Death Dis* **14**, 139 (2023).
https://doi.org/10.1038/s41419-023-05661-y

48 Tang, Y. *et al.* OsNSUN2-Mediated 5-Methylcytosine mRNA Modification Enhances Rice
Adaptation to High Temperature. *Dev Cell* **53**, 272-286.e277 (2020).
https://doi.org/10.1016/j.devcel.2020.03.009

49 Squires, J. E. *et al.* Widespread occurrence of 5-methylcytosine in human coding and
non-coding RNA. *Nucleic Acids Res* **40**, 5023-5033 (2012).
https://doi.org/10.1093/nar/gks144

50 Zhang, P., Wang, Y. & Gu, X. RNA 5-Methylcytosine Controls Plant Development and
Environmental Adaptation. *Trends Plant Sci* **25**, 954-958 (2020).
https://doi.org/10.1016/j.tplants.2020.07.004

51 Fletcher, K. *et al.* The WD40 domain of ATG16L1 is required for its non-canonical role in
lipidation of LC3 at single membranes. *EMBO J* **37** (2018).
https://doi.org/10.15252/embj.201797840

52 Guo, X. *et al.* A Rac/Cdc42-specific exchange factor, GEFT, induces cell proliferation,
transformation, and migration. *J Biol Chem* **278**, 13207-13215 (2003).
https://doi.org/10.1074/jbc.M208896200

53 Naim, V. & Rosselli, F. The FANC pathway and BLM collaborate during mitosis to prevent
micro-nucleation and chromosome abnormalities. *Nat Cell Biol* **11**, 761-768 (2009).
https://doi.org/10.1038/ncb1883

54 Anger, A. M. *et al.* Structures of the human and Drosophila 80S ribosome. *Nature* **497**, 80-85
(2013). https://doi.org/10.1038/nature12104

55 Liang, X. *et al.* Structural snapshots of human pre-60S ribosomal particles before and after
nuclear export. *Nat Commun* **11**, 3542 (2020). https://doi.org/10.1038/s41467-020-17237-x

56 Schaefer, M. RNA 5-Methylcytosine Analysis by Bisulfite Sequencing. *Methods Enzymol* **560**,
297-329 (2015). https://doi.org/10.1016/bs.mie.2015.03.007

57 Bourgeois, G. *et al.* Eukaryotic rRNA Modification by Yeast
5-Methylcytosine-Methyltransferases and Human Proliferation-Associated Antigen p120.
PLoS ONE **10**, e0133321 (2015). https://doi.org/10.1371/journal.pone.0133321

58 Gu, X. & Liang, Z. Transcriptome-Wide Mapping 5-Methylcytosine by m(5)C RNA
Immunoprecipitation Followed by Deep Sequencing in Plant. *Methods Mol Biol* **1933**,
389-394 (2019). https://doi.org/10.1007/978-1-4939-9045-0_24

59 Kim, K., Shi, A. B., Kelley, K. & Chen, X. S. Unraveling the Enzyme-Substrate Properties for
APOBEC3A-Mediated RNA Editing. *J Mol Biol* **435**, 168198 (2023).
https://doi.org/10.1016/j.jmb.2023.168198

60 Barka, A. *et al.* The Base-Editing Enzyme APOBEC3A Catalyzes Cytosine Deamination in
RNA with Low Proficiency and High Selectivity. *ACS Chem Biol* **17**, 629-636 (2022).
https://doi.org/10.1021/acschembio.1c00919

61 Pan, Y. *et al.* Nanoscale Characterization of Interaction of APOBEC3G with RNA.
Biochemistry **56**, 1473-1481 (2017). https://doi.org/10.1021/acs.biochem.6b01189

62 Kluesner, M. G. *et al.* EditR: A Method to Quantify Base Editing from Sanger Sequencing.
Crispr j **1**, 239-250 (2018). https://doi.org/10.1089/crispr.2018.0014

63 Bienert, S. *et al.* The SWISS-MODEL Repository-new features and functionality. *Nucleic*

678   *Acids Res* **45**, D313-d319 (2017). https://doi.org:10.1093/nar/gkw1132

679   64   Delano, W. L. The PyMol Molecular Graphics System. *Proteins Structure Function and*

680   *Bioinformatics* **30**, 442-454 (2002).

681   65   Liu, Q. *et al.* Hi-TOM: a platform for high-throughput tracking of mutations induced by

682   CRISPR/Cas systems. *Sci China Life Sci* **62**, 1-7 (2019).

683   https://doi.org:10.1007/s11427-018-9402-9

684   66   Olarerin-George, A. O. & Jaffrey, S. R. MetaPlotR: a Perl/R pipeline for plotting metagenes of

685   nucleotide modifications and other transcriptomic sites. *Bioinformatics* **33**, 1563-1564 (2017).

686   https://doi.org:10.1093/bioinformatics/btx002

687   67   Quinlan, A. R. & Hall, I. M. BEDTools: a flexible suite of utilities for comparing genomic

688   features. *Bioinformatics* **26**, 841-842 (2010). https://doi.org:10.1093/bioinformatics/btq033

689   68   Crooks, G. E., Hon, G., Chandonia, J. M. & Brenner, S. E. WebLogo: a sequence logo

690   generator. *Genome Res* **14**, 1188-1190 (2004). https://doi.org:10.1101/gr.849004

691   69   Sherman, B. T. *et al.* DAVID: a web server for functional enrichment analysis and functional

692   annotation of gene lists (2021 update). *Nucleic Acids Res* **50**, W216-w221 (2022).

693   https://doi.org:10.1093/nar/gkac194

     FIGURES LEGENDS

697
Fig.1: **Development of DRAM system for m^{5}C detection.**
(A) Schematic diagram of the DRAM assay. DRAM, DRAM^{mut} and Deaminase system were
transfected into HEK293T cells separately. After DRAM transfection, the deaminase was directed
by m^{5}C reader to the vicinity of the m^{5}C site and induce C-to-U/A-to-G mutations, whereas
transfection of the DRAM^{mut} or Deaminase system failed to effectively induce similar mutations
due to the absence of the m^{5}C-recognition-binding domain.
(B) The overall design of DRAM, DRAM^{mut} and Deaminase system.

Fig.2: DRAM detection system was assayed in an m^{5}C -dependent form.

(A, B) Two m^{5}C sites from RPSA (**A**) and AP5Z1 (**B**) mRNA detected by deep sequencing of bisulfite sequencing PCR in HEK293T cells. The m^{5}C sites are highlighted by red color. The m□C fraction of RPSA and AP5Z1 were 75.5% and 27.25% (The number of reads is greater than 1000).

(C, D) Sanger sequencing following RT-PCR verified two m^{5}C sites from RPSA (**C**) and AP5Z1 (**D**) mRNAs in DRAM-transfected HEK293T cells, respectively. HEK293T cells only expressing DRAM^{mut} or Deaminase were served as negative controls. The left panel illustrates the location of DRAM induced mutation sites, which is highlighted in red asterisk. The right panel shows the corresponding quantification of sanger sequencing.

(E, F) The knockout efficiency of NSUN2 (**E**) and NSUN6 (**F**) in HEK293T cell lines verified by Western blotting. The protein level of α-Tubulin and GAPDH were served as loading controls, separately.

719   **(G, H)** DRAM induced mutations close to m^{5}C sites in AP5Z1 (**G**) and RPSA (**H**) mRNAs after

720   NSUN2 and NSUN6 knockout in HEK293T cells. The left panel illustrates the location of DRAM

721   induced mutation sites, which is highlighted in red asterisk. The right panel shows the

722   corresponding quantification of sanger sequencing.

Fig.3: DRAM enables transcriptome-wide analysis of m5C methylation.

(**A**) Schematic of the DRAM-seq method.

(**B, C**) Integrative genomics viewer (IGV) browser traces of DRAM-seq data expressing the indicated constructs in RPSA (**B**, left panel), TARBP2 (**B**, right panel), AP5Z1(**C**, left panel), and TRAF7 (**C**, left panel) mRNAs. C-to-U or A-to-G mutations found in at least 10% of reads are indicated by coloring. The previously published RNA BS-seq datasets from two individual studies were displayed as panel "Yang et al." and "Zhang et al.". (n(DRAM)=3 independent samples, n(Deaminase)=2 independent samples, and n(DRAM[mut])=1 independent sample.)

(**D, E**) Integrative genomics viewer (IGV) browser traces of DRAM-seq data in wildtype and methyltranferases knockout cells in AP5Z1 (**D**) and RPSA (**E**) mRNAs. C-to-U or A-to-G mutations were found in at least 10% of reads are indicated by coloring. The previously published RNA BS-seq datasets from two individual studies were displayed as panel "Yang et al." and "Zhang et al.". n=3 independent samples.

(**F**) Screening process for DRAM-seq assays and principles for screening high-confidence genes.

(**G**) The pie chart shows the distribution of editing sites in different transcript region in cells expressing DRAM (n=3 independent samples).

(**H**) The density map showing the distribution of editing events across the mRNA transcripts detected by DRAM-seq.

(**I**) The frequency plot shows the distribution of the distances of edit events in DRAM-seq relative to the m5C sites from the published BS-seq datasets. The position of each m5C site of BS-seq is determined as 0, and the relative distance of each site to the nearest edit event in DRAM-seq is calculated and plotted. The plots are presented separately based on the cutoff of upstream and downstream 3000bp (above) and 80bp (below) windows.

748   **(J, K)** Motif analysis discovered within the ±20nt region around the C-to-U or A-to-G editing site

749   in cells expressing DRAM-CBE (**J**), APOBEC1(**J**), DRAM-ABE (**K**) and TadA-8e (**K**).

Fig.4: Stable and comprehensive cellular identification of m^{5}C loci by DRAM-seq.

(**A**) Comparison of the overall distribution of genes with m^{5}C modifications detected by DRAM-seq, Yang et al. and Zhang et al. on chromosomes. The mutation sites detected by DRAM-seq on each gene are categorized into dual-colored short lines, with positive strand mutations shown in orange and negative strand mutations in dark green. The line graph and kernel density plot in the inner ring represent the locations and distributions of overlapping genes detected by DRAM-seq (red), Yang et al. (blue) and Zhang et al. (light green).

(**B**) Venn diagram showing the overlap between DRAM-seq and Yang et al.'s edited genes.

(**D**) Venn diagram showing the overlap between DRAM-seq and Zhang et al.'s edited genes.

(**C, E**) Integrative genomics viewer (IGV) browser traces of DRAM-seq data expressing the indicated constructs in the ATG16L1(**B**), ARHGEF25(**B**), FANCD2(**D**), and RPL15(**D**) mRNAs. C-to-U/A-to-G mutations found in at least 10% of reads are indicated by coloring, and the m^{5}C site found by BS-seq is also labelled.

(**F**) Genes with DRAM-seq editing events were analyzed for KEGG bioprocess enrichment.

(**G**) GO biological processes enrichment analysis of genes with DRAM-seq editing events. Statistical analyses were performed using the DAVID tool.

Fig.5: Low-input m⁵C detection and transfection efficiency of DRAM system.

(**A**, **B**) DRAM analysis of RPSA (**A**) and AP5Z1 (**B**) mRNAs with 250 ng, 50 ng, and 10 ng of input RNA. Representative Sanger sequencing plots are shown on the left panel, with mutation sites marked with asterisks. The mutation rates are quantified on the right panel.

(**C**) Flowchart illustrating Cell viability analysis by CCK8 reagent after DRAM transfection in HEK293T cells.

(**D**) Quantitative comparison of the relative proliferative capacity of DRAM-expressing and untransfected cells.

(**E**, **F**) The expression levels of DRAM-CBE (**E**) and DRAM-ABE (**F**) systems different plasmid transfection concentrations were verified by Western blotting.

(**G**, **H**) Editing of RPSA (**G**) and AP5Z1 (**H**) mRNA at varying concentrations of DRAM protein expression. The left panels indicate Sanger sequencing results following RT-PCR, while the corresponding quantifications of DRAM-induced mutations are shown in the right panels.

785