Placental malaria is associated with a TLR–Endothelin-3–oxidative damage response in human placenta tissues

Samuel Chenge[1,2], Melvin Mbalitsi[1], Harrison Ngure[1], Moses Obimbo[3], Mercy Singoei[3], Mourine Kangogo[2], Bernard N. Kanoi[1], Jesse Gitaka[1,*], and Francis M. Kobia[1,*]

[1]Centre for Malaria Elimination, Mount Kenya University, Thika – Kenya.

[2]Department of Medical Microbiology, Jomo Kenyatta University of Agriculture and Technology, Nairobi – Kenya

[3]Department of Human Anatomy, University of Nairobi, Nairobi – Kenya

*Correspondence: Francis M. Kobia (fkobia@associates.mku.ac.ke), Center for Malaria Elimination, Mount Kenya University, P.O. Box 342-01000, Thika – Kenya. Jesse Gitaka (jgitaka@mku.ac.ke), Center for Malaria Elimination, Mount Kenya University, P.O. Box 342-01000, Thika – Kenya.

Abstract

Placental malaria, which is mainly caused by the sequestration of *Plasmodium falciparum*-

infected erythrocytes in the placenta, is an important driver of poor pregnancy outcomes, including

fetal growth restriction, preterm birth, and stillbirth. However, the mechanisms underlying its

adverse outcomes are unclear. Mouse models have shown that placental malaria triggers a

proinflammatory response in the placenta, which is accompanied by a fetal Toll-like receptor

(TLR)4-mediated innate immune response associated with improved fetal outcomes. Here, we

used hematoxylin and eosin staining to identify placental malaria positive and negative samples

in our biobank of placentas donated by women living in a malaria-endemic region of Kenya and

assessed the impact of placental malaria on the expression of TLRs, Endothelins, and oxidative

damage. RT-qPCR analysis revealed that placental malaria was associated with an upregulation

of TLR4, TLR7, and Endothelin-3. Moreover, immunohistochemistry showed that placental

malaria was associated with elevated expression levels of the oxidative DNA damage marker, 8-

hydroxy-2'-deoxyguanosine, while RT-qPCR revealed that this was accompanied by an

upregulation of p21, an inhibitor of cell cycle progression and marker of cellular response to DNA

damage. These findings allude to a novel mechanism of placental malaria pathogenesis driven

by a TLR–Endothelin-3–oxidative DNA damage signaling axis.

Keywords: Placental malaria, malaria in pregnancy, poor pregnancy outcomes, innate immunity,

Plasmodium falciparum

1. Introduction

According to the World Health Organization, globally, there were about 249 million malaria cases and 608,000 malaria-associated deaths in 2022, with sub-Saharan Africa accounting for most of the cases and deaths [1]. Pregnant women have a higher susceptibility to malaria infection [2] and it is estimated that in 2022, there were about 12.7 million cases of malaria in pregnancy in sub-Saharan Africa [1]. Malaria in pregnancy is associated with several adverse outcomes in the mother, fetus, and neonate [3]. For the fetus, malaria in pregnancy severely worsens pregnancy outcomes and frequently leads to fetal growth restriction (including low birthweight, small for gestational age, and intrauterine growth restriction) and may result in preterm birth and stillbirth [3–5].

The adverse effects of malaria in pregnancy on the fetus are attributable to malaria infection of the placenta [1], leading to placental malaria. Placental malaria is characterized by the sequestration of *Plasmodium*-infected erythrocytes in placental intervillous spaces. This phenomenon is most frequently associated with *Plasmodium falciparum* (*P. falciparum*), the species associated with the most severe form of malaria [6]. The sequestration of *P. falciparum*-infected erythrocytes in the placenta is mediated by the interaction between variant surface chondroitin surface antigen 2, a *Plasmodium falciparum* protein expressed on the surface of infected erythrocytes [7], and chondroitin sulfate A on the surface of the syncytiotrophoblast [7], the placental epithelial cell layer that contacts maternal blood [8].

The adverse impacts of placental malaria on fetal well-being most likely result from the negative effects of placental malaria on placental health and function, since the vertical transmission of malaria to the fetus is rare [9]. Indeed, placental malaria is reported to induce placental inflammation [10,11] and placental histological changes [12], which may contribute to placental insufficiency and poor pregnancy outcomes. However, the mechanisms underlying placental

malaria-driven placental pathobiology are not fully understood at the cellular and cell signaling

levels.

Innate immune factors, such as Toll-like receptor (TLR)4, 7, and 9, which respond to infection by

recognizing invading pathogens for clearance, are reported to respond to malaria infection [13],

although their role in placental malaria is unclear. Although several studies have investigated

maternal responses to placental malaria, few have studied how the fetus responds to parasite

sequestration in the placenta. Nonetheless, a mouse model revealed that placental malaria

triggers a TLR4-mediated innate immune reaction that adversely affects fetal outcomes, which is

countered by a fetal innate immune reaction that led to better pregnancy outcomes [14]. This

suggests the presence of TLR-mediated innate immune responses to placental malaria, although

this has not been reported in the context of human placental malaria. Here, considering that

mouse data show that TLR4 modulates endothelin-1 expression [15], malaria is inflammatory and

oxidative [16], oxidative DNA damage upregulates TLR4 [17], and TLR signaling is thought to

promote DNA repair [18], we used biobank placenta samples donated by women living in a

malaria-endemic region of Kenya to examine the hypothesis that human placental malaria triggers

a TLR–Endothelin–oxidative damage signaling response.

2. Materials and methods

2.1 The biobank

The study used biobank placenta samples donated by residents of Bungoma County, a malaria-

endemic region of Western Kenya. The biobank was established by a previous prospective study

as described recently [19]. This study did not require participant written informed consent since it

was granted during biobanking [19]. Participants with a known record of sexually transmitted

disease infection during pregnancy, those with pregnancy-associated noncommunicable

diseases (preeclampsia and gestational diabetes) during the current pregnancy (at placenta

donation), and those with twin pregnancies were excluded from analyses. Based on the

underlying biobank data, malaria in pregnancy was defined as having at least one episode of

hospital-diagnosed malaria during pregnancy. All data underlying the databank were deidentified.

Equal numbers of male and female placentas were analyzed. The characteristics of the biobank's

placenta donors and samples are summarized in Table 1.

2.2 Histological analysis

Formalin-fixed placenta tissues were embedded in paraffin blocks as previously described [20]

using an automated tissue embedding system (MediMeas). H&E analysis was used to confirm

placental malaria presence, which is indicated by the presence of infected erythrocytes in the

placenta. Briefly, formalin-fixed paraffin-embedded samples were sectioned onto charged

microscope slides (Dako, Cat No. K8020) at a 5-μm thickness, dried at 37 °C overnight on a slide

warmer, dewaxed in xylene (Finar Chemicals, Cat No. 21940LC250), rehydrated by dipping

across an alcohol gradient of absolute, 95%, 70%, and 50% ethanol (Scharlau, Cat No.

ET00052500), and then in distilled water. They were then submerged in hematoxylin (Loba

Chemie, Cat No. 04023) for seven minutes, rinsed with running water, and then destained through

10 dips in acid alcohol (1% hydrochloric acid in 70% ethanol). Next, they were submerged in eosin

(Griffchem, Cat No. 45380) for 45 seconds followed by dehydration in 95% ethanol and absolute

ethanol (five minutes each) and then cleared in xylene baths (10 minutes each) before being

cover-slipped using dibutylphthalate polystyrene xylene mountant (Finar Chemicals, Cat No.

10525LM250). The slides were then examined under a microscope (Richter Optica UX1, M2

Scientifics), followed by imaging in ≥10 fields of view per slide at a 40X magnification using a

Moticam microscope camera (Motic Scientific). Placental malaria was then diagnosed as

described before [21], based on the presence of infected erythrocytes. Placental malaria-

associated histopathological features were assessed by counting the number of syncytial knots

as described previously [22], and measuring the fibrin-occupied placental areas using imageJ.

These analyses were done in at least 10 fields of view per sample. The levels of placental malaria

burden in the placental malaria-positive samples were determined by counting the number of

identified infected erythrocytes in at least 10 fields of view per slide imaged at a magnification of

40X.

2.4 RNA extraction and reverse transcription quantitative PCR (RT-qPCR)

Total RNA was extracted from placenta tissue using a HigherPurity™ Tissue Total RNA

Purification kit as per the manufacturer's guidelines (Canvax, cat No. AN0152) and quantified

using a NanoDrop Microvolume Spectrophotometer (ThermoFisher Scientific) following the

manufacturer's instructions. For each sample, 500 ng of RNA were retrotranscribed into cDNA

using a LunaScript™ RT SuperMix cDNA Synthesis Kit (NEB, Cat. No. E3010L) using the

manufacturer's protocol. RT-qPCR analysis was done on a QuantStudio™ 5 Real-Time PCR

System in a final volume of 20 µl containing 10 µl of GoTaq qPCR Master Mix (Promega, Cat No.

PRA6001), 2 µl of the forward plus reverse primers (final primer concentration: 500 nM), 3 µl of

nuclease free water (Promega, Cat No. P119E), and 5 µl of cDNA using the following program:

50 °C for two minutes, 95 °C for 10 minutes, followed by 40 cycles at 95 °C for 15 seconds and

60°C for 30 seconds. Relative gene expression was determined using the $2^{-\Delta\Delta ct}$ method [23], using

β-actin as the reference gene. Primers were purchased from Macrogen and primer sequences

are provided in Table 3.

2.5 *P. falciparum* detection PCR

The presence of *P. falciparum* in placenta samples was evaluated using One Taq® Quick-Load®

2X Master Mix with Standard Buffer (NEB, Cat No. M0486L) and the following thermocycler

program: Initial denaturation at 95 °C for five minutes, followed by 35 cycles of denaturation at 95

°C for 30 seconds, annealing at 55 °C for 60 seconds, and extension at 72 °C for 75 seconds,

and then a final extension at 72 °C for five minutes. Primer sequences are shown in Table 3. The

PCR product was subjected to agarose (Sigma–Aldrich, Cat No. A9539) gel electrophoresis using

1X tris–borate–EDTA buffer alongside a 100 base pair ladder using SafeView™ Classic (Applied

Biological Materials, Cat No. G108) nucleic acid stain and Gel Loading Dye, Purple (6X) (NEB,

Cat No. B7024S). The bands were developed and imaged using a UVITEC Gel Documentation

System (Cleaver Scientific).

2.6 Immunohistochemistry

The sections were deparaffinized by warming at 55 °C for 15 minutes followed by dipping in three

xylene baths, about 10 dips each. They were then rehydrated and subjected to heat-induced

epitope retrieval by boiling for 30 minutes in Citrate Buffer, pH 6.0 (Sigma–Aldrich, cat. No.

C9999). They were then cooled to room temperature, rinsed with distilled water for five minutes

and then blocked with 0.3% Triton-X in 1X phosphate-buffered saline (PBST). Next, they were

blocked with 10% normal donkey serum (Abcam, cat. No. ab7475) in PBST for two hours followed

by overnight incubation (4 °C) with anti-DNA/RNA damage antibody [15A3] (Abcam, cat. No.

ab62623) at 1:2500 in blocking solution. Sections were then washed thrice (10 minutes each)

using PBST and then incubated at room temperature for two hours with horseradish peroxidase-

conjugated goat anti-mouse secondary antibody (Jackson ImmunoResearch, cat. No. 115-035-

003) at 1:5000 in blocking solution. The sections were then washed thrice (10 minutes each) using

PBST followed by signal development using an ImmPACT® DAB Substrate Kit (Vector, cat. No.

SK-4105) as per the manufacturer's protocol. They were then dehydrated using 95%, 95%, 100%,

and 100% ethanol (five minutes each), cleared by dipping in three xylene baths and then cover-

slipped using a xylene-based mountant and allowed to dry. They were then examined under a

light microscope and imaged at a magnification of 40X.

2.7 Data analyses

Statistical analyses were done using GraphPad Prism version 9. Data are presented as

percentages, raw values, or mean ± standard deviation. Differences between two groups were

compared using a t-test. Correlation analyses were done using nonparametric Spearman

correlation analysis. For each placenta, the placental malaria burden in placentas was indicated

by the total number of infected erythrocytes in the placenta section. The birthweight-to-placenta

weight ratio was obtained by dividing the birthweight (grams) with the corresponding placenta's

weight (grams). The correlation between placental malaria status and birthweight, placental

weight, and birthweight-to-placental weight ratio was assessed using GraphPad prism to examine

the impact of placental malaria on fetal outcomes. $P < 0.05$ indicates statistically significant

differences.

3. Results

3.1 Main characteristics of the placenta donors and donated placentas

All placenta donors were ≥18-years-old. The mean age, gravidity, birthweight, placental weight,

and birthweight-to-placental weight ratio of the cohort of placenta donors were 24.7 years, 2.69,

3077.64 g, 470.04 g, and 6.51, respectively (Table 1). Grouping the placenta donors into those

with a known history of malaria in pregnancy (MiP) and those without (NoMiP), revealed that in

the MiP vs NoMiP groups, maternal age and gravidity were not significantly different (mean age:

24.4 [range: 18–30] vs 25 [range: 18–40] years, $P = 0.45$; mean gravidity: 2.5 [range: 1–7] vs 2.9

[range: 1–7), $P = 0.12$). However, in the MiP vs NoMiP groups, birthweight (mean: 2870.5 [range:

1600–5000] vs 3272.3 [range: 2000–4500] g, placental weight (mean: 464.3 [range: 280.4–675)

vs 492.9 [range: 342.3–715] g, and birthweight-to-placental weight ratio (mean: 6.27 [range: 3.2–

10.2] vs 6.73 [range: 4.18–10], were significantly lower in the MiP group ($P < 0.0001$, = 0.009,

and = 0.03, respectively). The presence of placental malaria was determined using H&E analysis

as we described recently [19]. Of the placentas, 92 (51.4%) belonged to male babies, and eight

(4.5%) were preterm (Table 2). All downstream analyses were done on placenta samples that

were confirmed to be placental malaria-positive or placental malaria-negative using H&E staining.

3.2 Placental malaria correlates negatively with birthweight and birthweight-to-placenta

ratio

Representative images of placental malaria-negative and placental malaria-positive samples are

shown in Figure 1A–B, and the presence of *P. falciparum* in placental malaria-positive tissues

was confirmed using PCR (Figure 1C). Analysis of the data underlying the placental biobank

revealed that when compared with the placental malaria-negative group, birthweight was

significantly lower in the placental malaria-positive group, which had more low birthweight cases

(Figure 1A, *P* = 0.03, low birthweight: <2500 g), but placental weight did not differ between the

two groups (Figure 2B, *P* = 0.8). However, the birthweight-to-placental weight ratio was lower in

the placental malaria-positive group, although the difference did not reach statistical significance

(Figure 1F, *P* = 0.1). Next, we used the H&E images to determine the proportion (%) of infected

erythrocytes (IEs) in each placenta sample, and then used the obtained values to assess the

correlation between the placental malaria burden and birthweight, placental weight, and the

birthweight-to-placental weight ratio, i.e., the fetal weight obtained per gram of the placenta, which

is an indicator of placental efficiency, with higher birthweight-to-placental weight ratios indicating

greater placental efficiency [24]. This analysis revealed negative correlation between IEs

(proportion of infected erythrocytes [%]) and birthweight (correlation coefficient [rs]: -0.23, *P* <

0.005, 95% confidence interval [CI]: -0.37 − -0.08), and IEs (%) and birthweight-to-placental

weight ratio (rs: -0.20, *P* = 0.007, 95% CI: -0.35 − -0.05). Expectedly, placenta weight had a

positive correlation with birthweight (rs: 0.28, *P* < 0.001, 95% CI: 0.13 − 0.41) and a negative

correlation with birthweight-to-placental weight ratio (rs: -0.42, *P* < 0.001, 95% CI: -0.54 − -0.28).

However, placental malaria burden did not exhibit correlation with placenta weight (rs: 0.01, *P* =

0.93, 95% CI: -0.16 − 0.15). Taken together, these findings indicate that placental malaria impairs

placenta function, leading to low birthweight via reduced placenta efficiency, as indicated by the

negative correlation between placental malaria burden and the birthweight-to-placental weight

ratios of placental malaria-exposed neonates.

3.3 Placental malaria markedly alters placental histological features

Next, we assessed the impact of placental malaria on syncytial knotting and fibrin deposition in

our placenta samples. This analysis revealed that when compared with placental malaria-negative

samples (A), placental malaria-positive samples had more syncytial knots (B, yellow arrowhead)

and greater fibrin-occupied placental area (C, FD [fibrin deposit] in the broken line-demarcated

area). Quantification analyses revealed that when compared with placental malaria-negative

samples, the levels of these histological features were significantly higher in placental malaria-

positive samples (D, SK [syncytial knots], P = 0.047 and E, fibrin area, P < 0.0005). These

observations indicate that in our study cohort, placental malaria may have adversely affected fetal

outcomes by at least in part, altering normal placental histological features.

3.4 Placental malaria is associated with an upregulation of TLR4, TLR7, and Endothelin 3

We then sought to determine if placental malaria altered the expression of TLRs. To this end, we

focused on TLR4, TLR7, and TLR9, which have been associated with response to malaria

infection in mice [25,26], and with mouse placental malaria in the case of TLR4 [27], although this

has not been reported in human placental malaria. To evaluate the effect of placental malaria on

these innate immune system receptors, we assessed their expression levels using RT-qPCR. The

analysis revealed that when compared with placental malaria-negative controls, placental

malaria-positive samples expressed significantly higher levels of TLR4 and TLR7, but not TLR9

(Figure 3A–C, P = 0.002, 0.03, and 0.59, respectively). This is consistent with mouse data

showing that placental malaria upregulates TLR4-mediated expression of endothelin-1 [15]. We

therefore wondered if human placental malaria alters the expression of Endothelin genes. RT-

qPCR analysis of Endothelin-1 and -3 gene expression revealed that only Endothelin-3 was

detectable in our placenta samples and that when compared with placental malaria-negative

placentas, placental malaria-positive samples had significantly higher Endothelin-3 levels (Figure

3D, P = 0.004), indicating the presence of a TLR–Endothelin signaling axis in response to human

placental malaria.

3.5 Placental malaria is associated with high oxidative DNA damage

Since TLR4 was upregulated in placental malaria-positive placenta samples, we wondered whether placental malaria-driven TLR4 activation is associated with a dysregulation of other signaling processes that may underlie or contribute to placental malaria-mediated placental pathobiology. Because malaria is known to be strongly inflammatory and oxidative, which may drive host tissue damage [16], and because oxidative DNA damage is associated with TLR4 upregulation [17] and that TLR signaling is thought to promote DNA repair [18], we wondered if the TLR4 upregulation in the placental malaria-positive samples might be associated with placental oxidative DNA damage. To assess this possibility, we used immunohistochemistry to assess the levels of 8-hydroxy-2'-deoxyguanosine, a marker of oxidative DNA stress [28]. This analysis revealed that when compared with placental malaria-negative samples, placental malaria-positive samples express markedly higher levels of 8-hydroxy-2'-deoxyguanosine (Figure 4A). Staining the same samples with the secondary antibody only (without the primary antibody) confirmed signal specificity (Figure 4B). To further assess the effect of placental malaria on oxidative stress, we used RT-qPCR to examine the level of p21, a mediator of cell cycle arrest and indicator of cellular response to DNA damage [29]. This revealed that when compared with placental malaria-negative samples, placental malaria-positive samples had significantly higher levels of p21 (Figure 4C, P = 0.02). Taken together, these data indicate that placental malaria triggers markedly high levels of placental oxidative DNA stress, placenta tissue damage, and cellular response to DNA stress, which may contribute to placental malaria pathobiology, and that in response, TLR signaling may at least in part, be upregulated to counter these adverse effects through the promotion of DNA repair.

4. Discussion

Malaria in pregnancy often results in placental malaria, where erythrocytes that are infected with *P. falciparum*, the parasite that most frequently causes placental malaria, sequester in placental

intervillous spaces [7]. Placental malaria may cause various adverse fetal outcomes, including

stillbirth, preterm birth, and fetal growth restriction [3–5], and because *P. falciparum* rarely

undergoes vertical transmission [9], these effects are likely caused by placental malaria-driven

pathobiological effects in the placenta, which may impair placental function. However, although

studies have implicated effects like inflammation and histological changes in placental malaria

pathogenesis, the mechanisms underlying the adverse effects of human placental malaria on the

placenta are unclear. Moreover, because many malaria in pregnancy cases in malaria-endemic

regions are asymptomatic [30,31] and the placenta is inaccessible during pregnancy, there are

no ways of detecting and intervening against placental malaria during pregnancy. Thus, there is

an urgent need to better understand the placental pathobiology of placental malaria to guide the

development of effective diagnostic and therapeutic tools.

Mouse models indicate that placental malaria triggers innate immune responses (mainly via

TLR4) that are associated with poor fetal outcomes and that TLR4-mediated fetal responses to

placental malaria cause improved outcomes [14]. However, the effect of human placental malaria

on TLR-mediated immunity in the placenta has not been examined. In this study, we leveraged

our well-characterized biobank of placenta samples from a malaria endemic region of Kenya

(Table 1) and found that in our study cohort, placental malaria burden had a significant negative

correlation with birthweight and birthweight-to-placental weight ratio, that it was associated with

significantly higher placental histological lesions, higher levels of TLR4 and Endothelin-3

expression, and enhanced oxidative DNA damage when compared with samples without

placental malaria.

Consistent with previous findings implicating placental malaria in fetal growth restriction [32], we

observed that in our study cohort, relative to the placental malaria-negative cases, placental

malaria was associated with low birthweight. Moreover, we observed that placental malaria was

associated with lower birthweight-to-placental weight ratios, an indicator of placental efficiency in

which higher ratios indicate higher nutrient transfer for every gram of the placenta and vice versa

[24], but not with lower placental weight (Figure 1D–F), an observation that to our knowledge, has

not been previously reported in human placental malaria. Interestingly, our analyses also indicate

that the placental malaria burden (percentage of infected erythrocytes in a sample's intervillous

spaces) correlates negatively with birthweight but not with placental weight. Taken together, these

observations indicate that placental malaria contributes to fetal growth restriction primarily by

impairing placental function, and not via placental growth inhibition, although the precise

mechanisms remain unclear. This possibility is crucial considering that women in malaria-endemic

regions may experience multiple malaria reinfections throughout pregnancy. Therefore, further

studies, such as using *in vitro* and organoid systems, are needed to comprehensively investigate

this possibility.

Our analyses revealed that placental malaria-positive samples had markedly higher levels of fibrin

deposition and syncytial knotting than placental malaria-negative samples, which is in line with

earlier findings [33]. These changes, which indicate placental injury, and have been associated

with placental malperfusion and poor fetal outcomes, including fetal growth restriction [34,35],

may contribute to the low birthweight observed in our placental malaria cohort. However, studies

are needed to establish the mechanisms by which placental malaria alters placental histological

features, how these changes correlate with fetal outcomes and postnatal wellbeing, and whether

they can predict fetal wellbeing in postnatal life.

TLRs are key innate immunity factors that sense host invasion by pathogens and activate host

immune defenses [36]. Mouse models of malaria indicate that TLR4, TLR7, and TLR9 are

involved in detecting and responding to malaria infection [25,26]. Moreover, mouse models

indicate that at the fetal–maternal interface, placental malaria activates TLR4-mediated immune

responses that drive poor fetal outcomes, and that fetal TLR4-mediated counterresponses

improve pregnancy outcomes [14,27,37]. However, this observation has not been previously

made in human placental malaria. Here, we observed that the expression levels of TLR4 and

TLR7, but not TLR9, were significantly upregulated in placental malaria-positive samples,

indicating that placental infection triggers an innate immune reaction and that it is mainly driven

by TLR4 and TLR7, although the status of other TLRs during placental malaria warrants

investigation. Considering that TLRs are important drivers of inflammation [38], which is

implicated in placental malaria pathogenesis [39], taken together with the observed changes in

placental histological features, our findings indicate for the first time, that TLR-mediated

responses to placental malaria may contribute to local placental inflammation, which may underlie

the observed placental malaria-associated low birthweight and placental weight, although the

precise mechanisms remain unclear. Mouse data show that placental malaria-driven TLR4

expression drives placental endothelin-1 expression [15], and for the first time, our findings show

that placental malaria is also associated with the upregulation of Endothelin-3. The Endothelin

ligands 1, 2, and 3 are a family of vasoactive factors that influence a range of cellular processes,

such as vascular remodeling and angiogenesis [40]. Moreover, Endothelin-3 is reported to have

anti-inflammatory effects [41,42], suggesting that its upregulation in the context of placental

malaria-mediated TLR4 upregulation is a mechanism of countering TLR4-dependent placenta

inflammation. Collectively, these observations indicate that human placental malaria may activate

a TLR4–Endothelin-3 signaling axis, but further studies are needed to test this hypothesis and to

determine its implications in placental malaria pathobiology and fetal outcomes.

Based on reports that malaria is strongly oxidative [16], oxidative stress causes tissue damage

[43], oxidative DNA damage upregulates TLR4 [17], and that the TLR pathway might promote

DNA repair [18], we reasoned that our observation of TLR4 and TLR7 upregulation in placental

malaria-positive samples might be accompanied by placental oxidative DNA damage. This

hypothesis was confirmed by our immunohistochemistry data, which showed that 8-hydroxy-2'-

deoxyguanosine, a marker of oxidative DNA stress [28], was markedly upregulated in placental

malaria samples. Moreover, gene expression analysis revealed that these events were accompanied by a significant upregulation of p21, a cell cycle inhibitor and marker of cellular response to DNA damage [29]. These observations align with previous findings that in a mouse model, placental malaria is associated with placental oxidative damage [44]. Furthermore, p21 upregulation in the placenta may arrest the cell cycle to allow for oxidative damage resolution, which may contribute to the low placental weight observed in our cohort, but this possibility requires further investigation. Together, our data suggest the presence of a previously unknown TLR–Endothelin–oxidative damage axis in human placental malaria.

Finally, our data also show that in the placental malaria-negative and placental malaria-positive groups, the analyzed parameters varied across samples, which may reflect inter-individual biological variability [45]. In the future, our observations will be validated experimentally, such as by challenging cultured primary human trophoblasts with *P. falciparum* for detailed mechanistic interrogation.

5. Conclusion

Despite its heavy burden and adverse effects on maternal and fetal outcomes, the mechanisms underlying the placental pathobiology of placental malaria are unclear. Considering that malaria is rarely transmitted to the fetus [9], the adverse fetal outcomes of malaria in pregnancy are mainly driven by events that disrupt placenta physiology and function. Importantly, because of placental inaccessibility, placental malaria can only be confirmed via postnatal placental histopathology. These challenges highlight the urgent need to better understand the mechanisms underlying the placental pathobiology of placental malaria, which may inform the development of sensitive tools for diagnosing placental malaria during pregnancy as well as effective therapeutic interventions. Our findings that placental malaria may drive TLR-mediated responses in the placenta raise the possibility that modulating innate responses to placental malaria may improve fetal outcomes, as we previously discussed [13]. Moreover, our identification of an axis involving TLRs, Endothelins,

and oxidative DNA damage during placental malaria (Figure 5) highlights processes with the

potential for intervention against human placental malaria. However, further studies, such as using

primary human trophoblasts, human placental organoids, or human placental *ex vivo* systems are

needed to validate our observations. Such approaches can investigate the mechanisms of

placental malaria pathobiology more rigorously than can be done using term placentas.

Funding statement

This project has received funding from the EDCTP2 programme (TMA2019CDF-2736) supported

by the European Union and Novartis Global Health Basel Switzerland.

Acknowledgments

We thank our placenta donors and the staff at Webuye County and Mary Help of The Sick (Thika)

Hospitals for their generous support. We thank the staff of the histopathology and anatomy

departments at the University of Nairobi and Mount Kenya University for their help with

histopathology. We are grateful to Prof. Walter Jaoko, Prof. Omu Anzala, and Dr. Daniel Muema

of KAVI–ICR for kindly sharing laboratory space and resources. We are thankful to Prof. Roger

Smith and Dr. Kaushik Maiti, Mothers and Babies Research Centre, Hunter Medical Research

Institute, Newcastle, NSW – Australia, for sharing antibodies and other resources.

Conflict of interest

The authors declare no conflicts of interest.

Ethics statement

This study was approved by Mount Kenya University's ethics review committee (approval number

2516).

Data accessibility

The data underlying this study are contained in the article.

Figures

Figure 1. (A–B) Representative hematoxylin and eosin images of placental malaria (PM)-negative (A) and PM-positive samples (B). When compared with a PM-negative sample, the positive sample has malaria-infected erythrocytes (black arrowheads) in the placental intervillous space. (C) PCR confirmed the presence *of P. falciparum* in the PM-positive sample in B. L: 100 base pair ladder, N: PM-negative sample, S: PM-positive sample in B (histology), P: positive control (*P. falciparum* strain 3D7 genomic DNA). Expected PCR band size: 205 base pairs. (D–F) PM was associated with lower birthweight (BW) (D, $P = 0.03$) and lower birthweight-to-placental weight (BW:PW) ratio, although the difference did not reach statistical significance (F, $P = 0.1$), but not with lower placental weight (PW) (E, $P = 0.8$). In D–F, whiskers are drawn from the 10^{th} to 90^{th} percentile. (G) A correlation matrix shows that the proportion of infected erythrocytes, IEs (%), in the placenta correlated negatively with BW ($P < 0.005$) and the BW:PW ratio ($P = 0.007$), but it did not correlate with PW ($P = 0.94$). PW correlated positively with BW and negatively with the BW:PW ratio (both $P < 0.001$).

Figure 2. (A–C) In the samples from the biobank underlying our study, when compared with

placental malaria (PM)-negative samples (A), PM was associated with significantly higher rates

of syncytial knots (B, yellow arrowhead) and placental fibrin deposits (C; FD, marked with broken

line). Black arrowheads indicate infected erythrocytes. (D–E) Quantification revealed that the

levels of syncytial knots (SK [D], n = 21 and 38 for PM-neg and PM-pos, respectively; *P* = 0.047)

and the area of placenta intervillous space containing fibrin (E, n = 26 and 38 for PM-neg and PM-

pos, respectively, *P* < 0.0005) were significantly higher in the PM-positive (PM-pos) samples than

in the PM-negative (PM-neg) samples. Whiskers are drawn from the 10th to 90th percentile.

Figure 3. Placental malaria is associated with the upregulation of TLR4, TLR7, and Endothelin-3. (A–C) When compared with placental malaria (PM)-negative samples (PM-neg), PM-positive (PM-pos) samples expressed significantly higher levels of TLR4 and TLR7, but not TLR9 (P = 0.002, 0.03, and 0.59, respectively). (D) PM-positive samples also expressed higher levels of Endothelin-3 (P = 0.004).

Figure 4. Analysis of oxidative DNA damage in placental malaria (PM)-negative vs PM-positive

samples. (A–B) Immunohistochemistry revealed that when compared with PM-negative samples,

PM-positive tissues had markedly higher levels of 8-hydroxy-2'-deoxyguanosine (8-OHdG), an

indicator of oxidative damage. Staining the same samples with the secondary antibody only (B)

confirmed signal specificity for this marker. (C) RT-qPCR showed that PM-positive samples

express significantly higher levels of p21 (P = 0.02).

Figure 5. Schematic summary of the hypothesized TLR–Endothelin-3–oxidative stress axis in human placental malaria. Further investigation is needed to validate this axis and determine its potential for intervention against placental malaria.

442   **Tables**

443   **Table 1. Summary of placenta donors' demographics**

Group	Age (range)	Gravidity (range)	BW (g) (range)	PW (g) (range)	BW:PW (range)
All donors	24.7 years (18–40)	2.69 (1–7)	3077.64 (1600–5000)	479.04 (280.37–715)	6.51 (3.2–10.2)
MiP	24.4 years (18–30)	2.5 (1–7)	2870.5 (1600–5000) ****	464.3 (280.4–675) **	6.27 (3.2–10.2) *
NoMiP	25 years (18–40)	2.9 (1–7)	3272.3 (2000–4500)	492.9 (342.3–715)	6.73 (4.18–10)

Analysis of the placenta donors' data revealed that maternal age and gravidity were not

significantly different in the MiP (group with known history of malaria in pregnancy) vs the NoMiP

(group without known MiP history) groups (P = 0.45 and 0.12, respectively), whereas birthweight

(BW), placental weight (PW), and BW:PW ratios were significantly lower in the MiP group when

compared with the No MiP group (P = < 0.0001, 0.009, and 0.03, respectively). *, **, and ****

indicate P < 0.05, 0.005, and 0.0005, respectively.

Table 2. Main characteristics of the donated placentas

Placental malaria-positive placenta samples (infected erythrocytes present)	92 (51.4%)
Placental malaria-negative placenta samples	58 (32.4%)
Placentas with past placental malaria infection (hemozoin present)	29 (16.2%)
Placenta samples from male fetuses	92 (51.4%)
Placenta samples from female fetuses	87 (48.6 %)
Placenta samples from pre-term deliveries	8 (4.5%)

The general characteristics of the placentas used in this study are summarized. Placental malaria

status was determined using hematoxylin and eosin (H&E) staining of placental tissue. Placental

malaria: observation of at least one infected erythrocyte [46]. Past placental malaria: absence of

infected erythrocytes coupled with hemozoin in fibrin and pigmented macrophages [21]. Chronic

placental malaria: detection of infected erythrocytes, as well as hemozoin in fibrin and pigmented

macrophages (past placental malaria) [47].

457   **Table 3. List of primers used in the study**

Target	Forward primer	Reverse primer
TLR4	5'-AGACCTGTCCCTGAACCCTAT-3'	5'-CGATGGACTTCTAAACCAGCCA-3'
TLR9	5'-CTGCCTTCCTACCCTGTGAG-3'	5'-GGATGCGGTTGGAGGACAA-3'
TLR7	5'-TCCTTGGGGCTAGATGGTTTC-3'	5'-TCCACGATCACATGGTTCTTTG-3'
β-actin	5'-CATGTACGTTGCTATCCAGGC-3'	5'-CTCCTTAATGTCACGCACGAT-3'
p21	5'-TGTCCGTCAGAACCCATGC-3'	5'-AAAGTCGAAGTTCCATCGCTC-3'
Endothelin-3	5'-GGGACTGTGAAGAGACTGTGG-3'	5'-AGACACACTCCTTGTCCTTGTA-3'
P. falciparum	5'-TTAAACTGGTTTGGGAAACCAAATATATT-3'	5'-ACACAATGAACTCAATCATGACTACCCGTC-3'

References

1. World Health Organization. World malaria report 2023. 2023. Available: https://www.wipo.int/amc/en/mediation/

478
479

2. Rogerson SJ, Mwapasa V, Meshnick SR. Malaria in pregnancy: Linking immunity and pathogenesis to prevention. American Journal of Tropical Medicine and Hygiene. 2007;77: 14–22.

480
481
482

3. Bauserman M, Conroy AL, North K, Patterson J, Bose C, Meshnick S. An Overview of Malaria in Pregnancy. Semin Perinatol. 2019;43: 282. doi:10.1053/J.SEMPERI.2019.03.018

483
484
485

4. Fried M, Kurtis JD, Swihart B, Pond-Tor S, Barry A, Sidibe Y, et al. Systemic Inflammatory Response to Malaria During Pregnancy Is Associated With Pregnancy Loss and Preterm Delivery. Clin Infect Dis. 2017;65: 1729. doi:10.1093/CID/CIX623

486
487
488

5. Schmiegelow C, Matondo S, Minja DTR, Resende M, Pehrson C, Nielsen BB, et al. Plasmodium falciparum Infection Early in Pregnancy has Profound Consequences for Fetal Growth. J Infect Dis. 2017;216: 1601–1610. doi:10.1093/infdis/jix530

489
490
491

6. Trampuz A, Jereb M, Muzlovic I, Prabhu RM. Clinical review: Severe malaria. Crit Care. 2003;7: 315. doi:10.1186/CC2183

492
493

7. Ayres Pereira M, Mandel Clausen T, Pehrson C, Mao Y, Resende M, Daugaard M, et al. Placental Sequestration of Plasmodium falciparum Malaria Parasites Is Mediated by the Interaction Between VAR2CSA and Chondroitin Sulfate A on Syndecan-1. Rowe JA, editor. PLoS Pathog. 2016;12: e1005831. doi:10.1371/journal.ppat.1005831

494
495
496
497

8. Frank HG. Placental Development. Fetal and Neonatal Physiology, 2-Volume Set. 2017; 101–113. doi:10.1016/B978-0-323-35214-7.00010-X

498
499

9. Harrington WE, Duffy PE. Congenital malaria: Rare but potentially fatal. Ped Health. 2008;2: 235–248. doi:10.2217/17455111.2.2.235/ASSET/IMAGES/LARGE/GRAPHIC22.JPEG

500
501
502

503   10.   Fried M, Duffy PE. Malaria during Pregnancy. Cold Spring Harb Perspect Med. 2017;7:

504   a025551. doi:10.1101/cshperspect.a025551

505   11.   Rogerson SJ, Brown HC, Pollina E, Abrams ET, Tadesse E, Lema VM, et al. Placental

506   Tumor Necrosis Factor Alpha but Not Gamma Interferon Is Associated with Placental

507   Malaria and Low Birth Weight in Malawian Women. Infect Immun. 2003;71: 267.

508   doi:10.1128/IAI.71.1.267-270.2003

509   12.   Zakama AK, Ozarslan N, Gaw SL. Placental Malaria. Curr Trop Med Rep. 2020;7: 162–

510   171. doi:10.1007/S40475-020-00213-2

511   13.   Kobia FM, Maiti K, Obimbo MM, Smith R, Gitaka J. Potential pharmacologic interventions

512   targeting TLR signaling in placental malaria. Trends Parasitol. 2022;38: 513–524.

513   doi:10.1016/J.PT.2022.04.002

514   14.   Rodrigues-Duarte L, Pandya Y, Neres R, Penha-Gonçalves C. Fetal and maternal innate

515   immunity receptors have opposing effects on the severity of experimental malaria in

516   pregnancy: Beneficial roles for fetus-derived Toll-like receptor 4 and type I interferon

517   receptor 1. Infect Immun. 2018;86. doi:10.1128/IAI.00708-

518   17/SUPPL_FILE/ZII999092383S1.PDF

519   15.   Pandya Y, Marta A, Barateiro A, Bandeira CL, Dombrowski JG, Costa J, et al. TLR4-

520   Endothelin Axis Controls Syncytiotrophoblast Motility and Confers Fetal Protection in

521   Placental Malaria. Infect Immun. 2021;89. doi:10.1128/IAI.00809-20/ASSET/283CE60E-

522   5A6D-46F3-9439-7A889F86217A/ASSETS/IMAGES/LARGE/IAI.00809-20-F001.JPG

523   16.   Vasquez M, Zuniga M, Rodriguez A. Oxidative Stress and Pathogenesis in Malaria. Front

524   Cell Infect Microbiol. 2021;11. doi:10.3389/FCIMB.2021.768182

525   17.   Tawadros PS, Powers KA, Ailenberg M, Birch SE, Marshall JC, Szaszi K, et al. Oxidative

526   Stress Increases Surface Toll-Like Receptor 4 Expression in Murine Macrophages Via

527   Ceramide Generation. Shock. 2015;44: 157–165. doi:10.1097/SHK.0000000000000392

18.   Harberts E, Gaspari AA. TLR Signaling and DNA Repair: Are They Associated? J Invest

Dermatol. 2013;133: 296. doi:10.1038/JID.2012.288

19.   Mbalitsi M, Singoei M, Chenge S, Ngure HM, Angienda PO, Obimbo MM, et al. A biobank

resource for placental malaria and placental health research. Open Research Europe.

2024;4: 273. doi:10.12688/OPENRESEUROPE.18617.1

20.   Canene-Adams K. Preparation of Formalin-fixed Paraffin-embedded Tissue for

Immunohistochemistry. Methods Enzymol. 2013;533: 225–233. doi:10.1016/B978-0-12-

420067-8.00015-5

21.   Odongo CO, Odida M, Wabinga H, Obua C, Byamugisha J. Burden of Placental Malaria

among Pregnant Women Who Use or Do Not Use Intermittent Preventive Treatment at

Mulago Hospital, Kampala. Malar Res Treat. 2016;2016. doi:10.1155/2016/1839795

22.   Senagore PK, Holzman CB, Parks WT, Catov JM. Working towards a Reproducible Method

for Quantifying Placental Syncytial Knots. http://dx.doi.org/102350/15-08-1701-OA1.

2016;19: 389–400. doi:10.2350/15-08-1701-OA.1

23.   Livak KJ, Schmittgen TD. Analysis of Relative Gene Expression Data Using Real-Time

Quantitative PCR and the $2-\Delta\Delta CT$ Method. Methods. 2001;25: 402–408.

doi:10.1006/METH.2001.1262

24.   Hayward CE, Lean S, Sibley CP, Jones RL, Wareing M, Greenwood SL, et al. Placental

adaptation: What can we learn from Birthweight:placental weight ratio? Front Physiol.

2016;7: 177721. doi:10.3389/FPHYS.2016.00028/BIBTEX

25.   Zhang Y, Zhu X, Feng Y, Pang W, Qi Z, Cui L, et al. TLR4 and TLR9 signals stimulate

protective immunity against blood-stage Plasmodium yoelii infection in mice. Exp Parasitol.

2016;170: 73–81. doi:10.1016/J.EXPPARA.2016.09.003

26.   Baccarella A, Fontana MF, Chen EC, Kim CC. Toll-Like Receptor 7 Mediates Early Innate

Immune Responses to Malaria. Infect Immun. 2013;81: 4431. doi:10.1128/IAI.00923-13

27.   Barboza R, Lima FA, Reis AS, Murillo OJ, Peixoto EPM, Bandeira CL, et al. TLR4-Mediated

554   Placental Pathology and Pregnancy Outcome in Experimental Malaria. Sci Rep. 2017;7.

555   doi:10.1038/S41598-017-08299-X

556   28.   Valavanidis A, Vlachogianni T, Fiotakis C. 8-hydroxy-2' -deoxyguanosine (8-OHdG): A

557   critical biomarker of oxidative stress and carcinogenesis. J Environ Sci Health C Environ

558   Carcinog Ecotoxicol Rev. 2009;27: 120–139. doi:10.1080/10590500902885684

559   29.   Villeneuve NF, Sun Z, Chen W, Zhang DD. Nrf2 and p21 regulate the fine balance between

560   life and death by controlling ROS levels. Cell Cycle. 2009;8: 3255.

561   doi:10.4161/CC.8.20.9565

562   30.   Salgado C, Ayodo G, Macklin MD, Gould MP, Nallandhighal S, Odhiambo EO, et al. The

563   prevalence and density of asymptomatic Plasmodium falciparum infections among children

564   and adults in three communities of western Kenya. Malar J. 2021;20. doi:10.1186/S12936-

565   021-03905-W

566   31.   Idris ZM, Chan CW, Kongere J, Gitaka J, Logedi J, Omar A, et al. High and Heterogeneous

567   Prevalence of Asymptomatic and Sub-microscopic Malaria Infections on Islands in Lake

568   Victoria, Kenya. Scientific Reports 2016 6:1. 2016;6: 1–13. doi:10.1038/srep36958

569   32.   Menendez C, Ordi J, Ismail MR, Ventura PJ, Aponte JJ, Kahigwa E, et al. The Impact of

570   Placental Malaria on Gestational Age and Birth Weight. J Infect Dis. 2000;181: 1740–1745.

571   doi:10.1086/315449

572   33.   Crocker IP, Tanner OM, Myers JE, Bulmer JN, Walraven G, Baker PN. Syncytiotrophoblast

573   degradation and the pathophysiology of the malaria-infected placenta. Placenta. 2004;25:

574   273–282. doi:10.1016/j.placenta.2003.09.010

575   34.   Lampi K, Papadogiannakis N, Sirotkina M, Pettersson K, Ajne G. Massive perivillous fibrin

576   deposition of the placenta and pregnancy outcome: A retrospective observational study.

577   Placenta. 2022;117: 213–218. doi:10.1016/J.PLACENTA.2021.12.013

35.   Brink LT, Roberts DJ, Wright CA, Nel DG, Schubert PT, Boyd TK, et al. Placental pathology

in spontaneous and iatrogenic preterm birth: Different entities with unique pathologic

features. Placenta. 2022;126: 54. doi:10.1016/J.PLACENTA.2022.06.004

36.   Takeda K, Kaisho T, Akira S. Toll-like receptors. Annual Review of Immunology. Annual

Reviews 4139 El Camino Way, P.O. Box 10139, Palo Alto, CA 94303-0139, USA ; 2003.

pp. 335–376. doi:10.1146/annurev.immunol.21.120601.141126

37.   Pandya Y, Penha-Gonçalves C. Maternal-fetal conflict during infection: Lessons from a

mouse model of placental malaria. Front Microbiol. 2019;10: 1126.

doi:10.3389/FMICB.2019.01126/BIBTEX

38.   Fukata M, Vamadevan AS, Abreu MT. Toll-like receptors (TLRs) and Nod-like receptors

(NLRs) in inflammatory disorders. Semin Immunol. 2009;21: 242–253.

doi:10.1016/J.SMIM.2009.06.005

39.   Chua CLL, Khoo SKM, Ong JLE, Ramireddi GK, Yeo TW, Teo A. Malaria in Pregnancy:

From Placental Infection to Its Abnormal Development and Damage. Front Microbiol.

2021;12: 3452. doi:10.3389/FMICB.2021.777343/BIBTEX

40.   Rodríguez-Pascual F, Busnadiego O, Lagares D, Lamas S. Role of endothelin in the

cardiovascular system. Pharmacol Res. 2011;63: 463–472.

doi:10.1016/J.PHRS.2011.01.014

41.   Sato A, Ebina K. Common mechanism in endothelin-3 and PAF receptor function for anti-

inflammatory responses. Eur J Pharmacol. 2013;718: 30–33.

doi:10.1016/J.EJPHAR.2013.09.025

42.   Sato A, Ebina K. Endothelin-3 at low concentrations attenuates inflammatory responses

via the endothelin B2 receptor. Inflamm Res. 2013;62: 417–424. doi:10.1007/S00011-013-

0594-3

43.   Burton GJ, Jauniaux E. Oxidative stress. Best Pract Res Clin Obstet Gynaecol. 2011;25:

287. doi:10.1016/J.BPOBGYN.2010.10.016

44.   Sarr D, Cooper CA, Bracken TC, Martinez-Uribe O, Nagy T, Moore JM. Oxidative Stress:

605   A Potential Therapeutic Target in Placental Malaria. Immunohorizons. 2017;1: 29.

606   doi:10.4049/IMMUNOHORIZONS.1700002

607   45.   Hansen KD, Wu Z, Irizarry RA, Leek JT. Sequencing technology does not eliminate

608   biological variability. Nat Biotechnol. 2011;29: 572. doi:10.1038/NBT.1910

609   46.   Muehlenbachs A, Nabasumba C, McGready R, Turyakira E, Tumwebaze B, Dhorda M, et

610   al. Artemether-lumefantrine to treat malaria in pregnancy is associated with reduced

611   placental haemozoin deposition compared to quinine in a randomized controlled trial. Malar

612   J. 2012;11: 1–9. doi:10.1186/1475-2875-11-150/FIGURES/5

613   47.   Rogerson SJ, Hviid L, Duffy PE, Leke RF, Taylor DW. Malaria in pregnancy: pathogenesis

614   and   immunity.   Lancet   Infectious   Diseases.   Elsevier;   2007.   pp.   105–117.

615   doi:10.1016/S1473-3099(07)70022-1