Streptococcus pneumoniae infection of lung epithelial cells induces internalization of surface

GPI-anchored proteins through pneumolysin-mediated activation of host Rho GTPases

Nick Lee[1,#], Manmeet Bhalla [2,#], Joana M. Pereira[4,5,6], Stacie Clark [1,3], Sara E. Roggensack [1,3],

Alenka Lovy[7], Michael J. Pereira[1,3], Guillermo Arroyo[1], Walter Adams[8], Andrew Camilli[1], Rodney

K. Tweten[9], John M. Leong[1,10,*] Sandra Sousa[4,5,*] and Elsa N. Bou Ghanem[2,*]

[#]Co-first authors. *Co-corresponding authors. These authors contributed equally to this work.

[1]Department of Molecular Biology & Microbiology, Tufts University School of Medicine, Boston,

MA

[2]Department of Microbiology & Immunology, University at Buffalo School of Medicine, Buffalo, NY

[3]Graduate Program in Molecular Biology & Microbiology, Tufts Graduate School of Biomedical

Sciences, Boston, MA

[4]i3S-Instituto de Investigação e Inovação em Saúde, Universidade do Porto, Porto, Portugal

[5]Instituto de Biologia Molecular e Celular, Universidade do Porto, Porto, Portugal

[6]Molecular and Cellular (MC) Biology PhD Program, ICBAS - Instituto de Ciência Biomédicas

Abel Salazar, University of Porto, Porto, Portugal

[7]Center for Vision Research and the Department of Ophthalmology and Visual Sciences, SUNY

Upstate Medical University Syracuse, NY

[8]Department of Biological Sciences, San Jose State University, San Jose, CA

[9]Department of Microbiology & Immunology, The University of Oklahoma Health Sciences

Center, Oklahoma City, OK

[10]Stuart B. Levy Center for the Integrated Management of Antimicrobial Resistance, Tufts

University, Boston, MA

Running title: GPI-anchored proteins are internalized post pneumolysin intoxication.

Keywords: Pneumococcus, Lung Epithelial Cells, PLY, Glycosylphosphatidylinositol Anchored

Proteins, GPI-AP, RhoGTPases, RhoA, Cdc42, Membrane Repair, Cholesterol-Dependent

Cytolysins

*Please address correspondence to John M. Leong: John.Leong@tufts.edu, Sandra Sousa:

srsousa@i3s.up.pt, and Elsa N. Bou Ghanem: elsaboug@buffalo.edu

Abstract

A return to homeostasis after infection-associated cellular injury can be accelerated by a rapid

damage response. *S. pneumoniae*, a typically asymptomatic colonizer of the host upper

respiratory tract, can cause serious and life-threatening infections when it gains access to the

lungs and other organs. The cholesterol binding *S. pneumoniae* pore-forming toxin, pneumolysin

(PLY), is central to the induction of host cell damage. Here, we first found that mouse lung

infection by *S. pneumoniae* diminished pulmonary expression of CD73, a

glycosylphosphatidylinositol anchored protein (GPI-AP) that modulates inflammation. Infection of

the human pulmonary epithelial cell line H292 resulted in a PLY-dependent reduction of not only

cell surface CD73, but also the population of surface expressed GPI-APs. The decrease in cell

surface GPI-APs was rapid, required pore-forming activity, and could be recapitulated by purified

PLY and other cholesterol binding cytolysins. In response to PLY-mediated insult, GPI-APs were

not released from the surface of epithelial cells in extracellular vesicles but rather internalized by

a mechanism dependent on the Rho GTPases RhoA and Cdc42. Internalization of GPI-APs was

associated with lower levels of PLY-induced apoptosis and membrane permeabilization. These

findings suggest that internalization of GPI-APs from epithelial cell membranes may constitute a

rapid innate repair response to cell damage induced by PLY and other pore forming toxins that

could help bacteria evade host defenses as many GPI-APs have roles in immunity.

Author summary

Streptococcus pneumoniae causes serious infections that can result in mortality. The pore-

forming toxin, pneumolysin (PLY) produced by these bacteria is important for their ability to cause

disease. Understanding how the host responds to damage by this toxin can result in better

treatment against infection. In this study, we found that PLY-mediated injury results in decreased

expression of glycosylphosphatidylinositol anchored proteins (GPI-AP) from the cell surface by

internalization. GPI-AP co-localize in cholesterol-rich areas of the membrane where PLY inserts

to form pores and cells with decreased surface GPI-APs were associated with less of PLY-

induced cell death and membrane permeabilization. These results suggest that GPI-AP are

internalized as part of repair mechanisms activated in response to infection-induced cell injury.

As many GPI-APs have important roles in the immune response, their removal from the cell may

inadvertently help the bacteria establish better infection.

Introduction

Streptococcus pneumoniae (pneumococcus), although an asymptomatic colonizer of the

upper respiratory tract of healthy hosts, can, in susceptible populations, cause severe lower

respiratory tract infections with high mortality and morbidity rates [1, 2]. *S. pneumoniae*

predominantly causes pneumonia but can reach the bloodstream and cause serious

extrapulmonary diseases, such as septicemia, meningitis, and endocarditis [3, 4]. More than 100

different *S. pneumoniae* serotypes have been identified based upon capsular polysaccharide

content [5, 6], which plays a role in the severity of pneumococcal disease [7]. *S. pneumoniae*

harbors many other virulence factors that promote its transition from the colonizer state to an

infectious stage. These range from surface proteins involved in interaction with host pulmonary

epithelium to the expression of factors that mediate metabolic adaptation or immune evasion [8].

Among these bacterial factors is a 53-kDa heat-labile cholesterol-dependent cytotoxin

(CDC), pneumolysin (PLY), which is a highly conserved and important virulence factor that

promotes the establishment of infection and evasion of host immune responses [9, 10]. PLY

creates pores in the cell membrane [11-13] through a multistep process [14] in which 30-50 toxin

monomers insert in the cholesterol-rich domains of host cells, oligomerize to form a large pre-

pore complex [15, 16] that then undergoes conformational changes to produce a large 25-nm β-

barrel pore in the membrane [11, 17]. Numerous studies have highlighted the role of PLY in

damaging the pulmonary epithelium, activating the classical complement pathway, inducing

inflammation, and inhibiting neutrophil function [8, 12]. Strains lacking PLY show reduced

virulence [18].

Glycosylphosphatidylinositol-anchored proteins (GPI-APs) are expressed in highly

divergent organisms, e.g., from protozoa to humans, and mediate diverse functions, such as

environmental sensing, cell adhesion and migration, and immunomodulation [19-21]. GPI-APs

are attached to the outer leaflet of the cell membrane through a highly conserved

glycosylphosphatidylinositol (GPI) anchor, which is composed of a phospholipid portion and a

glycan chain [22]. The GPI anchor is covalently linked to the C-terminus of the protein in a multi-

step post-translational modification process that occurs in the endoplasmic reticulum (ER) [23,

24]. The GPI-APs then migrates to the cell surface via the secretory pathway, which involves

trafficking through the ER, Golgi apparatus, and vesicular transport mechanisms, steps that are

at least partially dictated by the GPI anchor [25]. Machinery that controls membrane fusion and

sorting drives GPI-AP insertion in the plasma membrane, and in polarized epithelial cells, GPI-

APs are selectively sorted to the apical domain [26] where they are organized into lipid rafts,

specialized microdomains enriched in cholesterol and sphingolipids [22, 27]. In polarized cells the

clusters are assembled in the Golgi apparatus and traffic to the apical domain, while in non-

polarized cells the clusters organize from GPI-APs monomers already expressed at the cell

membrane [28]. The expression of GPI-APs on the cell surface is highly regulated. GPI-APs can

be released from the cell surface by the action of specific phospholipases, such as phospholipase

C (PLC) [20], which cleave the GPI anchor resulting in the release of the protein into the

extracellular environment. They can also be internalized through endocytosis, sorted and recycled

back to the cell surface following vesicular trafficking pathways [29].

Rho GTPases are small GTP-binding proteins that act as molecular switches by cycling

between an inactive GDP-bound state and an active GTP-bound state [30]. When activated, Rho

GTPases interact with downstream effector proteins to initiate specific signaling cascades, which

ultimately regulate key cellular functions [30]. In particular, Rho GTPases are implicated in the

regulation of intracellular membrane trafficking processes such as endocytosis, exocytosis, and

vesicle transport, and they control the cytoskeleton remodeling required for vesicle movement

and fusion with target membranes [31]. The Rho GTPase Cdc42 has been proposed as a

regulator of GPI-APs uptake [29]. During GPI-APs endocytosis, Cdc42 activation and the level of

cholesterol at the cell membrane are coupled, with concomitant recruitment of actin

polymerization machinery [32]. Following their internalization through a pathway that is

cholesterol-sensitive and dependent on Cdc42-activation, GPI-APs traffic to the recycling

endosomes and are recycled back to the cell surface, thus replenishing the pool of proteins that

may have been removed or degraded.

Many GPI-APs have roles in immune modulation [20, 24]. Among them is CD73, the

enzyme required for production of extracellular adenosine (EAD). EAD is produced in response

to cellular damage whereby the ATP released in the extracellular milieu is sequentially converted

to adenosine monophosphate (AMP) and then to adenosine by two surface ecto-nucleotidases,

CD39, a transmembrane protein and the GPI-AP CD73, respectively [33]. CD73 is crucial for host

resistance to infection by *S. pneumoniae* [34] as mice lacking this enzyme are highly susceptible

to infection and cannot control bacterial numbers [34]. We previously reported that the absence

of CD73 on neutrophils diminished their ability to kill pneumococcus [34], thus favoring bacterial

growth. Interestingly, the expression of CD73 is significantly reduced on neutrophils migrating into

the lungs in response to pneumococcal infection [35], indicating that pneumococcal infection may

result in the modulation of CD73 on host cell surfaces.

In this study, we report that infection with *S. pneumoniae* reduces CD73 surface

expression on host pulmonary epithelial cells. This reduction reflects a general effect of PLY on

the surface GPI-AP expression and depends on its pore forming capacity. The reduction in CD73

expression was not due to its release from the cell surface but rather due to internalization of GPI-

APs in a RhoA- and Cdc42-dependent manner. Internalization of GPI-APs was associated with

reduced host cell damage in response to PLY. This study identifies a possible mechanism through

which repair of damage caused by S. *pneumoniae* helps the bacteria potentially escape the host

immune response.

Results

Infection with *S. pneumoniae* reduces the expression of GPI-APs on the surface of lung epithelial cells.

We previously found that the EAD-producing enzymes CD73 and CD39 were crucial for host resistance against pneumococcal pneumonia [34, 36, 37]. To study the effect of *S. pneumoniae* infection on the expression of CD73 and CD39, we intra-tracheally (i.t.) infected C57BL/6 mice with *S. pneumoniae* TIGR4 strain or mock-treated with PBS. Forty-eight hours post infection (hpi), the lungs were harvested and the expression of CD73 and CD39 on total lung cells quantified using flow cytometry. *S. pneumoniae* significantly reduced the percentage of lung cells that express CD73 compared to uninfected controls, while the expression of CD39 remained unchanged (Fig 1A). In addition, a reduction in the amount of surface CD73 was detected on lung epithelial (EPCAM⁺) cells (Fig 1B).

To further explore the mechanisms involved, we modeled the decreased expression of CD73 on lung epithelial cells *ex vivo* using H292 cells, a human pulmonary epithelial cell line extensively used to study the interaction between the host factors and respiratory pathogens [38-41]. Infection of H292 cells with increasing numbers of *S. pneumoniae* TIGR4 resulted in a dose-dependent decrease in the expression levels of CD73 (Fig 1C), mimicking the *in vivo* finding.

These data indicate that *S. pneumoniae* infection specifically regulates the expression of CD73, a GPI-AP, but does not affect the levels of CD39, a transmembrane protein [42]. We thus tested if infection with *S. pneumoniae* selectively targets the surface expression of GPI-APs in general. H292 cells infected with *S. pneumoniae* TIGR4 were labelled with antibodies recognizing the GPI-APs CD55 and CD59 or stained with fluorescently labeled inactive variant of the protein aerolysin (FLAER), which selectively binds GPI anchors at the cell membrane [43]. Flow cytometric analysis showed that *S. pneumoniae* infection significantly reduced the expression of both CD55 and CD59, as well as FLAER staining intensity (Fig 2A). This reduction was not

common to all surface proteins, as *S. pneumoniae* infection increased the expression of

transmembrane proteins such as CD47 and CD54 (Fig 2B). These findings indicate that

pneumococcus specifically reduces the surface expression of GPI-APs.

In vivo, epithelial cells are polarized, and GPI-APs mainly localize at the apical domain

[44]. To determine if the pneumococcus-induced reduction of GPI-APs surface exposure also

occurs on polarized epithelial cells, H292 cells were grown as polarized monolayers on Transwell

filters [39, 45]. The cells were then apically infected with *S. pneumoniae* TIGR4 for an hour and

the expression of GPI-APs was analyzed through flow cytometry. The expression levels of the

GPI-APs CD73, CD55 and CD59 were significantly reduced in infected cells as compared to the

uninfected controls (Fig S1). Overall, these data show that infection with *S. pneumoniae* reduces

the surface expression of GPI-APs on lung epithelial cells.

S. pneumoniae-induced reduction of GPI-APs is dependent on the pore forming ability of

PLY.

To determine if the reduced surface exposure of CD73 was strain dependent, H292 were

infected with D39, a serotype 2 strain of *S. pneumoniae*. Like TIGR4 infection, infection with D39

significantly reduced the expression of CD73 on H292 cells (Fig S2A), suggesting that the effect

of *S. pneumoniae* on CD73 surface expression is not strain specific. Infection of H292 cells with

Δ*cps* TIGR4, lacking the polysaccharide capsule, also reduced surface CD73 similarly to wild-

type TIGR4 (Fig S2A), thus indicating that the reduction of CD73 expression is independent from

the capsule. However, we found that heat-killed *S. pneumoniae* TIGR4 not only failed to reduce

CD73 expression on H292 cells but rather increased it (Fig S2B). This indicates that a heat-labile

bacterial factor is involved in reducing the expression of CD73 on lung epithelial cells.

S. pneumoniae pneumolysin (PLY) is heat-labile [12, 13], so we tested if the reduction in

CD73 is PLY-dependent. H292 cells infected with Δ*ply* TIGR4 did not show diminished CD73

expression on the cell surface compared to the uninfected controls (Fig 3A). Moreover, infection

of H292 cells with *Bacillus subtilis* expressing *ply* (*B. subtilis* PLY) significantly reduced levels of CD73 as compared to cells challenged with wild-type *B. subtilis* (Fig 3A). To test if PLY alone is sufficient for reducing the expression of CD73, H292 cells were treated with increasing concentrations of purified recombinant PLY (rPLY). The expression of CD73 was reduced upon incubation with rPLY in a dose-dependent manner (Fig 3A). Altogether, these findings demonstrate that PLY is required and sufficient for decreasing CD73 from the cell surface of pulmonary epithelial cells.

To determine if pore formation is required to decrease CD73 expression on host cell surface, we tested PLY mutants defective at different stages of pore formation. 2.5 µg/ml of wild-type PLY elicited more than 3-fold reduction in CD73 expression (Fig 3A and B) but the same concentration of PLY ^{T456G}, which is incapable of binding cholesterol [46], PLY $^{M-1}$, which is locked at an early stage of pore formation [11, 15, 47], or PLY $^{P-1}$, which is locked at late stage [11, 15, 47], failed to induce any change in CD73 expression on lung epithelial cells (Fig 3B). These results indicate that the phenotype observed requires the pore-forming ability of PLY.

As PLY pores may result in the death of lung epithelial cells during infection [48, 49], we tested if the reduced expression of GPI-APs could be a consequence of cell death. We tracked cell death following infection with *S. pneumoniae* TIGR4 by measuring apoptosis and found that infection resulted in rapid apoptosis of H292 cells (Annexin$^+$) as early as 10 minutes post-infection (Fig S3A). This depended on PLY, as infection with Δ*ply* *S. pneumoniae* TIGR4 was not associated with a significant effect on cell viability (Fig S3A). However, flow cytometric analysis showed that infection with *S. pneumoniae* induced a similar significant 2-fold reduction in the intensity of FLAER on both apoptotic (Annexin$^+$) and non-apoptotic (Annexin$^-$) cell populations compared to the uninfected control groups (Fig S3B), demonstrating that decrease in GPI-APs is not a result of cell death. We further confirmed that PLY-dependent reduction in GPI-APs could be observed at this early 10-minute timepoint, as 10-minutes post-infection of H292 cells with

wild-type but not Δ*ply* TIGR4 resulted in significant reduction of FLAER binding (Fig 3C).

Altogether, these data indicate that *S. pneumoniae*-induced reduction in GPI-APs on host lung

epithelial cells is PLY-mediated.

Cholesterol-dependent-cytolysins reduce the expression of GPI-APs at the surface of lung

epithelial cells.

PLY belongs to a large family of cholesterol-dependent-cytolysins (CDC) that share

structural and functional properties [12]. To determine if the PLY-mediated reduction of GPI-APs

expression is a common feature shared among different members of CDCs, H292 cells were

challenged with two other CDCs: Streptolysin O (SLO) and Intermedilysin (ILY), produced by

Group A *Streptococci* and *Streptococcus intermedius*, respectively [50]. As pore formation is

required for reduction of GPI-AP expression, we treated H292s at concentrations sufficient to

induce equal levels of pore-formation, but that did not result in H292 cell death [11]. We found

that incubation of cells with SLO and ILY with 5 units of pore forming activity caused a reduction

in the FLAER intensity like that induced by PLY (Fig 3D), indicating that the decrease of GPI-AP

expression is a feature common to different CDCs.

GPI-APs are not released from the cell surface upon infection with *S. pneumoniae*.

GPI-APs can be removed from the host cell surface through cleavage by

phosphatidylinositol phospholipase C (PI-PLC) [51]. In fact, *S. pneumoniae* that produces the

pneumococcal surface protein C (PspC) can activate PI-PLC [52]. However, we found that PspC-

deficient *S. pneumoniae*, which are incapable of activating host PI-PLC, showed no defect in

triggering a reduction of surface CD73 (Fig S4A). In addition, treatment of H292 cells with the

phospholipase C specific inhibitor U-73122 (or its inactive analogue U-73343 [53]) did not

abrogate the *S. pneumoniae*-mediated reduction in CD73 surface expression (Fig S4B). These

findings indicate that reduction in cell surface associated CD73 upon *S. pneumoniae* infection

does not depend on the activity of host PI-PLC.

As part of a membrane repair mechanism, host cells infected with *S. pneumoniae* can

release extracellular membrane vesicles through which they expel membrane-bound bacterial

products such as PLY [54, 55]. To determine if CD73 was shed in host-derived vesicles, H292

cells were treated with CellTracker Green dye CMFDA, which stains the cell cytoplasm and can

be used to track host-derived vesicles. Following challenge with *S. pneumoniae* TIGR4, cell free

supernatants were collected, stained with FLAER and the presence of vesicles simultaneously

positive for FLAER and CMFDA was analyzed through flow cytometry. Surprisingly, we observed

a significant reduction in the percentage of vesicles FLAER$^+$ CMFDA$^+$ released by H292 cells

infected by *S. pneumoniae* compared to the uninfected controls (Fig S5), suggesting that the

decrease of GPI-APs induced by *S. pneumoniae* infection is not due to the release of extracellular

vesicles by the infected cells. As injured cells can also release parts of injured membranes via

the formation of protrusions named blebs [56, 57], we tested the effect of treating H292 cells with

blebbistatin, a drug reported to inhibit blebbing [58]. We found that treatment with blebbistatin did

not affect the levels of GPI-APs (Fig S6). Collectively, these data indicate that the reduction in

GPI-APs following pneumococcal infection is not due to their release from the cell surface by the

activity of phospholipases, shedding in vesicles or blebs.

Infection with *S. pneumoniae* triggers the internalization of GPI-APs.

As we found that the decreased levels of GPI-APs at the cell surface are not related to

their extracellular release, we tested if infection of lung epithelial cells by *S. pneumoniae* triggers

their internalization. To do so, H292 cells were infected with *S. pneumoniae,* stained with FLAER

to identify GPI-APs, and imaged by confocal microscopy. In conditions where H292 cells were

treated with HBSS (uninfected controls) or were challenged with Δ*ply S. pneumoniae*, GPI-APs

were uniformly detected on the surface of the cells (Fig 4A), following the outline of the cell

membrane. However, in *S. pneumoniae* infected cells, GPI-APs lose their uniform association to

the cell cortex and appear in clusters inside cells (Fig 4A, white arrows). These data suggest that

infection with *S. pneumoniae* triggers the internalization of GPI-APs.

If GPI-APs are removed from the cell surface by internalization (vs. extracellular

shedding), the total amount of GPI-AP associated with cells is predicted to be unchanged. To test

that, H292 cells were stained with FLAER before infection by *S. pneumoniae* ("PRE"). After 1 h,

cell-associated GPI-APs was quantitated by flow cytometry. Indeed, we found that FLAER

staining remained unchanged, i.e., was indistinguishable from FLAER staining of uninfected cells

(Fig. 4B, "PRE"). To confirm that under these experimental conditions, *S. pneumoniae* resulted in

the removal of GPI-APs from the cell surface, in parallel we stained H292 only after the 1 h

infection ("POST"). As predicted, surface staining by FLAER after infection revealed decreased

surface GPI-APs (Fig. 4B, "POST"). These findings confirm that infection with *S. pneumoniae*

induces the internalization of GPI-APs.

To test whether PLY was sufficient to trigger this process, as well as to follow GPI-AP

internalization in living cells, HeLa cells (chosen due to their ease of imaging by microscopy) were

pre-labelled with FLAER and then challenged with purified PLY or mock-treated. FLAER-stained

GPI-APs were visualized by fluorescent microscopy for 45 minutes (Supplementary video 1).

FLAER intensity was rapidly reduced on the cell surface upon challenged with PLY (Fig 4C and

Supplementary video 1), in addition FLAER signal was detected intracellularly within 3 minutes of

incubation with PLY (Fig 4C and Supplementary video 1) in structures resembling intracellular

vesicles (Fig 4C and Fig S7). At 45 min after PLY incubation FLAER signal at the surface of the

cells appeared to increase (Figure 4C and S7), suggesting that internalized GPI-APs are recycled

back to the cell surface. Overall, these data show that the infection with *S. pneumoniae* triggers

the internalization of surface GPI-APs which are then found intracellularly in vesicle-like

structures.

Internalization of surface GPI-APs is mediated by PLY-induced activation of host Rho GTPases RhoA and Cdc42.

To explore the mechanism behind *S. pneumoniae*-mediated internalization of GPI-APs, we first assessed a possible role of the actin cytoskeleton, given that dynamic changes in the actin machinery are often required for endocytosis [59]. H292 cells were pre-treated with Cytochalasin D, an inhibitor of actin polymerization, at 2 μM, a concentration that prevented the actin-dependent cell-cell spread of *Listeria monocytogenes* ([60] and data not shown). Treatment with Cytochalasin D did not affect the reduction of FLAER intensity following *S. pneumoniae* infection (Fig S8A), suggesting an actin-independent mechanism for the internalization of GPI-APs. Similar results were observed when Cytochalasin D pre-treated H292 cells were challenged with PLY or SLO (Fig S8B), indicating that other CDCs also induce the internalization of GPI-APs through a mechanism independent of actin polymerization.

The host Rho family of guanosine 5'-triphosphate (GTP)-binding proteins (GTPases) are known to play important roles in protein trafficking, receptor-mediated endocytosis, and internalization of GPI-APs [61]. To examine the role of Rho GTPases in the internalization of GPI-APs induced by *S. pneumoniae* infection, H292 cells were treated either with *Clostridium difficile* toxin B (ToxB), which irreversibly inactivates Rho GTPases by glucosylation [62], or with vehicle control, before infection with *S. pneumoniae*. While infection with *S. pneumoniae* in vehicle-treated cells caused the expected reduction in FLAER intensity indicating the decreased surface expression of GPI-APs (Fig 5A), cell surface expression of GPI-APs was not affected in cells pre-treated with ToxB (Fig 5A), suggesting a role for Rho GTPases in PLY-mediated GPI-APs internalization. To confirm the role of Rho GTPases in the internalization of GPI-APs, we knocked down their expression using RNAi. H292 cells were transfected with siRNAs targeting Cdc42, RhoA and Rac-1 or with scrambled non-targeting siRNA as control. At 48 h post transfection, cells were challenged with *S. pneumoniae* at a MOI of 10 for 1 h and FLAER intensity was quantified

through flow cytometry. Depletion of Cdc42 and RhoA but not Rac-1 abrogated the expected

reduction in FLAER intensity usually observed in H292 cells infected by *S. pneumoniae* (Fig 5B).

This indicates that the internalization of pneumococcal-mediated GPI-APs depends on the Rho

GTPases Cdc42 and RhoA.

Rho GTPases were previously reported to be activated by PLY in neuronal cells [13, 63].

We assessed if Cdc42 and RhoA are activated by PLY in epithelial cells in the time frame of PLY-

mediated internalization of GPI-APs. We transfected HeLa cells with plasmids encoding the

fluorescence resonance energy transfer (FRET) probes (CFP/YFP FRET pairs) that reflect the

activation state of RhoA or Cdc42. Each probe comprises a truncated form of the GTPase and

the Rho-binding domain (RBD) of effectors, bound to the CFP or YFP, respectively. The binding

of GTP to the GTPase (resulting in activation) brings the YFP-bound GTPase domain and the

CFP-bound RBD into close proximity, allowing FRET from CFP to YFP [55, 56]. Transfected HeLa

cells were challenged with 0.25 nM of PLY and live-cell microscopy was performed to visualize

the activation status of RhoA and Cdc42. FRET ratio (FRET/CFP) was calculated specifically at

the cell cortex using Image J. Cells were imaged for 20 min before PLY was added and for

additional 10 min after PLY challenge. For both probes, the ratio of FRET/CFP remained stable

during the 20 min before PLY addition, showing that the levels of GTPase activation do not vary

in these conditions (Fig 5C). Shortly after the addition of PLY, we observed a modest but

consistent increase in the FRET/CFP ratio levels. The increase detected for the Cdc42 probe was

very rapid and of short duration (about 40 s), while the increase for the RhoA probe appeared

more sustained in time (140s) (Fig 5C). These data suggest that PLY triggers a fast and transient

activation of both RhoA (Supplementary video 2) and Cdc42 (Supplementary video 3) at the cell

cortex. The ratio of FRET/CFP, for both probes, rapidly decreased to values that tended to be

lower than those obtained before PLY incubation (Fig 5C), indicating that although the activity of

RhoA and Cdc42 is triggered during the very early response to PLY induced damage, it is rapidly

shutdown. Overall, these data indicate that PLY induces internalization of GPI-APs through the

rapid, spatially, and temporally regulated activation of the Rho GTPases RhoA and Cdc42.

GPI-APs are not required for PLY-mediated pore formation.

The PLY-related CDC ILY utilizes the human GPI-AP CD59 as a receptor to induce

oligomerization and pore formation [64] and the cytolytic pore forming toxin aerolysin binds the

GPI moiety of the GPI-anchored T-lymphocyte protein Thy-1 [19]. To test whether PLY requires

the expression of GPI-APs on the host cell surface to form pores, H292 cells were either treated

with recombinant PI-PLC to cleave GPI-APs from the cell surface [51], or mock-treated with PBS.

As expected, treatment of cells with PI-PLC caused a significant reduction in the percentage of

GPI$^+$ cells (FLAER$^+$) compared to the untreated control group (Fig S9A). Upon addition of PLY to

H292 cells, we detected the expected decrease in GPI$^+$ cells compared to the untreated control

group (Fig S9A). We did not see a further reduction in the number of cells expressing GPI-APs

when PLY was added to PI-PLC pre-treated cells (Fig S9A).

We then assessed whether reduced expression of GPI-APs impact PLY-mediated pore

formation using PI permeability assays. We observed that PLY challenge induced a 4-fold

increase in the number of permeabilized cells (PI$^+$) in both the untreated and PI-PLC pre-treated

conditions in comparison to the respective baseline controls (Fig S9B). Finally, the average

intensity (MFI) of PI staining was unaltered by PI-PLC treatment. These data indicate that PLY

does not require surface GPI-APs to bind and form pores on host cell membranes.

Internalization of GPI-APs is associated with lower levels of PLY-induced cellular damage.

PLY forms pores in cholesterol rich domains, where GPI-APs are also colocalized [12].

Depending on the extent of damage, host cells can repair the damaged membrane and revert to

homeostasis [65]. To couple the kinetics of the PLY-mediated removal of GPI-AP from the cell

surface with an analysis of the population of cells that have undergone this reduction, H292 cells

were challenged with *S. pneumoniae* or mock-treated with PBS for different time periods and the

kinetics of reduction in FLAER intensity was determined throughout infection. According to our

previous live-cell imaging experiments of cells challenged with rPLY (Fig 4C), we expected the

reduction to occur rapidly. Indeed, *S. pneumoniae* significantly reduced FLAER intensity within

10 minutes of infection compared to the PBS-treated cells and retained this reduced level for the

remainder of the 90-minute observation period (Fig 6A). To analyze the specific population of

cells that reduced GPI-AP, we utilized flow cytometry and found that by 10 minutes after infection

approximately 60% of the infected cells showed reduced levels of GPI-AP (termed "GPI^{low}" cells)

(Fig 6B), a percentage that also remained stable through the rest of the experiment.

To test if the reduction of GPI-APs from the cell surface affected cellular viability, we

measured the staining of the population of GPI^{low} or GPI^{high} H292 cells by Annexin V, which stains

apoptotic cells, and PI, which stains cells with permeable membranes. Interestingly, we found

that, throughout infection, the population of infected GPI^{high} H292 cells had a significantly higher

percentage of membrane damaged (PI$^+$) (Fig 6C) and apoptotic (Annexin V$^+$) cells than the

population of GPI^{low} cells (Fig 6D). These findings suggest that reduced GPI-APs surface

expression is associated with reduced levels of PLY-induced damage. Altogether, these findings

suggest that cells with reduced expression of GPI-APs are more resistant to PLY-induced toxicity

and that the internalization of GPI-anchored proteins seems to be part of membrane repair that

limits the extent of PLY-induced cellular damage.

Discussion

The outcome of host-pathogen interaction is proposed to be mediated by the damage response [66, 67]. This damage can be caused by the invading organism and/ or the host's immune response to infection. *S. pneumoniae*, a typically asymptomatic colonizer of the host upper respiratory tract, can cause serious and life-threatening infections when it gains access to the lungs and other organs and this is directly linked to its ability to cause organ damage and impair organ function [68-71]. One key component of this is the bacterial toxin PLY, found in all the clinical isolates of *S. pneumoniae* [72]. PLY induces host cell damage by making pores in cells [71]. However, host survival can be contingent on repairing/controlling infection-induced damage [73]. In this study, we found that PLY-induced pores are associated with the rapid removal of GPI-APs from the host cell surface, which can have profound effects on host response to infection. PLY activated host RhoGTPases RhoA and Cdc42, resulting in the internalization of GPI-APs (Figure 7). This internalization of GPI-APs was associated with reduced cellular damage thus, suggesting the induction of host repair mechanisms.

GPI-APs are a heterogenous group of proteins which lack transmembrane domains and are dependent on the GPI moiety for their expression on the cell surface [24]. Through their interactions with the sphingolipids and cholesterol in the lipid bilayer, the GPI moieties anchor the proteins in lipid-rich microdomains or membrane rafts [24, 74]. We found here that removal of GPI-APs from the cell surface was not only mediated by PLY, but also occurred upon treatment of host cells with other cholesterol dependent cytolysins (CDCs) including SLO and ILY. CDCs bind to the cholesterol moieties in the host cell surface through the undecapeptide domain of the peptide to mediate their cytolytic activity [14, 64]. Binding of the soluble monomeric form of the CDCs to the cholesterol-rich membrane results in a sequence of structural changes causing oligomerization of the monomers with subsequent formation of transmembrane β-barrel that results in pore formation in the host cell membrane [64, 75]. Therefore, CDCs and GPI-APs will

likely localize together in cholesterol-rich microdomains at the cell surface where the damage

occurs. Consequently, internalization of GPI-APs following CDC-induced damage may be part of

membrane repair. Prior work with SLO showed that pores formed by the toxin are endocytosed

as part of the membrane repair process [76]. Here, we found that removal of GPI-APs by PLY

was dependent on the pore forming ability of the toxin and that internalization of GPI-APs was

associated with reduced cellular damage, suggesting that the host can repair the damage at the

membrane.

The type and extent of host cell damage as well as the host response to CDCs is highly

concentration dependent [65]. At high concentration levels, PLY induces irreversible loss of cell

membrane integrity and cellular homeostasis, while at lower concentrations, it activates host cell

repair mechanisms to restore their membrane integrity and to survive [65]. Initiation of host cell

repair of the plasma membrane depends on a limited influx of Ca^{2+} that triggers cytoskeletal

rearrangements to plug the holes in the plasma membrane [13, 77]. This has been described to

be mediated by the activation of host GTPases, re-localization of host intracellular organelles and

recruitment of cytoplasmic Ca^{2+} responsive proteins including annexins that interact with the

plasma membrane phospholipids at the injury site [13, 77, 78]. Host Rho GTPases, including Rho,

Rac and Cdc42, play a major role as molecular switches that regulate key cellular signaling

pathways and link surface receptors to the downstream processes culminating in endocytosis,

exocytosis and actin polymerization that affect cell polarity, migration, cytokinesis and vesicle

trafficking [79]. While exploring the cellular mechanisms that drive GPI-APs reduction at the cell

surface, we found that removal of GPI-APs from the cell surface upon *S. pneumoniae* infection

was dependent on Cdc42 and RhoA while Rac1 was not involved in this process. These data fit

with what has been reported in the literature regarding the role of Rho GTPases in the turnover

of GPI-APs under homeostatic conditions. GPI-APs are normally recycled from the cell surface

through an endocytic process called potocytosis involving caveolae, which is independent of both

dynamin and clathrin-mediated pathways [29, 80]. Once internalized, GPI-APs within the cell can

be seen in tubular structures called GPI-enriched endosomal compartments [80, 81].

Internalization of GPI-APs via endocytosis is under the regulation of host GTPases [80, 81]. This

includes CD59, one of the GPI-APs for which we observed reduction in surface expression upon

infection, whose internalization via endosomes is reported to be regulated by Cdc42 [82].

Additionally, perturbation in cholesterol moieties in the lipid bilayer also leads to internalization of

GPI-APs via clathrin-and dynamin-independent endocytic pathway [80]. Therefore, CDC binding

to the cell surface and pore formation results in the perturbation of the cholesterol rich

microdomains where GPI-APs are localized resulting in initiation of the endocytic pathway which

may lead to the repair of the injured membrane.

One of the interesting findings of this study was that actin polymerization had no apparent

role in the internalization of GPI-APs despite the activation of Cdc42 and RhoA. Cdc42 is a key

regulator of actin polymerization through the regulation of N-WASP and Arp2/3 complex and

similarly, RhoA also regulates actin polymerization [79]. Several recent reports have described

actin-independent endocytic events in various cell types. Work on non-adherent K562 human

erythroleukemic cells or adherent Cos-7 cells showed a lack of actin polymerization role in the

endocytosis process in these cells [83]. In another study, treatment of A431 cells (adherent and

in suspension) with cytochalasin D did not affect the process of endocytosis indicating an actin-

independent mechanism of endocytosis in these cells [83]. Additionally, the rational that the

polymerization of actin is important to generate the mechanical force required for vesicle fission

during clathrin-mediated endocytosis was recently challenged. In mouse chromaffin cells,

inhibition of dynamin GTPase but not of actin polymerization resulted in changes in the fission-

pore conductance in the endocytic membranes indicating that the pinching off of the endocytic

vesicle from the plasma membrane is actin-independent [84]. The requirement of actin in

endocytosis could be spatially regulated as in polarized cells inhibition of actin polymerization

blocked clathrin-mediated endocytosis from the apical but not the basolateral surface of polarized

Madin-Darby Canine Kidney Cells (MDCK) [85]. This study observed PLY-mediated reduction in

GPI-AP expression in pulmonary epithelial cells from infected murine models as well as in non-

polarized and polarized H292 cells. Overall, our findings point to a signaling pathway whereby

PLY activated Cdc42 and RhoA initiate internalization of GPI-APs independent of actin

polymerization machinery. Other bacterial pathogens have evolved to use actin polymerization-

independent mechanisms to trigger host responses. For example, *Salmonella enterica* serovar

Typhimurium was reported to activate Rho kinases to mediate entry into the cell by activating

myosin II independent of actin polymerization [86]. Similarly, *Listeria monocytogenes* was shown

to manipulate host exocytosis independent of actin polymerization to enter the host cell [87].

Apart from stimulating membrane repair through annexins, injury to plasma membrane by

PLY, LLO or SLO also results in a phenomenon called shedding whereby the host cell releases

extracellular vesicles to not only remove the injured part of the membrane but also to remove the

membrane bound toxin [54, 88]. Blebbing is one of the ways to mediate the shedding of larger

vesicular structures whereby the cell undergoes actomyosin-dependent localized contraction in

the plasma membrane to create confined spaces or a localized bulge with elevated Ca^{2+} levels to

prevent leakage to cytoplasmic contents and proteins in the extracellular milieu [13, 65]. These

localized bulges in the plasma membrane are eventually released by the cells either to remove

the areas of pores in case of a low degree of plasma membrane injury or to initiate the processes

of regulated apoptosis and inflammation in case of extensive cell damage [56, 65]. Since shedding

and blebbing result in the loss of part of the plasma membrane, it could also result in the loss of

expression of GPI-APs. However, data presented here using CMFDA dye to track the vesicles

and using chemical inhibition of blebbing showed that this was not the case. Surprisingly, we

observed a significant reduction in the percentage of vesicles FLAER+ CMFDA+ released by

H292 cells infected by *S. pneumoniae* compared to the uninfected controls. We also found no

evidence for the enzymatic release of GPI-APs from the cell surface to the extracellular milieu.

Certain bacterial species including *S. pneumoniae* have evolved to induce the expression of host

lipolytic enzyme phospholipase [89]. Given that the anchors of GPIs interact with the lipid moieties

of the plasma membrane, the GPI-APs are therefore susceptible to cleavage by phospholipases

[20, 21]. The findings here with phospholipase C specific inhibitor U-73122 and using a Δ*pspC S.*

pneumoniae strain showed that the reduction in CD73 expression following PLY treatment was

not a consequence of cleavage of GPI-APs by the host phospholipase.

The removal of GPI-APs from the host cell surface can seriously affect the host's ability to

control infection. These GPI-bound proteins are essential for many host physiological functions

including but not limited to enzymatic activities, nutrient uptake, complement regulators, immune

antigens and adhesion molecules [20, 28, 34, 90, 91]. This work shows that treatment with PLY

reduces CD55, CD59 and CD73 levels on pulmonary epithelial cells, all of which are GPI-APs

and important regulators of host inflammatory response. CD73 is an ectonucleosidase that is

required for initiating host responses to damage [42]. The host can sense infection-induced

damage by recognizing several damage-associated molecular patterns (DAMPS), including direct

recognition of extracellular ATP and its breakdown product adenosine in the extracellular

environment [92]. CD73 is required to dephosphorylate ATP, which has leaked from damaged

cells, into adenosine [93]. We previously found that extracellular adenosine production by CD73

was required for host resistance against *S. pneumoniae* in part by modulating epithelial responses

to these bacteria [34-36]. Other groups have also found that CD73 expression on epithelial cells

is important in other infections where CD73 expression on intestinal epithelial cells was needed

to control *Salmonella enterica* serovar Typhimurium infection [94]. Apart from CD73, both CD55

and CD59 act as inhibitors of complement membrane attack complex (MAC) [95, 96] and have a

role in neutrophil trans-epithelial migration [97]. Therefore, PLY's downregulation of CD55 and

CD59 could also have important implications on host immune responses. In summary, our

findings suggest that repair mechanisms activated in response to infection-induced cellular

damage may inadvertently help *S. pneumoniae* and other bacterial pathogens evade host

defenses, reflecting the dual function of PLY during infection.

Materials and Methods

Ethics Statement: Mice were handled in accordance with the recommended Guide for the Care

and Use of Laboratory Animals published by the National Institutes of Health. All procedures were

done according to the protocol approved by the University at Buffalo Institutional Animal Care and

Use Committee (IACUC), approval number MIC33018Y.

Mice: All the animal work was performed in 8-12 weeks C57BL/6J male mice purchased from

Jackson Laboratories (Bar Harbor, ME) and housed at a specific-pathogen free facility at the

University at Buffalo.

Mammalian cell lines and media: Human epithelial cervical cancer cell line HeLa (ATTC) and

pulmonary mucoepidermoid carcinoma cell line NCL-H292 (ATCC) were grown in Dulbecco's

modified Eagle's medium (ThermoFisher Scientific) and RPMI 1640 media (ThermoFisher

Scientific), respectively, supplemented with 10% fetal bovine serum (FBS) (ThermoFisher

Scientific) and 100 U penicillin/streptomycin (ThermoFisher Scientific). For all the experiments

involving lifting and/or seeding, the cells were washed twice with dPBS (Gibco) and treated with

Trypsin/EDTA (Gibco).

Bacterial strains and growth conditions: The following bacterial strains were used: *S.*

pneumoniae TIGR4 and D39 [98], PLY-deletion (Δ*ply*) [98], capsule-deletion (Δ*cps*) [99], and

pneumococcal surface protein C-deletion (Δ*pspC*) mutant strains of *S. pneumoniae* TIGR4 [100]

B. subtilis and its mutant expressing PLY (*B. subtilis* PLY) [101]. The pneumococcal strains were

grown to mid-log phase in Todd–Hewitt broth (BD Biosciences) supplemented with Oxyrase

(Fisher) and 0.5 % yeast extract at 37 °C/5 % carbon dioxide. Bacterial aliquots were then frozen

at –80 °C in growth media with 20 % glycerol. Tryptic Soy Agar plates with 5 % sheep blood

(Northeast Laboratory Services, P1100) were used to confirm the titers of *S. pneumoniae* in the

aliquots. Before infecting animals or H292 cells, bacterial aliquots were thawed and diluted in PBS

to the desired number. *B. subtilis* strains were grown overnight at 37 °C in Luria-Bertani (LB)

Broth.

Toxin purification: Purification of SLO (Streptolysin O), ILY (intermedilysin) and PLY (PLY) along

with its mutant forms was carried out as previously described [11, 47]. Briefly, the cultures of *E.*

coli BL-XL-1 Blue expressing polyhistidine-tagged toxins were grown in sterile TB broth (with

1:100 of overnight culture) grown at 37 °C with shaking at 200 rpm supplemented with 100 µg/mL

ampicillin. Expression of toxins was induced by the addition of isopropyl β-D-

thiogalactopyranoside (IPTG, Fisher Scientific) to a final concentration of 1 mM when the OD_{600}

of the culture reached 0.5-0.6 after which the cultures were allowed to grow overnight. This was

followed by centrifugation, resuspending and lysing bacterial pellets in PBS supplemented with

halt protease inhibitor cocktail (Thermo Fisher Scientific) using microfluidizer. The cell lysates

were centrifuged at 10000 g for 30 min at 4 °C and the clear supernatant was collected and

passed through Ni-NTA agarose column (Qiagen). Contaminating proteins were washed away by

passing a 20–120 mM imidazole gradient through the column while the bound toxins were eluted

using PBS containing 500 mM imidazole. The purity of the toxin was confirmed by running the

collected fraction on SDS-PAGE. The toxin-containing fractions were mixed with 10% (v/v)

glycerol and 5µM Tris(2-carboxyethyl) phosphine hydrochloride (TCEP) and flash-frozen on dry

ice and then shifted to −80 °C until use. *C. difficile* toxin B (Tox B) was purchased from Sigma.

Unless indicated, we used 0.25 nM of toxin for experiments with purified PLY as this concentration

corresponded to 1 pore forming Unit which causes PI uptake in only 50% of the cell population

and was sufficient to induce low-grade cellular damage.

Toxin activity assay: The activity of purified toxins was measured as previously described [11]. Briefly, aliquots of different toxins were thawed on ice and centrifuged at 14,000 rpm at 4 °C for 10 min to remove protein precipitates. Bradford assay (Bio-Rad) was used to measure the concentrations of toxins per the manufacturer's instructions. Toxins at different concentrations were added in wells of non-adherent 96-well plates containing approximately 1 x 10^5 H292 cells and incubated for 10 min at 37 °C. Following incubation, cells were stained with propidium iodide (PI) (1 mg/mL), incubated in dark at RT for 15 min and analyzed through flow cytometry. The toxin concentration that caused permeabilization of 50 % of the H292 cells (PI+) was referred as "1 unit" (1U) of toxin activity. Cells treated with 5 % Triton X-100 acted as a positive control while HBSS was used as negative control.

Murine infection studies: Mice anesthetized under isoflurane treatment were intratracheally (it) instilled with *S. pneumoniae* TIGR4 at a dose of 2 x 10^6 CFU, as described previously [34, 102]. At 48 h post-infection, the mice were euthanized, lungs were harvested and single-cell suspension was obtained by digesting the tissue in RPMI 1640 containing 10 % FBS and 2 mg/mL of type 2 collagenase (Worthington) and 30 µL/mL of 10 mg/mL DNase (Worthington) stock prepared in dH_2O as previously described [102].

Preparing heat killed *S. pneumoniae* TIGR4: Frozen aliquots of *S. pneumoniae* TIGR4 were thawed on ice, centrifuged at 1700 g for 5 min and resuspended in PBS in the original volume. The bacterial suspension was then incubated at 65 °C for 2.5 h. Bacterial killing was confirmed by plating the suspension on blood agar plate.

Preparation of polarized H292 monolayers and infection with *S. pneumoniae*: Polarized H292 monolayers were prepared as previously described [45]. Briefly, the inverted side of the filter membrane of 24-well Transwells with a pore size of 3 µm were coated with 70µL of 30µg/mL of sterile collagen solution (3 mg/mL of collagen stock diluted in 60% ethanol) and allowed to dry for 3 to 4 h at RT under sterile conditions. Approximately 1 x 10^6 H292 cells resuspended in RPMI

1640 media (with FBS and Pen/Strep) were added in the volume of 70µL on the coated filter

membrane and allowed to grow overnight at 37 ºC with 5 % CO_2. The following day, the Transwells

with cell were inverted into a sterile 24-well plate containing 1mL of H292 complete media and

cells were allowed to grow and polarize for a week at 37 ºC with 5 % CO_2. Cells were then washed,

infected apically with *S. pneumonia,* treated with trypsin and type II collagenase, stained for GPI-

APs, and analyzed by flow cytometry.

Antibodies, inhibitors, and other reagents: The antibodies used in flow cytometry studies to

measure the expression of specific GPI-APs included the following antibodies: anti-CD73 (AD2,

eBioscience), anti-CD39 (eBioA1, eBioscience), anti-CD55 (JS11, Biolegend), anti-CD47

(CC2C6, Biolegend) and anti-CD54 (HA58, Biolegend). The primary antibodies used for Western

blotting included anti-Rac1 (Proteintech) (1:1000), anti-Cdc42 (Cell Signaling) (1:1000), anti-

RhoA (Cell Signaling) (1:1000) and anti-GAPDH (MA5-15738) (1:5000). Horseradish peroxidase

(HRP)-conjugated secondary antibodies were purchased from Jackson Immunolabs. The

following anti-mouse antibodies were also used: Anti-CD326 (EpCAM) (G8.8, eBioscience), anti-

CD39 (24DMS1, eBioscience) and anti-CD73 (eBioTY/11.8, eBioscience). The inhibitors

Cytochalasin D, PI-PLC, PI-PLC inhibitor U-73122 and its control U-73343 (Invitrogen,

ThermoFisher Scientific) and Blebbistatin were purchased from Sigma and dissolved in DMSO

(unless indicated). As appropriate, control groups were treated with diluted DMSO, PBS or HBSS.

Detection of plasma membrane integrity: *S. pneumoniae* or PLY-induced cellular damage was

determined by using ANNEXIN/PI staining (BioLegend), per the manufacturer's protocol. Briefly,

infected H292 cells were centrifuged at 1200 rpm at 4 °C, the supernatants were discarded, and

pellets were re-suspended in 100 µL of Ca^{2+} free staining buffer (supplied with the kit). 2.5 µL of

FITC Annexin V and 5 µL of Propidium Iodide were used per sample. The samples were then

incubated for 15 min at room temperature (RT) in the dark and immediately analyzed by flow

cytometry.

FLAER labeling: Following challenge with different bacterial strains or toxins, H292 cells were centrifuged at 1200 rpm at 4 °C, the supernatants were discarded, and pellets were re-suspended in 50 µL of FLAER (FL2S, Cederlane) diluted in FACS buffer (HBSS Ca/Mg$^+$ containing 1% FBS) at the final dilution of 1:100. The cells were then incubated for 30 min on ice in the dark. Following incubation, the cells were washed twice with 100 µL of FACS buffer, resuspended in 300 µL of FACS buffer and analyzed by flow cytometry.

Flow cytometry: Fluorescence intensities for different GPI-APs, Annexin V and Propidium Iodide were measured on a BD Fortessa cytometer. At least 50,000 events for lung tissue and 10,000 events for *ex vivo* assays with H292 cells were captured for analysis. For the analysis of FLAER-positive cell vesicles (Figure S5), the events were captured by legacy MoFlo cytometer. For the data in Figure S9, the BD Accuri C6 flow cytometer was used to capture the events. All the analysis was done using FlowJo software.

RNA-interference: Approximately, 1.5 x 10^4 H292 cells were seeded in 24-well plates (Falcon) and cells were transfected the following day in serum free media using 100 nM of short interfering RNA (siRNA), Opti-mem (Invitrogen) and the lipid reagent LF2000 (Invitrogen) as described previously [1]. Serum was added on day 3 at a final concentration of 10% in the transfected wells. On day 4, the cells were challenged with *S. pneumoniae* TIGR4 strain or lysed for Western blotting to confirm protein depletion. SMARTpool siRNAs from Dharmacon were used to deplete Cdc42 (M-005057-01-0005), Rac1 (M-003560-06-0005) and RhoA (M-003860-03-0005). A non-targeting NTC siRNA (D-001210-02-05) was used as a control.

Western blotting: To confirm protein depletion, approximately 48 h post-transfection H292 cells were lysed on ice for 30 min in ice-cold RIPA buffer (1% Triton X-100, 0.25% sodium deoxycholate, 0.05% SDS, 50 mM Tris-HCl [pH 7.5], 2 mM EDTA, 150 mM NaCl, 1 mM phenylmethylsulfonyl fluoride, 1 mM sodium orthovanadate, 10 mg/liter each of aprotinin and leupeptin), as previously described [103, 104]. The cell lysates were centrifuged at 12,000 rpm at

4 °C and bicinchoninic acid kit (Pierce) was used to determine the protein concentration in the

samples. An equal amount of protein (20 µg) was run on Mini-PROTEAN TGX Stain-Free Precast

Gels (BioRad) SDS-polyacrylamide gel and transferred to polyvinylidene difluoride (PVDF)

membranes. The membranes were blocked in 3% BSA for 1h at RT and incubated overnight at 4

°C with primary antibodies diluted in TBS/T with 3 % BSA. The membranes were washed thrice

with TBST (5 min per wash) followed by incubation with HRP-tagged secondary antibodies for 1

h at RT. The bands were detected using ClarityTM Western ECL substrate (BioRad) on the

ChemiDoc XRS+ system (BioRad).

Measuring phosphatidylinositol-specific phospholipase C (PI-PLC) and PLY-induced cell

damage: H292 cells were seeded in a 6-well plate (5 x10^5 cells/well) and 20 h later were pre-

treated with PI-PLC (0.1U) prepared in (Opti-MEM supplemented with 10mM Hepes, 1mM EDTA

and 0.1%BSA) [105]. Cells were then challenged for 10 min with PLY at 0.25 nM diluted in HBSS

(or HBSS only in the untreated condition). Cells were then harvested, centrifuged at 1200 rpm for

5 min at 4 °C and incubated for 30 min on ice in 50 µL of FACS buffer containing 0.5 µL of FLAER

(1:100). Following incubation, cells were washed and centrifuged at 1200 rpm for 5 min at 4° C

and resuspended in FACS buffer. Just before flow cytometry analysis, propidium iodide (PI) was

added at 2 ug/mL and incubated for 1 minute. Then, cells were analyzed for FLAER and PI

staining in BD Accuri C6 flow cytometer. At least 20,000 events per sample were collected. Data

obtained were analyzed using FlowJo software (Version 10.8).

Immunofluorescence microscopy: Approximately, 2 x 10^5 H292 cells were grown in a complete

medium on 13 mm diameter coverslips in 6-well plates. The day after, the cells were washed and

challenged with *S. pneumoniae* in serum-free RPMI 1640 medium. The plate was centrifuged at

1000 rpm for 2 min to enhance contact between bacteria and H292 cells and then incubated for

1 h at 37 °C in 5 % CO_2. Cells were washed in PBS, fixed in PBS with 4 % paraformaldehyde,

quenched with 20 mM NH4Cl (20 min), permeabilized with 0.1 % Triton X-100 (5 min) and blocked

with 3 % BSA in PBS (1 h). The antibodies were then diluted in PBS containing 1% BSA and cells

were labeled for GPI-AP nucleus and F-actin. GPI-AP using FLAER-Alexa 488 (Cedarlane)

(1:50), DAPI (Sigma) (1:1000) and Alexa fluor 680 phalloidin (Life technologies) (1:50),

respectively as described previously [106]. Cells were then washed and mounted with Vectashield

(Vector laboratories) and imaged using a 63 x 1.4 NA objective lens on a confocal laser-scanning

microscope (Leica SP8) and processed using ImageJ software.

RhoA and Cdc42 activation and FRET: HeLa cells seeded in Ibitreat 8-well dishes (4 x 10^4

cells/well) (Ibidi) were transfected approximately 20 h later with 300 ng of plasmid encoding FRET

probes (CFP/YFP FRET pairs) for RhoA or Cdc42 using jetPrime transfection reagent (Polyplus)

and incubated for 16–20 h to allow the expression of the recombinant proteins. FRET probes

used were described in [107, 108]. Cells were then kept in Opti-MEM (Gibco, without phenol red)

at 37 ℃ in 5 % CO_2 and imaged using Nikon TI microscope coupled with IRIS9 camera and using

60 x 1.4 NA objective. Cell imaging was started 20 min before PLY (0.25 nM) addition and

continued for 10 min after challenge with the toxin. CFP/FRET/YFP fluorescence data were

acquired after every 20 s. Image sequences were analyzed using ImageJ and FRET/CFP signal

ratio were calculated only at the cell cortex using a custom-made macro. A total number of 79

cells from 2 independent experiments were analyzed for RhoA activation and 74 cells were

analyzed from 2 independent experiments for Cdc42 activation.

Live cell imaging and signal quantification: HeLa cells were seeded in 8-well Ibidi plates

(1x10^5 cells/well) and 20 h later stained with FLAER (1:100) for 30 min. Cells were then imaged

using a Nikon Ti microscope coupled with IRIS9 camera with 40x 0.95 NA objective, for 15 min

with a time interval of 30 s before the addition of PLY (0.25 nM diluted in HBSS) or HBSS (used

as untreated control), and then for additional 45 min post PLY addition. Quantification of the

FLAER signal was obtained by image analysis using ImageJ. A total of 53 cells were analyzed.

The mean fluorescence intensity (MFI) was determined overtime in randomly selected regions of

interest (ROIs) at the cell membranes. These values were then normalized to the mean MFI

obtained from the same cells before these were challenged with PLY or HBSS.

Statistical analysis: Graph Pad Prism9 was used to perform all the statistical analysis. All the

data graphs represent the mean values +/- standard deviation (SD). Significant differences were

determined using One-way ANOVA followed by Tukey's or Dunnett's multiple comparison test or

Student's t-test or one sample t and Wilcoxon test, as indicated. p values of 0.05 or lower were

considered significant.

Acknowledgements

We want to thank Mafalda Sousa (ALM facility, i3S) for developing the macro for calculation of

the FRET/CFP signal ratio.

Conflict of interest

The authors have declared that no competing interests exist.

Author contributions

NL and MB conducted research, analyzed data, and wrote paper. JMP, SER, AL, MP, GA, and

WA conducted research and analyzed data. AK and RKT provided valuable reagents and

feedback. SS and JML designed the research and wrote the paper. ENBG designed research,

performed experiments, and wrote the paper. JML, SS and ENBG had responsibility for the final

content. All authors read and approved the final manuscript.

Figures and Legends

Figure 1: Pulmonary infection by *S. pneumoniae* significantly reduces CD73 expression on

the surface of lung epithelial cells. C57BL/6 (8-weeks old) male mice were infected intra-

tracheally with 2×10^6 CFU of *S. pneumoniae* TIGR4 strain or mock infected with PBS. Mice were

euthanized 48 hpi, the lungs harvested and digested into a single cell suspension, and surface

stained for CD73 and CD39. Flow cytometry was used to determine (A) the percentage of CD73$^+$

and CD39$^+$ total lung cells and (B) the expression (mean fluorescent intensity or MFI) of CD73 on

lung epithelial cells (gated on EPCAM+ cell population). The data shown are pooled from 4-5

mice per group with each box representing one mouse. *, $p < 0.05$ by Student's t-test indicate

significant differences from uninfected controls. (C) H292 cells were challenged with *S.*

pneumoniae TIGR4 strain at indicated MOIs for 1 h and MFI of surface exposed CD73 was

measured by flow cytometry. The panel on the right shows a representative histogram showing

the intensity of CD73 labeling comparing uninfected and infected conditions. Data are pooled from

four independent experiments. *, $p < 0.05$ by One-way ANOVA followed by Dunnet's test.

A. GPI-APs

B. Trans-membrane proteins

☐ Uninfected ■ + *S. pneumoniae*

Figure 2: ***S. pneumoniae*** **infection of lung epithelial cell lines decreases surface associated**

Glycosylphosphatidylinositol anchored proteins (GPI-APs). H292 cells were infected with *S.*

pneumoniae TIGR4 strain at an MOI of 10 for 1 h. Flow cytometry was used to measure the

expression (MFI) of (A) CD55 and CD59, GPI-APs detected using specific antibodies, and of bulk

GPI-APs detected by using the fluorescent-labeled mutant of aerolysin (FLAER) or (B) CD47 and

CD54, both transmembrane proteins detected by using specific antibodies. Data are pooled from

three independent experiments. *, $p < 0.05$ by Student's t-test indicates significant differences from

uninfected controls.

Figure 3: The decrease in cell surface expression of GPI-APs is dependent on PLY and its pore formation capacity. H292 cells were infected with the indicated bacterial strains at an MOI of 10 or treated with the indicated concentrations of recombinant toxins for 1h (A-B) or 10 min (C-D). The cells were then stained and the MFI of CD73 or FLAER was measured by flow cytometry. Data pooled from three independent experiments are shown. #, $p < 0.05$ indicates significant differences from uninfected controls and *, $p < 0.05$ indicates significant differences between the indicated groups by One-way ANOVA followed by Tukey's test (A-C) or Dunnet's test (D). Units for the indicated toxins were determined by their ability to induce pores in the cells. 1U of toxin is equivalent to the concentration of toxin that causes pores in 50% of the cells. rPLY (recombinant PLY), PLY (Pneumolysin), SLO (Streptolysin O) and ILY (intermedilysin). T459G, M-1 and P-1 refer to *S. pneumoniae* strains expressing mutant forms of PLY that cannot form fully functional pores (see Materials & Methods).

Figure 4: Infection with *S. pneumoniae* triggers the internalization of GPI-APs from the host cell surface. (A) H292 cells were infected with wildtype or *S. pneumoniae* Δ*ply* at an MOI of 20 for 1 h, fixed with PFA for 20 min, permeabilized with 0.1% Tx-100 and stained with phalloidin (to detect actin filaments**,** red) and FLAER (to detect GPI-APS, green). White arrows point GPI-APs concentrating intracellularly into vesicle-like structures. Insets highlight FLAER staining in the different conditions. Bars indicate a 20 µm scale. Representative images from four independent experiments are shown. (B) Left panel shows the schematic illustration of FLAER labeling of H292 cells either pre- or post-infection with *S. pneumoniae* TIGR4 strain (MOI of 10 for 1 h). The right panel shows the MFI of FLAER measured by flow cytometry. Data pooled from three independent experiments are shown. *, *p*< 0.05 indicates significant differences between the indicated groups by One-way ANOVA followed by Tukey's test. (C) The left panel shows sequential frames of time-lapse microscopy video of HeLa cells pre-labeled with FLAER and challenged with PLY (0.25 nM) or HBSS (control). PLY was added at time point 0 min. The inset

shows detailed distribution of GPI-APs up to 45 min post PLY challenge. The right panel shows

quantified MFI for the FLAER signal overtime. Image analysis was performed using ImageJ by

randomly selecting a region of interest (ROI) in the membrane of at least 53 cells per condition

and calculating MFI overtime. The values were then normalized to the mean MFI calculated

before adding PLY or HBSS. Scale bar for the middle lane is 10 μm and for the bottom lane is 1

μm. White arrows in the bottom lane indicate where GPI-APs appear in "vesicle-like" structures

likely inside the cells. *, $p<0.05$ by Student's paired t-test.

Figure 5: Removal of GPI-APs from the cell surface largely depends on host GTPases RhoA and Cdc42 activity. (A) H292 cells were pre-treated with 100 ng/ml *C. difficile* toxin B (Tox B) or vehicle control (VC) overnight and infected with *S. pneumoniae* TIGR4 strain at an MOI of 10 for 30 min, labeled with FLAER and the MFI was assessed by flow cytometry. Data are pooled from three independent experiments. *, $p< 0.05$ by One-way ANOVA followed by Tukey's test. (B) H292 cells were transfected with a control non-targeting (NTC) siRNA, or siRNA targeting GTPases Cdc42, RhoA and Rac1. At 48 h post-transfection, whole cell lysates were used to confirm the depletion of the target proteins through Western blotting. Detection of GAPDH was used as a loading control (left panel). Depleted and control cells were infected with *S. pneumoniae* TIGR4 at an MOI of 10 for 30 min, labeled with FLAER and the MFI assessed by flow cytometry (right panel). The MFI values of FLAER are displayed as fold changes with respect to the uninfected baseline for the respective siRNA condition. The data shown are the mean of three independent experiments done in duplicates. *, $p < 0.05$ indicates significantly different than 1 by

one-sample t-test. (C) HeLa cells over-expressing a FRET probe for RhoA or Cdc42 were imaged

for 20 min before and 10 min after incubation with 0.25 nM of PLY. The upper panel shows

representative sequential frames of time-lapse images of HeLa cells before and after incubation

with PLY. FRET/CFP ratio images are shown in blue-to-white pseudocolor scale with white being

high activation. Arrows show the regions where RhoA (left) or Cdc42 (right) are activated

(FRET/CFP signal is high). The lower panel shows the quantitative analysis for the ratio between

the FRET and CFP signal (FRET/CFP), which measures the RhoA and Cdc42 activation levels.

The quantification was performed using a custom-made macro to calculate FRET/CFP ratio

specifically at the cell membrane. 79 cells (for RhoA) and 74 cells (for Cdc42) were analyzed from

two independent experiments.

Figure 6: Internalization of GPI-APs is associated with lower levels of PLY-induced cellular damage. H292 cells were infected with *S. pneumoniae* TIGR4 strain at an MOI of 10 or left uninfected. The cells were then labeled with FLAER and flow cytometry was used to measure (A) the MFI of FLAER at indicated time points of infection and (B) calculate the percentage of H292 cells (left panel) with down-regulated FLAER expression (termed as GPI^low) based upon the gating strategy shown in the right panel. (C) H292 cells suspended in media containing propidium iodide (PI) were infected with *S. pneumoniae* TIGR4 strain at an MOI of 10 and labeled with FLAER. GPI^low vs GPI^hi cell populations were then gated on, as indicated in the left panel of (B) and the percentage of H292 cells within each population that incorporated the PI dye was assessed by flow cytometry. (D) H292 cells were infected with *S. pneumoniae* for the indicated times and stained with Annexin V and FLAER. GPI^low vs GPI^hi populations were gated (right panel in B) and the percentage of apoptotic (Annexin V +) cells within each population was determined through flow cytometry. Data pooled from three independent experiments are shown. * *p*< 0.05 indicate differences by one-way ANOVA comparison between the different groups at indicated time points.

Figure 7: Working Model. PLY inserts in cholesterol-rich microdomains where GPI-APs are colocalized. PLY-induced pores trigger membrane repair resulting in internalization of the GPI-AP containing damaged membrane in a RhoA/Cdc42-dependent manner.

References:

1. Estimates of the global, regional, and national morbidity, mortality, and aetiologies of lower respiratory infections in 195 countries, 1990-2016: a systematic analysis for the Global Burden of Disease Study 2016. Lancet Infect Dis. 2018;18(11):1191-210. Epub 2018/09/24. doi: 10.1016/s1473-3099(18)30310-4. PubMed PMID: 30243584; PubMed Central PMCID: PMCPMC6202443.

2. Organization WH. Pneumonia in children. . Accessed April 2, 2024. doi: https://www.who.int/news-room/fact-sheets/detail/pneumonia.

3. van der Poll T, Opal SM. Pathogenesis, treatment, and prevention of pneumococcal pneumonia. Lancet. 2009;374(9700):1543-56. Epub 2009/11/03. doi: 10.1016/S0140-6736(09)61114-4. PubMed PMID: 19880020.

4. Randle E, Ninis N, Inwald D. Invasive pneumococcal disease. Archives of disease in childhood - Education & practice edition. 2011;96(5):183-90. doi: 10.1136/adc.2010.191718.

5. Ganaie F, Saad Jamil S, McGee L, van Tonder Andries J, Bentley Stephen D, Lo Stephanie W, et al. A New Pneumococcal Capsule Type, 10D, is the 100th Serotype and Has a Large cps Fragment from an Oral Streptococcus. mBio. 11(3):e00937-20. doi: 10.1128/mBio.00937-20.

6. Weiser JN, Ferreira DM, Paton JC. Streptococcus pneumoniae: transmission, colonization and invasion. Nat Rev Microbiol. 2018;16(6):355-67. Epub 2018/03/31. doi: 10.1038/s41579-018-0001-8. PubMed PMID: 29559457; PubMed Central PMCID: PMCPMC5949087.

7. Obaro S, Adegbola R. The pneumococcus: carriage, disease and conjugate vaccines. Journal of medical microbiology. 2002;51(2):98-104. Epub 2002/02/28. doi: 10.1099/0022-1317-51-2-98. PubMed PMID: 11863272.

8. AlonsoDeVelasco E, Verheul AF, Verhoef J, Snippe H. Streptococcus pneumoniae: virulence factors, pathogenesis, and vaccines. Microbiol Rev. 1995;59(4):591-603. Epub

1995/12/01. doi: 10.1128/mr.59.4.591-603.1995. PubMed PMID: 8531887; PubMed Central

PMCID: PMCPMC239389.

9.   Rubins JB, Charboneau D, Fasching C, Berry AM, Paton JC, Alexander JE, et al. Distinct

roles for pneumolysin's cytotoxic and complement activities in the pathogenesis of pneumococcal

pneumonia. Am J Respir Crit Care Med. 1996;153(4 Pt 1):1339-46. Epub 1996/04/01. doi:

10.1164/ajrccm.153.4.8616564. PubMed PMID: 8616564.

10.   Mitchell TJ, Andrew PW, Saunders FK, Smith AN, Boulnois GJ. Complement activation

and antibody binding by pneumolysin via a region of the toxin homologous to a human acute-

phase protein. Mol Microbiol. 1991;5(8):1883-8. Epub 1991/08/01. doi: 10.1111/j.1365-

2958.1991.tb00812.x. PubMed PMID: 1766369.

11.   Adams W, Bhowmick R, Bou Ghanem EN, Wade K, Shchepetov M, Weiser JN, et al.

Pneumolysin Induces 12-Lipoxygenase–Dependent Neutrophil Migration during

Streptococcus pneumoniae Infection. The Journal of Immunology. 2020;204(1):101-

11. doi: 10.4049/jimmunol.1800748.

12.   Nishimoto AT, Rosch JW, Tuomanen EI. Pneumolysin: Pathogenesis and Therapeutic

Target. Frontiers in Microbiology. 2020;11. doi: 10.3389/fmicb.2020.01543.

13.   Pereira JM, Xu S, Leong JM, Sousa S. The Yin and Yang of Pneumolysin During

Pneumococcal Infection. Frontiers in Immunology. 2022;13. doi: 10.3389/fimmu.2022.878244.

14.   Tweten RK, Hotze EM, Wade KR. The Unique Molecular Choreography of Giant Pore

Formation by the Cholesterol-Dependent Cytolysins of Gram-Positive Bacteria. Annu Rev

Microbiol. 2015;69:323-40. Epub 2015/10/22. doi: 10.1146/annurev-micro-091014-104233.

PubMed PMID: 26488276; PubMed Central PMCID: PMCPMC7875328.

15.   Shatursky O, Heuck AP, Shepard LA, Rossjohn J, Parker MW, Johnson AE, et al. The

mechanism of membrane insertion for a cholesterol-dependent cytolysin: a novel paradigm for

pore-forming toxins. Cell. 1999;99(3):293-9. Epub 1999/11/11. doi: 10.1016/s0092-

8674(00)81660-8. PubMed PMID: 10555145.

16. Dang TX, Hotze EM, Rouiller I, Tweten RK, Wilson-Kubalek EM. Prepore to pore transition

of a cholesterol-dependent cytolysin visualized by electron microscopy. J Struct Biol.

2005;150(1):100-8. Epub 2005/03/31. doi: 10.1016/j.jsb.2005.02.003. PubMed PMID: 15797734.

17. van Pee K, Neuhaus A, D'Imprima E, Mills DJ, Kühlbrandt W, Yildiz Ö. CryoEM structures

of membrane pore and prepore complex reveal cytolytic mechanism of Pneumolysin. eLife.

2017;6:e23644. doi: 10.7554/eLife.23644.

18. Berry AM, Yother J, Briles DE, Hansman D, Paton JC. Reduced virulence of a defined

pneumolysin-negative mutant of Streptococcus pneumoniae. Infect Immun. 1989;57(7):2037-42.

Epub 1989/07/01. doi: 10.1128/iai.57.7.2037-2042.1989. PubMed PMID: 2731982; PubMed

Central PMCID: PMCPMC313838.

19. Diep DB, Nelson KL, Raja SM, Pleshak EN, Buckley JT. Glycosylphosphatidylinositol

anchors of membrane glycoproteins are binding determinants for the channel-forming toxin

aerolysin. J Biol Chem. 1998;273(4):2355-60. Epub 1998/01/27. doi: 10.1074/jbc.273.4.2355.

PubMed PMID: 9442081.

20. Fujihara Y, Ikawa M. GPI-AP release in cellular, developmental, and reproductive biology.

J Lipid Res. 2016;57(4):538-45. Epub 2015/11/26. doi: 10.1194/jlr.R063032. PubMed PMID:

26593072; PubMed Central PMCID: PMCPMC4808780.

21. Ikezawa H. Glycosylphosphatidylinositol (GPI)-anchored proteins. Biol Pharm Bull.

2002;25(4):409-17. Epub 2002/05/09. doi: 10.1248/bpb.25.409. PubMed PMID: 11995915.

22. Zurzolo C, Simons K. Glycosylphosphatidylinositol-anchored proteins: Membrane

organization and transport. Biochimica et Biophysica Acta (BBA) - Biomembranes.

2016;1858(4):632-9. doi: https://doi.org/10.1016/j.bbamem.2015.12.018.

23. Lopez S, Rodriguez-Gallardo S, Sabido-Bozo S, Muñiz M. Endoplasmic Reticulum Export

of GPI-Anchored Proteins. International Journal of Molecular Sciences. 2019;20(14):3506.

PubMed PMID: doi:10.3390/ijms20143506.

24.   Kinoshita T. Biosynthesis and biology of mammalian GPI-anchored proteins. Open
Biology. 2020;10(3):190290. doi: doi:10.1098/rsob.190290.

25.   Muñiz M, Zurzolo C. Sorting of GPI-anchored proteins from yeast to mammals--common
pathways at different sites? J Cell Sci. 2014;127(Pt 13):2793-801. Epub 2014/06/08. doi:
10.1242/jcs.148056. PubMed PMID: 24906797.

26.   Paladino S, Lebreton S, Tivodar S, Campana V, Tempre R, Zurzolo C. Different GPI-
attachment signals affect the oligomerisation of GPI-anchored proteins and their apical sorting. J
Cell Sci. 2008;121(Pt 24):4001-7. Epub 2008/12/06. doi: 10.1242/jcs.036038. PubMed PMID:
19056670.

27.   Raghupathy R, Anilkumar AA, Polley A, Singh PP, Yadav M, Johnson C, et al.
Transbilayer lipid interactions mediate nanoclustering of lipid-anchored proteins. Cell.
2015;161(3):581-94. Epub 2015/04/25. doi: 10.1016/j.cell.2015.03.048. PubMed PMID:
25910209; PubMed Central PMCID: PMCPMC4651428.

28.   Lebreton S, Zurzolo C, Paladino S. Organization of GPI-anchored proteins at the cell
surface and its physiopathological relevance. Crit Rev Biochem Mol Biol. 2018;53(4):403-19.
Epub 2018/07/25. doi: 10.1080/10409238.2018.1485627. PubMed PMID: 30040489.

29.   Sabharanjak S, Sharma P, Parton RG, Mayor S. GPI-Anchored Proteins Are Delivered to
Recycling Endosomes via a Distinct cdc42-Regulated, Clathrin-Independent Pinocytic Pathway.
Developmental Cell. 2002;2(4):411-23. doi: https://doi.org/10.1016/S1534-5807(02)00145-4.

30.   Hodge RG, Ridley AJ. Regulating Rho GTPases and their regulators. Nat Rev Mol Cell
Biol. 2016;17(8):496-510. Epub 2016/06/16. doi: 10.1038/nrm.2016.67. PubMed PMID:
27301673.

31.   Heasman SJ, Ridley AJ. Mammalian Rho GTPases: new insights into their functions from
in vivo studies. Nat Rev Mol Cell Biol. 2008;9(9):690-701. Epub 2008/08/23. doi:
10.1038/nrm2476. PubMed PMID: 18719708.

32.   Chadda R, Howes MT, Plowman SJ, Hancock JF, Parton RG, Mayor S. Cholesterol-
sensitive Cdc42 activation regulates actin polymerization for endocytosis via the GEEC pathway.
Traffic. 2007;8(6):702-17. Epub 2007/04/28. doi: 10.1111/j.1600-0854.2007.00565.x. PubMed
PMID: 17461795.

33.   Hasko G, Linden J, Cronstein B, Pacher P. Adenosine receptors: therapeutic aspects for
inflammatory and immune diseases. Nature reviews Drug discovery. 2008;7(9):759-70. Epub
2008/09/02. doi: 10.1038/nrd2638. PubMed PMID: 18758473; PubMed Central PMCID:
PMC2568887.

34.   Bou Ghanem EN, Clark S, Roggensack SE, McIver SR, Alcaide P, Haydon PG, et al.
Extracellular Adenosine Protects against Streptococcus pneumoniae Lung Infection by
Regulating Pulmonary Neutrophil Recruitment. PLoS pathogens. 2015;11(8):e1005126. Epub
2015/08/28. doi: 10.1371/journal.ppat.1005126. PubMed PMID: 26313746; PubMed Central
PMCID: PMC4552087.

35.   Siwapornchai N, Lee JN, Tchalla EYI, Bhalla M, Yeoh JH, Roggensack SE, et al.
Extracellular adenosine enhances the ability of PMNs to kill Streptococcus pneumoniae by
inhibiting IL-10 production. J Leukoc Biol. 2020. Epub 2020/02/06. doi: 10.1002/JLB.4MA0120-
115RR. PubMed PMID: 32017200.

36.   Bhalla M, Heinzinger LR, Morenikeji OB, Marzullo B, Thomas BN, Bou Ghanem EN.
Transcriptome profiling reveals CD73 and age-driven changes in neutrophil responses against
Streptococcus pneumoniae. Infect Immun. 2021:Iai0025821. Epub 2021/07/27. doi:
10.1128/iai.00258-21. PubMed PMID: 34310891.

37.   Bhalla M, Simmons SR, Abamonte A, Herring SE, Roggensack SE, Bou Ghanem EN.
Extracellular adenosine signaling reverses the age-driven decline in the ability of neutrophils to
kill Streptococcus pneumoniae. Aging Cell. 2020:e13218. Epub 2020/08/14. doi:
10.1111/acel.13218. PubMed PMID: 32790148.

38.   Bhalla M, Hui Yeoh J, Lamneck C, Herring SE, Tchalla EYI, Heinzinger LR, et al. A1

adenosine receptor signaling reduces Streptococcus pneumoniae adherence to pulmonary

epithelial cells by targeting expression of platelet-activating factor receptor. Cellular microbiology.

2020;22(2):e13141. Epub 2019/11/12. doi: 10.1111/cmi.13141. PubMed PMID: 31709673.

39.   Bhowmick R, Tin Maung NH, Hurley BP, Ghanem EB, Gronert K, McCormick BA, et al.

Systemic disease during Streptococcus pneumoniae acute lung infection requires 12-

lipoxygenase-dependent inflammation. J Immunol. 2013;191(10):5115-23. Epub 2013/10/04. doi:

10.4049/jimmunol.1300522. PubMed PMID: 24089193; PubMed Central PMCID: PMC3836588.

40.   Yonker LM, Pazos MA, Lanter BB, Mou H, Chu KK, Eaton AD, et al. Neutrophil-Derived

Cytosolic PLA2alpha Contributes to Bacterial-Induced Neutrophil Transepithelial Migration. J

Immunol. 2017;199(8):2873-84. Epub 2017/09/10. doi: 10.4049/jimmunol.1700539. PubMed

PMID: 28887431; PubMed Central PMCID: PMC5636676.

41.   Choi CH, Lee JS, Lee YC, Park TI, Lee JC. Acinetobacter baumannii invades epithelial

cells and outer membrane protein A mediates interactions with epithelial cells. BMC microbiology.

2008;8:216. Epub 2008/12/11. doi: 10.1186/1471-2180-8-216. PubMed PMID: 19068136;

PubMed Central PMCID: PMC2615016.

42.   Antonioli L, Pacher P, Vizi ES, Haskó G. CD39 and CD73 in immunity and inflammation.

Trends Mol Med. 2013;19(6):355-67. Epub 2013/04/23. doi: 10.1016/j.molmed.2013.03.005.

PubMed PMID: 23601906; PubMed Central PMCID: PMCPMC3674206.

43.   Brodsky RA, Mukhina GL, Li S, Nelson KL, Chiurazzi PL, Buckley JT, et al. Improved

detection and characterization of paroxysmal nocturnal hemoglobinuria using fluorescent

aerolysin. Am J Clin Pathol. 2000;114(3):459-66. Epub 2000/09/16. doi: 10.1093/ajcp/114.3.459.

PubMed PMID: 10989647; PubMed Central PMCID: PMCPMC4124633.

44.   Paladino S, Lebreton S, Lelek M, Riccio P, De Nicola S, Zimmer C, et al. GPI-anchored

proteins are confined in subdiffraction clusters at the apical surface of polarized epithelial cells.

Biochem J. 2017;474(24):4075-90. Epub 2017/10/20. doi: 10.1042/bcj20170582. PubMed PMID: 29046391; PubMed Central PMCID: PMCPMC5712066.

45.   Au - Kusek ME, Au - Pazos MA, Au - Pirzai W, Au - Hurley BP. In vitro Coculture Assay to Assess Pathogen Induced Neutrophil Trans-epithelial Migration. JoVE. 2014;(83):e50823. doi: doi:10.3791/50823.

46.   Farrand AJ, LaChapelle S, Hotze EM, Johnson AE, Tweten RK. Only two amino acids are essential for cytolytic toxin recognition of cholesterol at the membrane surface. Proceedings of the National Academy of Sciences. 2010;107(9):4341-6. doi: doi:10.1073/pnas.0911581107.

47.   Shepard LA, Heuck AP, Hamman BD, Rossjohn J, Parker MW, Ryan KR, et al. Identification of a membrane-spanning domain of the thiol-activated pore-forming toxin Clostridium perfringens perfringolysin O: an alpha-helical to beta-sheet transition identified by fluorescence spectroscopy. Biochemistry. 1998;37(41):14563-74. Epub 1998/10/17. doi: 10.1021/bi981452f. PubMed PMID: 9772185.

48.   Schmeck B, Gross R, N'Guessan PD, Hocke AC, Hammerschmidt S, Mitchell TJ, et al. Streptococcus pneumoniae-induced caspase 6-dependent apoptosis in lung epithelium. Infect Immun. 2004;72(9):4940-7. Epub 2004/08/24. doi: 10.1128/iai.72.9.4940-4947.2004. PubMed PMID: 15321985; PubMed Central PMCID: PMCPMC517413.

49.   Marriott HM, Mitchell TJ, Dockrell DH. Pneumolysin: a double-edged sword during the host-pathogen interaction. Curr Mol Med. 2008;8(6):497-509. Epub 2008/09/11. doi: 10.2174/156652408785747924. PubMed PMID: 18781957.

50.   Heuck AP, Moe PC, Johnson BB. The cholesterol-dependent cytolysin family of gram-positive bacterial toxins. Subcell Biochem. 2010;51:551-77. Epub 2010/03/10. doi: 10.1007/978-90-481-8622-8_20. PubMed PMID: 20213558.

51.   Paulick MG, Bertozzi CR. The Glycosylphosphatidylinositol Anchor: A Complex Membrane-Anchoring Structure for Proteins. Biochemistry. 2008;47(27):6991-7000. doi: 10.1021/bi8006324.

52. Asmat TM, Agarwal V, Räth S, Hildebrandt JP, Hammerschmidt S. Streptococcus

pneumoniae infection of host epithelial cells via polymeric immunoglobulin receptor transiently

induces calcium release from intracellular stores. J Biol Chem. 2011;286(20):17861-9. Epub

2011/04/02. doi: 10.1074/jbc.M110.212225. PubMed PMID: 21454571; PubMed Central PMCID:

PMCPMC3093861.

53. Takenouchi T, Ogihara K, Sato M, Kitani H. Inhibitory effects of U73122 and U73343 on

Ca2+ influx and pore formation induced by the activation of P2X7 nucleotide receptors in mouse

microglial cell line. Biochim Biophys Acta. 2005;1726(2):177-86. Epub 2005/08/27. doi:

10.1016/j.bbagen.2005.08.001. PubMed PMID: 16122875.

54. Letsiou E, Teixeira Alves LG, Fatykhova D, Felten M, Mitchell TJ, Müller-Redetzky HC, et

al. Microvesicles released from pneumolysin-stimulated lung epithelial cells carry mitochondrial

cargo and suppress neutrophil oxidative burst. Scientific Reports. 2021;11(1):9529. doi:

10.1038/s41598-021-88897-y.

55. Alves S, Pereira JM, Mayer RL, Gonçalves ADA, Impens F, Cabanes D, et al. Cells

Responding to Closely Related Cholesterol-Dependent Cytolysins Release Extracellular Vesicles

with a Common Proteomic Content Including Membrane Repair Proteins. Toxins (Basel).

2022;15(1). Epub 2023/01/21. doi: 10.3390/toxins15010004. PubMed PMID: 36668824; PubMed

Central PMCID: PMCPMC9865450.

56. Babiychuk EB, Monastyrskaya K, Potez S, Draeger A. Blebbing confers resistance against

cell lysis. Cell Death Differ. 2011;18(1):80-9. Epub 2010/07/03. doi: 10.1038/cdd.2010.81.

PubMed PMID: 20596076; PubMed Central PMCID: PMCPMC3131879.

57. Wolfmeier H, Radecke J, Schoenauer R, Koeffel R, Babiychuk VS, Drücker P, et al. Active

release of pneumolysin prepores and pores by mammalian cells undergoing a Streptococcus

pneumoniae attack. Biochim Biophys Acta. 2016;1860(11 Pt A):2498-509. Epub 2016/08/03. doi:

10.1016/j.bbagen.2016.07.022. PubMed PMID: 27481675.

58.   Limouze J, Straight AF, Mitchison T, Sellers JR. Specificity of blebbistatin, an inhibitor of

myosin II. J Muscle Res Cell Motil. 2004;25(4-5):337-41. Epub 2004/11/19. doi: 10.1007/s10974-

004-6060-7. PubMed PMID: 15548862.

59.   Yarar D, Waterman-Storer CM, Schmid SL. A Dynamic Actin Cytoskeleton Functions at

Multiple Stages of Clathrin-mediated Endocytosis. Molecular Biology of the Cell. 2005;16(2):964-

75. doi: 10.1091/mbc.e04-09-0774. PubMed PMID: 15601897.

60.   Pistor S, Chakraborty T, Niebuhr K, Domann E, Wehland J. The ActA protein of Listeria

monocytogenes acts as a nucleator inducing reorganization of the actin cytoskeleton. Embo j.

1994;13(4):758-63. Epub 1994/02/15. doi: 10.1002/j.1460-2075.1994.tb06318.x. PubMed PMID:

8112291; PubMed Central PMCID: PMCPMC394872.

61.   Mayorga LS, Diaz R, Stahl PD. Regulatory role for GTP-binding proteins in endocytosis.

Science. 1989;244(4911):1475-7. Epub 1989/06/23. doi: 10.1126/science.2499930. PubMed

PMID: 2499930.

62.   Aktories K, Schmidt G, Just I. Rho GTPases as Targets of Bacterial Protein Toxins.

2000;381(5-6):421-6. doi: doi:10.1515/BC.2000.054.

63.   Iliev AI, Djannatian JR, Nau R, Mitchell TJ, Wouters FS. Cholesterol-dependent actin

remodeling via RhoA and Rac1 activation by the <i>Streptococcus pneumoniae</i> toxin

pneumolysin. Proceedings of the National Academy of Sciences. 2007;104(8):2897-902. doi:

doi:10.1073/pnas.0608213104.

64.   Tweten RK. Cholesterol-dependent cytolysins, a family of versatile pore-forming toxins.

Infect Immun. 2005;73(10):6199-209. Epub 2005/09/24. doi: 10.1128/iai.73.10.6199-6209.2005.

PubMed PMID: 16177291; PubMed Central PMCID: PMCPMC1230961.

65.   Brito C, Cabanes D, Sarmento Mesquita F, Sousa S. Mechanisms protecting host cells

against bacterial pore-forming toxins. Cell Mol Life Sci. 2019;76(7):1319-39. Epub 2018/12/29.

doi: 10.1007/s00018-018-2992-8. PubMed PMID: 30591958; PubMed Central PMCID:

PMCPMC6420883.

66.   Casadevall A, Pirofski La. Host-Pathogen Interactions: The Attributes of Virulence. The

Journal of Infectious Diseases. 2001;184(3):337-44. doi: 10.1086/322044.

67.   Casadevall A, Pirofski LA. Host-pathogen interactions: basic concepts of microbial

commensalism, colonization, infection, and disease. Infect Immun. 2000;68(12):6511-8. Epub

2000/11/18. doi: 10.1128/iai.68.12.6511-6518.2000. PubMed PMID: 11083759; PubMed Central

PMCID: PMCPMC97744.

68.   Brown AO, Millett ER, Quint JK, Orihuela CJ. Cardiotoxicity during invasive pneumococcal

disease. Am J Respir Crit Care Med. 2015;191(7):739-45. Epub 2015/01/30. doi:

10.1164/rccm.201411-1951PP. PubMed PMID: 25629643; PubMed Central PMCID:

PMCPMC4407487.

69.   Adams W, Bhowmick R, Bou Ghanem EN, Wade K, Shchepetov M, Weiser JN, et al.

Pneumolysin Induces 12-Lipoxygenase–Dependent Neutrophil Migration during Streptococcus

pneumoniae Infection. The Journal of Immunology. 2020;204(1):101-11. doi:

10.4049/jimmunol.1800748.

70.   Reyes LF, Restrepo MI, Hinojosa CA, Soni NJ, Anzueto A, Babu BL, et al. Severe

Pneumococcal Pneumonia Causes Acute Cardiac Toxicity and Subsequent Cardiac Remodeling.

Am J Respir Crit Care Med. 2017;196(5):609-20. Epub 2017/06/15. doi: 10.1164/rccm.201701-

0104OC. PubMed PMID: 28614669; PubMed Central PMCID: PMCPMC5620668.

71.   Kadioglu A, Weiser JN, Paton JC, Andrew PW. The role of Streptococcus pneumoniae

virulence factors in host respiratory colonization and disease. Nature reviews Microbiology.

2008;6(4):288-301. Epub 2008/03/15. doi: 10.1038/nrmicro1871. PubMed PMID: 18340341.

72.   Price KE, Camilli A. Pneumolysin localizes to the cell wall of Streptococcus pneumoniae.

J Bacteriol. 2009;191(7):2163-8. Epub 2009/01/27. doi: 10.1128/jb.01489-08. PubMed PMID:

19168620; PubMed Central PMCID: PMCPMC2655535.

73.   Martins R, Carlos AR, Braza F, Thompson JA, Bastos-Amador P, Ramos S, et al. Disease

Tolerance as an Inherent Component of Immunity. Annual Review of Immunology.

2019;37(1):405-37. doi: 10.1146/annurev-immunol-042718-041739. PubMed PMID: 30673535.

74.   Varma R, Mayor S. GPI-anchored proteins are organized in submicron domains at the cell

surface. Nature. 1998;394(6695):798-801. Epub 1998/09/02. doi: 10.1038/29563. PubMed PMID:

9723621.

75.   Shepard LA, Shatursky O, Johnson AE, Tweten RK. The mechanism of pore assembly

for a cholesterol-dependent cytolysin: formation of a large prepore complex precedes the insertion

of the transmembrane beta-hairpins. Biochemistry. 2000;39(33):10284-93. Epub 2000/08/24. doi:

10.1021/bi000436r. PubMed PMID: 10956018.

76.   Corrotte M, Fernandes MC, Tam C, Andrews NW. Toxin pores endocytosed during

plasma membrane repair traffic into the lumen of MVBs for degradation. Traffic. 2012;13(3):483-

94. Epub 2012/01/04. doi: 10.1111/j.1600-0854.2011.01323.x. PubMed PMID: 22212686;

PubMed Central PMCID: PMCPMC3356596.

77.   Bouillot S, Reboud E, Huber P. Functional Consequences of Calcium Influx Promoted by

Bacterial Pore-Forming Toxins. Toxins (Basel). 2018;10(10). Epub 2018/09/28. doi:

10.3390/toxins10100387. PubMed PMID: 30257425; PubMed Central PMCID:

PMCPMC6215193.

78.   Ray S, Roth R, Keyel PA. Membrane repair triggered by cholesterol-dependent cytolysins

is activated by mixed lineage kinases and MEK. Science advances. 2022;8(11):eabl6367. Epub

2022/03/17. doi: 10.1126/sciadv.abl6367. PubMed PMID: 35294243; PubMed Central PMCID:

PMCPMC8926344.

79.   Hervé JC, Bourmeyster N. Rho GTPases at the crossroad of signaling networks in

mammals.   Small   GTPases.   2015;6(2):43-8.   Epub   2015/06/26.   doi:

10.1080/21541248.2015.1044811. PubMed PMID: 26110743; PubMed Central PMCID:

PMCPMC4601364.

80.   Lakhan SE, Sabharanjak S, De A. Endocytosis of glycosylphosphatidylinositol-anchored

proteins. Journal of Biomedical Science. 2009;16(1):93. doi: 10.1186/1423-0127-16-93.

81.   Aït-Slimane T, Galmes R, Trugnan G, Maurice M. Basolateral internalization of GPI-

anchored proteins occurs via a clathrin-independent flotillin-dependent pathway in polarized

hepatic cells. Mol Biol Cell. 2009;20(17):3792-800. Epub 2009/07/17. doi: 10.1091/mbc.e09-04-

0275. PubMed PMID: 19605558; PubMed Central PMCID: PMCPMC2735478.

82.   Naslavsky N, Weigert R, Donaldson JG. Characterization of a nonclathrin endocytic

pathway: membrane cargo and lipid requirements. Mol Biol Cell. 2004;15(8):3542-52. Epub

2004/05/18. doi: 10.1091/mbc.e04-02-0151. PubMed PMID: 15146059; PubMed Central PMCID:

PMCPMC491817.

83.   Fujimoto LM, Roth R, Heuser JE, Schmid SL. Actin Assembly Plays a Variable, but not

Obligatory Role in Receptor-Mediated Endocytosis. Traffic. 2000;1(2):161-71. doi:

https://doi.org/10.1034/j.1600-0854.2000.010208.x.

84.   Yao L-H, Rao Y, Bang C, Kurilova S, Varga K, Wang C-Y, et al. Actin Polymerization Does

Not Provide Direct Mechanical Forces for Vesicle Fission during Clathrin-Mediated Endocytosis.

The Journal of Neuroscience. 2013;33(40):15793-8. doi: 10.1523/jneurosci.2171-13.2013.

85.   Hyman T, Shmuel M, Altschuler Y. Actin is required for endocytosis at the apical surface

of Madin-Darby canine kidney cells where ARF6 and clathrin regulate the actin cytoskeleton. Mol

Biol Cell. 2006;17(1):427-37. Epub 2005/10/28. doi: 10.1091/mbc.e05-05-0420. PubMed PMID:

16251360; PubMed Central PMCID: PMCPMC1345679.

86.   Hänisch J, Kölm R, Wozniczka M, Bumann D, Rottner K, Stradal TE. Activation of a

RhoA/myosin II-dependent but Arp2/3 complex-independent pathway facilitates Salmonella

invasion.   Cell   Host   Microbe.   2011;9(4):273-85.   Epub   2011/04/20.   doi:

10.1016/j.chom.2011.03.009. PubMed PMID: 21501827.

87.   Van Ngo H, Bhalla M, Chen D-Y, Ireton K. A role for host cell exocytosis in InlB-mediated

internalisation of Listeria monocytogenes. Cellular Microbiology. 2017;19(11):e12768. doi:

https://doi.org/10.1111/cmi.12768.

88.   Alves S, Pereira JM, Mayer RL, Gonçalves ADA, Impens F, Cabanes D, et al. Cells

Responding to Closely Related Cholesterol-Dependent Cytolysins Release Extracellular Vesicles

with a Common Proteomic Content Including Membrane Repair Proteins. Toxins. 2023;15(1):4.

PubMed PMID: doi:10.3390/toxins15010004.

89.   Schmiel DH, Miller VL. Bacterial phospholipases and pathogenesis. Microbes and

Infection. 1999;1(13):1103-12. doi: https://doi.org/10.1016/S1286-4579(99)00205-1.

90.   Shichinohe N, Kobayashi D, Izumi A, Hatanaka K, Fujita R, Kinoshita T, et al. Sequential

hydrolysis of FAD by ecto-5′ nucleotidase CD73 and alkaline phosphatase is required

for uptake of vitamin B$_2$ into cells. Journal of Biological Chemistry. 2022;298(12).

doi: 10.1016/j.jbc.2022.102640.

91.   Hasko G, Csoka B, Koscso B, Chandra R, Pacher P, Thompson LF, et al. Ecto-5'-

nucleotidase (CD73) decreases mortality and organ injury in sepsis. J Immunol.

2011;187(8):4256-67. Epub 2011/09/16. doi: 10.4049/jimmunol.1003379. PubMed PMID:

21918191; PubMed Central PMCID: PMC3387540.

92.   Tanaka K, Choi J, Cao Y, Stacey G. Extracellular ATP acts as a damage-associated

molecular pattern (DAMP) signal in plants. Frontiers in Plant Science. 2014;5. doi:

10.3389/fpls.2014.00446.

93.   Regateiro FS, Cobbold SP, Waldmann H. CD73 and adenosine generation in the creation

of regulatory microenvironments. Clinical and experimental immunology. 2013;171(1):1-7. Epub

2012/12/04. doi: 10.1111/j.1365-2249.2012.04623.x. PubMed PMID: 23199317; PubMed Central

PMCID: PMC3530089.

94.   Kao DJ, Saeedi BJ, Kitzenberg D, Burney KM, Dobrinskikh E, Battista KD, et al. Intestinal

Epithelial Ecto-5'-Nucleotidase (CD73) Regulates Intestinal Colonization and Infection by

Nontyphoidal Salmonella. Infect Immun. 2017;85(10). Epub 2017/07/19. doi: 10.1128/IAI.01022-

16. PubMed PMID: 28717030; PubMed Central PMCID: PMC5607412.

95.   Dho SH, Lim JC, Kim LK. Beyond the Role of CD55 as a Complement Component.

Immune Netw. 2018;18(1):e11. Epub 2018/03/06. doi: 10.4110/in.2018.18.e11. PubMed PMID:

29503741; PubMed Central PMCID: PMCPMC5833118.

96.   Huang Y, Qiao F, Abagyan R, Hazard S, Tomlinson S. Defining the CD59-C9 binding

interaction.   J   Biol   Chem.   2006;281(37):27398-404.   Epub   2006/07/18.   doi:

10.1074/jbc.M603690200. PubMed PMID: 16844690.

97.   Lawrence DW, Bruyninckx WJ, Louis NA, Lublin DM, Stahl GL, Parkos CA, et al.

Antiadhesive role of apical decay-accelerating factor (CD55) in human neutrophil transmigration

across mucosal epithelia. The Journal of experimental medicine. 2003;198(7):999-1010. Epub

2003/10/08. doi: 10.1084/jem.20030380. PubMed PMID: 14530374; PubMed Central PMCID:

PMC2194216.

98.   Greene NG, Narciso AR, Filipe SR, Camilli A. Peptidoglycan Branched Stem Peptides

Contribute to Streptococcus pneumoniae Virulence by Inhibiting Pneumolysin Release. PLoS

pathogens.   2015;11(6):e1004996.   Epub   2015/06/27.   doi: 10.1371/journal.ppat.1004996.

PubMed PMID: 26114646; PubMed Central PMCID: PMC4483231.

99.   Shainheit MG, Mule M, Camilli A. The core promoter of the capsule operon of

Streptococcus pneumoniae is necessary for colonization and invasive disease. Infect Immun.

2014;82(2):694-705. Epub 2014/01/31. doi: 10.1128/IAI.01289-13. PubMed PMID: 24478084;

PubMed Central PMCID: PMC3911406.

100.   Yuste J, Khandavilli S, Ansari N, Muttardi K, Ismail L, Hyams C, et al. The effects of PspC

on complement-mediated immunity to Streptococcus pneumoniae vary with strain background

and   capsular   serotype.   Infect   Immun.   2010;78(1):283-92.   Epub   2009/11/04.   doi:

10.1128/IAI.00541-09. PubMed PMID: 19884335; PubMed Central PMCID: PMC2798213.

101.   Price KE, Greene NG, Camilli A. Export requirements of pneumolysin in Streptococcus

pneumoniae. J Bacteriol. 2012;194(14):3651-60. Epub 2012/05/09. doi: 10.1128/jb.00114-12.

PubMed PMID: 22563048; PubMed Central PMCID: PMCPMC3393478.

102.   Herring SE, Mao S, Bhalla M, Tchalla EYI, Kramer JM, Bou Ghanem EN. Mitochondrial

ROS production by neutrophils is required for host antimicrobial function against Streptococcus

pneumoniae and is controlled by A2B adenosine receptor signaling. PLOS Pathogens.

2022;18(11):e1010700. doi: 10.1371/journal.ppat.1010700.

103.   Bhalla M, Law D, Dowd GC, Ireton K. Host Serine/Threonine Kinases mTOR and Protein

Kinase C-α Promote InlB-Mediated Entry of Listeria monocytogenes. Infection and immunity.

2017;85(7):e00087-17. doi: 10.1128/IAI.00087-17. PubMed PMID: 28461391.

104.   Bhalla M, Law D, Dowd GC, Ireton K. Host Serine/Threonine Kinases mTOR and Protein

Kinase C-alpha Promote InlB-Mediated Entry of Listeria monocytogenes. Infect Immun.

2017;85(7). Epub 2017/05/04. doi: 10.1128/IAI.00087-17. PubMed PMID: 28461391; PubMed

Central PMCID: PMCPMC5478960.

105.   Hirata T, Mishra SK, Nakamura S, Saito K, Motooka D, Takada Y, et al. Identification of a

Golgi GPI-N-acetylgalactosamine transferase with tandem transmembrane regions in the

catalytic domain. Nature Communications. 2018;9(1):405. doi: 10.1038/s41467-017-02799-0.

106.   Dowd GC, Bhalla M, Kean B, Thomas R, Ireton K. Role of Host Type IA Phosphoinositide

3-Kinase Pathway Components in Invasin-Mediated Internalization of Yersinia enterocolitica.

Infection and Immunity. 2016;84(6):1826-41. doi: doi:10.1128/IAI.00142-16.

107.   Yoshizaki H, Ohba Y, Kurokawa K, Itoh RE, Nakamura T, Mochizuki N, et al. Activity of

Rho-family GTPases during cell division as visualized with FRET-based probes. J Cell Biol.

2003;162(2):223-32. Epub 2003/07/16. doi: 10.1083/jcb.200212049. PubMed PMID: 12860967;

PubMed Central PMCID: PMCPMC2172791.

108.   Itoh RE, Kurokawa K, Ohba Y, Yoshizaki H, Mochizuki N, Matsuda M. Activation of rac

and cdc42 video imaged by fluorescent resonance energy transfer-based single-molecule probes

in the membrane of living cells. Mol Cell Biol. 2002;22(18):6582-91. Epub 2002/08/23. doi:

10.1128/mcb.22.18.6582-6591.2002. PubMed PMID: 12192056; PubMed Central PMCID:

PMCPMC135619.

1242