Genetic purging of strongly deleterious mutations underlies black-necked crane's unusual escape from an extinction vortex

Ning Cui[1, #], Xuntao Ma[2, #], Heqi Wu[1, #], Xiaocheng Chen[3], Chih-Ming Hung[4], Lixun Zhang[5], Fumin Lei[6], Le Yang[7], Dao Yan[1], Xiaojun Yang[1], Feng Dong[1,*]

1. State Key Laboratory of Genetic Resources and Evolution & Yunnan Key Laboratory of Biodiversity and Ecological Conservation of Gaoligong Mountain, Kunming Institute of Zoology, Chinese Academy of Sciences, Kunming 650223, China
2. College of Animal Sciences and Technology, Sichuan Agricultural University, Chengdu 611130, China
3. Northwest Institute of Plateau Biology, Chinese Academy of Sciences, Xining 810001, China
4. Biodiversity Research Center, Academia Sinica, Taipei 116, Taiwan
5. College of Ecology, Lanzhou University, Lanzhou 730000, China
6. Key Laboratory of Zoological Systematics and Evolution, Institute of Zoology, Chinese Academy of Sciences, Beijing 100101, China
7. Tibet Plateau Institute of Biology, Lhasa 850000, China

# These authors contributed equally
* Corresponding authors:
Feng Dong, Tel: +86-871-65192023, E-mail: dongfeng@mail.kiz.ac.cn, Address: No.17 Longxin Road, Kunming, Yunnan, 650201, China

Abstract

Many species are undergoing rapid demographic declines, necessitating an examination of the

resulting genetic impacts. The prevailing small population paradigm posits an elevated

genetic load and extinction risk. However, instances of fast recovery from severe population

bottlenecks suggest alternative outcomes. To investigate this issue, we performed a

population genomic analysis on the black-necked crane, analyzing 42 modern and 11

historical genomes. This study revealed substantial evidence of large-effect allele purging

underlying the unexpectedly rapid population recovery following an abrupt bottleneck during

the 1980s. Nevertheless, forward simulations supposing a prolonged bottleneck (e.g., five

generations) predicted a reversion with negative prospects, implying that rapid population

recovery served as both the cause and consequence of the species escaping from an extinction

vortex. These findings shed light on a potential positive microevolutionary response to current

widespread population collapses and underscore the urgency of implementing active and

effective conservation strategies to reverse this trend before it becomes irreversible.

Keywords

Population bottleneck, genetic load, selection coefficients, genomic diversity, inbreeding

Introduction

Global biodiversity is currently experiencing an unprecedented crisis, characterized by alarming rates of species extinction [1,2], with even common species showing rapid population declines [3]. Global vertebrate populations, encompassing 4 392 species, have decreased by an average of 68% in the past five decades [4]. This decline raises a critical question about the genomic response of populations to rapid collapses. The lack of such knowledge prevents scientific evaluations of population viabilities and impedes efficient conservation policymaking.

In addition to increased inbreeding and decreased genetic diversity, the genetic consequences of population declines are particularly marked by the accumulation of deleterious mutations—genetic load [5]. In large, stable populations, genetic load is typically maintained at a relatively low level due to a balance between mutagenesis and purifying selection [6]. However, demographic contractions can disrupt this balance, exerting an enigmatic influence on genetic load [5], an area that remains contentious in the fields of evolutionary biology and conservation [6]. Nevertheless, recent studies suggest that the velocity of population decline is a critical factor [7, 8].

Gradual and prolonged declines are thought to reduce recessive detrimental alleles through intensive purifying selection—a process of genetic purging—which may facilitate long-term persistence of species, even in small populations [9-12]. In contrast, rapid and severe declines (i.e., bottlenecks) may accelerate the accumulation of deleterious mutations due to enhanced genetic drift and relaxed purifying selection [5]. This can induce inbreeding depression, resulting in a negative feedback on population size and ultimately elevating extinction risk by trapping the species in an 'extinction vortex', a concept central to the small population paradigm [9,13-15]. Nevertheless, there are several exceptional cases of rapid recovery from sudden and severe bottlenecks [16-18], suggesting complex genetic consequences of population declines. These cases serve as valuable resources for understanding the potential of genetic purging in rapidly declining populations, an area that remains to be empirically examined.

The black-necked crane (*Grus nigricollis*), a notable flagship bird species (Figure 1a, primarily breeds on the high-altitude Qinghai-Tibet Plateau (QTP) and winters at lower altitudes in southeast QTP as well as the western part of the Yunnan-Guizhou Plateau (Figure 1b; [19,20]). Despite its former ubiquity, the species is widely believed to have experienced a

severe population decline to only 100-300 individuals during the late 20th century [21,22],

before rebounding to 15□000 by 2020 (Figure 2a) [22]. This represents one of the fastest

recoveries observed in the animal kingdom, although little is known about the underlying

genetic mechanism driving it. In this study, we assembled the first chromosome-level genome

of the black-necked crane and compared population genomic characteristics before and after

the bottleneck, with the specific goal of investigating the role of genetic purging. By

comprehensively examining the genetic consequences in this recent bottleneck and fast

recovery, this study aims to shed light on potential conservation strategies amid the current

widespread and rapid declines in animal populations.

Results

Comprehensive genomic data

We constructed a high-quality genome assembly of the black-necked crane using second-

and third-generation sequencing and chromosome conformation capture-3C (Hi-C) techniques.

The final assembly (Gnig-Y8) was 1.35 Gb in length and showed high continuity, with an

N50 of 85.88 Mb. Notably, the top 34 largest scaffolds, out of a total of 314 scaffolds (> 4

Mbp; hereafter: pseudochromosomes) accounted for 96% of the genome (1.29 Gb), showing

remarkable improvement over the previous genome version [23]. Further analysis confirmed

high genomic completeness of this assembly, achieving a retrieval rate of 97% for complete

Benchmarking Universal Single-Copy Orthologs (BUSCO) among birds (Supplementary

Figure S1). The annotation process identified 20□963 gene models with 922□165 exons.

We resequenced a total of 52 genomes with an average coverage of 30.2x (15.9-83.8x;

Supplementary Table S1) and compiled four comprehensive population genomic datasets for

subsequent analyses. The comparative genomic dataset comprised 2□829□286 high-quality

autosomal biallelic single nucleotide polymorphisms (SNPs) derived from 41 modern black-

necked cranes spanning their extant distribution range and 11 historical samples predating the

recent bottleneck (Figure 1b), which were used to estimate genomic diversity and genetic load,

while a subset of 560□083 putatively unlinked SNPs was used for genetic clustering analysis.

We also compiled a dataset of 1□319□028 automatic biallelic SNPs from modern samples to

reconstruct recent demography and a subset of 21□590 putatively unlinked SNPs to estimate

contemporary effective population size (N_e).

Lack of population genetic structure

We conducted admixture and principal component analysis (PCA) of historical and modern black-necked cranes to examine potential genetic stratification. Admixture analysis indicated that the lowest cross-validation (CV) error occurred at *K (number of supposed genetic clusters)* =1 (Figure 1), suggesting a single breeding population origin for all samples. The PCA results further demonstrated markedly reduced heterogeneity in modern genomes compared to historical genomes (Figure 1d).

Recent sharp population bottleneck

We retrieved details on recent population trajectories for the black-necked crane using the optimal GONE algorithm [24, 25], which uses a linkage disequilibrium spectrum and provides high resolution within the last 100 generations (1 000 years here) [24]. Results revealed a sharp bottleneck during the 1980s, with an eight-fold N_e decline from approximately 1 000 to 122 diploids (Figure 2a). Similarly, independent estimation using NeEstimator [4] based on putatively unlinked loci suggested a contemporary N_e of 127 diploids (95% confidence interval: 112 to 137).

Decreased genomic diversity and increased inbreeding

To characterize genomic diversity before and after the bottleneck, we first calculated genome-wide genetic diversity (π), which was 1.2 times higher in the historical than in the modern genomes (Wilcoxon signed rank test, $P<0.001$; Supplementary Figure S2). Second, we estimated genome-wide heterogeneity (h; average likelihood of heterozygosity for each site), which was 1.1 times higher in the historical than in the modern genomes (Wilcoxon rank sum test, $P<0.001$; Figure 2b). Third, we measured inbreeding coefficients (F_{ROH}) as the proportion of the genome covered by runs of homogeneity (ROHs), finding a 2.6-fold increase in modern genomes compared to historical genomes (Wilcoxon signed rank test, $P<0.001$; Figure 2c).

Increased genetic load in homozygous state

We measured and compared genetic load in historical and modern genomes using three distinct approaches. First, we estimated genetic load by counting the number of derived

deleterious variants in coding regions and classified them into three types: those in

heterozygotes as masked genetic load, those in homozygotes as realized genetic load, and the

combined total of the first two types as total genetic load. We categorized the potential

impacts of genetic load as High and Moderate using SnpEff [26] and compared their

distributions between historical and modern genomes (see Methods and Materials). In the

High-impact category, we observed a 22.6% reduction in total genetic load in the modern

genomes relative to historical ones (Wilcoxon rank sum test, $P<0.001$; Figure 3a), with

masked genetic load showing a 3.2% reduction (Welch's two-sample t-test, $P<0.001$; Figure

3b) but realized genetic load showing a 26.6% increase (Welch's two-sample t-test, $P<0.001$;

Figure 3c). In the Moderate impact category, we found no difference in total genetic load

(Wilcoxon rank sum test, $P=0.053$; Supplementary Figure S3a), but a 2.4% decrease in

masked genetic load (Welch's two-sample t-test, $P<0.001$; Supplementary Figure S3b) and a

21.5% increase in realized genetic load in the modern genomes compared with the historical

genomes (Welch's two-sample t-test, $P<0.001$; Supplementary Figure S3c).

Second, we estimated and compared the relative frequency of derived alleles

($R_{modern/historical}$) for the High- and Moderate-impact variants in the historical and modern

genomes. The High-impact category showed an excess of deleterious derived alleles

($R_{modern/historical}=1.24$; Supplementary Figure S4) in homozygotes and a deficit in heterozygotes

($R_{modern/historical}=0.93$; Supplementary Figure S4) in the modern genomes, indicating higher

realized genetic load and lower masked genetic load. The Moderate-impact category showed a

congruent excess in both homozygotes and heterozygotes ($R_{modern/historical}=1.23$ and 1.04,

respectively; Figure 4).

Third, we compared the relative abundance of homozygous High-impact deleterious

variants within and outside ROHs in the modern genomes. We found a 40.2% lower presence

of these variants within ROHs (Welch's pairwise t-test, $P<0.001$; Figure 3d). Given that

deleterious variants within ROHs may reduce individual fitness more markedly than those

outside ROHs [5], these results suggest a signature of genetic purging in the modern

population.

Constant purifying selection over the bottleneck

We examined purifying selection signals between the historical and modern genomes.

First, we estimated genomic evolutionary rate profiling (GERP) scores for each site of the

genomes, where loci with higher values were assumed to be under higher purifying selection

pressure. We examined sites under the strongest evolutionary constraint (i.e., those with the

top 100 GERP scores) and referred to their scores as the maximum GERP (mGERP) scores.

We found lower mGERP scores in the modern genomes compared to the historical genomes

in heterozygotes (Welch's two-sample *t*-test, *P*=0.011; Figure 4a), but no significant

difference in homozygotes (Welch's two-sample *t*-test, *P*=0.751; Figure 4b). In addition, we

found significantly higher mGERP scores in heterozygotes than homozygotes in both

historical and modern genomes (Welch's pairwise *t*-test, *P*<0.001 for both; Figure 4c-d).

Second, we conducted forward simulations and compared the individual-based selection

coefficients of (partially) recessive deleterious mutations in homozygous sites between the

historical and modern genomes (e.g., dominance coefficient (h)=0 for fully recessive and

h=0.25 for partially recessive), a major genetic basis for inbreeding depression [8]. Given that

deleterious mutations present negative selection coefficients, lower values indicate more

relaxed purifying selection. We assumed a pre-bottleneck consensus population size (N_c) of

approximately 10 000 individuals given an empirical N_e/N_c ratio of 0.1 [27] and a pre-

bottleneck N_e of approximately 1 000 (Figure 2a). Recapitulation of the demographic history

of the black-necked crane indicated no significant temporal difference in the minimum

selection coefficient for both fully and partially recessive deleterious mutations (Wilcoxon

rank sum test, *P*=0.331 and *P*=0.256, respectively; Figure 5a-b). Assuming the bottleneck

lasted for five generations (approximately 50 years), we found that purifying selection was

significantly relaxed for both types of recessive deleterious mutations (Wilcoxon rank sum

test, *P*<0.001; Figure 5c-d).

Discussion

Based on comprehensive population genomic data, we identified an abrupt eight-fold

reduction in the effective population size of the black-necked crane during the 1980s (Figure

2a). This decline corresponds with historical records of a severe bottleneck, which reduced the

population to only 100-300 individuals, during the same period [21,22]. While this decline

was initially considered a potential artifact of inadequate investigation, subsequent intensive

field surveys challenged this view, showing a continual and rapid population expansion

(Figure 1a; [22]), indicating recovery from a preceding bottleneck. Support for the bottleneck

event is further provided by our observations of reduced genetic heterogeneity and elevated

inbreeding in modern genomes compared to historical genomes (Figure 1d and Figure 2c).

Despite asymmetric sample sizes between historical and modern genomes (e.g., 11 versus 41),

this sampling scheme is unlikely to introduce overt bias as both groups contained more than

the minimum sample size of 4-6 individuals required for accurate estimations of genomic

characteristics [28,29]. Our findings suggest that the black-necked crane remains at high

genetic risk, with a contemporary N_e (e.g., 127) well below the generally accepted safe

genetic threshold (e.g., 500; [30]), suggesting potential vulnerability to changing

environments [30]. These results stand in marked contrast to the recent removal of this

species from the threatened category of the IUCN Red List in 2020 [31].

According to the small population paradigm, the recent and sudden bottleneck likely

trapped the black-necked crane in an extinction vortex by accelerating the accumulation of

genetic load [13]. Indeed, we observed a post-bottleneck increase in realized genetic load

(homozygous sites) (Figure 3c) alongside a decrease in the masked genetic load

(heterozygous sites) (Figure 3b). This shift is expected as increased homozygosity, facilitated

by inbreeding, converts masked genetic load into realized genetic load [13]. Moreover, this

conversion is anticipated to release harmful variants hidden in heterogeneity in the historical

genomes, thereby intensifying the deleterious nature of realized genetic load in the modern

genomes and elevating the risk of inbreeding depression and extinction (Figure 4c-d).

In sharp contrast, the black-necked crane has demonstrated a rapid recovery following the

bottleneck, with population numbers rebounding to 15 000 individuals by 2020 [22],

indicating limited inbreeding depression. We propose that the rapid purging of strongly

deleterious mutations likely drove this unexpected population trajectory. Notably, our

observations revealed that homozygous High-impact alleles were proportionally less enriched

inside than outside ROHs in the modern genomes (Figure 3d), consistent with genetic purging

of strongly deleterious mutations during inbreeding. Furthermore, our empirical data (Figure

4b) and forward simulations (Figure 5a-b) demonstrated a temporally consistent level of

purifying selection on homozygotes pre- and post-bottleneck. This likely acted as a filter to

purge alleles with large recessive deleterious effects on fitness, a decisive source of

inbreeding depression [8] due to increased exposure to homozygotes. The purging of strongly

deleterious alleles by efficient purifying selection may have enabled the black-necked crane to

escape an expected extinction vortex, a mechanism that may also apply to other organisms

with similar rapid demographic recoveries [16,17].

Based on comprehensive examination of the potential genetic load dynamics of a sudden

and severe population collapse followed by an unexpectedly fast recovery, our study sheds

light on the genetic consequences of contemporary widespread and rapid population declines.

Our findings stand in contrast to the existing small population paradigm by demonstrating an

escape from a potential extinction vortex following drastic population decline. Nevertheless,

consistent with theoretical predictions [5], forward simulations suggested that the duration of

population bottlenecks may alter this genetic consequence. For example, a prolonged

bottleneck lasting for five generations may lead to relaxed purifying selection and the

accumulation of strongly deleterious mutations, thereby elevating the risk of inbreeding

depression and extinction (Figure 5d-e). Thus, rapid population recovery may serve as both

the cause and consequence of the unusual escape of the black-necked crane from an extinction

vortex. Overall, our study raises hope for the prospects of current widespread and rapid

population declines, while urging the need for active and effective conservation measures to

reverse the trend before it is too late.

Methods and Materials

Sampling and sequencing

We collected blood tissue from a rescued female black-necked crane in Yunnan Province,

China. We extracted high molecular weight genomic DNA using a QIAGEN® Genomic Kit

(Cat#13343, QIAGEN) for shotgun short-read and PacBio long-read sequencing. For shotgun

short-read data, we generated 151 bp paired-end reads from a 350 bp insert-size library using

the MGISEQ-T7 platform and obtained 47.13 Gb of clean reads after quality control using

fastp v0.12.6 (-n 0 -f 5 -F 5 -t 5 -T 5; [32]). For the PacBio long-read dataset, we sheared the

gross genomic DNA by g-TUBEs (Covaris, USA) and prepared SMRTbell libraries with an

average insert-size of 14.85 kb according to the standard PacBio protocols (Pacific

Biosciences, CA, USA). Raw polymerase reads were produced using the PacBio Sequel□

platform with a Sequel II Sequencing Kit 2.0 on Nextomics, resulting in 30.58 Gb of PacBio

subreads after cleaning raw reads with ccs (-min-passes 1 -min-rq 0.99 -min-length 100; [33]).

To assemble hybrid genomes into chromosomes, we generated Hi-C data for BioNano

mapping construction. The blood tissue was fixed through a 10 min incubation of the cross-

linking reaction by vacuum infiltration in a nuclear isolation buffer supplemented with 2%

formaldehyde. We terminated the crosslinking process with glycine and constructed a library

using 100 units of DpnII restriction endonuclease (GATC), followed by a sequencing

procedure with the same protocol as the second-generation sequencing data. Finally, we

obtained 76.21 Gb of indexed Hi-C clean reads after filtering using HiC-Pro v2.8.1 [34] and

fastp.

Genome assembly

We assembled a chromosome-level reference genome using the above three

complementary datasets. We initially assembled the PacBio clean reads into contigs using

Hifiasm (https://github.com/chhylp123/hifiasm), against which we mapped all PacBio clean

reads with BLASR [35] to correct potential sequencing errors. We then polished these contigs

with shotgun clean data using NextPolish (four iterations; [36]) with default parameters. To

further improve the accuracy of the initial assembly, we clustered, ordered, and oriented

contigs onto chromosomes using LACHESIS [37] with Hi-C clean data (-

cluster_min_re_sites=100, -cluster_max_link_density=2.5, -

cluster_noninformative_rative=1.4, -order_min_n_res_in_trunk=60, -

order_min_n_res_in_shreads=60). We assessed the completeness of the final assembly (Gnig-

Y8) by searching 8□338 single-copy orthologs in the aves_odb10 dataset with BUSCO v4.0.5

[38]. We preliminarily obtained the annotations by mapping genes of the well-annotated

chicken genome (GenBank assembly accession: GCA_000002315.5, hereafter: chicken

genome) onto Gnig-Y8 using Liftoff v1.6.1 [39] and then removed gene models with in-frame

STOP codons using gffread utility (-V option) from cufflinks v2.2.1 [40].

Resequencing and variant calling

The population genomic data involved 41 modern individuals sampled from 2010 to 2020

as well as 11 historical specimens collected from 1956 to 1976, a period predating the recent

bottleneck during the mid-1980s (see details in Supplementary Table S2). We resequenced

these samples with a 350 bp insert-size library and 151 bp read length, removed adapters with

Cutadapt v2.10 [41], masked low-quality reads (e.g., Phred score<15) with Seqtk [42], and

performed the mapping procedure (including alignment, sorting, and deduplication) using

Burrows-Wheeler Aligner v0.7.17 [43], SAMtools v1.7 [44], and Genome Analysis Tool Kit

(GATK) v4.0.3 [45]. We identified raw variants using Sambamba v0.6.6 [46] and

implemented various filtering procedures for downstream analyses.

Genetic clustering analysis

We conducted genetic clustering analysis using the likelihood-based algorithm in Admixture v1.3.0 (with ten runs of each $K=1$ to 5) [47] and PCA in GCTA v1.92.2beta [48]. To prepare the inputs, we extracted autosomal biallelic SNPs with VCFtools v0.1.17 ('-f Qual=20/MinMQ=30' & '-min-alleles 2 -max-alleles 2 -remove-indels -mac 2 -max-missing 1.0'; [49]), then filtered for potential linkage disequilibrium using PLINK v1.0.7 ('-indep 50 5 2'; [50]), finally obtaining 560 083 putatively unlinked SNPs.

Demographic reconstruction

We estimated the recent population history of the black-necked crane using GONE [51] based on linkage disequilibrium data from modern samples. To scale demographic events in calendar time, we assumed a generation length of 10 years [52] and estimated the recombination rate using the following procedure. We first calculated the population recombination rate ($\rho=4N_er$, where r is the recombination rate per generation) with a 50 kb nonoverlapping window using the R package FastEPRR [53]. We prepared inputs by filtering the raw variants from Sambamba using VCFtools with the following parameters: '-f Qual=20/MinMQ=30' & '-max-missing 1.0 -minDP 5 -min-alleles 2 -max-alleles 2 -remove-indels -maf 0.05', finally obtaining 1 319 028 automatic biallelic SNPs. We also calculated the population mutation rate ($\theta=4N_e\mu$, where μ is the recombination rate per generation) using ANGSD v0.928 ('-doSaf 1 -remove_bads -only_proper_pairs 1 -minMapQ 30 -minQ 20 -minInd 41 | realSFS | thetaStat'; [54]) with the same window size based on the mapped BAM files. We finally estimated the recombination rate as 3.42×10^{-8} using the equation $r=(\rho/\theta)\times\mu$, assuming an μ of 1.45×10^{-8} per site per generation [55].

We also estimated contemporary N_e using a linkage disequilibrium method in NeEstimator [56]. To address the reduced precision associated with large and linked loci, we thinned the dataset previously used in the GONE analysis to a minimum distance of 50 kb, yielding a set of 21 590 SNPs.

Genomic diversity and inbreeding estimation

We compared genetic diversity between the historical and modern samples using nucleotide diversity (π) and heterogeneity (h). We performed both analyses based on

2□829□286 biallelic data points, excluding singletons, with the VCFtools command: -max-
missing 1.0 -mac 2 -minDP 3 -min-alleles 2 -max-alleles 2 -remove-indels. We calculated
genome-wide π using VCFtools with a 50 kb window and 20 kb steps. We counted
heterogeneous sites for each individual and measured the *h* index by dividing the counts by
the autosomal genome length (1.17 Gb).

To assess the level of inbreeding in the black-necked crane, we calculated the length and
number of ROHs for modern and historical genomes with PLINK with the following
parameters: "-homozyg-window-snp 50 -homozyg-snp 50 -homozyg-window-missing 3 -
homozyg-kb 100 -homozyg-density 1000". We calculated the inbreeding coefficient (F_{ROH})
for each individual by dividing the sum length of ROHs by the autosomal genome length
(1.17 Gb). Higher F_{ROH} values are indicative of higher inbreeding levels.

Genetic load calculation
We estimated genetic load dynamics between the modern and historical genomes based
on functional effects of SNPs in coding regions and conservation scores under strict
evolutionary constraints. Both analyses were based on the same dataset as in the *Genomic*
diversity and inbreeding estimation section, with ancestral states polarized by genomic data of
the hooded crane (*Grus monachus*) (GenBank assembly accession: GCA_012487855.1), the
sister species of the black-necked crane [57].

First, we built a database based on the Gnig-Y8 annotation and categorized the effects of
coding variants into four impact levels using SnpEff v4.3.68 [26]: a) High: variants assumed
to have disruptive impact on the protein (e.g., loss-of-function and frameshift variants); b)
Moderate: nondisruptive variants that may change protein effectiveness (e.g.,
inframe_deletion); c) Low: variants mostly harmless or unlikely to change protein behavior
(e.g., synonymous variants); and d) Modifier: usually noncoding variants or variants affecting
noncoding genes (e.g., downstream variants). We counted the number of total variants and
respective proportions of homo- and heterozygotes (multiplied by 2 in homozygous positions)
in the High and Moderate categories and compared the total, realized, and masked genetic
loads between the historical and modern genomes.

Second, we estimated the relative risk of deleterious burden, $R_{modern/historical}$ [58], based on
the derived allele frequency of deleterious mutations from the historical and modern genomes.
We conducted this estimation for the realized, masked, and total genetic loads using a script

brrAB developed by Zheng et al. (2023) [59]. Here, R $_{modern/historical}$ equal to 1 corresponds to

no change in frequency between historical and modern genomes, while R $_{modern/historical}$<1 or >1

indicates an increase or a decrease in frequency in historical genomes relative to modern

genomes, respectively. The distributions in the R $_{modern/historical}$ variance were estimated based

on jack-knifing across chromosomes.

Third, to test the hypothesis that deleterious variants in ROH tracts may reduce individual

fitness with additive effects [5], where purging, if any, is expected to be more common than

in the rest of genome, we compared the ratio of High-impact (obtained from SnpEff output) to

all homozygous loci in ROH and non-ROH portions of each modern genome.

Selection coefficient estimation

We employed three approaches to compare purifying selection coefficients between

historical and modern genomes. First, we estimated the coefficients for each individual using

conservation scores. In brief, we downloaded GERP scores computed for 27 sauropsids by

multiple whole‑genome alignment [60] and converted chicken-genome-focused scores to the

black-necked crane genome using the LiftOver program (http://genome.ucsc.edu/cgi-

bin/hgLiftOver). We then compared the top 100 GERP (mGERP) scores in homo- and

heterozygotes between the historical and modern genomes to detect dynamics of purifying

selection coefficients over the bottleneck.

Second, we conducted forward simulations using SLiM v2.0.1 [61]. In brief, we modeled

individuals with a diploid genome of 2 Mb, composed of 2 000 genes, each 1 kb in length,

across 34 pseudochromosomes mirroring the relative proportions and positions observed in

the Gnig-Y8 assembly, assuming: a) a per-generation mutation rate of 1.45×10^{-8} [55]; b) a

recombination rate of 3.42×10^{-8}, as estimated in the previous *Demographic reconstruction*

section between genes and no intragenic recombination; c) a relative occurrence ratio of 7:3

for neutral and deleterious mutations; and d) a gamma distribution of fitness (mean=0.03,

standard deviation (sd)=0.1) used as selection coefficients of deleterious mutations [62]. We

simulated neutral and deleterious variation following a 100 000 generation burn-in period,

with three distinct scenarios: a) a pre-bottleneck population with an N_c of 10 000 individuals

given an empirical N_e/N_c ratio of 0.1 and a pre-bottleneck N_e level of approximately 1 000; b)

a post-bottleneck population recapitulating the demographic history of the black-necked crane,

decreasing from N_c=10 000 to 100 individuals in a generation and rebounding immediately

to 15 000 individuals over four generations; and c) a scenario similar to the second, but with

a prolonged bottleneck (e.g., with 100 individuals) lasting five generations. For each scenario,

we performed a range of dominance coefficients, including entirely recessive (h=0), partially

recessive (h=0.25), and entirely additive (h=0.5). We randomly subsampled 50 individuals

from the last simulation step and compared their mFEs of deleterious mutations for homo-

and heterozygous mutations.

References

1. Ceballos G, Ehrlich P R, Barnosky A D, et al. (2015). Accelerated modern human-induced

species losses: Entering the sixth mass extinction. Sci Adv, 1: e1400253.

2. Ceballos G, Ehrlich P R, Dirzo R. (2017). Biological annihilation via the ongoing sixth

mass extinction signaled by vertebrate population losses and declines. Proc Natl Acad Sci

U S A, 114: E6089-E6096.

3. Rosenberg K V, Dokter A M, Blancher P J, et al. (2019). Decline of the North American

avifauna. Science, 366: 120-124.

4. WWF. (2020). Living planet report 2020 bending the curve of biodiversity loss. (R. E. A.

Almond, M. Grooten, & T. Petersen, Eds.). WWF.

5. Bertorelle G, Raffini F, Bosse M, et al. (2022). Genetic load: genomic estimates and

applications in non-model animals. Nature Rev Genet, 23: 492-503.

6. Lynch M. (2010). Evolution of the mutation rate. Trends Genet, 26: 345-352。

7. Bortoluzzi C, Bosse M, Derks M F L, et al. (2020). The type of bottleneck matters: Insights

into the deleterious variation landscape of small managed populations. Evolutionary

applications, 13: 330-341.

8. Hedrick P W, Garcia-Dorado A. (2016). Understanding inbreeding depression, purging,

and genetic rescue. Trends Ecol Evol, 31: 940-952.

9. von Seth J, van der Valk T, Lord E, et al. (2022). Genomic trajectories of a near-extinction

event in the Chatham Island black robin. BMC genomics, 23: 1-17.

10. Caballero A, Bravo I, Wang J. (2017). Inbreeding load and purging: implications for the

short-term survival and the conservation management of small populations. Heredity, 118:

177-185.

11. Robinson J A, Brown C, Kim B Y, et al. (2018). Purging of strongly deleterious mutations

explains long-term persistence and absence of inbreeding depression in island foxes.

Current Biology, 28: 3487-3494.e4.

12. Robinson J A, Kyriazis C C, Nigenda-Morales S F, et al. (2022). The critically

endangered vaquita is not doomed to extinction by inbreeding depression. Science, 376:

635-639.

13. Ouborg N J, Pertoldi C, Loeschcke V, et al. (2010). Conservation genetics in transition to

conservation genomics. Trends Genet, 26: 177-187.

14. Feng S, Fang Q, Barnett R, et al. (2019). The genomic footprints of the fall and recovery

of the crested ibis. Curr Biol, 29: 340-9.e7.

15. van der Valk T, Díez-Del-Molino D, Marques-Bonet T, et al. (2019). Historical genomes

reveal the genomic consequences of recent population decline in eastern gorillas. Curr Biol,

29: 165-70.e6.

16. Xu G, Lv Z, Zhang S. (2022). 100+ biodiversity positive practices and actions around the

world. China Environment Publishing Group, Beijing.

17. Suckling K, Greenwald N, Curry T. (2012). On time, on target: how the Endangered

Species Act is saving America's wildlife. Center for Biological Diversity, Tucson, AZ.

18. Yu Y, He G, Li D Y, et al. (2022). Climate change challenge, extinction risk, and

successful conservation experiences for a threatened primate species in China: golden

snub-nosed monkey (*Rhinopithecus roxellana*). Zoological Research, 43: 940.

19. Archibald G W, Meine C D, Kirwan G M. (2020). Black-necked Crane (*Grus nigricollis*),

version 1.0. In Birds of the World (Hoyo J del, Elliott A, Sargatal J, Christie D A, and

Juana E de, Editors). Cornell Lab of Ornithology, Ithaca, NY, USA.

20. He P, Kong D J, Liu Q, Yu H Z, Zhao J L, Yang X J. (2011). Roosting-site characteristics

of wintering black-necked cranes (*Grus nigricollis*) at Napahai, Yunnan. Zoological

Research, 32: 150-156.

21. Matthiessen P. (1994). At the end of Tibet. Audubon, 96: 10-49.

22. China Wildlife Conservation Association, Bird Media, Ltd. Cranes in China. Posts and

Telecommunications Press, Beijing.

23. Zhou C, Yu H, Geng Y, et al. (2019). A high-quality draft genome assembly of the black-

necked crane (*Grus nigricollis*) based on nanopore sequencing. Genome biology and

evolution, 11: 3332-3340.

24. Santiago E, Novo I, Pardiñas A F, et al. (2020). Recent demographic history inferred by
high-resolution analysis of linkage disequilibrium. Molecular Biology and Evolution, 37:
3642-3653.
25. Reid B N, Pinsky M L. (2022). Simulation-based evaluation of methods, data types, and
temporal sampling schemes for detecting recent population declines. Integrative and
Comparative Biology, 62: 1849-1863.
26. Cingolani P, Platts A, Wang L L, et al. (2012). A program for annotating and predicting
the effects of single nucleotide polymorphisms, SnpEff: SNPs in the genome of Drosophila
melanogaster strain w1118; iso-2; iso-3. Fly, 6: 80-92.
27. Frankham, R. (1995). Effective population size/adult population size ratios in wildlife: a
review. Genetics Research, 66: 95-107.
28. Qu W M, Liang N, Wu Z K, et al. (2020). Minimum sample sizes for invasion genomics:
Empirical investigation in an invasive whitefly. Ecology and evolution, 10: 38-49.
29. Li H, Qu W, Obrycki J J, et al. (2020). Optimizing sample size for population genomic
study in a global invasive lady beetle, Harmonia axyridis. Insects, 11: 290.
30. Hoban S, Bruford M, Jackson J D U, et al. (2020). Genetic diversity targets and indicators
in the CBD post-2020 Global Biodiversity Framework must be improved. Biological
Conservation, 248: 108654.
31. BirdLife International. (2023). Species factsheet: *Grus nigricollis*. Downloaded from
http://datazone.birdlife.org/species/factsheet/black-necked-crane-grus-nigricollis on
28/07/2023.
32. Chen S, Zhou Y, Chen Y, et al. (2018) fastp: an ultra-fast all-in-one FASTQ preprocessor.
Bioinformatics, 34: i884-i890.
33. https://github.com/PacificBiosciences/ccs.
34. Servant N, Varoquaux N, Lajoie B R, et al. (2015). HiC-Pro: an optimized and flexible
pipeline for Hi-C data processing. Genome Biol, 16: 259-70.
35. Chaisson M J, Tesler G. (2012). Mapping single molecule sequencing reads using basic
local alignment with successive refinement (BLASR): application and theory. BMC
Bioinformatics, 13: 238.
36. Hu J, Fan J, Sun Z, et al. (2020). NextPolish: a fast and efficient genome polishing tool for
long-read assembly. Bioinformatics, 36: 2253-2255.

37. Burton J N, Adey A, Patwardhan R P, et al. (2013). Chromosome-scale scaffolding of de
novo genome assemblies based on chromatin interactions. Nature biotechnology, 31: 1119-
1125.

38. Simão F A, Waterhouse R M, Panagiotis P, et al. (2015). BUSCO: assessing genome
assembly and annotation completeness with single-copy orthologs. Bioinformatics, 31:
3210-3212.

39. Shumate A, Salzberg S L. (2021). Liftoff: accurate mapping of gene annotations.
Bioinformatics, 37: 1639-1643.

40. Trapnell C, Williams B A, Pertea G, et al. (2010). Transcript assembly and quantification
by RNA-Seq reveals unannotated transcripts and isoform switching during cell
differentiation. Nat Biotechnol, 28: 511-515.

41. Martin M. (2011). Cutadapt removes adapter sequences from high-throughput sequencing
reads. EMBnet. Journal, 17: 10-12.

42. Yun S, Yun S J. (2014). Masking as an effective quality control method for next-
generation sequencing data analysis. BMC Bioinformatics, 15: 382.

43. Li H, Durbin R. (2009). Fast and accurate short read alignment with Burrows-Wheeler
transform. Bioinformatics, 25: 1754-1760.

44. Li H, Handsaker B, Wysoker A, et al. (2009). 1000 Genome Project Data Processing
Subgroup. The Sequence Alignment/Map format and SAMtools. Bioinformatics, 25: 2078-
2079.

45. McKenna A, Hanna M, Banks E, et al. (2010). The Genome Analysis Toolkit: a map
reduce framework for analyzing next-generation DNA sequencing data. Genome Res, 20:
1297-1303.

46. Tarasov A, Vilella A J, Cuppen E, et al. (2015). Sambamba: fast processing of NGS
alignment formats. Bioinformatics, 31: 2032-2034.

47. Alexander D H, Novembre J, Lange K. (2009). Fast model-based estimation of ancestry in
unrelated individuals. Genome Res, 19: 1655-1664.

48. Yang J, Lee S H, Goddard M E, et al. (2011). GCTA: a tool for genome-wide complex
trait analysis. Am J Hum Genet, 88: 76-82.

49. Danecek P, Auton A, Abecasis G, et al. (2011). The variant call format and VCFtools.
Bioinformatics, 27: 2156-2158.

50. Purcell S, Neale B, Todd-Brown K, et al. (2007). PLINK: a tool set for whole-genome
association and population-based linkage analyses. American Journal of Human Genetics,
81: 559-75.

51. Santiago E, Novo I, Pardiñas A F, et al. (2020). Recent demographic history inferred by
high-resolution analysis of linkage disequilibrium. Molecular Biology and Evolution, 37:
3642-3653.

52. Bird J P, Martin R, Akçakaya H R, et al. (2020). Generation lengths of the world's birds
and their implications for extinction risk. Conserv. Biol, 34: 1252-1261.

53. Gao F, Ming C, Hu W, et al. (2016). New software for the fast estimation of population
recombination rates (FastEPRR) in the genomic era. G3: Genes, Genomes, Genetics, 6:
1563-1571.

54. Korneliussen T S, Albrechtsen A, Nielsen R. (2014). ANGSD: analysis of next generation
sequencing data. BMC bioinformatics, 15: 356.

55. Zhang G, Li C, Li Q, et al. (2014). Comparative genomics reveals insights into avian
genome evolution and adaptation. Science, 346: 1311-1320.

56. Do C, Waples R S, Peel D, et al. (2014). NeEstimator v2: re□implementation of software
for the estimation of contemporary effective population size (N_e) from genetic data.
Molecular ecology resources, 14: 209-214.

57. Krajewski C, Sipiorski J T, Anderson F E. (2010). Complete mitochondrial genome
sequences and the phylogeny of cranes (Gruiformes: Gruidae). The Auk, 127: 440-452.

58. Xue Y, Prado-Martinez J, Sudmant P H, et al. (2015). Mountain gorilla genomes reveal
the impact of long-term population decline and inbreeding. Science, 348: 242-245.

59. Zheng C, Chen Q, Huang S, et al. (2023). Chromosome-scale genome assembly and
conservation genomics reveals purifying selection and adaptation in a critically endangered
bird. Authorea Preprints.

60. https://ftp.ensembl.org/pub/current_compara/conservation_scores/27_sauropsids.gerp_con
servation_score/

61. Haller B C, Messer P W. (2017). SLiM 2: flexible, interactive forward genetic simulations.
Mol. Biol. Evol, 34: 230-240.

62. Kim B Y, Huber C D, Lohmueller K E. (2017). Inference of the distribution of selection
coefficients for new nonsynonymous mutations using large samples. Genetics, 206: 345-
361.

Data availability

The scripts used in the paper were deposited in the Science Data Bank (doi:10.57760/sciencedb.10054). Genomic data were archived in the Genome Warehouse (accession number: GWHDTVR00000000) and Genome Sequence Archive (accession numbers: CRA012191 and CRA012234).

Acknowledgments

The authors would like to thank Dr. Qiang Liu, Dr. Dejun Kong, Dr. Kai Wang, Dr. Peng He, Mr. Weixiong Luo, and Ms. Mengyin An for efforts in collecting DNA samples. This work was supported by the National Key R&D Program of China (2022YFC2601200) and Second Tibetan Plateau Scientific Expedition and Research (STEP) program (2019QZKK0501)

Author contributions

F.D. conceived the ideas; N.C., X.T.M., F.D., and H.Q.W conducted the analyses; X.C.H., H.Q.W., L.X.Z., F.M.L., L.Y., D.Y., and X.J.Y. collected DNA samples; F.D. led the writing, with substantial input from C.M.H., which was approved by all other authors.

Competing financial interest statement

The authors declare no competing financial interests.

Figure legends

Figure 1 Sampling locations and population structure of black-necked crane

a: Ecological photograph (credit to Zhaxi). b: Sampling localities for historical and modern

samples. c: Scatter diagrams for cross-validation (cv) errors in admixture analyses, with K

(number of supposed genetic clusters) from 1 to 5. d: Principal component analysis (PCA)

plots for all historical and modern samples.

Figure 2 Demographic history, genomic diversity, and inbreeding estimation for black-

necked crane

a: Population trajectory of recent census population size (N_c; orange, data retrieved from ref[22])

and effective population size (N_e; blue) with time in unit of generations. b: Boxplot of

individual genome-wide heterozygosity. c: Individual inbreeding coefficients estimated as

F_{ROH}. For (b) and (c), horizontal lines within boxplots and bounds of boxes represent 1st and

3rd quartiles (box), median (line inside box) outliers (open circle), and 1.5× interquartile

range.

Figure 3 Genetic load estimation on coding regions for black-necked crane

A: Total genetic load; B: Masked genetic load; C: Realized genetic load; and D: Relative

genetic load inside and outside ROHs based on High-impact category from SnpEff analysis.

Boxplot shows 1st and 3rd quartiles (box), median (line inside box), outliers (open circle),

and 1.5× interquartile range.

Figure 4 Selection coefficient estimation based on mGERP scores for the black-necked

crane

Comparisons were conducted for (A) heterozygotes and (B) homozygotes between historical

and modern genomes, as well as for individual genomes among (C) historical and (D) modern

genomes. Boxplot comprises 1st and 3rd quartiles (box), median (line inside box), outliers

(open circle), and 1.5× interquartile range.

Figure 5 Forward simulations on selection coefficient (s) for deleterious mutations in

homozygotes of black-necked crane

Simulations recapitulated demographic history (A-B) and supposed a prolonged bottleneck

(C-D) under fully recessive (A and C) and partially recessive (B and D). Boxplot comprises

1st and 3rd quartiles (box), median (line inside box), and 1.5× interquartile range.