Rhizobacteria inoculation of plants for reducing insect pests: A meta-analysis on insect behaviour and fitness

Sharon E Zytynska[1,2], Megan Parker[1], Oriana Sanchez-Mahecha[2]

1. Department of Evolution, Ecology, and Behaviour, Institute of Infection, Veterinary and Ecological Sciences, University of Liverpool, UK. 2. Technical University of Munich, Terrestrial Ecology Research Group, Department for Life Science Systems, Germany

Abstract

Rhizobacteria inoculation of plants has shown promising potential for enhancing resistance against insect pests by reducing herbivore fitness and altering herbivore-natural enemy interactions. Understanding interactions among specific rhizobacterial species and their plants in the rhizosphere is crucial for developing effective strategies to harness these benefits for pest management in agriculture. We present a meta-analysis examining the impact of rhizobacteria inoculation on herbivore interactions with plants. The findings indicate that rhizobacteria inoculation generally reduces herbivore fitness and host choice behaviours. Rhizobacteria inoculation may also enhance the recruitment of natural enemies of herbivores, thus increasing top-down predator control. The effects on herbivores varied significantly depending on the rhizobacterial species, with *Bacillus* spp. showing stronger effects compared to other commonly studied *Pseudomonas* spp. Rhizobacteria notably reduced traits such as host choice, leaf consumption, survival, and reproduction of chewing herbivores, while primarily impacting sucking herbivores by reducing reproduction. Single-strain inoculants tended to perform better, especially for sucking herbivores, suggesting potential strain incompatibility issues with multi-strain inoculants. Furthermore, field trials showed less impact on insect fitness reduction compared to experiments under controlled conditions, possibly due to soil diversity and environmental factors affecting inoculant persistence. These results underscore the need for considering broader environmental interactions when developing effective rhizobacteria-based pest management strategies. Understanding specific and generalist rhizosphere interactions can aid in developing synthetic microbial communities with broad protective functions across various plants and environments.

Introduction

The rhizosphere is the narrow region of soil directly influenced by plant roots, hosting a dynamic and diverse microbial community with up to 10^{11} microbial cells per gram root and more than 30,000 prokaryotic species (Berendsen, Pieterse & Bakker 2012). This zone plays a vital role in sustaining plant health by fostering interactions between roots and a myriad of microorganisms. Plants influence the rhizosphere microbiome by releasing root exudates and altering soil conditions, selectively recruiting distinct microbial communities tailored to a plant's unique physiological and biochemical traits (Berendsen, Pieterse & Bakker 2012; Trivedi *et al.* 2020). Recruitment of rhizosphere bacteria can occur over short-time frames, enabling a plant to efficiently respond to changes in the environment, e.g. from drought or nutrient stress

(Bakker *et al.* 2013). Recent syntheses examining the effect of rhizobacterial inoculation on plant responses to drought (Rubin, van Groenigen & Hungate 2017; Zhao *et al.* 2022), temperature (Zhang *et al.* 2023), salt (Pan *et al.* 2019), phosphate (De Zutter *et al.* 2022) and across different experimental factors (Zeffa *et al.* 2020; Li *et al.* 2022; da Silva *et al.* 2024) have demonstrated the strong and broad impact that different rhizobacteria can have on plant fitness and yield.

Plants can also respond to herbivory through recruitment of different rhizobacteria communities, that enhance plant defences responses, thereby decreasing insect population growth rates (Pineda *et al.* 2010). Even before herbivory, plants can be primed to respond more quickly upon insect feeding through application of various plant hormones but also inoculation with defence-inducing rhizobacteria (Pieterse *et al.* 2013; Pieterse *et al.* 2014; Selosse, Bessis & Pozo 2014). Priming of a plant involves structural changes in the histones that fold DNA, leading to a quicker initiation of RNA transcription and faster synthesis of defence-related compounds. The mechanisms underlying this vary across plant and herbivore species, with variable induction of defences by various rhizobacteria. Plant responses to chewing insects, e.g. caterpillars, differ from responses to sucking insects, e.g. aphids or whitefly(Wu & Baldwin 2010; Schuman & Baldwin 2016). These different plant responses also lead to an expectation, and finding, that not all rhizobacteria will be effective against all insect herbivores (Blubaugh *et al.* 2018; Friman *et al.* 2021).

Here we present a meta-analysis to synthesise published data measuring effects of bacterial inoculation of plants on insect behaviour and fitness, comparing chewing and sucking insects. In addition to defence induction, microbial inoculation and herbivory can also trigger changes in plant volatiles that can recruit natural enemies of the herbivores. While Herbivore-Induced Plant Volatiles (HIPVs) in plants are well-known to attract natural enemies, the consequences of microbial-induced plant volatiles on herbivore or natural enemy behaviour is relatively unknown. We also examined parasitoid wasp and predator behaviour in response to bacterial-inoculation of plants. This meta-analysis aims to describe general effects of experimental microbial inoculation of plants on insect behaviour and fitness, with a focus on rhizosphere bacteria effects on chewing and sucking insects.

Our main hypotheses are:
1. The effect of microbial inoculation on insect responses varies across insect feeding groups
2. The type of inoculation (single or community) as well as the main bacterial genus in the inoculant will impact insect responses
3. The experimental environment (controlled vs field) will alter the strength of response of the insect

Methods

We used two broad and inclusive search approaches to maximise identification of papers for inclusion. The first search was compiled in ISI Web of Science up to September 2023 using the following keywords combinations: (Rhizo* AND aphid*), (Rhizo* AND Whitefly*), (Rhizo* AND cater*), (Rhizo* AND lepido*), (Rhizo* AND herbivor*), resulting in 428 papers. We used separate terms for aphid, whitefly, caterpillars, and lepidoptera as these are well-known species studied in this area, and the term herbivore was used to encompass a wider range.

The second search used three sub-searches using terms 1. ("Insect*" AND ("volatile*" OR "terpen*" OR "sesquiterp*" OR "floral volatile*" OR "leaf volatile*") AND ("microb*" OR "bacteri*" OR "rhizo*" OR "fungi*") AND ("behav*" OR "attract*" OR "repel*" OR "pref*" OR "oviposit*" OR "feed*"), 2. ("Insect*" OR "herbiv*" OR "behav*" OR "parsit*") AND ("volatile*" OR "terpen*" OR "sesquiterp*" OR "floral volatile*" OR "leaf volatile*") AND ("microb*" OR "bacteri*" OR "fungi*") AND ("plant*") AND ("behav*" OR "attract*" OR "repel*" OR "pref*"), 3. ("Insect*" OR "herbiv*" OR "behave*" OR "parsit*") AND ("volatile*" OR "terpen*" OR "sesquiterp*" OR "floral volatile*" OR "leaf volatile*" OR "secondary metabol*") AND ("microb*" OR "bacteri*" OR "fungi*") AND ("behav*" OR "attract*" OR "repel*" OR "pref*") resulting in 1615 papers. Following assessment of title and abstract, we identified 180 potential papers of interest, from which 70 provided suitable data for inclusion in the meta-analysis (Fig. 1).

We screened the studies using the following inclusion criteria: (1) data must come from experimental or field studies that tested the bacteria effect on insects comparing a bacteria-treated with control untreated plants; (2) data are presented to enable extraction of mean values, a measure of variance, and sample size; (c) measured variables are related to insect behaviour (preference or consumption rate) or fitness (insect survival, development time, body size, fecundity, or lifespan. We extracted the mean, variance, and sample size across the treated and untreated groups. We also extracted experimental information on bacteria inoculum (single strain or multi-strain community), microbial genus/species (multi-genus noted for mixed-genera community inoculations), insect feeding type (chewing herbivore, sucking herbivore, parasitoid wasps or predator), insect family/species, plant family/species, inoculation method (soil, seed, root, leaf), and study environment (field, greenhouse, or growth chamber). Following in-depth screening, 44 papers were removed due to not containing primary data, 66 removed due to lack of informative data across treatments and/or controls, resulting in a final data set of 344 data points extracted from 70 papers published between 1997 and 2023 fulfilling all the inclusion criteria.

Fig. 1. Summary of selection process for final papers for inclusion in the meta-analysis, as based on PRISMA guidelines

In studies with multiple genotypes, strains, or varieties, data were pooled within species due to lack of replication to examine specific effects. We selected only data from the "wildtype" microbial strain where data were presented also on lab-generated mutants. For controlled environment time-series experiments we chose the time that related best to the other studies in the dataset; for example, the number of offspring produced by aphids is commonly measured at day 10 or 14. For field data, we averaged across multiple sampling time points.

Data analysis

The meta-analysis was performed in RStudio version 2023.12.1 (RStudio Team 2020) using R version 4.3.0 (R Core Team 2024) using the package metaphor (Viechtbauer 2010). Effect sizes were calculated using the standardized mean difference with an unbiased estimate of the sampling variances (SMDH, giving Hedges' g). This measurement compares the treated plants (inoculated with the bacteria and exposed to the herbivore insect) to the control plants (that were just exposed to the herbivore insect). A priori power analyses (medium effect size d=0.5) were calculated in R following Quintana and Tiebel (2019) for the main insect variables within the analysed data subset. For the power analysis, we used the number of effect sizes from the extracted data (data points), the average number of replicates used to calculate each effect size (control vs treated plant), against a predicted effect size of d = 0.5 (medium) and low heterogeneity (h = 0.33).

The effect of the inoculated bacteria on the insect traits was analysed with a meta-analytic linear mixed effect model (rma.mv). Study was included as a random effect to account for multiple data points across insect, plant, and bacteria species within each study. In each insect subset (sucking and chewing herbivores), insect responses (levels: host choice, feeding, mortality, development, body size, fecundity, lifespan) were treated as fixed effect moderators. Additional models were used to determine the effect across different plant and insect families, as well as across experimental manipulations (field vs controlled pot experiments). Importantly, only results are presented when there were at least three data points from three independent studies for the comparisons. We evaluated publication bias in the dataset by assessing the funnel-plot asymmetry and Eggers test. For the data visualisation, figures show both the mean effect size and 95% intervals obtained from the linear mixed effect models output, as well as the individual-level data as a bubble plot with sizes relative to the precision of the estimate (1/SE), where larger bubbles show a higher certainty of the data.

Results

Effects of rhizobacteria inoculation of plants on associated insects

The herbivore final dataset contained 344 data points from 66 papers, representing 215 data points (38 papers) from chewing herbivores and 129 data points (31 papers) from sucking herbivores, with data on all responses identified. Across all measured response traits in all herbivore species, rhizobacteria inoculation of plants had a negative effect on insect herbivores (hedges g = -0.407, P<0.001, n=348). Yet, the strength and significance of these effects varied between chewing (hedges g = -0.594, P<0.001, n=219) and sucking herbivores (hedges g = -0.182, P=0.209, n=129), as well as across the different response traits measured (test of moderators QM=147.4, df=14, P<0.001; Fig.1a,b). A large residual heterogeneity in this model (QM=1445.9, df=366, P<0.001) indicated the need to include additional factors in

the model (rhizobacteria/plant/insect species or system information) and therefore in following sections we examined chewing and sucking herbivore datasets separately.

We also extracted 31 data points from eight papers with information on rhizobacterial effects on insect natural enemies (predators and parasitoid, examining variables including host choice, feeding behaviour, and fecundity; Fig. 1c). While there were no general significant effects due to a lack of data (hedges g = 0.416, P=0.385, n=31), there is a potential pattern of increased host choice to inoculated plants from pot and field studies (hedges g = 0.564, P=0.322, n=19).

Fig. 2. Effect of rhizobacterial inoculation across response traits measured for (a) chewing herbivores, (b) sucking herbivores and (c) natural enemies (parasitoid wasps and predators). Effect size mean and 95% confidence intervals from the linear mixed effect models output, individual study data shown by coloured bubble plots with sizes relative to the precision of the estimate (1/SE), where larger bubbles show a higher certainty of the data.

Effect of rhizobacteria inoculation on chewing insects

The chewing insects belonged to six difference families (Lepidoptera n=178, Chrysomelidae n=19, Anthomyiidae n=15, Cecidomyiidae n=1, Gryllotalpidae n=1, Sciaridae n=1) and the Lepidoptera represented nine genera, including *Spodoptera* sp. (n=48 data points, from 13 papers) and *Plutella xylostella* (n=67 data points, from 4 papers); with a tendency for a small number of papers to examine a broad range of rhizobacteria leading to a loss of independence in these effect estimates.

Across the different response traits measured for chewing herbivores, body size was the most measured variable but showed no strong significant reduction due to rhizobacteria inoculation of plants (hedges g = -0.170, P=0.248, n=72, studies=24). However, *Bacillus* inoculations had a significant negative impact on body size (hedges g = -0.718, P<0.001, n=24, studies=6), while *Pseudomonas* and 'Other' inoculations (representing nine genera) did not (*Pseudomonas*: hedges g = -0.074, P=0.671, n=39, studies=13; Other: hedges g = -0.062,

P=0.804, n=9, studies=6; Fig. 3a). Similarly, only *Bacillus* inoculants reduced insect survival but examined in just two studies (hedges g = -0.602, P=0.028, n=11, studies=2; Fig. 3a). Insect development time was positively affected by *Bacillus* inoculation of the host plant, representing shorter development time, again from only two studies (hedges g = 0.561, P=0.026, n=19, studies=2). Effects on insect behaviour were observed across the range of bacterial inoculants leading to overall significant reductions in the number of insects landing on an inoculated host plant (host choice: hedges g = -0.636, P=0.006, n=25, studies=8) with subsequent reduction in leaf consumption (feeding: hedges g = -0.965, P<0.001, n=26, studies=8) (Fig 3a).

Fig. 3. Effect of rhizobacterial inoculation on chewing herbivores, separated by *Bacillus* spp inoculants (red), *Pseudomonas* spp. inoculants (blue), and other rhizobacteria (grey) across (a) response traits (b) main insect families, and within Lepidoptera and (c) host plant family. Effect size mean and 95% confidence intervals from the linear mixed effect model outputs, individual data shown as bubble plots with sizes relative to the precision of the estimate (1/SE), where larger bubbles show a higher certainty of the data.

Of the six chewing herbivore families studied, rhizobacteria inoculation significantly reduced chewing herbivore fitness within Chrysomelidae (hedges g = -0.635, P=0.018, n=19, studies =8) and Lepidoptera (hedges g = -0.382, P=0.008, n=178, studies =26) across a diverse set of microbial inoculants. Chrysomelidae leaf beetles were negatively affected by non-*Bacillus* bacterial inoculants of their host plants (Other: hedges g = -0.839, P=0.021, n=5, studies=4; *Pseudomonas*: hedges g = -0.625, P=0.094, n=5, studies=2; *Bacillus* hedges g = -0.431, P=0.210, n=9, studies=4; Fig. 3a). Whereas, within Lepidoptera the strongest effects were from *Bacillus* inoculants (Spodoptera: hedges g = -0.969, P=0.043, n=26, studies=3; Other Lepidoptera: hedges g = -1.00, P=0.073, n=20, studies=3; Fig. 3b), with close to significant

effects of *Pseudomonas* inoculants on Spodoptera (hedges g = -0.525, P=0.069, n=18, studies=4; Fig. 3b). There were no significant effects on Plutella Lepidopterans (Fig. 3b).

Chewing insects were most often studied on Brassicaceae (120/215) or Poaceae (73/215) plants with both Chrysomelidae and Lepidoptera studied within both plant families. *Bacillus* inoculants within both plant families had negative effects on insects (Poaceae: hedges g = -0.818, P=0.006, n=40, studies=5; Brassicaceae: hedges g = -0.663, P=0.051, n=51, studies=4; Fig. 3c). Other microbial inoculants (including Alcaligenes, Enterobacter, and Kluyvera) also showed significant effects on chewing insects but data were collected from a single paper and thus robustness is low (hedges g = 0.826, P=0.030, n=20, studies=1; Fig. 3c). *Pseudomonas* inoculants had no overall effect within Brassicaceae (hedges g = -0.161, P=0.405, n=49, studies=12; Fig. 2c) or Poaceae (hedges g = 0.152, P=0.738, n=5, studies=3; Fig. 3c).

Effect of rhizobacteria inoculation on sucking herbivores

The data were dominated by studies on Aphididae (86/129 data points from 8 species of aphid); these included *Brevicoryne brassicae* (n=23), *Acyrthosiphon pisum* (n=18), *Myzus persicae* (n=14), *Rhopalosiphum padi* (n=13), *Lipaphis erysimi* (n=18), *Aphis glycines* (n=5), undescribed aphids (n=4), *Aphis gossypii* (n=3), and *Sitobion avenae* (n=2). The remaining data were from Tetranychidae spider-mites (n=28 on *Tetranychus urticae*), Aleyrodidae whitefly (n=7 from Bemisia tabaci and n=4 from *Aleyrodes proletella),* Triozidae psyllids (n=3, from *Bactericera cockerelli*), and n=1 from the Cicadellidae leafhopper *Amrasca biguttula.*

The data were dominated by studies assessing sucking herbivore fecundity (number of offspring produced) (73/129) which was significantly reduced by rhizobacteria inoculation of the host plants (hedges g = -0.624, P<0.001, n=73, studies=27). With combined data across all inoculants, we also found significant effects on lifespan (hedges g = -0.413, P=0.030, n=10, studies=5), body size (hedges g = -0.539, P=0.046, n=4, studies=3), and a potential negative impact on development time to adult (hedges g = -0.308, P=0.090, n=19, studies=8). Similar to the chewing herbivores, the majority of experiments on sucking herbivores used either *Bacillus* (n=71/129) or *Pseudomonas* strains (n=33/129). We found substantial variation in the number of studies and strength of effects across these different inoculants (Fig. 4a). Insect fecundity was significantly reduced by *Bacillus* (hedges g = -0.567, P=0.050, n=42, studies=12) and other inoculant species (hedges g = -0.889, P=0.006, n=14, studies=10), but not *Pseudomonas* spp. (hedges g = -0.500, P=0.111, n=17, studies=9). There was insufficient data to conduct similar comparisons for the other response traits.

The three insect families, with sufficient data for analysis, were all negatively impacted by rhizobacterial inoculation of the host plants (Aphididae: hedges g = -0.469, P=0.005, n=86, studies=24; Aleyrodidae: hedges g = -0.767, P=0.027, n=11, studies=5; Tetranychidae: hedges g = -0.607, P=0.059, n=28, studies=3). In particular, Aleyrodidae insects (whitefly) were significantly impacted by *Bacillus* inoculants (hedges g = -1.086, P=0.016, n=8, studies=3; Fig 4b), while Aphididae (aphids) were most negatively affected by non-*Bacillus* inoculants ('Other': hedges g = -0.589, P=0.027, n=17, studies=10; *Pseudomonas*: hedges g = -0.514, P=0.097, n=23, studies=8; Fig 4b). For aphid studies, the other inoculants included *Rhizobium* sp, *Bradyrhizobium* sp, *Sinorhizobium* sp, and an *Acidovorax* sp.

Sucking insects were studied across five different plant families (Fig 4c). We found significant negative effects of rhizobacterial inoculation on sucking insects feeding on plants within Brassicaceae (hedges g = -0.755, P<0.006, n=39, studies=10), driven by effects of *Bacillus* spp (hedges g = -0.931, P=0.006, n=26, studies=4) and other inoculants (hedges g = -0.933, P=0.022, n=3, studies=2), but not *Pseudomonas* (hedges g = -0.355, P=0.277, n=10, studies=4) (Fig 4c). *Bacillus* spp inoculants also significantly impacted sucking insects on Solanaceae plants (hedges g = -0.753, P=0.011, n=20, studies=5; Fig. 4c).

Fig. 4. Effect of rhizobacterial inoculation on sucking herbivores, separated by *Bacillus* spp inoculants (red), *Pseudomonas* spp. inoculants (blue), and other rhizobacteria (grey) across (a) response traits (b) main insect families, and (c) host plant family. Effect size mean and 95% confidence intervals from the linear mixed effect model outputs, individual data shown as bubble plots with sizes relative to the precision of the estimate (1/SE), where larger bubbles show a higher certainty of the data.

Effect of rhizobacteria community inoculations (multi-*Bacillus* and multi-genus inoculants)

Chewing herbivores were slightly more negatively affected when feeding on plants inoculated with multiple rhizobacteria (hedges g = -0.383, P=0.043, n=42, studies=9) than single inoculants (hedges g = -0.273, P=0.051, n=173, studies=34) (community vs single: QM= 5.04, df=2, P=0.081). This difference was driven by the majority of community inoculations using multiple *Bacillus* strains (hedges g = -0.653, P=0.002, n=39, studies=7) rather than multi-genus communities, for which there was insufficient information for a robust analysis (hedges g = -0.511, P=0.330, n=3, studies=2). The effect size of multi-strain *Bacillus* inoculants on chewing herbivores was comparable to single-strain *Bacillus* inoculations (hedges g = -0.598, P=0.002, n=63, studies=10; Fig 5a). Single-strain inoculants from *Pseudomonas* had only minimal overall effect on chewing herbivores (hedges g = -0.296, P=0.064, n=56, studies=16; Fig 5a).

In contrast, sucking insects were more negatively affected by single (hedges g = -0.552, P=0.003, n=113, studies=28) than less-studied community inoculations (hedges g = -0.348, P=0.094, n=16, studies=8). Community inoculations were dominated by those using multi-strain *Bacillus* inoculants that, surprisingly, had no significant effect on insect fitness (hedges g = -0.236, P=0.393, n=10, studies=5) despite single-strain *Bacillus* inoculants having a strong effect (hedges g = -0.624, P=0.007, n=59, studies=10). Of particular note, multi-genus inoculants were able to reduce sucking insect fitness yet examined across very few studies (hedges g = -1.016, P=0.010, n=6, studies=3). Multi-genus inoculants contained various combinations of *Bradyrhizobium* spp, *Azospirillium* spp, *Delftia* spp, *Azotobacter* spp, as well as *Bacillus* spp and *Pseudomonas* spp.

Effect of controlled pot experiments compared to field trials

For chewing herbivores, only the host-choice of insects to *Bacillus*-inoculated plants could be analysed to compare field choice and controlled pot experiments since other variables were only measured in controlled pot experiments. Similarly, for sucking insects only insect population sizes were measured (fecundity) across both environments, with multiple bacterial inoculants. We found that for these variables that the significant effect of rhizobacteria in controlled pot experiments (chewers: hedges g = -1.244, P=0.023, n=9, studies=3; suckers: hedges g = -0.748, P=0.004, n=44, studies=20) was lost in field trials (chewers: hedges g = -0.646, P=0.235, n=9, studies=5; suckers: hedges g = -0.449, P=0.256, n=29, studies=8).

Fig. 5. Effect of rhizobacterial inoculation using single and community inoculations for (a) chewing or (b) sucking herbivores, and comparing controlled pot and field trials for (c) chewing herbivore host-choice, and (d) sucking herbivore fecundity. Effect size mean and 95% confidence intervals from the linear mixed effect model outputs, individual data shown as bubble plots with sizes relative to the precision of the estimate (1/SE), where larger bubbles show a higher certainty of the data.

Discussion

Our meta-analysis showed an overall reduction in herbivore fitness and avoidance behaviour on rhizobacteria inoculated plants. In contrast, there is potential for rhizobacteria inoculation to increase host choice of natural enemies of herbivore insects, increasing top-down predator control. We found significant effects of rhizobacterial inoculation of host plants for chewing herbivores through reduced host choice, feeding (leaf consumption), survival and reproduction (fecundity), while sucking herbivores only were significantly impacted through reduced reproduction (fecundity).

There were strong differences across the rhizobacteria species, with *Bacillus* spp (Blake, Christensen & Kovács 2021) inoculants providing stronger general effects than the commonly studied *Pseudomonas* spp (Pieterse *et al.* 2021). We grouped all other rhizobacteria species since alone they had insufficient data, but they also significantly impacted insects across various traits. These effects were not restricted to specific insect or plant families also highlighting the generalisability of rhizobacterial inoculation for plant protection (Biere & Bennett 2013; Gruden *et al.* 2020).

Single-strain inoculants generally performed better (stronger reduction of insect traits) but this may be due to the choice of multi-strain inoculant communities as opposed to loss of effect. Interestingly, for the sucking herbivores, single-strain *Bacillus* inoculants performed better than multi-strain *Bacillus* inoculants, indicating potential strain incompatibility could affect outcomes(Díaz *et al.* 2023). Lastly, we also observed a loss of insect fitness reduction under field trial conditions compared to controlled pots. This could be due to the restricted number of studied and traits measured in both environments, or conceivably the increased native microbiota of field soils and environmental variability may reduce the ability of inoculants to colonise and persist among plant roots (Gruden *et al.* 2020; Frew *et al.* 2022; He *et al.* 2024).

There is great potential for using rhizobacteria to increase plant resistance to insect pests but these meta-analysis results show that we need to consider the broader interacting environment when developing effective inoculants (Song *et al.* 2020; Shayanthan, Ordoñez & Oresnik 2022). This will involve understanding which rhizosphere are important for specific plants (He *et al.* 2024) but also those that are generalist across multiple host families. With this knowledge, we can develop synthetic microbial communities with broad acting functions across multiple plants, crops and environments.

Acknowledgements

This work was supported by a BBSRC (UKRI) David Phillips Fellowship BB/S010556/1 to SEZ, BBSRC NLD studentship to MP, and German Research Council, Deutsche Forschungsgemeinschaft (DFG; project number 397565003) funding to SZ and OSM.

References

Bakker, P.A., Berendsen, R.L., Doornbos, R.F., Wintermans, P.C. & Pieterse, C.M. (2013) The rhizosphere revisited: root microbiomics. *Frontiers in plant science,* **4,** 165.

Berendsen, R.L., Pieterse, C.M. & Bakker, P.A. (2012) The rhizosphere microbiome and plant health. *Trends in Plant Science,* **17,** 478-486.

Biere, A. & Bennett, A.E. (2013) Three-way interactions between plants, microbes and insects. *Functional Ecology,* **27,** 567-573.

Blake, C., Christensen, M.N. & Kovács, Á.T. (2021) Molecular aspects of plant growth promotion and protection by Bacillus subtilis. *Molecular Plant-Microbe Interactions,* **34,** 15-25.

Blubaugh, C.K., Carpenter-Boggs, L., Reganold, J.P., Schaeffer, R.N. & Snyder, W.E. (2018) Bacteria and Competing Herbivores Weaken Top-Down and Bottom-Up Aphid Suppression. *Front Plant Sci,* **9,** 1239.

da Silva, F.B., Barbosa, J.Z., Tiecher, T., Borin, J.B.M., Treichel, B. & de Sá, E.L.S. (2024) The species-dependent effect of PGPR co-inoculation in legume plants: A global meta-analysis. *Rhizosphere,* 100869.

De Zutter, N., Ameye, M., Bekaert, B., Verwaeren, J., De Gelder, L. & Audenaert, K. (2022) Uncovering new insights and misconceptions on the effectiveness of phosphate solubilizing rhizobacteria in plants: a meta-analysis. *Frontiers in plant science,* **13,** 858804.

Díaz, P.R., Merlo, F., Carrozzi, L., Valverde, C., Creus, C.M. & Maroniche, G.A. (2023) Lettuce growth improvement by Azospirillum argentinense and fluorescent Pseudomonas co-inoculation depends on strain compatibility. *Applied Soil Ecology,* **189,** 104969.

Frew, A., Antunes, P.M., Cameron, D.D., Hartley, S.E., Johnson, S.N., Rillig, M.C. & Bennett, A.E. (2022) Plant herbivore protection by arbuscular mycorrhizas: a role for fungal diversity? *New Phytologist,* **233,** 1022-1031.

Friman, J., Pineda, A., van Loon, J.J. & Dicke, M. (2021) Bidirectional plant-mediated interactions between rhizobacteria and shoot-feeding herbivorous insects: a community ecology perspective. *Ecological Entomology,* **46,** 1-10.

Gruden, K., Lidoy, J., Petek, M., Podpečan, V., Flors, V., Papadopoulou, K.K., Pappas, M.L., Martinez-Medina, A., Bejarano, E., Biere, A. & Pozo, M.J. (2020) Ménage à Trois: Unraveling the Mechanisms Regulating Plant–Microbe–Arthropod Interactions. *Trends in Plant Science,* **25,** 1215-1226.

He, X., Wang, D., Jiang, Y., Li, M., Delgado-Baquerizo, M., McLaughlin, C., Marcon, C., Guo, L., Baer, M. & Moya, Y.A. (2024) Heritable microbiome variation is correlated with source environment in locally adapted maize varieties. *Nature Plants,* 1-20.

Li, J., Wang, J., Liu, H., Macdonald, C.A. & Singh, B.K. (2022) Application of microbial inoculants significantly enhances crop productivity: a meta-analysis of studies from 2010 to 2020. *Journal of Sustainable Agriculture and Environment,* **1,** 216-225.

Pan, J., Peng, F., Xue, X., You, Q., Zhang, W., Wang, T. & Huang, C. (2019) The growth promotion of two salt-tolerant plant groups with PGPR inoculation: a meta-analysis. *Sustainability,* **11,** 378.

Pieterse, C.M., Berendsen, R.L., de Jonge, R., Stringlis, I.A., Van Dijken, A.J., Van Pelt, J.A., Van Wees, S., Yu, K., Zamioudis, C. & Bakker, P.A. (2021) Pseudomonas simiae WCS417: star track of a model beneficial rhizobacterium. *Plant and soil,* **461,** 245-263.

Pieterse, C.M., Poelman, E.H., Van Wees, S.C. & Dicke, M. (2013) Induced plant responses to microbes and insects. *Frontiers in plant science,* **4**.

Pieterse, C.M.J., Zamioudis, C., Berendsen, R.L., Weller, D.M., Van Wees, S.C.M. & Bakker, P. (2014) Induced Systemic Resistance by Beneficial Microbes. *Annual Review of Phytopathology,* **52,** 347-375.

Pineda, A., Zheng, S.-J., van Loon, J.J.A., Pieterse, C.M.J. & Dicke, M. (2010) Helping plants to deal with insects: the role of beneficial soil-borne microbes. *Trends in Plant Science,* **15,** 507-514.

Quintana, D. & Tiebel, J. (2019) An R script and excel file to calculate statistical power for your meta-analysis. https://osf.io/5c7uz/.

R Core Team (2024) R: A Language and Environment for Statistical Computing. R Foundation for Statistical Computing, Vienna, Austria.

RStudio Team (2020) RStudio: Integrated Development for R. RStudio, Inc., Boston, MA.

Rubin, R.L., van Groenigen, K.J. & Hungate, B.A. (2017) Plant growth promoting rhizobacteria are more effective under drought: a meta-analysis. *Plant and soil,* **416,** 309-323.

Schuman, M.C. & Baldwin, I.T. (2016) The layers of plant responses to insect herbivores. *Annual Review of Entomology,* **61,** 373-394.

Selosse, M.-A., Bessis, A. & Pozo, M.J. (2014) Microbial priming of plant and animal immunity: symbionts as developmental signals. *Trends in microbiology,* **22,** 607-613.

Shayanthan, A., Ordoñez, P.A.C. & Oresnik, I.J. (2022) The role of synthetic microbial communities (syncom) in sustainable agriculture. *Frontiers in Agronomy,* **4,** 58.

Song, C., Zhu, F., Carrion, V.J. & Cordovez, V. (2020) Beyond Plant Microbiome Composition: Exploiting Microbial Functions and Plant Traits via Integrated Approaches. *Front Bioeng Biotechnol,* **8,** 896.

Trivedi, P., Leach, J.E., Tringe, S.G., Sa, T. & Singh, B.K. (2020) Plant-microbiome interactions: from community assembly to plant health. *Nat Rev Microbiol,* **18,** 607-621.

Viechtbauer, W. (2010) Conducting meta-analyses in R with the metafor package. *Journal of Statistical Software,* **36,** 1-48.

Wu, J. & Baldwin, I.T. (2010) New insights into plant responses to the attack from insect herbivores. *Annual review of genetics,* **44,** 1-24.

Zeffa, D.M., Fantin, L.H., Koltun, A., de Oliveira, A.L., Nunes, M.P., Canteri, M.G. & Gonçalves, L.S. (2020) Effects of plant growth-promoting rhizobacteria on co-inoculation with Bradyrhizobium in soybean crop: a meta-analysis of studies from 1987 to 2018. *PeerJ,* **8,** e7905.

Zhang, Z., Li, Y., Williams, R.A., Chen, Y., Peng, R., Liu, X., Qi, Y. & Wang, Z. (2023) Responses of soil respiration and its sensitivies to temperature and precipitation: A meta-analysis. *Ecological Informatics*, 102057.

Zhao, X., Tian, P., Sun, Z., Liu, S., Wang, Q. & Zeng, Z. (2022) Rhizosphere effects on soil organic carbon processes in terrestrial ecosystems: A meta-analysis. *Geoderma,* **412,** 115739.