**Iron transport pathways in the human malaria parasite *Plasmodium falciparum***

**revealed by RNA-sequencing**

Juliane Wunderlich[1,2]*, Vadim Kotov[2], Lasse Votborg-Novél[1], Christina Ntalla[1], Maria

Geffken[3], Sven Peine[3], Silvia Portugal[1], Jan Strauss[2#]

[1]Max Planck Institute for Infection Biology (MPIIB), Berlin, Germany
[2]Centre for Structural Systems Biology (CSSB), Hamburg, Germany
[3]University Medical Center Hamburg-Eppendorf (UKE), Institute of Transfusion Medicine,
Hamburg, Germany
10

*Correspondence:

Juliane Wunderlich

juliane.wunderlich@mail.mcgill.ca, Current address: Robert Koch Institute, Department of

Infectious Disease Epidemiology, Berlin, Germany

[#]Current address: German Maritime Centre, Hamburg, Germany

ORCiD:

Juliane Wunderlich          0000-0002-5818-5488

Lasse Votborg-Novél        0000-0003-0381-0669

Sven Peine                  0000-0002-6365-9916

Silvia Portugal             0000-0003-4567-9101

Jan Strauss                 0000-0002-6208-791X

**Keywords: *Plasmodium falciparum*, malaria, drug target, iron deficiency, transporters,**

**nutrient uptake, gene expression, AlphaFold.**

## ABSTRACT

Host iron deficiency is protective against severe malaria as the human malaria parasite *Plasmodium falciparum* depends on bioavailable iron from its host to proliferate. The essential pathways of iron acquisition, storage, export, and detoxification in the parasite differ from those in humans, as orthologs of the mammalian transferrin receptor, ferritin, or ferroportin, and a functional heme oxygenase are absent in *P. falciparum*. Thus, the proteins involved in these processes may be excellent targets for therapeutic development, yet remain largely unknown. Here, we show that parasites cultured in erythrocytes from an iron-deficient donor displayed significantly reduced growth rates compared to those grown in red blood cells from healthy controls. Sequencing of parasite RNA revealed diminished expression of genes involved in overall metabolism, hemoglobin digestion, and metabolite transport under low-iron versus control conditions. Supplementation with hepcidin, a specific ferroportin inhibitor, resulted in increased labile iron levels in erythrocytes, enhanced parasite replication, and transcriptional upregulation of genes responsible for merozoite motility and host cell invasion. Through endogenous GFP tagging of differentially expressed putative transporter genes followed by confocal live-cell imaging, proliferation assays with knockout and knockdown lines, and protein structure predictions, we identified six proteins that are likely required for ferrous iron transport in *P. falciparum*. Of these, we localized *Pf*VIT and *Pf*ZIPCO to cytoplasmic vesicles, *Pf*MRS3 to the mitochondrion, and the novel putative iron transporter *Pf*E140 to the plasma membrane for the first time in *P. falciparum*. *Pf*NRAMP/*Pf*DMT1 and *Pf*CRT were previously reported to efflux $Fe^{2+}$ from the digestive vacuole. Our data support a new model for parasite iron homeostasis, in which *Pf*E140 is involved in iron uptake across the plasma membrane, *Pf*MRS3 ensures non-redundant $Fe^{2+}$ supply to the mitochondrion as the main site of iron utilization, *Pf*VIT transports excess iron into cytoplasmic vesicles, and *Pf*ZIPCO exports $Fe^{2+}$ from these organelles in case of iron scarcity. These results provide new insights into the parasite's response to differential iron availability in its environment and into the mechanisms of iron transport in *P. falciparum* as promising candidate targets for future antimalarial drugs.

## INTRODUCTION

Iron is an essential micronutrient for all living organisms and has been associated with virulence of many pathogens. Iron abundance increases the replication of human immunodeficiency virus (HIV) (1) and *Mycobacterium tuberculosis* (2), and promotes biofilm formation in *Pseudomonas aeruginosa* (3). A "fight for iron" has been described between bacteria and the human host in the gastrointestinal tract (4), where the metal skews the composition of the gut microbiome by facilitating the growth of enteropathogenic *Escherichia coli* and *Salmonella* (5). Similarly, cancer cells require more iron compared to healthy cells (6) and higher ferritin levels in individuals diagnosed with COVID-19 were associated with increased disease severity and lethality (7).

Host iron deficiency is known to be protective against severe malaria (8-11) and iron chelators have cytocidal effects on the human malaria parasite *Plasmodium falciparum* (12). This obligate intracellular parasite depends on bioavailable iron for its proliferation and relies entirely on the host to meet its nutrient requirements (13). Furthermore, *P. falciparum* senses environmental fluctuations (14-16) and modulates its virulence in response (16). While iron is crucial for DNA replication and repair, mitochondrial electron transport, and redox regulation, it becomes toxic when in excess, as it is a source of damaging reactive oxygen species (17). Importantly for therapeutic development, the mechanisms of iron acquisition, storage, detoxification, and export in the parasite are different from those in humans, as orthologs of the mammalian transferrin receptor, ferritin, or ferroportin, and a functional heme oxygenase are absent in *Plasmodium* (18).

While human blood plasma contains between 10 and 30 µM total iron and an erythrocyte carries approximately 20 mM Fe (19), only 3 µM labile iron is present in the cytosol of uninfected red blood cells, and 1.6 µM in *P. falciparum*-infected ones (20). An estimated total iron concentration of 500 mM (21) is reached within the parasite's digestive vacuole (DV), where iron-containing hemoglobin (Hb) is digested and the released heme is detoxified by biocrystallization into hemozoin (22). However, *P. falciparum* cannot access this iron source and is thought to acquire bioavailable $Fe^{2+}$ from the host cell (23). Over-elevated ferrous iron levels likely compromise the integrity of the DV membrane and cytosolic iron also needs to be regulated to prevent oxidative stress (18). Iron detoxification in the parasite can be achieved by translocating the metal ion into dynamic intracellular $Fe^{2+}$ stores, which may include acidocalcisomes – cytoplasmic vesicles that contain high concentrations of phosphate, calcium, iron, and zinc (24). In contrast to *Trypanosoma brucei* (24), no transport proteins have yet been experimentally shown to localize to the acidocalcisome membrane in *P. falciparum* (25, 26). Like the DV, these organelles are thought to be acidified by the plant-

like V-ATPase and their low internal pH may fuel secondary active transport processes (22,

27).

*P. falciparum* encodes approximately 200 transmembrane or membrane-associated

transport proteins (channels, pores, carriers, and pumps), many of which are essential for

parasite growth and lack human homologs (28). For instance, the vacuolar iron transporter

*Pf*VIT (PF3D7_1223700), an ortholog of *Arabidopsis thaliana* VIT1 (expect value (E) = 5 x

$10^{-29}$, 30.5% identity, 87% coverage, as determined by position-specific iterated BLAST

(29)), is a $Fe^{2+}/H^+$ exchanger that plays a role in iron detoxification (30-32). While its

orthologs were localized to the endoplasmic reticulum in *Plasmodium berghei* (30) and to the

vacuolar compartment in *Toxoplasma gondii* (33), the subcellular localization in *P.*

*falciparum* had not been investigated experimentally prior to this study. Similarly, the Zrt-, Irt-

like protein domain-containing protein (ZIPCO) was suggested to import $Fe^{2+}$ and $Zn^{2+}$ into

the cytosol and localized to the parasite plasma membrane (PPM) in *P. berghei* sporozoites

in indirect immunofluorescence assays (34), but *Pf*ZIPCO (PF3D7_1022300) had not been

studied yet. The chloroquine resistance transporter *Pf*CRT (PF3D7_0709000) and the

natural resistance-associated macrophage protein *Pf*NRAMP (PF3D7_0523800, also called

*Pf*DMT1 for divalent metal transporter 1, although this abbreviation is already in use for the

drug/metabolite transporter 1) have been detected at the digestive vacuolar (DV) membrane

(35, 36). Both proteins were proposed to export $Fe^{2+}$ into the cytosol in symport with protons

on the basis of transport assays using *Xenopus* oocytes (37) and proliferation assays with a

conditional knockdown line under different iron conditions (38), respectively.

In *Saccharomyces cerevisiae*, a model organism for eukaryotic iron homeostasis, the

mitochondrial carrier protein MRS3 (mitochondrial RNA-splicing protein 3) was shown to

ensure $Fe^{2+}$ supply to the mitochondrion (39-41) and its ortholog *Tg*MIT (mitochondrial iron

transporter) was detected at the same organelle in *T. gondii*. (33). The mitochondrion of *P.*

*falciparum* is also the focal point for cellular iron metabolism and contains iron-dependent

proteins implicated in the biosynthesis of heme and iron-sulfur clusters, redox reactions, and

electron transport (18). Because of sequence similarity (35.1% identity with the yeast

ortholog, E = 3 x $10^{-14}$, 26% coverage), it was proposed that *Pf*MRS3 (also known as

mitoferrin (*Pf*MFRN), PF3D7_0905200) mediates $Fe^{2+}$ import into the mitochondrion in *P.*

*falciparum* (42). However, no experimental evidence was collected and it is known that not

only the localization but also the structure and function of homologous proteins can vary in

related apicomplexan parasites (42, 43).

Despite the importance of iron for *P. falciparum* virulence, understanding of the molecular
mechanisms of iron sensing, acquisition, utilization, and regulation in the parasite remains
limited. The goal of this exploratory study was to dissect how the parasite responds to
differences in iron availability in its environment and to identify putative iron transporters as
potential new antimalarial drug targets. We investigated growth and gene expression of the
laboratory *P. falciparum* strain 3D7 under control (iron-replete), high-iron and low-iron
conditions, and in the presence of the iron-regulatory peptide hormone hepcidin. In the
human body, hepcidin is produced to reduce the concentration of serum iron when it rises
above a certain threshold. The hormone binds specifically to ferroportin on the surface of
many cell types including erythrocytes (44) and can sterically inhibit the transporter's iron
export activity, thereby increasing intracellular and decreasing serum iron levels (45, 46).
Here, whole-transcriptome sequencing was used to identify putative iron transport proteins
on the basis of differential gene expression patterns between high vs. low-iron conditions.
We then further studied these proteins by analyzing their subcellular localization in live
parasites as well as their predicted 3D structures and by determining growth rates of the
respective knockout or knockdown parasite lines under various conditions.
144
**RESULTS**
**Elevated erythrocyte labile iron levels promote *P. falciparum* proliferation in vitro**
To investigate whether labile iron levels in the erythrocyte correlate with parasite replication
rates, we established different iron conditions in vitro. The first approach was to culture *P.*
*falciparum* 3D7 parasites in 0 Rh+ erythrocytes from voluntary blood donations by
Caucasians aged 18 to 21 at the University Medical Center Hamburg-Eppendorf in
Germany. Therefore, samples from a person with an elevated ferritin level (greater than 200
µg/L (47), in this case 231 µg/L ferritin, 18.2 g/dL Hb, 51.5% hematocrit), an iron-deficient
individual (serum ferritin < 12 ng/ml (48), here: 3 µg/L ferritin, 11.4 g/dL Hb, 36.3%
hematocrit) and a healthy donor (21 µg/L ferritin, 15.0 g/dL Hb, 42.3% hematocrit) were
used. Secondly, infection of red blood cells from other healthy individuals with or without the
addition of 0.7 µM hepcidin to the culture medium was compared. This concentration was
chosen as it had the strongest effect on parasite proliferation in preliminary experiments, and
is expected to increase intracellular $Fe^{2+}$ as it is twice as high as the hepcidin level needed to
reduce $^{55}$Fe export from preloaded mature erythrocytes by 30% within one hour of incubation
(44).
161
Relative labile iron levels in uninfected erythrocytes were estimated by determining the mean
fluorescence intensity (MFI) of the iron-sensitive dye Phen Green SK in 100,000 cells per

replicate using flow cytometry. As binding of ferrous iron to the metal-binding moiety causes

fluorescence quenching of the fluorophore, a reduction in fluorescence intensity indicates

higher labile iron levels (49). In erythrocytes from the iron-deficient donor, the Phen Green

SK MFI was 43% higher relative to control, confirming reduced labile iron levels (Fig. 1A).

The parasite replication rate after one intraerythrocytic developmental cycle (IDC) decreased

by 16% (Fig. 1B), the DNA content of late schizonts by 19% (Fig. 1C) and the number of

merozoites counted per late schizont by 14% (Fig. 1D). In contrast, labile iron levels of

erythrocytes from the donor with higher iron status were only slightly increased (without

statistical support, two-tailed unpaired $t$ tests with Welch's correction for unequal variances

and adjusted with the Holm-Šídák method for multiple comparisons, $P = 0.25$) relative to

blood with normal iron level (healthy control) – as were the parasite proliferation rate, the

DNA content and the merozoite number of mature schizonts (Fig. 1). To further increase

intracellular labile iron levels, we incubated parasites with 0.7 µM hepcidin during one IDC,

resulting in 11% reduced Phen Green SK MFI compared to control (Fig. 1A). Under these

conditions, the parasite growth rate increased by 57% (Fig. 1B), the DNA content per

schizont by 16% (Fig. 1C), and the number of merozoites produced per schizont by 15%

(Fig. 1D).

Taken together, these data show that parasites grown in erythrocytes from an iron-deficient

donor displayed significantly reduced growth rates compared to healthy control. Our in vitro

results also demonstrate that hepcidin treatment of control erythrocytes elevated intracellular

$Fe^{2+}$ concentrations and promoted parasite proliferation.

**RNA-sequencing reveals differential expression of putative iron transporters**

To identify iron-regulated mechanisms and putative iron transporters in *P. falciparum*, we

carried out whole-transcriptome profiling using bulk RNA-sequencing (Fig. 2). *P. falciparum*

3D7 parasites were cultured either using erythrocytes from a donor with higher, control

(healthy) or low iron status (experiment 1); or with red blood cells from another healthy donor

in the presence or absence of 0.7 µM hepcidin (experiment 2). Samples from three biological

replicates per condition were harvested at the ring and trophozoite stage (6 – 9 and 26 – 29

hours post invasion, hpi) during the second IDC under the conditions specified.

To exclude the possibility that differences in mRNA abundance were a consequence of

divergent progression through the IDC under different nutritional conditions, we assessed

the average developmental age of the parasites in each sample on the basis of a statistical

maximum likelihood estimation (MLE) method of transcriptional patterns according to

Lemieux et al. (50). The general transcriptional patterns of parasites were highly similar at

individual time points and consistent across different experimental treatments, corresponding

to those of a 3D7 reference strain (51) at approximately 10 hpi and 35 hpi (Fig. 2A). This

indicates that differences in mRNA abundance of parasites were not caused by divergent

progression through the IDC but by direct effects of the experimental treatments. As the 3D7

strain we used for the experiments had a reduced total IDC length of 44 h instead of 48 h,

possibly because of gene deletions that may have occurred during long-term culturing (52,

53), it progresses through the cycle faster than the 3D7 reference strain (51). This may

explain why the calculated MLEs of parasite age were higher than the actual values of 6 – 9

hpi and 26 – 29 hpi (Fig. 2A).

Using a threshold of 1.5 for the fold change (FC) in gene expression ($\log_2$ FC of 0.585 or -

0.585) yielded twelve significantly upregulated and 175 downregulated genes in ring-stage

parasites under high vs. low-iron conditions ($P < 0.05$, exact test for the negative binomial

distribution with Benjamini-Hochberg correction (54)). As differences in transporter gene

transcription are typically small (55, 56), we examined the 351 upregulated and 770

downregulated genes with a significant expression change and a minimum absolute value of

the $\log_2$ FC of 0.2 for this comparison (Fig. 2B). The full RNA-sequencing datasets are

available in the BioStudies repository (57) under accession number E-MTAB-13411

(https://www.ebi.ac.uk/biostudies/studies/E-MTAB-13411) and differential gene expression

test results for individual genes are shown in Supplementary Tables S1 and S2. The highly

polymorphic *var*, *stevor*, and *rifin* gene families were excluded from downstream analyses

because of their significant sequence diversity between parasites of the same strain during

mitotic growth (58, 59). Functional Gene Ontology (GO), Kyoto Encyclopedia of Genes and

Genomes (KEGG) and Reactome (REAC) term enrichment analyses of differentially

expressed genes (DEGs) were performed using the g:Profiler web server (60).

Under high vs. low-iron conditions at the ring stage (6 – 9 hpi), the GO term for biological

process GO:0055085 "transmembrane transport" was 2.8-fold enriched ($P = 0.007$,

hypergeometric test) among significantly upregulated parasite genes (Fig. 2C). Using the

recently updated *P. falciparum* transporter list (28), all genes with differential expression

levels at the ring stage were then screened for transport proteins and all of the five genes

previously proposed as iron transporters in *Plasmodium* (VIT, ZIPCO, NRAMP/DMT1, CRT,

MRS3/MFRN (18, 43)) were found differentially expressed (Table 1). Other significantly

enriched functional terms at the ring stage under high-iron conditions were GO:0009056

"catabolic process", GO:0020020 "food vacuole", KEGG:01100 "metabolic pathways", and

GO:0005737 "cytoplasm" (Fig. 2C). Among downregulated genes under high vs. low-iron

conditions at the ring stage, KEGG:03440 "homologous recombination", KEGG:03410 "base

excision repair", GO:0007049 "cell cycle", and GO:0015630 "microtubule cytoskeleton" were
enriched (Fig. 2C). At the more metabolically active trophozoite stage (26 – 29 hpi),
processes related to mRNA splicing and protein production were overrepresented in
upregulated genes, as indicated by the 2.9-fold enrichment ($P = 0.00006$) of the
KEGG:03040 pathway "spliceosome" and the 2.5-fold enrichment ($P < 0.05$) of the
GO:0015934 term "large ribosomal subunit" (Fig. 2C).
244
In contrast, hepcidin treatment resulted in reduced metabolism compared to control
conditions, as KEGG:00040 "pentose and glucuronate interconversions", REAC:R-PFA-
71291 "metabolism of amino acids and derivatives", GO:0005737 "cytoplasm", and
GO:0015934 "large ribosomal subunit" were significantly enriched in downregulated genes
during the parasite ring stage at 6 – 9 hpi (Fig. 2E). Among significantly upregulated genes
in the presence vs. absence of hepcidin, the terms GO:0070258 "inner membrane pellicle
complex" ($P < 0.05$), KEGG:03430 "mismatch repair" ($P = 0.04$), and GO:0015630
"microtubule cytoskeleton" ($P = 0.04$) were enriched at the ring stage. GO:0044409 "entry
into host" ($P = 0.00008$) and GO:0052126 "movement in host environment" ($P = 0.0001$)
were overrepresented at the trophozoite stage (Fig. 2E), possibly linked to the observed
increase in parasite proliferation (Fig. 1B).
256
Our RNA-sequencing data also revealed the differential expression of genes involved in
epigenetic, transcriptional, translational, and post-translational regulation. Under high vs.
low-iron conditions, histone deacetylation and chromatin organization processes as well as
GO:1990904 "ribonucleoprotein complex" were significantly enriched in upregulated genes
at the trophozoite stage, and GO:000370 "DNA binding transcription factor activity" in
downregulated genes at the ring stage (Fig. 2C). Furthermore, the known iron-regulatory
protein *Pf*IRP or aconitate hydratase (61, 62) was upregulated during the ring stage under
high vs. low-iron conditions ($\log_2 FC = 0.49$, $P = 0.00003$) and downregulated in the
presence of hepcidin ($\log_2 FC = -0.27$, $P = 0.01$) as compared to control. Many protein
kinases involved in post-translational modifications and endocytosis were also upregulated
at 26 – 29 hpi at high vs. low iron levels, as indicated by the enriched terms GO:0043170
"macromolecule metabolic process" and KEGG:04070 "phosphatidylinositol signaling
system" (Fig. 2C).
270
**Localization of putative iron transporters in *P. falciparum***
On the basis of transcriptomic profiles and the *P. falciparum* transporter list (28), six proteins
with a potential role in iron transport were identified (Table 1 and 2). The subcellular
localization of the four proteins that had not yet been localized in *P. falciparum* (*Pf*MRS3,

*Pf*VIT, *Pf*ZIPCO, and *Pf*E140) was then examined by endogenous tagging with GFP and

confocal imaging of live parasites under physiological control conditions. At least two cell

lines were generated per transporter candidate with consistent results and representative

example images are shown in Fig. 3. Diagnostic PCRs confirmed the fusion of *gfp* to the

respective gene of interest and the absence of parental DNA at the original locus

(Supplementary Fig. S2). Only the *Pf*MRS3 reporter cell line still contained wild-type DNA of

the parental parasites even after prolonged WR99210/neomycin selection and limiting

dilution cloning (Supplementary Fig. S2), indicating the importance of this mitochondrial

transporter for asexual parasite growth during the blood stage.

The GFP-tagged mitochondrial carrier protein *Pf*MRS3 exclusively localized to the

mitochondrion, as determined by colocalization with MitoTracker Red (Fig. 3A,

Supplementary Video S1). *Pf*VIT-GFP displayed a punctate fluorescence pattern within the

cytoplasm (Fig. 3B, Supplementary Video S2), resembling that of *Pf*ZIPCO-GFP (ZIP

domain-containing protein, PF3D7_1022300, Fig. 3C, Supplementary Video S3). These

structures did not colocalize with ER Tracker Red in live cells (Fig. 3B, Supplementary Video

S2). For both *Pf*VIT-GFP and *Pf*ZIPCO-GFP, the number of cytoplasmic foci increased as

the parasites matured from the ring to the late schizont stage (Fig. 3B and C). To test

whether these could be acidocalcisomes, we employed LysoTracker Deep Red, commonly

used to visualize small acidic organelles in *T. brucei* (24). However, the fluorescent dye only

stained the DV in *P. falciparum* (Fig. 3C, Supplementary Video S3) and no acidocalcisome-

specific marker is currently available for this parasite.

GFP-tagged *Pf*E140 (PF3D7_0104100), also known as conserved *Plasmodium* membrane

protein or CPMP (63), localized to the parasite plasma membrane, as evidenced by the ring-

like fluorescence pattern around newly formed merozoites (Fig. 3D, Supplementary Video

S4). The fluorescence intensity was very low at the ring and early trophozoite stage

compared to schizonts. Because of amino acid sequence similarity (E = $9 \times 10^{-5}$, 22.5%

identity, 66% coverage) to the essential apicoplast transporter *Pf*DER1-2 (29, 64), we also

investigated the potential colocalization with the apicoplast marker *Pf*ACP (acyl carrier

protein), which could not be detected (Fig. 3D, Supplementary Video S4).

**The role of *Pf*VIT, *Pf*ZIPCO and *Pf*E140 for asexual parasite growth**

To study the function of the putative transport proteins identified, we used targeted gene

disruption (TGD) by selection-linked integration (SLI) to generate the corresponding

knockout parasite lines for the putative iron transporters that are non-essential during *P.*

*falciparum* blood stage: *Pf*VIT and *Pf*ZIPCO (Fig. 4A, Supplementary Fig. S2). As GFP was

cloned in frame with the truncated version of the respective transporter (the N-terminal 143

amino acids (aa) of 274-aa *Pf*VIT or 117 of the 325 aa of *Pf*ZIPCO), the subcellular

localization of the resulting GFP fusion protein was also assessed. *Pf*VIT(1-143)-GFP

localized to cytoplasmic structures and *Pf*ZIPCO(1-117)-GFP to the DV and cytoplasmic

vesicles (Fig. 4A). Proliferation assays were then performed to determine the importance of

the respective transporter for parasite growth. While the *Pf*VIT knockout had no effect on

parasite growth under standard conditions, addition of hepcidin reduced the growth rate of

the ΔVIT line by 30% (Fig. 4B). Of note, hepcidin generally had a smaller effect after two

cycles of incubation (Fig. 4B) than after one cycle compared to the first IDC (Fig. 1B).

Unexpectedly, knocking out *Pf*ZIPCO led to a growth rate increase by 42% after two IDCs

relative to wild-type 3D7 parasites (Fig. 4B).

*Pf*E140 is predicted to be essential (65) and the only putative iron transporter identified that

localized to the PPM (Fig. 3), thus potentially important for iron uptake in *P. falciparum*. For

an inducible knockdown, a *glmS* ribozyme sequence (66) was introduced upstream of the 3'

untranslated region in the pSLI plasmid, allowing for conditional mRNA degradation by

adding 2.5 mM glucosamine (GlcN) to the culture medium (Fig. 4C, Supplementary Fig. S2).

The knockdown led to a 61% decrease in total parasite fluorescence intensity after 36 hours

of GlcN treatment (Fig. 4D) without affecting parasite size compared to untreated control

(Fig. 4E). Addition of GlcN also caused a 38% growth rate reduction of the *Pf*E140-GFP-

*glmS* line, which was rescued by hepcidin treatment to a proliferation level that was not

significant different from that under standard culture conditions ($P$ = 0.25, Fig. 4F). The

generation of a *Pf*MRS3-knockdown line was not successful after four independent attempts,

supporting the essentiality of the gene for asexual growth (65).

**Characterization and functional implications of predicted protein structures**

We next took advantage of the recent progress in protein structure prediction and generated

models of the putative iron transport proteins identified (Table 2) using AlphaFold2 (67, 68)

and AlphaFold2-multimer (69). The transmembrane regions of the proteins typically

exhibited the highest confidence score, while some other protein portions appeared

unstructured (Fig. 5A). Regions that are likely located within a membrane were validated by

inspecting the molecular lipophilicity potential of the protein surfaces (Fig. 5B). A clear

hydrophobic belt was observed for all proteins and their orientation in the membrane was

determined on the basis of orthologous proteins. As transport cavities with negatively

charged residues are a hallmark of heavy metal ion transporters, we analyzed the

distribution of charge on the surface of the proteins and looked for negatively charged

regions to assess the capacity to bind cations like $Fe^{2+}$ (Fig. 5C). To gain further insights into

the functions of the proteins identified, we also compared the predicted structures with those

of well-characterized homologs from *S. cerevisiae*, *Eucalyptus grandis, Bordetella*

*bronchiseptica* and *Staphylococcus capitis* (Table 2, Supplementary Fig. S3 and S4).

The outer surface of *Pf*MRS3 (transport classification (TC): 2.A.29, mitochondrial carrier

family) is positively charged (Fig. 5C) and there is a clear negatively charged patch in the

putative binding pocket facing the mitochondrial intermembrane space. We compared the

predicted *Pf*MRS3 structure with that of *S. cerevisiae* MRS3, which is known to import

ferrous iron into the mitochondrial matrix across the inner membrane (39-41). The predicted

structures of *Pf*MRS3 and *S. cerevisiae* MRS3 were superimposed with an average root

mean square deviation of Cα atoms (Cα RMSD) of the 205 matched residues of 2.3 Å

(Supplementary Fig. S3A and S4A). The conserved histidine residues His[48] and His[105] that

were required for $Fe^{2+}$ transport by *S. cerevisiae* MRS3 in reconstituted liposomes (39) are

also present in *Pf*MRS3 and the three functionally relevant histidine residues identified in

yeast are in a similar functional context in both structures (Supplementary Fig. S3A and

S4A). This suggests that MRS3 may elicit similar molecular functions in *S. cerevisiae* and *P.*

*falciparum.*

*Pf*VIT is highly similar to VIT1 from *E. grandis* (E = 7 x 10[-27], 30.3% identity, 84% coverage),

for which an experimental structure is available (Protein Data Bank (PDB) identifier: 6IU4).

The plant protein crystallized as a homodimer (70), and the same oligomeric state was

suggested for the vacuolar iron transporter family (TC: 2.A.89) protein in *P. falciparum* (31).

A *Pf*VIT monomer also has five transmembrane domains and comprises a negatively

charged region facing the cytosol that may enable cation transport (Fig. 5C). In agreement

with this, one $Fe^{2+}$ ion and two $Zn^{2+}$ ions were bound by a strongly charged region on the

cytosolic side of the *E. grandis* VIT1 monomer (70) and a highly similar putative binding

pocket is present in the parasite protein (Supplementary Fig. S4B). In the structural

alignment, 219 of the 227 residues of the experimental *Eg*VIT1[23-249] structure are within 5 Å

of the predicted structure of *Pf*VIT with an average Cα RMSD of 1.9 Å and the key residues

in the metal-binding domain (Glu[102], Glu[105], Glu[113], Glu[116], using *Eg*VIT1[23-249] numbering) are

placed in a similar molecular context in the predicted structure of *Pf*VIT (Supplementary Fig.

S3B and S4B). The residues in the transmembrane domain that are in the vicinity of the $Co^{2+}$

ion in the *Eg*VIT1[23-249] structure (Met[80] and Asp[43]) are also conserved (Supplementary Fig.

S3B), which is in line with a similar function of *Pf*VIT and *Eg*VIT1.

*Pf*ZIPCO contains seven transmembrane domains and was modeled as a homodimer (Fig.

5A), as it is part of the zinc ($Zn^{2+}$)-iron ($Fe^{2+}$) permease (ZIP) family (TC: 2.A.5), whose

members usually function as homo- or heterodimers (71). The negatively charged patch in

each binding pocket facing the vesicle lumen (Fig. 5C) may be involved in cation transport to

the cytosolic side. In an overlay of the *Pf*ZIPCO model with the cryo-EM structure (PDB:

8GHT) of a ZIP transporter from *B. bronchiseptica* in the presence of either $Zn^{2+}$ or $Cd^{2+}$ ions

(72), the average Cα RMSD of 140 sequence-aligned residues was 2.0 Å (Supplementary

Fig. S3C and S4C). Several key residues of the metal binding site M1 of *Bb*ZIP (Met[99],

His[177], Glu[181], Glu[211]) were also found in *Pf*ZIPCO, whereas others (Asn[178], Gln[207], Asp[208],

Glu[240]) were different (Supplementary Fig. S3C), possibly resulting in divergent substrate

specificity.

*Pf*NRAMP (TC: 2.A.55, metal ion ($Mn^{2+}$-iron) transporter family) is a homolog of the human

endosomal $Fe^{2+}$ transporter 2/DMT1 (73) and contains twelve transmembrane domains (Fig.

5A). Like *Pf*CRT (TC 2.A.7, drug/metabolite exporter family), for which a recent cryo-EM

structure (PDB: 6UKJ) is available (74), the predicted structure possesses a negatively

charged region within its binding pocket facing the digestive vacuolar lumen (Fig. 5C). This is

consistent with binding of cations such as $Fe^{2+}$. The *Pf*NRAMP model was superimposed on

the solved crystal structure of *S. capitis* NRAMP/DMT (PDB: 5M95, E = 1 x $10^{-32}$, 26.3%

identity, 60% coverage), which was shown to bind $Mn^{2+}$, $Fe^{2+}$, $Co^{2+}$, $Ni^{2+}$, $Cd^{2+}$ and $Pb^{2+}$ (75).

In the overlay, the average Cα RMSD of the 349 matched residues was 1.6 Å and the

negatively charged cavity of *Pf*NRAMP was in close proximity to the $Mn^{2+}$ ion bound to *S.*

*capitis* NRAMP (Supplementary Fig. S3D and Supplementary Fig. S4D). Two of the four key

residues required for ion coordination in the binding pocket of the bacterial protein (Asn[52]

and Asp[49]) are present in *Pf*NRAMP, whereas the two other residues (Met[226] and Ala[223]) are

changed to serine (75). While potential effects of these differences on substrate specificity

and / or transport activity remain to be elucidated, *Pf*NRAMP is likely to perform cation

transport from the DV into the cytosol.

*Pf*E140 is predicted to be anchored in the parasite plasma membrane by a bundle of five

transmembrane domains (76) and forms a coiled coil with a hydrophilic region that displays

negatively charged patches exposed to the extracellular side (Fig. 5A and B). No human

orthologs could be identified for this highly conserved *Plasmodium* protein (29). As there is

no obvious channel or cavity in the transmembrane region of the *Pf*E140 monomer (Fig. 5B

and C), the helical bundles may form a dimer to enable ion transport. However, we were not

able to obtain a *Pf*E140 dimer model with AlphaFold2-multimer because of its sequence

length. To predict functional residues on the basis of the amino acid sequence and the

AlphaFold2 structure of *Pf*E140, we used DeepFRI graph convolutional network (77), which

has significant denoising capability and can reliably assign GO terms to residues in the

protein. In particular, the terms GO:0022857 "transmembrane transporter activity" (DeepFRI

gradCAM score 0.94), GO:0015075 "monoatomic ion transmembrane transporter activity"

(score 0.78), and GO:0046873 "metal ion transmembrane transporter activity" (score 0.67)

were assigned to a putative transmembrane region of *Pf*E140 with high confidence

(Supplementary Fig. S5). We thus speculate that the protein is a transporter of metal ions.

**DISCUSSION**

Here, we studied the role of iron in growth and transcription of *P. falciparum* by using blood

from individuals of different iron status and by adding hepcidin as an iron-regulatory

hormone and ferroportin inhibitor. Overall, our data demonstrate the importance of $Fe^{2+}$ in

parasite replication and development and highlight areas for further study. We showed that

in vitro growth rates of *P. falciparum* 3D7 and the number of merozoites formed per schizont

were reduced within erythrocytes that contain lower concentrations of labile iron, while

culturing in blood from an individual with higher iron status did not lead to a significant

increase in labile iron levels within erythrocytes or in parasite growth relative to control (Fig.

1). Consistent with this, reduced propagation of *P. falciparum* 3D7, Dd2, and FCR3-FMG

was reported when erythrocyte samples from iron-deficient individuals used for parasite

culture (10, 78). This effect was eliminated after these donors were iron-supplemented,

whereas supplementation of healthy (iron-replete) donors did not significantly promote

parasite growth (10). The strong increase in parasite replication in the presence of hepcidin

relative to control conditions (Fig. 1B) may have been a result of enhanced invasion

efficiency in addition to the increased number of merozoites formed (Fig. 1D). Earlier studies

also found that higher hepcidin levels in blood samples were associated with elevated *P.*

*falciparum* growth rates in vitro (78) and severe malaria in vivo (79), however, the effect of

experimental hepcidin addition on parasite growth had not been assessed previously.

To identify putative iron transporters and iron-regulated processes, we carried out RNA-

sequencing analyses of *P. falciparum* during the ring (6 – 9 hpi) and trophozoite (26 – 29

hpi) stages cultured under the different iron conditions described above. A higher number of

biological processes and pathways were significantly enriched among DEGs when

erythrocytes from donors with different iron status were used for parasite culture (Fig. 2C)

compared to red blood cells from the same healthy donor in the presence vs. absence of

hepcidin (total of 28 vs. 13 functional terms, Fig. 2E). This may reflect greater differences in

the culture conditions; for instance, blood from the donor with high serum ferritin and Hb

levels may have also contained more glucose or copper (80, 81), potentially explaining the

more diverse physiological response of the parasite. Including erythrocyte samples from

more individuals in the growth experiments and RNA-sequencing analysis would have

provided further insights, however, the provision of sufficient blood from iron-deficient donors

is limited by ethical constraints.

The availability of additional nutrients likely resulted in increased endocytosis and digestion

of host cell contents in the DV of the parasite, leading to enhanced metabolism, mRNA

splicing, and protein production. Interestingly, the terms KEGG:01100 "metabolic pathways"

and GO:0005737 "cytoplasm" were also found to be enriched in upregulated parasite genes

in children with high vs. low parasitemia (82, 83). RNA binding and mRNA splicing

processes were previously reported to be overrepresented in upregulated genes in severe

malaria linked to high parasite density (82-84). Hence, an increase in overall parasite fitness

under high vs. low-iron conditions may explain the increase in parasite multiplication (Fig.

1B) and could be associated with higher parasitemia and disease severity. Consistent with

the observed upregulation of transmembrane transporters at the parasite ring stage (6 – 9

hpi) under high vs. low-iron conditions, Mancio-Silva et al. found that the functional term "ion

transporter activity" was enriched in *P. berghei* genes that were downregulated under caloric

restriction at 6 and 10 hpi (16). Thus, transmembrane transporter genes may need to be

transcribed at the beginning of the IDC to ensure that the appropriate level of transport

proteins is available for nutrient acquisition and metabolite efflux during the subsequent

metabolically active trophozoite and schizont stages.

Hepcidin plays a central role in mammalian iron homeostasis and reduces serum iron

concentrations (85). It is also known that hepcidin levels are elevated in *P. falciparum*-

infected individuals, especially those with high parasitemia (79, 86), and that malaria causes

iron deficiency (79). The transcription profile of parasites treated with 0.7 µM hepcidin

showed similarities to those cultured in erythrocytes from the iron-deficient donor compared

to standard conditions in terms of downregulated catabolic and translation processes as well

as transport protein regulation (Fig. 2, Table 1). This may be related to the fact that an

aberrant hepcidin increase causes systemic iron deficiency as a result of restricted iron

availability (87). The upregulation of genes involved in merozoite motility (*Pf*MTIP, *Pf*GAP45,

and various inner membrane complex proteins) and host cell entry (such as *Pf*AMA1,

*Pf*MSP3, *Pf*MSP7, and *Pf*EBA181) when hepcidin was present (Fig. 2) may suggest an

improved ability of the released merozoites to invade erythrocytes. Thus, the addition of the

peptide hormone to the culture media could be a signal for the parasite to reduce metabolic

processes and to increase its invasion efficiency.

In addition to roles in parasite proliferation and development, different levels of labile iron

may induce regulatory processes at various levels. Under high-iron conditions, the observed

upregulation of histone deacetylation (Fig. 2C) may lead to the condensation and thus

deactivation of certain chromatin regions (88). Similarly, iron-mediated regulation of mRNA

translation by iron-regulatory proteins has been described in yeast, trypanosomes and

mammals (89-91). The binding sites and target genes of the differentially expressed

transcription factors and of *Pf*IRP remain to be identified in *P. falciparum*. Moreover, protein

phosphorylation may play a role in iron-dependent regulatory mechanisms. As a

serine/threonine kinase (KIN) serves as a nutrient sensor in *P. berghei*, driving a fast

response that leads to increased parasite multiplication and virulence (16), a similar kinase

may sense iron and lead to increased replication in *P. falciparum*.

On the basis of our RNA-sequencing results (Fig. 2 and Table 1) and the *P. falciparum*

transporter list (28), we identified six proteins that are likely involved in iron transport in the

parasite (Table 2 and Figure 6) and analyzed their subcellular localization (Fig. 3), their

importance for growth (Fig. 4), and their predicted structures (Fig. 5). *Pf*MRS3 transcription

was upregulated at the ring stage under high vs. low-iron conditions ($\log_2$ FC = 0.33, $P$ =

0.002, Fig. 2B), and fluorescence of the GFP-tagged protein was exclusively detected at the

mitochondrion (Fig. 3A). As a disruption of the gene was reported to fail (65), we were not

able to generate a knockdown line after four independent attempts, and parental DNA of the

original gene locus was still present in the GFP reporter line (Supplementary Fig. S2),

*Pf*MRS3 is likely essential for asexual growth like PBANKA_041620 (E = 1 x 10$^{-69}$, 71.4%

identity, 25% coverage) in *P. berghei* (92). The orthologous mitochondrial iron transporter

(*Tg*MIT, TGME49_277090, E = 7 x 10$^{-19}$, 26.0% identity, 28% coverage) also localized to the

mitochondrion in *T. gondii* and was upregulated at the protein level upon iron overload in

consequence of a *Tg*VIT knock out in the related apicomplexan parasite (33). Our structural

analyses (Fig. 5, Supplementary Fig. S3A and S4A) further support that *Pf*MRS3 imports

ferrous iron into the mitochondrion, the main iron user of the cell, thereby reducing the

cytosolic $Fe^{2+}$ concentration (Fig. 6) as a means of detoxification, which has been reported

for yeast (93). The protein's substrate specificity as well as iron binding and transport activity

remain to be confirmed experimentally.

Complementation assays in *S. cerevisiae* indicated a role for *Pf*VIT in iron detoxification (30,

31) and we observed that the expression of the gene was upregulated under high vs. low-

iron conditions in *P. falciparum* ($\log_2$ FC = 0.29, $P$ = 0.02, Fig. 2B). The fluorescence pattern

of *Pf*VIT-GFP in live cells (Fig. 3B) was consistent with cytoplasmic vesicles that may be

acidocalcisomes, as described for *T. brucei* VIT1 (24). An increase in the number of

fluorescent punctate structures during parasite development (Fig. 3B) was also observed for

VIT in *T. gondii* (33). *Pf*VIT shares 47.0% identity with *Tg*VIT (E = 8 x 10$^{-84}$, 95% coverage)

and 36.9% identity with *Tb*VIT1 (E = 9 x $10^{-39}$, 98% coverage). In contrast, *P. berghei* VIT

(PBANKA_143860, E = 3 x $10^{-160}$, 79.3% identity, 98% coverage) was shown to localize to

the ER in indirect immunofluorescence assays (30). This may be explained by differences

between species or variation in methodology such as fixation, permeabilization, and

immunolabeling techniques as opposed to live-cell imaging (94-96).

Transport assays using inverted vesicles that were prepared using recombinant *Pf*VIT

expressed in *E. coli* demonstrated that the protein is a $Fe^{2+}/H^+$ antiporter (32). The

translocation of $Fe^{2+}$ in exchange for $H^+$ is likely fueled by the pH gradient across the

membrane of the acidic vesicles and the high similarity of the putative $Fe^{2+}$-binding pocket at

the cytosolic side of the predicted *Pf*VIT structure with that of experimentally characterized

*Eg*VIT1 (Fig. 5, Supplementary Fig. S4B) provide further evidence for our hypothesis. While

not essential during asexual blood stages (65), a knockout of VIT resulted in reduced liver

stage development in *P. berghei* (30) and increased sensitivity to high iron levels in both *P.*

*berghei* (30) and *T. gondii* (33). Similarly, growth of the ΔVIT *P. falciparum* line was not

affected under standard conditions, whereas the addition of hepcidin – which increases

intracellular labile iron levels (Fig. 1A) – compromised parasite proliferation in our study (Fig.

4B). Thus, we hypothesize that the transporter sequesters $Fe^{2+}$ into cytoplasmic vesicles,

which is important for iron detoxification under high-iron conditions. While ΔVIT *P. falciparum*

is more sensitive to elevated intracellular $Fe^{2+}$ concentrations (Fig. 4B) as a consequence of

impaired removal of excess iron from the cytosol, *Pf*MRS3 may compensate for a loss of

*Pf*VIT under standard conditions by transporting ferrous iron from the cytosol into the

mitochondrion (Fig. 6).

In contrast to the PPM staining of *P. berghei* sporozoites in immunofluorescence assays,

*Pf*ZIPCO-GFP expression resulted in a punctate fluorescence pattern in the cytoplasm of

live blood-stage *P. falciparum* (Fig. 3C), similar to that of *Pf*VIT-GFP (Fig. 3B). Whereas the

*Pf*ZIPCO knockout caused a growth increase under standard conditions (Fig. 4B), ΔZIPCO

*P. berghei* parasites displayed normal blood-stage development but impaired sporozoite

infectivity as well as reduced replication at the liver stage in mice (34). Interestingly, the

ortholog TGME49_225530 is also dispensable in *T. gondii* tachyzoites with a phenotype

score of –2.94 (values below -1.5 are considered non-essential (97)). Hence, $Fe^{2+}$ efflux from

cytoplasmic vesicles (potentially acidocalcisomes) into the cytosol via *Pf*ZIPCO may be

dispensable in *P. falciparum* under iron-replete conditions during the blood stage because of

the redundancy with iron import mechanisms into the parasite, and the production of the

protein may result in a fitness cost. In contrast, liver-stage parasites in low-iron environments

may rely on the transporter's activity when the demand for iron is high during schizogony. As

the transcription of $Pf$ZIPCO was upregulated at low vs. control iron levels ($log_2$ FC = 0.55, $P$

= 0.04, Table 1) and in response to hepcidin treatment ($log_2$ FC = 0.63, $P$ = 0.006, Fig. 2D,

Table 1), the transport protein may release $Fe^{2+}$ and $Zn^{2+}$ ions from intracellular stores, in

this case cytoplasmic vesicles (Fig. 6), in case of scarcity, thereby increasing cytosolic ion

levels like other ZIP transporters (43). While our analyses of the predicted structure and its

alignment with $Bb$ZIP indicate that $Pf$ZIPCO likely has the capacity to bind and transport

cations like $Fe^{2+}$ or $Zn^{2+}$ (Fig. 5, Supplementary Fig. S3C and S4C), its substrate specificity

can only be conclusively established by characterizing the purified protein. Liposomal assays

with the putative zinc transporter $Pf$ZIP1 (PF3D7_0609100, 24.5% identity with $Pf$ZIPCO, E

= 1 x $10^{-19}$, 78% coverage), which localized to the plasma membrane in schizonts,

demonstrated that this ZIP transporter preferentially binds $Zn^{2+}$ over $Fe^{2+}$ (98). Interestingly,

the preference was abolished if the histidine-rich loop at the C-terminus of $Pf$ZIP1, which is

not present in $Pf$ZIPCO, was truncated. As mRNA levels of $Pf$ZIP1 were enhanced at low

cytosolic $Zn^{2+}$ levels (98) but not differentially regulated under various iron conditions

(Supplementary Tables S1 and S2), it may play a role in zinc rather than iron homeostasis

under physiological conditions.

As the highest intracellular iron concentration in $P. falciparum$ is reached within the DV (21,

99), the free form of the metal may need to be exported from this compartment under high-

iron conditions to prevent damage to the DV membrane (Fig. 6). This function may be

fulfilled by $Pf$CRT (37) and / or $Pf$NRAMP (73), which were both upregulated under high vs.

low-iron conditions in our RNA-sequencing analysis ($log_2$ FC = 0.26, $P$ = 0.007 and $log_2$ FC =

0.28, $P$ = 0.003, respectively, Fig. 2B, Table 1) and are essential in asexual parasites (35,

36, 65). The predicted structure of $Pf$NRAMP (Fig. 5) reflects the state that is open towards

the cytosol as in the crystal structure of NRAMP from $Deinococcus$ $radiodurans$ (100). While

a negatively charged cavity inside the protein is clearly visible in the $Pf$NRAMP model, the

proposed outward-facing permeation pathway for metal ions is likely occluded in this

conformation (Fig. 5C). It is conceivable that $Fe^{2+}$ ions permeate through this pathway from

the DV lumen and bind to the charged cavity like the $Mn^{2+}$ ion to $S. capitis$ NRAMP/DMT

(Supplementary Fig. S4D). $Pf$NRAMP might function similarly to its ortholog in $D.$

$radiodurans$, which was shown to mediate pH-dependent transport of $Fe^{2+}$ and $Mn^{2+}$ in

symport with $H^+$ using uptake assays in $E. coli$, HEK293T cells, and proteoliposomes (100,

101).

Expression of the surface protein $Pf$E140 was upregulated when iron levels were low

compared to standard conditions ($log_2$ FC = 0.65, $P$ = 0.0006, Table 1) and the GFP fusion

protein localized to the PPM only, as evidenced by the fluorescent edges of free merozoites

(Fig. 3D). This observation is consistent with the fact that the extracellular portions of this

protein are highly polymorphic because of their exposure to the immune system at the

sporozoite stage (76). Interestingly, vaccines targeting *Py*E140 in *Plasmodium yoelii* were

reported to induce up to 100% sterile protection mediated by antibodies in mice (102). The

reduced parasite replication rate upon its conditional knockdown demonstrates the

importance of *Pf*E140 for parasite growth and the rescue of the *Pf*E140 knockdown by

hepcidin treatment support a role of this putative transporter in iron uptake (Fig. 4F). Its

predicted essential nature (65), in addition to the absence of orthologs in humans, make it an

excellent drug target candidate. While our *P. falciparum* gene expression data (Fig. 2B and

D) point towards a role of *Pf*E140 in iron homeostasis, its precise function is still unclear and

it remains to be clarified whether the large coiled-coil domain exposed to the extracellular

space (Fig. 5) can mediate dimerization upon substrate binding. Given our experimental

results and the functional annotations (Fig. 5C, Supplementary Fig. S5), we hypothesize that

*Pf*E140 is a plasma membrane transporter for inorganic cations such as metal ions.

In conclusion, this is the first study to investigate *P. falciparum* transcriptomics under

different iron conditions and to determine the subcellular localization of the known and

putative iron transport proteins *Pf*MRS3, *Pf*VIT, *Pf*ZIPCO and *Pf*E140 as well as the growth

effects of a *Pf*VIT or *Pf*ZIPCO knockout and an inducible *Pf*E140 knockdown. Our results

reveal how the human malaria parasite reacts to alterations in host iron status and provide

new insights into the mechanisms of iron transport in *P. falciparum* in addition to offering

avenues for the development of novel therapeutic strategies against malaria. We propose a

model for the regulation of iron homeostasis in the *P. falciparum*-infected erythrocyte with a

series of six organelle-specific iron transport proteins in the parasite (Fig. 6): One route of

iron uptake into the parasite is through the release of $Fe^{2+}$ upon hemoglobin digestion in the

DV and the efflux of the ion into the cytosol mediated by *Pf*NRAMP (38) and / or *Pf*CRT (37).

Ferrous iron likely also enters the parasite cytosol across the PPM via *Pf*E140 and this

pathway may be particularly important during schizogony, when the putative transporter

gene is abundantly transcribed and new merozoites without a DV are formed (51). Once

inside the cytosol, iron concentrations need to be tightly regulated to avoid toxicity, which

could be achieved by $Fe^{2+}$ import into the mitochondrion as the main site of iron utilization

via *Pf*MRS3 and through transport into (*Pf*VIT) and out of (*Pf*ZIPCO) cytoplasmic vesicles

functioning as labile iron pools. To confirm the hypotheses of our exploratory study, transport

assays with purified proteins like those performed with recombinant *Pf*VIT (32) are required

for the formal demonstration of substrate specificities and activities of the other transporters

in addition to further functional characterization of the proteins during the mosquito, liver and

asexual blood stages of the parasite. As no ortholog of the essential *Pf*E140 and only a

distant homolog of the non-redundant mitochondrial iron importer *Pf*MRS3 (42) are present

in humans, these provide candidate targets for urgently needed new antimalarial drugs.

Furthermore, dissecting how *P. falciparum* senses changes in micronutrient availability in its

environment and how it modulates its virulence accordingly is an area of considerable

interest for future investigation, as iron is an essential regulatory signal for virulence factors

in many pathogens.

**MATERIALS AND METHODS**

***P. falciparum* culture and proliferation assays**

The *P. falciparum* strain 3D7 was cultured according to modified standard procedures (103)

at 5% hematocrit using human 0 Rh+ erythrocytes from the University Medical Center

Hamburg-Eppendorf (UKE), Germany, at 1% $O_2$, 5% $CO_2$ and 94% $N_2$. RPMI 1640 medium

was supplemented with 0.5% (w/v) AlbuMAX II, 20 µg/mL gentamicin and 100 µM

hypoxanthine (Thermo Fisher Scientific). Mature schizonts were obtained by treating

schizonts at 40 hpi with 1 mM compound 2 (4-[7-[(dimethylamino)methyl]-2-(4-

fluorphenyl)imidazo[1,2-α]pyridine-3-yl]pyrimidin-2-amine, LifeArc) for 8 h. To count the

number of merozoites per mature schizont, Giemsa-stained blood smears were analyzed by

light microscopy. Only single-infected cells with one digestive vacuole were taken into

account.

To assess parasite proliferation over six days, a previously described assay on the basis of

flow cytometry was employed (104). Parasites were synchronized to a 3-h age window by

isolating late schizonts from a 60% Percoll (GE Healthcare) gradient and culturing these for

3 h with fresh erythrocytes (105), followed by controlled elimination of advanced parasite

stages using 5% (w/v) D-sorbitol (Carl Roth) for 10 min at 37°C (106). The growth assay was

started at 0.1% parasitemia using the resulting ring-stage parasites at 0 – 3 hpi. The

parasitemia was determined at the trophozoite stage every two days by flow cytometry and

culture media with the respective supplements were exchanged daily.

**Flow cytometry**

To determine parasitemia, 20 µL of resuspended parasite culture was added to 80 µL culture

medium and stained with 5 µg/mL SYBR Green I (Thermo Fisher Scientific) and 4.5 µg/mL

dihydroethidium (DHE, Sigma-Aldrich) in the dark for 20 min at room temperature. Stained

cells were washed with PBS three times and analyzed with an ACEA NovoCyte flow

cytometer and NovoExpress Software (version 1.6.1, Agilent). Forward and side scatter

gating was used to identify erythrocytes and SYBR Green I fluorescence intensity to

determine the number of parasitized cells per 100,000 events recorded for each replicate.

For Phen Green SK measurements, uninfected erythrocytes were washed with PBS and

incubated with 10 µM Phen Green SK in PBS at 37°C for 60 min. DHE at 4.5 µg/mL was

added during the last 20 min of incubation. After three washes with PBS, the cells were

analyzed as described above.

**Cloning of DNA constructs**

For generating the GFP reporter lines, a homologous region of approximately 800 bp at the

3' end of the respective gene was amplified without the stop codon from 3D7 gDNA using

Phusion high fidelity DNA polymerase (New England Biolabs). A homology region of about

400 bp at the 5' end of the respective gene was used for targeted gene disruption. The

fragments were then inserted into pSLI-GFP (107) using Not*I* and Avr*II* restriction sites. For

*glmS* constructs, pSLI-GFP-*glmS* (108) was used as a vector instead. All oligonucleotides

and plasmids used in this study are listed in Supplementary Table S3.

**Transfection of *P. falciparum***

As described previously (109), parasites at the late schizont stage were purified using 60%

Percoll (105) and electroporated with 50 µg DNA of the respective plasmid in a 0.2-cm gap

cuvette (Bio-Rad Laboratories) using Amaxa Nucleofector 2b (Lonza). Either 4 nM WR99210

(Jacobus Pharmaceuticals) or 2 µg/mL blasticidin S (Life Technologies) was used for

selecting transfectants. For the selection of parasites that were genomically modified using

the SLI system (107), 400 µg/mL G418 (ThermoFisher Scientific) was added to the culture

medium once the parasitemia reached 5%. After the selection of modified parasites,

genomic DNA was isolated with the QIAamp DNA Mini Kit (Qiagen) and diagnostic tests for

correct integration into the genome were performed as specified earlier (107).

**Confocal live-cell microscopy**

Erythrocytes infected with parasites at different stages at 3 – 6% parasitemia were incubated

in culture medium with 20 nM MitoTracker Red, 200 nM ER Tracker Red or 100 nM

LysoTracker Deep Red (Invitrogen, if applicable) at 37°C for 20 min. Then, 200 nM Hoechst-

33342 (Invitrogen) was added for 10 min prior to washing the cells with Ringer's solution

(122.5 mM NaCl, 5.4 mM KCl, 1.2 mM CaCl$_2$, 0.8 mM MgCl$_2$, 11 mM D-glucose, 25 mM

HEPES, 1 mM NaH$_2$PO$_4$, pH 7.4) prewarmed to 37°C and seeding on a chambered No. 1.5

polymer cover slip (Ibidi). After 5 min, unbound erythrocytes were removed by washing with

Ringer's solution and the sample was placed into an incubation chamber that maintained the

microscope work area including the objective at 37°C. Images and videos were acquired

using an SP8 confocal microscope system with a 63x oil-corrected lens (C-Apochromat,

numerical aperture = 1.4) and Lightning deconvolution software (Leica), and processed

using ImageJ version 2.9.0/1.53t (110). If fluorescence intensities were to be quantified, no

averaging or deconvolution software was applied.

**Conditional knockdown mediated by *glmS* ribozyme**

For *glmS*-based knockdown induction (66), highly synchronous parasites at the early ring

stage were cultured with or without supplementation with 2.5 mM glucosamine (GlcN,

Sigma-Aldrich). The knockdown was quantified by confocal live-cell microscopy using

schizonts 36 h post GlcN treatment initiation. Images of parasites of similar size were

acquired with the same settings and background-corrected fluorescence intensities

(integrated density) as well as the size of the region of interest were determined using

ImageJ version 2.9.0/1.53t (110), and the data visualized using Graph Pad Prism version

9.4.1.

**Sample collection for RNA extraction and RNA sequencing**

Parasites were synchronized to a three-hour window after invasion of erythrocytes from the

respective donor as described above. Samples for RNA-sequencing were prepared in

triplicate for each condition and time point, i.e., three separate parasite cultures each were

grown in parallel for a total of at least two weeks. During the second IDC, two 10-mL dishes

each were harvested for parasites at the ring stage (6 – 9 hpi) and one 10-mL dish each for

trophozoites (26 – 29 hpi). Samples were collected by centrifuging the culture for 5 min at

800 g and 37°C and dissolving the erythrocyte pellet using 5 mL TRIzol (ThermoFisher

Scientific) prewarmed to 37°C, followed by immediate transfer to -80°C for storage. The

parasitemia was 0.3% at the start of the experiments with high, control and low-iron donor

blood and 2 – 3% at the time of harvest. Parasite cultures treated with 0.7 μM hepcidin

(Bachem) had a starting parasitemia of 0.6% and untreated cultures 1% to reach a

parasitemia of 4 – 5% during the second cycle. For each experiment, the parasitemia was

kept consistent at the point of harvest as high parasite densities can affect transcription (15).

For RNA extraction, the samples frozen in TRIzol were thawed, mixed thoroughly with 0.1

volume cold chloroform, and incubated at room temperature for 3 min. Following

centrifugation at 20,000 g and 4°C for 30 min, the supernatants were transferred to fresh

vials and combined with 70% ethanol of equal volume. RNA was purified using the RNeasy

MinElute Kit (Qiagen) by on-column DNase I digest for 30 min and elution with 14 μL water.

The GLOBINclear Human Kit (ThermoFisher Scientific) was then employed to deplete

human globin mRNA in all samples. The Qubit RNA HS Assay Kit and Qubit 3.0 fluorometer

(ThermoFisher Scientific) were used for RNA quantification. Upon arrival at the EMBL

Genomics Core Facility (GeneCore Heidelberg, Germany), the RNA quality of each sample
was evaluated using the RNA 6000 Nano kit and Bioanalyzer 2100 (Agilent). The median
RNA integrity number (RIN) of all samples was 7.30 (IQR: 6.85 – 8.15, Supplementary Fig.
S1). Individually barcoded strand-specific libraries for mRNA sequencing were prepared
from total RNA samples of high quality (approximately 150 ng per sample) using the
NEBNext® RNA Ultra II Directional RNA Library Prep Kit (New England Biolabs) for 12 PCR
cycles on the liquid handler Biomek i7 (Beckman Coulter) at GeneCore. Libraries that
passed quality control were pooled in equimolar amounts, and a 2 pM solution of this pool
was sequenced unidirectionally on a NextSeq® 500 System (Illumina) at GeneCore,
resulting in about 500 million reads of 85 bases each.
765

**RNA sequencing read mapping and data analysis**
Following successful initial quality control of the RNA-sequencing reads with FastQC version
0.11.8 (111), sequencing adapters were trimmed using Cutadapt version 2.10 (112). A
genome index was generated using the FASTA sequence file of the *P. falciparum* 3D7
genome release 46 (PlasmoDB-46_Pfalciparum3D7_Genome.fasta) and the GFF3
annotation file (PlasmoDB-46_Pfalciparum3D7.gff), both obtained from PlasmoDB (113),
with STAR version 2.7.5c (114). The same R package was used to align reads to
the genome with a maximum of three allowed mismatches (--outFilterMismatchNmax 3). To
consolidate the results obtained with FastQC and STAR alignments, a single report file was
created using MultiQC version 1.9 (115).
776

The mapped reads were then summarized in Sequence Alignment/Map (SAM) format using
featureCounts (116) from the R package Rsubread version 2.2.1 (117). For counting
mapped reads per gene using featureCounts, fragments with a minimum length of 50 bases
were considered (minFragLength = 50). Therefore, gene IDs and lengths of transcripts were
extracted from PlasmoDB-46_Pfalciparum3D7_AnnotatedTranscripts.fasta with SAMtools
faidx version 1.10.2 (118). The R package edgeR 3.30.3 (119) was used to compute RPKM
values (reads per kilobase per million mapped reads) and for differential gene expression
analysis. Gene annotations were retrieved from PlasmoDB (113) and PhenoPlasm (120).
The results of these analyses were visualized with volcano plots using the R package
Enhanced Volcano version 1.15.0 (121). The raw and processed data (FASTA files, RPKM
values and results of the differential gene expression analysis) can be accessed at
https://www.ebi.ac.uk/biostudies/studies/E-MTAB-13411.
789

The highly polymorphic *var*, *stevor*, and *rifin* gene families were excluded from downstream
analyses because of their great sequence diversity between parasites of the same strain

during mitotic growth (58, 59). Genes that were significantly regulated (defined as $P < 0.05$

according to the exact test for the negative binomial distribution with Benjamini-Hochberg

correction (54) and an absolute value of $\log_2 FC \geq 0.2$) were subjected to functional

enrichment analysis with g:Profiler (https://biit.cs.ut.ee/gprofiler/gost (60), accessed on

August 17, 2022). The resulting GO, KEGG and REAC terms were summarized using

REVIGO (http://revigo.irb.hr/) with the similarity value set to 0.5 (122) and visualized as in

Thomson-Luque et al. (83) using the scientific color map "roma" (123). To estimate parasite

age, an algorithm developed by Avi Feller and Jacob Lemieux (50) was adapted to use

expression data from Broadbent et al. (51) with the time points 6, 14, 20, 24, 28, 32, 36, 40,

44, and 48 hpi as reference. The code and data used were deposited to Zenodo with the

record ID 7996302 (https://zenodo.org/record/7996302).

**Protein structure prediction**

Structure predictions for monomeric proteins were obtained from AlphaFold Protein

Structure Database version 3 (67, 68) and homodimeric proteins were predicted using

AlphaFold2-multimer version 2.2.2, database version 2.2.0 (69) deployed at the EMBL

Hamburg computer cluster. Molecular visualization was performed with UCSF ChimeraX

version 1.3 (124). UCSF Chimera MatchMaker and Match → Align tools with default settings

were used for structural comparison of the predicted structures of *P. falciparum* proteins with

putative orthologs and sequence alignments were generated using the Match → Align tool

(125). The DeepFRI server (https://beta.deepfri.flatironinstitute.org) was used to identify

possible functional residues with the DeepFRI graph convolutional network (77).

**REFERENCES**

1.    Drakesmith H, Prentice A. Viral infection and iron metabolism. Nat Rev Microbiol.
2008;6(7):541-52.
2.    Zhang L, Hendrickson RC, Meikle V, Lefkowitz EJ, Ioerger TR, Niederweis M.
Comprehensive analysis of iron utilization by *Mycobacterium tuberculosis*. PLoS Pathog.
2020;16(2):e1008337.
3.    Kang D, Kirienko NV. Interdependence between iron acquisition and biofilm formation in
*Pseudomonas aeruginosa*. J Microbiol. 2018;56(7):449-57.
4.    Sousa Gerós A, Simmons A, Drakesmith H, Aulicino A, Frost JN. The battle for iron in enteric
infections. Immunology. 2020;161(3):186-99.
5.    Paganini D, Zimmermann MB. The effects of iron fortification and supplementation on the gut
microbiome and diarrhea in infants and children: a review. Am J Clin Nutr. 2017;106(Suppl 6):1688s-
93s.
6.    Brown RAM, Richardson KL, Kabir TD, Trinder D, Ganss R, Leedman PJ. Altered iron
metabolism and impact in cancer biology, metastasis, and immunology. Front Oncol. 2020;10:476.
7.    Taneri PE, Gómez-Ochoa SA, Llanaj E, Raguindin PF, Rojas LZ, Roa-Díaz ZM, et al. Anemia
and iron metabolism in COVID-19: a systematic review and meta-analysis. Eur J Epidemiol.
2020;35(8):763-73.
8.    Nyakeriga AM, Troye-Blomberg M, Dorfman JR, Alexander ND, Bäck R, Kortok M, et al. Iron
deficiency and malaria among children living on the coast of Kenya. J Infect Dis. 2004;190(3):439-47.

9.      Gwamaka M, Kurtis JD, Sorensen BE, Holte S, Morrison R, Mutabingwa TK, et al. Iron
deficiency protects against severe *Plasmodium falciparum* malaria and death in young children. Clin
Infect Dis. 2012;54(8):1137-44.
10.    Clark MA, Goheen MM, Fulford A, Prentice AM, Elnagheeb MA, Patel J, et al. Host iron status
and iron supplementation mediate susceptibility to erythrocytic stage Plasmodium falciparum. Nature
Communications. 2014;5:4446.
11.    Brabin L, Roberts SA, Tinto H, Gies S, Diallo S, Brabin B. Iron status of Burkinabé adolescent
girls predicts malaria risk in the following rainy season. Nutrients. 2020;12(5).
12.    Thipubon P, Uthaipibull C, Kamchonwongpaisan S, Tipsuwan W, Srichairatanakool S.
Inhibitory effect of novel iron chelator, 1-(N-acetyl-6-aminohexyl)-3-hydroxy-2-methylpyridin-4-one
(CM1) and green tea extract on growth of *Plasmodium falciparum*. Malaria J. 2015;14(1):382.
13.    Scholl PF, Tripathi AK, Sullivan DJ. Bioavailable iron and heme metabolism in *Plasmodium
falciparum*. Curr Top Microbiol Immunol. 2005;295:293-324.
14.    Brancucci NMB, Gerdt JP, Wang C, De Niz M, Philip N, Adapa SR, et al.
Lysophosphatidylcholine regulates sexual stage differentiation in the human malaria parasite
*Plasmodium falciparum*. Cell. 2017;171(7):1532-44.e15.
15.    Chou ES, Abidi SZ, Teye M, Leliwa-Sytek A, Rask TS, Cobbold SA, et al. A high parasite
density environment induces transcriptional changes and cell death in *Plasmodium falciparum* blood
stages. FEBS J. 2018;285(5):848-70.
16.    Mancio-Silva L, Slavic K, Grilo Ruivo MT, Grosso AR, Modrzynska KK, Vera IM, et al. Nutrient
sensing modulates malaria parasite virulence. Nature. 2017;547(7662):213-6.
17.    Sigala PA, Goldberg DE. The peculiarities and paradoxes of *Plasmodium* heme metabolism.
Annu Rev Microbiol. 2014;68:259-78.
18.    Mach J, Sutak R. Iron in parasitic protists – from uptake to storage and where we can
interfere. Metallomics. 2020;12(9):1335-47.
19.    Egan TJ, Combrinck JM, Egan J, Hearne GR, Marques HM, Ntenteni S, et al. Fate of haem
iron in the malaria parasite *Plasmodium falciparum*. Biochemical Journal. 2002;365(2):343-7.
20.    Loyevsky M, John C, Dickens B, Hu V, Miller JH, Gordeuk VR. Chelation of iron within the
erythrocytic *Plasmodium falciparum* parasite by iron chelators. Mol Biochem Parasitol. 1999;101(1-
2):43-59.
21.    Becker K, Tilley L, Vennerstrom JL, Roberts D, Rogerson S, Ginsburg H. Oxidative stress in
malaria parasite-infected erythrocytes: host–parasite interactions. Int J Parasitol. 2004;34(2):163-89.
22.    Wunderlich J, Rohrbach P, Dalton JP. The malaria digestive vacuole. Front Biosci (Schol Ed).
2012;4:1424-48.
23.    Kloehn J, Harding CR, Soldati-Favre D. Supply and demand - heme synthesis, salvage and
utilization by Apicomplexa. FEBS J. 2020;288(2):382-404.
24.    Huang G, Ulrich PN, Storey M, Johnson D, Tischer J, Tovar JA, et al. Proteomic analysis of
the acidocalcisome, an organelle conserved from bacteria to human cells. PLoS Pathog.
2014;10(12):e1004555.
25.    Ruiz FA, Luo S, Moreno SN, Docampo R. Polyphosphate content and fine structure of
acidocalcisomes of *Plasmodium falciparum*. Microsc Microanal. 2004;10(5):563-7.
26.    Magowan C, Brown JT, Liang J, Heck J, Coppel RL, Mohandas N, et al. Intracellular
structures of normal and aberrant *Plasmodium falciparum* malaria parasites imaged by soft x-ray
microscopy. Proc Natl Acad Sci U S A. 1997;94(12):6222-7.
27.    de Oliveira LS, Alborghetti MR, Carneiro RG, Bastos IMD, Amino R, Grellier P, et al. Calcium
in the backstage of malaria parasite biology. Front Cell Infect Microbiol. 2021;11:708834.
28.    Wunderlich J. Updated list of transport proteins in *Plasmodium falciparum*. Front Cell Infect
Microbiol. 2022;12:926541.
29.    Altschul SF, Madden TL, Schäffer AA, Zhang J, Zhang Z, Miller W, et al. Gapped BLAST and
PSI-BLAST: a new generation of protein database search programs. Nucleic Acids Res.
1997;25(17):3389-402.
30.    Slavic K, Krishna S, Lahree A, Bouyer G, Hanson KK, Vera I, et al. A vacuolar iron-
transporter homologue acts as a detoxifier in *Plasmodium*. Nat Commun. 2016;7:10403.
31.    Sharma P, Tóth V, Hyland EM, Law CJ. Characterization of the substrate binding site of an
iron detoxifying membrane transporter from *Plasmodium falciparum*. Malar J. 2021;20(1):295.
32.    Labarbuta P, Duckett K, Botting CH, Chahrour O, Malone J, Dalton JP, et al. Recombinant
vacuolar iron transporter family homologue *Pf*VIT from human malaria-causing *Plasmodium
falciparum* is a $Fe^{2+}/H^+$ exchanger. Sci Rep. 2017;7:42850.
33.    Aghabi D, Sloan M, Gill G, Hartmann E, Antipova O, Dou Z, et al. The vacuolar iron
transporter mediates iron detoxification in *Toxoplasma gondii*. Nat Commun. 2023;14(1):3659.

34. Sahu T, Boisson B, Lacroix C, Bischoff E, Richier Q, Formaglio P, et al. ZIPCO, a putative
metal ion transporter, is crucial for *Plasmodium* liver□stage development. EMBO Mol Med.
2014;6(11):1387-97.
35. Waller KL, Muhle RA, Ursos LM, Horrocks P, Verdier-Pinard D, Sidhu AB, et al. Chloroquine
resistance modulated *in vitro* by expression levels of the *Plasmodium falciparum* chloroquine
resistance transporter. J Biol Chem. 2003;278(35):33593-601.
36. Wichers JS, Mesén-Ramírez P, Fuchs G, Yu-Strzelczyk J, Stäcker J, von Thien H, et al.
PMRT1, a *Plasmodium*-specific parasite plasma membrane transporter, is essential for asexual and
sexual blood stage development. mBio. 2022;13(2):e00623-22.
37. Bakouh N, Bellanca S, Nyboer B, Moliner Cubel S, Karim Z, Sanchez CP, et al. Iron is a
substrate of the *Plasmodium falciparum* chloroquine resistance transporter *Pf*CRT in Xenopus
oocytes. J Biol Chem. 2017;292(39):16109-21.
38. Loveridge KM, Sigala PA. Unraveling mechanisms of iron acquisition in malaria parasites.
bioRxiv. 2024:2024.05.10.587216.
39. Brazzolotto X, Pierrel F, Pelosi L. Three conserved histidine residues contribute to
mitochondrial iron transport through mitoferrins. Biochem J. 2014;460(1):79-89.
40. Mühlenhoff U, Stadler JA, Richhardt N, Seubert A, Eickhorst T, Schweyen RJ, et al. A specific
role of the yeast mitochondrial carriers Mrs3/4p in mitochondrial iron acquisition under iron-limiting
conditions. J Biol Chem. 2003;278(42):40612-20.
41. Froschauer EM, Schweyen RJ, Wiesenberger G. The yeast mitochondrial carrier proteins
Mrs3p/Mrs4p mediate iron transport across the inner mitochondrial membrane. Biochim Biophys Acta.
2009;1788(5):1044-50.
42. Mather MW, Henry KW, Vaidya AB. Mitochondrial drug targets in apicomplexan parasites.
Curr Drug Targets. 2007;8(1):49-60.
43. Sloan MA, Aghabi D, Harding CR. Orchestrating a heist: uptake and storage of metals by
apicomplexan parasites. Microbiology (Reading). 2021;167(12).
44. Zhang D-L, Wu J, Shah BN, Greutélaers KC, Ghosh MC, Ollivierre H, et al. Erythrocytic
ferroportin reduces intracellular iron accumulation, hemolysis, and malaria risk. Science.
2018;359(6383):1520-3.
45. Aschemeyer S, Qiao B, Stefanova D, Valore EV, Sek AC, Ruwe TA, et al. Structure-function
analysis of ferroportin defines the binding site and an alternative mechanism of action of hepcidin.
Blood. 2018;131(8):899-910.
46. Billesbølle CB, Azumaya CM, Kretsch RC, Powers AS, Gonen S, Schneider S, et al. Structure
of hepcidin-bound ferroportin reveals iron homeostatic mechanisms. Nature. 2020.
47. Marfil-Rivera LJ. Iron overload. Medicina Universitaria. 2015;17(69):240-2.
48. Knovich MA, Storey JA, Coffman LG, Torti SV, Torti FM. Ferritin for the clinician. Blood Rev.
2009;23(3):95-104.
49. Petrat F, Rauen U, de Groot H. Determination of the chelatable iron pool of isolated rat
hepatocytes by digital fluorescence microscopy using the fluorescent probe, phen green SK.
Hepatology. 1999;29(4):1171-9.
50. Lemieux JE, Gomez-Escobar N, Feller A, Carret C, Amambua-Ngwa A, Pinches R, et al.
Statistical estimation of cell-cycle progression and lineage commitment in *Plasmodium falciparum*
reveals a homogeneous pattern of transcription in *ex vivo* culture. Proc Natl Acad Sci U S A.
2009;106(18):7559-64.
51. Broadbent KM, Broadbent JC, Ribacke U, Wirth D, Rinn JL, Sabeti PC. Strand-specific RNA
sequencing in *Plasmodium falciparum* malaria identifies developmentally regulated long non-coding
RNA and circular RNA. BMC Genomics. 2015;16(1):454.
52. Wichers JS, Scholz JAM, Strauss J, Witt S, Lill A, Ehnold LI, et al. Dissecting the gene
expression, localization, membrane topology, and function of the *Plasmodium falciparum* STEVOR
protein family. mBio. 2019;10(4).
53. Stewart LB, Diaz-Ingelmo O, Claessens A, Abugri J, Pearson RD, Goncalves S, et al. Intrinsic
multiplication rate variation and plasticity of human blood stage malaria parasites. Commun Biol.
2020;3(1):624.
54. Benjamini Y, Hochberg Y. Controlling the false discovery rate: a practical and powerful
approach to multiple Testing. J Royal Stat Soc Ser B. 1995;57(1):289-300.
55. Abrahamian M, Ah-Fong AM, Davis C, Andreeva K, Judelson HS. Gene expression and
silencing studies in *Phytophthora infestans* reveal infection-specific nutrient transporters and a role for
the nitrate reductase pathway in plant pathogenesis. PLoS Pathog. 2016;12(12):e1006097.

56.  Küpper H, Kochian LV. Transcriptional regulation of metal transport genes and mineral
nutrition during acclimatization to cadmium and zinc in the Cd/Zn hyperaccumulator, *Thlaspi*
*caerulescens* (Ganges population). New Phytol. 2010;185(1):114-29.
57.  Sarkans U, Gostev M, Athar A, Behrangi E, Melnichuk O, Ali A, et al. The BioStudies
database - one stop shop for all data supporting a life sciences study. Nucleic Acids Res.
2018;46(D1):D1266-D70.
58.  Bozdech Z, Llinás M, Pulliam BL, Wong ED, Zhu J, DeRisi JL. The transcriptome of the
intraerythrocytic developmental cycle of *Plasmodium falciparum*. PLoS Biol. 2003;1(1):E5.
59.  Kidgell C, Volkman SK, Daily J, Borevitz JO, Plouffe D, Zhou Y, et al. A systematic map of
genetic variation in *Plasmodium falciparum*. PLoS Pathog. 2006;2(6):e57.
60.  Raudvere U, Kolberg L, Kuzmin I, Arak T, Adler P, Peterson H, et al. g:Profiler: a web server
for functional enrichment analysis and conversions of gene lists (2019 update). Nucleic Acids Res.
2019;47(W1):W191-W8.
61.  Loyevsky M, LaVaute T, Allerson CR, Stearman R, Kassim OO, Cooperman S, et al. An IRP-
like protein from *Plasmodium falciparum* binds to a mammalian iron-responsive element. Blood.
2001;98(8):2555-62.
62.  Hodges M, Yikilmaz E, Patterson G, Kasvosve I, Rouault TA, Gordeuk VR, et al. An iron
regulatory-like protein expressed in *Plasmodium falciparum* displays aconitase activity. Mol Biochem
Parasitol. 2005;143(1):29-38.
63.  Lerch A, Koepfli C, Hofmann NE, Messerli C, Wilcox S, Kattenberg JH, et al. Development of
amplicon deep sequencing markers and data analysis pipeline for genotyping multi-clonal malaria
infections. BMC Genomics. 2017;18(1):864.
64.  Spork S, Hiss JA, Mandel K, Sommer M, Kooij TW, Chu T, et al. An unusual ERAD-like
complex is targeted to the apicoplast of *Plasmodium falciparum*. Eukaryot Cell. 2009;8(8):1134-45.
65.  Zhang M, Wang C, Otto TD, Oberstaller J, Liao X, Adapa SR, et al. Uncovering the essential
genes of the human malaria parasite *Plasmodium falciparum* by saturation mutagenesis. Science.
2018;360(6388):eaap7847.
66.  Prommana P, Uthaipibull C, Wongsombat C, Kamchonwongpaisan S, Yuthavong Y, Knuepfer
E, et al. Inducible knockdown of *Plasmodium* gene expression using the *glmS* ribozyme. PLoS One.
2013;8(8):e73783.
67.  Jumper J, Evans R, Pritzel A, Green T, Figurnov M, Ronneberger O, et al. Highly accurate
protein structure prediction with AlphaFold. Nature. 2021;596(7873):583-9.
68.  Varadi M, Anyango S, Deshpande M, Nair S, Natassia C, Yordanova G, et al. AlphaFold
Protein Structure Database: massively expanding the structural coverage of protein-sequence space
with high-accuracy models. Nucleic Acids Res. 2022;50(D1):D439-D44.
69.  Evans R, O'Neill M, Pritzel A, Antropova N, Senior A, Green T, et al. Protein complex
prediction with AlphaFold-Multimer. bioRxiv. 2022.
70.  Kato T, Kumazaki K, Wada M, Taniguchi R, Nakane T, Yamashita K, et al. Crystal structure of
plant vacuolar iron transporter VIT1. Nat Plants. 2019;5(3):308-15.
71.  Saier MH, Jr., Reddy VS, Tsu BV, Ahmed MS, Li C, Moreno-Hagelsieb G. The Transporter
Classification Database (TCDB): recent advances. Nucleic Acids Res. 2016;44(D1):D372-D9.
72.  Pang C, Chai J, Zhu P, Shanklin J, Liu Q. Structural mechanism of intracellular autoregulation
of zinc uptake in ZIP transporters. Nat Commun. 2023;14(1):3404.
73.  Martin RE, Henry RI, Abbey JL, Clements JD, Kirk K. The 'permeome' of the malaria parasite:
an overview of the membrane transport proteins of *Plasmodium falciparum*. Genome Biol.
2005;6(3):R26.
74.  Kim J, Tan YZ, Wicht KJ, Erramilli SK, Dhingra SK, Okombo J, et al. Structure and drug
resistance of the *Plasmodium falciparum* transporter PfCRT. Nature. 2019.
75.  Ehrnstorfer IA, Geertsma ER, Pardon E, Steyaert J, Dutzler R. Crystal structure of a SLC11
(NRAMP) transporter reveals the basis for transition-metal ion transport. Nat Struct Mol Biol.
2014;21(11):990-6.
76.  Meerstein-Kessel L, Venhuizen J, Garza D, Proellochs NI, Vos EJ, Obiero JM, et al. Novel
insights from the *Plasmodium falciparum* sporozoite-specific proteome by probabilistic integration of
26 studies. PLoS Comput Biol. 2021;17(4):e1008067.
77.  Gligorijević V, Renfrew PD, Kosciolek T, Leman JK, Berenberg D, Vatanen T, et al. Structure-
based protein function prediction using graph convolutional networks. Nat Commun. 2021;12(1):3168.
78.  Goheen MM, Bah A, Wegmüller R, Verhoef H, Darboe B, Danso E, et al. Host iron status and
erythropoietic response to iron supplementation determines susceptibility to the RBC stage of
falciparum malaria during pregnancy. Sci Rep. 2017;7(1):17674.

79.  Muriuki JM, Mentzer AJ, Mitchell R, Webb EL, Etyang AO, Kyobutungi C, et al. Malaria is a
cause of iron deficiency in African children. Nat Med. 2021;27(4):653-8.
80.  Kim CH, Kim HK, Bae SJ, Park JY, Lee KU. Association of elevated serum ferritin
concentration with insulin resistance and impaired glucose metabolism in Korean men and women.
Metabolism. 2011;60(3):414-20.
81.  Newhouse IJ, Clement DB, Lai C. Effects of iron supplementation and discontinuation on
serum copper, zinc, calcium, and magnesium levels in women. Med Sci Sports Exerc.
1993;25(5):562-71.
82.  Milner DA, Jr., Pochet N, Krupka M, Williams C, Seydel K, Taylor TE, et al. Transcriptional
profiling of *Plasmodium falciparum* parasites from patients with severe malaria identifies distinct low
vs. high parasitemic clusters. PLoS One. 2012;7(7):e40739.
83.  Thomson-Luque R, Votborg-Novél L, Ndovie W, Andrade CM, Niangaly M, Attipa C, et al.
*Plasmodium falciparum* transcription in different clinical presentations of malaria associates with
circulation time of infected erythrocytes. Nat Commun. 2021;12(1):4711.
84.  Lee HJ, Georgiadou A, Walther M, Nwakanma D, Stewart LB, Levin M, et al. Integrated
pathogen load and dual transcriptome analysis of systemic host-pathogen interactions in severe
malaria. Sci Transl Med. 2018;10(447).
85.  Muckenthaler MU, Rivella S, Hentze MW, Galy B. A red carpet for iron metabolism. Cell.
2017;168(3):344-61.
86.  Cercamondi CI, Egli IM, Ahouandjinou E, Dossa R, Zeder C, Salami L, et al. Afebrile
*Plasmodium falciparum* parasitemia decreases absorption of fortification iron but does not affect
systemic iron utilization: a double stable-isotope study in young Beninese women. Am J Clin Nutr.
2010;92(6):1385-92.
87.  Ginzburg YZ. Hepcidin-ferroportin axis in health and disease. Vitam Horm. 2019;110:17-45.
88.  Duraisingh MT, Skillman KM. Epigenetic variation and regulation in malaria parasites. Annu
Rev Microbiol. 2018;72:355-75.
89.  Ramos-Alonso L, Romero AM, Martínez-Pastor MT, Puig S. Iron regulatory mechanisms in
*Saccharomyces cerevisiae*. Front Microbiol. 2020;11:582830.
90.  Gilabert Carbajo C, Cornell LJ, Madbouly Y, Lai Z, Yates PA, Tinti M, et al. Novel aspects of
iron homeostasis in pathogenic bloodstream form *Trypanosoma brucei*. PLOS Pathogens.
2021;17(6):e1009696.
91.  Wang J, Pantopoulos K. Regulation of cellular iron metabolism. Biochem J. 2011;434(3):365-
81.
92.  Bushell E, Gomes AR, Sanderson T, Anar B, Girling G, Herd C, et al. Functional profiling of a
*Plasmodium* genome reveals an abundance of essential genes. Cell. 2017;170(2):260-72.e8.
93.  Li L, Murdock G, Bagley D, Jia X, Ward DM, Kaplan J. Genetic Dissection of a Mitochondria-
Vacuole Signaling Pathway in Yeast Reveals a Link between Chronic Oxidative Stress and Vacuolar
Iron Transport. J Biol Chem. 2010;285(14):10232-42.
94.  Schnell U, Dijk F, Sjollema KA, Giepmans BN. Immunolabeling artifacts and the need for live-
cell imaging. Nat Methods. 2012;9(2):152-8.
95.  Mathew R, Wunderlich J, Thivierge K, Cwiklinski K, Dumont C, Tilley L, et al. Biochemical and
cellular characterisation of the *Plasmodium falciparum* M1 alanyl aminopeptidase (*Pf*M1AAP) and
M17 leucyl aminopeptidase (*Pf*M17LAP). Sci Rep. 2021;11(1):2854.
96.  Schembri L, Dalibart R, Tomasello F, Legembre P, Ichas F, De Giorgi F. The HA tag is
cleaved and loses immunoreactivity during apoptosis. Nat Methods. 2007;4(2):107-8.
97.  Sidik SM, Huet D, Ganesan SM, Huynh MH, Wang T, Nasamu AS, et al. A genome-wide
CRISPR screen in *Toxoplasma* identifies essential apicomplexan genes. Cell. 2016;166(6):1423-
35.e12.
98.  Shrivastava D, Jha A, Kabrambam R, Vishwakarma J, Mitra K, Ramachandran R, et al.
*Plasmodium falciparum* ZIP1 is a zinc-selective transporter with stage-dependent targeting to the
apicoplast and plasma membrane in erythrocytic parasites. ACS Infect Dis. 2024;10(1):155-69.
99.  Rohrbach P, Friedrich O, Hentschel J, Plattner H, Fink RH, Lanzer M. Quantitative calcium
measurements in subcellular compartments of *Plasmodium falciparum*-infected erythrocytes. J Biol
Chem. 2005;280(30):27960-9.
100. Bozzi AT, Bane LB, Weihofen WA, Singharoy A, Guillen ER, Ploegh HL, et al. Crystal
structure and conformational change mechanism of a bacterial Nramp-family divalent metal
transporter. Structure. 2016;24(12):2102-14.
101. Bozzi AT, Bane LB, Zimanyi CM, Gaudet R. Unique structural features in an Nramp metal
transporter impart substrate-specific proton cotransport and a kinetic bias to favor import. J Gen
Physiol. 2019;151(12):1413-29.

102.    Smith EC, Limbach KJ, Rangel N, Oda K, Bolton JS, Du M, et al. Novel malaria antigen
*Plasmodium yoelii* E140 induces antibody-mediated sterile protection in mice against malaria
challenge. PLoS One. 2020;15(5):e0232234.
103.    Trager W, Jensen JB. Human malaria parasites in continuous culture. Science.
1976;193(4254):673-5.
104.    Malleret B, Claser C, Ong AS, Suwanarusk R, Sriprawat K, Howland SW, et al. A rapid and
robust tri-color flow cytometry assay for monitoring malaria parasite development. Sci Rep.
2011;1:118.
105.    Rivadeneira EM, Wasserman M, Espinal CT. Separation and concentration of schizonts of
*Plasmodium falciparum* by Percoll gradients. J Protozool. 1983;30(2):367-70.
106.    Lambros C, Vanderberg JP. Synchronization of *Plasmodium falciparum* erythrocytic stages in
culture. J Parasitol. 1979;65(3):418-20.
107.    Birnbaum J, Flemming S, Reichard N, Soares AB, Mesén-Ramírez P, Jonscher E, et al. A
genetic system to study *Plasmodium falciparum* protein function. Nat Methods. 2017;14(4):450-6.
108.    Burda PC, Crosskey T, Lauk K, Zurborg A, Söhnchen C, Liffner B, et al. Structure-based
identification and functional characterization of a lipocalin in the malaria parasite *Plasmodium
falciparum*. Cell Rep. 2020;31(12):107817.
109.    Moon RW, Hall J, Rangkuti F, Ho YS, Almond N, Mitchell GH, et al. Adaptation of the
genetically tractable malaria pathogen *Plasmodium knowlesi* to continuous culture in human
erythrocytes. Proc Natl Acad Sci U S A. 2013;110(2):531-6.
110.    Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, et al. Fiji: an
open-source platform for biological-image analysis. Nat Methods. 2012;9(7):676-82.
111.    Andrews S. FastQC: a quality control tool for high throughput sequence data. Babraham
Bioinformatics, Babraham Institute, Cambridge, UK; 2010.
112.    Martin M. Cutadapt removes adapter sequences from high-throughput sequencing reads.
EMBnet J. 2011;17(1):10-2.
113.    Aurrecoechea C, Brestelli J, Brunk BP, Dommer J, Fischer S, Gajria B, et al. PlasmoDB: a
functional genomic database for malaria parasites. Nucleic Acids Res. 2009;37(Database
issue):D539-D43.
114.    Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, et al. STAR: ultrafast
universal RNA-seq aligner. Bioinformatics. 2013;29(1):15-21.
115.    Ewels P, Magnusson M, Lundin S, Käller M. MultiQC: summarize analysis results for multiple
tools and samples in a single report. Bioinformatics. 2016;32(19):3047-8.
116.    Liao Y, Smyth GK, Shi W. featureCounts: an efficient general purpose program for assigning
sequence reads to genomic features. Bioinformatics. 2014;30(7):923-30.
117.    Liao Y, Smyth GK, Shi W. The Subread aligner: fast, accurate and scalable read mapping by
seed-and-vote. Nucleic Acids Res. 2013;41(10):e108.
118.    Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, et al. The Sequence
Alignment/Map format and SAMtools. Bioinformatics. 2009;25(16):2078-9.
119.    Robinson MD, McCarthy DJ, Smyth GK. edgeR: a Bioconductor package for differential
expression analysis of digital gene expression data. Bioinformatics. 2010;26(1):139-40.
120.    Sanderson T, Rayner JC. PhenoPlasm: a database of disruption phenotypes for malaria
parasite genes. Wellcome Open Res. 2017;2:45.
121.    Blighe K, Rana S, Lewis M. EnhancedVolcano: Publication-ready volcano plots with
enhanced colouring and labeling. R package version. 2022;1.15.
122.    Supek F, Bošnjak M, Škunca N, Šmuc T. REVIGO summarizes and visualizes long lists of
gene ontology terms. PLoS One. 2011;6(7):e21800.
123.    Crameri F, Shephard GE, Heron PJ. The misuse of colour in science communication. Nat
Commun. 2020;11(1):5444.
124.    Goddard TD, Huang CC, Meng EC, Pettersen EF, Couch GS, Morris JH, et al. UCSF
ChimeraX: Meeting modern challenges in visualization and analysis. Protein Sci. 2018;27(1):14-25.
125.    Meng EC, Pettersen EF, Couch GS, Huang CC, Ferrin TE. Tools for integrated sequence-
structure analysis with UCSF Chimera. BMC Bioinformatics. 2006;7:339.
126.    Birnbaum J, Scharf S, Schmidt S, Jonscher E, Hoeijmakers WAM, Flemming S, et al. A
Kelch13-defined endocytosis pathway mediates artemisinin resistance in malaria parasites. Science.
2020;367(6473):51-9.
127.    Ward DM, Kaplan J. Ferroportin-mediated iron transport: Expression and regulation.
Biochimica et Biophysica Acta (BBA) - Molecular Cell Research. 2012;1823(9):1426-33.

128. Mesén-Ramírez P, Bergmann B, Elhabiri M, Zhu L, von Thien H, Castro-Peña C, et al. The
parasitophorous vacuole nutrient channel is critical for drug access in malaria parasites and
modulates the artemisinin resistance fitness cost. Cell Host Microbe. 2021.
129. Sassmannshausen J, Pradel G, Bennink S. Perforin-like proteins of apicomplexan parasites.
Front Cell Infect Microbiol. 2020;10:578883.
130. Edayé S, Georges E. Characterization of native PfABCG protein in *Plasmodium falciparum*.
Biochem Pharmacol. 2015;97(2):137-46.
131. Jonker JW, Buitelaar M, Wagenaar E, Van Der Valk MA, Scheffer GL, Scheper RJ, et al. The
breast cancer resistance protein protects against a major chlorophyll-derived dietary phototoxin and
protoporphyria. Proc Natl Acad Sci U S A. 2002;99(24):15649-54.
132. Ahiya AI, Bhatnagar S, Morrisey JM, Beck JR, Vaidya AB. Dramatic consequences of
reducing erythrocyte membrane cholesterol on *Plasmodium falciparum*. Microbiol Spectr.
2022;10(1):e0015822.
133. Schmidt O, Pfanner N, Meisinger C. Mitochondrial protein import: from proteomics to
functional mechanisms. Nat Rev Mol Cell Biol. 2010;11(9):655-67.
134. Sheiner L, Soldati-Favre D. Protein trafficking inside *Toxoplasma gondii*. Traffic.
2008;9(5):636-46.
135. Moonah S, Sanders NG, Persichetti JK, Sullivan DJ, Jr. Erythrocyte lysis and *Xenopus laevis*
oocyte rupture by recombinant *Plasmodium falciparum* hemolysin III. Eukaryot Cell.
2014;13(10):1337-45.
136. Lim L, Linka M, Mullin KA, Weber AP, McFadden GI. The carbon and energy sources of the
non-photosynthetic plastid in the malaria parasite. FEBS Lett. 2010;584(3):549-54.
137. Hayashi M, Yamada H, Mitamura T, Horii T, Yamamoto A, Moriyama Y. Vacuolar $H^+$-ATPase
localized in plasma membranes of malaria parasite cells, *Plasmodium falciparum*, is involved in
regional acidification of parasitized erythrocytes. J Biol Chem. 2000;275(44):34353-8.
138. Friedrich O, Reiling SJ, Wunderlich J, Rohrbach P. Assessment of *Plasmodium falciparum*
PfMDR1 transport rates using Fluo-4. J Cell Mol Med. 2014;18(9):1851-62.
139. Marapana DS, Dagley LF, Sandow JJ, Nebl T, Triglia T, Pasternak M, et al. Plasmepsin V
cleaves malaria effector proteins in a distinct endoplasmic reticulum translocation interactome for
export to the erythrocyte. Nat Microbiol. 2018;3(9):1010-22.
140. Blume M, Hliscs M, Rodriguez-Contreras D, Sanchez M, Landfear S, Lucius R, et al. A
constitutive pan-hexose permease for the *Plasmodium* life cycle and transgenic models for screening
of antimalarial sugar analogs. Faseb j. 2011;25(4):1218-29.
141. Spillman Natalie J, Allen Richard JW, McNamara Case W, Yeung Bryan KS, Winzeler
Elizabeth A, Diagana Thierry T, et al. $Na^+$ regulation in the malaria parasite *Plasmodium falciparum*
involves the cation ATPase *Pf*ATP4 and is a target of the spiroindolone antimalarials. Cell Host
Microbe. 2013;13(2):227-37.
142. Boucher MJ, Ghosh S, Zhang L, Lal A, Jang SW, Ju A, et al. Integrative proteomics and
bioinformatic prediction enable a high-confidence apicoplast proteome in malaria parasites. PLoS
Biol. 2018;16(9):e2005895.
143. van Esveld SL, Meerstein-Kessel L, Boshoven C, Baaij JF, Barylyuk K, Coolen JPM, et al. A
prioritized and validated resource of mitochondrial proteins in *Plasmodium* identifies unique biology.
mSphere. 2021;6(5):e0061421.
144. Wichers JS, van Gelder C, Fuchs G, Ruge JM, Pietsch E, Ferreira JL, et al. Characterization
of Apicomplexan Amino Acid Transporters (ApiATs) in the malaria parasite *Plasmodium falciparum*.
mSphere. 2021;6(6):e00743-21.
145. Gardner MJ, Hall N, Fung E, White O, Berriman M, Hyman RW, et al. Genome sequence of
the human malaria parasite *Plasmodium falciparum*. Nature. 2002;419(6906):498-511.
146. Martin RE. The transportome of the malaria parasite. Biol Rev Camb Philos Soc.
2020;95(2):305-32.
147. Carter NS, Ben Mamoun C, Liu W, Silva EO, Landfear SM, Goldberg DE, et al. Isolation and
functional characterization of the *Pf*NT1 nucleoside transporter gene from *Plasmodium falciparum*. J
Biol Chem. 2000;275(14):10683-91.
148. Kenthirapalan S, Waters AP, Matuschewski K, Kooij TW. Copper-transporting ATPase is
important for malaria parasite fertility. Mol Microbiol. 2014;91(2):315-25.
149. Park DJ, Lukens AK, Neafsey DE, Schaffner SF, Chang HH, Valim C, et al. Sequence-based
association and selection scans identify drug resistance loci in the *Plasmodium falciparum* malaria
parasite. Proc Natl Acad Sci U S A. 2012;109(32):13052-67.

150.    Sayers CP, Mollard V, Buchanan HD, McFadden GI, Goodman CD. A genetic screen in
rodent malaria parasites identifies five new apicoplast putative membrane transporters, one of which
is essential in human malaria parasites. Cell Microbiol. 2018;20(1):e12789.
151.    Cowell AN, Istvan ES, Lukens AK, Gomez-Lorenzo MG, Vanaerschot M, Sakata-Kato T, et al.
Mapping the malaria parasite druggable genome by using *in vitro* evolution and chemogenomics.
Science. 2018;359(6372):191-9.
152.    Gupta Y, Sharma N, Singh S, Romero JG, Rajendran V, Mogire RM, et al. The multistage
antimalarial compound calxinin perturbates *P. falciparum* $Ca^{2+}$ homeostasis by targeting a unique ion
channel. Pharmaceutics. 2022;14(7):1371.
153.    Nozawa A, Ito D, Ibrahim M, Santos HJ, Tsuboi T, Tozawa Y. Characterization of
mitochondrial carrier proteins of malaria parasite *Plasmodium falciparum* based on *in vitro* translation
and reconstitution. Parasitol Int. 2020;79:102160.
154.    Agrawal S, Striepen B. More membranes, more proteins: complex protein import mechanisms
into secondary plastids. Protist. 2010;161(5):672-87.
155.    Schureck MA, Darling JE, Merk A, Shao J, Daggupati G, Srinivasan P, et al. Malaria parasites
use a soluble RhopH complex for erythrocyte invasion and an integral form for nutrient uptake. eLife.
2021;10:e65282.
156.    Chitale M, Hawkins T, Park C, Kihara D. ESG: extended similarity group method for
automated protein function prediction. Bioinformatics. 2009;25(14):1739-45.
157.    Stanway RR, Bushell E, Chiappino-Pepe A, Roques M, Sanderson T, Franke-Fayard B, et al.
Genome-scale identification of essential metabolic processes for targeting the *Plasmodium* liver
stage. Cell. 2019;179(5):1112-28.e26.
158.    Kuhn Y, Sanchez CP, Ayoub D, Saridaki T, Van Dorsselaer A, Lanzer M. Trafficking of the
phosphoprotein PfCRT to the digestive vacuolar membrane in *Plasmodium falciparum*. Traffic.
2010;11(2):236-49.
159.    Ecker A, Lakshmanan V, Sinnis P, Coppens I, Fidock DA. Evidence that mutant *Pf*CRT
facilitates the transmission to mosquitoes of chloroquine-treated *Plasmodium* gametocytes. J Infect
Dis. 2011;203(2):228-36.
1216

**Figure legends:**

**Figure 1: Effects of the iron status of the blood donor and of hepcidin on (A)**

**erythrocyte labile iron levels, (B) *P. falciparum* 3D7 growth rates, (C) DNA content per**

**mature schizont, and (D) the number of merozoites per mature schizont.**

The relative labile iron level and DNA content per cell were assessed in the presence or

absence of 0.7 µM hepcidin (Hep) by measuring the mean fluorescence intensity (MFI) of

Phen Green SK or SYBR Green I compared to control (Ctrl, untreated, normal hemoglobin

level) using flow cytometry. Therefore, 100,000 cells were analyzed per replicate. Parasite

growth rates refer to the fold change in parasitemia after one intraerythrocytic developmental

cycle in vitro relative to control as determined by flow cytometry with SYBR Green I (104).

Mature schizonts were obtained by treating schizonts at 40 hpi with 1 mM compound 2 (4-[7-

[(dimethylamino)methyl]-2-(4-fluorphenyl)imidazo[1,2-α]pyridin-3-yl]pyrimidin-2-amine) for 8

h. To count the number of merozoites, Giemsa-stained blood smears were analyzed

microscopically and only single-infected cells with one digestive vacuole were considered.

Means and 95% confidence intervals (indicated by error bars) are shown. Statistical

significance was calculated with two-tailed unpaired *t* tests with Welch's correction for

unequal variances and adjusted with the Holm-Šídák method for multiple comparisons

except for merozoite numbers, which were compared using Mann-Whitney test. N

represents the number of parasites and n the number of independent experiments.

1236

**Figure 2: Differential expression of *P. falciparum* 3D7 genes under various iron**

**conditions.**

Parasites were cultured with erythrocytes from an individual with high, medium (healthy) or

low iron status (experiment 1) or with red blood cells from another healthy donor in the

presence or absence of 0.7 µM hepcidin (experiment 2). Samples were harvested at the ring

and trophozoite stage (6 – 9 and 26 – 29 hours post invasion, hpi) with three biological

replicates per time point and condition. The maximum likelihood estimation (MLE) of the

average developmental age of the parasites for each condition and time point (**A**) was

calculated using an algorithm developed by Avi Feller and Jacob Lemieux (50). CI,

confidence interval. The volcano plots (**B** and **D**) show transcriptional changes of all parasite

genes. Red dots indicate significantly ($P < 0.05$, exact test for negative binomial distribution)

upregulated genes ($\log_2$ (fold change) ≥ 0.2), blue dots stand for significantly downregulated

genes ($\log_2$ (fold change) ≤ -0.2), while grey dots represent genes that did not significantly

differ in transcription under the conditions described ($P ≥ 0.05$ and / or -0.2 < $\log_2$ (fold

change) < 0.2). Differentially expressed genes encoding putative iron transporters (see

Table 1) are labeled. Panels **C** and **E** show the enrichment of Gene Ontology (GO), Kyoto

Encyclopedia of Genes and Genomes (KEGG) and Reactome (REAC) terms among

significantly regulated genes excluding *var, stevor* and *rifin* gene families at the two time

points. The functional terms were summarized using REVIGO (122) to remove

redundancies, represented by circles and plotted according to the significance of their

enrichment (-$\log_{10}$ (adjusted *P*), hypergeometric test). The size of the circle is proportional to

the number of differentially regulated genes in the dataset that are associated with the

respective term, while the color stands for the fold enrichment. The gray dashed line

indicates the threshold of the adjusted *P* value (-$\log_{10}$ 0.05 = 1.3).

1261

**Figure 3: Subcellular localization of known and putative iron transport proteins.**

Representative erythrocytes infected with *P. falciparum* 3D7 parasites endogenously

expressing GFP-tagged *Pf*MRS3 (**A**), *Pf*VIT (**B**), *Pf*ZIPCO (**C**) or *Pf*E140 (**D**) were

additionally stained with the fluorescent dyes Hoechst-33342, MitoTracker Red, ER Tracker

Red and/or LysoTracker Deep Red. Co-transfection with a construct that encodes the 60 N-

terminal amino acids of acyl carrier protein (*Pf*ACP) tagged with mCherry (126) resulted in

red fluorescence of the apicoplast. Live-cell images were taken under physiological

conditions at 37°C using an SP8 confocal laser-scanning microscope (Leica). DIC,

differential interference contrast. Scale bar, 2 µm.

1271

**Figure 4: *Pf*VIT and *Pf*E140 are important for *P. falciparum* growth and may be involved in intracellular iron homeostasis.**

**A** Representative erythrocytes infected with *P. falciparum* 3D7 parasites that endogenously express a truncated version of *Pf*VIT or *Pf*ZIPCO tagged with GFP (green). **B** Growth rates of knockout parasite lines generated. **C** Reduction of *Pf*E140-GFP fluorescence (green) in live 3D7 parasites caused by *glmS*-mediated knockdown that was induced by treatment with 2.5 mM glucosamine (GlcN) for 36 h compared to untreated control (Ctrl). **D** Total parasite fluorescence intensities were quantified as background-corrected integrated densities using ImageJ version 2.9.0/1.53t (110) and compared using Mann-Whitney test. **E** The size of the parasites was measured as the area of the region of interest and compared using equal variance unpaired *t* test. **F** Conditional knockdown of *Pf*E140 induced by treatment with 2.5 mM GlcN results in a growth defect during asexual blood stage development. Live parasites were stained with Hoechst-33342 (blue) and imaged under physiological conditions at 37°C using an SP8 confocal laser-scanning microscope (Leica). DIC, differential interference contrast. Scale bar, 2 µm. Error bars represent 95% confidence intervals of the mean, N the number of parasites analyzed, n the number of independent experiments and Hep treatment with 0.7 µM hepcidin. Growth rates refer to the fold change in parasitemia after two intraerythrocytic developmental cycles in vitro relative to untreated wild-type 3D7 parasites (WT) as determined by flow cytometry with SYBR Green I (104). Statistical significance of growth differences was calculated with two-tailed unpaired *t* tests with Welch's correction for unequal variances and adjusted with the Holm-Šídák method for multiple comparisons.

**Figure 5: Structures of known and putative *P. falciparum* iron transporters as viewed from the membrane plane.**

**A** Predicted protein structures with per-residue pLDDT (predicted local distance difference test) confidence scores on a scale from 0 to 100, where blue represents high and red low confidence, respectively. The experimentally determined structure of *Pf*CRT is shown in gray. **B** Molecular lipophilicity potential of the protein surfaces as implemented in UCSF ChimeraX; tan is hydrophobic and cyan hydrophilic. Dashed lines above and below the tan regions of all proteins indicate the respective membrane and disordered loops were removed for clarity. **C** Surface charge of the proteins with positively charged areas colored blue and negatively charged ones red. Putative cation-binding site are indicated with an asterisk and transport directions by arrows. *Pf*E140 likely forms a dimer but is shown as a monomer, as no predicted dimer structure could be obtained using AlphaFold2-multimer. The putative cation-binding sites for this protein are based on DeepFRI gradCAM scores for the functional term GO:0015075 "monoatomic ion transmembrane transporter activity" (Supplementary Fig. S5).

1309

**Figure 6: Iron homeostasis in a *P. falciparum*-infected erythrocyte.**

The human blood plasma contains between 10 and 30 μM total iron and the erythrocyte

cytosol approximately 20 mM (19). However, the labile iron pool is only 3 μM in an

uninfected erythrocyte and 1.6 μM in a *P. falciparum*-infected one (20). Human ferroportin

(FPN) at the host cell surface (erythrocyte plasma membrane, EPM) exports ferrous iron

from the erythrocyte (127) and the nutrient pore formed by *Pf*EXP1 and *Pf*EXP2 allows the

passage of ions through the parasitophorous vacuole membrane (PVM) (128). *Pf*E140 at the

parasite plasma membrane (PPM) may mediate iron uptake into the parasite cytosol and the

mitochondrial carrier protein *Pf*MRS3 likely translocates $Fe^{2+}$ into the mitochondrion, a site of

*de novo* heme biosynthesis (this study). We propose that the vacuolar iron transporter

(*Pf*VIT) is involved in iron detoxification by transporting excess $Fe^{2+}$ from the cytosol into

cytoplasmic vesicles that may be acidocalcisomes, whereas *Pf*ZIPCO releases $Fe^{2+}$ from

these organelles under low-iron conditions. The digestive vacuole (DV) contains a high

amount of total iron as it is the site of hemoglobin degradation and hemozoin formation (21).

The chloroquine resistance transporter (*Pf*CRT) and the natural resistance-associated

macrophage protein (*Pf*NRAMP, also called *Pf*DMT1 for divalent metal transporter 1) were

suggested to mediate proton-coupled export of $Fe^{2+}$ from the DV into the parasite cytosol

(37, 73). Both acidocalcisomes and the DV are likely acidified by the plant-like $H^+$-pump V-

ATPase, which can fuel secondary active transport processes (22, 27). Parasite-encoded

proteins are shown in orange and human-encoded transporters in blue.

1330

**Tables:**

 **Table 1: *P. falciparum* transport proteins with differential gene expression under various iron conditions.**

Putative and known transporter genes were filtered from differentially expressed genes in the described RNA-sequencing experiments using a list of *P. falciparum* transport proteins (28). The $\log_2$ (fold change) of gene expression at the ring stage (6 – 9 hours post invasion) and known or proposed functions are indicated for significantly regulated genes (exact $P < 0.05$). The identified (potential) iron transport proteins are highlighted in red. DV, digestive vacuole; EPM, erythrocyte plasma membrane; PPM, parasite plasma membrane.

| Gene product and ID | $\log_2$(fold change) high vs. low Fe | $\log_2$(fold change) low Fe vs. control | $\log_2$(fold change) hepcidin vs. control | Known or putative function |
|---|---|---|---|---|
| PLP5 (PF3D7_0819200) | + 0.49 ($P = 0.0002$) | n.s. | - 0.29 ($P = 0.003$) | Host cell permeabilization and rupture (129) |
| ABCG (PF3D7_1426500) | + 0.45 ($P = 0.02$) | - 0.57 ($P = 0.006$) | n.s. | Putative metabolite exporter at PPM (130), human ortholog ABCG2 exports heme (131) |
| VP1 (PF3D7_1456800) | + 0.39 ($P = 0.0006$) | - 0.34 ($P = 0.004$) | n.s. | Active $H^+$ export across PPM (132) |
| TOM7 (PF3D7_0823700) | + 0.38 ($P = 0.02$) | n.s. | n.s. | Protein import across outer mitochondrial membrane (133, 134) |
| HlyIII (PF3D7_1455400) | + 0.38 ($P = 0.0009$) | n.s. | n.s. | Forms pore (~3.2 nm) for solutes and ions in EPM (135) |
| TPT (PF3D7_0508300) | + 0.37 ($P = 0.001$) | - 0.40 ($P = 0.002$) | n.s. | Imports phosphoenolpyruvate, dihydroxyacetone, and 3-phosphoglycerate across outer apicoplast membrane (136) |
| $V_o$ c (PF3D7_0519200) | + 0.36 ($P = 0.0006$) | n.s. | - 0.16 ($P = 0.007$) | V-ATPase subunit: active $H^+$ export from cytosol (137) |
| MDR1 (PF3D7_0523000) | + 0.34 ($P = 0.0009$) | - 0.23 ($P = 0.03$) | n.s. | Active drug and solute import into DV (138) |
| NT3 (PF3D7_1469400) | + 0.34 ($P = 0.03$) | n.s. | n.s. | Putative nucleoside transporter (73) |
| SEC61α (PF3D7_1346100) | + 0.33 ($P = 0.0006$) | - 0.22 ($P = 0.04$) | - 0.17 ($P = 0.005$) | ER import of proteins destined for export (139) |
| <span style="color:red">MRS3 (PF3D7_0905200)</span> | <span style="color:red">+ 0.33 ($P = 0.002$)</span> | <span style="color:red">- 0.19 ($P = 0.04$)</span> | <span style="color:red">n.s.</span> | <span style="color:red">Putative $Fe^{2+}$ importer into mitochondrial matrix (40)</span> |
| HT1 (PF3D7_0204700) | + 0.33 ($P = 0.002$) | - 0.30 ($P = 0.006$) | n.s. | Imports glucose and fructose across PPM (140) |
| ATP4 (PF3D7_1211900) | + 0.33 ($P = 0.005$) | - 0.40 ($P = 0.002$) | n.s. | $H^+$ import, $Na^+$ export across PPM (141) |
| MCT2 (PF3D7_0926400) | + 0.32 ($P = 0.005$) | n.s. | n.s. | Exports organic solutes from apicoplast, imports $H^+$ (142) |
| $V_1$ B (PF3D7_0406100) | + 0.30 ($P = 0.001$) | - 0.25 ($P = 0.008$) | - 0.13 ($P = 0.04$) | V-ATPase subunit: active $H^+$ export from cytosol (137) |
| $V_1$ A (PF3D7_1311900) | + 0.30 ($P = 0.001$) | - 0.28 ($P = 0.003$) | - 0.20 ($P = 0.001$) | V-ATPase subunit: active $H^+$ export from cytosol (137) |
| TIM16 (PF3D7_0513500) | + 0.29 ($P = 0.03$) | n.s. | n.s. | Protein import across inner mitochondrial membrane (133, 143) |
| <span style="color:red">VIT (PF3D7_1223700)</span> | <span style="color:red">+ 0.29 ($P = 0.02$)</span> | <span style="color:red">n.s.</span> | <span style="color:red">n.s.</span> | <span style="color:red">$Fe^{2+}$ sequestration from cytosol in exchange for $H^+$ (30-32)</span> |
| ApiAT2 (PF3D7_0914700) | + 0.28 ($P = 0.006$) | n.s. | n.s. | Putative amino acid transporter at PPM (144) |
| <span style="color:red">NRAMP (PF3D7_0523800)</span> | <span style="color:red">+ 0.28 ($P = 0.003$)</span> | <span style="color:red">n.s.</span> | <span style="color:red">n.s.</span> | <span style="color:red">Symport of $Mn^{2+}$ or $Fe^{2+}$ with $H^+$ from DV into cytosol (22, 36)</span> |
| AMC1 (PF3D7_0108800) | + 0.26 ($P = 0.03$) | n.s. | n.s. | Putative mitochondrial transporter (145) |
| <span style="color:red">CRT (PF3D7_0709000)</span> | <span style="color:red">+ 0.26 ($P = 0.007$)</span> | <span style="color:red">n.s.</span> | <span style="color:red">n.s.</span> | <span style="color:red">Symport of positively charged dipeptides or $Fe^{2+}$ with $H^+$ from DV into cytosol (37)</span> |
| $V_1$ H (PF3D7_1306600) | + 0.26 ($P = 0.01$) | n.s. | n.s. | V-ATPase subunit: active $H^+$ export from cytosol (137) |
| AAC2 (PF3D7_1004800) | + 0.25 ($P = 0.03$) | n.s. | - 0.21 ($P = 0.04$) | Mitochondrial ADP/ATP antiporter (143) |
| PiT (PF3D7_1340900) | + 0.24 ($P = 0.02$) | - 0.21 ($P = 0.04$) | n.s. | Imports phosphate and $Na^+$ into cytosol across PPM (73) |
| $V_o$ d (PF3D7_1464700) | + 0.24 ($P = 0.02$) | - 0.23 ($P = 0.03$) | - 0.15 ($P = 0.02$) | V-ATPase subunit: active $H^+$ export from cytosol (137) |
| $V_o$ c" (PF3D7_1354400) | + 0.23 ($P = 0.03$) | n.s. | n.s. | V-ATPase subunit: active $H^+$ export from cytosol (137) |
| ATP10 (PF3D7_0727800) | + 0.23 ($P = 0.01$) | n.s. | n.s. | Active apicoplast $Mn^{2+}$ transporter (146) |
| SulP (PF3D7_1471200) | + 0.23 ($P = 0.02$) | n.s. | n.s. | Inorganic anion antiporter at PPM (73) |
| ATP2 (PF3D7_1219600) | + 0.22 ($P = 0.02$) | - 0.22 ($P < 0.05$) | - 0.14 ($P = 0.03$) | Putative phospholipid flippase at PPM (73) |
| NT1 (PF3D7_1347200) | + 0.22 ($P = 0.02$) | n.s. | n.s. | Purin base import across PPM (147) |
| AAT1 (PF3D7_0629500) | - 0.26 ($P = 0.007$) | n.s. | n.s. | Putative amino acid transporter at PPM and DV (145) |
| MIT1 (PF3D7_1120300) | - 0.27 ($P = 0.02$) | n.s. | n.s. | Putative mitochondrial magnesium/nickel/cobalt ion channel (143) |
| AAT2 (PF3D7_1208400) | - 0.29 ($P = 0.009$) | n.s. | n.s. | Putative amino acid transporter at PPM (73) |
| MDR7 (PF3D7_1209900) | - 0.34 ($P = 0.04$) | n.s. | n.s. | Active efflux of peptides from mitochondrion (42, 73) |
| CuTP (PF3D7_0904900) | - 0.36 ($P = 0.008$) | + 0.52 ($P = 0.004$) | n.s. | Active $Cu^{2+}$ export from cytoplasmic vesicle (73, 148) |

| | | | | |
|---|---|---|---|---|
| GCβ (PF3D7_1360500) | - 0.36 ($P = 0.007$) | n.s. | n.s. | Putative phospholipid flippase at PPM (73) |
| TPC (PF3D7_1368700) | - 0.37 ($P = 0.01$) | + 0.33 ($P = 0.02$) | n.s. | Thiamine pyrophosphate import, nucleotide export from mitochondrion (146) |
| $F_1$ γ (PF3D7_1311300) | - 0.38 ($P = 0.03$) | n.s. | + 0.32 ($P = 0.03$) | Subunit of mitochondrial $H^+$-importing ATP synthase (145) |
| PF3D7_0614900 | - 0.43 ($P = 0.005$) | + 0.72 ($P = 0.0001$) | + 0.29 ($P = 0.0003$) | Putative ABC transporter at PPM (28) |
| MFS4 (PF3D7_1203400) | - 0.44 ($P = 0.0008$) | + 0.56 ($P = 0.0005$) | + 0.27 ($P = 0.02$) | Putative transporter (73) |
| $F_o$ d (PF3D7_0311800) | - 0.46 ($P = 0.004$) | + 0.34 ($P = 0.02$) | n.s. | Subunit of mitochondrial $H^+$-importing ATP synthase (146) |
| PF3D7_1004600 | - 0.47 ($P = 0.01$) | + 0.58 ($P = 0.003$) | + 0.28 ($P = 0.03$) | Putative ABC transporter linked to drug resistance (149) |
| MPC2 (PF3D7_1470400) | - 0.47 ($P = 0.03$) | + 0.66 ($P = 0.005$) | n.s. | Subunit of mitochondrial putative pyruvate:$H^+$ importer (146) |
| <span style="color:red">E140 (PF3D7_0104100)</span> | <span style="color:red">- 0.47 ($P = 0.03$)</span> | <span style="color:red">+ 0.65 ($P = 0.006$)</span> | <span style="color:red">n.s.</span> | <span style="color:red">Putative transport protein at PPM (28)</span> |
| MDR4 (PF3D7_0302600) | - 0.55 ($P = 0.001$) | + 0.63 ($P = 0.001$) | + 0.33 ($P = 0.02$) | Active drug export across innermost apicoplast membrane (150, 151) |
| <span style="color:red">ZIPCO (PF3D7_1022300)</span> | <span style="color:red">- 0.57 ($P = 0.03$)</span> | <span style="color:red">+ 0.55 ($P = 0.04$)</span> | <span style="color:red">+ 0.63 ($P = 0.006$)</span> | <span style="color:red">$Fe^{2+}$/$Zn^{2+}$ import into cytosol (34)</span> |
| $F_1$ δ (PF3D7_1147700) | - 0.59 ($P = 0.004$) | + 0.63 ($P = 0.005$) | n.s. | Subunit of mitochondrial $H^+$-importing ATP synthase (145) |
| TRP-ML (PF3D7_1313500) | - 0.59 ($P = 0.01$) | + 0.64 ($P = 0.005$) | n.s. | Putative $Ca^{2+}$ channel (152) |
| $F_1$ ε (PF3D7_0715500) | - 0.67 ($P = 0.04$) | + 0.94 ($P = 0.007$) | n.s. | Subunit of mitochondrial $H^+$-importing ATP synthase (145) |
| MFS3 (PF3D7_0919500) | - 0.85 ($P = 0.01$) | + 0.72 ($P = 0.01$) | n.s. | Putative sugar transporter (145) |
| ATP11 (PF3D7_1468600) | n.s. | + 1.14 ($P = 0.005$) | n.s. | Putative phospholipid flippase at PPM (28, 73) |
| NGT (PF3D7_0505300) | n.s. | + 0.91 ($P = 0.00008$) | + 0.79 ($P = 0.00004$) | UDP-N-acetylglucosamine import, UMP export from Golgi (73) |
| SAMC (PF3D7_1241600) | n.s. | + 0.77 ($P = 0.03$) | + 0.68 ($P = 0.02$) | S-adenosylmethionine import into mitochondrion (153) |
| PF3D7_0614900 | n.s. | + 0.72 ($P = 0.0001$) | + 0.29 ($P = 0.0003$) | Putative active transporter at PPM (28) |
| GPH (PF3D7_0529200) | n.s. | + 0.70 ($P = 0.03$) | n.s. | Putative sugar:cation symporter (73) |
| ATP9 (PF3D7_1348800) | n.s. | + 0.62 ($P = 0.04$) | n.s. | Active $Ca^{2+}$ import into DV? (146) |
| CTR2 (PF3D7_1421900) | n.s. | + 0.52 ($P = 0.004$) | n.s. | Putative apicoplast copper channel (146) |
| MDR5 (PF3D7_1339900) | n.s. | + 0.51 ($P = 0.02$) | n.s. | Active solute export across PPM (146) |
| TIC20 (PF3D7_1144700) | n.s. | + 0.49 ($P = 0.007$) | n.s. | Protein import across innermost apicoplast membrane (154) |
| PF3D7_0924500 | n.s. | + 0.47 ($P = 0.002$) | n.s. | Putative $Na^+$/$H^+$ exchanger (71) |
| PPT (PF3D7_0530200) | n.s. | + 0.46 ($P = 0.02$) | n.s. | Imports phosphoenolpyruvate, dihydroxyacetone, and 3-phosphoglycerate across inner apicoplast membrane (136) |
| CLAG3.1 (PF3D7_0302500) | n.s. | + 0.45 ($P < 0.05$) | n.s. | Purine, amino acid, sugar, and vitamin import across EPM (155) |
| CDF (PF3D7_0715900) | n.s. | + 0.44 ($P = 0.02$) | n.s. | Putative $Zn^{2+}$ importer (43) into cytoplasmic vesicles (36) |
| ATP7 (PF3D7_0319000) | n.s. | + 0.44 ($P = 0.007$) | n.s. | Putative phospholipid flippase at PPM (28, 73) |
| RhopH3 (PF3D7_0905400) | n.s. | + 0.42 ($P = 0.04$) | n.s. | Purine, amino acid, sugar, and vitamin import across EPM (155) |
| AQP2 (PF3D7_0810400) | n.s. | + 0.34 ($P = 0.02$) | - 0.19 ($P < 0.05$) | Water channel at PPM (156) |
| MFS2 (PF3D7_0916000) | n.s. | + 0.33 ($P = 0.04$) | n.s. | Putative sugar transporter (73) |
| TFP1 (PF3D7_0206200) | n.s. | + 0.33 ($P = 0.02$) | n.s. | Putative metabolite transporter at PPM (146) |
| ATP6 (PF3D7_0106300) | n.s. | - 0.22 ($P = 0.03$) | n.s. | Active $Ca^{2+}$ import into ER for storage (146) |
| $V_1$ G (PF3D7_1323200) | n.s. | - 0.23 ($P = 0.04$) | - 0.12 ($P = 0.04$) | V-ATPase subunit: active $H^+$ export from cytosol (137) |
| FNT (PF3D7_0316600) | n.s. | - 0.27 ($P = 0.02$) | - 0.16 ($P = 0.004$) | Lactate/formate and $H^+$ release from cytosol (146) |
| MFR1 (PF3D7_0614300) | n.s. | - 0.33 ($P = 0.004$) | - 0.14 ($P = 0.02$) | Putative organic anion transporter (73) |
| PMRT1 (PF3D7_1135300) | n.s. | - 0.36 ($P = 0.009$) | n.s. | Putative transporter at PPM (36) |
| PLP3 (PF3D7_0923300) | n.s. | - 0.42 ($P = 0.015$) | n.s. | Host cell permeabilization and rupture (129) |
| OSCP (PF3D7_1310000) | n.s. | n.s. | + 0.58 ($P = 0.01$) | Subunit of mitochondrial $H^+$-importing ATP synthase (145) |
| GFT (PF3D7_0212000) | n.s. | n.s. | + 0.54 ($P = 0.03$) | GDP-fucose import, GMP export from Golgi (73) |
| SEC61γ (PF3D7_0210000) | n.s. | n.s. | - 0.23 ($P = 0.001$) | ER import of proteins destined for export (139) |
| SEC61β (PF3D7_0821800) | n.s. | n.s. | - 0.31 ($P = 0.0005$) | ER import of proteins destined for export (139) |
| DTC (PF3D7_0823900) | n.s. | n.s. | - 0.34 ($P = 0.004$) | Imports dicarboxylate, exports tricarboxylate from mitochondrion (145) |

1337

**Table 2: Proteins identified by RNA-sequencing that may be involved in iron transport in *P. falciparum*.**

The classification of the proteins identified is indicated according to the Transport Classification Database (71). Data on human orthologs was retrieved using the NCBI position-specific iterated (PSI) BLAST with default settings at https://blast.ncbi.nlm.nih.gov/Blast.cgi (29). DV, digestive vacuole; E, expect value; EM, electron microscopy; hpi, hours post invasion; MIS, mutagenesis index score; PDB, Protein Data Bank; PPM, parasite plasma membrane.

| Gene product, gene ID | Blood stage tran-scription peak (51) | Classifi-cation (71) | Solved structure of a similar protein | Potential human ortholog(s) | Localization in *P. falciparum* | Transport assays and proposed function | Mutability in *P. falciparum* | Essentiality in *P. berghei* |
|---|---|---|---|---|---|---|---|---|
| **MRS3,** PF3D7_0905200 | 20 hpi | 2.A.29.5.9 | None | Mitoferrin-1 (E = 7 x $10^{-10}$, 27.0% identity, 26% coverage); Mitoferrin-2 (E = 6 x $10^{-8}$, 27.9% identity, 12% coverage) | Mitochondrion (this study) | Liposomal transport assays with *S. cerevisiae* MRS3 (39): $Fe^{2+}$ import into mitochondrial matrix across inner membrane | Non-disruptable, MIS: 0.135 (65) | PBANKA_041620: essential at asexual blood stage (92) |
| **VIT,** PF3D7_1223700 | 36 hpi | 2.A.89.1.13 | Crystal structure of VIT1 from *Eucalyptus grandis*: PDB: 6IU9 (70) | None | Cytoplasmic vesicles (this study) | Transport assays using inverted vesicles with *Pf*VIT (32): $Fe^{2+}$ export from cytosol into cytoplasmic vesicles (this study) in exchange for $H^+$ | Disruptable, MIS: 0.903 (65) | PBANKA_143860: dispensable at asexual blood stage with growth rate of 1 (95% CI: 0.94 – 1.05) (92), and at mosquito and liver stage (157); reduced growth in blood and liver in mice (30) |
| **ZIPCO,** PF3D7_1022300 | 32 hpi | 2.A.5.3.12 | Cryo-EM structure of ZIP from *Bordetella bronchiseptica*, PDB: 8GHT (72) | ZIP1 (E = 2 x $10^{-5}$, 21.7% identity, 48% coverage) | Cytoplasmic vesicles (this study) | $Zn^{2+}$ uptake assays with *Bb*ZIP in *Escherichia coli* (72), Liposomal assays with *Pf*ZIP1 (98): $Fe^{2+}/Zn^{2+}$ import into cytosol from cytoplasmic vesicles (this study) | Disruptable, MIS: 1 (65), growth increase at asexual blood stage (this study) | PBANKA_050650: growth rate of 0.86 (95% CI: 0.74 – 0.98) at asexual blood stage (92); dispensable at blood and mosquito stages but reduced sporozoite infectivity and inefficient liver schizogony in mice (34) |
| **NRAMP,** PF3D7_0523800 | 14 hpi | 2.A.55.2.25 | Crystal structure of NRAMP/DMT from *Staphylo-coccus capitis*, PDB: 5M95 (75) | NRAMP1 (E = 2 x $10^{-38}$, 28.4% identity, 56% coverage); NRAMP2 (E = 2 x $10^{-38}$, 27.6% identity, 65% coverage) | DV (36) | Uptake assays with *Deinococcus radiodurans* NRAMP in *E. coli* and proteoliposomes (100, 101): pH-dependent symport of $Mn^{2+}/Fe^{2+}$ with $H^+$ into cytosol from DV | Non-disruptable, MIS: 0.123 (65) | PBANKA_123860: no mutants generated |
| **CRT,** PF3D7_0709000 | 14 hpi | 2.A.7.20.1 | Cryo-EM structure of *Pf*CRT from strain 7G8, PDB: 6UKJ (74) | None | DV (158) | Transport assays using *Xenopus* oocytes with *Pf*CRT (37): symport of $Fe^{2+}$ with $H^+$ into cytosol from DV | Non-disruptable, MIS: 0.127 (65), essential at asexual blood stage | PBANKA_121950: essential at asexual blood stage (92, 159) |

| | | | | | | | |
|---|---|---|---|---|---|---|---|
| | | | | | | | (35) |
| **E140, PF3D7_0104100** | 40 hpi | Unknown | None | None | PPM (this study) | Putative Fe$^{2+}$ importer at PPM (this study) | Non-disruptable, MIS: 0.119 (65) | PBANKA_0209000: growth rate of 0.48 (95% CI: 0.32 – 0.63) at asexual blood stage (92), dispensable in mosquito and liver stages (157) |

**Supplementary Material**

<u>**Supplementary Figures**</u>

**Supplementary Figure S1:** RNA quality.

**Supplementary Figure S2:** Cloning strategy and confirmation of correct DNA integration

into the genome of the cell lines generated.

**Supplementary Figure S3:** Sequence alignments of *P. falciparum* proteins with functionally

characterized homologs.

**Supplementary Figure S4:** Alignments of functional sites in the predicted *P. falciparum*

protein structures with those of functionally characterized homologs.

1351

<u>**Supplementary Videos**</u>

**Supplementary Video S1:** Parasite endogenously expressing *Pf*MRS3-GFP stained with

MitoTracker Red and Hoechst-33342.

**Supplementary Video S2:** Parasite endogenously expressing *Pf*VIT-GFP stained with ER

Tracker Red and Hoechst-33342.

**Supplementary Video S3:** Parasite endogenously expressing *Pf*ZIPCO-GFP stained with

LysoTracker Deep Red and Hoechst-33342.

**Supplementary Video S4:** Parasite expressing *Pf*E140-GFP endogenously, *Pf*ACP(1-60)-

mCherry episomally and stained with Hoechst-33342.

1361

<u>**Supplementary Tables**</u>

**Supplementary Table S1:** Differentially expressed genes of *P. falciparum* 3D7 cultured with

erythrocytes from a donor with higher, control or low iron status.

**Supplementary Table S2:** Differentially expressed genes of *P. falciparum* 3D7 cultured with

erythrocytes from a healthy donor with or without addition of 0.7 µM hepcidin.

**Supplementary Table S3:** Oligonucleotides (A) and plasmids (B) used in this study.

1368

1369 <u>**Supplementary PDB files**</u>

1370 Top-scoring PDB files for the multimers that were computed at EMBL:

1371 **PfVIT.pdb**

1372 **PfZIPCO.pdb**

1373

**Acknowledgements**

The authors thank the Genomics Core Facility at EMBL Heidelberg, especially Vladimir

Benes, for the RNA-sequencing service, and EMBL Hamburg for the provision of research

and technical support as well as access to research infrastructures. Grzegorz Chojnowski

and the group of Jan Kosinski at EMBL Hamburg enabled the AlphaFold2 workflow at the
EMBL Hamburg computer cluster. The Advanced Light and Fluorescence Microscopy
Facility at CSSB Hamburg, in particular Roland Thünauer, supported microscopy
experiments and the Bernhard Nocht Institute for Tropical Medicine (BNITM) provided lab
space. We gratefully acknowledge Tobias Spielmann for pSLI-GFP and pARL-*Pf*ACP(1-60)-
mCherry, Paul Burda for pSLI-GFP-*glmS*, Jacobus Pharmaceuticals for WR99210, Anna
Bachmann and Mayka Sánchez for helpful advice, Eileen Devaney, Katharina Jungnickel,
and Samuel Pažický for critical reading of the manuscript, and Heidrun von Thien, Yannick
Höppner, and Gabriela Guédez for technical assistance.

1387

## Funding

This work was supported by a Boehringer Ingelheim Foundation Exploration Grant, the
Partnership for Innovation, Education and Research (PIER) of Hamburg University and
DESY (project PIF-2018-87) and the European Molecular Biology Laboratory (EMBL). JW
was additionally funded by the European Research Council under the European Union's
Horizon 2020 Research and Innovation Programme (grant agreement 759534) and VK by a
research fellowship from the EMBL Interdisciplinary Postdoc (EIPOD) Programme under
Marie Curie Cofund Actions MSCA-COFUND-FP (grant agreement 847543). The funders
had no role in study design, data collection and analysis, decision to publish, or preparation
of the manuscript.

## Author contributions

JS and JW designed the study; MG and SPe recruited the blood donors; JW and CN
performed the experiments; JW, JS, VK and LVN analyzed the data; VK generated the
structural models, and JW wrote the manuscript with contributions from VK, JS and SPo. All
authors read and approved the submitted version.

**Figure 1: Effects of the iron status of the blood donor and of hepcidin on (A) erythrocyte labile iron levels, (B) *P. falciparum* 3D7 growth rates, (C) DNA content per mature schizont, and (D) the number of merozoites per mature schizont.**

The relative labile iron level and DNA content per cell were assessed in the presence or absence of 0.7 µM hepcidin (Hep) by measuring the mean fluorescence intensity (MFI) of Phen Green SK or SYBR Green I compared to control (Ctrl, untreated, normal hemoglobin level) using flow cytometry. Therefore, 100,000 cells were analyzed per replicate. Parasite growth rates refer to the fold change in parasitemia after one intraerythrocytic developmental cycle in vitro relative to control as determined by flow cytometry with SYBR Green I (104). Mature schizonts were obtained by treating schizonts at 40 hpi with 1 mM compound 2 (4-[7-[(dimethylamino)methyl]-2-(4-fluorphenyl)imidazo[1,2-α]pyridine-3-yl]pyrimidin-2-amine) for 8 h. To count the number of merozoites, Giemsa-stained blood smears were analyzed microscopically and only single-infected cells with one digestive vacuole were considered. Means and 95% confidence intervals (indicated by error bars) are shown. Statistical significance was calculated with two-tailed unpaired t tests with Welch's correction for unequal variances and adjusted with the Holm-Šídák method for multiple comparisons except for merozoite numbers, which were compared using Mann-Whitney test. The number of independent experiments is represented as n and the number of parasites imaged in three independent experiments as N.

**Figure 2: Differential expression of *P. falciparum* 3D7 genes under various iron conditions.**
Parasites were cultured with erythrocytes from an individual with high, medium (healthy) or low iron status (experiment 1) or with red blood cells from another healthy donor in the presence or absence of 0.7 µM hepcidin (experiment 2). Samples were harvested at the ring and trophozoite stage (6 – 9 and 26 – 29 hours post invasion, hpi) with three biological replicates per time point and condition. The maximum likelihood estimation (MLE) of the average developmental age of the parasites for each condition and time point (**A**) was calculated using an algorithm developed by Avi Feller and Jacob Lemieux (50). CI, confidence interval. The volcano plots (**B** and **D**) show transcriptional changes of all parasite genes. Red dots indicate significantly ($P$ < 0.05, exact test for negative binomial distribution) upregulated genes ($\log_2$ (fold change) ≥ 0.2), blue dots stand for significantly downregulated genes ($\log_2$ (fold change) ≤ -0.2), while grey dots represent genes that did not significantly differ in transcription under the conditions described ($P$ ≥ 0.05 and / or -0.2 < $\log_2$ (fold change) < 0.2). Differentially expressed genes encoding putative iron transporters (see Table 1) are labeled. Panels **C** and **E** show the enrichment of Gene Ontology (GO), Kyoto Encyclopedia of Genes and Genomes (KEGG) and Reactome (REAC) terms among significantly regulated genes excluding *var*, *stevor* and *rifin* gene families at the two time points. The functional terms were summarized using REVIGO (122) to remove redundancies, represented by circles and plotted according to the significance of their enrichment (-$\log_{10}$ (adjusted $P$), hypergeometric test). The size of the circle is proportional to the number of differentially regulated genes in the dataset that are associated with the respective term, while the color stands for the fold enrichment. The gray dashed line indicates the threshold of the adjusted $P$ value (-$\log_{10}$ 0.05 = 1.3).

**Figure 3: Subcellular localization of known and putative iron transport proteins.**
Representative erythrocytes infected with *P. falciparum* 3D7 parasites endogenously expressing GFP-tagged
*Pf*MRS3 (**A**), *Pf*VIT (**B**), *Pf*ZIPCO (**C**) or *Pf*E140 (**D**) were additionally stained with the fluorescent dyes
Hoechst-33342, MitoTracker Red, ER Tracker Red and/or LysoTracker Deep Red. Co-transfection with a
construct that encodes the 60 N-terminal amino acids of acyl carrier protein (*Pf*ACP) tagged with mCherry
(125) resulted in red fluorescence of the apicoplast. Live-cell images were taken under physiological
conditions at 37°C using an SP8 confocal laser-scanning microscope (Leica). DIC, differential interference
contrast. Scale bar, 2 µm.

**Figure 4: *Pf*VIT and *Pf*E140 are important for *P. falciparum* growth and may be involved in intracellular iron homeostasis.**

**A** Representative erythrocytes infected with *P. falciparum* 3D7 parasites that endogenously express a truncated version of *Pf*VIT or *Pf*ZIPCO tagged with GFP (green). **B** Growth rates of knockout parasite lines generated. **C** Reduction of *Pf*E140-GFP fluorescence (green) in live 3D7 parasites caused by *glmS*-mediated knockdown that was induced by treatment with 2.5 mM glucosamine (GlcN) for 36 h compared to untreated control (Ctrl). **D** Total parasite fluorescence intensities were quantified as background-corrected integrated densities using ImageJ version 2.9.0/1.53t (110) and compared using Mann-Whitney test. **E** The size of the parasites was measured as the area of the region of interest and compared using equal variance unpaired t test. **F** Conditional knockdown of *Pf*E140 induced by treatment with 2.5 mM GlcN results in a growth defect during asexual blood stage development. Live parasites were stained with Hoechst-33342 (blue) and imaged under physiological conditions at 37°C using an SP8 confocal laser-scanning microscope (Leica). DIC, differential interference contrast. Scale bar, 2 μm. Error bars represent 95% confidence intervals of the mean, N the number of parasites analyzed, n the number of independent experiments and Hep treatment with 0.7 μM hepcidin. Growth rates refer to the fold change in parasitemia after two intraerythrocytic developmental cycles in vitro relative to untreated wild-type 3D7 parasites (WT) as determined by flow cytometry with SYBR Green I (104). Statistical significance of growth differences was calculated with two-tailed unpaired t tests with Welch's correction for unequal variances and adjusted with the Holm-Šídák method for multiple comparisons.

**Figure 5: Structures of known and putative *P. falciparum* iron transporters as viewed from the membrane plane.**
**A** Predicted protein structures with per-residue pLDDT (predicted local distance difference test) confidence scores on a scale from 0 to 100, where blue represents high and red low confidence, respectively. The experimentally determined structure of *Pf*CRT is shown in gray. **B** Molecular lipophilicity potential of the protein surfaces as implemented in UCSF ChimeraX; tan is hydrophobic and cyan hydrophilic. Dashed lines above and below the tan regions of all proteins indicate the respective membrane and disordered loops were removed for clarity. **C** Surface charge of the proteins with positively charged areas colored blue and negatively charged ones red. Putative cation-binding site are indicated with an asterisk and transport directions by arrows. *Pf*E140 likely forms a dimer but is shown as a monomer, as no predicted dimer structure could be obtained using AlphaFold2-multimer. The putative cation-binding sites for this protein are based on DeepFRI gradCAM scores for the functional term GO:0015075 "monoatomic ion transmembrane transporter activity" (Supplementary Fig. S5).

**Figure 6: Iron homeostasis in a *P. falciparum*-infected erythrocyte.**
The human blood plasma contains between 10 and 30 μM total iron and the erythrocyte cytosol approximately 20 mM (19). However, the labile iron pool is only 3 μM in an uninfected erythrocyte and 1.6 μM in a *P. falciparum*-infected one (20). Human ferroportin (FPN) at the host cell surface (erythrocyte plasma membrane, EPM) exports ferrous iron from the erythrocyte (127) and the nutrient pore formed by *Pf*EXP1 and *Pf*EXP2 allows the passage of ions through the parasitophorous vacuole membrane (PVM) (128). *Pf*E140 at the parasite plasma membrane (PPM) may mediate iron uptake into the parasite cytosol and the mitochondrial carrier protein *Pf*MRS3 likely translocates $Fe^{2+}$ into the mitochondrion, a site of *de novo* heme biosynthesis (this study). We propose that the vacuolar iron transporter (*Pf*VIT) is involved in iron detoxification by transporting excess $Fe^{2+}$ from the cytosol into cytoplasmic vesicles that may be acidocalcisomes, whereas *Pf*ZIPCO releases $Fe^{2+}$ from these organelles under low-iron conditions. The digestive vacuole (DV) contains a high amount of total iron as it is the site of hemoglobin degradation and hemozoin formation (21). The chloroquine resistance transporter (*Pf*CRT) and the natural resistance-associated macrophage protein (*Pf*NRAMP, also called *Pf*DMT1 for divalent metal transporter 1) were suggested to mediate proton-coupled export of $Fe^{2+}$ from the DV into the parasite cytosol (37, 73). Both acidocalcisomes and the DV are likely acidified by the plant-like $H^+$-pump V-ATPase, which can fuel secondary active transport processes (22, 27). Parasite-encoded proteins are shown in orange and human-encoded transporters in blue.