SON-dependent nuclear speckle rejuvenation alleviates proteinopathies.

William Dion[1, #], Yuren Tao[2, #], Maci Chambers[1], Shanshan Zhao[2], Riley K. Arbuckle[3,4], Michelle Sun[1], Syeda Kubra[1], Imran Jamal [1], Yuhang Nie[2], Megan Ye[1], Mads B. Larsen[1], Daniel Camarco[1], Eleanor Ickes[1], Claire DuPont[1], Haokun Wang[1], Bingjie Wang[3], Silvia Liu[5,6], Shaohua Pi[3], Bill B Chen[1,7], Yuanyuan Chen [3,8] *, Xu Chen[2] *, and Bokai Zhu[1,5,9] *

[1] Aging Institute of UPMC, University of Pittsburgh School of Medicine, Pittsburgh, PA, U.S.A.

[2] Department of Neuroscience, School of Medicine, University of California, San Diego, CA, U.S.A.

[3] Department of Ophthalmology, University of Pittsburgh School of Medicine, PA, U.S.A.

[4] Department of Human Genetics, University of Pittsburgh Graduate School of Public Health, Pittsburgh, PA, USA.

[5] Pittsburgh Liver Research Center, University of Pittsburgh, Pittsburgh, PA, U.S.A.

[6] Department of Pathology, University of Pittsburgh School of Medicine, Pittsburgh, PA, U.S.A.

[7] Division of Pulmonary, Allergy and Critical Care Medicine, Department of Medicine, University of Pittsburgh School of Medicine, Pittsburgh, PA, U.S.A.

[8] Department of Pharmacology and Chemical Biology, University of Pittsburgh School of Medicine, PA, U.S.A.

[9] Division of Endocrinology and Metabolism, Department of Medicine, University of Pittsburgh School of Medicine, Pittsburgh, PA, U.S.A.

# These authors contribute equally.

Corresponding authors*:

Yuanyuan Chen: cheny1@pitt.edu

Xu Chen: x1chen@health.ucsd.edu

Bokai Zhu: bzhu@pitt.edu

Abstract

Current treatments targeting individual protein quality control have limited efficacy in alleviating proteinopathies, highlighting the prerequisite for a common upstream druggable target capable of global proteostasis modulation. Building on our prior research establishing nuclear speckles as a pivotal membrane-less organelle responsible for global proteostasis transcriptional control, we aim to alleviate proteinopathies through nuclear speckle rejuvenation. We identified pyrvinium pamoate as a small-molecule nuclear speckle rejuvenator that enhances protein quality control while suppressing YAP1 signaling via decreasing the surface/interfacial tension of nuclear speckle condensates through interaction with the intrinsically disordered region of nuclear speckle scaffold protein SON. In pre-clinical models, nanomolar pyrvinium pamoate alleviated retina degeneration and reduced tauopathy by promoting autophagy and ubiquitin-proteasome system in a SON-dependent manner without causing cellular stress. Aberrant nuclear speckle morphology, reduced protein quality control and increased YAP1 activity were also observed in human tauopathies. Our study uncovers novel therapeutic targets for tackling protein misfolding disorders within an expanded proteostasis framework encompassing nuclear speckles and YAP1.

Introduction:

Proteinopathies are diseases associated with the accumulation of misfolded proteins, which often arise from a decline in proteostasis pathways, including the ubiquitin-proteasome system (UPS), the ER-Golgi protein secretory pathways, and autophagy lysosomal pathway (ALP) [1, 2]. However, therapies targeting singular pathways have limited efficacy, indicating an incomplete understanding of disease mechanisms.

We recently discovered that under physiological conditions, the network of proteostasis pathways manifests as cell-autonomous 12-hour (12h) ultradian rhythms, regulated by a dedicated 12h oscillator, independent from the 24h circadian clock and the cell cycle [3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13]. By studying this 12h oscillator, we uncovered an unexpected role of nuclear speckles in global proteostasis control [9]. Nuclear speckles are membrane-less organelles important for mRNA processing and gene regulation [14, 15, 16, 17, 18], and their liquid-liquid phase separation (LLPS) dynamics dictate the global transcriptional capacity of proteostasis genes [9]. Moderate overexpression of the nuclear speckle scaffolding protein SON is sufficient to decrease nuclear speckle sphericity, increase the recruitment of nuclear speckles to chromatin, amplify proteostasis gene expression, and reduce protein aggregation [9]. Conversely, reducing SON level leads to much more spherical and stagnant speckles, sequesters nuclear speckles away from chromatin, blunts proteostasis gene expression and subsequently elevates intracellular protein aggregates [19]. More importantly, SON expression decreases with age in various tissues in mice, concomitant with more spherical and smaller nuclear speckles (**Supplementary Fig. 1a-f**). In addition, reduced SON expression was also observed in aging human lungs, and in brain tissues of human subjects with Alzheimer's disease (**Supplementary Fig. 1g-i**) [9, 20]. Based upon these results, we herein propose that enhancing SON expression or function, either genetically or pharmacologically, under aging and disease conditions, could potentially restore nuclear speckle morphology and function. This, in turn, would bolster the entire protein quality control system, thereby delaying or even reversing the progression of both aging-related and inherited proteinopathies. We introduce this concept as SON-dependent 'nuclear speckle rejuvenation' (**Fig. 1a**).

In this study, we initially characterized the comprehensive transcriptome changes resulting
from SON-mediated nuclear speckle rejuvenation and unexpectedly discovered a broader
proteostasis framework that incorporates SON-mediated nuclear speckles condensation,
unfolded protein response (UPR) transcription factors (TF)-mediated proteostasis gene activation
and repression of YAP1 transcriptional activity. Via a high-throughput drug screen, we identified
pyrvinium pamoate (PP) as a small molecule nuclear speckle rejuvenator that recapitulates the
entire transcriptome changes elicited by SON overexpression. Mechanistically, via a cell-free
nuclear speckle reconstitution system, we demonstrated that PP exerts transcriptional
reprogramming via reducing the surface/interfacial tension of nuclear speckle condensates and
promoting their wetting of genomic DNA via targeting the intrinsically disordered region (IDR) of
SON. In preclinical models, PP exhibited strong efficacy in protecting against both tauopathy and
retinal degeneration at nanomolar concentration without inducing cellular stress. Lastly, we
showed that both the reduction of protein quality control gene expression and increase of YAP1
transcriptional activity is associated with retinal degeneration in mice and tauopathy in humans.

Result:

Genetic rejuvenation of nuclear speckles transcriptionally reprograms global proteostasis
and YAP1 activity in an opposing manner.

To determine the extent by which SON transcriptionally reprograms gene expression under both
basal and proteotoxic stress conditions, we performed bulk mRNA-Seq on immortalized mouse
embryonic fibroblasts (MEFs) with either SON knockdown (KD) by siRNA or overexpression (OE)
via CRISPRa[21], in the absence or presence of the ER stress inducer tunicamycin (Tu) as
previously described (**Supplementary Table1**) [9]. Principal component analysis (PCA) on total
mRNA level indicated that while SON manipulation has little effects on global gene expression
under basal condition, SON OE and KD significantly amplified and dampened the global
transcriptional response to ER stress, respectively (**Fig. 1b**). These include 461 genes that are
normally induced, and 901 genes repressed by Tu under normal SON expression condition (**Fig.**
1c-e, Supplementary Fig. 2a). For both groups of genes, we further observed a strong
correlation between the relative fold induction or repression for each gene under SON OE and
KD conditions (**Supplementary Fig. 2b**), further demonstrating the robustness of bidirectional
control on proteostasis gene expression by SON.

As expected, gene ontology (GO) analysis revealed that those ER stress-induced 461 genes
are strongly enriched in protein quality control pathways, including protein folding (such as *Pdia3*,
Dnajb11 and *Manf*), ER/Golgi quality control (such as *Sec23b*, *Hyou1* and *Hspa5*), tRNA
aminoacylation (such as *Gars*, *Iars* and *Eprs*), ER-associated protein degradation (ERAD) (such
as *Edem1*, *Syvn1*, *Sel1l* and *Ube2g2*), and autophagy (such as *Sqstm1* and *Atg13*) (**Fig. 1f,**
Supplementary Fig. 2a). To shed light on the mechanisms by which SON transcriptionally
amplifies gene activation in response to ER stress, we performed both Landscape In Silico
deletion Analysis (LISA) [22] and motif analysis to infer the transcriptional regulators that may
mediate nuclear speckles interactions with chromatin. Both analyses revealed basic leucine
zipper (bZIP) transcription factors (TFs), including ATF6, XBP1, ATF4 and CREB1 as the top
candidates (**Fig. 1g and Supplementary Fig. 2c**). ChIP-qPCR further showed increased
recruitment of nuclear speckles to the 3' regions of selective proteostasis genes in response to
SON OE under both basal and ER stress conditions, concomitant with increased recruitment of
XBP1s to the promoter regions of the same genes (**Supplementary Fig. 2d, e**). To corroborate
our *in vitro* findings, we further examined a recently published murine *in vivo* hepatic XBP1s

interactome dataset [10], and found proteins involved in mRNA splicing and processing are very
strongly enriched in the XBP1s interactome at CT8, a time when hepatic SON expression peaks
[9], with PRPF8, SNRNP200 and DHX9 being the top three most abundant proteins detected in the
entire XBP1s interactome at CT8 (**Fig. 1h, i**). By contrast, at CT0 when SON expression is the
lowest, the amount of splicing proteins that interact with XBP1s is markedly reduced (**Fig. 1i**). The
observed decreased recruitment of splicing proteins to XBP1s at CT0 is not due to reduced XBP1s
level itself, as the hepatic XBP1s expression at CT0 is in fact higher compared to CT8 [3, 23]. These
results thus reinforce the notion that nuclear speckles rejuvenation by SON OE is sufficient to
amplify the global proteostasis transcriptional activation, likely via facilitating physical interactions
between nuclear speckles and UPR TF like XBP1s.

Compared to induced genes, much less is known about the gene programs that are repressed
under ER stress. GO analysis indicate that those 901 ER stress-repressed genes are strongly
enriched in Hippo-YAP1 signaling that regulates the diverse biological processes of angiogenesis,
axon guidance, epithelial to mesenchymal transition (EMT), wound healing, cell adhesion, cell
migration, and extra cellular matrix organization, with examples of canonical YAP1 target genes
Bmp4, *Tuba1a*, *Fzd2*, *Tgfb3* and *Yap1* itself and its transcriptional partner *Tead2* [24, 25, 26] (**Fig. 1c-**
f and Supplementary Fig. 2a). YAP1 and TEAD2 were further predicted to be transcriptional
regulators of these 901 ER stress-repressed genes via both LISA and motif analysis (**Fig. 1j and**
Supplementary Fig. 2c) and the nuclear YAP1 level was significantly reduced in response to
either Tu or SON OE and further decreased upon the combination of the two (**Fig. 1k, l**). We
further performed TEAD luciferase reporter assay and found that both SON OE and Tu
significantly reduced the TEAD response element-driven luciferase activity in MEFs, with the
lowest observed in SON OE cells under ER stress (**Supplementary Fig. 3a**). Scratch assays
further confirmed that both SON OE and Tu significantly reduced cell migration in MEFs, with the
lowest observed in SON OE cells under ER stress (**Fig. 1m, n**). To rule out the possibility that the
global repression of YAP1 transcriptional output during ER stress is specific to MEFs or Tu, we
further analyzed a recent transcriptome dataset in the human astrocytoma-derived LN-308 cell
line in response to both Tu and thapsigargin (Thap) (another ER stress inducer) treatments [27],
and observed a strong downregulation of genes involved in YAP1 signaling under both treatments
that progressed with time (**Supplementary Fig. 3b-d**). Together, our data indicates that the
downregulation of YAP1 transcriptional output is an integral component of the global
transcriptional response to proteotoxic stress, and it is also under nuclear speckles control.

Given the established roles of nuclear speckles in mRNA processing [28, 29, 30], we next
investigated whether SON also regulates mRNA splicing dynamics. While SON manipulation has
no effects on the overall transcriptional state of mature mRNA under basal DMSO condition
(**Supplementary Fig. 4a bottom**) (consistent with total mRNA shown in **Fig. 1a**), it exerted
profound effects on the pre-mRNA level (**Supplementary Fig. 4a top**), indicating a change in
splicing dynamics. By either estimating the relative splicing rates among different groups under
basal condition using a simple first-order kinetic model of transcription (see Materials and
Methods) or quantifying global intron retention events using the iRead algorithm [31], we found that
SON increases the splicing rates and improves the splicing fidelity of genes involved in
proteostasis and RNA metabolism, and negatively regulates those involved in YAP1-related
processes of cell migration, axon guidance, cell adhesion and EMT (**Supplementary Figs. 4b-e,**
5a-d and 6a-d). The significant enrichment in mRNA processing genes themselves under SON
control is consistent with known potent autoregulation of splicing factors [32, 33]. Furthermore, we
also found that SON can activate and repress the mature mRNA expression of a select set of

proteostasis (albeit very modestly) and YAP1 target genes, respectively, under basal DMSO
conditions (**Supplementary Fig. 7a-c**). Finally, we observed that SON can also activate an anti-
viral response gene signature under basal DMSO condition (**Supplementary Fig. 7b**), suggesting
a potential broader implication of nuclear speckle rejuvenation in boosting innate immunity.
Collectively, our results demonstrated that genetically rejuvenating nuclear speckles reprograms
global proteostasis and YAP1 transcriptional output in an opposing manner, under both basal and
proteotoxic stress conditions.

Our results thus far demonstrated a tripartite network where nuclear speckle rejuvenation by
SON boosts proteostasis and suppresses YAP1. Two possible topologies of the network exist. In
model one, nuclear speckles can signal both proteostasis and YAP1 signaling directly
(**Supplementary Fig. 8a, model 1**), while in model 2, nuclear speckles repress YAP1
downstream of increased proteostasis gene program (**Supplementary Fig. 8a, model 2**). To
distinguish between the two models, we examined a recently published RNA-seq dataset in
HEK293T cells treated with DMSO, Thap or a very specific XBP1s small molecule activator IXA4
[34, 35, 36]. While IXA4 can induce a robust proteostasis gene signature similar to that of Thap, it
failed to repress YAP1 transcriptional output genes as Thap did (**Supplementary Fig. 8b-d**).
Collectively these results support the first model where nuclear speckles can program
proteostasis gene expression and YAP1 transcriptional output in parallel, likely via promoting
physical interaction between nuclear speckles and XBP1s for the former, and triggering YAP1
nuclear exclusion for the latter (**Supplementary Fig. 8e**). We speculate the opposing changes in
proteostasis and YAP1 signaling may reflect an energetic trade-off between proteostasis and the
control of cell dynamics under proteotoxic stress (**Supplementary Fig. 8f**).

High-throughput screen (HTS) identified pyrvinium pamoate (PP) as a small-molecule
rejuvenator of the nuclear speckle.

Having established the proof-of-principle of nuclear rejuvenation via SON OE, we next explore
the feasibility of rejuvenating nuclear speckles pharmacologically, for two reasons. First,
compared to the challenging implementation of SON-based gene therapy, which faces obstacles
due to the considerable size (~7kb) of SON's open reading frame, using small molecules to boost
nuclear speckle activity may offer superior therapeutic potential. Secondly, we wanted to utilize
an orthogonal approach to further demonstrate the opposing transcriptional changes of
proteostasis and YAP1 signaling following nuclear speckle rejuvenation. Since SON OE and KD
reduced and increased the sphericity of nuclear speckles [19], respectively, putative nuclear speckle
rejuvenators are expected to reduce the sphericity (and increase the diffuseness or irregularity)
of speckles.

We started with a library of over 2500 FDA-approved drugs and ran a primary HTS on our
previously described EGFP::SC35 (SRSF2) MEFs (EGFP was knocked in to the N-terminus of
endogenous *Srsf2* locus, which is a well-established marker for nuclear speckles [9]) to identify
compounds that could alter nuclear speckles sphericity (**Fig. 2a**). As a quality control, our primary
screen successfully identified four histone deacetylase inhibitors that produced much more
spherical nuclear speckles approaching and/or exceeding r=0.9 (**Supplementary Fig. 9a, b**)**,** in
alignment with a previous study [37]. In the end, we identified five compounds - the tyrosine kinase
inhibitors nintedanib (NB) and ponatinib (PB), the anti-microbial proflavine hemisulfate (PH) and
proflavine, and the anthelmintic pyrvinium pamoate (PP) – having the ability to both decrease
nuclear speckles sphericity and amplify *Perk*-promoter driven dGFP expression (*Perk* is a UPR
target and exhibits 12h rhythms of expression [3]) in a dose-dependent manner (**Fig. 2b, and**

Supplementary Fig. 9c-e). For PP, a clear dose response in reducing the nuclear speckle
sphericity was observed between 0 and 0.3µM, and further increasing its concentration failed to
further decrease the sphericity (**Fig. 2b and Supplementary Fig. 9c**). PP also increased the
perimeter of nuclear speckles (**Fig. 2c**), indicating that PP can increase the surface area of
nuclear speckles in the three-dimensional space of the nucleus.

To determine which of these drugs are *bona fide* nuclear rejuvenators, we performed mRNA-
Seq on MEFs treated with NB, PB, PH or PP for 24 hours and compared the transcriptome
signature of each drug (**Supplementary Fig. 10a,** and **Supplementary Table2**) with those of
SON OE and KD under both DMSO and Tu conditions. Gene set enrichment analysis (GSEA) [38]
using either those 461 Tu-induced or 901 Tu-repressed genes (depicted in **Fig. 1e**) revealed PP
as the only drug triggering a transcriptional response with strong resemblance to SON OE cells
in response to ER stress (**Fig. 2d, e and Supplementary Fig. 10b**). These included both
upregulated genes implicated in protein quality control and downregulated genes involved in the
regulation of cell dynamics (**Fig. 2f, g and Supplementary Fig. 10c, d**). LISA analysis on
differentially expressed genes by PP revealed bZIP TFs ATF4 and YAP1 among top
transcriptional regulators of upregulated and downregulated genes, respectively (**Supplementary
Fig. 10e**). 1µm PP increased the expression of UPR and integrated stress response (ISR) TFs –
XBP1s and ATF4 - at both the mRNA and protein level (**Supplementary Fig. 10f, g)** and induced
nuclear exclusion of YAP1 (**Supplementary Fig. 10h)**, while not affecting SON level
(**Supplementary Fig. 10i**).

Our analysis so far suggested the PP induced a transcriptional signature that resembles a
mixture of responding to SON OE and Tu treatment. We performed additional comparative
transcriptome analysis to further validate this conclusion. First, when comparing the fold induction
or repression of gene expression by PP and Tu, the signature of PP is more similar to that of Tu
under SON OE compared to under SON KD condition (p= 0.00195 by Chow tests)
(**Supplementary Fig. 11a**). Secondly, similar to SON OE (**Supplementary Fig. 7b**), PP also
induced expression of genes involved in anti-viral response (**Supplementary Fig. 11b-f**), distinct
from those induced under ER stress. Thirdly, GSEA indicated a strong resemblance of gene
signatures repressed by PP and SON OE that are enriched in the control of cell dynamics, under
basal conditions in the absence of ER stress (**Supplementary Fig. 12a-e**). Lastly, like SON
(**Supplementary Figs. 5d and 6d**), PP also improves the splicing fidelity of splicing genes
themselves (**Supplementary Fig 13a-f**), again reflecting autoregulation of splicing factors. Taken
together, these results indicate that PP is a bona fide nuclear speckle rejuvenator that at 1µM
induces a transcriptional signature in MEFs highly similar to that of SON OE cells under both
basal and proteotoxic stress conditions.

PP reduces the surface tension of nuclear speckle condensates via targeting SON IDR.

To confirm that PP rejuvenates nuclear speckles in a SON-dependent manner, we knocked down
Son via siRNA in MEFs. Son knockdown leads to smaller and more spherical speckles, consistent
with our previous study [19] (**Fig. 2h**). Importantly, PP's ability to reduce nuclear speckle sphericity
is abolished in *Son* knockdown MEFs (**Fig. 2h**)**.** Subsequently, *Son* knockdown impaired PP's
ability to both activate protein quality control gene expression and repress YAP1 transcriptional
output (**Fig. 2i**). To determine whether PP can physically interact with SON in MEFs, we
performed cellular thermal shift assay (CETSA), which is based on ligand-induced thermal
stabilization of target proteins, whereas unbound proteins denature, aggregate and precipitate at
elevated temperatures, ligand-bound proteins remain soluble due to increased stability [39, 40]. Using

a SON-specific antibody (**Supplementary Fig. 14a**), we found that PP induced a thermal shift of
SON with a direction consistent with stabilization (**Fig. 3a**). As negative controls, we found that
PP does not stabilize SRSF2 (SC35), another nuclear speckle protein, or the ISR/UPR TF ATF4,
whose expression is nonetheless significantly increased by PP (**Fig. 3b and Supplementary Fig.**
14b).

SON is the central scaffold protein of nuclear speckles [41, 42, 43], and its ~12h rhythmic
concentration fluctuation drives ~12h rhythmic nuclear speckles LLPS dynamics and chromatin
binding alternating between either a diffuse and chromatin-associated state or a punctate and
chromatin-dissociated state [9]. By contrast, SRSF2(SC35) is one of the critical subunits of the
spliceosomes, which have a broader spatial distribution also occupying the periphery of nuclear
speckles, particularly at the interface between nuclear speckles and the nucleoplasm or chromatin
and is not essential for speckle formation [44, 45]. In MEFs, PP generated a more diffuse and irregular
(less spherical) nuclear speckles with larger surface area (**Fig. 2b, c**), suggesting that PP could
influence nuclear speckle condensates material properties, likely via reducing the surface tension
of speckles (surface tension is the tendency of liquid droplets to minimize the total surface area,
therefore an increased surface area is suggestive of reduced surface tension[46]). Given the
CETSA data indicating PP can bind to SON directly, we next tested whether PP can directly
impact the condensates formation of two nuclear speckle protein, SRSF2 and SON, using an *in*
vitro droplet formation assay [47]. Using different computational algorithms to search for intrinsically
disordered region (IDR) [48, 49, 50], we identified two IDRs at the N and C terminals of mouse SON,
and long stretches of IDRs spanning two-thirds of mouse SRSF2 (**Fig. 3c and Supplementary**
Fig. 14c). We separately cloned the regions encoding both SON IDRs and the full-length SRSF2
into the C-terminal of mCherry and purified recombinant proteins from *E. coli* (**Supplementary**
Fig. 14d). Purified recombinant proteins were added to buffers containing 10% crowding reagents
PEG-8000 as previously described [47]. Confocal fluorescence microscopy of the different protein
solutions revealed mCherry positive, micron-sized spherical droplets freely moving in solution and
wetting the surface of the glass coverslip (**Supplementary Movies 1-3**). All proteins droplets were
highly spherical, exhibited fusion/coalescence behaviors (**Supplementary Movies 1-3**), and
scaled in size and number positively with increasing concentration of proteins and negatively with
increasing salt concentration (**Supplementary Fig. 14e, f**), all properties expected for liquid-like
droplets [47].

Due to surface tension, small droplets will eventually morph into a fewer number of large
droplets, resulting in a net decrease of surface area, either via coalescence or Ostwald ripening
[51] (**Fig. 3d**), which was seen for all protein droplets after 20 minutes of time lapse imaging (**Fig.**
3e, f, and Supplementary Fig. 14g). Addition of PP to SON IDR2, but not SON IDR1 and SRSF2
protein solutions, significantly reduced the kinetics of this process in a dose-dependent manner,
resulting in a negligible decrease of relative surface area after 20 minutes (**Fig. 3d-f, and**
Supplementary Fig. 14g, h). The significance of SON IDR2 is reinforced by the substantial
evolutionary conservation of its sequences across seven distinct species spanning wide
phylogenetic distances from zebra fish to humans (**Supplementary Fig. 14i**).

To better recapitulate the complex compositions of nuclear speckles in the cell-free system,
we further supplemented recombinant mCherry-SON IDR2 with nuclear extract (NE) from Hela
cells that include all active components of transcription and splicing factors (**Supplementary Fig.**
15a) [47]. Mass spectrometry confirmed that Hela NE-supplemented mCherry-SON IDR2
condensates preferentially compartmentalized splicing factors (including SRSF2), with twelve of

the top fifteen enriched proteins previously identified in nuclear speckles [52, 53] (**Supplementary Fig. 15b, c**). By contrast, other nuclear proteins like proteasome subunits, DNA repair factors or general transcription factors were not enriched in the reconstituted condensates (**Supplementary Fig. 15b, c**). These condensates further exhibited less spherical morphology, had increased number and total size (**Supplementary Fig. 15d, e**), features expected from nuclear speckle-like condensates with heterogeneous viscoelastic properties [54]. Importantly, the addition of PP reduced both the sphericity and surface tension of Hela NE-supplemented SON IDR2 condensates in a dose-dependent manner (**Fig. 3g-i**), consistent with the observed effects of PP on nuclear speckle morphology in cells. Together, these data suggest that PP can reduce the surface tension to stabilize both homotypic and heterotypical NE-supplemented SON condensates in a cell-free system, via interacting with SON C-terminal IDR2.

To investigate whether PP may affect the relative spatial distribution of SON and SRSF2 within nuclear speckles, we further supplemented recombinant mCherry-SON IDR2 with NE from GFP::SRSF2-expressing MEFs (**Fig. 3j**). The resulting nuclear speckle-like condensates recapitulated the anticipated spatial distribution of SON and SRSF2 proteins, with the former located at the center, and the latter exhibiting a broader distribution with its highest concentration often observed at 350nm away from the SON IDR2 center (**Supplementary Fig. 15f and Supplementary Movie 4**). While PP does not alter the relative spatial distribution of SON IDR2 and SRSF2, it reduced their sphericity, and markedly increased SRSF2 content at the periphery of nuclear speckle-like condensates (**Fig. 3k-m**). To determine whether PP may also influence the wetting of genomic DNA by nuclear speckles [55], we further added mouse genomic DNA into the droplet solution. In accordance with observations in intact cells, nuclear speckle condensates largely don't mix with but can wet the DNA (**Supplementary Fig. 15g**). Interestingly, the addition of PP more than doubled the wetting of genomic DNA by the reconstituted nuclear speckles (**Fig. 3n, o**).

To validate that PP can alter the nuclear speckle LLPS properties *in vitro* in the context of intact cells, we further performed 1,6 hexanediol sensitivity assay [56] using the same EGFP::SRSF2 MEFs [9]. A short term 1,6 hexanediol treatment preferentially dissolves liquid but not solid condensates, thus a change in the sensitivity to 1,6 hexanediol reflects an alteration in the LLPS property of a given condensates [56]. As demonstrated in **Supplementary Fig. 16a, b**, PP desensitized SRSF2 to the increasing concentrations of 1,6 hexanediol. This effect was not observed on two other biomolecular condensates, the nuclear MED1 [47] and cytosolic GW182 present in P-bodies [57] (**Supplementary Fig. 16c, d**). Taken together, these results suggest a mechanism where PP can reduce the surface/interfacial tension of nuclear speckles via targeting SON IDR, leading to larger surface areas with increased SRSF2 content at the periphery, increased wetting of genomic DNA and subsequently a higher portion of spliceosomes stably engaging in active RNA processing and transcription elongation of proteostasis genes. Since no active transcription occurs in the *in vitro* droplet formation assay, these results demonstrate that PP-mediated nuclear speckle LLPS change is a cause, rather than a consequence of or response to, global transcriptional reprogramming.

PP reduces both pathological Tau and rhodopsin levels by boosting ALP and UPS, at the expense of YAP1 signaling.

Since PP treatment leads to a global increase of protein quality control gene expression, we went on to determine the effects of PP on global protein synthesis, and degradation via UPS and ALP in MEFs under two different concentrations: 0.1 and 1µM. Using puromycin incorporation assay

[58], we found that 0.1 µM of PP did not alter global protein synthesis (**Fig. 4a, b**). The lack of effects
of 0.1 µM of PP on global translation was further confirmed by the lack of changes of p-eIF2α and
ATF4 level (**Supplementary Fig. 17a, b**). To quantify the UPS activity, we treated MEFs with PP
alone or in combination with the proteasome inhibitor MG132 and blotted for high molecular
weight poly-ubiquitinated proteins. 0.1 µM PP led to a significant reduction of poly-ubiquitinated
protein levels (**Fig. 4c, d**), and co-treatment with MG132 restored the level of poly-ubiquitinated
proteins to a level similar to that of MG132 treatment alone (**Fig. 4c, d**). We further directly
measured the activity of 20S proteasome core and found 0.1 µM PP significantly increased the
20S proteasome activity in a SON-dependent manner (**Supplementary Fig. 17c**). These data
collectively indicated that 0.1 µM PP increases UPS-mediated degradation of poly-ubiquitinated
protein. To quantify the autophagic flux, we utilized a tandem LC3 reporter mCherry-GFP-LC3
where an increase in the number of red-fluorescent cytosolic puncta indicates increased
autolysosome formation and autophagic flux (**Fig. 4e**) [59]. 0.1 µM PP markedly increased the
formation of autolysosomes (**Fig. 4f**). To verify this result, we further blocked autophagy at the
late stage autophagosome-lysosome fusion step using Bafilomycin A1 (BafA) [60], and quantified
the level of LC3I and LC3II with or without PP. 0.1 µM PP treatment resulted in an increased ratio
of LC3II/I and reduced level of LC3II, both of which were significantly increased by BafA co-
treatment to a level similar or higher than what was observed under DMSO condition
(**Supplementary Fig. 17d,e**). These results collectively indicate that 0.1 µM PP also augments
autophagic flux. Compared to 0.1µM PP, 1µM PP leads to a global translation repression
(**Supplementary Fig. 18a, b**) concomitant with ATF4 induction (**Supplementary Fig. 10g**), but
is still able to promote both UPS and ALP (**Supplementary Fig. 18c-f**). In sum, these findings
demonstrate that nanomolar concentrations of PP effectively rejuvenate nuclear speckles,
enhancing both the UPS and ALP without inducing cellular stress. However, at higher micromolar
concentrations, PP not only acts as a nuclear speckle rejuvenator but also induces cellular stress,
likely due to its known inhibitory effects on mitochondrial activity[61, 62].

Decline of both ALP and UPS are associated with proteinopathies such as Alzheimer's
disease (AD), frontotemporal dementia (FTD), Parkinson's disease (PD) and a subtype of Retinitis
pigmentosa (RP) with *RHODOPSIN* (*RHO*) mutations [63]. To determine whether PP can protect
against proteinopathies via boosting UPS and ALP, we focused on two different diseases, a
genetic form of RP with a mutant RHO, and tauopathy common in both AD and FTD. RP causes
blindness via the primary loss of rod cells and secondary loss of cone cells. Proline to histidine at
codon 23 (P23H) is the most common mutation in RHO protein, resulting in autosomal dominant
RP in humans [64]. The heterozygous $Rho^{P23H/+}$ knock-in mouse develops progressive retinal
degeneration that resembles the RP phenotype in patients [65, 66]. Recently, several studies
suggested that boosting ERAD can protect against mouse models of RP by increasing elimination
of the mutant RHO^{P23H} protein [65, 67, 68, 69]. To determine if PP can also reduce RHO^{P23H} level, we
used a NIH3T3 cell line ectopically expressing RHO^{P23H} protein [70]. Treating this cell line with 0.1
µM of PP for 24 hours led to a reduction of both monomer and dimer forms of RHO^{P23H} protein in
a SON-dependent manner (**Fig. 4g, h, and Supplementary Fig. 19a, b**). Blocking autophagy
with ULK1/2 inhibitor SBI-0206965 or BafA, and UPS with MG132, respectively, abolished the
effect of PP on reducing RHO^{P23H} protein level (**Fig. 4i, j, and Supplementary Fig. 19c-f**),
suggesting that both increased ALP and UPS are responsible for the increased elimination of
RHO^{P23H} by PP.

Both UPS and ALP are also involved in the degradation of tau protein in tauopathies [71, 72, 73,
74]. To test the effect of PP on tau proteostasis in mouse primary neuronal cultures, we

overexpressed human Tau carrying P301S mutation - an FTD-causing mutation in the human
MAPT gene (Tau) [75]. After treatment with increasing concentrations of PP for 24 hours, a decline
in both total and phosphorylated Tau (Ser396/404) was observed in a dose-dependent manner,
with approximately 50% and 65% reduction in total and p-Tau, respectively, detected at 100nM
PP (**Fig. 4k**). It is noteworthy that these reductions in Tau level occurred without any observable
signs of cellular toxicity (**Supplementary Fig. 19g**). PP also promoted autophagic flux in neurons,
as demonstrated by an increase of LC3 II/I ratio, decreased level of p62/SQSTM1 and increased
autolysosome formation (**Fig. 4k and Supplementary Fig. 19h, i**) [76]. Blocking autophagic flux
with BafA dampened the effects of PP on reducing cellular Tau level (**Fig. 4l, m**). As observed in
fibroblasts, PP also promoted UPS activity in neurons; however, inhibiting proteasome activity
with MG132 has minimal effects on PP's ability to reduce Tau level (**Supplementary Fig. 19j, k**).
Consistent with PP's ability to boost UPS and ALP, PP increased the expression of genes involved
in both pathways in P301S hTau-expressing neurons (**Fig. 4n**). Collectively, these results indicate
that increased autophagy flux largely underlies PP's effect in reducing Tau burden in mouse
primary neurons.

Nuclear speckle rejuvenation by SON OE or PP increases global protein quality control at the
cost of reduced YAP1 signaling in both MEFs (**Fig. 2i**) and neurons (**Fig. 4n**), raising an
interesting question of whether the downregulation of YAP1 signaling also contributed to PP's
efficacy in alleviating proteinopathy. To address this question, we restored YAP1 signaling with
previously published YAP1 activators XMU-MP-1 and/or TRULI [77, 78,79]. We found that while XMU-
MP-1 antagonized the downregulation of YAP1 target genes by PP as expected, it also potently
dampened the upregulation of proteostasis genes (**Fig. 4o**). In addition, both XMU-MP-1 and
TRULI negated PP's effect on reducing RHO[P23H] level in NIH3T3 cells (**Supplementary Fig. 20a-**
d). In addition, knocking down MST1, a kinase that inhibits YAP1 activity [80], similarly blocked the
ability of PP to reduce RHO[P23H] level (**Supplementary Fig. 20e, f**). Similarly, TRULI also blocked
PP's effect on reducing Tau level in primary neurons (**Supplementary Fig. 20g, h**). These results
suggest that for nuclear speckle rejuvenation to achieve the maximum effectiveness in alleviating
proteinopathy, it is essential to simultaneously uphold heightened protein quality control and
reduced YAP1 activity.

PP protects against mouse retina degeneration *ex vivo* and alleviates tauopathy in
***Drosophila*.**

Next, we assessed the effectiveness of PP in ameliorating proteinopathies by utilizing animal
models of RP and tauopathy. To determine whether PP has the potential to restore gene
expression changes in the retina of *Rho^{P23H/+}* mice, we performed RNA-seq in the retina of one
and three months old wild-type and *Rho^{P23H/+}* mice and compared the gene signatures of *Rho^{P23H/+}*
retina with that of PP (**Fig. 5a-e**). In the retina of one month-old *Rho^{P23H/+}* mice, we observed a
significant downregulation of protein transport and autophagy gene expression that showed large
convergence with those upregulated by PP (**Fig. 5a, b**). This includes *Reep6* gene, which
regulates protein trafficking in the ER and its loss-of-function mutation causes autosomal-
recessive RP in both humans and mice (**Fig. 5e**) [81]. By three months, the downregulation of
proteostasis gene expression persists in *Rho^{P23H/+}* mice retina, concomitant with a significant
increase of YAP1-mediated cell dynamics gene expression that also overlaps with PP-
downregulated genes (**Fig. 5c-e**). To directly test the efficacy of PP in protecting against RP, we
treated retina explants isolated from *Rho^{P23H/+}* mice with nanomolar range of PP for 10 days, and
visible light optical coherence tomography (vis-OCT) imaging [82] revealed a remarkable efficacy

of PP in safeguarding the mouse $Rho^{P23H/+}$ retina explants from degeneration. Notably, the cell
counts in the outer nuclear layer closely resembled that of the WT retina explant control (**Fig. 5f-**
j), indicating the protective potential of PP against degenerative processes. Moreover, no
noticeable indications of toxicity were observed throughout the entire duration of the experiment,
further highlighting the safety of PP as a promising therapeutic candidate.

Pan-neuronal expression of wildtype human *MAPT* gene in *Drosophila* recapitulates essential
features of tauopathies, including hyperphosphorylated and misfolded tau, age-dependent neuron
loss, and reduced life span [83], and SON IDR2 sequence is also conserved in flies
(**Supplementary Fig 14i**). Thus, we next tested the efficacy of PP in ameliorating tauopathy in
male flies that express 2N4R isoform of human Tau (*MAPT*) pan-neuronally [*elav*^{c155}-Gal4: UAS-
hTau1.13 (C155>UAS-hTau1.13)] as well as in control *elav* ^{c155}-Gal4 (C155) flies [84]. Both
C155>UAS-hTau1.13 and C155 flies were fed with either standard diet or diet supplemented with
25μM PP, which did not affect the normal development and growth of flies despite its effect in
attenuating WNT and YAP1 signaling [85]. We quantified disease progression at different stages of
fly development with both larval crawling and adult fly climbing assay at 14 and 21 days of age.
PP feeding preserved motor function in C155>UAS-hTau1.13 third instar larvae and adult flies,
with their locomotor performance restored to a level similar to or even slightly higher than control
C155 flies fed with a standard diet (**Fig. 5k, l**). Notably, PP also enhanced the motor function of
adult wild-type control (C155) flies at 21 days of age (**Fig. 5l**). This improvement is likely linked to
PP's ability to promote overall proteostasis, particularly protein turnover rates, a process known
to prolong health and lifespan in flies [86]. PP further extended the median lifespan of C155>UAS-
hTau1.13 flies by 16% from 51 to 59 days (**Fig. 5m**). Consistent with the overall phenotypes, PP
significantly reduced the level of p-Tau and to a lesser extent total Tau in the brains of 21 days-
old C155>UAS-hTau1.13 flies (**Fig. 5n**). Since p-Tau are prone to misfolding and aggregation, it
supports the notion that PP is increasing the overall capacity of protein quality control to remove
misfolded and aggregated proteins, while having negligible effects on normal protein functions.

PP has the potential for treating tauopathy in humans.

To determine the potential of PP for treating tauopathy in humans, we studied whether gene
expression signatures that are opposite of PP can be observed in brain regions of human subjects
with AD. We initially performed a post-hoc analysis of a total of nineteen bulk RNA-seq datasets
encompassing hippocampus, entorhinal cortex, temporal cortex and frontal cortex regions in
control and AD human subjects [87], and found that genes repressed by PP have increased
expression in all four brain regions of human subjects with AD (such as *YAP1, TEAD1 and AMOT*)
(**Supplementary Figs. 21 and 22a**). By contrast, genes that were upregulated by PP displayed
significantly decreased expression in temporal cortex (such as genes involved in ERAD: *EDEM1,*
SEL1L, autophagy: *ATG13*, protein folding: *HYOU1*, and tRNA aminoacylation: *GARS, IARS*)
(**Supplementary Figs. 21 and 22a**). To validate these findings, we further analyzed an
independent single-nucleus RNA-seq (snRNA-Seq) dataset in the prefrontal cortex regions of
human individuals with varying degrees of AD pathology (**Supplementary Fig. 22b**) [88]. We found
that proteostasis genes upregulated by PP are consistently downregulated in all cell types with
strong prominence in neurons and oligodendrocytes in both early and late-stage human AD
subjects (**Supplementary Fig. 22c, d**). Genes that are downregulated by PP (those enriched in
regulation of cell dynamics by YAP1) are initially downregulated in all cell types in the early stage
but significantly upregulated during the late stage of AD in all cell types but inhibitory neurons
(**Supplementary Fig. 22e, f**). This early to late AD progression is concomitant with strong

increase of tauopathy, but not the amyloid burden in these individuals (**Supplementary Fig. 22b**).
Consistent with *in vivo* data, we also observed a significant decrease of proteostasis gene
expression as well as an increase of YAP1-TEAD2 target gene expression in human induced
pluripotent stem cells (iPSC)-derived neurons that express the P301S 4R-Tau when compared to
wild-type 4R-Tau control cells (**Fig. 6a-c**) [74]. These upregulated YAP1-TEAD2 target genes also
overlap with those repressed by PP (**Fig. 6b**).

Finally, to directly test the efficacy of PP in reducing tauopathy in human, we utilized human
iPSC-neurons harboring homozygous FTD-causing MAPT V337M mutation (herein referred to as
V337M) and isogenic wild-type control cells [89] and treated both cell lines with nanomolar range of
PP for 12~24 hours. After 24 hours of treatment with 500 nM PP, iPSC neurons exhibited no signs
of toxicity, as confirmed by the LDH release assay (**Supplementary Fig. 23a, b**). Moreover, the
nanomolar PP treatment did not trigger cellular stress, as indicated by the unchanged p-eIF2α
levels (**Supplementary Fig. 23c**). Immunofluorescence against nuclear speckles marker SRRM2
revealed that compared to controls, V337M iPSC-neurons exhibited aberrant nuclear speckle
morphology characterized by smaller size and more spherical shape, and 12 hours of 100nM PP
treatment fully restored nuclear speckle morphology to normal size and diffuseness (**Fig. 6d, e**).
Consequently, PP markedly reduced the level of V337M p-Tau, concomitant with increased
autophagic flux (**Fig. 6d-h**). To confirm that PP rejuvenates nuclear speckles and reduces V337M
p-Tau in a SON-dependent manner, we knocked down SON using lentiviral shRNA and repeated
the experiment. As demonstrated in **Fig. 6i-k and Supplementary Fig. 23d**, PP failed to
rejuvenate nuclear speckles or reduce p-Tau level in the presence of SON knockdown in either
wild-type or V337M Tau-expressing iPSC neurons. Together, these findings indicate that PP
treatment has great potential to normalize gene expression patterns and reduce Tau burden in
AD/ADRD-affected humans with severe tauopathy. Further, these results provide strong support
for the decline of nuclear speckles LLPS dynamics (becoming smaller and more spherical) as a
driver of tauopathy.

Discussion

Several recent studies indicate that both the decline of nuclear speckle functions and
dysregulated mRNA splicing are associated with proteinopathies in humans, including tauopathy,
RP and amyotrophic lateral sclerosis (ALS) [90, 91, 92, 93]. For example, two studies showed that
elevated Tau aggregates have the capability to relocate to the nucleus, thereby directly modifying
the characteristics of nuclear speckles [92, 94]. In agreement with this study, a recently published
Tau interactome in human iPSC neurons [95] also revealed strong enrichment of Tau-interacting
proteins involved in regulating RNA splicing and RNA stability. Additionally, it was previously
shown that cryptic splicing errors are associated with neurofibrillary tangle burden in human AD
subjects [96]. In humans, among the 12 autosomal dominant RP genes identified, four encode
ubiquitously expressed proteins involved in pre-mRNA splicing (including *PRPF31*, *PRPF8*,
PRPF3 and *RP9*), demonstrating the important roles of RNA processing in the pathogenesis of
retinal degeneration [93]. These studies thus provide the rationale for nuclear speckle rejuvenation
as a strategy for counteracting various proteinopathies.

Exploring the therapeutic potential of targeting biomolecule condensates represents an
exciting avenue for research and drug development [97, 98]. Our study is proof of principle
demonstrating that nuclear speckle LLPS can also be therapeutically targeted. Manipulating
nuclear speckle LLPS through SON overexpression is a conceptually viable approach. However,
the practical implementation of this strategy presents significant challenges due to the large size

of the human SON open reading frame, making it technically difficult to design gene therapy
targeting SON. That said, we cannot rule out the possibility that overexpression of specific
truncated SON domains, such as SON IDR2, may be sufficient to rejuvenate nuclear speckles,
and future efforts will be directed toward exploring such possibilities.

Through a high-throughput drug screen, we identified PP as a small nuclear speckle
rejuvenator by directly interacting with SON and modulating nuclear speckle LLPS dynamics.
Interestingly, pyrvinium is enriched with cationic amines and aromatic motifs, chemical features
that were predicted to partition into various nuclear condensates [99]. PP was originally developed
as an anthelmintic drug effective for treating pinworm infections [100]. Moreover, it has gained strong
recent interest as an anti-cancer reagent due to its ability to inhibit WNT signaling [100]. Our current
study further expands its therapeutic values to proteinopathies, including both reducing tauopathy
in neurons and flies and protecting against retina degeneration in an *ex vivo* mouse RP model.
Future efforts should be directed toward testing the toxicity and efficacy of this drug in mouse
models of neurodegenerative diseases.

While we don't yet know the full detailed mechanisms by which PP modulates nuclear
speckles dynamics and boosts proteostasis gene transcription, several lines of evidence suggest
that it does so in part by reducing the surface tension and consequently increasing the surface
areas of nuclear speckles via an SON-dependent manner. Our in vitro reconstitution system
further showed that reduced nuclear speckles surface tension by PP further facilitates nuclear
speckles wetting of chromatin [46]. Thus, since spliceosomes reside at the interfacial boundary
between nuclear speckles and nucleoplasm/chromatin[44], larger surface areas also entails a
higher probability of spliceosome stably engaging in mRNA processing and transcription
elongation. The mechanism by which PP reduces the surface tension of nuclear speckles is not
yet fully understood. While the straightforward explanation would be that PP acts as a surfactant,
this seems unlikely due to its lack of hydrophilic moieties. Alternatively, it is plausible that by
interacting with SON, PP weakens the intermolecular attractive interactions among different SON
proteins, and/or SON and IDRs of other nuclear speckle proteins due to screening effects, leading
to an overall reduction of surface tension of nuclear speckles, similar to what is previously
described for the effect of increasing salt concentration on reducing surface tension for protein
condensates [101, 102]. The positive charge carried by pyrvinium adds to the allure of this hypothesis.
Further research is needed to elucidate the precise mechanisms by which PP influences the
surface tension of nuclear speckles.

Genetic and pharmacological rejuvenation of nuclear speckles by SON and pyrvinium share
similar transcriptome signatures, including upregulation of extensive protein quality control gene
expression, and intriguingly, downregulation of YAP1-regulated genes involved in cell migration,
cell proliferation, would healing, and extracellular matrix organization. The contrasting changes in
proteostasis and YAP1-mediated cell dynamics gene expression are observed in various cell lines
under proteostatic stress. Consequently, both SON overexpression and ER stress inhibits
YAP/TEAD transcriptional activity and impedes cell migration. These findings demonstrate that
YAP1 signaling is an inherent component of global transcriptional control of proteostasis. One
possible explanation for this phenomenon is that cells need to allocate their energy towards
enhancing overall proteostasis, which may come at the cost of cell proliferation, migration, and
extracellular matrix organization. Therefore, an energetic trade-off between proteostasis and cell
dynamics control could be crucial for maintaining cellular functions when faced with fluctuating
environments, such as proteotoxic stress.

Nuclear speckles play a vital role in coordinating the opposing changes observed in
proteostasis and cell dynamics regulation. Elevated SON expression facilitates increased
physical interactions between nuclear speckles and XBP1s, leading to augmented transcription
of proteostasis genes. Since the number of proteostasis genes under the control of SON
surpasses those directly regulated by XBP1s, we postulate that nuclear speckles can be recruited
to additional proteostasis bZIP TFs upon rejuvenation, possibly via increased wetting between
nuclear speckle and TF-mediated condensates [46]. Future work with unbiased profiling of nuclear
speckle composition (via proximity labeling for example) can unveil the detailed molecular
mechanisms through which nuclear speckle rejuvenation globally activates the proteostasis gene
program. On the other hand, much less is clear on how nuclear speckle rejuvenation represses
YAP1 transcription activity. Upon SON overexpression and pyrvinium treatment, we found a
significantly reduced level of nuclear YAP1 protein and a lower nucleus/cytosol ratio, indicating
an active nuclear exclusion of YAP1 protein. Like nuclear speckles, YAP1 can also form
biomolecular condensates, and a recent study reported that YAP1 nuclear condensates and the
nuclear speckles showed limited nuclear co-localization [103], suggesting a low level of wetting of
these two condensates under normal physiological conditions. Pending further investigation, we
speculate herein that nuclear speckles rejuvenation may further reduce the wetting of these two
condensates, resulting in the alteration of YAP condensate composition, and ultimately its nuclear
exclusion.

While both proteostasis and YAP1 signaling are downstream of nuclear speckles, direct
antagonistic reciprocal interactions between these two are likely to be present as well. A recent
study reported that in *Drosophila*, the proteostasis output gene Bip can sequester the fly YAP1
ortholog Yorkie, in the cytoplasm to restrict Yorkie transcription output [104]. Conversely, in
undifferentiated pleomorphic sarcoma, YAP1 can suppress PERK and ATF6-mediated UPR
target expression, and treatment with the YAP1 inhibitor Verteporfin upregulated the UPR and
autophagy [105]. The latter is further consistent with our findings showing that restoring YAP1 activity
dampened the efficacy of PP on activating protein quality control gene expression and reducing
proteinopathies. These findings indicate that in order to maximize the effectiveness of nuclear
speckle rejuvenation, it is crucial to maintain elevated levels of protein quality control while
simultaneously reducing YAP1 activity. Thus, a delicate balance between protein quality control
and YAP1 activity appears essential for effective nuclear speckle rejuvenation. This observation
may also explain why therapies merely aimed at activating protein quality control pathways often
have limited efficacies. Supporting this notion, while reduced expression of genes involved in
ERAD and autophagy are observed in the brains of individuals with AD, these subjects also exhibit
elevated gene expression of *YAP1*, *TEAD1,* and other YAP1 target genes, consistent with
previous studies [87, 106]. Increased YAP1 target gene expression was further observed in human
iPSC tauopathy model as well as in the retina of Rho [P23H/+] mice. These results collectively
highlight the importance of suppressing YAP1 signaling as a potential strategy for managing both
AD and RP.

In conclusion, our study makes substantial conceptual contributions to the broader
proteostasis framework by incorporating nuclear speckle LLPS and YAP1 signaling as critical
components. From a translational perspective, our research unveils the promising therapeutic
potential of nuclear speckle rejuvenation in tackling proteinopathies, achieved by simultaneous
activation of protein quality control and inhibition of YAP1 activity. Additionally, our findings
underscore the significance of harnessing the 12-hour oscillator to unveil hidden principles of
proteostasis control.

Materials and methods

Mice

For retinal explant studies, wildtype C57BL/6J and Rho$^{P23H/+}$ knock-in mice (Jackson Laboratory Strain #017628) were euthanized by CO2 and retinae were isolated for culture. The animal studies were carried out in accordance with the National Institutes of Health guidelines and were granted formal approval by the University of Pittsburgh's Institutional Animal Care and Use Committee (approval numbers IS00013119 and IS00023112).

Larva crawling assay

PP solubilized in DMSO were diluted directly into the fly medium at the final concentration of 25 µM and vortexed extensively to obtain homogeneous culture. Crawling assays were performed on 1.5% agarose plates made with a 2.3:1 combination of grape juice and water. A sample size of 10 to 15 larvae were selected for each genotype and assays were done using larvae in the third instar state. The larvae were first removed from vials and gently placed into a petri dish containing deionized water to allow for residual food to be washed off the body. After 15 seconds, the larvae were transferred to a petri dish containing the 1.5% agarose mixture and were given one minute to rest. They were then transferred to a second dish filled with the 1.5% agarose mixture and timed immediately for one minute, during which their crawling performance was measured. A transparent plastic lid was placed on top of the plates and the crawling path of the larvae were traced. Observations of the crawling activity were done under a light microscope. The brightness and distance of the light source above the plates were kept constant across all trials and genotypes. The crawling paths of the larvae were measured using FIJI ImageJ and the average distance traveled was taken for each genotype.

Adult fly climbing assay

Male adult Drosophila melanogaster flies at 14 and 21 days of age fed with a normal diet or diet supplemented with 25 µM PP were used for assessing climbing ability. Flies were grouped into cohorts of the same sex, pre-mated, and age-matched, with a maximum of 20 individuals per vial (usually 5-15). All flies used in each trial were hatched within a 3-day window. The evening prior to each assay, flies were gently transferred to fresh tubes to allow for grooming and access to food. To ensure consistent conditions, assays were conducted at approximately the same time of day with a consistent ambient light setting. A custom climbing vial was employed, divided into six compartments, each labeled with a number (1 to 6) to denote climbing speed. The vial was positioned against a white background to enhance visibility during photography. Flies were transferred from their housing vial to the climbing vial, which was covered with a plastic plate on top. To initiate the assay, the flies were gently tapped to the bottom of the vial and allowed 10 seconds to climb. A cell phone camera was used to capture a photograph of the vial. Care was taken to ensure the camera was level with the vial, all flies were visible, and the background was free from stains or spots. The number of flies in each compartment of the climbing vial was counted at each time point and recorded on a dedicated worksheet. Each cohort of flies underwent five consecutive trials, with approximately 1 minute of rest between each trial. The average score of each cohort was determined by dividing the total score by the total number of flies.

Fibroblast cell culture and drug treatment

MEFs and NIH 3T3 cells were cultured at 37°C and 5% CO_2 in Dulbecco's Modified Eagle's
Medium (DMEM, glucose 4.5 g/L with phenol red) and supplemented with 10% fetal bovine serum
(FBS), 1 mM sodium pyruvate (Gibco), and penicillin (100 U/mL)-streptomycin (100 µg/mL)
(Gibco). Methods for the manipulation of *Son* (transient knockdown or constitutive
overexpression) and validation of changes to protein (SON) levels with regards to the mRNA-Seq
data are previously described in [9]. For Tu treatment, 100 ng/mL Tu (in DMSO) for six hours was
used unless otherwise noted. NB (HY-50904), PB (HY-12047), PH (HY-B0883), PP (HY-A0293),
MG-132 (HY-13259), SBI-0206965 (HY-16966) and XMU-MP-1 (HY-100526) were purchased
through MedChemExpress and BafA (1334) were purchased from Tocris. All drugs were handled
per manufacturer instruction.

siRNA Transient Transfections

MEFs were transfected with 10µM of different siRNAs for 24~48 hours with Lipofectamine
RNAiMAX reagents (Life technologies) per the manufacturer's instructions. Sources of siRNA are
as follows: siGENOME non-targeting siRNA pool (Dharmacon, D-001206-1305), siGENOME
SMARTpool son siRNA (Dharmacon, L-059591-01-0005), siGENOME SMARTpool Stk3/Mst1
siRNA (Dharmacon, L-040440-00-0005), and siGENOME SMARTpool Stk4/Mst2 siRNA
(Dharmacon, L-059385-00-0005).

Primary neuron cell culture and P301S-Tau virus infection

The cerebral cortices of 3-4 neonate mice (P0) were dissected on ice, the meninges were
removed and placed in the cold dissection medium (DM), consisting of 6 mM MgCl2 (Sigma
M1028-100 ml), 0.25 mM $CaCl_2$ (Sigma C7902), 10 mM HEPEs (100X), 0.9% Glucose, 20 µM D-
AP5 (Cayman, NC1368401), and 5 µM NBQX (Tocris Bioscience, 10-441-0). After dissection, the
brain tissues were washed with DM 1~2 times and incubated with 13mL of DM containing papain
(Worthington, LK003176) in 37°C water bath for 20 min. The suspension was shaken every 5
min. 10mL media containing 18 ml DM + 2 ml low OVO + 133 ul DNase I (dilute 10X low OVO
and 150x DNase I to DM) were added into the suspension to stop the digestion in 37°C water
bath for 5 min. The solution was taken off and 10 ml fresh solution was added in. Then the tissues
were triturated until there were no visible chunks, and the solution was filtered through the 70 µm
cell strainer. The cell solution was then centrifuged at 1000 rpm for 10 min and the supernatant
was discarded. The cell pellet was gently resuspended in 20 ml B27/NBM/High glucose media,
and the suspension was centrifuged at 850 rpm for 5 min. After that, the supernatant was taken
off and B27/NBM (1 ml/mouse brain) was added to resuspend the cells until single cell solution.
The cells were counted and plated onto the coverslips at 250k in 24-well plates for imaging or
800k in 12-well plates for qRT-PCR or Western blots. AAV-P301S hTau (Viro-vek) were infected
at DIV1 at 100 MOI.

Human iPSC-derived neurons culture

Human iPSC-derived neurons were pre-differentiated and differentiated as described [89]. Briefly,
iPSCs were pre-differentiated in Matrigel-coated plates or dishes in N2 Pre-Differentiation
Medium containing the following: KnockOut DMEM/F12 as the base, 1× MEM non-essential
amino acids, 1× N2 Supplement (Gibco/Thermo Fisher Scientific, cat. no. 17502-048), 10 ng/ml
of NT-3 (PeproTech, cat. no. 450-03), 10 ng/ml of BDNF (PeproTech, cat. no. 450-02), 1 µg/ml of
mouse laminin (Thermo Fisher Scientific, cat. no. 23017-015), 10 nM ROCK inhibitor and 2 µg
/mlof doxycycline to induce expression of mNGN2. After 3 d, on the day referred to hereafter as
Day 0, pre-differentiated cells were re-plated into BioCoat poly-D-lysine-coated plates or dishes

(Corning, assorted cat. no.) in regular neuronal medium, which we refer to as +AO neuronal
medium, containing the following: half DMEM/F12 (Gibco/Thermo Fisher Scientific, cat. no.
11320-033) and half neurobasal-A (Gibco/Thermo Fisher Scientific, cat. no. 10888-022) as the
base, 1× MEM non-essential amino acids, 0.5× GlutaMAX Supplement (Gibco/Thermo Fisher
Scientific, cat. no. 35050-061), 0.5× N2 Supplement, 0.5× B27 Supplement (Gibco/Thermo Fisher
Scientific, cat. no. 17504-044), 10 ng/ml of NT-3, 10 ng/ml of BDNF and 1 µg/ml of mouse laminin.
Neuronal medium was half-replaced every week.

Efficacy test of PP in Retina explant culture

Wild type and $Rho^{P23H/+}$ mice were euthanized at P15, and retina explants were isolated and
cultured as previously described[107, 108]. Briefly, eyeballs were enucleated and incubated in Ames
solution containing 0.22 mM L-cysteine (Sigma-Aldrich) and 20 U papain (Worthington, Freehold
NJ, USA) at 37 °C for 30 min. The digestion was stopped by transferring the eyes to Dulbecco's
modified Eagle's medium (DMEM; Gibco) containing 10% fetal calf serum (FCS; Gibco) and
penicillin &streptomycin antibiotics (1x, GenClone) at 4 °C for 5 min. The eye cup was made by
gently removing the cornea, iris and lens. Each eye cup was flattened by four radio cuts and the
sclera was then carefully peeled off from the retina:RPE complex. The retina:RPE explant was
transferred to a trans well insert with 0.4-micron pore polycarbonate membrane (ThermoFisher)
sitting on the surface of 1.5 mL of neurobasal-A plus medium (Gibco) containing 2% B27
supplement (Gibco) in a 6-well cell culture plate, and the RPE layer was facing down the transwell
membrane. The retinal explants were cultured at 37 °C with 5% CO_2.The medium was replaced
with fresh medium containing 0.5 µM PP after 24 h, which was replaced again every 2 days until
10 days in culture (DIV). A visible light optical coherence tomography (vis-OCT) prototype [109] was
utilized to monitor the explants noninvasively at day 0 and day 10. Retinal layers were segmented
automatically using a deep learning method and then manually corrected by a customized
software to calculate the retinal thickness. Retina explants were collected at 10 DIV and
processed for fixation, dehydration, paraffin embedding, cross-sections, dewaxing, rehydration
and hematoxylin and eosin (H&E) staining[110]. H&E-stained slides were imaged by regular light
microscopy with a color camera, and the number of nuclei in the outer nuclear layer (ONL) was
calculated manually.

Autophagy reporter assay

To express the LC3 reporter in the neurons, the primary mouse neuronal cultures were infected
with the homemade lentivirus-mCherry-GFP-LC3 for 7 days. The florescent signal from the
vacuoles at different stages were acquired by confocal imaging. The mCherry-GFP-LC3
fluorescence images were acquired with a Leica TCS SP8 confocal system using 63x oil-
immersion objective. 488 nm and 568 nm laser were used to excite the GFP and mCherry,
respectively. Images were taken with the same confocal settings. Minor image adjustment
(brightness and/or contrast) was performed in ImageJ. The GFP and mCherry signal collected
were merged into one image to quantify the red, green, and yellow vacuoles for quantification of
different types of vacuoles. The different colored fluorescent signal was manually counted in each
cell, and each point represents the average number of the specific vacuole for one cultured cell.
For autophagy reporter assay in MEFs, pCDH-EF1a-mCherry-EGFP-LC3B was a gift from Sang-
Hun Lee (Addgene plasmid # 170446 ; http://n2t.net/addgene:170446 ; RRID:Addgene_170446)
[111] and purchased from addgene. Lentivirus was packaged from HEK293T cells as previously
described [19] and was used to infect MEFs with a MOI of 3 three times. The quantification was
performed essentially the same way as in neurons.

TEAD luciferase reporter assay

MEFs with the CRISPRa system either overexpressing *Son* or serving as controls used in this assay were previously described in [112]. Briefly, cells were seeded at a density of 7000 cells per well in a 96-well plate with a clear bottom and white walls. Cells were then transfected using the Lipofectamine 3000 transfection kit (ThermoFisher #L3000015) for 22 hours with the 8xGTIIC-luciferase plasmid, a gift from Stefano Piccolo (Addgene plasmid # 34615 ; http://n2t.net/addgene:34615 ; RRID:Addgene_34615) [113]. The Dual-Glo Luciferase Assay System (Promega #E2920) was used with a SpectraMax i3x plate reader (Molecular Devices) to measure firefly luciferase signal (500ms integration time, 1mm from the plate read height).

Proteasome 20S activity assay

The proteasome 20S activity assay was performed per manufacturer's instruction (Sigma Aldrich, MAK172). Briefly, MEFs transfected with scrambled control or *Son* siRNA were treated with DMSO or 100nM PP for 24 hours. After treatment, cells were washed with PBS and cultured in phenol-free DMEM. Assay reagents were added directly to the cells, and fluorescent signals were measured by a fluorescent plate reader 2 hours later at $\lambda ex = 490$ nm and and $\lambda em = 525$ nm. The final signal was corrected by subtracting the fluorescence of the background blank (medium without cells) from the fluorescence of all test wells.

Scratch assay

Cells were grown until they were 100% confluent, ER stress was induced as previously described, and then a single scratch was performed with a pipette tip per well. Cells were imaged immediately after scratching (0hr) and then after 23hr. The Cell Profiler [114] "Wound Healing" pipeline (https://cellprofiler.org/examples) was used to measure the "Percentage of Gap Filled".

Immunoblot

Different cells were harvested and fractionated to produce cytosolic and nuclear lysates using the NE-PER kit (Thermo Fisher Scientific). For whole cell lysates, cells were lysed in RIPA buffer. Both protease and phosphatase inhibitors were included in the respective lysis buffer. ~47 µg of protein was separated on a 4%-15% gradient SDS-polyacrylamide gel (Bio-Rad) which were transferred to nitrocellulose membranes, stained with Ponceau S stain, washed, blocked with 5% non-fat milk, and incubated overnight at 4°C with the following primary antibodies: anti-α-Tubulin (Cell Signaling Technology (C.S.T.) #2144), anti-Lamin A/C (C.S.T. #4777), anti-SON (Abcam #121033 and LSBio LS-C803664), anti-YAP1 (C.S.T. #12395), anti-GFP (C.S.T. #2956), anti-Tau (Sigma-Aldrich, #A0024), anti-p-Tau (a gift from Dr. Peters Davies), anti-β-actin (C.S.T. #4970), anti-puromycin (BioLegend 381502), anti-ubiquitin (C.S.T. #58395), anti-LC3-I/II (C.S.T. #2775), anti-p62 (C.S.T. #23214), anti-ATF4 (C.S.T. #11815), anti-ATF6 (Novus 70B1413.1), anti-MST1 (C.S.T. #3682), anti-MST2 (C.S.T. #3952), anti-eIF2α (C.S.T. #5324), anti-p-eIF2α (Ser51) (C.S.T. #3597) and anti-XBP1s (BioLegend 658802). The 1D4 anti-rhodopsin antibody[115] was obtained as a gift from Dr. Krzysztof Palczewski's laboratory. Membranes were treated with the appropriate secondary antibody conjugated to horseradish peroxidase the following day and then ECL Prime Western Blotting Detection Reagent (Cytiva) was applied. A Bio-Rad ChemiDoc MP Imaging System was used to visualize the signal, and signal intensities were determined with ImageJ [116]. For anti-Rhodopsin western blot, the protein samples were not boiled before loading.

Cellular thermal shift assay (CETSA)

EGFP::SC35 MEFs with EGFP knocked into the N-terminal of mouse *Srsf2* locus (previously
described in [9]) were treated with either DMSO or 3μM PP for 50 minutes at 37°C. Cells were then
trypsinized and resuspended in PBS with either DMSO or 3μM PP and 100 μL of the suspensions
were distributed to PCR tubes for the thermal shift assay (three minutes at a range of
temperatures). The temperatures used were: 42.0°C, 42.5°C, 43.9°C, 46.2°C, 49.3°C, 53.3°C,
57.9°C, 62.1°C, 65.2°C, 67.8°C, 69.2°C, and 70.0°C. After the samples were heated, they sat at
20°C for three minutes, were snap frozen in liquid nitrogen and thawed for three cycles to lyse
the cells, and then spun at 20,000 x g for 20 minutes at 4°C. The supernatant was then removed,
and immunoblotting was performed using anti-GFP (C.S.T. 2956), anti-ATF4 (C.S.T. #11815),
and anti-SON (Lifespan Biosciences #LS-C803664-100), followed by appropriate secondary
antibody. Band intensity on the blots were relative to the intensity of the 42°C band and were
normalized so that this band's (42°C) intensity was set equal to 1.

Protein purification and *in vitro* droplet formation assay

Regions of SON and the entire SRSF2 protein were fused to mCherry. cDNA encoding the SON-
IDR N terminal (region 1), SON-IDR C-terminal (region 2), and SRSF2 were each cloned into the
expression vector pET21a (+)-Histag-mCherry (Addgene plasmid # 70719) (Niederholtmeyer et
al., 2015). The plasmids obtained were transformed into C3013 *E. Coli* (NEB C3013I). Fresh
bacterial colonies were inoculated into LB media containing ampicillin and grown overnight at
37°C. Overnight cultures were diluted in 500mL of LB broth with ampicillin and grown at 37°C until
reaching OD 0.6. IPTG was then added to 2mM, and growth continued for 3h at 37°C. The cells
were pelleted and stored frozen at -80°C. Bacterial pellets were resuspended in 15mL of Buffer
A (50 mM Tris-HCl, 500 mM NaCl) containing protease inhibitors (Pierce, A32965) and 10mM
imidazole. The suspension was sonicated on ice for 15 cycles of 30 sec on, 30 sec off. The lysate
was centrifuged for 40 minutes at 15,000 RPM at 4°C to clear debris, then added to 2mL of
preequilibrated Ni-NTA agarose beads (Qiagen cat no. 30210). The agarose lysate slurry
incubated for 1.5hrs at 4°C while rocking, then allowed to flow through the column. The packed
agarose was washed with 15mL of Buffer A with 10mM imidazole. Protein was eluted with 5mL
of Buffer A containing 15mM imidazole, 10mL Buffer A containing 100mM imidazole, then 10mL
Buffer A containing 200mM imidazole. All elutions were collected in 1mL fractions. Aliquots of the
collected fractions were run on an SDS-PAGE gel and stained with Imperial Protein Stain to verify
the amount and purity of the protein. Fractions containing protein were combined and dialyzed
against Dialysis Buffer (50mM Tris-HCl, 500mM NaCl, 10% glycerol, 1mM DTT). Recombinant
mCherry fusion proteins were concentrated and desalted to 200uM protein concentration and
125mM NaCl using Amicon Ultra centrifugal filters (MilliporeSigma cat no. UFC801024) following
manufacturer's instructions. 20uM of recombinant protein was added to Droplet Buffer (50mM
Tris-HCl, 10% glycerol. 1mM DTT, 10% PEG) containing indicated final salt and pyrvinium
concentrations. For droplet formation assay with Hela nuclear extract supplementation, NE was
added to different concentration of SON IDR2 at the final concentration of 1.5mg/ml in Droplet
buffer (20mM HEPES, pH 7.9, 20% glycerol, 125mM KCl, 0.2mM EDTA, 0.5mM DTT, 10% DEG).
For droplet formation assay with GFP::SRSF2 MEF nuclear extract supplementation, NE was
added to 10uM SON IDR2 at the final concentration of 0.6mg/ml in Droplet buffer. A custom
imaging chamber was created by placing strips of tape on a glass coverslip, forming a square.
The protein solution was immediately loaded onto the center of the square and covered with a
second glass coverslip. Slides were then imaged with a Leica confocal microscope with a 63x oil
objective. The image series were taken over a 20-minute time span with 1 image every 30
seconds.

Mass spectrometry to profile condensates composition

Hela nuclear extract samples were thawed at room temperature, vortexed for 10 minutes, bath sonicated for 5 minutes and centrifuged at 13000g for 10 minutes at room temperature prior to quantification of total protein by a Pierce 660 Protein Assay (Thermo Scientific #22660). Condensates were obtained by centrifuging at 10,000xg for 10 minutes and resuspended in 5%SDS in 50 mM TEAB prior to total protein quantification. Protein digestion was carried out on 10 μg of protein from each sample on S-trap micro columns (Protifi) according to the manufacturer's protocol. Following digestion, peptide samples were then dried in a speedvac and resuspended in a solution of 3% acetonitrile and 0.1% TFA and desalted using Pierce Peptide Desalting Spin Columns (Thermo Scientific # 89851). Eluants were dried in a speedvac and resusupended in a solution of 3% acetonitrile and 0.1% formic acid to a final concentration of 0.5 μg/μL. Mass spectrometry analysis was conducted on a Thermo Fisher QE-HFX coupled to a Vanquish Neo UHPLC. Approximately 1 μg of each sample was loaded onto an EASY-Spray PepMap RSLC C18 column (2 μm, 100A, 75μm x 50 cm) and eluted at 300 nl/min over a 120-minute gradient. MS1 spectra were collected at 120,000 resolution with a full scan range of 350 – 1400 m/z, a maximum injection time of 50ms and the automatic gain control (AGC) set to 3e6. The precursor selection window was 1.4 m/z and fragmentation were carried out with HCD at 28% NCE. MS2 were collected with a resolution of 30,000, a maximum injection time of 50ms and the AGC set to 1e5 and the dynamic exclusion time set to 90s. The collected MS data were analyzed using MSFragger V4.0[1] and searched against the human SwissProt database. The search parameters were set as follows: strict trypsin digestion, missing cleavage up to 2, carbamidomethylation of cysteine as static modification, oxidization of methionine and protein N-terminal acetylation as variable modification, a maximal mass tolerance of 20 ppm for the precursor ions and 20ppm for the fragment ions, and false detection rate (FDR) was set to be 1%.

1,6-hexanediol treatment to examine effects of PP on LLPS

A 10% (w/v) 1,6-hexanediol (1,6-HD, MilliporeSigma) solution was prepared in Dulbecco's Modified Eagle's Medium (DMEM, glucose 4.5 g/L with phenol red) supplemented with 10% fetal bovine serum (FBS), 1 mM sodium pyruvate (Gibco), and penicillin (100 U/mL)-streptomycin (100 μg/mL) (Gibco). To examine NS LLPS dynamics, EGFP::SC35 MEFs (previously described in [9]) were treated with DMSO or 1 μM PP for 30 minutes and then treated with 0, 1, 2, or 10% 1,6-HD for 20 minutes. Cells were fixed in 2% paraformaldehyde, stained with bisBenzimide H 33258 (Hoechst), and then imaged. Image analysis was completed in Cell Profiler; briefly, the process was to image the cells in the 405 (Hoescht), 488 (GFP), and 555 (high intensity to image whole cells) and then use the 405 channel to determine the nuclei boundaries, 488 to determine EGFP::SC35, and 555 to determine the area of the whole cells. We differentiated nuclear and cytosolic areas by subtracting the 405 signals from the 555 signals. 488 signal was then quantified in the aforementioned nuclear area and cytosolic area and compared. 20+ cells were measured for each condition. The average Manders coefficient was determine with ImageJ [116] by averaging the tM1 and tM2 values.

This same process was used to examine the effects of PP on GW182 and MED1 except in wildtype MEFs. The signal was identified by immunofluorescence (IF). Briefly, IF was performed by fixing cells with 4% paraformaldehyde in PBS, permeabilizing with 0.2% Triton X-100 in PBS, blocked with 2% bovine serum albumin in PBS, and then incubated with primary antibody (GW182 (ab156173, Abcam) or MED1 (ab60950, Abcam)) diluted per manufacturer's recommendation overnight at 4°C. Cells were then treated with the appropriate 1:1000 secondary antibody

883 overnight at 4°C, stained with Hoechst the following day, and then mounted with ProLong Gold
884 Antifade (Invitrogen). Signal (either IF or endogenous GFP) sphericity was determined as
885 previously described in [9]. 20+ cells were measured for each condition.

Reverse transcription quantitative polymerase chain reaction (RT-qPCR)

For Reverse transcription-quantitative polymerase chain reaction (RT-qPCR), cDNA was
produced using the Superscript III (Thermo Fisher) kit and qPCR was completed using the SYBR
Green system (Thermo Fisher) in a CFX384 Real-Time System (Bio-Rad). Endogenous *β-actin*
levels were used as controls. The qPCR primer sequences were as follows:

β-actin forward: AAGGCCAACCGTGAAAAGAT
β-actin reverse: GTGGTACGACCAGAGGCATAC
Amot forward: CTGGAAGCAGATATGACCAAGT
Amot reverse: GGTGTTAGGAGAGTGGCTAATG
Atf4 forward: CCACTCCAGAGCATTCCTTTAG
Atf4 reverse: CTCCTTTACACATGGAGGGATTAG
Atg4c forward: GTGCGGAATGAGGCTTATCA
Atg4c reverse: CCAGACTTCTTCCCAAACTCTATC
Bmp4 forward: AACGTAGTCCCAAGCATCAC
Bmp4 reverse: CGTCACTGAAGTCCACGTATAG
Ern1 forward: TCCTAACAACCTGCCCAAAC
Ern1 reverse: TCTCCTCCACATCCTGAGATAC
Fdz1 forward: GAGATCCACCTTCCAGCTTTAT
Fzd1 reverse: CACTCCCTCTGAACAACTTAGG
Hyou1 forward: GAGGCGAAACCCATTTTAGA
Hyou1 reverse: GCTCTTCCTGTTCAGGTCCA
Manf forward: GACAGCCAGATCTGTGAACTAAAA
Manf reverse: TTTCACCCGGAGCTTCTTC
Rnf166 forward: GAAGACACACTCCCGCTTTA
Rnf166 reverse: CTGAGACCAACTCTCCTTGTG
Sirt2 forward: CATAGCCTCTAACCACCATAGC
Sirt2 reverse: GTAGCCTGTTGTCTGGGAATAA
Sqstm1 forward: AACAGATGGAGTCGGGAAAC
Sqstm1 reverse: AGACTGGAGTTCACCTGTAGA
Tgfb3 forward: CCACGAACCTAAGGGTTACTATG
Tgfb3 reverse: CTGGGTTCAGGGTGTTGTATAG
Ube2q2 forward: TTCCTAAGCACCTGGATGTTG
Ube2q2 reverse: CTCCTCCTCTTCCTCTTCTTCT
Xbp1 forward: GGGTCTGCTGAGTCC
Xbp1 reverse: CAGACTCAGAATCTGAAGAGG
Cul5 forward: GAACACAGGCACCCTCATATT
Cul5 reverse: AGTTACACTCTCGTCGTGTTTC
Psmg1 forward: CCAGTGGTTGGAGAAGGTTT
Psgm1 reverse: GGGTCTTGTAGTCTGTGATGTG
Atg14 forward: CATTCCCTGGATGGGCTAAA
Atg14 reverse: CCTCAGGAACAAGAAGGAAGAG
Yap1 forward: CCAATAGTTCCGATCCCTTTCT

Yap1 reverse: TGGTGTCTCCTGTATCCATTTC

Chromatin Immunoprecipitation (ChIP)

ChIP for SC35 was performed using anti-SC35 antibody (ab11826, Abcam) as previously
described [117]. Briefly, mouse liver samples were submerged in PBS + 1% formaldehyde, cut into
small (~1 mm3) pieces with a razor blade and incubated at room temperature for 15 minutes.
Fixation was stopped by the addition of 0.125 M glycine (final concentration). The tissue pieces
were then treated with a TissueTearer and finally spun down and washed twice in PBS. Chromatin
was isolated by the addition of lysis buffer, followed by disruption with a Dounce homogenizer.
The chromatin was enzymatically digested with MNase. Genomic DNA (Input) was prepared by
treating aliquots of chromatin with RNase, Proteinase K and heated for reverse-crosslinking,
followed by ethanol precipitation. Pellets were resuspended and the resulting DNA was quantified
on a NanoDrop spectrophotometer. An aliquot of chromatin (10 µg) was precleared with protein
A agarose beads (Invitrogen). Genomic DNA regions of interest were isolated using 4 µg of
antibody. Complexes were washed, eluted from the beads with SDS buffer, and subjected to
RNase and proteinase K treatment. Crosslinking was reversed by incubation overnight at 65 °C,
and ChIP DNA was purified by phenol-chloroform extraction and ethanol precipitation. ChIP-
qPCR for MEFs were essentially performed the same way as previously described with anti-SC35
(ab11826, Abcam) and anti-XBP1s antibody (Biolegend 658802), except that the MEFs were
directly fixed with 1% formaldehyde before subject to nuclei isolation and chromatin
immunoprecipitation. The primers used for ChIP-qPCR are as follows:

Gene desert forward primer: GCAACAACAACAGCAACAATAAC
Gene desert reverse primer: CATGGCACCTAGAGTTGGATAA
Xbp1 promoter region forward primer: GGCCACGACCCTAGAAAG
Xbp1 promoter region reverse primer: GGCTGGCCAGATAAGAGTAG
Xbp1 gene body region forward primer: CTTTCTCCACTCTCTGCTTCC
Xbp1 gene body region reverse primer: ACACTAGCAAGAAGATCCATCAA
Manf promoter region forward primer: ACAGCAGCAGCCAATGA
Manf promoter region reverse primer: CAGAAACCTGAGCTTCCCAT
Manf gene body region forward primer: CAACCTGCCACTAGATTGAAGA
Manf gene body region reverse primer: AGGCATCCTTGTGTGTCTATTT
Hyou1 promoter region forward primer: GACTTCGCAATCCACGAGAG
Hyou1 promoter region reverse primer: GACTTCTGCCAGCATCGG
Hyou1 gene body region forward primer: TGGAAGAGAAAGGTGGCTAAAG
Hyou1 gene body region reverse primer: TCCCAAGTGCTGGGATTAAAG

HTS of FDA-approved drugs and quantitative imaging analysis

EGFP::SC35 MEFs were seeded in black 384 well plate with glass bottom (Cellvis). The FDA-
approved compound library (100nL per drug) was stamped to 384-well tissue culture plates using
CyBio Well vario (Analytik Jena). Compound solutions were then added to the cell plate at the
final concentrations of 10 µM using a BRAVO liquid handler. After 18 hours of treatment, culture
media was removed, and cells were fixed with 4% PFA followed by DAPI staining. Cells were
imaged using a GE INCELL 2200 with 60x lens. Maximum intensity projection images were
captured, and nuclear speckles sphericity was quantified with CellProfiler as previously described
in [9]. Briefly, for speckle i the sphericity is defined as equation 1:

971
$$\text{Sphericity } i = 2\sqrt{\pi} * \sqrt{\text{area } i} \div \text{circumference } i \qquad (1)$$

972   So that a perfect circle will have a sphericity of 1, and a line will have a sphericity of 0. To calculate
973   the average sphericity of a given image that has k total speckles, we calculated the area-weighted
974   average as described in equation 2.

975
$$Average\ sphericity/image = \sum_1^k Sphericity\ i \times area\ i / \sum_1^k area\ i \qquad (2)$$

For the secondary screening to determine if compounds affected nuclear speckles morphology in
a dose-dependent manner, specific compounds were selected using a TTP Mosquito X1 followed
by serial dilutions of compounds that were prepared using a Bravo automated liquid handling
platform (Agilent). Cells were treated, imaged, and analyzed according to the same protocol
described above. For the tertiary screen we used MEFs with a *Perk* promoter-driven dGFP as
previously described [118]. Cells were treated with drugs of interest and either DMSO or Tu at the
same time as described above and both GFP/cell and cell number were determined with
CellProfiler as previously described [3]. For both dose-response and *Perk* promoter-driven dGFP
experiments, eight biological replicates were performed per drug per dose.

Immunofluorescence

Immunofluorescence was performed as previously described [119]. Briefly, liver OCT sections or
cells cultured in chamber slide were fixed in cold acetone for 10 mins at -20 ºC. The sections were
then air dried, rehydrated with PBS and permeabilized with PBS+ 0.1% Triton X-100. The sections
were then blocked with 10% goat serum at room temperature for 1 hour. For mouse liver tissues,
primary antibodies against SC35/SRRM2 (ab11826, Abcam) were conjugated to Alexa-488,
respectively per manufacture's protocol and added to the OCT section at 1:1000 dilution overnight
at 4 ºC. Next day, sections were washed five times with PBS and counterstained with DAPI before
mounting (with ProlongGold Glass) and imaging using Leica SP8 lightening confocal microscope
(Leica Microsystems). For cell culture experiment, after incubation with anti-SRRM2 primary
antibody, Alex488-conjugated secondary antibodies were added and the rest was performed as
essentially the same way.

mRNA-seq and transcriptome analysis

For all mRNA-seq or RT-qPCR, total mRNA was isolated and purified from MEFs using the
PureLink RNA Mini Kit (Thermo Fisher). For mRNA-seq, samples were submitted to the UPMC
Genome Center for quality control, mRNA library preparation (Truseq Stranded mRNA (poly-A
pulldown), and sequencing (paired-end 101 bp reads and ~40 million reads per sample). The
sequencing was performed on a NextSeq 2000 sequencer.

For the *son* overexpression or knockdown (OE/KD) sequencing data, the raw RNA-seq FASTQ
files were analyzed by FastQC for quality control. Adaptors and low-quality reads were filtered by
Trimmomatic [120]. Then the processed reads were aligned by HISAT2 [121] against mouse reference
mm10. For gene-level intron/exon quantification, bedtools software [122] was used to collect and
count reads that aligned to any intron/exon of the given gene. If one read spans across multiple
exons of the same gene, it will only be counted once. If one read spans intron/exon junction, it
will only be counted as intron. The intron/exon count was normalized by gene length and total
reads for FPKM normalization.

The analysis of drug-treated samples was done using Galaxy [123]: Trim Galore was used for quality
control [124] and Salmon [125] was used to normalize the paired-end reads (TPM method). The online

3D RNA-Seq [126] pipeline was used to determine upregulated and downregulated genes, in
addition to generating the PCA plot.

For RNA-seq of *Rho*^{*P23H/+*} and wild-type mice, total RNA was isolated from retinae isolated from
mice at 1, 3, and 6 months of age using TRIzol organic extraction. RNA-seq was performed by
QuickBiology Inc. RNA integrity was checked by Agilent Bioanalyzer. Libraries for RNA-seq were
prepared with KAPA Stranded mRNA-Seq poly(A) selected kit (KAPA Biosystems, Wilmington,
MA) using 250 ng toal RNAs for each sample. Paired end sequencing was performed on Illumina
HighSeq 4000 (Illumina Inc., San Diego, CA).

The reads were first mapped to the latest UCSC transcript set using Bowtie2 version 2.1.0[127] and
the gene expression level was estimated using RSEM v1.2.15[128]. TMM (trimmed mean of M-
values) was used to normalize the gene expression. Differentially expressed genes were
identified using the edgeR program[129]. Genes showing altered expression with p < 0.05 and more
than 1.5-fold changes were considered differentially expressed. Goseq was uesd to perform the
GO enrichment analysis and Kobas was used to perform the pathway analysis.

mRNA splicing rates analysis

The mRNA processing rate was estimated by the simple kinetic model (equation 3) where pre-
mRNA was converted to mature mRNA with the mRNA processing rate *Kp* and the mature mRNA
is subject to decay with a constant decay rate *Kd*.

1031
$$Pre - mRNA \xrightarrow{Kp} mature\ mRNA \xrightarrow{Kd} \tag{3}$$

1032 Under the basal condition (DMSO), we assume a steady state of mature mRNA expression whose
1033 level does not change over time; thus, we have:

1034
$$\frac{d\ C(mature\ mRNA)}{dt} = Kp \times\ C(pre - mRNA) - C(mature\ mRNA) \times Kd = 0$$

1035
$$Kp = \frac{C\ (mature\ mRNA) \times Kd}{C\ (pre - mRNA)} \tag{4}$$

1036 Under *Son* overexpression or knocking down condition (condition 2), if we assume the mature
1037 mRNA degradation rate does not change with *Son* OE/KD (*K1*d=*K2*d), then the ratio of splicing
1038 rates between basal condition 1 and *Son* OE/KD condition 2 is given by:

1039
$$\frac{Kp1}{Kp2} = \frac{C1\ (mature\ mRNA) \times C2(pre - mRNA)}{C1\ (pre - mRNA) \times C2\ (mature\ mRNA)} \tag{5}$$

Gene set enrichment analysis (GSEA)

GSEA was performed with software version 4.1.0. TPM quantification of transcriptome under
different drugs or DMSO control was used as input for gene expression. Parameters used for the
analysis: 1000 permutations, permutation type: gene set.

Intron retention detection

Intron retention events were detected by iREAD [31]. Intron retention events are selected with
default settings T>=20, J>=1, FPKM>=2.

Gene ontology analysis

DAVID (Version 2021) [130] (https://david.ncifcrf.gov) was used to perform Gene Ontology analyses.
Briefly, gene names were first converted to DAVID-recognizable IDs using Gene Accession
Conversion Tool. The updated gene list was then subject to GO analysis using all Homo Sapiens
as background and with Functional Annotation Chart function. GO_BP_DIRECT,
KEGG_PATHWAY or UP_KW_BIOLOGICAL_PROCESS was used as GO categories. Only GO
terms with a p value less than 0.05 were included for further analysis.

Motif analysis

1055 Motif analysis was performed with the SeqPos motif tool (version 0.590) [131] embedded in Galaxy
1056 Cistrome using all motifs within the homo sapiens reference genome hg19 as background. LISA
1057 analysis was performed using webtool (http://lisa.cistrome.org/).

Statistical Analysis

1059 Data was analyzed and presented with GraphPad Prism software. Plots show individual data
1060 points and bars at the mean and ± the standard error of the mean (SEM). Individual tests used
1061 were provided at the end of each figure legend.

Author contributions

1063 Conceptualization, Y.C., X.C., and B.Z.; Methodology, W.D., B.B.C., S.L., S.P., Y.C., X.C., and
1064 B.Z.; Investigation, W.D., Y.T., S.K., S.Z., M.C., R.K.A., M.S., Y.N., M.Y., M.B.L., D.C., E.I., C.D.,
1065 H.W., S.L., S.P., B.B.C., Y.C., X.,C, and B.Z.; Writing – Original Draft, W.D., Y.C., X.C., and B.Z.;
1066 Writing – Review & Editing, all authors; Funding Acquisition, W.D., B.B.C., Y.C., X.C., and B.Z.;
1067 Resources, B.B.C. S.P., Y.C., X.C. and B.Z; Supervision, Y.C., X.C., and B.Z.

Acknowledgements

We would like to thank Drs. Yvonne Eisele, Yuan Liu and Toren Finkel (University of Pittsburgh
School of Medicine, Pittsburgh, PA, U.S.A.) for their technical assistance and/or comments on
the manuscript. We thank Dr. Krzysztof Palczewski who generously shared the 1D4 anti-
rhodopsin antibody. W.D. was supported by grant T32 HL082610 through the National Institutes
of Health (NIH), the Diana Jacobs Kalman/AFAR Scholarship for Research in the Biology of Aging
through the American Federation for Aging Research, and fellowship F31 AG080998. B.B.C. was
supported by grant 1R35HL139860 through the NIH. Y.C was supported by the R01 EY030991.
X.C. was supported by the National Institutes of Health grants R01AG074273 and R01AG078185.
B. Zhu was supported by grants 1DP2GM140924 and 1R21AG071893 through the NIH, and a
grant from Richard King Mellon foundation. This research was supported in part by the University
of Pittsburgh Center for Research Computing through the resources provided. Specifically, this
work used the HTC cluster, which is supported by NIH award number S10OD028483. This
research project was supported in part by the Pittsburgh Liver Research Centre supported by
NIH/NIDDK Digestive Disease Research Core Center grant P30DK120531, the Ophthalmology
and Visual Sciences Research Center core grant P30 EY08098, the Eye and Ear Foundation of
Pittsburgh and an unrestricted grant from Research to Prevent Blindness.

Declaration of interests

1086 B.B.C. is Co-founder of Koutif Therapeutic Inc., Co-founder and VP of Drug Discovery for
1087 Generian Pharmaceuticals, and Co-founder and C.S.O. of Coloma Therapeutics Inc..

Data availability

All raw and processed sequencing data generated in this study have been submitted to the NCBI Gene Expression Omnibus (GEO; http://www.ncbi.nlm.nih.gov/geo/) under accession numbers GSE224275. The mass spectrometry proteomics data have been deposited to the ProteomeXchange Consortium via the PRIDE partner repository with the dataset identifier PXD050371. All data needed to evaluate the conclusions in the paper are present in the paper and/or the Supplementary Materials.

References and Notes

1. Gestwicki JE, Garza D. Protein quality control in neurodegenerative disease. *Prog Mol Biol Transl Sci* **107**, 327-353 (2012).

2. Golde TE, Borchelt DR, Giasson BI, Lewis J. Thinking laterally about neurodegenerative proteinopathies. *J Clin Invest* **123**, 1847-1855 (2013).

3. Zhu B*, et al.* A Cell-Autonomous Mammalian 12 hr Clock Coordinates Metabolic and Stress Rhythms. *Cell Metab* **25**, 1305-1319 e1309 (2017).

4. Zhu B, Dacso CC, O'Malley BW. Unveiling "Musica Universalis" of the Cell: A Brief History of Biological 12-Hour Rhythms. *J Endocr Soc* **2**, 727-752 (2018).

5. Meng H*, et al.* XBP1 links the 12-hour clock to NAFLD and regulation of membrane fluidity and lipid homeostasis. *Nature Communications* **11**, 6215 (2020).

6. Pan Y*, et al.* 12-h clock regulation of genetic information flow by XBP1s. *PLOS Biology* **18**, e3000580 (2020).

7. Ballance H, Zhu B. Revealing the hidden reality of the mammalian 12-h ultradian rhythms. *Cellular and Molecular Life Sciences*, (2021).

8. Asher G, Zhu B. Beyond circadian rhythms: Emerging roles of ultradian rhythms in control of liver functions. *Hepatology* **n/a**, (2022).

9. Dion W*, et al.* Four-dimensional nuclear speckle phase separation dynamics regulate proteostasis. *Science Advances* **8**, eabl4150 (2022).

10. Meng H*, et al.* Defining the mammalian coactivation of hepatic 12-h clock and lipid metabolism. *Cell Reports* **38**, 110491 (2022).

11. Scott MR*, et al.* Twelve-hour rhythms in transcript expression within the human dorsolateral prefrontal cortex are altered in schizophrenia. *PLOS Biology* **21**, e3001688 (2023).

1130   12.   Zhu B, *et al.* Evidence for conservation of a primordial 12-hour ultradian gene program in
1131   humans. *bioRxiv*, 2023.2005.2002.539021 (2023).

1132
1133   13.   Zhu B, Liu S. Preservation of ~12-h ultradian rhythms of gene expression of mRNA and
1134   protein metabolism in the absence of canonical circadian clock. *Frontiers in Physiology* **14**,
1135   (2023).

1136
1137   14.   Spector DL, Lamond AI. Nuclear speckles. *Cold Spring Harb Perspect Biol* **3**, (2011).

1138
1139   15.   Hasenson SE, Shav-Tal Y. Speculating on the Roles of Nuclear Speckles: How RNA-
1140   Protein Nuclear Assemblies Affect Gene Expression. *Bioessays* **42**, e2000104 (2020).

1141
1142   16.   Liao SE, Regev O. Splicing at the phase-separated nuclear speckle interface: a model.
1143   *Nucleic Acids Research* **49**, 636-645 (2020).

1144
1145   17.   Alexander KA, *et al.* Nuclear speckles regulate HIF-2α programs and correlate with
1146   patient survival in kidney cancer. *bioRxiv*, 2023.2009.2014.557228 (2023).

1147
1148   18.   Alexander KA, *et al.* p53 mediates target gene association with nuclear speckles for
1149   amplified RNA expression. *Mol Cell* **81**, 1666-1681 e1666 (2021).

1150
1151   19.   Dion W, *et al.* Four-dimensional nuclear speckle phase separation dynamics regulate
1152   proteostasis. *Science Advances* **8**, eabl4150 (2022).

1153
1154   20.   Jia M, *et al.* Transcriptional changes of the aging lung. *Aging Cell* **22**, e13969 (2023).

1155
1156   21.   Chavez A, *et al.* Comparison of Cas9 activators in multiple species. *Nature methods* **13**,
1157   563-567 (2016).

1158
1159   22.   Qin Q, *et al.* Lisa: inferring transcriptional regulators through integrative modeling of
1160   public chromatin accessibility and ChIP-seq data. *Genome Biology* **21**, 32 (2020).

1161
1162   23.   Cretenet G, Le Clech M, Gachon F. Circadian clock-coordinated 12 Hr period rhythmic
1163   activation of the IRE1alpha pathway controls lipid metabolism in mouse liver. *Cell Metab* **11**,
1164   47-57 (2010).

1165
1166   24.   Pocaterra A, Romani P, Dupont S. YAP/TAZ functions and their regulation at a glance.
1167   *Journal of Cell Science* **133**, jcs230425 (2020).

1168
1169   25.   Panciera T, Azzolin L, Cordenonsi M, Piccolo S. Mechanobiology of YAP and TAZ in
1170   physiology and disease. *Nature Reviews Molecular Cell Biology* **18**, 758-770 (2017).

1171

1172 26.   Piccolo S, Dupont S, Cordenonsi M. The Biology of YAP/TAZ: Hippo Signaling and
1173   Beyond. *Physiological Reviews* **94**, 1287-1312 (2014).

1174
1175 27.   Reich S*, et al.* A multi-omics analysis reveals the unfolded protein response regulon and
1176   stress-induced resistance to folate-based antimetabolites. *Nature Communications* **11**, 2936
1177   (2020).

1178
1179 28.   Lu X, Ng H-H, Bubulya PA. The role of SON in splicing, development, and disease.
1180   *Wiley Interdiscip Rev RNA* **5**, 637-646 (2014).

1181
1182 29.   Ahn EY*, et al.* SON controls cell-cycle progression by coordinated regulation of RNA
1183   splicing. *Mol Cell* **42**, 185-198 (2011).

1184
1185 30.   Kim J-H*, et al.* SON drives oncogenic RNA splicing in glioblastoma by regulating
1186   PTBP1/PTBP2 switching and RBFOX2 activity. *Nature Communications* **12**, 5551 (2021).

1187
1188 31.   Li H-D, Funk CC, Price ND. iREAD: a tool for intron retention detection from RNA-seq
1189   data. *BMC Genomics* **21**, 128 (2020).

1190
1191 32.   Sureau A, Gattoni R, Dooghe Y, Stévenin J, Soret J. SC35 autoregulates its expression
1192   by promoting splicing events that destabilize its mRNAs. *Embo j* **20**, 1785-1796 (2001).

1193
1194 33.   Ding F, Su CJ, Edmonds KK, Liang G, Elowitz MB. Dynamics and functional roles of
1195   splicing factor autoregulation. *Cell Rep* **39**, 110985 (2022).

1196
1197 34.   Grandjean JMD*, et al.* Pharmacologic IRE1/XBP1s activation confers targeted ER
1198   proteostasis reprogramming. *Nat Chem Biol* **16**, 1052-1061 (2020).

1199
1200 35.   Madhavan A*, et al.* Pharmacologic IRE1/XBP1s activation promotes systemic adaptive
1201   remodeling in obesity. *Nature Communications* **13**, 608 (2022).

1202
1203 36.   Grandjean JMD, Wiseman RL. Small molecule strategies to harness the unfolded
1204   protein response: where do we go from here? *Journal of Biological Chemistry* **295**, 15692-
1205   15711 (2020).

1206
1207 37.   Schor IE*, et al.* Perturbation of chromatin structure globally affects localization and
1208   recruitment of splicing factors. *PloS one* **7**, e48084 (2012).

1209
1210 38.   Subramanian A*, et al.* Gene set enrichment analysis: A knowledge-based approach for
1211   interpreting genome-wide expression profiles. *Proceedings of the National Academy of
1212   Sciences* **102**, 15545-15550 (2005).

1213

39. Molina DM, *et al.* Monitoring Drug Target Engagement in Cells and Tissues Using the Cellular Thermal Shift Assay. *Science* **341**, 84-87 (2013).

40. Jafari R, *et al.* The cellular thermal shift assay for evaluating drug target interactions in cells. *Nature Protocols* **9**, 2100-2122 (2014).

41. Ilik İA, Malszycki M, Lübke AK, Schade C, Meierhofer D, Aktaş T. SON and SRRM2 are essential for nuclear speckle formation. *eLife* **9**, e60579 (2020).

42. Fei J, *et al.* Quantitative analysis of multilayer organization of proteins and RNA in nuclear speckles at super resolution. *Journal of Cell Science* **130**, 4180-4192 (2017).

43. Sharma A, Takata H, Shibahara K, Bubulya A, Bubulya PA. Son is essential for nuclear speckle organization and cell cycle progression. *Mol Biol Cell* **21**, 650-663 (2010).

44. Liao SE, Regev O. Splicing at the phase-separated nuclear speckle interface: a model. *Nucleic Acids Res* **49**, 636-645 (2021).

45. Xu S, Lai SK, Sim DY, Ang WSL, Li HY, Roca X. SRRM2 organizes splicing condensates to regulate alternative splicing. *Nucleic Acids Res* **50**, 8599-8614 (2022).

46. Gouveia B, Kim Y, Shaevitz JW, Petry S, Stone HA, Brangwynne CP. Capillary forces generated by biomolecular condensates. *Nature* **609**, 255-264 (2022).

47. Sabari BR, *et al.* Coactivator condensation at super-enhancers links phase separation and gene control. *Science* **361**, eaar3958 (2018).

48. Chen Z, *et al.* Screening membraneless organelle participants with machine-learning models that integrate multimodal features. *Proceedings of the National Academy of Sciences* **119**, e2115369119 (2022).

49. Romero, Obradovic, Dunker K. Sequence Data Analysis for Long Disordered Regions Prediction in the Calcineurin Family. *Genome Inform Ser Workshop Genome Inform* **8**, 110-124 (1997).

50. Walsh I, Martin AJM, Di Domenico T, Tosatto SCE. ESpritz: accurate and fast prediction of protein disorder. *Bioinformatics* **28**, 503-509 (2011).

51. Santos J, Calero N, Trujillo-Cayado LA, Garcia MC, Muñoz J. Assessing differences between Ostwald ripening and coalescence by rheology, laser diffraction and multiple light scattering. *Colloids and Surfaces B: Biointerfaces* **159**, 405-411 (2017).

52.   Dopie J, Sweredoski MJ, Moradian A, Belmont AS. Tyramide signal amplification mass spectrometry (TSA-MS) ratio identifies nuclear speckle proteins. *Journal of Cell Biology* **219**, (2020).

53.   Saitoh N, Spahr CS, Patterson SD, Bubulya P, Neuwald AF, Spector DL. Proteomic analysis of interchromatin granule clusters. *Mol Biol Cell* **15**, 3876-3890 (2004).

54.   Riback JA*, et al.* Viscoelasticity and advective flow of RNA underlies nucleolar form and function. *Molecular Cell* **83**, 3095-3107.e3099 (2023).

55.   Pancholi A*, et al.* RNA polymerase II clusters form in line with liquid phase wetting of chromatin. *bioRxiv*, 2021.2002.2003.429626 (2021).

56.   Liu X*, et al.* Time-dependent effect of 1,6-hexanediol on biomolecular condensates and 3D chromatin organization. *Genome Biology* **22**, 230 (2021).

57.   Liu J, Rivas FV, Wohlschlegel J, Yates JR, 3rd, Parker R, Hannon GJ. A role for the P-body component GW182 in microRNA function. *Nat Cell Biol* **7**, 1261-1266 (2005).

58.   Ravi V, Jain A, Mishra S, Sundaresan NR. Measuring Protein Synthesis in Cultured Cells and Mouse Tissues Using the Non-radioactive SUnSET Assay. *Curr Protoc Mol Biol* **133**, e127 (2020).

59.   Pakos-Zebrucka K, Koryga I, Mnich K, Ljujic M, Samali A, Gorman AM. The integrated stress response. *EMBO Rep* **17**, 1374-1395 (2016).

60.   Mauvezin C, Neufeld TP. Bafilomycin A1 disrupts autophagic flux by inhibiting both V-ATPase-dependent acidification and Ca-P60A/SERCA-dependent autophagosome-lysosome fusion. *Autophagy* **11**, 1437-1438 (2015).

61.   Harada Y, Ishii I, Hatake K, Kasahara T. Pyrvinium pamoate inhibits proliferation of myeloma/erythroleukemia cells by suppressing mitochondrial respiratory complex I and STAT3. *Cancer Letters* **319**, 83-88 (2012).

62.   Ishii I, Harada Y, Kasahara T. Reprofiling a classical anthelmintic, pyrvinium pamoate, as an anti-cancer drug targeting mitochondrial respiration. *Frontiers in Oncology* **2**, (2012).

63.   Bayer TA. Proteinopathies, a core concept for understanding and ultimately treating degenerative disorders? *Eur Neuropsychopharmacol* **25**, 713-724 (2015).

64.   Hartong DT, Berson EL, Dryja TP. Retinitis pigmentosa. *Lancet* **368**, 1795-1809 (2006).

1298   65.   Lee E-J, Chan P, Chea L, Kim K, Kaufman RJ, Lin JH. ATF6 is required for efficient
1299   rhodopsin clearance and retinal homeostasis in the P23H rho retinitis pigmentosa mouse
1300   model. *Scientific Reports* **11**, 16356 (2021).

1301

1302   66.   Vats A*, et al.* Nonretinoid chaperones improve rhodopsin homeostasis in a mouse model
1303   of retinitis pigmentosa. *JCI Insight* **7**, (2022).

1304

1305   67.   Chiang W-C*, et al.* Robust Endoplasmic Reticulum-Associated Degradation of
1306   Rhodopsin Precedes Retinal Degeneration. *Molecular Neurobiology* **52**, 679-695 (2015).

1307

1308   68.   Chiang W-C, Hiramatsu N, Messah C, Kroeger H, Lin JH. Selective Activation of ATF6
1309   and PERK Endoplasmic Reticulum Stress Signaling Pathways Prevent Mutant Rhodopsin
1310   Accumulation. *Investigative Ophthalmology & Visual Science* **53**, 7159-7166 (2012).

1311

1312   69.   Lobanova ES*, et al.* Increased proteasomal activity supports photoreceptor survival in
1313   inherited retinal degeneration. *Nat Commun* **9**, 1738 (2018).

1314

1315   70.   Liu X*, et al.* Pharmacological clearance of misfolded rhodopsin for the treatment of RHO-
1316   associated retinitis pigmentosa. *The FASEB Journal* **34**, 10146-10167 (2020).

1317

1318   71.   Lee MJ, Lee JH, Rubinsztein DC. Tau degradation: The ubiquitin–proteasome system
1319   versus the autophagy-lysosome system. *Progress in Neurobiology* **105**, 49-59 (2013).

1320

1321   72.   Limanaqi F, Biagioni F, Gambardella S, Familiari P, Frati A, Fornai F. Promiscuous
1322   Roles of Autophagy and Proteasome in Neurodegenerative Proteinopathies. *Int J Mol Sci* **21**,
1323   (2020).

1324

1325   73.   Samelson AJ*, et al.* CRISPR screens in iPSC-derived neurons reveal principles of tau
1326   proteostasis. *bioRxiv*, 2023.006.2016.545386 (2023).

1327

1328   74.   Bravo CP*, et al.* Human iPSC 4R tauopathy model uncovers modifiers of tau
1329   propagation. *bioRxiv*, 2023.006.2019.544278 (2023).

1330

1331   75.   Bugiani O*, et al.* Frontotemporal dementia and corticobasal degeneration in a family with
1332   a P301S mutation in tau. *Journal of neuropathology and experimental neurology* **58**, 667-677
1333   (1999).

1334

1335   76.   Chen X*, et al.* Promoting tau secretion and propagation by hyperactive p300/CBP via
1336   autophagy-lysosomal pathway in tauopathy. *Molecular Neurodegeneration* **15**, 2 (2020).

1337

1338   77.   Triastuti E*, et al.* Pharmacological inhibition of Hippo pathway, with the novel kinase
1339   inhibitor XMU-MP-1, protects the heart against adverse effects during pressure overload. *Br J*
1340   *Pharmacol* **176**, 3956-3971 (2019).

78.	Hao X*, et al.* XMU-MP-1 attenuates osteoarthritis via inhibiting cartilage degradation and chondrocyte apoptosis. *Frontiers in Bioengineering and Biotechnology* **10**, (2022).

79.	Kastan N*, et al.* Small-molecule inhibition of Lats kinases may promote Yap-dependent proliferation in postmitotic mammalian tissues. *Nature Communications* **12**, 3100 (2021).

80.	Qin F, Tian J, Zhou D, Chen L. Mst1 and Mst2 kinases: regulations and diseases. *Cell & Bioscience* **3**, 31 (2013).

81.	Arno G*, et al.* Mutations in REEP6 Cause Autosomal-Recessive Retinitis Pigmentosa. *Am J Hum Genet* **99**, 1305-1315 (2016).

82.	Pi S*, et al.* Fully automated OCT-based tissue screening system. *Opt Lett* **49**, 4481-4484 (2024).

83.	Wittmann CW*, et al.* Tauopathy in <i>Drosophila</i>: Neurodegeneration Without Neurofibrillary Tangles. *Science* **293**, 711-714 (2001).

84.	Berger C, Renner S, Lüer K, Technau GM. The commonly used marker ELAV is transiently expressed in neuroblasts and glial cells in the Drosophila embryonic CNS. *Developmental Dynamics* **236**, 3562-3568 (2007).

85.	Hirabayashi S, Baranski TJ, Cagan RL. Transformed Drosophila cells evade diet-mediated insulin resistance through wingless signaling. *Cell* **154**, 664-675 (2013).

86.	Wang L*, et al.* JNK modifies neuronal metabolism to promote proteostasis and longevity. *Aging Cell* **18**, e12849 (2019).

87.	Xu M*, et al.* A systematic integrated analysis of brain expression profiles reveals YAP1 and other prioritized hub genes as important upstream regulators in Alzheimer's disease. *Alzheimers Dement* **14**, 215-229 (2018).

88.	Mathys H*, et al.* Single-cell transcriptomic analysis of Alzheimer's disease. *Nature* **570**, 332-337 (2019).

89.	Wang C*, et al.* Scalable Production of iPSC-Derived Human Neurons to Identify Tau-Lowering Compounds by High-Content Screening. *Stem Cell Reports* **9**, 1221-1233 (2017).

90.	Wu R*, et al.* Disruption of nuclear speckle integrity dysregulates RNA splicing in C9ORF72-FTD/ALS. *Neuron*, (2024).

91. Lester E, *et al.* Tau aggregates are RNA-protein assemblies that mislocalize multiple nuclear speckle components. *Neuron* **109**, 1675-1691.e1679 (2021).

92. McMillan PJ, *et al.* Pathological tau drives ectopic nuclear speckle scaffold protein SRRM2 accumulation in neuron cytoplasm in Alzheimer's disease. *Acta Neuropathologica Communications* **9**, 117 (2021).

93. Mordes D, *et al.* Pre-mRNA splicing and retinitis pigmentosa. *Mol Vis* **12**, 1259-1271 (2006).

94. Lester E, *et al.* Tau aggregates are RNA-protein assemblies that mislocalize multiple nuclear speckle components. *Neuron* **109**, 1675-1691 e1679 (2021).

95. Tracy TE, *et al.* Tau interactome maps synaptic and mitochondrial processes associated with neurodegeneration. *Cell* **185**, 712-728.e714 (2022).

96. Hsieh YC, *et al.* Tau-Mediated Disruption of the Spliceosome Triggers Cryptic RNA Splicing and Neurodegeneration in Alzheimer's Disease. *Cell Rep* **29**, 301-316 e310 (2019).

97. Wheeler RJ. Therapeutics-how to treat phase separation-associated diseases. *Emerg Top Life Sci* **4**, 307-318 (2020).

98. Conti BA, Oppikofer M. Biomolecular condensates: new opportunities for drug discovery and RNA therapeutics. *Trends in Pharmacological Sciences* **43**, 820-837 (2022).

99. Kilgore HR, *et al.* Distinct chemical environments in biomolecular condensates. *Nature Chemical Biology*, (2023).

100. Schultz CW, Nevler A. Pyrvinium Pamoate: Past, Present, and Future as an Anti-Cancer Drug. *Biomedicines* **10**, 3249 (2022).

101. Jawerth LM, *et al.* Salt-Dependent Rheology and Surface Tension of Protein Condensates Using Optical Traps. *Physical Review Letters* **121**, 258101 (2018).

102. Spruijt E, Sprakel J, Cohen Stuart MA, van der Gucht J. Interfacial tension between a complex coacervate phase and its coexisting aqueous phase. *Soft Matter* **6**, 172-178 (2010).

103. Cai D, Sukenik S, Feliciano D, Gruebele M, Lippincott-Schwartz J. Phase Separation of YAP Reprograms Cells for Long-term YAP Target Gene Expression. *bioRxiv*, 438416 (2018).

104. Yang S, *et al.* Bip-Yorkie interaction determines oncogenic and tumor-suppressive roles of Ire1/Xbp1s activation. *Proceedings of the National Academy of Sciences* **119**, e2202133119 (2022).

105. Rivera-Reyes A*, et al.* YAP1 enhances NF-κB-dependent and independent effects on clock-mediated unfolded protein responses and autophagy in sarcoma. *Cell Death Dis* **9**, 1108 (2018).

106. Min Y*, et al.* Cross species systems biology discovers glial DDR2, STOM, and KANK2 as therapeutic targets in progressive supranuclear palsy. *Nature Communications* **14**, 6801 (2023).

107. Vallazza-Deschamps G*, et al.* Excessive activation of cyclic nucleotide-gated channels contributes to neuronal degeneration of photoreceptors. *Eur J Neurosci* **22**, 1013-1022 (2005).

108. Vats A*, et al.* Nonretinoid chaperones improve rhodopsin homeostasis in a mouse model of retinitis pigmentosa. *JCI Insight* **7**, (2022).

109. Pi S*, et al.* Retinal capillary oximetry with visible light optical coherence tomography. *Proceedings of the National Academy of Sciences of the United States of America* **117**, 11658-11666 (2020).

110. Fanning S, Selkoe D, Dettmer U. Parkinson's disease: proteinopathy or lipidopathy? *NPJ Parkinsons Dis* **6**, 3 (2020).

111. Wulansari N*, et al.* Neurodevelopmental defects and neurodegenerative phenotypes in human brain organoids carrying Parkinson's disease-linked DNAJC6 mutations. *Sci Adv* **7**, (2021).

112. Dion W*, et al.* Four-dimensional nuclear speckle phase separation dynamics regulate proteostasis. *Sci Adv* **8**, eabl4150 (2022).

113. Dupont S*, et al.* Role of YAP/TAZ in mechanotransduction. *Nature* **474**, 179-183 (2011).

114. Stirling DR, Swain-Bowden MJ, Lucas AM, Carpenter AE, Cimini BA, Goodman A. CellProfiler 4: improvements in speed, utility and usability. *BMC bioinformatics* **22**, 1-11 (2021).

115. Liu X*, et al.* Pharmacological clearance of misfolded rhodopsin for the treatment of RHO-associated retinitis pigmentosa. *Faseb j* **34**, 10146-10167 (2020).

116. Schneider CA, Rasband WS, Eliceiri KW. NIH Image to ImageJ: 25 years of image analysis. *Nature methods* **9**, 671-675 (2012).

117. Zhu B*, et al.* Coactivator-Dependent Oscillation of Chromatin Accessibility Dictates Circadian Gene Amplitude via REV-ERB Loading. *Mol Cell* **60**, 769-783 (2015).

1469 118.   Zhu B*, et al.* A Cell-Autonomous Mammalian 12 hr Clock Coordinates Metabolic and
1470   Stress Rhythms. *Cell Metabolism* **25**, 1305-1319.e1309 (2017).

1471
1472 119.   Zhu B*, et al.* Peroxisome proliferator-activated receptor beta/delta cross talks with E2F
1473   and attenuates mitosis in HRAS-expressing cells. *Mol Cell Biol* **32**, 2065-2082 (2012).

1474
1475 120.   Bolger AM, Lohse M, Usadel B. Trimmomatic: a flexible trimmer for Illumina sequence
1476   data. *Bioinformatics* **30**, 2114-2120 (2014).

1477
1478 121.   Kim D, Langmead B, Salzberg SL. HISAT: a fast spliced aligner with low memory
1479   requirements. *Nat Methods* **12**, 357-360 (2015).

1480
1481 122.   Quinlan AR, Hall IM. BEDTools: a flexible suite of utilities for comparing genomic
1482   features. *Bioinformatics* **26**, 841-842 (2010).

1483
1484 123.   Blankenberg D*, et al.* Galaxy: A Web-Based Genome Analysis Tool for Experimentalists.
1485   *Current Protocols in Molecular Biology* **89**, 19.10.11-19.10.21 (2010).

1486
1487 124.   Krueger F. Trim Galore.) (2021).

1488
1489 125.   Patro R, Duggal G, Love MI, Irizarry RA, Kingsford C. Salmon provides fast and bias-
1490   aware quantification of transcript expression. *Nature methods* **14**, 417-419 (2017).

1491
1492 126.   Guo W*, et al.* 3D RNA-seq: a powerful and flexible tool for rapid and accurate differential
1493   expression and alternative splicing analysis of RNA-seq data for biologists. *RNA biology* **18**,
1494   1574-1587 (2021).

1495
1496 127.   Langmead B, Salzberg SL. Fast gapped-read alignment with Bowtie 2. *Nat Methods* **9**,
1497   357-359 (2012).

1498
1499 128.   Li B, Dewey CN. RSEM: accurate transcript quantification from RNA-Seq data with or
1500   without a reference genome. *BMC Bioinformatics* **12**, 323 (2011).

1501
1502 129.   Robinson MD, McCarthy DJ, Smyth GK. edgeR: a Bioconductor package for differential
1503   expression analysis of digital gene expression data. *Bioinformatics* **26**, 139-140 (2010).

1504
1505 130.   Huang da W, Sherman BT, Lempicki RA. Systematic and integrative analysis of large
1506   gene lists using DAVID bioinformatics resources. *Nat Protoc* **4**, 44-57 (2009).

1507
1508 131.   He HH*, et al.* Nucleosome dynamics define transcriptional enhancers. *Nature Genetics*
1509   **42**, 343-347 (2010).

1510

1511 132. Zhang MJ, Pisco AO, Darmanis S, Zou J. Mouse aging cell atlas analysis reveals global
1512 and cell type-specific aging signatures. *Elife* **10**, (2021).

1513
1514 133. A single-cell transcriptomic atlas characterizes ageing tissues in the mouse. *Nature* **583**,
1515 590-595 (2020).

1516
1517 134. Jiang J, Wang C, Qi R, Fu H, Ma Q. scREAD: A Single-Cell RNA-Seq Database for
1518 Alzheimer's Disease. *iScience* **23**, 101769 (2020).

1519

1520
1521

1522

1523

1524 **<u>Fig. legends</u>**

1525

Fig. 1. Genetic rejuvenation of nuclear speckle transcriptionally reprograms global proteostasis and YAP1 signaling in an opposing manner. (**a**) The diagram of the approach of

nuclear speckle rejuvenation to alleviate proteinopathy. (**b**) PCA of global transcriptional response

to Tu in the presence of SON OE (via CRISPRa) or KD (via siRNA). (**c, d**) Relative expression

(R.E.) of representative proteostasis genes (**c**) and YAP1 target genes (**d**) in response to Tu in

the presence of SON OE/KD (n=4 for SON KD, and n=3 for SON OE). (**e**) Heat map of relative

fold change of gene expression by Tu in SON OE/KD cells compared to control cells. Only those

genes with at least 1.4-fold induction by Tu ($\log_2 > 0.5$) with a p value smaller than 0.05 in control

condition are included. Among these genes are 461 genes whose induction by Tu are further

amplified by SON OE (induced more) and dampened (induced less) *by* SON KD; and 901 genes

whose repression by Tu are further amplified by *Son* OE (repressed further) and dampened

(repressed less) by SON KD. (**f**) GO analysis of those 461 and 901 genes showing enriched

KEGG pathways. (**g**) Enriched XBP1s binding motif ACGTCA on the promoters of 461 genes. (**h**)

Top enriched GO terms for top 500 most abundant proteins that are detected in hepatic XBP1s

interactome at CT8, as previously reported [10]. (**i**) Heatmap showing relative abundance of 45

proteins involved in mRNA splicing and processing within the XBP1s interactome at CT0 and

CT8, respectively. (**j**) Enriched TEAD2 binding motif GGCGG on the promoters of 901 genes.

Western blot (**k**) and quantification (**l**) of nuclear and cytosolic level of YAP1 in control and SON

OE cells in response to Tu (n=2). Scratch assay with representative images (**m**) and quantification

(**n**) of cell migration rate in control and SON OE cells in response to Tu (n=5). All data mean ±

standard error of the mean (S.E.M.). Statistical tests used: unpaired one-tailed Student's t-test for

c, d, l, and n. Paired one-tailed Student's t-test for i.

Fig. 2. HTS identifies pyrvinium as a SON-dependent nuclear speckle rejuvenator. (**a**)
Workflow detailing our initial drug screen and subsequent steps to identify compounds that affect
the nuclear speckle morphology in a dose-dependent manner and amplifies the UPR. (**b**) Dose-

dependent effect on nuclear speckles morphology by PP, with a representative image of nuclear
speckles under DMSO or 0.1μM of PP (n=25~57). (**c**) Quantification of total area-normalized
perimeter of nuclear speckles in control and 1μM PP condition per cell (n=26 for DMSO and n=35
for PP). (**d**) GSEA analysis showing a similar transcriptome signature between PP-upregulated
genes and 461 genes further amplified by SON OE during ER stress. (**e**) GSEA analysis showing
a similar transcriptome signature between PP-downregulated genes and 901 genes further
blunted by SON OE during ER stress. (**f**) Volcano plot showing fold change by PP versus log
transformed p values. Genes induced or repressed by at least 1.41-fold with a p value smaller
than 0.05 are boxed. (**g**) GO analysis of differentially expressed genes by PP. (**h**) Representative
images and quantification of sphericity of GFP signal from GFP::SRSF2 MEFs with scrambled or
Son siRNA treated with DMSO or increasing concentration of PP for 25 hours (n=31~81). (**i**) $\log_2$
normalized fold change in response to PP treatment (0.3 μm) for 24 hours in control and SON KD
MEFs with p values shown for one tailed t-test (n=3). All data mean ± S.E.M. Statistical tests used:
unpaired one-tailed Student's t-test for c and i. Ordinary one-way ANOVA for b and h.

1566

1567

1568

1569

1570

1571

1572

1573

1574

1575

1576

1577

1578

1579

1580

1581

1582

1583

1584

1585

1586

Fig. 3. PP targets SON to reduce the interfacial tension of nuclear speckles. CETSA of SON (**a**) and GFP::SRSF2 (**b**) with 3µM PP. Both representative blot and quantification from independent replicates are shown. (SON: n=6 for DMSO and n=5 for PP; GFP::SRSF2: n=3 for both DMSO and PP). (**c**) Computational prediction of IDR in mouse SON. (**d**) Diagram illustrating how surface tension influences droplets coalescence kinetics. (**e, f**) Representative images of droplet formation assay with different recombinant proteins (**e**) and quantification (**f**) of area-normalized perimeter changes in the time span of 20 minutes with 125mM NaCl (n=2~3). (**g-i**) Representative images of droplet formation with NE-supplemented SON IDR2 with increasing concentration of PP (**g**) and quantification of sphericity (n=12 for with NE and n=42 for without NE) (**h**) and area-normalized perimeter changes (n=2) (**i**) in the time span of 20 minutes. (**j-m**) Diagram of droplet formation assay where SON IDR2 is expected to compartmentalize splicing factors, including GFP::SRSF2 into the nuclear speckle-like condensates. GFP::SRSF2 is expected to exhibit a broader spatial distribution than the SON IDR2 core (**j**). Representative images (**k**) and quantification of sphericity (**l**) and spatial distribution of mCherry::SON IDR2 and GFP::SRSF2 (**m**). (**n, o**) Mouse genomic DNA was further added to the solution. Representative images (**n**) and quantification (**o**) of spatial distribution of mCherry::SON IDR2, GFP::SRSF2 and DNA. All data mean ± S.E.M. Statistical tests used: mixed-effects analysis for a, b and m. Ordinary one-way ANOVA for f, h, I and l. Two-way RM ANOVA for o.

1609

Fig. 4. PP reduces pathological Tau and Rhodopsin level by boosting autophagy and UPS

activity. (**a-d**) MEFs were treated with DMSO or 0.1μM PP for ~24 hours and then co-treated

with or without puromycin (10 μg/mL for 30 minutes), or MG132 (10μM for 110 minutes) (n=4 for

all samples). Western blot and quantification of puromycin-incorporated proteins (**a, b**), and poly-

ubiquitinated protein (**c, d**). (**e, f**) MEFs were treated with DMSO or 0.1μM PP for ~24 hours and

autophagic flux was measured via a mCherry::GFP::LC3 fusion protein reporter. Chimeric

proteins comprising LC3B fused with both GFP and mCherry offer a method to track autophagic

flux. Autophagosomes (APG) marked by mCherry::GFP::LC3 exhibit both mCherry and GFP

signals. Following fusion with lysosome to form autolysosome (AL), GFP signals diminish
significantly in the acidic environment, while mCherry signals remain relatively stable. (**e**).
Quantification of the number of APG, AL and total vesicles (**f**). n=23 for DMSO and 29 for PP. (**g,**
h) NIH3T3 RHO^{P23H} cells were treated with 0.1µM PP for 24 hours and Western blot (**g**) and
quantification (**h**) of RHO^{P23H} level (n=3). (**i, j**) 0.1µM PP-treated NIH3T3 RHO^{P23H} cells were co-
treated with or without 1µM SBI-0206965 for 24 hours. Western blot (**i**) and quantification (**j**) of
RHO^{P23H} level (n=3). (**k-m**) Tau (P301S)-expressing primary neurons were treated with increasing
concentration of PP for 24 hours, and western blot and quantification of different proteins (n=4 for
HT7, PHF1 and LC3, and n=6 for P62) (**k**) or treated with 0.1µM PP for 12h hours in the presence
or absence of BafA (50nM) in the last hour and western blot (**l**) and quantification (n=4 for total
Tau and P62 and n=8 for p-Tau) (**m**) of different proteins. (**n**) Tau P301S-expressing neurons
were treated with DMSO or 0.1µM PP for 12 hours and qPCR of selective proteostasis and YAP1
target genes (n=6~11). (**o**) MEFs were treated with DMSO, PP (1µM), XMU-MP-1 (1µM) or XMU-
MP-1+PP for 24 hours, and qPCR of protein quality control and YAP1s output gene expression
quantified as log transformed fold change under DMSO or XMU-MP-1 condition by PP (n=3). All
data mean ± S.E.M. Statistical tests used: unpaired one-tailed Student's t-test for b, c, f, h, j, m, n
and o. Ordinary one-way ANOVA for k.

1635

1636

1637

1638

1639

1640

1641

1642

1643

1644

1645

1646

1647

1648

1649

1650

1651

1652

1653

1654

Fig. 5. PP alleviates proteinopathies in preclinical models. (**a-e**) RNA-seq was performed in the retina of 1 and 3 months-old wild-type and Rho [P23H/+] mice. GO of differentially expressed genes (DEG) (FDR<0.1) (**a, c**) and GSEA comparing these DEG with that of PP (**b, d**). Heatmap of selective genes (**e**). (**f-j**) Retina explants isolated either from *Rho*[P23H/+] mice P15 and cultured with PP or DMSO vehicle control or from wild-type mice cultured for 10 days *ex vivo*. (**f**) The morphology retinae were imaged and scanned before (Day 0) and after treatment (Day 10) by a webcam (top) and visible light optical coherence tomography (vis-OCT) with tissue thickness

shown as a heatmap with a color legend indicating thickness from 0-300 μm (bottom). Scale, 1
mm. (**g**) and (**h**) are bar plots of retinal thickness and volume, respectively, measured from the
vis-OCT scanning data. n=4~5. (**i**) Representative retinal histology images of the retina explants
cultured for 10 days with black bars showing the outer nuclear layer (ONL). (**j**) The nuclei count
in the outer nuclei layer (ONL) along six horizontal positions at peripheral-central-peripheral
positions across each cross-section in (**i**). n=3-4. (**k-n**) Male C155>UAS-hTau1.13 and C155 flies
were fed with either standard diet or diet supplemented with 25μM PP. Quantification of distance
travelled from larval crawling assay (n=10) (**k**), climbing index score from adult fly climbing assay
at 14 days and 21 days (n=5) (**l**) of age, and lifespan assay (**m**) (n=32 for C155-Ctrl, n=65 for
hTau-Ctrl and n=44 for hTau-PP). (**n**) Western blot and quantification of p-Tau (AT8 antibody)
and total Tau (DAKO antibody) level in 21 days C155>UAS-hTau1.13 flies fed with control or PP
diet n=4. All data mean ± S.E.M. Statistical tests used: two-way ANOVA and Turkey multiple
comparison for g and h, two-way ANOVA for j, unpaired one-tailed Student's t-test for k, l (day
21) and n. Log-rank (Mantel-Cox) test for m.

1676

1677

1678

1679

1680

1681

1682

1683

1684

1685

1686

1687

1688

1689

1690

1691

Fig. 6. PP rejuvenates nuclear speckles and alleviates tau burden in human iPSC-neurons expressing mutant Tau. (**a**) GO analysis of up and downregulated genes in 4R-Tau P301S iPSC neurons reported in [74]. (**b**) GSEA comparing genes upregulated in 4R-Tau P301S iPSC neurons with those downregulated by PP. (**c**) Motif analysis of promoters of genes upregulated in 4R-Tau

P301S iPSC neurons compared to 4R-Tau (FDR<0.05) revealed top enriched motif of TEAD2. (**d, e**) Wild-type and V337M Tau-expressing iPSC-neurons were treated with DMSO or PP (10nM) for 12 hours, and IF against nuclear speckle (Ab11826 against SRRM2), p-Tau (Ser422) and chromatin (DAPI) were performed. Representative images (**d**) and quantification of nuclear speckle sphericity and area, intensity of nuclear, cytosol and total level of p-Tau (Ser422) and ratio of cytosol versus nuclear level of p-Tau (**e**). (**f, g**) V337M Tau-expressing iPSC-neurons were treated with DMSO or increasing concentrations of PP for 12 hours. Representative western blot (**f**) and quantification (**g**) of total and p-Tau (Ser202/Thr205) (n=3). (**h**) V337M Tau-expressing iPSC-neurons were treated with DMSO or 100nM PP for 12 hours, and autophagy flux was quantified by the autophagy reporter. (**i-k**) Wild-type and V337M Tau-expressing iPSC-neurons were infected with scrambled shRNA or SON shRNA-encoding lentivirus and treated with DMSO or PP (100nM) for 12 hours, and IF against nuclear speckle (Ab11826 against SRRM2), p-Tau (Ser422) and chromatin (DAPI) were performed. Quantification of nuclear speckle area (**i**) sphericity (**j**), intensity of total level of p-Tau (Ser422) (**k**) were shown. Data: Mean ± SEM.

Supplementary Figures.

1731

Supplementary Figure 1. SON expression is reduced during aging in mice and humans. (a)
Normalized counts of *Son* mRNA in different mouse tissues with different ages (WBC; White blood

cells) according to Tabula Muris Senis database (n=3~6) [132, 133]. (**b**) qPCR of *Son* level in liver of
mice with different ages (n=3~7). (**c, d**) Western blot (**c**) and quantification (**d**) of SON level in
liver of mice with different ages (n=5~6). (**e, f**) Representative immunofluorescence images of
SRRM2 and DAPI in the liver of 3 and 22 months old male mice (**e**) and quantification of the
sphericity of nuclear speckles (**f**). n=25 for both age groups. (**g**) Single cell RNA-seq data of *SON*
mRNA level in different cell types in lung tissues of young and aging humans, reported from [20].
(**h**) Counts per million normalized expressions of *SON* in brains of human AD compiled from RNA-
seq data from the AMP-AD consortium. (**i**) Heat map showing relative expression of *SRSF2* and
SON in different cell types in the cortex of human AD subjects [134]. Blue color indicates lower level
in AD subjects compared to controls. Data: Mean ± S.E.M. Statistical tests used: two-way ANOVA
and Turkey multiple comparison for h, unpaired one-tailed Student's t-test for a, b, d, and f.
Wilcoxon test for g.

1746

Supplementary Figure 2. Genetic rejuvenation of nuclear speckles by SON. (**a**) Relative
expression (R.E.) of representative proteostasis genes (top) and YAP1 target genes (bottom) in

response to Tu in the presence of SON OE/KD (n=4 for SON KD and n=3 for SON OE). (**b**)
Scatter plot showing relative fold change by *Son* KD versus SON OE for both Tu-induced and Tu-
repressed genes. (**c**) Top predicted transcription regulators of 461 and 901 genes by the LISA
Cistrome DB TR ChIP-Seq models. (**d, e**) Selected genes aligned for SC35 and XBP1s ChIP-seq
signal from CT12 in XBP1*Flox* mice [6] (**d**) and ChIP-qPCR of XBP1s and SC35 on selected regions
(indicated by red bars) (n=2) (**e**). Data: Mean ± S.E.M. Statistical tests used: unpaired one-tailed
Student's t-test for a and e. Linear regression for b.

1756

1757

1758

1759

1760

1761

1762

1763

1764

1765

1766

Supplementary Figure 3. YAP1 transcriptional output is repressed during ER stress. (**a**) TEAD luciferase reporter assay in control and SON OE MEFs in response to Tu (n=12 for all groups). (**b**) Heatmap showing transcriptomes that are significantly downregulated (log $_2$-fold change smaller than -0.5) with a p value less than 0.05) either under Tu or Thap treatment at 6h. (**c**) GO analysis of genes that are significantly downregulated either under Tu or Thap treatment at 6h. (**d**) Heatmap of representative YAP1 target genes as in **b**. Data: Mean ± S.E.M. Statistical tests used: unpaired one-tailed Student's t-test for a.

1774

1775

Supplementary Figure 4. SON regulates mRNA splicing rates under both basal and ER stress conditions (**a**) PCA analysis of global transcriptional response to Tu in the presence of SON OE/KD. Both pre-mRNA (top) and mature mRNA (bottom) are shown. (**b, c**) Volcano plot of mRNA splicing rates changes in SON KD (**b**) or OE (**c**) MEFs under basal condition (DMSO). (**d**) Heat map of fold change of RNA splicing rate, pre and mature mRNA level in SON OE/KD MEFs compared to control MEFs under vehicle (DMSO) condition. Four clusters of genes are shown. (**e**) GO analysis of genes in four clusters showing enriched KEGG pathways.

1783

1784

1785

1786

1787

1788

1789

1790

1791

1792

1793

1794

1795

Supplementary Figure 5. SON knockdown increases intron retention of proteostasis and mRNA metabolism genes. Heat map (**a**) and quantification (**b**) of intron retention events in MEFs with control or SON KD under basal (DMSO) and Tu conditions. Four clusters are shown. (**c**) The Integrative Genome Viewer representation of intron retention in selected genes. (**d**) GO analysis of genes in four clusters showing enriched KEGG pathways. Data: box and whiskers with minimum to maximum. Statistical tests used: unpaired one-tailed Student's t-test for b.

1802

1803

1804

1805

1806

1807

1808

1809

1810

1811

Supplementary Figure 6. SON overexpression decreases intron retention of protein processing and mRNA metabolism genes. Heat map (**a**) and quantification (**b**) of intron retention events in MEFs with control or SON overexpression under basal (DMSO) and Tu conditions. Three clusters are shown. (**c**) The Integrative Genome Viewer representation of intron retention in selected genes. (**d**) GO analysis of genes in three clusters showing enriched KEGG pathways. Data: box and whiskers with minimum to maximum. Statistical tests used: unpaired one-tailed Student's t-test for b.

1827

Supplementary Figure 7. SON reprograms opposing proteostasis and YAP1 transcriptional output under basal conditions. (**a**) Heat map of fold change of mature mRNA level in *Son* OE/KD MEFs compared to control MEFs under vehicle (DMSO) condition. All mature mRNAs in this heatmap are statistically differentially expressed (P<0.05) in both SON OE/KD conditions, compared to their respective controls. (**b**) GO analysis of these 481 and 501 genes. (**c**) Representative mature mRNA expression of proteostasis and YAP1 target genes (n=4 for Ctrl and *Son* siRNA and n=3 for Ctrl and *Son* sgRNA). Data: Mean ± S.E.M. Statistical tests used: unpaired one-tailed Student's t-test for c.

1836

1837

1838

Supplementary Figure 8. Nuclear speckle LLPS dictates opposing proteostasis and YAP1 signaling. (**a**) Two models explaining the relationship among nuclear speckles LLPS dynamics, proteostasis and YAP1 transcriptional output. Our results support model 1. (**b**) Heatmap demonstrates relative fold change of gene expression relative to DMSO control in IXA4, or Thap treated HEK293T cells. All genes induced or repressed by at least 1.41-fold with p value smaller than 0.05 in Thap condition (left), and representative YAP1-related genes (right). (**c, d**) GO analysis of all upregulated (**c**) or downregulated (**d**) genes in either IXA4 or Thap treatment by at least 1.41-fold with a p-value smaller than 0.05. (**e**) An expanded model of how the LLPS of nuclear speckles can dictate proteostasis and YAP1 transcriptional output. Please see the main text for details. (**f, g**) Diagram showing temporal changes of NS' LLPS (black), proteostasis (red) and YAP1 transcriptional output (blue) signal during ER stress (**f**).

1854

Supplementary Figure 9. HTS identifies compounds that alter nuclear speckle morphology and the UPR. (a) Compounds in the FDA-approved library ranked from lowest to highest on their ability to reduce NS sphericity. **(b)** Five drugs are shown to have a dose-dependent effect on increasing sphericity of NS (n=16). **(c)** Dose-dependent effects of drugs on decreasing NS sphericity (n=16). **(d, e)** GFP/cell **(d)** or cell number **(e)** measured in *Perk* promoter-driven dGFP reporter MEFs in the presence of Tu for four or eight hours after pre-treatment of different concentrations of drugs or DMSO for 24 hours (n=4). Data: Mean ± S.E.M.

1862

1863

1864

Supplementary Figure 10. PP is a bona fide nuclear speckle rejuvenator. (**a**) MEFs were treated with 3 µM NB, 1 µM PB, 3 µM PH, and 1 µM PP for 24 hours and RNA-seq was performed. PCA of global transcriptional response to drug treatments. (**b**) For each of the GSEA analysis, genes further activated by SON OE or further repressed by SON OE are compared to the transcriptome signatures of MEFs under different drug treatments. (**c, d**) Gene expression of select protein quality control (**c**) and YAP1 target genes (**d**) genes determined through mRNA-Seq (n=3). (**e**) LISA analysis listing log transformed p values for top predicted transcription regulators for genes upregulated (x-axis) and downregulated (y-axis) by PP. (**f**) Gene expression of select UPR genes determined through mRNA-Seq under different drugs treatment (n=3). (**g-i**) Western blot and quantification of UPR TFs (**g**), YAP1 nuclear and cytosol (**h**) and SON (**i**) level in response to 1 µM PP for 24 hours (n=3). Data: Mean ± S.E.M. Statistical tests used: unpaired one-tailed Student's t-test for g-i.

Supplementary Figure 11. Comparison of upregulated genes by PP and SON OE. (a) Scatter plot comparing the fold change of gene expression by PP (x-axis) and by Tu (y-axis) under SON OE or SON KD condition. Correlation coefficient and p value are shown for each plot. Chow test indicates statistically significant coefficients between the two linear regressions with p=0.00195. (**b, c**) Venn diagram showing (**b**) and GO analysis of (**c**) specific and commonly upregulated genes by PP and Tu in SON OE MEFs. (**d**) Venn diagram showing specific and common upregulated genes by PP and SON in MEFs. (**e**) GO analysis of common 101 genes. (**f**) GSEA analysis comparing genes upregulated by SON with those regulated by PP.

1888

Supplementary Figure 12. Comparison of downregulated genes by PP and SON OE. (**a-b**) Venn diagram showing (**a**) and GO analysis of (**b**) specific and common downregulated genes by PP and Tu in SON OE MEFs. (**c**) Venn diagram showing specific and common downregulated genes by PP and SON in MEFs. (**d**) GO analysis of 186 common genes. (**e**) GSEA analysis comparing genes downregulated by SON with those regulated by PP.

1894

Supplementary Figure 13. Comparison of intron retention (IR) events among PP, SON OE and SON KD. (**a**) Quantification of IR under DMSO and PP condition (n=3). (**b**) Heatmap showing RPKM normalized level of retained introns in DMSO and PP condition. (**c, d**) Venn diagram comparing genes with specific or common IRs between different conditions. (**e**) Genome browser view of selective genes with reduced IR by PP. (**f**) GO analysis of genes with increased or reduced IR by PP. Data: Mean ± S.E.M. Statistical tests used: unpaired one-tailed Student's t-test for a.

1901

Supplementary Figure 14. PP directly targets SON to modulate nuclear speckle LLPS dynamics. (**a**) Western blots of SON with siRNA-mediated knockdown and SON OE. (**b**) CETSA of ATF4 with 3µM PP. Both representative blot and quantification from independent replicates are shown (n=2 for DMSO and n=3 for PP). (**c**) Computational prediction of IDR in mouse SRSF2 (SC35) protein. (**d**) Diagram illustrating the constructs for droplet formation assay (**e**) Representative images of droplet formation assay with different concentrations of recombinant protein at 125mM NaCl. (**f**) Representative images of droplet formation assay with different salt concentrations with 20µM recombinant proteins. (**g, h**) Representative images of droplet formation assay with different recombinant proteins (**g**) and quantification (**h**) of area-normalized perimeter changes in the time span of 20 minutes with 50mM NaCl (n=3). (**i**) Alignment of protein sequences of SON orthologs in seven different species. SON IDR2 is located within the most conserved region (highlighted by light yellow). Data: Mean ± S.E.M. Statistical tests used: Two-way ANOVA for b and one-way ANOVA for h.

Supplementary Figure 15. PP modulates nuclear speckle LLPS dynamics in a cell-free system. (**a**) Diagram showing NE-supplemented SON IDR2 condensates are expected to compartmentalize splicing factors and exhibit less spherical morphology. (**b**) NE-supplemented SON IDR2 condensates are spun down and subject to mass spectrometry. Top 15 proteins mostly enriched in SON IDR2-compartmentzlied condensates with p value<0.05, and the status of whether these proteins have been identified in nuclear speckles in cells in two datasets [52, 53] (**c**) GO of top enriched biological pathways of proteins depleted or enriched in SON IDR2-compartmentzlied condensates. (**d, e**) Representative images of droplet formation assay with

1946 increasing concentration of SON IDR2 with or without NE supplementation (**e**) and quantification
1947 (**e**) of the number, sphericity and total areas of droplets (n=6). (**f**) Representative images and
1948 quantification of spatial distribution of mCherry::SON IDR2 and GFP::SRSF2. (**g**) Representative
1949 images and quantification of spatial distribution of mCherry::SON IDR2, GFP::SRSF2 and mouse
1950 genomic DNA. Data: Mean ± S.E.M. Statistical tests used: unpaired one-tailed Student's t-test for
1951 e.

1952

1953

1954

1955

1956

1957

1958

1959

1960

1961

1962

Supplementary Figure 16. PP alters the sensitivity of different condensates to 1,6 hexanediol. (**a**) 1,6 hexanediol sensitivity assay with representative images and quantification (n=20~95) of the ratio of cytosol over nuclear intensity of GFP::SRSF2 signal. (**b**) 1,6 hexanediol sensitivity assay with representative images and quantification of sphericity (n=24~50) of GFP::SRSF2 signal and Manders' coefficient (n=10~14) of signals of GFP and Hoechst. (**c**) 1,6 hexanediol sensitivity assay with representative images and quantification of ratio of cytosol to nuclear MED1 signal (n=19~29) and sphericity of MED1 signal (n=31~75). (**d**) 1,6 hexanediol sensitivity assay with representative images and quantification of ratio of cytosol to nuclear GW182 signal (n=16~54) and sphericity (n=19~97) of GW182 signal. Data: Mean ± S.E.M. Statistical tests used: unpaired one-tailed Student's t-test for all data.

1973

1974

1975

1976

1977

Supplementary Figure 17. Nanomolar concentration of PP promotes UPS and ALP without inducing cellular stress. (**a, b**) MEFs were treated with DMSO or 100nM PP for 24 hours. Representative western blot (**a**) and quantification (**b**) of different proteins (n=3). (**c**) MEFs were transiently transfected with scrambled or Son siRNA for 24 hours before treated with DMSO or 100nM PP for another 24 hours. 20S proteasome activity assay was then performed (n=10~13). (**d, e**) MEFs were treated with vehicle control or 1μM PP for ~22 hours and then co-treated with or without Baf A (100nM for 22 hours) (n=2~4). Representative western blot image (**d**) and quantification (**e**) of LC3II and LC3II/LC3I ratio. Data: Mean ± S.E.M. Statistical tests used: unpaired one-tailed Student's t-test for all data.

1987

Supplementary Figure 18. Micromolar PP promotes autophagy and UPS activity and represses translation. MEFs were treated with vehicle control or 1μM PP for ~24 hours (22 hours for **e** and **f**) and then co-treated with or without puromycin (10 μg/mL for 30 minutes), MG132 (10μM for 110 minutes) or Baf A (100nM for 22 hours) (n=3 for all samples). Western blot and quantification of puromycin-incorporated proteins (**a, b**), poly-ubiquitinated protein (**c, d**) and LC3II and LC3II/LC3I ratio (**e, f**). Data: Mean ± S.E.M. Statistical tests used: unpaired one-tailed Student's t-test for all data.

Supplementary Figure 19. Pyrvinium pamoate reduces pathological Tau and Rhodopsin level by boosting autophagy and UPS activity. (**a, b**) NIH3T3 RHO[P23H] cells were transfected with scrambled or *Son* siRNA for 24 hours before treated with DMSO or 0.1µM PP for another 24 hours. Western blot (**a**) and quantification (**b**) of RHO[P23H] level (n=4). (**c, d**) NIH3T3 RHO[P23H] cells

2000 were treated with 0.1µM PP and co-treated with or without BafA (100nM) for 24 hours. Western
2001 blot (**c**) and quantification (**d**) of RHO^{P23H} level (n=3). (**e, f**) NIH3T3 RHO^{P23H} cells were co-treated
2002 with or without MG132 (10µM for 120 minutes). Western blot (**e**) and quantification (**f**) of RHO^{P23H}
2003 level (n=3). (**g**) Representative images of primary mouse neurons treated with DMSO or 0.5 µM
2004 PP for 24 hours. (**h, i**) Representative images showing an increase of the number of mCherry
2005 positive puncta in primary neurons cultured in the presence of 0.1µM PP for 12 hours, with
2006 zoomed in images of regions marked with white rectangles (**h**). Quantification of the number of
2007 total vacuoles, autophagosome and autolysosomes (n=7~12) (**i**). (**j, k**) Tau P301S-expressing
2008 primary neurons were co-treated with vehicle or 0.1µM PP in the presence or absence of MG132
2009 (10µM) for 12 hours and western blot (**j**) and quantification (**k**) of different proteins (n=3~5). All
2010 data mean ± S.E.M. Statistical tests used: unpaired one-tailed Student's t-test for all data.

2011

2012

2013

2014

2015

2016

2017

2018

2019

2020

2021

2022

2023

2024

2025

Supplementary Figure 20. Pyrvinium pamoate reduces pathological Rhodopsin level in a manner that depends on reduced YAP1 activity. (**a, b**) NIH3T3 RHO^P23H cells were treated with 0.1µM PP for 24 hours and co-treated with or without XMU-MP-1 (1µM). Western blot (**a**) and quantification (**b**) of RHO^P23H level (n=3). (**c, d**) NIH3T3 RHO^P23H cells were treated with 0.1µM PP for 24 hours and co-treated with or without TRULI (1µM). Western blot (**c**) and quantification (**d**) of RHO^P23H level (n=3). (**e, f**) NIH3T3 RHO^P23H cells were transiently transfected with scrambled or Mst1/Mst2 siRNAs for 24 hours and then treated with DMSO or 0.1µM PP for another 24 hours. Western blot (**e**) and quantification (**f**) of MST1/2 and RHO^P23H level (n=3). (**g, h**) Tau P301S-expressing primary neurons were co-treated with vehicle or 0.1µM PP in the presence or absence of YAP1 activator TRULI (10µM) for 12 hours and western blot (**g**) and quantification (**h**) of different proteins (n=4). All data: mean ± S.E.M. Statistical tests used: unpaired one-tailed Student's t-test for all data.

2038

2039

2040

Genes upregulated by PP

Genes downregulated by PP

Supplementary Figure 21. Gene signatures in human AD subjects are opposite from those regulated by PP revealed by bulk RNA-Seq. Relative gene expressions in different brain regions of human AD subjects normalized to control subjects as reported in [87].

2059

Supplementary Figure 22. Gene signatures in late-stage human AD subjects with severe tauopathy are opposite from those regulated by PP. (**a**) Log $_2$ transformed values of fold change of gene expression of different individual or combined brain regions of AD versus control human subjects for top genes that were either upregulated or downregulated by PP (with a log$_2$ fold change > 1.5) in MEFs. (**b**) Phenotypic clustering of 48 individuals (columns) using seven clinicopathological variables as reported and adapted from [88]. (**c-f**) Log $_2$ transformed values of fold change of mean gene expression of different cell types of early or late AD versus no pathology human subjects for top genes that were either upregulated (**c**) or downregulated by PP in MEFs (**e**). Log 2 transformed values of fold change of mean gene expression of different cell types of late AD versus early AD human subjects for top genes that were either upregulated (**d**) or downregulated by PP in MEFs (**f**). Data: Mean ± S.E.M. Statistical tests used: unpaired one-tailed Student's t-test for a. One sample t-test (one-tailed). * p<0.05, ** p<0.01, *** p<0.001, **** p<0.0001 for c-f.

2073

2074

2075

2076

Supplementary Figure. 23. Nanomolar PP rejuvenates nuclear speckles and alleviates tau burden in human iPSC-neurons expressing mutant Tau in a SON-dependent manner without causing cellular stress. (**a-b**) WT and V337M Tau-expressing iPSC neurons were treated with 500 nM PP for 24 hours and LDH release assay were performed. LDH enzyme activity (**a**) and normalized cytotoxicity (**b**) were shown. (**c**) WT and V337M Tau-expressing iPSC neurons were treated with increasing concentration of PP for 12 hours and western blot of eIF2α and p-eIF2α were performed. The ratio of p-eIF2α to total eIF2α were calculated. (**d**) Wild-type and V337M Tau-expressing iPSC-neurons were infected with scrambled shRNA or SON shRNA-encoding lentivirus and treated with DMSO or PP (100nM) for 12 hours, and IF against nuclear speckle (Ab11826 against SRRM2), p-Tau (Ser422) and chromatin (DAPI) were performed.

2087

2088

2089

2090

2091

2092

2093

2094

2095

Supplemental Movie legends:

2097   **Movie S1.** Time lapse imaging of droplet formation with 20μm SON IDR1 in 125mM NaCl.

2098   **Movie S2.** Time lapse imaging of droplet formation with 20μm SON IDR2 in 125mM NaCl.

2099   **Movie S3.** Time lapse imaging of droplet formation with 20μm SRSF2 in 125mM NaCl.

2100   **Movie S4.** Time lapse imaging of droplet formation with 10μm SON IDR2 supplemented with
2101   0.6mg/ml GFP::SRSF2 MEF NE.

Supplemental Table legends:

2103   **Table S1.** FPKM normalization of RNA-seq of SON OE and KD cells in the absence or presence
2104   of Tunicamycin in MEFs.

2105   **Table S2.** TPM normalization of RNA-seq of different chemical treatments in MEFs.

2106

2107

2108

2109