Computational stabilization of a non-heme iron enzyme enables efficient

evolution of new function

Authors

Brianne R. King, Kiera H. Sumida, Jessica L. Caruso, David Baker, and Jesse G. Zalatan*

Corresponding Author

Jesse G. Zalatan − Department of Chemistry, University of Washington, Seattle, Washington

98195, United States; Email: zalatan@uw.edu

Authors

Brianne R. King − Department of Chemistry, University of Washington, Seattle, Washington

98195, United States

Kiera H. Sumida − Department of Chemistry and Institute for Protein Design, University of

Washington, Seattle, Washington 98195, United States

Jessica L. Caruso − Department of Chemistry, University of Washington, Seattle, Washington

98195, United States

David Baker − Institute for Protein Design, Department of Biochemistry, and Howard Hughes

Medical Institute, University of Washington, Seattle, Washington 98195, United States

Abstract

Directed evolution has emerged as a powerful tool for engineering new biocatalysts. However, introducing new catalytic residues can be destabilizing, and it is generally beneficial to start with a stable enzyme parent. Here we show that the deep learning-based tool ProteinMPNN can be used to redesign Fe(II)/αKG superfamily enzymes for greater stability, solubility, and expression while retaining both native activity and industrially-relevant non-native functions. For the Fe(II)/αKG enzyme tP4H, we performed site-saturation mutagenesis with both the wild-type and stabilized design variant and screened for activity increases in a non-native C-H hydroxylation reaction. We observed substantially larger increases in non-native activity for variants obtained from the stabilized scaffold compared to those from the wild-type enzyme. ProteinMPNN is user-friendly and widely-accessible, and straightforward structural criteria were sufficient to obtain stabilized, catalytically-functional variants of the Fe(II)/αKG enzymes tP4H and GriE. Our work suggests that stabilization by computational sequence redesign could be routinely implemented as a first step in directed evolution campaigns for novel biocatalysts.

Introduction

Directed evolution is a powerful method to generate enzymes for new chemical transformations.[1,2] However, catalytic functional groups often have destabilizing effects on protein structure, and altering active site groups for new reactions can lead to unstable, non-functional proteins.[3–11] Initiating a directed evolution campaign from a stabilized variant can be an effective way to overcome this problem.[12,13] Typically, a starting point for directed evolution is obtained by screening a library of candidate enzymes for a desired promiscuous activity. If a thermostable homolog with similar catalytic properties is identified, it can then be used as a starting point for evolution of the desired function.[14,15] Alternatively, there are a variety of strategies to produce stable variants using directed evolution,[16,17] ancestral reconstruction,[14] protein recombination,[18,19] or computational engineering.[20–22] These methods are often time- and resource-intensive, highlighting the need for simple and accessible alternatives.

An important class of enzymes where stabilization would be useful is the non-heme iron(II) α-ketoglutarate-dependent oxygenase (Fe(II)/αKG) superfamily. These enzymes have emerged as a rich source of potential new biocatalysts. Fe(II)/αKG enzymes can perform asymmetric C-H oxyfunctionalization reactions on small molecule substrates, which could allow expedient and sustainable diversification of simple building blocks to a range of complex polyfunctional compounds.[23–26] The advantages offered by this enzyme family include a high degree of chemical flexibility in the iron-containing active site due to multiple open coordination sites, the utilization of benign molecular oxygen as an oxidant, and use of the inexpensive and readily available co-factor αKG. However, Fe(II)/αKGs can be relatively unstable,[27,28] which may limit their practical applications in organic synthesis.

The recent use of an Fe(II)/αKG in an industrial-scale drug biosynthesis pathway highlights

both the potential advantages and drawbacks of this family for biocatalysis. An engineered

Fe(II)/αKG was used to catalyze an enantioselective C-H hydroxylation to produce a key

intermediate for the anti-cancer drug belzutifan.[28] The reaction could be performed on kilogram-

scale and bypassed five steps of the pre-existing chemical synthesis route. Notably, this effort

required an extensive, large-scale directed evolution campaign. Furthermore, early rounds of

screening yielded stabilizing mutations before significant improvements in turnover could be

obtained in later rounds. These findings highlight the potential utility of Fe(II)/αKG for practical,

industrial-scale green chemistry but also the importance of enzyme stability in the evolution of

new function.

Recent applications of deep learning to protein design have provided new and relatively

straightforward methods to stabilize protein scaffolds,[22,29,30] and there is broad interest in applying

these approaches to directed evolution.[31] Here we demonstrate that the deep learning-based tool

ProteinMPNN[29,30] enables more efficient optimization of a synthetically-relevant, non-native C-H

hydroxylation reaction in an Fe(II)/αKG family member. A critical step was restricting the

redesign from modifying active site and adjacent residues, which would otherwise be readily

mutated to stabilize the enzyme. With a stabilized starting point for site-saturation mutagenesis,

we observed substantially larger increases in non-native activity compared to the same mutations

in the wild-type parent enzyme. We suggest that this designed stabilization approach should be

routinely used in future directed evolution campaigns with the Fe(II)/αKG superfamily and will

likely be effective in a broad range of other enzyme families.

Results and Discussion

Fe(II)/αKGs with promiscuous activity for C-H hydroxylation of free carboxylate substrates

Within the Fe(II)/αKG enzyme superfamily, free amino acid hydroxylases are attractive

candidates for engineering new reactions.[23–26] Because Fe(II)/αKG amino acid hydroxylases

already have catalytic machinery to interact with amine and carboxylate functional groups in

amino acids, we hypothesized that they might have promiscuous activity for substrates containing

only an amine or only a carboxylate. These molecules are important feedstocks for early-stage

oxyfunctionalization reactions in multi-step syntheses. Selectivity for these reactions has been

historically difficult to achieve with traditional transition metal catalysis, and a biocatalytic process

could offer improved regio- and stereoselectivity.[32,33]

Figure 1. Initial whole-cell reaction screen data with a panel of Fe(II)/αKG amino acid hydroxylases and free
acid substrate analogues. Reactions were performed in whole cell from 50 mL expression cultures where
whole cell volume was 1/20th the expression volume. Reactions were carried out in MOPS (pH 7.0, 50 mM)
with 20 mM substrate, 60 mM αKG (as disodium salt), 1 mM ferrous ammonium sulfate, and 1 mM L-
ascorbic acid.

We initially screened a panel of 12 Fe(II)/αKG amino acid hydroxylases for the ability to

hydroxylate free carboxylates (Figure 1). We chose a set of candidate carboxylate substrates (**1-3**)

that are structurally analogous to the native amino acid substrates L-pipecolic acid, L-proline, and

L-leucine. We used whole-cell biocatalysis and liquid chromatography-mass spectrometry (LC-

MS) to detect products. We confirmed that the native amino acid reaction products are detectable

with all 12 members of the enzyme panel (Table S6). We then screened for promiscuous activity

with carboxylates, and observed that one enzyme, tP4H,[34] has detectable activity with substrate **1**

(Figure 1). The total turnover number (TTN) with this substrate was ~5 after 24 hr incubation with

10 μM enzyme, ~130-fold lower than the TTN for the corresponding native amino acid substrate.

The reaction of tP4H with substrate **1** gives the *trans* product with a *d.r.* of 4:1 (Figure S4). tP4H

produces exclusively *trans* product with its native substrate L-pipecolic acid,[34] suggesting that the

free carboxylate substrate **1** and the native substrate are positioned similarly in the enzyme active

site with respect to the iron center. To confirm that tP4H and not a contaminating enzyme was

responsible for the observed non-native activity, we mutated active site residues that are involved

in Fe(II), substrate, or αKG binding. Because tP4H does not have an experimental structure, we

identified these active site residues using an Alphafold2[35] model (Figure S9) and comparisons to

structures of the highly homologous Fe(II)/αKG enzyme GriE.[36,37] In all cases, active site

mutations produced activity decreases for the non-native substrate **1** (Figure 2). We also observed

increased product yield with increasing wild-type tP4H concentration (Figure 2). Together these

results confirm that tP4H is responsible for the non-native reaction.

Figure 2. Validation of tP4H activity with free acid **1**. (A) Reaction of tP4H with substrate **1** to form *trans*-4-hydroxycyclohexane carboxylic acid **4**. (B) Yield of **4** after reaction of **1** with tP4H variants, as well as negative control reaction with Fe(II) and bovine serum albumin (BSA). The Fe 1 mM control was run in the absence of added enzyme. Purified enzyme concentration was varied between 10-40 μM with 20 mM **1**, 40 mM αKG, 1 mM ferrous ammonium sulfate, and 1 mM L-ascorbic acid in MES buffer (50 mM, pH 6.8). Reactions were carried out for 24 hours at 25 °C and quantified with analytical LC-MS. (C) Structural model of the tP4H showing key active site residues. Fe(II), αKG, and L-Pipecolic acid were modeled in Chimera.[38]

Stabilization of tP4H with ProteinMPNN

To improve tP4H activity towards substrate **1**, we began a directed evolution campaign but quickly encountered limitations due to poor enzyme stability. First, we found that tP4H variants were difficult to express and purify due to enzyme insolubility. Additionally, we found that the parent wild-type tP4H enzyme loses activity with time (Figure 3). These observations are consistent with prior reports on tP4H behavior.[34] It is possible for enzyme stability to improve during directed evolution, whether by selecting more stable variants during each round or by chance. For example, the evolved Fe(II)/αKG PsEFE had slight stability improvements compared to wild-type, despite the researchers not selecting for improved stability.[39] In another case, the Fe(II)/αKG UbP4H was successfully screened for improved stability after initial screening rounds destabilized the enzyme.[27] However, the benefits of starting with a highly stable enzyme for directed evolution are well-established.[12–14] A stabilized protein scaffold could potentially increase

the population of active, properly-folded protein or provide access to other mutants that otherwise

do not fold.

We used the deep learning-based tool ProteinMPNN[29,30] to generate stabilized variants of

tP4H through sequence design. We first used ProteinMPNN to redesign the entire tP4H sequence.

Unsurprisingly, we found that the predicted sequences eliminated key active site residues, which

is likely to disrupt enzymatic activity (Figure S10). This behavior is consistent with the well-

established propensity for catalytic active site residues to be destabilizing.[3–10] To preserve catalytic

function, we fixed the active site residues in all subsequent design efforts. We defined the active

site as any residues that contact the amino acid substrate, Fe(II), or the αKG cofactor, based on our

Alphafold2 model (Figure S9) and comparisons to the structure of the closely-related enzyme GriE

bound to L-leucine.[37] Because other residues throughout the protein could also be important, we

also tested four additional strategies using either sequence conservation or distance metrics. To

identify important conserved residues, we constructed a multiple sequence alignment (MSA) and

selected tP4H residues conserved in at least 35%, 70%, or 95% of sequences (Methods and Table

S4). Alternatively, we fixed any residues with side chains within a 10 Å sphere from the substrate

binding pocket. Using these five starting points (fix active site only, active site + 35%/70%/95%

conservation, active site + 10 Å sphere), we generated 48 ProteinMPNN sequences per method

and selected 4 each (20 total) for activity screens. Selection was based on calculated top-ranked

Cα-RMSD values matched to the input tP4H structure. We obtained only one variant that had any

detectable activity, with catalytic efficiency ~35-fold lower than wild-type tP4H (Figure S11). This

variant was designed from the sequence where >35% conserved residues are fixed, which

constrains more residues than the >70% or >95% cutoffs. This result suggests that even weakly

conserved residues may need to be fixed to maintain activity. Further analysis of the variant with

detectable activity revealed a ~3.5-fold increase in the K_M for the αKG cofactor (Figure S11). We

identified two residues in proximity to αKG, L228 and V230, that were mutated in the redesigned

sequence. These sequence changes may have contributed to improved stability at the expense of

cofactor binding and positioning, leading to the decrease in activity. Notably, L228 and V230 were

fixed in the designs generated from fixed active site + 10 Å sphere, but none of these designs had

detectable activity. Taken together, these findings suggest that additional criteria will be needed to

identify critical functional residues that should be fixed prior to sequence redesign.

To generate stabilized variants that maintain catalytic activity, we performed another set

of ProteinMPNN sequence redesigns with three new strategies to fix important residues. In each

case, we fixed the active site as defined above plus residues L228 and V230. For the first approach,

we fixed all residues at tP4H positions conserved in 35% of the MSA. We chose this cutoff because

it was the only one from our initial set that produced a stabilized variant with any detectable

activity, and we expected that fixing L228 or V230 could further improve these designs. For the

second and third approaches, we identified highly conserved positions regardless of whether the

wild-type tP4H residue is the most highly conserved amino acid. These strategies were based on

previous work suggesting that more stringent constraints are necessary to maintain activity in

ProteinMPNN redesign.[30] Every tP4H amino acid position was ranked based on the %

conservation of the most frequent amino acid present in the MSA, and the top 50% or 70% were

fixed as the wild-type tP4H residue. Together with fixed active site residues, these criteria resulted

in 148/272 (54%) or 198/272 (73%) fixed residues across the entire 272 amino acid protein. Using

these three strategies, we selected 32 designs each of 48 generated for a total of 96 sequences

(Supplementary spreadsheet – ProteinMPNN sequences_metrics). Of these designs, 69 expressed

detectable quantities of protein by SDS-PAGE and 11 had detectable activity above background

for the native substrate. For the active enzymes we proceeded to measure thermostability and

kinetic parameters for the native L-pipecolic acid substrate and the promiscuous carboxylate

substrate **1**. The variant with the highest k_{cat} for L-pipecolic acid was R2_11 (Table S7), with a k_{cat}

of 0.16 s^{-1} compared to 0.17 s^{-1} for wild-type. R2_11 was designed from the method where the top

70% ranked conserved residues were fixed, and had 44 designed mutations compared to the wild-

type sequence (Figure 3). R2_11 has modestly slower (~3-fold) non-native carboxylate

hydroxylase activity compared to wild-type tP4H and exhibits a 20 °C increase in thermostability

by temperature-dependent circular dichroism (CD) spectroscopy (Figure 3). When activity is

measured as a function of time, R2_11 maintains activity over a timescale of days, which is a

substantial improvement compared to wild-type tP4H (Figure 3).

Figure 3. Stability and activity of wild-type tP4H and ProteinMPNN design R2_11. (A) tP4H structure
(Alphafold2 model, Figure S9) color-coded to show sites fixed in the design process (blue, Supplementary

spreadsheet – ProteinMPNN sequences_metrics) and sites mutated in the ProteinMPNN R2_11 redesign
(orange-red). Sites colored black were neither fixed nor redesigned in the R2_11 variant. Side chains for
first shell active site residues (Table S4) are shown in blue. (B) Temperature-dependent CD spectroscopy
of wild-type tP4H and R2_11. (C) Activity-stability analysis of wild-type tP4H. (D) Activity-stability analysis
of R2_11. Relative activity was determined using PBP assay described in Supplementary Information.
202
203
Stabilization of GriE with ProteinMPNN

After successfully identifying sequence constraints for ProteinMPNN-mediated

stabilization of tP4H while maintaining catalytic function, we evaluated whether the same

approach would be effective with a second Fe(II)/αKG enzyme, GriE. This enzyme could benefit

from stabilization because, although it expresses well and is soluble, it loses activity at room

temperature over 24 hours (Figure S12). As with tP4H, we fixed the top 70% ranked conserved

residues along with catalytic residues identified from the GriE crystal structure (Supplementary

spreadsheet – ProteinMPNN sequences_metrics).[37] We generated 32 redesigned sequences and

found that 29 were expressed as soluble enzyme, and 27 showed activity with the GriE native

substrate L-leucine. The top design based on stability and kinetic parameters, GM_A9, showed a

catalytic efficiency (k_{cat}/K_M) 4-fold lower than wild-type GriE (Figure S12 & S13). One design,

GM_A11, had a 2-fold faster initial rate with L-leucine compared to GM_A9 but this design was

unstable by temperature-dependent CD and thus was not chosen for further analysis. For GM_A9,

the decrease in catalytic efficiency arises from a lower k_{cat} of 0.38 s^{-1} compared to our measured

k_{cat} of 2.1 s^{-1} for wild-type GriE. A decrease in catalytic efficiency is not surprising given that

increased stability could reduce conformational flexibility and negatively impact catalytic

function.[4,6,7]

We next screened the stabilized GriE redesign GM_A9 for substrate promiscuity.

Previously, wild-type GriE has been shown to accept substrates with increased substrate chain

lengths but has weaker activity towards substrates with substitution at C3.[40] We chose two

previously identified non-native substrates to test: L-norleucine and L-allo-isoleucine. L-

norleucine was chosen as a representative substrate with increased chain length compared to L-

leucine. L-allo-isoleucine was chosen because it has a methyl group substitution at C3. We

observed detectable activity with L-norleucine but not for L-allo-isoleucine (Table S8). Similar to

wild-type GriE, the GM_A9 variant maintained a preference for the extended chain L-norleucine

substrate over the C3-substituted L-allo-isoleucine substrate. The GM-A9 reactions with L-leucine

and L-norleucine were 11- and 4-fold slower than wild-type GriE reactions, respectively (Table

S8). These results suggest that our ProteinMPNN protocol can be readily applied to other

Fe(II)/αKG enzymes to stabilize proteins while maintaining synthetically-relevant catalytic

function that can be a foothold for further optimization by directed evolution.

Directed evolution of wild-type tP4H for carboxylate C-H hydroxylation activity

We next sought to improve the non-native carboxylate hydroxylase activity through

directed evolution. We prioritized tP4H because carboxylate hydroxylase activity was detectable

in both the wild-type and ProteinMPNN-stabilized variant, which allows for direct comparisons.

We conducted three rounds of directed evolution for both enzymes by varying first- and second-

shell substrate binding residues identified in the active site from our Alphafold2 structural model.

We defined the first shell as any residues that contact the amino acid substrate, based on

comparisons to the structure of ligand-bound GriE.[37] We defined the second shell as any residues

that make contacts with first shell residues. We used the 22c-trick method for single site-saturation

mutagenesis at each target position.[41]

We first performed directed evolution with wild-type tP4H. For the first screening round,

we chose three tP4H active site residues based on their potential role in substrate specificity: H58,

F114, and L174. Based on our tP4H structural model (Figure S9B), H58 likely contacts the amine

of native amino acid substrates and is presumably not needed or detrimental for carboxylate

substrates that lack an amine. F114 likely provides a substrate hydrophobic contact, and L174 is

part of a loop that could affect substrate binding. We screened whole cell biocatalysis reactions in

96 well plates for improved TTN and 80% *trans* selectivity in reactions with substrate **1**. Based on

production of hydroxylated product **4**, the top 5% of mutants were chosen for validation with

purified enzymes. We obtained several variants with modest activity improvements, and the best

performer was mutant H58L with a TTN of 7 (Figure 4A). In a second round starting from H58L,

we rescreened mutants at F114 and L174 and screened an additional 14 first and second shell

residues (Table S1). We identified the improved variant H58L/W170Q with a TTN of 15. In a

third round starting from H58L/W170Q, we screened 9 residues that showed activity increases in

previous rounds and identified the improved variant H58L/W170Q/E118K with a TTN of 31

(Figure 4A & Table S1). Overall, after three rounds of directed evolution for improved carboxylate

hydroxylase activity with wild-type tP4H we obtained a 6-fold improvement in TTN and

maintained >80% selectivity for the *trans* reaction product (Figure S4).

262
263
Figure 4. (A) Reaction scheme for C-H hydroxylation of substrate **1** with tP4H, R2_11, and associated
variants. (B) Directed evolution of wild-type tP4H. (C) Directed evolution of stabilized variant R2_11.
Reactions were carried out for 24 hr at 25 °C using purified enzyme (10-20 μM) in MES buffer (50 mM, pH
6.8), with 20 mM cyclohexane carboxylic acid **1**, 40 mM αKG, 1 mM ferrous ammonium sulfate, and 1 mM

ascorbic acid. Concentration of **4** in quenched reaction samples was quantified by analytical LC-MS
analysis.
270
271 *Directed evolution of ProteinMPNN redesign R2_11 for carboxylate C-H hydroxylation activity.*
272
To optimize carboxylate hydroxylase activity in ProteinMPNN-stabilized tP4H, we

conducted directed evolution using a similar strategy to our approach with wild-type tP4H, with

minor modifications. In the first round of site saturation mutagenesis, we started with a larger pool

of 19 first- and second-shell residues including the three sites from prior round one (H58, F114,

and L174) and the 16 additional sites from prior round 2 (Table S1). As before, we screened whole

cell biocatalysis reactions for improved TTN and >80% *trans* selectivity with carboxylic acid

substrate **1**. The top hit was H58F, which is the same position but a different mutant than the

previous round one winner, H58L. R2_11_H58F displayed a 27-fold increase in TTN relative to

the parent R2_11 (Figure 4B). This effect is substantially bigger than the <2-fold improvement

obtained with H58L relative to wild-type tP4H. Notably, the 27-fold increase in a single round

from stabilized R2_11 was already larger than the total 6-fold improvement from three rounds of

directed evolution from wild-type tP4H. Given the strong performance of the H58F mutant, we

also evaluated its effect in the wild-type tP4H background and observed a small, <2-fold increase

in TTN, similar to the effect of H58L on wild-type tP4H (Figure S14). Thus, the strong, 27-fold

improvement with the H58F mutant depends on the context of the stabilized R2_11 backbone.

In a second round of screening from R2_11_H58F, we selected the 18 residues that were

screened in prior round two from wild-type tP4H (Table S1). We retained this large pool of

residues to ensure a direct comparison to the tP4H directed evolution workflow. This round

identified improved variant L174G. R2_11_H58F/L174G has a TTN of 72. This TTN is a 1.6-fold

improvement from parent R2_11_H58F and outstrips any variant obtained from the wild-type

tP4H backbone (Figure 4B). In a third round of screening from R2_11_H58F/L174G, we selected

9 residues that were screened in prior round 3 from wild-type tP4H (Table S1). This round

identified the improved variant V57H with a TTN of 138, a 1.7-fold improvement from the

previous round.

Overall, the ProteinMPNN-stabilized tP4H directed evolution campaign produced an 80-

fold improvement in TTN from the base R2_11 redesign, compared to a modest 6-fold

improvement in the wild-type tP4H evolutionary trajectory. Although the R2_11 parent starts ~3-

fold slower than wild-type tP4H, the much larger improvement over three rounds of directed

evolution produced an R2_11 triple mutant with a 4.5-fold higher TTN than the triple mutant

obtained from wild-type tP4H (Figure 4).

In addition to a more efficient directed evolution trajectory, the R2_11 triple mutant

maintains high stability relative to the wild-type tP4H and associated variants from directed

evolution (Figure 5A, Figure S15). Higher stability allows reactions to be run more efficiently,

both at higher temperatures and for less time. For example, after 6 hours at 35 °C, the R2_11 triple

mutant reaches a mean TTN of 142 for the non-native reaction with carboxylate **1** to form product

4 with 4:1 selectivity for the *trans* reaction product (Figure 5B). The TTN after 6 hours at 35 °C

is comparable to the TTN after 24 hours at 25 °C. In contrast, the tP4H triple mutant shows a slight

decrease in TTN at 35 °C, likely due to enzyme instability at higher temperatures (Figure 5B). The

stability profile of the R2_11 triple mutant suggests that this enzyme will be more robust towards

further engineering compared to the tP4H triple mutant. Future engineering efforts with the R2_11

mutant could include improvements to key reaction metrics like turnover, selectivity, and

increased substrate scope.

Figure 5 A) Temperature dependent CD of the R2_11 and tP4H triple mutants. B) TTN for formation of **4**
(*d.r.* 4:1, Figure S4) with the R2_11 and tP4H triple mutants at two different temperatures. Reactions were
carried out for 6 hrs. at 25 °C and 35 °C using purified enzyme (15 µM) in MES buffer (50 mM, pH 6.8),
with 20 mM cyclohexane carboxylic acid **1**, 40 mM αKG, 1 mM ferrous ammonium sulfate, and 1 mM
ascorbic acid.

Conclusions

Directed evolution is a powerful tool to engineer enzymes for new-to-nature reactions.

However, many enzyme starting points for evolution may lack the stability required to reach user-

defined optimum fitness after multiple rounds of mutagenesis. Here we show that the deep

learning-based tool ProteinMPNN can be used to stabilize the Fe(II)/αKG enzyme superfamily

members tP4H and GriE with straightforward sequence constraints to maintain catalytic activity.

Consistent with previous results using ProteinMPNN,[30] the top tP4H design was identified by

using the most conservative of our chosen methods for fixing residues during sequence redesign.

Applying the same method to the related enzyme GriE readily produced stabilized variants with

catalytic activity.

Wild-type and redesigned tP4H both exhibit novel reactivity towards remote C-H

hydroxylation of a free carboxylic acid substrate. We directly compared evolutionary trajectories

of wild-type tP4H with the stabilized variant R2_11 and demonstrated superior performance of the

stabilized redesign variant. Future work will determine if this design method is generalizable to

optimize directed evolution for other enzymes and enzyme families. User-friendly deep learning-

based tools like ProteinMPNN and MutCompute are rapidly emerging,[22,29,30] and our work

suggests that these tools should be routinely incorporated into enzyme engineering workflows to

efficiently optimize catalytic fitness for new biocatalysts.[31]

Supplementary Information

Materials, experimental and analytical methods, compound characterization data (Figure S4-S7),

and enzyme characterization data (Figure S3, Figure S8, Figure S11-S15 and Tables S6-S8) (PDF).

Accession numbers for all Fe(II)/αKG enzymes used in this work, full nucleotide and amino acid

sequences for all reported wild-type and enzyme variants, oligonucleotide sequences for cloning

and for all ProteinMPNN designs, ProteinMPNN design screen criteria results (XLSX).

Acknowledgements

We thank Dr. Wolfgang Hüttel at the University of Freiburg and Hans Renata at Rice University

for the donation of the wild-type tP4H and GriE expression vectors, respectively, and for their

advice on tP4H and GriE reactions in various formats. We also thank Jonathan Zhang and Susanna

Vazquez Torres for their early contributions to Fe(II)/αKG reaction screening, and Dr. Martin

Sadilek in University of Washington Mass Spectrometry Facility for his continued support and

helpful advice in analytical method development. This work was supported by U.S. National

Institutes of Health grants T32 GM008268 (B.R.K., J.L.C.) and R35 GM124773 (J.G.Z.), and by

the Open Philanthropy Project Improving Protein Design Fund (K.H.S., D.B.).

Competing Interests

The authors declare that they have no competing interests.

References

(1) Renata, H.; Wang, Z. J.; Arnold, F. H. Expanding the Enzyme Universe: Accessing Non-Natural Reactions by Mechanism-Guided Directed Evolution. *Angew. Chem. Int. Ed.* **2015**, *54* (11), 3351–3367. https://doi.org/10.1002/anie.201409470.

(2) Zeymer, C.; Hilvert, D. Directed Evolution of Protein Catalysts. *Annu. Rev. Biochem.* **2018**, *87* (1), 1–27. https://doi.org/10.1146/annurev-biochem-062917-012034.

(3) Meiering, E. M.; Serrano, L.; Fersht, A. R. Effect of Active Site Residues in Barnase on Activity and Stability. *J. Mol. Biol.* **1992**, *225* (3), 585–589. https://doi.org/10.1016/0022-2836(92)90387-y.

(4) Shoichet, B. K.; Baase, W. A.; Kuroki, R.; Matthews, B. W. A Relationship between Protein Stability and Protein Function. *Proc. Natl. Acad. Sci.* **1995**, *92* (2), 452–456. https://doi.org/10.1073/pnas.92.2.452.

(5) Giver, L.; Gershenson, A.; Freskgard, P.-O.; Arnold, F. H. Directed Evolution of a Thermostable Esterase. *Proc. Natl. Acad. Sci.* **1998**, *95* (22), 12809–12813. https://doi.org/10.1073/pnas.95.22.12809.

(6) Beadle, B. M.; Shoichet, B. K. Structural Bases of Stability–Function Tradeoffs in Enzymes. *J. Mol. Biol.* **2002**, *321* (2), 285–296. https://doi.org/10.1016/s0022-2836(02)00599-5.

(7) Nagatani, R. A.; Gonzalez, A.; Shoichet, B. K.; Brinen, L. S.; Babbitt, P. C. Stability for Function Trade-Offs in the Enolase Superfamily "Catalytic Module". *Biochemistry* **2007**, *46* (23), 6688–6695. https://doi.org/10.1021/bi700507d.

(8) Tokuriki, N.; Stricher, F.; Serrano, L.; Tawfik, D. S. How Protein Stability and New Functions Trade Off. *PLoS Comput. Biol.* **2008**, *4* (2), e1000002. https://doi.org/10.1371/journal.pcbi.1000002.

(9) Tokuriki, N.; Jackson, C. J.; Afriat-Jurnou, L.; Wyganowski, K. T.; Tang, R.; Tawfik, D. S. Diminishing Returns and Tradeoffs Constrain the Laboratory Optimization of an Enzyme. *Nat. Commun.* **2012**, *3* (1), 1257. https://doi.org/10.1038/ncomms2246.

(10) Goldsmith, M.; Tawfik, D. S. Enzyme Engineering: Reaching the Maximal Catalytic Efficiency Peak. *Curr. Opin. Struct. Biol.* **2017**, *47*, 140–150. https://doi.org/10.1016/j.sbi.2017.09.002.

(11) Tsuboyama, K.; Dauparas, J.; Chen, J.; Laine, E.; Behbahani, Y. M.; Weinstein, J. J.; Mangan, N. M.; Ovchinnikov, S.; Rocklin, G. J. Mega-Scale Experimental Analysis of Protein Folding Stability in Biology and Design. *Nature* **2023**, *620* (7973), 434–444. https://doi.org/10.1038/s41586-023-06328-6.

(12) Bloom, J. D.; Labthavikul, S. T.; Otey, C. R.; Arnold, F. H. Protein Stability Promotes
Evolvability. *Proc. Natl. Acad. Sci.* **2006**, *103* (15), 5869–5874.
https://doi.org/10.1073/pnas.0510098103.

(13) Gumulya, Y.; Baek, J.-M.; Wun, S.-J.; Thomson, R. E. S.; Harris, K. L.; Hunter, D. J. B.;
Behrendorff, J. B. Y. H.; Kulig, J.; Zheng, S.; Wu, X.; Wu, B.; Stok, J. E.; Voss, J. J. D.; Schenk,
G.; Jurva, U.; Andersson, S.; Isin, E. M.; Bodén, M.; Guddat, L.; Gillam, E. M. J. Engineering
Highly Functional Thermostable Proteins Using Ancestral Sequence Reconstruction. *Nat. Catal.*
2018, *1* (11), 878–888. https://doi.org/10.1038/s41929-018-0159-5.

(14) Trudeau, D. L.; Tawfik, D. S. Protein Engineers Turned Evolutionists—the Quest for the
Optimal Starting Point. *Curr. Opin. Biotechnol.* **2019**, *60*, 46–52.
https://doi.org/10.1016/j.copbio.2018.12.002.

(15) Besenmatter, W.; Kast, P.; Hilvert, D. Relative Tolerance of Mesostable and Thermostable
Protein Homologs to Extensive Mutation. *Proteins: Struct., Funct., Bioinform.* **2007**, *66* (2),
500–506. https://doi.org/10.1002/prot.21227.

(16) Socha, R. D.; Tokuriki, N. Modulating Protein Stability – Directed Evolution Strategies for
Improved Protein Function. *FEBS J.* **2013**, *280* (22), 5582–5595.
https://doi.org/10.1111/febs.12354.

(17) Stimple, S. D.; Smith, M. D.; Tessier, P. M. Directed Evolution Methods for Overcoming
Trade-offs between Protein Activity and Stability. *AIChE J.* **2020**, *66* (3).
https://doi.org/10.1002/aic.16814.

(18) Li, Y.; Drummond, D. A.; Sawayama, A. M.; Snow, C. D.; Bloom, J. D.; Arnold, F. H. A
Diverse Family of Thermostable Cytochrome P450s Created by Recombination of Stabilizing
Fragments. *Nat. Biotechnol.* **2007**, *25* (9), 1051–1056. https://doi.org/10.1038/nbt1333.

(19) Heinzelman, P.; Snow, C. D.; Wu, I.; Nguyen, C.; Villalobos, A.; Govindarajan, S.;
Minshull, J.; Arnold, F. H. A Family of Thermostable Fungal Cellulases Created by Structure-
Guided Recombination. *Proc. Natl. Acad. Sci.* **2009**, *106* (14), 5610–5615.
https://doi.org/10.1073/pnas.0901417106.

(20) Goldenzweig, A.; Goldsmith, M.; Hill, S. E.; Gertman, O.; Laurino, P.; Ashani, Y.; Dym,
O.; Unger, T.; Albeck, S.; Prilusky, J.; Lieberman, R. L.; Aharoni, A.; Silman, I.; Sussman, J. L.;
Tawfik, D. S.; Fleishman, S. J. Automated Structure- and Sequence-Based Design of Proteins for
High Bacterial Expression and Stability. *Mol. Cell* **2016**, *63* (2), 337–346.
https://doi.org/10.1016/j.molcel.2016.06.012.

(21) Musil, M.; Konegger, H.; Hon, J.; Bednar, D.; Damborsky, J. Computational Design of
Stable and Soluble Biocatalysts. *ACS Catal.* **2019**, *9* (2), 1033–1054.
https://doi.org/10.1021/acscatal.8b03613.

(22) Lu, H.; Diaz, D. J.; Czarnecki, N. J.; Zhu, C.; Kim, W.; Shroff, R.; Acosta, D. J.; Alexander,
B. R.; Cole, H. O.; Zhang, Y.; Lynd, N. A.; Ellington, A. D.; Alper, H. S. Machine Learning-
Aided Engineering of Hydrolases for PET Depolymerization. *Nature* **2021**, *604* (7907), 662–
667. https://doi.org/10.1038/s41586-022-04599-z.

(23) Islam, M. S.; Leissing, T. M.; Chowdhury, R.; Hopkinson, R. J.; Schofield, C. J. 2-
Oxoglutarate-Dependent Oxygenases. *Annu. Rev. Biochem.* **2018**, *87*, 585–620.
https://doi.org/10.1146/annurev-biochem-061516-044724.

(24) Herr, C. Q.; Hausinger, R. P. Amazing Diversity in Biochemical Roles of Fe(II)/2-
Oxoglutarate Oxygenases. *Trends Biochem Sci* **2018**, *43* (7), 517–532.
https://doi.org/10.1016/j.tibs.2018.04.002.

(25) Papadopoulou, A.; Meyer, F.; Buller, R. M. Engineering Fe(II)/α-Ketoglutarate-Dependent
Halogenases and Desaturases. *Biochemistry* **2022**. https://doi.org/10.1021/acs.biochem.2c00115.

(26) Zwick, C. R.; Renata, H. Harnessing the Biocatalytic Potential of Iron- and α-Ketoglutarate-
Dependent Dioxygenases in Natural Product Total Synthesis. *Nat. Prod. Rep.* **2020**, *37* (8),
1065–1079. https://doi.org/10.1039/c9np00075e.

(27) Liu, C.; Zhao, J.; Liu, J.; Guo, X.; Rao, D.; Liu, H.; Zheng, P.; Sun, J.; Ma, Y.
Simultaneously Improving the Activity and Thermostability of a New Proline 4-Hydroxylase by
Loop Grafting and Site-Directed Mutagenesis. *Appl. Microbiol. Biotechnol.* **2019**, *103* (1), 265–
277. https://doi.org/10.1007/s00253-018-9410-x.

(28) Cheung-Lee, W. L.; Kolev, J. N.; McIntosh, J. A.; Gil, A. A.; Pan, W.; Xiao, L.; Velásquez,
J. E.; Gangam, R.; Winston, M. S.; Li, S.; Abe, K.; Alwedi, E.; Dance, Z. E. X.; Fan, H.; Hiraga,
K.; Kim, J.; Kosjek, B.; Le, D. N.; Marzijarani, N. S.; Mattern, K.; McMullen, J. P.; Narsimhan,
K.; Vikram, A.; Wang, W.; Yan, J.; Yang, R.; Zhang, V.; Zhong, W.; DiRocco, D. A.; Morris,
W. J.; Murphy, G. S.; Maloney, K. M. Engineering Hydroxylase Activity, Selectivity, and
Stability for a Scalable Concise Synthesis of a Key Intermediate to Belzutifan. *Angew. Chem.*
Int. Ed. **2024**, *63* (13), e202316133. https://doi.org/10.1002/anie.202316133.

(29) Dauparas, J.; Anishchenko, I.; Bennett, N.; Bai, H.; Ragotte, R. J.; Milles, L. F.; Wicky, B.
I. M.; Courbet, A.; Haas, R. J. de; Bethel, N.; Leung, P. J. Y.; Huddy, T. F.; Pellock, S.; Tischer,
D.; Chan, F.; Koepnick, B.; Nguyen, H.; Kang, A.; Sankaran, B.; Bera, A. K.; King, N. P.;
Baker, D. Robust Deep Learning–Based Protein Sequence Design Using ProteinMPNN. *Science*
2022, *378* (6615), 49–56. https://doi.org/10.1126/science.add2187.

(30) Sumida, K. H.; Núñez-Franco, R.; Kalvet, I.; Pellock, S. J.; Wicky, B. I. M.; Milles, L. F.;
Dauparas, J.; Wang, J.; Kipnis, Y.; Jameson, N.; Kang, A.; Cruz, J. D. L.; Sankaran, B.; Bera, A.
K.; Jiménez-Osés, G.; Baker, D. Improving Protein Expression, Stability, and Function with
ProteinMPNN. *J. Am. Chem. Soc.* **2024**. https://doi.org/10.1021/jacs.3c10941.

(31) Yang, J.; Li, F.-Z.; Arnold, F. H. Opportunities and Challenges for Machine Learning-
Assisted Enzyme Engineering. *ACS Cent. Sci.* **2024**, *10* (2), 226–241.
https://doi.org/10.1021/acscentsci.3c01275.

(32) Roduner, E.; Kaim, W.; Sarkar, B.; Urlacher, V. B.; Pleiss, J.; Gläser, R.; Einicke, W.;
Sprenger, G. A.; Beifuß, U.; Klemm, E.; Liebner, C.; Hieronymus, H.; Hsu, S.; Plietker, B.;
Laschat, S. Selective Catalytic Oxidation of C-H Bonds with Molecular Oxygen. *ChemCatChem*
2013, *5* (1), 82–112. https://doi.org/10.1002/cctc.201200266.

(33) He, C.; Whitehurst, W. G.; Gaunt, M. J. Palladium-Catalyzed C(Sp3)–H Bond
Functionalization of Aliphatic Amines. *Chem* **2019**, *5* (5), 1031–1058.
https://doi.org/10.1016/j.chempr.2018.12.017.

(34) Klein, C.; Hüttel, W. A Simple Procedure for Selective Hydroxylation of L-Proline and L-
Pipecolic Acid with Recombinantly Expressed Proline Hydroxylases. *Adv. Synth. Catal.* **2011**,
353 (8), 1375–1383. https://doi.org/10.1002/adsc.201000863.

(35) Mirdita, M.; Schütze, K.; Moriwaki, Y.; Heo, L.; Ovchinnikov, S.; Steinegger, M.
ColabFold: Making Protein Folding Accessible to All. *Nat. Methods* **2022**, *19* (6), 679–682.
https://doi.org/10.1038/s41592-022-01488-1.

(36) Hu, X.; Huang, X.; Liu, J.; Zheng, P.; Gong, W.; Yang, L. Structures of L-Proline Trans-
Hydroxylase Reveal the Catalytic Specificity and Provide Deeper Insight into AKG-Dependent
Hydroxylation. *Acta Crystallogr. Sect. D* **2023**, *79* (Pt 4), 318–325.
https://doi.org/10.1107/s2059798323001936.

(37) Lukat, P.; Katsuyama, Y.; Wenzel, S.; Binz, T.; König, C.; Blankenfeldt, W.; Brönstrup, M.;
Müller, R. Biosynthesis of Methyl-Proline Containing Griselimycins, Natural Products with
Anti-Tuberculosis Activity. *Chem Sci* **2017**, *8* (11), 7521–7527.
https://doi.org/10.1039/c7sc02622f.

(38) Pettersen, E. F.; Goddard, T. D.; Huang, C. C.; Couch, G. S.; Greenblatt, D. M.; Meng, E.
C.; Ferrin, T. E. UCSF Chimera—A Visualization System for Exploratory Research and
Analysis. *J. Comput. Chem.* **2004**, *25* (13), 1605–1612. https://doi.org/10.1002/jcc.20084.

(39) Goldberg, N. W.; Knight, A. M.; Zhang, R. K.; Arnold, F. H. Nitrene Transfer Catalyzed by
a Non-Heme Iron Enzyme and Enhanced by Non-Native Small-Molecule Ligands. *J. Am. Chem.*
Soc. **2019**, *141* (50), 19585–19588. https://doi.org/10.1021/jacs.9b11608.

(40) Zwick, C. R.; Renata, H. Remote C–H Hydroxylation by an α-Ketoglutarate-Dependent
Dioxygenase Enables Efficient Chemoenzymatic Synthesis of Manzacidin C and Proline
Analogs. *J. Am. Chem. Soc.* **2018**, *140* (3), 1165–1169. https://doi.org/10.1021/jacs.7b12918.

(41) Kille, S.; Acevedo-Rocha, C. G.; Parra, L. P.; Zhang, Z.-G.; Opperman, D. J.; Reetz, M. T.;
Acevedo, J. P. Reducing Codon Redundancy and Screening Effort of Combinatorial Protein

Libraries Created by Saturation Mutagenesis. *ACS Synth. Biol.* **2013**, *2* (2), 83–92.
https://doi.org/10.1021/sb300037w.

503