Global signal regression reduces connectivity patterns related to physiological signals and does not alter EEG-derived connectivity

Alba Xifra-Porxas [a], Michalis Kassinopoulos [a], Prokopis Prokopiou [b], Marie-Hélène Boudrias [c,d], Georgios. D. Mitsis [e]

[a] Graduate Program in Biological and Biomedical Engineering, McGill University, Montréal, Canada
[b] Integrated Program in Neuroscience, McGill University, Montréal, Canada
[c] School of Physical and Occupational Therapy, McGill University, Montréal, Canada
[d] Center for Interdisciplinary Research in Rehabilitation of Greater Montreal (CRIR), Montréal, Canada
[e] Department of Bioengineering, McGill University, Montréal, Canada

ABSTRACT

Functional brain connectivity measures extracted from resting-state functional magnetic resonance
imaging (fMRI) scans have generated wide interest as potential noninvasive biomarkers. In this
context, performing global signal regression (GSR) as a preprocessing step remains controversial.
Specifically, while it has been shown that a considerable fraction of global signal variations is
associated with physiological and motion sources, GSR may also result in removing neural
activity. Here, we address this question by examining the fundamental sources of resting global
signal fluctuations using simultaneous electroencephalography (EEG)-fMRI data combined with
cardiac and breathing recordings. Our results suggest that systemic physiological fluctuations
account for a significantly larger fraction of global signal variability compared to
electrophysiological fluctuations. Furthermore, we show that GSR reduces artifactual connectivity
due to heart rate and breathing fluctuations, but preserves connectivity patterns associated with
electrophysiological activity within the alpha and beta frequency ranges. Overall, these results
provide evidence that the neural component of resting-state fMRI-based connectivity is preserved
after the global signal is regressed out.

Keywords: global signal, simultaneous EEG-fMRI, physiological confounds, functional
connectivity, fMRI preprocessing, brain networks

Corresponding author: Georgios D. Mitsis, Department of Bioengineering, McGill University.
Montréal, Canada. Email: georgios.mitsis@mcgill.ca

1. INTRODUCTION

Studying the brain at rest has become a powerful tool towards revealing intrinsic characteristics of functional brain organization. Even in the absence of overt behaviour, brain activity fluctuates in an organized fashion in the form of large-scale brain networks that resemble those observed during behavioral tasks (Biswal et al., 1995; Fox and Raichle, 2007; Gratton et al., 2018; Smith et al., 2009). Functional magnetic resonance imaging (fMRI) has been widely used for studying the resting-state activity of temporally correlated and spatially distributed brain regions, and has shown promise to uncover mechanisms underlying neurological disorders (Gratton et al., 2019; Kassinopoulos et al., 2021; Xia et al., 2018). However, the blood-oxygenated level dependent (BOLD) signal typically employed in fMRI studies is only partially attributed to neural activity. Specifically, it has been shown that BOLD signal fluctuations are also driven by thermal noise as well as physiological and motion artifacts (Liu, 2016; Murphy et al., 2013).

The BOLD fMRI signal relies on changes in local cerebral blood flow (CBF) to detect neural activity, which constitutes an important caveat, as several additional processes may induce fluctuations in CBF. These include changes in heart rate and breathing patterns (Birn et al., 2006; Shmueli et al., 2007), variations in CO_2 (Prokopiou et al., 2019; Wise et al., 2004), as well as arterial blood pressure (Whittaker et al., 2019a). Furthermore, the neurally driven fraction of the BOLD fMRI signal is not a direct measurement of neural activity, but an indirect measurement determined by neurovascular coupling mechanisms (Iadecola, 2017; Logothetis, 2008; Logothetis et al., 2001). It is also worth noting that the characteristics of physiological response functions that describe the dynamic effects of the aforementioned physiological signals (heart rate, respiration, arterial CO_2 etc.) on the fMRI signal are highly similar to those of the hemodynamic response function (Prokopiou et al., 2022; Wu and Marinazzo, 2016), which further complicates disentangling the neural and physiological fMRI signal sources.

The aggregate nature of the BOLD fMRI signal described above renders the removal of its non-neural components a crucial and challenging task, which is particularly exacerbated in resting-state studies, where there is no *a priori* assumption for the temporal pattern of the underlying neural activity. Although there have been many advances in resting-state fMRI denoising (Caballero-Gaudes and Reynolds, 2017; Ciric et al., 2018), there is still no gold standard for resting-state fMRI preprocessing. In particular, global signal regression (GSR), which involves regressing out the average fMRI signal across the whole brain (global signal) from every voxel, has been proposed as a pre-processing step (Aguirre et al., 1998; Fox et al., 2009; Macey et al., 2004). Yet, as the processes underpinning the global signal are still poorly understood, GSR has turned out to be one of the most contentious preprocessing steps in fMRI denoising (Liu et al., 2017; Murphy and Fox, 2017; Power et al., 2017a). The rationale behind GSR is that the global signal mostly encompasses non-neuronal processes arising from physiological sources, head motion and scanner artifacts (Kassinopoulos and Mitsis, 2021; Power et al., 2017b). Therefore, GSR represents an effective data-driven approach for removing these global fluctuations, as they may result in artifactual functional connectivity patterns (Burgess et al., 2016; Ciric et al., 2017; Parkes et al., 2018; Xifra-Porxas et al., 2020). Indeed, GSR has been shown to increase the similarity of functional connectivity estimates across modalities (Keller et al., 2013). In addition, recent studies have reported that GSR improves the identifiability of well-established resting-state networks (Kassinopoulos and Mitsis, 2022), connectome fingerprinting accuracy (Xifra-Porxas et al., 2021), as well as the association between resting-state functional connectivity and behavioral measures (Li et al., 2019b), which suggests the potential benefits of performing GSR.

Yet, there is also converging evidence from simultaneous electrophysiological-fMRI studies that
neural activity is linked to the global signal. Earlier investigations, albeit not directly examining
the global signal, showed that fluctuations in local field potentials exhibited fairly widespread
correlations with fMRI activity over the macaque brain (Scholvinck et al., 2010). More recently,
fluctuations of the global signal have been linked to electrophysiological indices of arousal (C. W.
Wong et al., 2016; Wong et al., 2013) and glucose metabolism (Thompson et al., 2016).
Furthermore, global resting-state fluctuations were found to at least partially stem from the basal
forebrain (Liu et al., 2018; Turchi et al., 2018). Gutierrez-Barragan and colleagues showed that
brain states occur at specific phases of global fMRI signal fluctuations in the mouse brain
(Gutierrez-Barragan et al., 2019). Finally, individual variations in global signal topography have
been associated with behavioral measures (Li et al., 2019a). All these reports suggest that GSR
may remove neuronal-related fluctuations of interest in functional connectivity studies.

Until now, the vast majority of studies investigating the processes underpinning the global signal
have probed its physiological or neural origins separately. The neurally-related fraction of the
global signal has been associated to fluctuations in arousal and vigilance, likely regulated by the
autonomic nervous system (Oken et al., 2006; Olbrich et al., 2011). However, these apparent
neural fluctuations could also be associated with changes in systemic physiological quantities such
as heart rate and breathing, which can in turn influence the fMRI signal. Hence, there is potentially
a closed loop path through which neural autonomic activity indirectly contributes to the global
signal through changes in physiological signals. Overall, the above observations suggest that
physiological and neural processes should be simultaneously taken into consideration to fully
elucidate the nature of the resting-state global signal, as well as quantify the extent to which neural
and physiological mechanisms generate similar or distinct contributions to global signal
fluctuations.

While GSR is a more prevalent technique in resting-state studies, it was originally developed for
task-based studies (Zarahn et al., 1997), and it is still frequently applied in this setting (e.g. (Finn
et al., 2015; Marek et al., 2018)). Glasser et al. (2018) found that application of GSR to task fMRI
data led to reduced statistical sensitivity for detecting activations, suggesting that the global signal
in task-based studies contains relevant neural signal. However, the contributions of neurally-
related fluctuations to the global signal during behavioral tasks have not yet been elucidated, as
previous simultaneous electrophysiological-fMRI investigations of the global signal only
examined resting-state conditions.

In the present study, we used simultaneously acquired electroencephalography (EEG)-fMRI data,
as well as physiological recordings (heart rate and respiration), to quantify the unique and shared
contributions of physiological and neural processes to the global fMRI signal, both at rest and
during a hand-grip task. Furthermore, we generated two synthetic fMRI datasets that consisted of
systemic and electrophysiological fluctuations, respectively, and evaluated the similarity between
connectivity estimates extracted from the synthetic and experimental fMRI datasets. This allowed
us to examine the effects of GSR on connectivity estimates and address explicitly whether
eliminating the bias introduced by physiological processes inadvertently removes connectivity
patterns related to electrophysiological activity.

2. MATERIALS AND METHODS

2.1 Participants

A total of 12 healthy volunteers (25.1 ± 2.9 years; 4 female) participated in the study. All subjects were right-handed according to the Edinburgh Handedness Inventory (Oldfield, 1971) and had no history of neurological or psychiatric disorders. The study was approved by the McGill University Ethical Advisory Committee. All participants signed a written informed consent and were compensated for their participation.

2.2 Experimental paradigm

The protocol carried out inside the MR scanner consisted of two 15-min resting-state runs, with a 13-min hand-grip fMRI task interleaved between them (Figure 1a). During the resting-state periods, subjects were instructed to stare at a white fixation cross displayed on a dark background and not to think of anything in particular. After the first rest period, the maximum voluntary contraction (MVC) was obtained for each participant, using the same hand gripper subsequently employed for the motor task. The motor task was a unimanual isometric right hand-grip, during which the subjects had to apply force to track a ramp target as accurately as possible (Larivière et al., 2019; Xifra-Porxas et al., 2019). At the onset of the trial, an orange circle appeared on the screen, and the subjects had 2 s to increase their force to reach a white target block at 15% of their MVC. This force was held for 3 s. Following this, participants tracked a linear increase of the force to reach 30% of their MVC over a 3-s period, during which they had to maintain the circle inside the white target block, followed by a 3-s hold at 30% of their MVC (Figure 1b). A single trial lasted 11 s and the inter-trial interval was jittered between 3 and 5 s, during which subjects stared at a white cross. The task consisted of 50 trials, resulting in a total duration of about 13 min.

2.3 Data acquisition: EEG-fMRI data and physiological recordings

All experiments were conducted at the McConnell Brain Imaging Centre (BIC) of the Montreal Neurological Institute (MNI), McGill University, using a 3T Siemens MAGNETOM Prisma fit MRI scanner (Siemens AG, Germany). A 32-channel head coil was used to acquire whole-brain T2*-weighted functional gradient echo planar (EPI) data ($3\times3\times4$ mm^3 voxels, TR=2120 ms, TE=30 ms, flip angle=90°, FOV=192×192 mm^2, anterior-posterior phase encoding direction). Volumes were recorded in 35 transverse slices in descending order. A high-resolution structural volume was also acquired using a T1-weighted magnetization-prepared rapid acquisition gradient echo (MPRAGE) sequence (TR=2300 ms, TE=2.32 ms, flip angle=8°, FOV=240×240 mm^2, 0.9 mm^3 isotropic voxels).

EEG data were simultaneously recorded with an MR-compatible 64-channel system with Ag/AgCl ring-type electrodes (BrainAmp MR, Brain Products GmbH, Germany), sampled at 5000 Hz. Electrode impedances were maintained below 20 kΩ and verified both during setup and at the end of each scanning session. An equidistant electrode layout was used with AFz and Cz as ground and online recording reference, respectively. The EEG acquisition clock was synchronised with the MR scanner clock through a device that sent triggers to the EEG recording system every time an fMRI volume was acquired (TriggerBox, Brain Products GmbH, Germany). The electrodes were precisely localized using a 3-D electromagnetic digitizer (Polhemus Isotrack, USA).

Cardiac and breathing measurements were continuously recorded throughout the experiment using
a pulse oximeter and respiratory belt (BIOPAC Systems, Inc., USA). An MR-compatible hand
clench dynamometer (BIOPAC Systems, Inc., USA) was used to measure the subjects' hand-grip
strength during the motor paradigm. The pulse oximeter, respiratory belt and dynamometer were
connected to an MP150 data acquisition system (BIOPAC Systems, Inc., USA) from which the
signals were transferred to a computer, sampled at 1000 Hz.

2.4 Preprocessing

2.4.1 fMRI data

The fMRI data were preprocessed using FSL (Jenkinson et al., 2012). Automated brain extraction
using BET, motion correction via volume realignment using MCFLIRT, spatial smoothing (5 mm
FWHM Gaussian kernel) and high-pass temporal filtering (100 s cutoff) were performed.
Additional preprocessing included motion censoring based on the frame-wise displacement (FD)
and root mean square intensity change of BOLD signal across the whole brain (DVARS) measures
(Power et al., 2012). Motion censoring was applied by discarding volumes with FD>0.25 mm or
when DVARS exceeded its median absolute deviation by a factor of 3, as well as their adjacent
volumes. Volumes with subthreshold values of FD and DVARS were also discarded if they were
preceded and followed by flagged volumes. All scans from one participant were excluded from
the analysis, since more than 40% of volumes were identified as being contaminated by motion
(Suppl. Fig. 1). Therefore, results from a total of 11 subjects are presented below.

The censored fMRI data were co-registered to each subject's structural image and normalized to
MNI space (2 mm). Subsequently, the preprocessed data were parcellated into 300 regions of
interest (ROIs) using the Seitzman atlas (Seitzman et al., 2020). In that work, ROIs were assigned
to large-scale functional networks using the InfoMap community detection algorithm (Rosvall and
Bergstrom, 2008), applied to resting-state functional connectivity data. The fifteen ROIs that were
not assigned to any large-scale network in Seitzman et al. (2020) were excluded here from further
analysis, resulting in a final set of 285 ROIs.

We examined three sets of experimental fMRI data that differed in the set of nuisance regressors
removed:

(i) minimally preprocessed (MinPrep) fMRI data, with no additional denoising;

(ii) preprocessed fMRI data where white matter denoising was applied; and

(iii) preprocessed fMRI data with global signal regression (GSR), where both white matter
denoising and GSR was performed.

In the second dataset, white matter denoising was performed using principal component analysis
(PCA) on the parcellated data to mitigate the effects of head motion not corrected by motion
censoring as well as the effects of cardiac pulsatility and breathing-related motion (Behzadi et al.,
2007; Kassinopoulos and Mitsis, 2022; Muschelli et al., 2014). Specifically, 10 white matter PCA
components were removed from the data, as this number was found to maximize the functional
connectivity contrast (FCC; Figure 2), which is a quality control metric for evaluating the
identifiability of well-established large-scale networks (Kassinopoulos and Mitsis, 2022). In the

third dataset, the global signal and the same 10 white matter PCA components were simultaneously
regressed out.

2.4.2 EEG data

EEG data were corrected offline for gradient and ballistocardiogram (BCG) artifacts using sliding
average template subtraction in BrainVision Analyzer 2 (Brain Products GmbH, Germany). The
data were subsequently downsampled to 200 Hz and further preprocessed using independent
component analysis to remove remaining non-neural components such as ocular and muscle
artifacts, as well as gradient and BCG residuals. Non-neural components were identified through
visual inspection by three of the paper's authors (AXP, MK, PP), based on classification guidelines
provided in the ICLabel tutorial (https://labeling.ucsd.edu/tutorial/labels). The preprocessed data
were re-referenced using the average signal from all channels.

Time-frequency spectrograms were calculated for each EEG channel using the continuous wavelet
transform (CWT) with analytic Morlet wavelets in Matlab environment. Matlab's default settings
were used, which apply logarithmically spaced frequencies and 10 voices per octave. The
frequency range was automatically determined based on the signal duration and sampling rate (i.e.
200 Hz). A global spectrogram was then obtained by calculating the root mean square across all
channel-specific spectrograms. Then, we extracted the instantaneous power time-series through
averaging of the global spectrogram within the following frequency bands: delta (1.5-4 Hz), theta
(4-8 Hz), alpha (8-15 Hz) and beta (15-26 Hz). These global EEG power time-series were
subsequently convolved with a canonical double-gamma haemodynamic response function and
downsampled to the frequency of image volume sampling (TR=2.12 s). The censored frames
flagged during fMRI preprocessing were removed from the convolved EEG power time-series.

2.4.3 Physiological recordings

Beat-to-beat intervals were detected from the pulse oximeter signal, and the heart rate signal was
computed as the inverse of the time differences between pairs of adjacent peaks and converted to
beats-per-minute (bpm). Heart rate traces were visually examined to identify outliers, and an
outlier replacement filter was used to eliminate spurious changes in heart rate. The breathing signal
from the respiratory belt was detrended linearly, visually inspected and corrected for outliers using
a replacement filter, low-pass filtered at 5 Hz, and z-scored. The respiratory flow, proposed in
Kassinopoulos and Mitsis (2019) as a robust measure of the absolute flow of inhalation and
exhalation of a subject at each time point, was extracted by further smoothing the breathing signal
(moving average filter using a 1.5 sec window) and computing the square of the derivative of the
smoothed breathing signal. Both heart rate and respiratory flow signals were resampled to 10 Hz.

Following this, we modelled the effect of systemic low-frequency oscillations (SLFOs) associated
with heart rate and breathing patterns using a recently developed method that computes scan-
specific physiological response functions (PRFs) (Kassinopoulos and Mitsis, 2019; scripts
available on https://github.com/mkassinopoulos/PRF_estimation/). Briefly, this algorithm
estimates PRF curves so that the convolution between heart rate and respiratory flow with their
corresponding PRFs optimizes the fit on the global signal of the same scanning session, while
ensuring that the shapes of the PRF curves are physiologically plausible. The heart rate and
respiratory flow were convolved with their respective PRFs and added to obtain a time-series that
reflects the total effect of SLFOs.

It is important to note that SLFOs reflect slow physiological modulations occurring at low
frequencies up to ~0.15 Hz, such as fluctuations in heart rate and breathing patterns. These
processes unfold over time scales of ~10 seconds and are well captured by our sampling rate [1/TR
= 1/(2.120 s) ≈ 0.47 Hz], which satisfies the Nyquist criterion. While cardiac pulsatility artifacts
occur at higher frequencies (~1 Hz) and are subject to aliasing at typical TRs, our study does not
aim to model such high-frequency effects. Instead, the PRF-based approach targets the slower
systemic fluctuations that influence the BOLD signal more gradually and are not significantly
impacted by the TR used in our acquisition.

2.5 Data analysis

We initially quantified the variance of the fMRI global signal explained by SLFOs and EEG power
using partial correlation. Specifically, the contribution of SLFOs to the global signal was computed
controlling for the EEG power time-series and vice versa. To assess the significance of these
contributions to the global signal, surrogate data were generated via inter-subject surrogates
(Lancaster et al., 2018), using physiological and EEG power time-series from different subjects.
The contributions obtained using the experimental data from the same subject were compared to
the ones obtained from the surrogate data using the Wilcoxon rank-sum test. The significance level
was set to 0.05, and the p-values for the EEG bands were adjusted for multiple comparisons using
the false discovery rate (FDR) method.

In addition, we generated three synthetic fMRI datasets, namely the SLFOs synthetic fMRI, alpha
power synthetic fMRI, and beta power synthetic fMRI, following the framework employed in
Xifra-Porxas et al. (2021). In brief, the ROI fMRI time-series $y(t)$ in each of the 285 atlas-based
ROIs was modelled using multiple linear regression, as follows:

$$y(t) = \beta_0 + \beta_{SLFOs}x_{SLFOs}(t) + \boldsymbol{\beta}_{EEG}\boldsymbol{x}_{EEG}(t) + \varepsilon(t) \qquad (1)$$
$$\boldsymbol{\beta}_{EEG} = [\beta_{alpha} \quad \beta_{beta}]$$
$$\boldsymbol{x}_{EEG}(t) = \begin{bmatrix} x_{alpha}(t) \\ x_{beta}(t) \end{bmatrix}$$

where $x_{SLFOs}(t)$ is the physiological regressor related to SLFOs, $\boldsymbol{x}_{EEG}(t)$ are the two EEG power
time-series, $\{\beta_0, \beta_{SLFOs}, \boldsymbol{\beta}_{EEG}\}$ are the unknown parameters estimated for each of the three
experimental datasets, and ε is the error.

Then, for each of the three processes of interest (i.e. SLFOs, alpha and beta EEG power), the
estimated value $\hat{\beta}$ was multiplied by its corresponding regressor to obtain the associated
fluctuations ($\hat{y}_{PI}(t)$) within a specific ROI, and a 'clean' time-series ($\hat{y}_{PI+res}(t)$) was calculated
via removal of all other regressors, as follows:

$$\text{SLFOs: } \hat{y}_{PI}(t) = \hat{\beta}_{SLFOs}\, x_{SLFOs}(t) \qquad (2)$$
$$\hat{y}_{PI+res}(t) = y(t) - \hat{\beta}_0 - \hat{\beta}_{alpha}x_{alpha}(t) - \hat{\beta}_{beta}x_{beta}(t) \qquad (3)$$

$$\text{Alpha EEG power: } \hat{y}_{PI}(t) = \hat{\beta}_{alpha}\, x_{alpha}(t) \qquad (4)$$
$$\hat{y}_{PI+res}(t) = y(t) - \beta_0 - \hat{\beta}_{SLFOs}\, x_{SLFOs}(t) - \hat{\beta}_{beta}x_{beta}(t) \qquad (5)$$

$$\text{Beta EEG power: } \hat{y}_{PI}(t) = \hat{\beta}_{beta}\, x_{beta}(t) \qquad (6)$$

$$\hat{y}_{PI+res}(t) = y(t) - \beta_0 - \hat{\beta}_{SLFOs} x_{SLFOs}(t) - \hat{\beta}_{alpha} x_{alpha}(t) \tag{7}$$

261 In this manner, we generated 'cleaned' ROI time-series ($\hat{y}_{PI+res}(t)$) in which all regressors were
262 removed except the one corresponging to the specific process being evaluated. Subsequently, the
263 contribution of the latter to the remaining fluctuations within each ROI was quantified. To achieve
264 this, the estimated process signal was correlated to the 'clean' ROI time-series:

$$r_{PI} = corr\left(\hat{y}_{PI}(t), \hat{y}_{PI+res}(t)\right) \tag{8}$$

265 Afterwards, the estimated process signal was subtracted from the 'clean' ROI time-series to obtain
266 the 'residual' time-series. These time-series were correlated to the 'clean' time-series to quantify
267 the contribution of the 'residual' variations to the 'clean' ROI time-series:

$$\hat{y}_{res}(t) = \hat{y}_{PI+res}(t) - \hat{y}_{PI}(t) \tag{9}$$

$$r_{res} = corr\left(\hat{y}_{res}(t), \hat{y}_{PI+res}(t)\right) \tag{10}$$

268 Finally, synthetic datasets were constructed for each process by combining the normalized process-
269 specific signal and a first-order autoregressive (AR(1)) process ($\psi(t)$) scaled by their respective
270 correlation coefficients:

$$y_{PI}(t) = r_{PI} Z\left[\hat{y}_{PI}(t)\right] + r_{res} Z[\psi(t)] \tag{11}$$

where Z[·] denotes normalization to zero mean and unit variance.

Using the framework described above, we constructed synthetic fMRI datasets in which the
fluctuations attributed to the process of interest within each ROI was preserved while the
remaining fraction of fluctuations was replaced with uncorrelated random signal of equal
amplitude (Xifra-Porxas et al., 2021). We only considered the alpha and beta power among the
EEG bands because they were found to be the only ones contributing to the global signal (Figure
3; $p < 0.05$).

We then computed the static functional connectivity matrices (i.e. pairwise correlations between
time-series of all ROIs) for each of the three experimental datasets described earlier: minimally
preprocessed experimental fMRI, preprocessed experimental fMRI with white matter denoising,
and preprocessed experimental fMRI with both white matter denoising and GSR; as well as for
the three synthetic datasets: SLFOs synthetic fMRI, alpha power synthetic fMRI, and beta power
synthetic fMRI. Finally, we quantified the similarity between the SLFOs, alpha power and beta
power connectivity matrices with respect to the fMRI connectivity matrices derived from the three
experimental datasets, by computing the correlations between their upper triangular values. High
similarity is expected when a process, whether SLFOs or EEG band power, explains a substantial
portion of BOLD variance across multiple ROIs, resulting in elevated inter-regional correlations
among those regions and imposing a spatial covariance structure reflected in both the synthetic
and empirical functional connectivity matrices. The derived correlations were utilized to assess the
effect of GSR on the similarity between SLFOs and fMRI connectivity, as well as EEG power and
fMRI connectivity.

3. RESULTS

3.1 Association of SLFOs and EEG power with the fMRI global signal

All signals examined (i.e. fMRI global signal, SLFOs, and EEG band power signals) exhibited prominent low-frequency fluctuations below 0.15 Hz, as shown in Supplementary Figure 2. In line with previous studies (Birn et al., 2006; Kassinopoulos and Mitsis, 2019; Power et al., 2017a; Shmueli et al., 2007), SLFOs were strongly correlated with the global signal, both at rest and during the motor task (Figure 3). Regarding the EEG bands, alpha power was negatively correlated with the global signal both at rest and during the motor task (Figure 3), and beta power was negatively correlated with the global signal during the motor task (Figure 3b). Illustrative individual time series from representative subjects can be found in Suppl. Fig. 2. It is worth noting that the correlation strength of alpha and beta fluctuations with the global signal, albeit significant, was considerably weaker compared to the correlation between SLFOs and the global signal. The mean correlation value across subjects for the full model containing both SLFOs and significant EEG power bands was equal to 0.64 ± 0.12 (resting-state 1), 0.64 ± 0.09 (motor task), and 0.59 ± 0.09 (resting-state 2). Moreover, alpha power fluctuations were found to be negatively correlated with SLFOs ($R = -0.09, p = 0.02$) during resting-state 1. This correlation vanished when alpha fluctuations and SLFOs were controlled for global signal fluctuations (R=0.02, p=0.84), which may suggest a common component between those two variables that is reflected on the global signal, possibly reflecting a direct or indirect effect of autonomic activity. Yet, we were unable to replicate this result during the motor task or resting-state 2 (i.e. alpha power was not significantly correlated with SLFOs). Finally, delta and theta power fluctuations did not yield a consistent contribution to the global signal across subjects for any of the scans.

3.2 Effect of GSR on connectome patterns associated with SLFOs and EEG power fluctuations

For each subject and scanning session, connectivity matrices were derived using the synthetic dataset generated from SLFOs, enabling us to identify the connectivity edges more prone to exhibit artifactual connectivity attributable to SLFOs. We found that fluctuations due to SLFOs artifactually increased the connectivity across most brain networks and particularly within the visual network, as well as between the visual network and other brain networks (Figure 4a), consistent with previous observations using a larger cohort (Xifra-Porxas et al., 2021). Furthermore, the effect of SLFOs on the visual network were found to be spatially homogeneous at rest, whereas during the motor task, some edges were more affected than others (Figure 4a). Notably, the contribution of SLFOs to functional connectivity appeared to be attenuated during the second resting-state session compared to the first, suggesting session-related variability in SLFOs-induced artifacts. Consistent with our earlier study (Xifra-Porxas et al., 2021), WM denoising led to a significant reduction in similarity between the SLFOs' and the fMRI connectivity matrices both at rest and during the motor task (Figure 4b; $p < 0.001$). Further, GSR yielded a small additional reduction in similarity for all three scans ($p < 0.05$). This result supports the extensive evidence that, at least with respect to mitigating nuisance processes, GSR is effective.

Subsequently, we sought to determine whether applying GSR also removed substantial EEG power fluctuations. First, we derived connectivity matrices using the fMRI synthetic datasets generated using the BOLD fluctuations attributed to alpha and beta power activity, thus highlighting the edges underpinning connectivity patterns of likely neural origin. At rest, alpha power variations mostly contributed to connectivity within the visual network and dorsal attention network as well

as between these two networks (Figure 5a). On the other hand, during the motor task, alpha power
variations mostly contributed to connectivity within the default mode network (Figure 5a), whereas
beta power variations mainly contributed to connectivity within some edges of the visual network
(Figure 5c).

Finally, we evaluated whether performing GSR on the preprocessed fMRI data significantly
removed the connectivity signature of these neural processes. To do this, we compared the
similarity of the connectome patterns associated with EEG activity with the connectivity patterns
of the three experimental fMRI datasets. WM denoising was found to increase the similarity
between modalities for all scans (Figure 5b & 5d; $p < 0.05$). As seen in Figure 5b & 5d, GSR did
not have any significant effect on the similarity between the EEG-related connectivity signatures
and fMRI connectivity matrices, neither for alpha nor beta power variations.

4. DISCUSSION

GSR is a widely used preprocessing step to remove global artifacts (mostly heart rate and breathing effects, collectively termed SLFOs) from fMRI data (Ciric et al., 2017; Parkes et al., 2018; Power et al., 2018, 2017b; Kassinopoulos & Mitsis, 2021), but remains controversial because it may also discard neural signals (Liu et al., 2017; Murphy and Fox, 2017). Up to date, the vast majority of studies investigating the origins of the global signal and evaluating the potential effects of GSR have examined fMRI data concurrently with either physiological or electrophysiological recordings, but not both at the same time. In the present study, we used simultaneous EEG-fMRI data, as well as cardiac and breathing recordings, to examine the processes underpinning the global signal and the impact of GSR on measures of brain activity and connectivity related to neural and systemic physiological fluctuations. Our results suggest that the global signal is strongly associated with physiological processes (R ~ 0.6) and more weakly associated with EEG power fluctuations (R ~ -0.1). We further demonstrated that GSR effectively removes the connectome patterns induced by physiological processes (SLFOs), but preserves the connectome patterns associated with EEG alpha and beta power fluctuations. These results provide evidence that, in the context of connectivity analyses, GSR improves the denoising of fMRI data and does not seem to alter the connectivity profiles associated with electrophysiological activity.

4.1 Association of SLFOs and EEG power with the fMRI global signal

We first evaluated the unique contributions of SLFOs and EEG activity to the global signal. SLFOs were found to explain a considerable fraction of global signal variance at rest (Figure 3a,3c), consistent with several earlier studies (Birn et al., 2006; Chang and Glover, 2009; Erdoğan et al., 2016; Kassinopoulos and Mitsis, 2019; Power et al., 2017b), as well as during the motor task (Figure 3b). Alpha power fluctuations were found to be negatively correlated with the global signal for all scans (Figure 3a-c), consistent with earlier work (Wong et al., 2013; Liu et al., 2017; Han et al., 2019), and beta power was found to be negatively correlated with the global signal during the motor task (Figure 3b). Our results suggest that contributions of electrophysiological origin to the global signal, albeit significant, were substantially weaker than physiological contributions.

A possible explanation for this is that the sign of correlation between fMRI fluctuations and EEG activity may not be the same across brain regions and, thus, when averaging fMRI signals across the whole brain, the contributions of these electrophysiological fluctuations may cancel out to some extent. Several studies have reported that, during resting conditions, alpha activity is negatively correlated with sensorimotor areas but positively correlated with default-mode regions (Mantini et al., 2007; Mayhew and Bagshaw, 2017; Scheeringa et al., 2012). These trends were also observed in our data, resulting in the presence of both positively and negatively correlated activity in Figure a. Therefore, it is indeed likely that the weak relationship between global signal and EEG activity may be due to differences in the polarity of associated fMRI activity across regions.

4.2 Association between SLFOs and alpha power

We observed that alpha power and SLFOs were negatively correlated during resting-state 1, albeit weakly. This finding was not replicated during the motor task and resting-state 2. The weak association between SLFOs and alpha activity may be due to the fact that subjects had their eyes open, which is in agreement to previous work that has demonstrated an association between

393 respiration and alpha power during an eyes closed condition but not during eyes open (Yuan et al.,
394 2013). Independent of this, shared contributions from SLFOs and alpha activity to the global
395 signal, which could potentially reflect a direct or indirect effect of autonomic activity, were not
396 consistently observed in our data. Therefore, our results seem to indicate that the major contributor
397 to the global signal was physiological in origin.

4.3 Effect of GSR on connectome patterns associated with SLFOs and EEG power

We initially assessed the systematic effect of SLFOs on estimates of functional connectivity. The
grand-averaged functional connectivity matrices calculated from the SLFOs synthetic datasets
exhibited a heterogeneous pattern, characterized by stronger correlations within and between
sensory cortices, including the visual cortex, somatosensory cortex and auditory cortex, as well as
subcortical regions such as the thalamus and cerebellum (Figure 4a). This heterogeneity among
brain regions is not surprising as it has been reported that global signal fluctuations are non-
uniformly distributed across the brain (Fox et al., 2009; Kassinopoulos and Mitsis, 2019; Power et
al., 2017b). Furthermore, the artifactual connectivity patterns due to SLFOs observed in the present
study were similar to those reported in our recent study where a large number of healthy subjects
from the Human Connectome Project (HCP) dataset was used for cortical regions (Xifra-Porxas
et al., 2020). However, our results also revealed artifactual connectivity patterns within and
between subcortical regions such as the thalamus and cerebellum, which we were unable to
observe in the HCP dataset, likely due to poor signal-to-noise ratio in the subcortex in the HCP
data (Ji et al., 2018; Seitzman et al., 2020).

Importantly, we observed a task- and session-related modulation of SLFOs-induced connectivity
patterns (Figure 4a). Specifically, contributions of SLFOs were weaker during the motor task and
the second resting-state period compared to the first. This observation may seem paradoxical as
the contribution of SLFOs to the global signal did not significantly decrease across scans (Figure
3), but could be explained by the fact that global signal fluctuations were found to be reduced
during the motor task and the second resting-state period compared to the first resting-state period
(Supp. Fig. 3). It may also indicate that subjects were more alert during and after the motor task
(Wong et al., 2013; Yeo et al., 2015). This interpretation is further supported by the decreased
variability in heart rate observed during the second resting-state period compared to the first (p =
0.04; Supp. Fig. 5), suggesting a possible reduction in autonomic fluctuations. Finally, it could
reflect the effect of circadian/ultradian physiological fluctuations, given that scanning started at 6
pm for all subjects and it has been reported that the amplitude of global signal fluctuations
decreases as the day progresses (Orban et al., 2020).

The grand-averaged functional connectivity matrices calculated from the fraction of BOLD
variance explained by alpha and beta power fluctuations also revealed structured patterns (Figure
5a, 5c). At rest, fluctuations in alpha power were associated with correlated activity within and
between visual and dorsal attention networks (Figure 5a). Fluctuations in alpha power were also
associated with anticorrelations of visual and dorsal attention networks with the default mode
network and several subcortical regions. These results are consistent with earlier work reporting a
positive association of alpha power fluctuations with fMRI activity in the default mode network
and several subcortical regions, as well as a negative association between alpha power and fMRI
activity in sensory regions (Bowman et al., 2017; Jann et al., 2009; Mantini et al., 2007; Mayhew
and Bagshaw, 2017; Mo et al., 2013; Moosmann et al., 2003; Scheeringa et al., 2012). During the

motor task, we found that alpha power was positively correlated with fMRI connectivity within
the default mode network (Figure 5a), possibly reflecting the analogy between task-related
synchronization/desynchronization alpha patterns and activation/deactivation default mode
fluctuations (Mayhew et al., 2013; Mo et al., 2013). Furthermore, fluctuations in beta power were
mostly associated with positively and negatively correlated activity within the visual and default
mode networks, but surprisingly not in the somatosensory network (Figure 5c).

Regarding the effect of GSR on the estimates of functional connectivity, our recent work (Xifra-
Porxas et al., 2020) provided evidence that GSR is effective in terms of removing systematic biases
on measures of resting-state functional connectivity arising due to SLFOs. However, since we did
not have electrophysiological recordings in that study, we were unable to assess whether GSR also
removed any signal of interest. Here, we show that, in addition to mitigating the effect of sLFOs
on resting-state connectivity patterns, GSR yielded similar results during the motor task (Figure
4b). It is worth noting that even though we used a relatively aggressive preprocessing pipeline (10
PCA components from white matter) (Kassinopoulos and Mitsis, 2022), performing GSR was
found to be beneficial. Considering that in a large number of fMRI studies only the average signals
from white matter and cerebrospinal fluid compartments are regressed out, our results suggest that
the effectiveness of GSR with respect to artifact removal would have been even more pronounced
for milder preprocessing pipelines. Furthermore, we found that EEG power fluctuations within the
alpha and beta bands were correlated to the global signal (Figure 3) but the extent of shared
variance was small and, thus, GSR did not have any significant effect on the fMRI connectivity
patterns attributed to alpha (Figure 5b) and beta (Figure 5d) electrophysiological power, neither at
rest nor during the motor task. This is to some extent supported by the findings reported in a study
involving macaque monkeys, where it was shown that, while inactivation of the basal forebrain
led to selective suppression of ipsilateral global components, resting-state brain networks
preserved their distinctive topography (Turchi et al., 2018). Overall, our results support the view
that the benefits of GSR in terms of fMRI denoising outweigh the potential loss of neural
information (Li et al., 2019a, 2019b).

4.4 Limitations

A key limitation of the present study is the relatively small sample size (n = 11), which we
acknowledge may constrain the generalizability of our findings and may also contribute to the non-
significant effects of GSR on the connectivity patterns related to neural activity. However, the
primary aim of this work was to establish a methodological proof-of-concept regarding the
differential impact of global signal regression on connectivity patterns linked to physiological
signals and neural oscillations. Despite the known inter-subject variability in EEG-fMRI studies,
the consistency of our results across subjects and scanning sessions suggests that the observed
effects are robust. Notably, we were able to replicate previous findings from a large cohort with
regards to the effect of GSR on connectivity patterns induced by SLFOs (Xifra-Porxas et al., 2021),
which lends support to the validity of our results. Further research with larger sample sizes will be
essential to provide further insights into the nuanced impact of GSR on neural connectivity
patterns.

Furthermore, despite the large fraction of the global signal variance that was explained by SLFOs,
and to a lesser extent by EEG power (Figure 3), there was still variance unaccounted for. The EEG
signal is not a perfect reflection of neural activity and is known to be considerably noisy in the MR
environment, as well as mostly blind to deeper sources. Therefore, it is likely that there was an

additional fraction of neural activity that was present in the global signal that we were unable to
detect in the EEG data. Likewise, there are other physiological factors not considered here that are
known to give rise to SLFOs and are reflected on the global signal, such as finger skin vascular
tone (Kassinopoulos and Mitsis, 2021; Özbay et al., 2019), arterial CO2 changes (Prokopiou et al.,
2019; Wise et al., 2004) and arterial blood pressure (Dagenais and Mitsis, 2023; Whittaker et al.,
2019b). Future studies using more direct surrogates of neuronal activity (e.g. intracranial EEG) are
needed to confirm whether crucial neural information is being removed through GSR.

Another limitation concerns the simplicity of the EEG indices used in this study. We focused on
global EEG power averaged across channels to remain consistent with prior work (e.g., Wong et
al., 2013; 2016), and to avoid increasing model complexity given our modest sample size.
However, we acknowledge that more spatially resolved EEG metrics, such as alpha asymmetry or
region-specific alpha spindles, have been widely applied in cognitive and affective neuroscience
(Ismail and Karwowski, 2020; Giannakakis et al., 2022), and may reveal stronger associations with
the fMRI global signal. These metrics have not yet been systematically examined in resting-state
EEG-fMRI studies. Future research with larger cohorts and improved spatial modeling will be
needed to clarify their relevance to global signal fluctuations.

Moreover, while our results suggest that GSR effectively mitigates physiological confounds
without substantially altering EEG-derived connectivity patterns, we cannot exclude the
possibility that GSR also suppresses neural signal components that are spatially widespread or
temporally synchronized across regions. Prior studies have shown that GSR can introduce anti-
correlations and reduce the amplitude of functional connectivity estimates (Murphy et al., 2009),
and that the global signal may contain meaningful neural fluctuations related to arousal and
vigilance (Liu et al., 2017). Therefore, it remains possible that subtle neural contributions to the
BOLD signal, particularly those not captured by global EEG power, are attenuated. Future studies
incorporating more sensitive electrophysiological markers, such as phase-based metrics or cross-
frequency coupling, may help clarify the extent to which GSR affects neural signal components.

Finally, we examined resting conditions and a hand-grip task, but results could be different during
other conditions, such as sleep (Duyn et al., 2020) or when investigating diurnal variations in large-
scale spontaneous brain activity (Orban et al., 2020).

5. CONCLUSIONS

Using simultaneous EEG-fMRI recordings, we demonstrated that the global BOLD fMRI signal
exhibits a significant association with SLFOs as well as EEG alpha and beta power fluctuations,
during resting conditions and motor task execution. In both cases, the effect of SLFOs was found
to be more pronounced. We also provide evidence that GSR effectively removes confounds in
functional connectivity induced by SLFOs, without severely disrupting the functional connectivity
patterns attributed to EEG alpha and beta activity, suggesting that it overall yields a beneficial
effect in terms of fMRI-based connectivity assessment.

FUNDING

This work was supported by funds from Fonds de la Recherche du Québec – Nature et Technologies (FRQNT) Team Grants 2016-191780 [MHB & GDM] and 2018-256480 [GDM], the Natural Sciences and Engineering Research Council of Canada (NSERC) Discovery Grant RGPIN-2019-06638 [GDM], the Canadian Foundation for Innovation grant numbers 34277 [MHB] and 34362 [GDM], and the Canada First Research Excellence Fund (awarded to McGill University for the Healthy Brains for Healthy Lives (HBHL) initiative) [MK]. AXP and MK received financial support through funding from McGill University and the Québec Bio-imaging Network (QBIN).

ACKNOWLEDGMENTS

We would like to thank the MRI technicians at the Montreal Neurological Institute for their patience and assistance collecting the data.

DECLARATION OF CONFLICTING INTERESTS

The authors declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

STATEMENTS AND DECLARATIONS

Not applicable.

AUTHORS' CONTRIBUTIONS

AXP: conceptualization, study design, participant recruitment, data acquisition, preprocessing, formal analysis, figure design, writing & revision; MK: conceptualization, experimental paradigm implementation, data acquisition setup, data acquisition, preprocessing assistance, result interpretation, revision; PP: data acquisition, preprocessing assistance, result interpretation; MHB: study design, revision. GDM: conceptualization, supervision, revision.

DATA AVAILABILITY STATEMENT

The data that support the findings of this study are available from the corresponding author upon reasonable request.

ETHICAL APPROVAL AND INFORMED CONSENT

The study was approved by the McGill University Ethical Advisory Committee. All participants provided written informed consent before participation and were compensated for their participation.

SUPPLEMENTARY MATERIAL

Supplementary material for this article has been submitted and will be available online upon publication.

REFERENCES

Behzadi, Y., Restom, K., Liau, J., Liu, T.T., 2007. A component based noise correction method (CompCor) for BOLD and perfusion based fMRI. Neuroimage 37, 90–101. https://doi.org/10.1016/j.neuroimage.2007.04.042

Birn, R.M., Diamond, J.B., Smith, M.A., Bandettini, P.A., 2006. Separating respiratory-variation-related fluctuations from neuronal-activity-related fluctuations in fMRI. Neuroimage 31, 1536–1548. https://doi.org/10.1016/j.neuroimage.2006.02.048

Biswal, B., Yetkin, F.Z., Haughton, V.M., Hyde, J.S., 1995. Functional connectivity in the motor cortex of resting human brain using echo-planar MRI. Magn Reson Med. https://doi.org/10.1002/mrm.1910340409

Bowman, A.D., Griffis, J.C., Visscher, K.M., Dobbins, A.C., Gawne, T.J., DiFrancesco, M.W., Szaflarski, J.P., 2017. Relationship Between Alpha Rhythm and the Default Mode Network. Journal of Clinical Neurophysiology 34, 527–533. https://doi.org/10.1097/WNP.0000000000000411

Burgess, G.C., Kandala, S., Nolan, D., Laumann, T.O., Power, J.D., Adeyemo, B., Harms, M.P., Petersen, S.E., Barch, D.M., 2016. Evaluation of Denoising Strategies to Address Motion-Correlated Artifacts in Resting-State Functional Magnetic Resonance Imaging Data from the Human Connectome Project. Brain Connect 6, 669–680. https://doi.org/10.1089/brain.2016.0435

Caballero-Gaudes, C., Reynolds, R.C., 2017. Methods for cleaning the BOLD fMRI signal. Neuroimage 154, 128–149. https://doi.org/10.1016/j.neuroimage.2016.12.018

Chang, C., Glover, G.H., 2009. Effects of model-based physiological noise correction on default mode network anti-correlations and correlations. Neuroimage 47, 1448–1459. https://doi.org/10.1016/j.neuroimage.2009.05.012

Chang, C., Leopold, D.A., Schölvinck, M.L., Mandelkow, H., Picchioni, D., Liu, X., Ye, F.Q., Turchi, J.N., Duyn, J.H., 2016. Tracking brain arousal fluctuations with fMRI. Proceedings of the National Academy of Sciences 113, 4518–4523. https://doi.org/10.1073/pnas.1520613113

Ciric, R., Rosen, A.F.G., Erus, G., Cieslak, M., Adebimpe, A., Cook, P.A., Bassett, D.S., Davatzikos, C., Wolf, D.H., Satterthwaite, T.D., 2018. Mitigating head motion artifact in functional connectivity MRI. Nat Protoc 13, 2801–2826. https://doi.org/10.1038/s41596-018-0065-y

Ciric, R., Wolf, D.H., Power, J.D., Roalf, D.R., Baum, G.L., Ruparel, K., Shinohara, R.T., Elliott, M.A., Eickho, S.B., Davatzikos, C., Gur, R.C., Gur, R.E., Bassett, D.S., 2017. Benchmarking of participant-level confound regression strategies for the control of motion artifact in studies of functional connectivity. Neuroimage 154, 174–187. https://doi.org/10.1016/j.neuroimage.2017.03.020

Dagenais, R., Mitsis, G.D., 2023. Non-invasive estimation of arterial blood pressure fluctuations using a peripheral photoplethysmograph inside the MRI scanner. Annu Int Conf IEEE Eng Med Biol Soc 2023, 1–4. https://doi.org/10.1109/EMBC40787.2023.10340020

Duyn, J.H., Ozbay, P.S., Chang, C., Picchioni, D., 2020. Physiological changes in sleep that affect
fMRI inference. Curr Opin Behav Sci 33, 42–50.
https://doi.org/10.1016/j.cobeha.2019.12.007

Erdoğan, S.B., Tong, Y., Hocke, L.M., Lindsey, K.P., deB Frederick, B., 2016. Correcting for
blood arrival time in global mean regression enhances functional connectivity analysis of
resting state fMRI-BOLD signals. Front Hum Neurosci 10, 1–22.
https://doi.org/10.3389/fnhum.2016.00311

Giannakakis, G., Grigoriadis, D., Giannakaki, K., Simantiraki, O., Roniotis, A., Tsiknakis, M.,
2022. Review on Psychological Stress Detection Using Biosignals. IEEE Trans. Affect.
Comput. 13, 440–460.Falahpour, M., Chang, C., Wong, C.W., Liu, T.T., 2018. Template-
based prediction of vigilance fluctuations in resting-state fMRI. Neuroimage 174, 317–327.
https://doi.org/10.1016/j.neuroimage.2018.03.012

Finn, E.S., Shen, X., Scheinost, D., Rosenberg, M.D., Huang, J., Chun, M.M., Papademetris, X.,
Constable, R.T., 2015. Functional connectome fingerprinting: Identifying individuals using
patterns of brain connectivity. Nat Neurosci 18, 1664–1671. https://doi.org/10.1038/nn.4135

Fox, M.D., Raichle, M.E., 2007. Spontaneous fluctuations in brain activity observed with
functional magnetic resonance imaging. Nat Rev Neurosci 8, 700–711.
https://doi.org/10.1038/nrn2201

Fox, M.D., Zhang, D., Snyder, A.Z., Raichle, M.E., 2009. The global signal and observed
anticorrelated resting state brain networks. J Neurophysiol 101, 3270–83.
https://doi.org/10.1152/jn.90777.2008

Glasser, M.F., Coalson, T.S., Bijsterbosch, J.D., Harrison, S.J., Harms, M.P., Anticevic, A., Van
Essen, D.C., Smith, S.M., 2018. Using temporal ICA to selectively remove global noise while
preserving global signal in functional MRI data. Neuroimage 181, 692–717.
https://doi.org/10.1016/j.neuroimage.2018.04.076

Gratton, C., Koller, J.M., Shannon, W., Greene, D.J., Maiti, B., Snyder, A.Z., Petersen, S.E.,
Perlmutter, J.S., Campbell, M.C., 2019. Emergent Functional Network Effects in Parkinson
Disease. Cerebral Cortex 29, 2509–2523. https://doi.org/10.1093/cercor/bhy121

Gratton, C., Laumann, T.O., Nielsen, A.N., Greene, D.J., Gordon, E.M., Gilmore, A.W., Nelson,
S.M., Coalson, R.S., Snyder, A.Z., Schlaggar, B.L., Dosenbach, N.U.F., Petersen, S.E., 2018.
Functional Brain Networks Are Dominated by Stable Group and Individual Factors, Not
Cognitive or Daily Variation. Neuron 439–452. https://doi.org/10.1016/j.neuron.2018.03.035

Gutierrez-Barragan, D., Basson, M.A., Panzeri, S., Gozzi, A., 2019. Infraslow State Fluctuations
Govern Spontaneous fMRI Network Dynamics. Current Biology 29, 2295-2306.e5.
https://doi.org/10.1016/j.cub.2019.06.017

Iadecola, C., 2017. The Neurovascular Unit Coming of Age: A Journey through Neurovascular
Coupling in Health and Disease. Neuron 96, 17–42.
https://doi.org/10.1016/j.neuron.2017.07.030

Ismail, L.E., Karwowski, W., 2020. Applications of EEG indices for the quantification of human
cognitive performance: A systematic review and bibliometric analysis, PLoS ONE.
https://doi.org/10.1371/journal.pone.0242857

Jann, K., Dierks, T., Boesch, C., Kottlow, M., Strik, W., Koenig, T., 2009. BOLD correlates of
EEG alpha phase-locking and the fMRI default mode network. Neuroimage 45, 903–916.
https://doi.org/10.1016/j.neuroimage.2009.01.001

Jenkinson, M., Beckmann, C.F., Behrens, T.E.J., Woolrich, M.W., Smith, S.M., 2012. FSL.
Neuroimage 62, 782–790. https://doi.org/10.1016/j.neuroimage.2011.09.015

Ji, J.L., Kulkarni, K., Repovš, G., Spronk, M., Cole, M.W., Anticevic, A., 2018. Mapping the
human brain's cortical-subcortical functional network organization. Neuroimage 185, 35–57.
https://doi.org/10.1016/j.neuroimage.2018.10.006

Kassinopoulos, M., Harper, R.M., Guye, M., Lemieux, L., Diehl, B., 2021. Altered Relationship
Between Heart Rate Variability and fMRI-Based Functional Connectivity in People With
Epilepsy. Front Neurol 12, 1–13. https://doi.org/10.3389/fneur.2021.671890

Kassinopoulos, M., Mitsis, G.D., 2022. A multi-measure approach for assessing the performance
of fMRI preprocessing strategies in resting-state functional connectivity. Magn Reson
Imaging 85, 228–250. https://doi.org/10.1016/j.mri.2021.10.028

Kassinopoulos, M., Mitsis, G.D., 2021. Physiological noise modeling in fMRI based on the
pulsatile component of photoplethysmograph. Neuroimage 242, 118467.
https://doi.org/10.1016/j.neuroimage.2021.118467

Kassinopoulos, M., Mitsis, G.D., 2019. Identification of physiological response functions to
correct for fluctuations in resting-state fMRI related to heart rate and respiration. Neuroimage
202. https://doi.org/10.1101/512855

Keller, C.J., Bickel, S., Honey, C.J., Groppe, D.M., Entz, L., Craddock, R.C., Lado, F.A., Kelly,
C., Milham, M., Mehta, A.D., 2013. Neurophysiological investigation of spontaneous
correlated and anticorrelated fluctuations of the BOLD signal. Journal of Neuroscience 33,
6333–6342. https://doi.org/10.1523/JNEUROSCI.4837-12.2013

Lancaster, G., Iatsenko, D., Pidde, A., Ticcinelli, V., Stefanovska, A., 2018. Surrogate data for
hypothesis testing of physical systems. Phys Rep 748, 1–60.
https://doi.org/10.1016/j.physrep.2018.06.001

Li, J., Bolt, T., Bzdok, D., Nomi, J., Yeo, B.T.T., Spreng, R.N., Uddin, L.Q., 2019a. Topography
and behavioral relevance of the global signal in the human brain. Sci Rep 10, 1–5.
https://doi.org/10.1038/s41598-019-50750-8

Li, J., Kong, R., Liégeois, R., Orban, C., Tan, Y., Sun, N., Holmes, A.J., Sabuncu, M.R., Ge, T.,
Yeo, B.T.T., 2019b. Global signal regression strengthens association between resting-state
functional connectivity and behavior. Neuroimage 196, 126–141.
https://doi.org/10.1016/j.neuroimage.2019.04.016

Liu, T.T., 2016. Noise contributions to the fMRI signal: An overview. Neuroimage 143, 141–151.
https://doi.org/10.1016/j.neuroimage.2016.09.008

Liu, T.T., Nalci, A., Falahpour, M., 2017. The global signal in fMRI: Nuisance or Information?
Neuroimage 150, 213–229. https://doi.org/10.1016/j.neuroimage.2017.02.036

Liu, X., De Zwart, J.A., Schölvinck, M.L., Chang, C., Ye, F.Q., Leopold, D.A., Duyn, J.H., 2018.
Subcortical evidence for a contribution of arousal to fMRI studies of brain activity. Nat
Commun 9, 1–10. https://doi.org/10.1038/s41467-017-02815-3

Logothetis, N.K., 2008. What we can do and what we cannot do with fMRI. Nature 453, 869–878.

Logothetis, N.K., Pauls, J., Augath, M., Trinath, T., Oeltermann, A., 2001. Neurophysiological
investigation of the basis of the fMRI signal. Nature 412, 150–157.
https://doi.org/10.1038/35084005

Mantini, D., Perrucci, M.G., Gratta, D., C., R., G.l, Corbetta, M., Del Gratta, C., Romani, G., 2007.
Electrophysiological signatures of resting state networks in the human brain. Proceedings of
the National Academy of Sciences 104, 13170.

Marek, S., Siegel, J.S., Gordon, E.M., Raut, R. V., Gratton, C., Newbold, D.J., Ortega, M.,
Laumann, T.O., Miller, D.B., Zheng, A., Lopez, K.C., Berg, J.J., Coalson, R.S., Nguyen,
A.L., Dierker, D., Van, A.N., Hoyt, C.R., McDermott, K.B., Norris, S.A., Shimony, J.S.,
Snyder, A.Z., Nelson, S.M., Barch, D.M., Schlaggar, B.L., Raichle, M.E., Petersen, S.E.,
Greene, D.J., Dosenbach, N.U.F., 2018. Spatial and Temporal Organization of the Individual
Human Cerebellum. SSRN Electronic Journal 100, 977-993.e7.
https://doi.org/10.2139/ssrn.3188429

Mayhew, S.D., Bagshaw, A.P., 2017. Dynamic spatiotemporal variability of alpha-BOLD
relationships during the resting-state and task-evoked responses. Neuroimage 155, 120–137.
https://doi.org/10.1016/j.neuroimage.2017.04.051

Mayhew, S.D., Ostwald, D., Porcaro, C., Bagshaw, A.P., 2013. Spontaneous EEG alpha oscillation
interacts with positive and negative BOLD responses in the visual-auditory cortices and
default-mode network. Neuroimage 76, 362–372.
https://doi.org/10.1016/j.neuroimage.2013.02.070

Mo, J., Liu, Y., Huang, H., Ding, M., 2013. Coupling between visual alpha oscillations and default
mode activity. Neuroimage 68, 112–118. https://doi.org/10.1016/j.neuroimage.2012.11.058

Moosmann, M., Ritter, P., Krastel, I., Brink, A., Thees, S., Blankenburg, F., Taskin, B., Obrig, H.,
Villringer, A., 2003. Correlates of alpha rhythm in functional magnetic resonance imaging
and near infrared spectroscopy. Neuroimage 20, 145–158. https://doi.org/10.1016/S1053-
8119(03)00344-6

Murphy, K., Birn, R.M., Handwerker, D.A., Jones, T.B., Bandettini, P.A., 2009. The impact of
global signal regression on resting state correlations: Are anti-correlated networks
introduced? Neuroimage 44, 893–905. https://doi.org/10.1016/j.neuroimage.2008.09.036

Murphy, K., Birn, R.M., Bandettini, P.A., 2013. Resting-state fMRI confounds and cleanup.
Neuroimage 80, 349–359. https://doi.org/10.1016/j.neuroimage.2013.04.001

Murphy, K., Fox, M.D., 2017. Towards a consensus regarding global signal regression for resting
state functional connectivity MRI. Neuroimage 154, 169–173.
https://doi.org/10.1016/j.neuroimage.2016.11.052

Muschelli, J., Nebel, M.B., Caffo, B.S., Barber, A.D., Pekar, J.J., Mostofsky, S.H., 2014. Reduction of motion-related artifacts in resting state fMRI using aCompCor. Neuroimage 96, 22–35. https://doi.org/10.1016/j.neuroimage.2014.03.028

Oken, B.S., Salinsky, M.C., Elsas, S.M., 2006. Vigilance, alertness, or sustained attention: physiological basis and measurement. Clinical Neurophysiology 117, 1885–1901. https://doi.org/10.1016/j.clinph.2006.01.017

Olbrich, S., Sander, C., Matschinger, H., Mergl, R., Trenner, M., Schönknecht, P., Hegerl, U., 2011. Brain and body: Associations between EEG-vigilance and the autonomous nervous system activity during rest. J Psychophysiol 25, 190–200. https://doi.org/10.1027/0269-8803/a000061

Oldfield, R.C., 1971. The assessment and analysis of handedness: The Edinburgh inventory. Neuropsychologia 9, 97–113. https://doi.org/10.1016/0028-3932(71)90067-4

Orban, C., Kong, R., Li, J., Chee, M.W.L., Yeo, B.T.T., 2020. Time of day is associated with paradoxical reductions in global signal fluctuation and functional connectivity. PLoS Biol 18, e3000602. https://doi.org/10.1371/journal.pbio.3000602

Özbay, P.S., Chang, C., Picchioni, D., Mandelkow, H., Chappel-Farley, M.G., van Gelderen, P., de Zwart, J.A., Duyn, J., 2019. Sympathetic activity contributes to the fMRI signal. Commun Biol 2, 1–9. https://doi.org/10.1038/s42003-019-0659-0

Parkes, L., Fulcher, B., Yücel, M., Fornito, A., 2018. An evaluation of the efficacy, reliability, and sensitivity of motion correction strategies for resting-state functional MRI. Neuroimage 171, 415–436. https://doi.org/10.1016/j.neuroimage.2017.12.073

Power, J.D., Barnes, K.A., Snyder, A.Z., Schlaggar, B.L., Petersen, S.E., 2012. Spurious but systematic correlations in functional connectivity MRI networks arise from subject motion. Neuroimage 59, 2142–2154. https://doi.org/10.1016/j.neuroimage.2011.10.018

Power, J.D., Laumann, T.O., Plitt, M., Martin, A., Petersen, S.E., 2017a. On Global fMRI Signals and Simulations. Trends Cogn Sci 21, 911–913. https://doi.org/10.1016/j.tics.2017.09.002

Power, J.D., Plitt, M., Gotts, S.J., Kundu, P., Voon, V., Bandettini, P.A., Martin, A., 2018. Ridding fMRI data of motion-related influences: Removal of signals with distinct spatial and physical bases in multiecho data. Proc Natl Acad Sci U S A 115, E2105–E2114. https://doi.org/10.1073/pnas.1720985115

Power, J.D., Plitt, M., Laumann, T.O., Martin, A., 2017b. Sources and implications of whole-brain fMRI signals in humans. Neuroimage 146, 609–625. https://doi.org/10.1016/j.neuroimage.2016.09.038

Prokopiou, P.C., Pattinson, K.T.S., Wise, R.G., Mitsis, G.D., 2019. Modeling of dynamic cerebrovascular reactivity to spontaneous and externally induced CO_2 fluctuations in the human brain using BOLD-fMRI. Neuroimage 186, 533–548. https://doi.org/10.1016/j.neuroimage.2018.10.084

Prokopiou, P.C., Xifra-Porxas, A., Kassinopoulos, M., Boudrias, M.H., Mitsis, G.D., 2022. Modeling the Hemodynamic Response Function Using EEG-fMRI Data During Eyes-Open

Resting-State Conditions and Motor Task Execution. Brain Topogr 35, 302–321.
https://doi.org/10.1007/s10548-022-00898-w

Rosvall, M., Bergstrom, C.T., 2008. Maps of random walks on complex networks reveal community
structure. Proc. Natl. Acad. Sci. U. S. A. 105, 1118–1123.
https://doi.org/10.1073/pnas.0706851105

Scheeringa, R., Petersson, K.M., Kleinschmidt, A., Jensen, O., Bastiaansen, M.C.M., 2012. EEG
Alpha Power Modulation of fMRI Resting-State Connectivity. Brain Connect 2, 254–264.
https://doi.org/10.1089/brain.2012.0088

Scholvinck, M.L., Maier, A., Ye, F.Q., Duyn, J.H., Leopold, D.A., 2010. Neural basis of global
resting-state fMRI activity. Proceedings of the National Academy of Sciences 107, 10238–
10243. https://doi.org/10.1073/pnas.0913110107

Seitzman, B.A., Gratton, C., Marek, S., Raut, R. V., Dosenbach, N.U.F., Schlaggar, B.L., Petersen,
S.E., Greene, D.J., 2020. A set of functionally-defined brain regions with improved
representation of the subcortex and cerebellum. Neuroimage 206, 116290.
https://doi.org/10.1016/j.neuroimage.2019.116290

Shmueli, K., van Gelderen, P., de Zwart, J.A., Horovitz, S.G., Fukunaga, M., Jansma, J.M., Duyn,
J.H., 2007. Low-frequency fluctuations in the cardiac rate as a source of variance in the
resting-state fMRI BOLD signal. Neuroimage 38, 306–320.
https://doi.org/10.1016/j.neuroimage.2007.07.037

Smith, S.M., Fox, P.T., Miller, K.L., Glahn, D.C., Fox, P.M., Mackay, C.E., Filippini, N., Watkins,
K.E., Toro, R., Laird, A.R., Beckmann, C.F., 2009. Correspondence of the brain's functional
architecture during activation and rest. Proceedings of the National Academy of Sciences
106, 13040–13045. https://doi.org/10.1073/pnas.0905267106

Thompson, G.J., Riedl, V., Grimmer, T., Drzezga, A., Herman, P., Hyder, F., 2016. The Whole-
Brain "Global" Signal from Resting State fMRI as a Potential Biomarker of Quantitative State
Changes in Glucose Metabolism. Brain Connect 6, 435–447.
https://doi.org/10.1089/brain.2015.0394

Turchi, J., Chang, C., Ye, F.Q., Russ, B.E., Yu, D.K., Cortes, C.R., Monosov, I.E., Duyn, J.H.,
Leopold, D.A., 2018. The Basal Forebrain Regulates Global Resting-State fMRI
Fluctuations. Neuron 97, 940-952.e4. https://doi.org/10.1016/j.neuron.2018.01.032

Whittaker, J.R., Driver, I.D., Venzi, M., Bright, M.G., Murphy, K., 2019a. Cerebral autoregulation
evidenced by synchronized low frequency oscillations in blood pressure and resting-state
fMRI. Front Neurosci 13, 1–12. https://doi.org/10.3389/fnins.2019.00433

Whittaker, J.R., Driver, I.D., Venzi, M., Bright, M.G., Murphy, K., 2019b. Cerebral autoregulation
evidenced by synchronized low frequency oscillations in blood pressure and resting-state
fMRI. Front Neurosci 13, 1–13. https://doi.org/10.3389/fnins.2019.00433

Wise, R.G., Ide, K., Poulin, M.J., Tracey, I., 2004. Resting fluctuations in arterial carbon dioxide
induce significant low frequency variations in BOLD signal. Neuroimage 21, 1652–1664.
https://doi.org/10.1016/j.neuroimage.2003.11.025

Wong, C.K., Zotev, V., Misaki, M., Phillips, R., Luo, Q., Bodurka, J., 2016. Automatic EEG-
assisted retrospective motion correction for fMRI (aE-REMCOR). Neuroimage 129, 133–
147. https://doi.org/10.1016/j.neuroimage.2016.01.042

Wong, C.W., DeYoung, P.N., Liu, T.T., 2016. Differences in the resting-state fMRI global signal
amplitude between the eyes open and eyes closed states are related to changes in EEG
vigilance. Neuroimage 124, 24–31. https://doi.org/10.1016/j.neuroimage.2015.08.053

Wong, C.W., Olafsson, V., Tal, O., Liu, T.T., 2013. The amplitude of the resting-state fMRI global
signal is related to EEG vigilance measures. Neuroimage 83, 983–990.
https://doi.org/10.1016/j.neuroimage.2013.07.057

Wu, G.R., Marinazzo, D., 2016. Sensitivity of the resting-state haemodynamic response function
estimation to autonomic nervous system fluctuations. Philosophical Transactions of the Royal
Society A: Mathematical, Physical and Engineering Sciences 374.
https://doi.org/10.1098/rsta.2015.0190

Xia, C.H., Ma, Z., Ciric, R., Gu, S., Betzel, R.F., Kaczkurkin, A.N., Calkins, M.E., Cook, P.A.,
García de la Garza, A., Vandekar, S.N., Cui, Z., Moore, T.M., Roalf, D.R., Ruparel, K., Wolf,
D.H., Davatzikos, C., Gur, R.C., Gur, R.E., Shinohara, R.T., Bassett, D.S., Satterthwaite,
T.D., 2018. Linked dimensions of psychopathology and connectivity in functional brain
networks. Nat Commun 9, 1–14. https://doi.org/10.1038/s41467-018-05317-y

Xifra-Porxas, A., Kassinopoulos, M., Mitsis, G.D., 2021. Physiological and motion signatures in
static and time-varying functional connectivity and their subject identifiability. Elife 10,
e62324. https://doi.org/10.7554/eLife.62324

Xifra-Porxas, A., Kassinopoulos, M., Mitsis, G.D., 2020. Physiological and head motion
signatures in static and time-varying functional connectivity and their subject
discriminability. bioRxiv.

Yeo, B.T.T., Tandi, J., Chee, M.W.L., 2015. Functional connectivity during rested wakefulness
predicts vulnerability to sleep deprivation. Neuroimage 111, 147–158.
https://doi.org/10.1016/j.neuroimage.2015.02.018

Yuan, H., Zotev, V., Phillips, R., Bodurka, J., 2013. Correlated slow fluctuations in respiration,
EEG, and BOLD fMRI. Neuroimage 79, 81–93.
https://doi.org/10.1016/j.neuroimage.2013.04.068

Zarahn, E., Aguirre, G.K., D'Esposito, M., 1997. Empirical Analyses of BOLD fMRI Statistics.
Neuroimage 5, 199–212. https://doi.org/10.1006/nimg.1997.0264

Behzadi, Y., Restom, K., Liau, J., Liu, T.T., 2007. A component based noise correction method
(CompCor) for BOLD and perfusion based fMRI. Neuroimage 37, 90–101.
https://doi.org/10.1016/j.neuroimage.2007.04.042

Birn, R.M., Diamond, J.B., Smith, M.A., Bandettini, P.A., 2006. Separating respiratory-variation-
related fluctuations from neuronal-activity-related fluctuations in fMRI. Neuroimage 31,
1536–1548. https://doi.org/10.1016/j.neuroimage.2006.02.048

827 Biswal, B., Yetkin, F.Z., Haughton, V.M., Hyde, J.S., 1995. Functional connectivity in the motor
828   cortex of resting human brain using echo-planar MRI. Magn Reson Med.
829   https://doi.org/10.1002/mrm.1910340409

830 Bowman, A.D., Griffis, J.C., Visscher, K.M., Dobbins, A.C., Gawne, T.J., DiFrancesco, M.W.,
831   Szaflarski, J.P., 2017. Relationship Between Alpha Rhythm and the Default Mode Network.
832   Journal of Clinical Neurophysiology 34, 527–533.
833   https://doi.org/10.1097/WNP.0000000000000411

834 Burgess, G.C., Kandala, S., Nolan, D., Laumann, T.O., Power, J.D., Adeyemo, B., Harms, M.P.,
835   Petersen, S.E., Barch, D.M., 2016. Evaluation of Denoising Strategies to Address Motion-
836   Correlated Artifacts in Resting-State Functional Magnetic Resonance Imaging Data from the
837   Human Connectome Project. Brain Connect 6, 669–680.
838   https://doi.org/10.1089/brain.2016.0435

839 Caballero-Gaudes, C., Reynolds, R.C., 2017. Methods for cleaning the BOLD fMRI signal.
840   Neuroimage 154, 128–149. https://doi.org/10.1016/j.neuroimage.2016.12.018

841 Chang, C., Glover, G.H., 2009. Effects of model-based physiological noise correction on default
842   mode network anti-correlations and correlations. Neuroimage 47, 1448–1459.
843   https://doi.org/10.1016/j.neuroimage.2009.05.012

844 Chang, C., Leopold, D.A., Schölvinck, M.L., Mandelkow, H., Picchioni, D., Liu, X., Ye, F.Q.,
845   Turchi, J.N., Duyn, J.H., 2016. Tracking brain arousal fluctuations with fMRI. Proceedings
846   of the National Academy of Sciences 113, 4518–4523.
847   https://doi.org/10.1073/pnas.1520613113

848 Ciric, R., Rosen, A.F.G., Erus, G., Cieslak, M., Adebimpe, A., Cook, P.A., Bassett, D.S.,
849   Davatzikos, C., Wolf, D.H., Satterthwaite, T.D., 2018. Mitigating head motion artifact in
850   functional connectivity MRI. Nat Protoc 13, 2801–2826. https://doi.org/10.1038/s41596-
851   018-0065-y

852 Ciric, R., Wolf, D.H., Power, J.D., Roalf, D.R., Baum, G.L., Ruparel, K., Shinohara, R.T., Elliott,
853   M.A., Eickho, S.B., Davatzikos, C., Gur, R.C., Gur, R.E., Bassett, D.S., 2017. Benchmarking
854   of participant-level confound regression strategies for the control of motion artifact in studies
855   of functional connectivity. Neuroimage 154, 174–187.
856   https://doi.org/10.1016/j.neuroimage.2017.03.020

857 Dagenais, R., Mitsis, G.D., 2023. Non-invasive estimation of arterial blood pressure fluctuations
858   using a peripheral photoplethysmograph inside the MRI scanner. Annu Int Conf IEEE Eng
859   Med Biol Soc 2023, 1–4. https://doi.org/10.1109/EMBC40787.2023.10340020

860 Duyn, J.H., Ozbay, P.S., Chang, C., Picchioni, D., 2020. Physiological changes in sleep that affect
861   fMRI inference. Curr Opin Behav Sci 33, 42–50.
862   https://doi.org/10.1016/j.cobeha.2019.12.007

863 Erdoğan, S.B., Tong, Y., Hocke, L.M., Lindsey, K.P., deB Frederick, B., 2016. Correcting for
864   blood arrival time in global mean regression enhances functional connectivity analysis of
865   resting state fMRI-BOLD signals. Front Hum Neurosci 10, 1–22.
866   https://doi.org/10.3389/fnhum.2016.00311

Falahpour, M., Chang, C., Wong, C.W., Liu, T.T., 2018. Template-based prediction of vigilance
fluctuations   in   resting-state   fMRI.   Neuroimage   174,   317–327.
https://doi.org/10.1016/j.neuroimage.2018.03.012

Finn, E.S., Shen, X., Scheinost, D., Rosenberg, M.D., Huang, J., Chun, M.M., Papademetris, X.,
Constable, R.T., 2015. Functional connectome fingerprinting: Identifying individuals using
patterns of brain connectivity. Nat Neurosci 18, 1664–1671. https://doi.org/10.1038/nn.4135

Fox, M.D., Raichle, M.E., 2007. Spontaneous fluctuations in brain activity observed with
functional   magnetic   resonance   imaging.   Nat   Rev   Neurosci   8,   700–711.
https://doi.org/10.1038/nrn2201

Fox, M.D., Zhang, D., Snyder, A.Z., Raichle, M.E., 2009. The global signal and observed
anticorrelated   resting   state   brain   networks.   J   Neurophysiol   101,   3270–83.
https://doi.org/10.1152/jn.90777.2008

Glasser, M.F., Coalson, T.S., Bijsterbosch, J.D., Harrison, S.J., Harms, M.P., Anticevic, A., Van
Essen, D.C., Smith, S.M., 2018. Using temporal ICA to selectively remove global noise while
preserving global signal in functional MRI data. Neuroimage 181, 692–717.
https://doi.org/10.1016/j.neuroimage.2018.04.076

Gratton, C., Koller, J.M., Shannon, W., Greene, D.J., Maiti, B., Snyder, A.Z., Petersen, S.E.,
Perlmutter, J.S., Campbell, M.C., 2019. Emergent Functional Network Effects in Parkinson
Disease. Cerebral Cortex 29, 2509–2523. https://doi.org/10.1093/cercor/bhy121

Gratton, C., Laumann, T.O., Nielsen, A.N., Greene, D.J., Gordon, E.M., Gilmore, A.W., Nelson,
S.M., Coalson, R.S., Snyder, A.Z., Schlaggar, B.L., Dosenbach, N.U.F., Petersen, S.E., 2018.
Functional Brain Networks Are Dominated by Stable Group and Individual Factors, Not
Cognitive or Daily Variation. Neuron 439–452. https://doi.org/10.1016/j.neuron.2018.03.035

Gutierrez-Barragan, D., Basson, M.A., Panzeri, S., Gozzi, A., 2019. Infraslow State Fluctuations
Govern Spontaneous fMRI Network Dynamics. Current Biology 29, 2295-2306.e5.
https://doi.org/10.1016/j.cub.2019.06.017

Han, H.H., Gu, Y., Liu, X., 2019. A neurophysiological event of arousal modulation may underlie
fMRI-EEG correlations. Front. Neurosci. 13, 1–5. https://doi.org/10.3389/fnins.2019.00823

Iadecola, C., 2017. The Neurovascular Unit Coming of Age: A Journey through Neurovascular
Coupling   in   Health   and   Disease.   Neuron   96,   17–42.
https://doi.org/10.1016/j.neuron.2017.07.030

Jann, K., Dierks, T., Boesch, C., Kottlow, M., Strik, W., Koenig, T., 2009. BOLD correlates of
EEG alpha phase-locking and the fMRI default mode network. Neuroimage 45, 903–916.
https://doi.org/10.1016/j.neuroimage.2009.01.001

Jenkinson, M., Beckmann, C.F., Behrens, T.E.J., Woolrich, M.W., Smith, S.M., 2012. FSL.
Neuroimage 62, 782–790. https://doi.org/10.1016/j.neuroimage.2011.09.015

Ji, J.L., Kulkarni, K., Repovš, G., Spronk, M., Cole, M.W., Anticevic, A., 2018. Mapping the
human brain's cortical-subcortical functional network organization. Neuroimage 185, 35–57.
https://doi.org/10.1016/j.neuroimage.2018.10.006

Kassinopoulos, M., Harper, R.M., Guye, M., Lemieux, L., Diehl, B., 2021. Altered Relationship
Between Heart Rate Variability and fMRI-Based Functional Connectivity in People With
Epilepsy. Front Neurol 12, 1–13. https://doi.org/10.3389/fneur.2021.671890

Kassinopoulos, M., Mitsis, G.D., 2022. A multi-measure approach for assessing the performance
of fMRI preprocessing strategies in resting-state functional connectivity. Magn Reson
Imaging 85, 228–250. https://doi.org/10.1016/j.mri.2021.10.028

Kassinopoulos, M., Mitsis, G.D., 2021. Physiological noise modeling in fMRI based on the
pulsatile component of photoplethysmograph. Neuroimage 242, 118467.
https://doi.org/10.1016/j.neuroimage.2021.118467

Kassinopoulos, M., Mitsis, G.D., 2019. Identification of physiological response functions to
correct for fluctuations in resting-state fMRI related to heart rate and respiration. Neuroimage
202. https://doi.org/10.1101/512855

Keller, C.J., Bickel, S., Honey, C.J., Groppe, D.M., Entz, L., Craddock, R.C., Lado, F.A., Kelly,
C., Milham, M., Mehta, A.D., 2013. Neurophysiological investigation of spontaneous
correlated and anticorrelated fluctuations of the BOLD signal. Journal of Neuroscience 33,
6333–6342. https://doi.org/10.1523/JNEUROSCI.4837-12.2013

Lancaster, G., Iatsenko, D., Pidde, A., Ticcinelli, V., Stefanovska, A., 2018. Surrogate data for
hypothesis testing of physical systems. Phys Rep 748, 1–60.
https://doi.org/10.1016/j.physrep.2018.06.001

Li, J., Bolt, T., Bzdok, D., Nomi, J., Yeo, B.T.T., Spreng, R.N., Uddin, L.Q., 2019a. Topography
and behavioral relevance of the global signal in the human brain. Sci Rep 10, 1–5.
https://doi.org/10.1038/s41598-019-50750-8

Li, J., Kong, R., Liégeois, R., Orban, C., Tan, Y., Sun, N., Holmes, A.J., Sabuncu, M.R., Ge, T.,
Yeo, B.T.T., 2019b. Global signal regression strengthens association between resting-state
functional connectivity and behavior. Neuroimage 196, 126–141.
https://doi.org/10.1016/j.neuroimage.2019.04.016

Liu, T.T., 2016. Noise contributions to the fMRI signal: An overview. Neuroimage 143, 141–151.
https://doi.org/10.1016/j.neuroimage.2016.09.008

Liu, T.T., Nalci, A., Falahpour, M., 2017. The global signal in fMRI: Nuisance or Information?
Neuroimage 150, 213–229. https://doi.org/10.1016/j.neuroimage.2017.02.036

Liu, X., De Zwart, J.A., Schölvinck, M.L., Chang, C., Ye, F.Q., Leopold, D.A., Duyn, J.H., 2018.
Subcortical evidence for a contribution of arousal to fMRI studies of brain activity. Nat
Commun 9, 1–10. https://doi.org/10.1038/s41467-017-02815-3

Logothetis, N.K., 2008. What we can do and what we cannot do with fMRI. Nature 453, 869–878.

Logothetis, N.K., Pauls, J., Augath, M., Trinath, T., Oeltermann, A., 2001. Neurophysiological
investigation of the basis of the fMRI signal. Nature 412, 150–157.
https://doi.org/10.1038/35084005

Mantini, D., Perrucci, M.G., Gratta, D., C., R., G.l, Corbetta, M., Del Gratta, C., Romani, G., 2007.
Electrophysiological signatures of resting state networks in the human brain. Proceedings of
the National Academy of Sciences 104, 13170.

Marek, S., Siegel, J.S., Gordon, E.M., Raut, R. V., Gratton, C., Newbold, D.J., Ortega, M.,
Laumann, T.O., Miller, D.B., Zheng, A., Lopez, K.C., Berg, J.J., Coalson, R.S., Nguyen,
A.L., Dierker, D., Van, A.N., Hoyt, C.R., McDermott, K.B., Norris, S.A., Shimony, J.S.,
Snyder, A.Z., Nelson, S.M., Barch, D.M., Schlaggar, B.L., Raichle, M.E., Petersen, S.E.,
Greene, D.J., Dosenbach, N.U.F., 2018. Spatial and Temporal Organization of the Individual
Human Cerebellum. SSRN Electronic Journal 100, 977-993.e7.
https://doi.org/10.2139/ssrn.3188429

Mayhew, S.D., Bagshaw, A.P., 2017. Dynamic spatiotemporal variability of alpha-BOLD
relationships during the resting-state and task-evoked responses. Neuroimage 155, 120–137.
https://doi.org/10.1016/j.neuroimage.2017.04.051

Mayhew, S.D., Ostwald, D., Porcaro, C., Bagshaw, A.P., 2013. Spontaneous EEG alpha oscillation
interacts with positive and negative BOLD responses in the visual-auditory cortices and
default-mode network. Neuroimage 76, 362–372.
https://doi.org/10.1016/j.neuroimage.2013.02.070

Mo, J., Liu, Y., Huang, H., Ding, M., 2013. Coupling between visual alpha oscillations and default
mode activity. Neuroimage 68, 112–118. https://doi.org/10.1016/j.neuroimage.2012.11.058

Moosmann, M., Ritter, P., Krastel, I., Brink, A., Thees, S., Blankenburg, F., Taskin, B., Obrig, H.,
Villringer, A., 2003. Correlates of alpha rhythm in functional magnetic resonance imaging
and near infrared spectroscopy. Neuroimage 20, 145–158. https://doi.org/10.1016/S1053-
8119(03)00344-6

Murphy, K., Birn, R.M., Bandettini, P.A., 2013. Resting-state fMRI confounds and cleanup.
Neuroimage 80, 349–359. https://doi.org/10.1016/j.neuroimage.2013.04.001

Murphy, K., Fox, M.D., 2017. Towards a consensus regarding global signal regression for resting
state functional connectivity MRI. Neuroimage 154, 169–173.
https://doi.org/10.1016/j.neuroimage.2016.11.052

Muschelli, J., Nebel, M.B., Caffo, B.S., Barber, A.D., Pekar, J.J., Mostofsky, S.H., 2014.
Reduction of motion-related artifacts in resting state fMRI using aCompCor. Neuroimage 96,
22–35. https://doi.org/10.1016/j.neuroimage.2014.03.028

Oken, B.S., Salinsky, M.C., Elsas, S.M., 2006. Vigilance, alertness, or sustained attention:
physiological basis and measurement. Clinical Neurophysiology 117, 1885–1901.
https://doi.org/10.1016/j.clinph.2006.01.017

Olbrich, S., Sander, C., Matschinger, H., Mergl, R., Trenner, M., Schönknecht, P., Hegerl, U.,
2011. Brain and body: Associations between EEG-vigilance and the autonomous nervous
system activity during rest. J Psychophysiol 25, 190–200. https://doi.org/10.1027/0269-
8803/a000061

Oldfield, R.C., 1971. The assessment and analysis of handedness: The Edinburgh inventory.
Neuropsychologia 9, 97–113. https://doi.org/10.1016/0028-3932(71)90067-4

Orban, C., Kong, R., Li, J., Chee, M.W.L., Yeo, B.T.T., 2020. Time of day is associated with
paradoxical reductions in global signal fluctuation and functional connectivity. PLoS Biol 18,
e3000602. https://doi.org/10.1371/journal.pbio.3000602

Özbay, P.S., Chang, C., Picchioni, D., Mandelkow, H., Chappel-Farley, M.G., van Gelderen, P.,
de Zwart, J.A., Duyn, J., 2019. Sympathetic activity contributes to the fMRI signal. Commun
Biol 2, 1–9. https://doi.org/10.1038/s42003-019-0659-0

Parkes, L., Fulcher, B., Yücel, M., Fornito, A., 2018. An evaluation of the efficacy, reliability, and
sensitivity of motion correction strategies for resting-state functional MRI. Neuroimage 171,
415–436. https://doi.org/10.1016/j.neuroimage.2017.12.073

Power, J.D., Barnes, K.A., Snyder, A.Z., Schlaggar, B.L., Petersen, S.E., 2012. Spurious but
systematic correlations in functional connectivity MRI networks arise from subject motion.
Neuroimage 59, 2142–2154. https://doi.org/10.1016/j.neuroimage.2011.10.018

Power, J.D., Laumann, T.O., Plitt, M., Martin, A., Petersen, S.E., 2017a. On Global fMRI Signals
and Simulations. Trends Cogn Sci 21, 911–913. https://doi.org/10.1016/j.tics.2017.09.002

Power, J.D., Plitt, M., Gotts, S.J., Kundu, P., Voon, V., Bandettini, P.A., Martin, A., 2018. Ridding
fMRI data of motion-related influences: Removal of signals with distinct spatial and physical
bases in multiecho data. Proc Natl Acad Sci U S A 115, E2105–E2114.
https://doi.org/10.1073/pnas.1720985115

Power, J.D., Plitt, M., Laumann, T.O., Martin, A., 2017b. Sources and implications of whole-brain
fMRI signals in humans. Neuroimage 146, 609–625.
https://doi.org/10.1016/j.neuroimage.2016.09.038

Prokopiou, P.C., Pattinson, K.T.S., Wise, R.G., Mitsis, G.D., 2019. Modeling of dynamic
cerebrovascular reactivity to spontaneous and externally induced CO_2 fluctuations in the
human brain using BOLD-fMRI. Neuroimage 186, 533–548.
https://doi.org/10.1016/j.neuroimage.2018.10.084

Prokopiou, P.C., Xifra-Porxas, A., Kassinopoulos, M., Boudrias, M.H., Mitsis, G.D., 2022.
Modeling the Hemodynamic Response Function Using EEG-fMRI Data During Eyes-Open
Resting-State Conditions and Motor Task Execution. Brain Topogr 35, 302–321.
https://doi.org/10.1007/s10548-022-00898-w

Scheeringa, R., Petersson, K.M., Kleinschmidt, A., Jensen, O., Bastiaansen, M.C.M., 2012. EEG
Alpha Power Modulation of fMRI Resting-State Connectivity. Brain Connect 2, 254–264.
https://doi.org/10.1089/brain.2012.0088

Scholvinck, M.L., Maier, A., Ye, F.Q., Duyn, J.H., Leopold, D.A., 2010. Neural basis of global
resting-state fMRI activity. Proceedings of the National Academy of Sciences 107, 10238–
10243. https://doi.org/10.1073/pnas.0913110107

Seitzman, B.A., Gratton, C., Marek, S., Raut, R. V., Dosenbach, N.U.F., Schlaggar, B.L., Petersen,
S.E., Greene, D.J., 2020. A set of functionally-defined brain regions with improved
representation of the subcortex and cerebellum. Neuroimage 206, 116290.
https://doi.org/10.1016/j.neuroimage.2019.116290

Shmueli, K., van Gelderen, P., de Zwart, J.A., Horovitz, S.G., Fukunaga, M., Jansma, J.M., Duyn, J.H., 2007. Low-frequency fluctuations in the cardiac rate as a source of variance in the resting-state fMRI BOLD signal. Neuroimage 38, 306–320. https://doi.org/10.1016/j.neuroimage.2007.07.037

Smith, S.M., Fox, P.T., Miller, K.L., Glahn, D.C., Fox, P.M., Mackay, C.E., Filippini, N., Watkins, K.E., Toro, R., Laird, A.R., Beckmann, C.F., 2009. Correspondence of the brain's functional architecture during activation and rest. Proceedings of the National Academy of Sciences 106, 13040–13045. https://doi.org/10.1073/pnas.0905267106

Thompson, G.J., Riedl, V., Grimmer, T., Drzezga, A., Herman, P., Hyder, F., 2016. The Whole-Brain "Global" Signal from Resting State fMRI as a Potential Biomarker of Quantitative State Changes in Glucose Metabolism. Brain Connect 6, 435–447. https://doi.org/10.1089/brain.2015.0394

Turchi, J., Chang, C., Ye, F.Q., Russ, B.E., Yu, D.K., Cortes, C.R., Monosov, I.E., Duyn, J.H., Leopold, D.A., 2018. The Basal Forebrain Regulates Global Resting-State fMRI Fluctuations. Neuron 97, 940-952.e4. https://doi.org/10.1016/j.neuron.2018.01.032

Whittaker, J.R., Driver, I.D., Venzi, M., Bright, M.G., Murphy, K., 2019a. Cerebral autoregulation evidenced by synchronized low frequency oscillations in blood pressure and resting-state fMRI. Front Neurosci 13, 1–12. https://doi.org/10.3389/fnins.2019.00433

Whittaker, J.R., Driver, I.D., Venzi, M., Bright, M.G., Murphy, K., 2019b. Cerebral autoregulation evidenced by synchronized low frequency oscillations in blood pressure and resting-state fMRI. Front Neurosci 13, 1–13. https://doi.org/10.3389/fnins.2019.00433

Wise, R.G., Ide, K., Poulin, M.J., Tracey, I., 2004. Resting fluctuations in arterial carbon dioxide induce significant low frequency variations in BOLD signal. Neuroimage 21, 1652–1664. https://doi.org/10.1016/j.neuroimage.2003.11.025

Wong, C.K., Zotev, V., Misaki, M., Phillips, R., Luo, Q., Bodurka, J., 2016. Automatic EEG-assisted retrospective motion correction for fMRI (aE-REMCOR). Neuroimage 129, 133–147. https://doi.org/10.1016/j.neuroimage.2016.01.042

Wong, C.W., DeYoung, P.N., Liu, T.T., 2016. Differences in the resting-state fMRI global signal amplitude between the eyes open and eyes closed states are related to changes in EEG vigilance. Neuroimage 124, 24–31. https://doi.org/10.1016/j.neuroimage.2015.08.053

Wong, C.W., Olafsson, V., Tal, O., Liu, T.T., 2013. The amplitude of the resting-state fMRI global signal is related to EEG vigilance measures. Neuroimage 83, 983–990. https://doi.org/10.1016/j.neuroimage.2013.07.057

Wu, G.R., Marinazzo, D., 2016. Sensitivity of the resting-state haemodynamic response function estimation to autonomic nervous system fluctuations. Philosophical Transactions of the Royal Society A: Mathematical, Physical and Engineering Sciences 374. https://doi.org/10.1098/rsta.2015.0190

Xia, C.H., Ma, Z., Ciric, R., Gu, S., Betzel, R.F., Kaczkurkin, A.N., Calkins, M.E., Cook, P.A., García de la Garza, A., Vandekar, S.N., Cui, Z., Moore, T.M., Roalf, D.R., Ruparel, K., Wolf, D.H., Davatzikos, C., Gur, R.C., Gur, R.E., Shinohara, R.T., Bassett, D.S., Satterthwaite,

T.D., 2018. Linked dimensions of psychopathology and connectivity in functional brain
networks. Nat Commun 9, 1–14. https://doi.org/10.1038/s41467-018-05317-y

Xifra-Porxas, A., Kassinopoulos, M., Mitsis, G.D., 2021. Physiological and motion signatures in
static and time-varying functional connectivity and their subject identifiability. Elife 10,
e62324. https://doi.org/10.7554/eLife.62324

Xifra-Porxas, A., Kassinopoulos, M., Mitsis, G.D., 2020. Physiological and head motion
signatures in static and time-varying functional connectivity and their subject
discriminability. bioRxiv.

Yeo, B.T.T., Tandi, J., Chee, M.W.L., 2015. Functional connectivity during rested wakefulness
predicts vulnerability to sleep deprivation. Neuroimage 111, 147–158.
https://doi.org/10.1016/j.neuroimage.2015.02.018

Yuan, H., Zotev, V., Phillips, R., Bodurka, J., 2013. Correlated slow fluctuations in respiration,
EEG, and BOLD fMRI. Neuroimage 79, 81–93.
https://doi.org/10.1016/j.neuroimage.2013.04.068

Zarahn, E., Aguirre, G.K., D'Esposito, M., 1997. Empirical Analyses of BOLD fMRI Statistics.
Neuroimage 5, 199–212. https://doi.org/10.1006/nimg.1997.0264

1078

1079

FIGURE CAPTIONS

Figure 1. (a) Experimental paradigm. Participants underwent two resting-state scans with eyes open, alternated by a motor task. The maximum voluntary contraction (MVC) of each participant was measured before performing the motor task. **(b) Hand-grip task.** Participants performed a unimanual right hand-grip task. During each trial, they initially fixated on a crosshair for a few seconds. This was followed by the appearance of an orange circle on the screen, whereby participants had 2 s to apply force to reach 15% of their MVC. A steady grip was then maintained for 3 s, which was followed by a guided ramp period where participants had to apply force to reach 30% of their MVC and sustain this grip strength for another 3 s.

Figure 2. Optimal number of PCA white matter components used for fMRI denoising. (a) Calculation of the functional connectivity contrast (FCC) measure, which quantifies the difference in correlation values among edges within and between-networks and serves in assessing the identifiability of well-established large-scale networks. **(b-d)** Group-averaged FCC measure as a function of the number of white matter PCA components used for denoising for **(b)** resting-state 1, **(c)** motor task, and **(d)** resting-state 2. The black line denotes mean across subjects and the shaded area denotes standard error. Based on these traces, 10 PCA components were used for denoising, as this number yielded a high FCC value for all three conditions.

Figure 3. Partial correlation of SLFOs and EEG power time-series with the fMRI global signal across subjects for (a) resting-state 1, (b) motor task, and (c) resting-state 2. Significance testing was performed against surrogate data (Wilcoxon rank-sum test, $*$ $p < 0.05$, $**$ $p < 0.001$). The p-values for the four EEG bands were corrected for multiple comparisons using FDR. SLFOs were highly correlated with the global signal, whereas alpha and beta power fluctuations were found to be weakly correlated with global signal variations.

Figure 4. Effect of GSR on connectome patterns induced by SLFOs. (a) Group averaged FC matrices computed using the synthetic dataset associated with systemic low frequency fluctuations (SLFOs) for each scanning session. **(b)** Similarity between the SLFOs' FC matrices and the FC matrices extracted from the minimally preprocessed (MinPrep) data, the preprocessed data (after

regressing out 10 PCA white matter components; WM_{10}), and the preprocessed data after global

signal regression (WM_{10} & GSR). While WM regression reduced the similarity between the

minimally preprocessed (MinPrep) and SLFOs connectivity matrices, suggesting a reduction in

connectivity bias, GSR further reduced this similarity significantly (Wilcoxon rank-sum test, * p

< 0.05, ** $p < 0.005$). Error bars denote standard deviation.

Figure 5. Effect of GSR on connectome patterns associated with EEG power. (a) Group

averaged FC matrices computed using the fMRI synthetic dataset associated with fluctuations in

alpha power, for each scanning session. **(b)** Similarity between the alpha band EEG-based FC

matrices and the FC matrices extracted from the minimally preprocessed data, the preprocessed

data (i.e. fMRI data after regressing out 10 PCA white matter components; WM_{10}), and the

preprocessed data after global signal regression (WM_{10} & GSR). **(c)** Group averaged FC matrices

computed using the fMRI synthetic dataset associated with fluctuations in beta power, for the

motor task. **(d)** Similarity between the beta band EEG-based FC matrices and the FC matrices

extracted from the minimally preprocessed data (MinPrep), the preprocessed data (WM_{10}), and the

preprocessed data after GSR (WM_{10} & GSR). GSR did not significantly alter the connectivity

related to EEG activity (Wilcoxon rank-sum test). In all subplots, error bars denote standard

deviation.

1125

(a)

Resting-state 1
15 min

MVC
30 s

Motor Task
~13 min (50 trials)

Resting-state 2
15 min

(b)

Motor Task

Intertrial
Jittering from 3 to 5 s

Adapt
2 s

15% MVC
3 s

Ramp
3 s

30% MVC
3 s

(a) **Functional Connectivity Contrast (FCC)**

(b) **Resting-state 1**

(c) **Motor task**

(d) **Resting-state 2**

Alpha power

(a)

Resting-state 1   z(r)

Motor task   z(r)

Resting-state 2   z(r)

(b)

Resting-state 1

Motor task

Resting-state 2

Beta power (Motor task)

(c)

z(r)

(d)