Novel multiparametric bulk and single EV pipeline for adipose cell-specific biomarker discovery in paired human biospecimens

Paola Loreto Palacio[1], Jacelyn Greenwald[2], Kim Truc Nguyen[3], Dharti Shantaram[4], Bradley L. Butsch[1], Yonseok Kim[2], Mangesh Hade Dattu[1], Sabrena Noria[5], Stacy A. Brethauer[5], Bradley J. Needleman[5], Vicky Wysocki[2], Willa Hsueh[4], Eduardo Reátegui[3], Setty M. Magaña[1]*

[1]Department of Pediatrics, Center for Clinical and Translational Research, Nationwide Children's Hospital, Columbus, Ohio.

[2]Department of Chemistry and Biochemistry, The Ohio State University, Columbus, Ohio.

[3]Department of Chemical and Biomedical Engineering, The Ohio State University, Columbus, Ohio.

[4]Diabetes and Metabolism Research Center, The Ohio State University, Columbus, Ohio.

[5]Center for Minimally Invasive Surgery, Department of Surgery, The Ohio State University College of Medicine, Columbus, Ohio.

Running Title: Human adipose extracellular vesicles

*Corresponding Author: Setty Magaña, MD, PhD

Address: Nationwide Children's Hospital

                 Department of Pediatrics, Division of Neurology

                 700 Children's Drive

                 Columbus, OH 43205

Telephone: 614-722-4625

Fax: 614-722-4633

Email: Setty.Magana@nationwidechildrens.org

Abstract

Obesity is a global health crisis that contributes to morbidity and mortality worldwide. Obesity's comorbid association with a variety of diseases, from metabolic syndrome to neurodegenerative disease, underscores the critical need to better understand the pathobiology of obesity. Adipose tissue, once seen as an inert storage depot, is now recognized as an active endocrine organ, regulating metabolic and systemic homeostasis. Recent studies spotlight the theranostic utility of extracellular vesicles (EVs) as novel biomarkers and drivers of disease, including obesity-related complications. Adipose-derived EVs (ADEVs) have garnered increased interest for their roles in diverse diseases, however robust isolation and characterization protocols for human, cell-specific EV subsets are limited. Herein, we directly address this technical challenge by establishing a multiparametric analysis framework that leverages bulk and single EV characterization, mRNA phenotyping and proteomics of human ADEVs directly from paired visceral adipose tissue, cultured mature adipocyte conditioned media, and plasma from obese subjects undergoing bariatric surgery. Importantly, rigorous EV phenotyping at the tissue and cell-specific level identified top 'adipose liquid biopsy' candidates that were validated in circulating plasma EVs from the same patient. In summary, our study paves the way toward a tissue and cell-specific, multiparametric framework for studying tissue and circulating adipose EVs in obesity-driven disease.

Introduction

Obesity has emerged as a global health crisis, reaching epidemic proportions and posing significant challenges to public health systems worldwide[1]. Obesity is associated with numerous metabolic disorders, as well as cancer, autoimmunity, and neurodegenerative diseases[2-6]. Adipose tissue, once considered an inert energy storage depot, is now recognized as an active endocrine organ that plays a crucial role in metabolic homeostasis and overall health[7]. Adipose tissue dysfunction, characterized by altered adipokine secretion, chronic inflammation, and impaired adipocyte differentiation, is a hallmark of obesity and a pivotal contributor to the development of obesity-related comorbidities[8,9].

Recent studies have revealed that obesity is also associated with enhanced production of adipose-derived extracellular vesicles (ADEVs)[10,11]. Extracellular vesicles (EVs) are small, membrane-bound vesicles released by cells into the extracellular space and represent a novel method of intercellular communication[12]. ADEVs and other EVs have emerged as potential players in the pathogenesis of obesity and its associated metabolic complications[13-15]. EVs harbor and transmit bioactive cargo (i.e., proteins, lipids, nucleic acids) to recipient cells, causing phenotypic and genotypic changes[16,17]. This ability has positioned EVs as central orchestrators of various processes, including normal cellular biology, immune regulation, tissue repair, and disease progression[12,18,19].

Two subsets of ADEVs include adipose tissue-derived EVs (ATEVs) and mature adipocyte-derived EVs (aEVs)—each with unique pathobiological roles and cell-specific phenotypic repertoires. ATEVs, which consist of EVs originating from a heterogenous composite of cells such as pre-adipocytes, immune cells, and stromal support cells, reflect the complex microenvironment of adipose tissue[10,20] On the other hand, aEVs are released specifically from mature adipocytes and provide a glimpse into the molecular messages that adipocytes utilize to communicate with other cells and tissues. Furthermore, ADEVs serve as a useful biological 'reference source' in the development of an adiposity-associated EV liquid biopsy.

Despite the growing interest in EVs and ADEVs, their full translational potential is hindered by several methodological and biological challenges, including a lack in standardized, reproducible, and robust isolation and cell-specific characterization protocols[21,22]. While a handful of studies have investigated human ADEVs[11,23,24], most ADEV studies have focused on rodent models[25-27], despite recognized cross-species differences in adipose markers between

rodents and humans[28]. This cross-species variation highlights the critical need for human-based

obesity EV paradigms.

To address the current methodological and biological challenges hindering the

translational advancement of human ADEVs in the clinic, we established a human-based,

multiparametric workflow for the isolation and characterization of ADEVs from paired

biospecimens derived from bariatric surgical patients. Our objectives were to: i) establish

standardized protocols for EV isolation from paired human specimens (i.e. visceral adipose

tissue, cultured mature adipocyte-conditioned media, and plasma); ii) employ cutting-edge

technologies to optimize single and bulk adipose EV characterization; iii) characterize the

adipose EV proteome in order to inform the development of an adipose EV liquid biopsy; iv)

validate the tissue and cell-specific adiposity EV signature in circulating plasma EVs from the

same patient, and iv) perform in silico prediction of ADEV functional roles in order to gain

insights into the biological relevance of ADEVs.

Results

Multiparametric isolation and characterization of obesity-associated EVs

Figure 1 illustrates our multiparametric pipeline for EV isolation and characterization from

bariatric patient adipose tissue and paired plasma. Most human EV biomarker studies have

focused on biofluids or in vitro EV sources, however technical challenges must be overcome for

tissue-derived EVs in order to maintain EV integrity[29-31]. To ensure our isolation method

produced intact, tissue-derived adipose EVs, we performed rigorous bulk and single EV

characterization, as described below.

Bulk EV characterization of paired human adipose tissue EV (ATEV), mature adipocyte

EV (aEV) and plasma EV (bEV) characteristics

Microfluidic resistive pulse sensing (MRPS) was used to examine the size distribution and

particle concentration of EVs from three adipose sources (ATEV, aEV, bEV) derived from the

same individual (**Fig. 2,** n = 3). MRPS is an electric sensing platform that utilizes cartridges with

pre-specified upper and lower limits of detection. We were interested in studying small EVs and

therefore utilized the C400 cartridge (size range 65 nm to 400 nm). MRPS revealed greater

particle size distribution for ATEVs compared to pure aEVs and bEVs. Median particle diameter

was 86.12 nm for ATEVs, 82.26 nm for aEVs, and 78.60 nm for bEVs. Mean particle

concentration was 1.27e+13 particles/mL for ATEVs, 4.09e+12 particles/mL for aEVs, and
4.13e+12 particles/mL for bEVs (**Fig. 2A-C**).

Figure 1. Workflow for EV isolation (A) and characterization (B) of ATEVs, aEVs, and
bEVs. (A) Top: Visceral adipose tissue (2-10 grams) was enzymatically digested, filtered, and
centrifuged, resulting in three phases: pelleted stromal vascular fraction (SVF), SVF supernatant
(SVF-SN) and floating mature adipocytes. Mature adipocytes were cultured for 24 hours (see
Methods), and the cell-conditioned media (CMM) was removed for EV isolation (designated
aEVs) via tangential flow filtration (TFF). SVF-SN was also collected for ATEV isolation via TFF.
Lower: Paired plasma was collected in EDTA tubes and subjected to centrifugation for
separation of plasma and cellular components. To avoid contamination from platelet EVs,
platelet depletion was performed before EV isolation using size exclusion chromatography.
Fractions 1-4 were pooled and concentrated. **(B)** Multiparametric characterization of EVs by
immunoblotting, microfluidic resistive pulse sensing (MRPS), transmission electron microscopy
(TEM), and single EV multiplex analysis.

TEM imaging showed heterogenous particle populations, including small EVs with classic 'cup-
shaped' fixation artefacts (**Fig. 2D-F**). Immunoblot analysis revealed heterogenous protein
expression of typical EV/non-EV and adipose markers across biological replicates of ATEVs,
aEVs and bEVs (**Fig. 2G-I**). Notably, while calnexin is typically considered a non-EV marker,
calnexin was detected in ATEVs and aEVs (**Fig. 2G, H**), consistent with previous reports of
endoplasmic reticulum proteins associated with tissue-derived EVs[32-34]. The presence of
adiponectin in early SEC fractions suggests its association with EVs rather than free proteins,
as previously reported[11].

Figure 2. Human adipose EVs exhibited nanoscale size, cup-shaped morphology, classic
EV and adipose markers. (A-C) Particle size distribution and concentration analysis via MRPS
(D-F) TEM images displaying cup-shaped morphology of isolated EVs. **(G-I)** Immunoblotting
analysis for EV (CD9 and CD63), non-EV (CALX) and adipocyte markers (ADIPOQ). (n = 3
biological replicates). ATEVs: Adipose Tissue-derived Extracellular Vesicles, aEVs: Mature
Adipocyte-derived Extracellular Vesicles. CD9: Cluster of Differentiation 9, CD63: Cluster of
Differentiation 63, CALX: Calnexin, ADIPOQ: Adiponectin.

Human ATEVs and aEVs exhibit unique and shared proteomes and biological relevance to obesity-associated metabolic and immune processes

Next, we aimed to develop an 'adipose EV signature,' by first performing bulk proteomics (**Fig. 3**) to explore the unique and shared proteomes of adipose tissue EVs (reflecting the complex adipose tissue microenvironment) and mature adipocyte EVs (reflecting the 'purest' adipose reference material). Three-way Venn diagram revealed 548 unique ATEV proteins, 116 unique aEV proteins, and 1233 shared proteins (**Fig. 3A**). The overlapping proteins suggest potential functional similarities and shared signaling pathways, while the unique proteins signify specific molecular characteristics associated with each group. We then generated a heat map to visualize the top 50 most abundant proteins in ATEVs and aEVs (**Fig. 3B**). This data set informed our design of single EV immunocapture probes against adipose-enriched candidates (see **Fig. 5**). **Supplementary Table 1** lists the most abundant unique and shared proteins.

To investigate canonical biological processes and molecular functions associated with the identified proteins in both ATEVs and aEVs, we performed Gene Ontology (GO) pathway analysis (**Supplementary Fig. 1A,B**). Biological processes identified in our analysis highlighted several significant pathways, including genomic stability and DNA processing and chromatin remodeling. Our study revealed the involvement of ATEVs and aEVs in processes such as pre-replicative complex assembly, regulation of cellular amino acid metabolic processes, cell adhesion, and regulation of transcription from RNA polymerase II promoter in response to hypoxia. Pathways responsible for antigen processing and presentation, cell adhesion, and protein-protein interactions were also identified.

KEGG pathway analysis provided further insights into the potential pathobiological relevance of ATEVs and aEVs (**Supplementary Fig. 1C**). Prominent pathways were related to protein processing, quality control and degradation (e.g. proteasome, protein degradation, and endoplasmic reticulum). Additional pathways included neutrophil extracellular trap formation, focal adhesion, metabolic pathways, regulation of actin cytoskeleton, complement and coagulation cascades, leukocyte transendothelial migration, phagosome formation, glycolysis/gluconeogenesis, ECM-receptor interaction, and N-glycan biosynthesis. These pathways imply potential roles of ATEVs and aEVs in processes related to immune responses, cell adhesion, cellular metabolism, cytoskeletal regulation, and intercellular communication. Moreover, pathways associated with cholesterol metabolism, carbon metabolism, fatty acid degradation, platelet activation, endocytosis, antigen processing and presentation, PPAR signaling, and fatty acid metabolism were also identified.

Figure 3. Proteomics analysis of paired human adipose EVs. EVs isolated from paired visceral adipose tissue (ATEVs) and cultured mature adipocytes (aEVs) underwent proteomics analysis. **(A)** Three-way Venn diagram revealed proteins identified in both ATEVs and aEVs (n=1233), with 548 proteins exclusively found in ATEVs and 116 proteins exclusively found in aEVs. **(B)** Heat maps of the top 50 enriched proteins in ATEVs (orange) and aEVs (purple). (n = 3 biological replicates). ATEVs: adipose tissue-derived extracellular vesicles, aEVs: mature adipocyte extracellular vesicles

Multiparametric single EV analysis of human adipose-derived EVs

Orthogonal and complementary single EV methods were utilized to characterize EV size, tetraspanin distribution, EV and adipose protein targets and select adipose mRNA transcripts in paired ATEVs and aEVs from the same individual.

Single ADEV analysis leveraging single-particle interferometric reflectance imaging sensor (SP-IRIS)

SP-IRIS[35] with fluorescence was employed to gain insights into the complexity and

heterogeneity of tetraspanin distribution across biologically similar but distinct adipose-enriched

EV subsets at single EV resolution (**Fig. 4**). Single EVs (sEVs) were immunologically

immobilized/captured by surface tetraspanin proteins CD63, CD81, and CD9 printed on a

microchip surface. Fluorescently-labelled anti-CD63, anti-CD81 and anti-CD9 antibodies were

added for detection. Capture events were counted (mean particle count) and sEVs were sized

by light interference (lower limit of detection of 50 nm). SP-IRIS revealed a particle size scatter

of 50 nm to 200 nm for both ATEVs and aEVs (data not shown), which was consistent with

MRPS size analysis. Quantification of events per capture chip revealed a notable presence of

particles, indicating successful ATEV (**Fig. 4A, C**) and aEV (**Fig. 4B, D**) capture. Both ATEVs

and aEVs demonstrated the most total capture events with CD63 capture and greater CD63

detection from the CD63 capture spot. CD81 capture resulted in similar expression patterns for

ATEVs and aEVs while CD9 capture revealed higher CD9 detection for aEVs but not ATEVs

(**Fig. 4A, B**).

Tetraspanin colocalization analysis revealed distinct patterns among the CD63, CD9,

and CD81 capture spots **(Fig. 4C,D; Supplementary Fig. 2)**. CD63 was the most prevalent

single-positive tetraspanin detected on both ATEVs and aEVs immobilized on the CD63 capture

spot, as visualized by red fluorescence signal (**Fig. 4C,D**). ATEVs and aEVs captured by CD81

both had the highest detection for CD9+ single positive EVs, as well as CD63+/CD9+ double

positive EVs. CD9 capture differed between ATEVs and aEVs, with greater CD81+ single

positive ATEVs and slightly greater CD9+ single positive aEVs. These results indicate that

CD63 is the predominant surface marker expressed in both ATEVs and aEVs. Representative

images obtained with the SP-IRIS instrument visually confirmed these expression patterns (**Fig.

4C,D**). In summary, sEV tetraspanin phenotyping provides valuable insights into the tetraspanin

expression profiles and molecular heterogeneity of ATEVs and aEVs.

Figure 4. Single ADEV multiplex analysis showed comparable classic EV markers in ATEVs and aEVs. Beeswarm plots showing tetraspanin profiles for ATEVs (**A**) and aEVs (**B**). The x-axis reflects tetraspanin detection antibody and the top label indicates the tetraspanin capture antibody. Both ATEVs and aEVs showed similar but not identical tetraspanin event frequencies (see manuscript text for details; n =3). SP-IRIS representative images per capture spot CD63, CD81, CD9 for ATEVs (**C**) and aEVs (**D**). Scale 10 um. All events were detected above mouse IgG isotype control levels (data not shown).

Total internal reflection fluorescence (TIRF) microscopy allows simultaneous detection of human adipose-derived EV protein and mRNA cargo at single EV resolution

To establish an adipose liquid biopsy for the interrogation of circulating plasma EVs, we next applied our adipose-enriched proteomics candidates to the development of adipose-enriched surface capture targets and molecular beacons for mRNA detection cargo at single EV

resolution. After immobilization of EVs using anti-CD63 and CD9 hybridized surface[36], two

adipocyte markers (adipocyte plasma membrane protein, APMAP; and perilipin) were selected

for detection of adipocyte-specific surface proteins on plasma EVs. Furthermore, we examined

the presence of specific mRNA molecules using molecular beacons designed against PPARG

and ADIPOQ, which are known to be highly expressed in adipocytes[37-38]. The results of the

protein detection revealed the presence of APMAP, perilipin, and CD63 on both single ATEVs

and aEVs (**Fig. 5**). Similarly, detection of the molecular beacons confirmed the presence of

PPARG and ADIPOQ mRNA within the immobilized vesicles. These results confirm our ability to

characterize plasma circulating, adipose-specific single EVs with simultaneous surface protein

and mRNA cargo.

Figure 5. Single ADEV cargo and surface marker analysis defines CD63+/APMAP+/PPARG+ as an adipose-specific EV signature for liquid biopsy development. (A) Schematic showing how TIRF works. **(B)** ATEV and aEV (**C**) TIRF analysis with simultaneous protein (APMAP, perilipin, CD63; left) and mRNA cargo (ADIPOQ, PPARG; right). (n = 3 paired biological replicates). ADIPOQ: Adiponectin, APMAP: adipocyte plasma membrane-associated protein, PPARG: peroxisome proliferator-activated receptor gamma, ADEV: adipose-derived EVs, ATEVs: adipose tissue-derived EVs, aEVs: mature adipocyte derived EVs

Multiparametric single particle analysis of plasma EVs and validation of an adipose liquid biopsy

Lastly, to explore the liquid biopsy utility of our adipose-derived EV analysis, we applied two complementary single EV methods to the plasma EVs (bEVs) that corresponded to the matched ATEVs and aEVs from the same individual (**Fig. 1, Fig. 4, Fig. 5**). SP-IRIS analysis was used to explore the tetraspanin distribution of bEVs and unlike the matched ATEV and aEV analyses, CD81 capture, rather than CD63 capture, resulted in the greatest total mean particle count and of CD81+ single positive EVs (**Fig. 6A,B**). Colocalization analysis demonstrated a high percentage of CD63+/CD81+ double-positive bEVs (**Supplementary Fig. 3**). These results underscore the heterogeneity of EV subsets across various biospecimens.

TIRF analysis of bEVs (**Fig. 6C,D**) and neat plasma (**Fig. 6E,F**) was employed to validate our ability to capture and characterize adipose-derived EVs in the circulation of bariatric patients. We successfully captured single, plasma-derived EVs that displayed EV (CD63) and adipocyte surface markers (APMAP, perilipin) and contained adipocyte mRNA cargo (ADIPOQ, PPARG). We next sought to evaluate the sensitivity of our sEV assay to detect low-abundant ADEVs in the circulation of bariatric patients by performing TIRF analysis on neat plasma (i.e. without EV isolation) from the same patients. We were again able to capture sEVs that displayed EV (CD63) and adipocyte surface markers (APMAP, perilipin) and contained adipocyte mRNA cargo (ADIPOQ, PPARG).

Figure 6. Multiparametric single EV analysis of plasma EVs with simultaneous adipose target protein and mRNA cargo characterization. **(A, B)** SP-IRIS tetraspanin profile characterization and colocalization representative images of bEVs (scale 10um). TIRF analysis on paired SEC-isolated plasma EVs (**C-D**) and neat plasma (**E-F**) from the same individual.

243   Simultaneous adipocyte and EV protein (APMAP, perilipin, CD63; left) and adipocyte mRNA

244   cargo (ADIPOQ, PPARG; right) detection in single EVs from bariatric patient plasma (n = 3

245   paired biological replicates).

Discussion

The dual roles of adipose-derived EVs (ADEVs) as biomarkers and pathogenic

conveyors of disease across a diversity of pathologies have recently garnered increased

attention[4,25,26,39]. Through the transfer of diverse bioactive cargoes, ADEVs can influence

disease mechanisms locally and systemically[26,40-42], particularly at the interface between

adipose tissue and systemic metabolic regulation[15]. Despite the growing interest in ADEVs, their

full translational potential has been hindered by methodological and biological challenges.

Firstly, standardized, reproducible and robust isolation protocols for tissue-derived EVs are

lacking[29]. Secondly, most ADEV studies have focused on rodent models[25-27], despite known

cross-species differences in adipose biology between rodents and humans[28]. Lastly, cell-

specific EV analysis is often technically challenging and also requires a priori knowledge of

known cell-specific markers for EV immunocapture[21]. Therefore, innovative approaches for

human adipocyte-specific EV analysis are needed to move the translational potential of ADEVs

beyond the laboratory and into the clinic.

Herein, we present a novel and systematic approach to develop a sensitive and robust

human adiposity-associated EV liquid biopsy. By utilizing paired biospecimens from the same

individual, we are the first to successfully interrogate paired human adipose EVs from tissue,

autologous cells, and plasma at both the bulk and single EV level. While there was expected

inter-individual heterogeneity across patients, remarkably, we observed intra-individual

similarities in the ADEVs from the same individual, irrespective of EV source (e.g., VAT,

cultured adipocytes, plasma). In this proof-of-concept study, we establish the feasibility of

analyzing low-abundant ADEVs in the circulation of bariatric patients and thus open new

avenues for studying ADEVs across diverse conditions known to be adversely associated with

obesity, such as cardiovascular disease[2], cancer[43], autoimmunity[44] and neurodegenerative

diseases[6].

Biological insights from adipose-derived extracellular vesicle proteomics

Adipose-derived EVs (such as ATEVs and aEVs) are central players in the growing field

of adipose tissue interorgan crosstalk and provide added complexity and scope to the adipose

secretome[15,42,45]. To gain insights into the potential novel contributions of EVs to the adipose

secretome, we performed bulk proteomics, KEGG and GO enrichment analysis on ATEVs and

aEVs. Our results confirm previous studies that EV biology, as reflected by the EV proteins

ANXA2[46], ANXA1[47], and CAV1[48], is a prominent aspect of the adipose milieu[15,20,45]. Additionally,

while ATEVs and aEVs exhibited shared proteomes, the analysis also revealed distinct proteins

between ATEVs and aEVs, reflecting diverse functional domains. Adipocyte EVs were primarily

enriched in histone proteins (H2BC26, MACROH2A1)[49], suggesting a more prominent role of

aEVs compared to ATEVs in chromatin remodeling and gene regulation. ATEVs were enriched

in proteins involved in proteasomal degradation (PSMD12, PSMD3) that have been implicated

in several cancers and neurodevelopmental disorders[50-52]. ATEVs were also enriched in the

autophagosomal regulator, SACM1L, which interestingly is the only phosphatidylinositol-4-

phosphate-specific phosphatase in humans[53,54].

A secondary objective of profiling the ATEV and aEV proteomes was to screen for

adipose-enriched candidates for subsequent adiposity liquid biopsy analysis from plasma. The

concept of liquid biopsy originates from the cancer field and encompasses minimally invasive

sampling of biofluids for circulating nanoparticles that reflect a tumor or tissue's molecular

signature—thereby obviating the need for tissue acquisition[55]. While early liquid biopsy studies

focused on circulating, cell-free tumor DNA or RNA, the discovery of EVs being found in every

biofluid studied has expanded the liquid biopsy field to include EVs as versatile and robust liquid

biopsy sources[21,56,57]. Toward this end, we leveraged our ADEV proteomics results to select

APMAP and perilipinas candidate adipose-enriched proteins for our plasma single EV analysis

and adiposity liquid biopsy development (see below). Both proteins are known to be involved in

diverse cellular functions, including adipocyte differentiation[58,59], adipogenesis[60,61], and insulin

resistance[60,62]. Notably, APMAP is also highly expressed in the brain where it appears to

complex with secretases involved in amyloid beta regulation. APMAP is additionally involved in

neuronal differentiation, mRNA splicing and the autophagy-lysosomal system[63,64]. In summary,

these results highlight the pathobiological relevance and biomarker potential of the human

ADEV proteome.

Single ADEV multiplex analysis as a next-generation tool for adiposity liquid biopsy

development

We leveraged two complementary, next-generation microchip-based platforms to

comprehensively phenotype paired human ADEVs from multiple sources at single EV

resolution[21]. SP-IRIS[35] and TIRF microscopy have emerged as valuable tools for the

visualization, quantification, and characterization of individual EVs[65,66]. SP-IRIS analysis allows

tetraspanin phenotyping at the single EV level. While CD9 and CD63 are commonly considered

pan-EV markers, it is known that tetraspanin profiles are heterogenous in presence and

quantity, depending on EV source, and can offer insights into EV subtypes[35]. In our paired

biospecimens, CD63 was the predominant tetraspanin expressed in ATEVs and aEVs but not in

plasma EVs from the same individual. These results highlight the molecular heterogeneity of EV

subsets from different biospecimens and underscore the need to use multiple EV surface

markers for immunocapture and/or characterization[35].

We employed TIRF microscopy as an ultrasensitive, high-resolution technique for the

simultaneous detection of specific proteins and mRNA molecules within immobilized paired

ATEVs, aEVs, bEVs and neat plasma from the same individual[66]. Starting with EVs from the

'primary source' (i.e. ATEVs and aEVs), TIRF microscopy defined our adiposity EV signature as

CD63+/APMAP+/PPARG+/ADIPOQ^{low}. Next, we technically validated the

CD63+/APMAP+/PPARG+/ADIPOQ^{low} adiposity EV signature in bEVs and neat plasma from

the same biological replicates. EVs isolated from paired plasma (bEVs), compared to matched

ATEVs and aEVs, demonstrated greater variable expression of CD63, APMAP, ADIPOQ and

PPARG. Notably, neat (unprocessed) plasma largely maintained the 'primary source' adiposity

EV signature with high CD63+/APMAP+/PPARG+/ADIPOQ^{low}. This observation bolsters the

utility of neat plasma EVs as suitable for tissue and cell-specific liquid biopsy applications.

Challenges and future direction

Tissue-derived EVs constitute only 0.2% of the total plasma EV pool, underscoring the

technical challenges in studying low abundance, tissue-derived and cell-specific EVs. Despite

this signal-to-noise challenge, adipose tissue is the most prominent contributor (>82%) of

tissue-derived EVs in human plasma[67], allowing us to employ our next generation, ultrasensitive

single EV methods to define an adiposity EV signature for liquid biopsy application. In this

innovative study, we are the first describe a multiparametric pipeline to characterize bulk and

single ADEVs from paired human visceral adipose tissue, autologous mature adipocytes, and

plasma. Importantly, this research pioneers the direct detection of an adipose-specific EV

signature from bariatric patient unprocessed plasma. Our study paves the way toward an

obesity EV liquid biopsy and broadens the therapeutic landscape of novel, human adipose-

enriched EV candidates worthy of future translational application.

Methods

Our study adheres to all applicable ethical regulations. The OSU Institutional Review
Board (IRB) approved the study, and all participants provided written informed consent. While
the study involved human subjects, they were not prospectively assigned to an intervention to
evaluate the effect on health-related biological outcomes or compare diets or treatments. As a
result, the study is exempt from registration in Clinical Trials.gov.

Study participants: VAT biopsies and plasma samples were obtained from patients with
obesity (n = 11, average age: 41.6 years old SD: 12.98, BMI: 40.97, SD: 6.00) who underwent
elective bariatric surgery at The Ohio State University (OSU) Center for Minimally Invasive
Surgery. Clinical data is summarized in **Supplementary Table 2**. Subjects received no
compensation for participation.

Adipose tissue dissociation and mature adipocyte culture for EV isolation: Adipose tissue
samples were obtained during bariatric surgery and dissociated within an hour from
procurement using collagenase digestion. The resulting cell suspension was centrifuged at 500
x g for 10 minutes to separate the mature adipocytes, the stromal vascular fraction (SVF), and
the dissociation solution. The mature adipocytes were cultured for 24 hours in Dulbecco's
Modified Eagle Medium/Ham's Nutrient Mixture F-12 (DMEM/F12), supplemented with 40%
M199 basal medium with Earle's salts and L-glutamine (MCDB201 Media), 2% Fetal Bovine
Serum (FBS), 1x Penicillin/Streptomycin, 1 nanomolar (nM) dexamethasone, L-ascorbic acid 2-
phosphate (LAAP) 0.1 millimolar (mM), Insulin-Transferrin-Selenium (ITS) MIX 1x, Linoleic acid-
Albumin 1x, insulin 5 micrograms per microliter (ug/uL), and Triiodothyronine (T3) 1 nM.

ADEV isolation via tangential flow filtration (TFF): Adipose tissue dissociation media and
mature adipocyte culture media were first filtered through 0.2um filters. Media was subsequently
concentrated into 5 ml and diafiltrated using TFF with a 500 kDa filter for purification at a flow
rate of 35 ml/min as previously described[68]. After TFF, the retentates were concentrated using
centrifugal units (30 kDa MWCO, MilliporeSigma Amicon Ultra Centrifugal Filter Unit, Fisher
Scientific) at 4000 × g for 30 min to a final volume of 100uL.

Plasma EV isolation via size exclusion chromatography (SEC): Whole blood was collected
in (ethylenediaminetetraacetic acid) EDTA tubes and centrifuged at 1500 x g, 10min at RT to
separate the plasma from blood cells. The plasma was then transferred to a new tube and
subjected to two sequential centrifugations at 2,500 x g for 15 min at RT. The resulting platelet-
depleted plasma was collected and subjected to EV isolation via SEC using the qEV single EV

columns. A total of 8 fractions were collected, and fractions 1-4 were used for EV

characterization. Briefly, 150 μL of the platelet-depleted plasma was loaded onto the top of the

qEV column, and EVs were separated based on their size and surface properties. The column

was eluted with phosphate-buffered saline, and 200 μL fractions were collected. Fractions 1-4

were pooled and concentrated using centrifugal filters units (30 kDa MWCO, MilliporeSigma

Amicon Ultra Centrifugal Filter Unit, Fisher Scientific) at 4000 × g for 30 min to a final volume of

100uL.

Microfluidic resistive pulse sensing (MRPS): Particle number concentrations and size

distributions were measured by microfluidic resistive pulse sensing using the Spectradyne nCS1

instrument (Spectradyne, Torrence, CA) equipped with C-400 polydimethylsiloxane cartridges to

cover a size range of approximately 65-400nm in vesicle diameters. Data analysis was

performed with the nCS1 Data Analyzer (Spectradyne).

Western blotting: EV samples were lysed in radioimmunoprecipitation assay (RIPA) buffer

(Thermo Scientific, Cat# J62524AD) with the addition of Thermo Scientific Pierce Protease and

Phosphatase Inhibitor (Thermo Scientific, Cat#A32959) for 15 min on ice. Protein

concentrations were quantified using a Micro BCA™ Protein Assay Kit (Thermo Scientific, Cat#

23235). Equivalent amounts of sample proteins in Laemmli buffer (With reducing agent, 2-

Mercaptoethanol (Sigma-Aldrich, CAT# 8057400005) were electrophoresed on 4-20% Mini-

PROTEAN® TGX Stain-Free gels (Bio-rad, Cat#4568096) and then transferred onto a

polyvinylidene fluoride (PVDF) membrane (Biorad, Cat#1620177). Membranes were blocked

and then probed with primary antibody diluted in TBS-T overnight at 4°C and then with

horseradish peroxidase conjugated secondary antibody for 1 hour at RT. Immunoreactivity was

determined using enhanced chemiluminescence (ECL) solutions (Biorad, Cat# 1705060) and

visualized using a Biorad ChemiDoc™ MP imaging system.

Single-particle interferometric reflectance imaging sensing (SP-IRIS): Unchained Labs

(Boston, MA, USA) silicone chips coated with tetraspanin-specific capture spots (CD63, CD9,

and CD81) and a negative control spot (mouse IgG isotype). Chips were incubated for 1 hour at

RT with 1 × 10E8 ADEVs and 1x10E7 plasma EVs diluted in a final volume of 60 μL of

incubation buffer A at room temperature. After the incubation, chips were washed 3 times for 3

min on an orbital plate shaker with wash solution B. The chips were scanned with the SP-IRIS™

R200 reader (Unchained labs) by the ExoScanner software (Unchained labs, Boston, MA,

USA). The data was analyzed using SP-IRISer software (Unchained labs, Boston, MA, USA).

Transmission electron microscopy: Two 20-μL droplets of deionized (DI) water and two droplets of UranyLess EM contrast stain (Electron Microscopy Science) were placed on a parafilm surface. Prior to that, TEM grids underwent a 1-minute plasma treatment. Subsequently, 10 μL of ATEVs, aEVs, and bEVs were deposited onto the plasma-treated surface through drop-casting. The samples were allowed to incubate on the TEM surface for 1 minute, followed by gentle blotting with filter paper to remove excess liquid. The TEM grids were promptly washed by dipping them into a DI water droplet, followed by blotting with filter paper. This washing step was repeated with another DI water droplet. The same technique was then repeated for the contrast stain, with each incubation lasting 22 seconds. The TEM grids were carefully stored in a grid box overnight to ensure complete drying before imaging. TEM imaging was performed using a Tecnai TF-20 microscope operating at 200kV.

Single EV analysis using Total internal reflection fluorescence microscopy

Antibody functionalization of the biochip: In accordance with the methodology delineated in the preceding publication, a biochip featuring a gold-coated substrate was painstakingly prepared and subsequently functionalized using biotin-PEG-SH[36]. The functionalization of the biochip with antibodies entailed a series of specific steps. A uniform working volume of 20 μL was employed in each well to introduce various solutions. To ensure an even distribution of solutions within the wells, all incubation stages were conducted on a shaker. Before the antibody functionalization, a thorough washing of each well was conducted by pipetting DI water ten times. Following this, a solution of NeutrAvidin (NA; Thermo Fisher Scientific) at a concentration of 50 μg/mL, diluted in phosphate-buffered saline (PBS; Thermo Fisher Scientific), was added to each well. This mixture was allowed to incubate at room temperature for 1 hour to bind with the biotin motifs on the gold surface. Excess NeutrAvidin was subsequently removed by pipetting PBS ten times, and this rinsing process was repeated three times.

Introduction of biotinylated capture antibodies: For the introduction of biotinylated capture antibodies, the antibodies were first biotinylated using the EZ-Link micro Sulfo-NHS-biotinylation kit (Thermo Fisher Scientific). Each antibody was then diluted to a concentration of 10 μg/mL in a 1% (w/v) solution of bovine serum albumin (BSA; Sigma-Aldrich) in PBS. The prepared biotinylated capture antibody mixture was added to each well and allowed to incubate at room temperature for 1 hour. To remove excess protein, the wells were washed by pipetting PBS up and down ten times, with this rinsing process being repeated three times.

Molecular beacons hybridization: The molecular beacons (MBs) were diluted down to a concentration of 5 μM for each individual MB, using 12.5 × Tris EDTA (TE) buffer (Sigma-Aldrich), which was diluted in deionized (DI) water. This dilution served the dual purpose of stabilizing the MBs and permeabilizing the RNA-encasing membrane. The resulting mixture of MBs was subsequently diluted by a factor of 25 within the biofluid sample, and the biochip was then subjected to a 2-hour incubation period at 37°C, creating an environment devoid of light to facilitate the hybridization of the MBs with the target RNA.

Capture of EVs on biochip surface: In each well, a solution of BSA in PBS, with a concentration of 3% (w/v), underwent a 1-hour incubation at room temperature. This step aimed to prevent nonspecific EV capture and maintain single-vesicle resolution. Following the removal of BSA, the biofluid samples were then introduced into the wells and allowed to incubate for 2 hours at room temperature under dark conditions. PBS was introduced into the wells for a 5-minute incubation period, followed by pipetting PBS up and down ten times to eliminate excessive EVs and unhybridized MBs. This rinsing process was repeated four times within a dimly lit environment.

Detection antibodies targeting membrane proteins: Initially, a solution of BSA at a concentration of 3% (w/v) in PBS was added to each well and incubated at room temperature for one hour. This step aimed to prevent the nonspecific binding of the fluorophore-conjugated antibodies (detection antibodies). After the BSA solution was removed, a solution containing detection antibodies at a concentration of 1 μg/mL was prepared in 10% (w/v) normal goat serum (NGS; Thermo Fisher Scientific). Subsequently, the wells were subjected to a one-hour incubation at room temperature in a dark setting. To eliminate excess detection antibodies, PBS was added to the wells and allowed to incubate for 5 minutes. Following this, gentle pipetting up and down of the solution was carried out ten times. The rinsing procedure was repeated thrice within a dimly lit environment.

EV proteomics digestion: EVs were digested using Protifi S-trapTM and the published protocol was adapted to fit the project needs. (1) Briefly, samples were digested in lysis buffer consisting of 10% Sodium Dodecyl Sulfate (SDS) with 100 millimolar (mM) tetraethylammonium bromide (TEAB). Samples were sonicated and then clarified by centrifugation. Protein concentration was determined using NanoDrop A280 (Thermo Fisher Scientific). Disulfide bonds were reduced by addition of dithiothreitol (DTT), final concentration 20 mM, and incubation at 95 ºC for 15 minutes. Free cysteines were alkylated by the addition of iodoacetamide at a final concentration of 40 mM, followed by incubation in the dark for 30

minutes at room temperature (RT). Unreacted iodoacetamide was removed by centrifugation,

after which the supernatant was collected. Samples were mixed with binding buffer in a 1:6 ratio

of sample to buffer volume. The resulting solution was loaded onto a Protifi S-trapTM micro spin

column in increments of 300 µL, and centrifuged until all protein was bound to the column. 20

µL of digestion buffer was added, containing Trypsin Gold protease, mass spectrometry grade

(Promega, ref V528A) in a 1:10 enzyme-to-protein ratio for each sample. Trypsin was

resuspended in 50mM triethylammonium bicarbonate (TEAB) at a 1mg/mL concentration for the

stock solution (Sigma ref T7408). Samples were kept at 37ºC overnight and eluted with

sequential washes of 50 mM TEAB in water, 0.2% formic acid in water, and 0.2% formic acid in

50:50 water/acetonitrile. Samples were dried and resuspended in 0.1% formic acid in water for

downstream analysis.

LC-MS/MS analysis: Bottom-up proteomics data were collected on a Bruker Scientific

(Billerica, MA) commercial timsTOF Pro instrument coupled to a Bruker Scientific nanoElute LC.

For each sample, 200 ng was loaded. Samples were run in data-dependent acquisition (DDA)

mode, using parallel accumulation-serial fragmentation (PASEF). The LC column was 25 cm

long with 1.7 µm C18 particles (IonOpticks), and samples were run at a rate of 0.401 µL/min,

using a linear reverse phase 120-minute gradient with a starting concentration of 2.0%

acetonitrile (ACN) with 0.1% formic acid (FA) and ending concentration of 80.0% ACN with

0.1% FA over a period of 110 minutes, with a 10-minute ending hold. The equilibration time was

estimated at 3.60 minutes. For MS/MS analysis, the mass range was set as 100 to 1700 m/z,

with 10 MS/MS scans, dynamic exclusion time of 0.4 min, and collision-induced dissociation

(CID) energy starting at 20.00 eV and ending at 59.00 eV.

Proteomics data processing and analysis: TimsTOF PASEF datasets were processed using

FragPipe (v. 19.1) in conjunction with the MS/MS search engine MSFragger (v. 3.7) and

IonQuant (v. 1.8.10). The Uniport human reviewed database was downloaded and used as the

input fasta file for spectral mapping. Common contaminants and decoys were added using

FragPipe software. The fraction of decoys was set to 50% and was generated by reversing

sequences from the target database. This target-decoy search was used to control the false

discovery rate (FDR). (https://doi.org/10.48550/arXiv.1501.00537) The precursor mass

tolerance was 20 ppm, and the enzyme was set to trypsin with 2 missed cleavages. Peptide

length was set to 7 – 50, and peptide mass range was 500 – 5,000. Variable modifications were

set to include methionine oxidation (+15.9949) and N-terminal acetylation (+42.0106).

Carbamidomethylation on cysteine residues from iodoacetamide treatment (+57.021464) was

set as a fixed modification. The number of variable modifications on a given peptide was set to

3. Percolator was used for peptide-spectrum match (PSM) validation, and MS1 quantification

was run using match between runs (MBR) with an FDR of 0.01. Intensity was normalized across

runs, using a log-ratio approach where the run with the highest number of ions is used as a

reference. Peptide-protein uniqueness was set to unique + razor, where unique peptide refers to

the number of PSMs that only map to the given protein, and razor refers to the total number of

PSMs that in support of a given protein.

FragPipe results were exported as TSV (tab separated value) files, compatible with

Excel for downstream processing. For identified proteins, precursor/MS1 intensity values were

used as quantitative markers of protein expression. Intensity is defined by IonQuant software as

the summed volume of the 0, +1 and +2 isotopes, where the volume is from ion mobility and

retention time information. Missing values were replaced with an appropriate low number

(5000). For individual protein identifications, values that were > 50% missing from either ATEV

or aEV groups were removed. Values that were not consistent (i.e., sometimes present,

sometimes missing) across a single group (ATEV or aEV) were also removed. The data were

log base 10 transformed and quantile normalized. A two-sided t test was used to compare

ATEVs to aEVs. P-values were FDR-corrected using the Benjamini-Hochberg method. A one-

sided t-test was used to look at the ratios of donor-matched aEV. FDR of 0.05 was considered

significant.

Proteome FragPipe outputs were further processed using the DAVID (Database for

Annotation, Visualization, and Integrated Discovery) bioinformatics tool[69,70].The GO (Gene

Ontology) pathway and KEGG (Kyoto Encyclopedia of Genes and Genomes) pathway analyses

were conducted. The DAVID tool allowed us to select the statistical analysis method, and we

opted to use the FDR as a threshold for identifying significant pathway enrichments.

Data availability

All data generated or analyzed during this study are included in this manuscript and its

supplementary information file. Further inquiries can be directed to the corresponding author.

Author contributions

P.L.P., W.H., E.R., S.M.M, conceptualization.

E.R., S.M.M. study design and funding acquisition.

P.L.P., D.S., J.G., K.T.N., B.L.B., J.K. S.N., S.A.B., B.J.N., V.W., W.H. investigation and
performed research.

P.L.P., J.G., K.T.N., J.K. organization and data analysis.

P.L.P., S.M.M completed the original draft of the manuscript.

E.R., S.M.M. data interpretation, revision, and correction of the manuscript. All authors
approved the final version of the manuscript.

Competing interests

The authors declare no competing interests.

Additional information

Correspondence and requests for materials should be addressed to Senior Corresponding
author.

Acknowledgements

We would like to thank William Ray for his illustrative media expertise in designing the figures
for this project. We also thank Ms. Mary Shows for editorial and proofreading assistance. Drs.
Kristy Townsend and Kristin Stanford provided technical input for adipose tissue processing.
SMM was supported by intramural funding from the Abigail Wexner Research Institute at
Nationwide Children's Hospital and NINDS (5K12NS098482-05). ER was supported by National
Institutes of Health (NIH) grants UG3/UH3TR002884 and U18TR003807. The content is solely
the responsibility of the authors and does not necessarily represent the official views of the
National Institutes of Health.

References

Seidell, J. C. & Halberstadt, J. The global burden of obesity and the challenges of prevention. *Ann Nutr Metab* **66 Suppl 2**, 7-12, doi:10.1159/000375143 (2015).

Powell-Wiley, T. M. *et al.* Obesity and Cardiovascular Disease: A Scientific Statement From the American Heart Association. *Circulation* **143**, e984-e1010, doi:10.1161/CIR.0000000000000973 (2021).

Avgerinos, K. I., Spyrou, N., Mantzoros, C. S. & Dalamaga, M. Obesity and cancer risk: Emerging biological mechanisms and perspectives. *Metabolism* **92**, 121-135, doi:10.1016/j.metabol.2018.11.001 (2019).

Picone, P., Di Carlo, M. & Nuzzo, D. Obesity and Alzheimer's disease: Molecular bases. *Eur J Neurosci* **52**, 3944-3950, doi:10.1111/ejn.14758 (2020).

Brenton, J. N. *et al.* Body mass index trajectories in pediatric multiple sclerosis. *Dev Med Child Neurol* **61**, 1289-1294, doi:10.1111/dmcn.14233 (2019).

Parimisetty, A. *et al.* Secret talk between adipose tissue and central nervous system via secreted factors-an emerging frontier in the neurodegenerative research. *J Neuroinflammation* **13**, 67, doi:10.1186/s12974-016-0530-x (2016).

Kershaw, E. E. & Flier, J. S. Adipose tissue as an endocrine organ. *J Clin Endocrinol Metab* **89**, 2548-2556, doi:10.1210/jc.2004-0395 (2004).

Longo, M. *et al.* Adipose Tissue Dysfunction as Determinant of Obesity-Associated Metabolic Complications. *Int J Mol Sci* **20**, doi:10.3390/ijms20092358 (2019).

Goossens, G. H. & Blaak, E. E. Adipose tissue dysfunction and impaired metabolic health in human obesity: a matter of oxygen? *Front Endocrinol (Lausanne)* **6**, 55, doi:10.3389/fendo.2015.00055 (2015).

Isaac, R., Reis, F. C. G., Ying, W. & Olefsky, J. M. Exosomes as mediators of intercellular crosstalk in metabolism. *Cell Metab* **33**, 1744-1762, doi:10.1016/j.cmet.2021.08.006 (2021).

Kranendonk, M. E. *et al.* Human adipocyte extracellular vesicles in reciprocal signaling between adipocytes and macrophages. *Obesity (Silver Spring)* **22**, 1296-1308, doi:10.1002/oby.20679 (2014).

Kalluri, R. & LeBleu, V. S. The biology, function, and biomedical applications of exosomes. *Science* **367**, doi:10.1126/science.aau6977 (2020).

Kumar, V., Kiran, S., Kumar, S. & Singh, U. P. Extracellular vesicles in obesity and its associated inflammation. *Int Rev Immunol* **41**, 30-44, doi:10.1080/08830185.2021.1964497 (2022).

Kwan, H. Y., Chen, M., Xu, K. & Chen, B. The impact of obesity on adipocyte-derived extracellular vesicles. *Cell Mol Life Sci* **78**, 7275-7288, doi:10.1007/s00018-021-03973-w (2021).

Bond, S. T., Calkin, A. C. & Drew, B. G. Adipose-Derived Extracellular Vesicles: Systemic Messengers and Metabolic Regulators in Health and Disease. *Front Physiol* **13**, 837001, doi:10.3389/fphys.2022.837001 (2022).

van Niel, G., D'Angelo, G. & Raposo, G. Shedding light on the cell biology of extracellular vesicles. *Nat Rev Mol Cell Biol* **19**, 213-228, doi:10.1038/nrm.2017.125 (2018).

La Torre, M. E. *et al.* Extracellular Vesicles Cargo in Modulating Microglia Functional Responses. *Biology (Basel)* **11**, doi:10.3390/biology11101426 (2022).

Tkach, M. & Thery, C. Communication by Extracellular Vesicles: Where We Are and Where We Need to Go. *Cell* **164**, 1226-1232, doi:10.1016/j.cell.2016.01.043 (2016).

Yang, E. *et al.* Exosome-mediated metabolic reprogramming: the emerging role in tumor microenvironment remodeling and its influence on cancer progression. *Signal Transduct Target Ther* **5**, 242, doi:10.1038/s41392-020-00359-5 (2020).

Zhao, R., Zhao, T., He, Z., Cai, R. & Pang, W. Composition, isolation, identification and function of adipose tissue-derived exosomes. *Adipocyte* **10**, 587-604, doi:10.1080/21623945.2021.1983242 (2021).

Shami-Shah, A., Travis, B. & Walt, D. R. Advances in extracellular vesicle isolation methods: a path towards cell-type specific EV isolation. *Extracell Vesicles Circ Nucleic Acids* **4**, 447-460, doi:10.20517/evcna.2023.14

 (2023).

Crewe, C. The challenges of interrogating adipose tissue extracellular vesicle functions in physiology. *Commun Biol* **5**, 581, doi:10.1038/s42003-022-03511-9 (2022).

Akbar, N., Pinnick, K. E., Paget, D. & Choudhury, R. P. Isolation and Characterization of Human Adipocyte-Derived Extracellular Vesicles using Filtration and Ultracentrifugation. *J Vis Exp*, doi:10.3791/61979 (2021).

Connolly, K. D. *et al.* Evidence for Adipocyte-Derived Extracellular Vesicles in the Human Circulation. *Endocrinology* **159**, 3259-3267, doi:10.1210/en.2018-00266 (2018).

Kulaj, K. *et al.* Adipocyte-derived extracellular vesicles increase insulin secretion through transport of insulinotropic protein cargo. *Nat Commun* **14**, 709, doi:10.1038/s41467-023-36148-1 (2023).

Gesmundo, I. *et al.* Adipocyte-derived extracellular vesicles regulate survival and function of pancreatic beta cells. *JCI Insight* **6**, doi:10.1172/jci.insight.141962 (2021).

Durcin, M. *et al.* Characterisation of adipocyte-derived extracellular vesicle subtypes identifies distinct protein and lipid signatures for large and small extracellular vesicles. *J Extracell Vesicles* **6**, 1305677, doi:10.1080/20013078.2017.1305677 (2017).

Borgeson, E., Boucher, J. & Hagberg, C. E. Of mice and men: Pinpointing species differences in adipose tissue biology. *Front Cell Dev Biol* **10**, 1003118, doi:10.3389/fcell.2022.1003118 (2022).

Vella, L. J. *et al.* A rigorous method to enrich for exosomes from brain tissue. *J Extracell Vesicles* **6**, 1348885, doi:10.1080/20013078.2017.1348885 (2017).

Freshney, R. *Culture of animal cells: a manual of basic technique and specialized applications*. 6th edn, (Wiley-Blackwell, 2010).

Hearnden, R., Sandhar, B., Vyas, V. & Longhi, M. P. Isolation of stromal vascular fraction cell suspensions from mouse and human adipose tissues for downstream applications. *STAR Protoc* **2**, 100422, doi:10.1016/j.xpro.2021.100422 (2021).

Cianciaruso, C. *et al.* Molecular Profiling and Functional Analysis of Macrophage-Derived Tumor Extracellular Vesicles. *Cell Rep* **27**, 3062-3080 e3011, doi:10.1016/j.celrep.2019.05.008 (2019).

Cianciaruso, C. *et al.* Primary Human and Rat beta-Cells Release the Intracellular Autoantigens GAD65, IA-2, and Proinsulin in Exosomes Together With Cytokine-Induced Enhancers of Immunity. *Diabetes* **66**, 460-473, doi:10.2337/db16-0671 (2017).

Kowal, J. *et al.* Proteomic comparison defines novel markers to characterize heterogeneous populations of extracellular vesicle subtypes. *Proc Natl Acad Sci U S A* **113**, E968-977, doi:10.1073/pnas.1521230113 (2016).

Breitwieser, K. *et al.* Detailed Characterization of Small Extracellular Vesicles from Different Cell Types Based on Tetraspanin Composition by ExoView R100 Platform. *Int J Mol Sci* **23**, doi:10.3390/ijms23158544 (2022).

Nguyen, L. T. H. *et al.* An immunogold single extracellular vesicular RNA and protein ((Au) SERP) biochip to predict responses to immunotherapy in non-small cell lung cancer patients. *J Extracell Vesicles* **11**, e12258, doi:10.1002/jev2.12258 (2022).

Ma, X., Wang, D., Zhao, W. & Xu, L. Deciphering the Roles of PPARgamma in Adipocytes via Dynamic Change of Transcription Complex. *Front Endocrinol (Lausanne)* **9**, 473, doi:10.3389/fendo.2018.00473 (2018).

Kern, P. A., Di Gregorio, G. B., Lu, T., Rassouli, N. & Ranganathan, G. Adiponectin expression from human adipose tissue: relation to obesity, insulin resistance, and tumor necrosis factor-alpha expression. *Diabetes* **52**, 1779-1785, doi:10.2337/diabetes.52.7.1779 (2003).

Garcia-Contreras, M. & Thakor, A. S. Human adipose tissue-derived mesenchymal stem cells and their extracellular vesicles modulate lipopolysaccharide activated human microglia. *Cell Death Discov* **7**, 98, doi:10.1038/s41420-021-00471-7 (2021).

Blandin, A. *et al.* Lipidomic analysis of adipose-derived extracellular vesicles reveals specific EV lipid sorting informative of the obesity metabolic state. *Cell Rep* **42**, 112169, doi:10.1016/j.celrep.2023.112169 (2023).

Niu, Q. *et al.* Adipose-derived mesenchymal stem cell-secreted extracellular vesicles alleviate non-alcoholic fatty liver disease via delivering miR-223-3p. *Adipocyte* **11**, 572-587, doi:10.1080/21623945.2022.2098583 (2022).

Thomou, T. *et al.* Adipose-derived circulating miRNAs regulate gene expression in other tissues. *Nature* **542**, 450-455, doi:10.1038/nature21365 (2017).

Pati, S., Irfan, W., Jameel, A., Ahmed, S. & Shahid, R. K. Obesity and Cancer: A Current Overview of Epidemiology, Pathogenesis, Outcomes, and Management. *Cancers (Basel)* **15**, doi:10.3390/cancers15020485 (2023).

Kwiat, V. R., Reis, G., Valera, I. C., Parvatiyar, K. & Parvatiyar, M. S. Autoimmunity as a sequela to obesity and systemic inflammation. *Front Physiol* **13**, 887702, doi:10.3389/fphys.2022.887702 (2022).

Crewe, C. & Scherer, P. E. Intercellular and interorgan crosstalk through adipocyte extracellular vesicles. *Rev Endocr Metab Disord* **23**, 61-69, doi:10.1007/s11154-020-09625-x (2022).

Hagiwara, K., Katsuda, T., Gailhouste, L., Kosaka, N. & Ochiya, T. Commitment of Annexin A2 in recruitment of microRNAs into extracellular vesicles. *FEBS Lett* **589**, 4071-4078, doi:10.1016/j.febslet.2015.11.036 (2015).

Jeppesen, D. K. *et al.* Reassessment of Exosome Composition. *Cell* **177**, 428-445 e418, doi:10.1016/j.cell.2019.02.029 (2019).

Crewe, C. *et al.* An Endothelial-to-Adipocyte Extracellular Vesicle Axis Governed by Metabolic State. *Cell* **175**, 695-708 e613, doi:10.1016/j.cell.2018.09.005 (2018).

Singh, R., Bassett, E., Chakravarti, A. & Parthun, M. R. Replication-dependent histone isoforms: a new source of complexity in chromatin structure and function. *Nucleic Acids Res* **46**, 8665-8678, doi:10.1093/nar/gky768 (2018).

Fararjeh, A. S. *et al.* Proteasome 26S Subunit, non-ATPase 3 (PSMD3) Regulates Breast Cancer by Stabilizing HER2 from Degradation. *Cancers (Basel)* **11**, doi:10.3390/cancers11040527 (2019).

Del Valle, L. *et al.* Insulin-like growth factor I receptor activity in human medulloblastomas. *Clin Cancer Res* **8**, 1822-1830 (2002).

Ma, J., Zhou, W., Yuan, Y., Wang, B. & Meng, X. PSMD12 interacts with CDKN3 and facilitates pancreatic cancer progression. *Cancer Gene Ther* **30**, 1072-1083, doi:10.1038/s41417-023-00609-y (2023).

Liu, K., Kong, L., Graham, D. B., Carey, K. L. & Xavier, R. J. SAC1 regulates autophagosomal phosphatidylinositol-4-phosphate for xenophagy-directed bacterial clearance. *Cell Rep* **36**, 109434, doi:10.1016/j.celrep.2021.109434 (2021).

Zhang, H. *et al.* PtdIns4P restriction by hydrolase SAC1 decides specific fusion of autophagosomes with lysosomes. *Autophagy* **17**, 1907-1917, doi:10.1080/15548627.2020.1796321 (2021).

Connolly, I. D., Li, Y., Gephart, M. H. & Nagpal, S. The "Liquid Biopsy": the Role of Circulating DNA and RNA in Central Nervous System Tumors. *Curr Neurol Neurosci Rep* **16**, 25, doi:10.1007/s11910-016-0629-6 (2016).

Yu, D. *et al.* Exosomes as a new frontier of cancer liquid biopsy. *Mol Cancer* **21**, 56, doi:10.1186/s12943-022-01509-9 (2022).

S, E. L. A., Mager, I., Breakefield, X. O. & Wood, M. J. Extracellular vesicles: biology and emerging therapeutic opportunities. *Nat Rev Drug Discov* **12**, 347-357, doi:10.1038/nrd3978 (2013).

Albrektsen, T., Richter, H. E., Clausen, J. T. & Fleckner, J. Identification of a novel integral plasma membrane protein induced during adipocyte differentiation. *Biochem J* **359**, 393-402, doi:10.1042/0264-6021:3590393 (2001).

Ambele, M. A., Dessels, C., Durandt, C. & Pepper, M. S. Genome-wide analysis of gene expression during adipogenesis in human adipose-derived stromal cells reveals novel patterns of gene expression during adipocyte differentiation. *Stem Cell Res* **16**, 725-734, doi:10.1016/j.scr.2016.04.011 (2016).

Ma, Y. *et al.* Identification of a Novel Function of Adipocyte Plasma Membrane-Associated Protein (APMAP) in Gestational Diabetes Mellitus by Proteomic Analysis of Omental Adipose Tissue. *J Proteome Res* **15**, 628-637, doi:10.1021/acs.jproteome.5b01030 (2016).

Konige, M., Wang, H. & Sztalryd, C. Role of adipose specific lipid droplet proteins in maintaining whole body energy homeostasis. *Biochim Biophys Acta* **1842**, 393-401, doi:10.1016/j.bbadis.2013.05.007 (2014).

Bialesova, L. *et al.* Epigenetic Regulation of PLIN 1 in Obese Women and its Relation to Lipolysis. *Sci Rep* **7**, 10152, doi:10.1038/s41598-017-09232-y (2017).

Gerber, H. *et al.* The APMAP interactome reveals new modulators of APP processing and beta-amyloid production that are altered in Alzheimer's disease. *Acta Neuropathol Commun* **7**, 13, doi:10.1186/s40478-019-0660-3 (2019).

Zhang, F. *et al.* Clusterin facilitates stress-induced lipidation of LC3 and autophagosome biogenesis to enhance cancer cell survival. *Nat Commun* **5**, 5775, doi:10.1038/ncomms6775 (2014).

Arab, T. *et al.* Characterization of extracellular vesicles and synthetic nanoparticles with four orthogonal single-particle analysis platforms. *J Extracell Vesicles* **10**, e12079, doi:10.1002/jev2.12079 (2021).

Nguyen, L. T. H. *et al.* An immunogold single extracellular vesicular RNA and protein (AuSERP) biochip to predict responses to immunotherapy in non-small cell lung cancer patients. *Journal of Extracellular Vesicles* **11**, e12258, doi:https://doi.org/10.1002/jev2.12258 (2022).

Li, Y. *et al.* EV-origin: Enumerating the tissue-cellular origin of circulating extracellular vesicles using exLR profile. *Comput Struct Biotechnol J* **18**, 2851-2859, doi:10.1016/j.csbj.2020.10.002 (2020).

Zhang, J. *et al.* Immunomagnetic sequential ultrafiltration (iSUF) platform for enrichment and purification of extracellular vesicles from biofluids. *Sci Rep* **11**, 8034, doi:10.1038/s41598-021-86910-y (2021).

Sherman, B. T. *et al.* DAVID: a web server for functional enrichment analysis and functional annotation of gene lists (2021 update). *Nucleic Acids Res* **50**, W216-W221, doi:10.1093/nar/gkac194 (2022).

Huang da, W., Sherman, B. T. & Lempicki, R. A. Systematic and integrative analysis of large gene lists using DAVID bioinformatics resources. *Nat Protoc* **4**, 44-57, doi:10.1038/nprot.2008.211 (2009).