1

Dependence of Contextual Modulation in Macaque V1 on Interlaminar Signal Flow

Shude Zhu[1], Yu Jin Oh[1], Ethan B. Trepka[1], Xiaomo Chen[1,2], Tirin Moore[1*]

Author affiliations:

[1]Department of Neurobiology and Howard Hughes Medical Institute, Stanford University School

of Medicine, Stanford, CA 94305, USA.

[2]Center for Neuroscience, Department of Neurobiology, Physiology, and Behavior, UC Davis,

Davis, CA 95618, USA.

*Correspondence: tirin@stanford.edu

Abstract

In visual cortex, neural correlates of subjective perception can be generated by modulation of

activity from beyond the classical receptive field (CRF). In macaque V1, activity generated by

nonclassical receptive field (nCRF) stimulation involves different intracortical circuitry than

activity generated by CRF stimulation, suggesting that interactions between neurons across V1

layers differ under CRF and nCRF stimulus conditions. Using Neuropixels probes, we measured

border ownership modulation within large, local populations of V1 neurons. We found that

neurons in single columns preferred the same side of objects located outside of the CRF. In

addition, we found that cross-correlations between pairs of neurons situated across

feedback/horizontal and input layers differed between CRF and nCRF stimulation. Furthermore,

independent of the comparison with CRF stimulation, we observed that the magnitude of border

ownership modulation increased with the proportion of information flow from feedback/horizontal

layers to input layers. These results demonstrate that the flow of signals between layers covaries

with the degree to which neurons integrate information from beyond the CRF.

Introduction

The classical receptive field (CRF) defines the region of the sensory periphery where appropriate stimulation evokes a spiking response in a neuron (Hartline, 1938; Alonso and Chen, 2009). Those evoked responses are largely driven by feedforward inputs from earlier stages of sensory processing. In macaque primary visual cortex (V1), neuronal activity is largely determined by ascending input from the dorsal lateral geniculate nucleus (dLGN) which arrives principally in layers 4Cα and 4Cβ (Hubel and Wiesel, 1972; Hendrickson et al., 1978; Blasdel and Lund, 1983; Callaway, 1998; Callaway, 2004). However, considerable evidence has established that the responses of sensory cortical neurons are also robustly influenced by stimulation concurrently presented outside of the CRF (Allman et al., 1985; Fitzpatrick, 2000; Albright and Stoner, 2002; Roelfsema, 2006; Angelucci et al., 2017). Specifically, whereas stimulation outside of the CRF alone fails to evoke spiking responses, such stimulation nonetheless alters the responses evoked by stimulation within the CRF. In visual cortex, these nonclassical receptive field (nCRF) effects often contribute to the neural correlates of visual perceptual phenomena, such as illusory contours (von der Heydt et al., 1984; Ramsden et al., 2001), visual salience and boundary segmentation (Knierim and van Essen, 1992; Sillito et al., 1995; Yan et al., 2018; Lee et al., 2002; Nothdurft et al., 1999), contour integration (Nelson and Frost, 1985; Kapadia et al., 1995; Li et al., 2006), figure-ground segregation (Lamme, 1995; Zipser et al., 1996; Poort et al., 2012), and border ownership (Zhou et al., 2000; von der Heydt, 2015; von der Heydt, 2023).

In contrast to the feedforward mechanisms underlying the CRF, nCRF effects are thought to be generated either by feedback from neurons with larger CRFs in higher areas (Hupe et al., 1998; Bullier et al., 2001; Bair et al., 2003; Nassi et al., 2013; Chen et al., 2017; Klink et al., 2017; Zhang et al., 2014; Nurminen et al., 2018; Keller et al., 2020; Pak et al., 2020; Vangeneugden et

al., 2019; Gieselmann and Thiele, 2022; Chen et al., 2014; van Kerkoerle et al., 2014), or through intracortical horizontal connections (Adesnik et al., 2012; Chisum et al., 2003; Stettler et al., 2002), or by both (Angelucci and Bressloff, 2006; Schwabe et al., 2006; Ichida et al., 2007; Angelucci et al., 2017; Liang et al., 2017; Self et al., 2013). In contrast to feedforward inputs, feedback and horizontal inputs to V1 both avoid layer 4C and terminate predominantly in superficial (layers 1-3) and deep (layers 5-6) layers (Rockland and Lund, 1983; Rockland and Pandya, 1979; Rockland and Virga, 1989; Anderson and Martin, 2009; Markov et al., 2014; Federer et al., 2021; Gilbert and Wiesel, 1983; Shmuel et al., 2005; Siu et al., 2021). These differences between feedforward and feedback/horizontal circuitry suggest that the flow of signals across V1 layers should covary with the type of visual stimulation.

Newly developed, high-density Neuropixels probes have enabled recordings from a large, dense population of neurons (Jun et al., 2017; Zhu et al., 2024; Trautmann et al., 2025), and dramatically increase the quantity of identifiable functional interactions between pairs of neurons, particularly in non-human primates (Trepka et al., 2022). Recently, this approach was used to recapitulate known circuit properties, such as the pairwise lead-lag relationship between simple and complex cells and the canonical laminar input-output relationship in macaque V1 (Trepka et al., 2022), as well as the visual hierarchy in mouse visual cortical (Siegle et al., 2021; Jia et al., 2022) and subcortical areas (Sibille et al., 2022).

Here, using Neuropixels probes, we recorded the activity of hundreds of neurons simultaneously from single V1 columns in anesthetized macaques during CRF and nCRF stimulation. We leveraged the high yields obtained from different layers to examine the organization and circuitry underlying contextual modulation, specifically border ownership (B_{own}). B_{own} is a form of contextual modulation in which neurons signal the occluding border between the

background and a foreground object using stimulus information beyond the CRF. This function

appears to be crucial for natural scene segmentation and object recognition (Nakayama et al.,

1995). It has been shown that many neurons in early visual areas, including V1, respond differently

to identical edges (borders) of objects when those objects lie at different locations outside the CRFs

(Zhou et al., 2000; Franken and Reynolds, 2021; Hesse and Tsao, 2016; Hesse and Tsao, 2023).

B_{own} emerges rapidly after visual response onset (Sugihara et al., 2011), yet the relative

contributions of feedforward, feedback, and horizontal circuits to B_{own} remain unknown. Also

unknown is whether B_{own} is present under anesthesia. Several other types of contextual modulation

have been demonstrated in anesthetized animals (Ramsden et al., 2001; Hupe et al., 1998; Bair et

al., 2003; Allman et al., 1985; Gilbert and Wiesel, 1990; Nothdurft et al., 1999; Webb et al., 2003;

Zarella and Ts'o, 2016; Muller et al., 2003; Henry et al., 2013; Bijanzadeh et al., 2018; although

see Lamme et al., 1998) and can involve inter-cortical feedback (Nurminen et al., 2018). Moreover,

B_{own} has been shown to occur independently of attention (Qiu et al., 2007; O'Herron and von der

Heydt, 2009), and before object shape recognition (Williford and von der Heydt, 2016; Ko and

von der Heydt, 2018), suggesting it may not rely on higher cognitive control or object recognition

areas. Rather, B_{own} may emerge through pre-attentive, automatic grouping mechanism in

low/middle level visual areas, signifying a representation of "proto-objects" in the early visual

cortex (von der Heydt, 2015; von der Heydt, 2023; Self et al., 2019).

Results

Classical RF stimulation in single V1 columns

Neuronal activity was recorded from two anesthetized rhesus macaques (M1 and M2, *Macaca mulatta*) using high-density, multi-contact Neuropixels probes (version 3A, IMEC Inc., Belgium) (Figure 1A) (Methods). These animals had no prior experience with awake experiments or exposure to the stimuli used in this study. Neuropixels probes were inserted into the lateral operculum of V1 with the aid of a surgical microscope at angles nearly perpendicular to the cortical surface. Prior to insertion, the probes were coated with a DiI derivative for subsequent histological visualization of the tracks (Methods) (Supplementary Figure S1). Four Neuropixels probe penetrations were made either in the opercular surface (M1, *pen 1-2*, ~4-6° eccentricities) or within the underlying calcarine sulcus (M2, *pen 3-4*, ~6-10° eccentricities) using the programmable channel selection feature of the Neuropixels probe. The boundaries of laminar compartments were estimated by combining current-source density (CSD) measurements and histological data (Figure 1B; Supplementary Figure S1) (Methods) (Zhu et al., 2024; Carr et al., 2025). Each recorded neuron was assigned to one of four laminar compartments, specifically 5/6, 4C, 4A/B, 2/3 (mean thickness: 489, 281, 311, 650 μm, respectively, consistent with previous anatomical (O'Kusky and Colonnier, 1982; Lund, 1973) and CSD (Self et al., 2013; Chen et al., 2017) studies). We measured the responses from a total of 677 visually driven neurons (N = 159, 181, 210, and 127 per recording) to classical receptive field (CRF) stimulation, and 621 neurons (N = 142, 179, 152, and 148 per recording) to nonclassical receptive field (nCRF) stimulation (Methods). As a result of the nearly perpendicular penetrations, the visual CRFs of V1 neurons were largely overlapping across the cortical depth (Figure 1C).

115   During each recording session, neurons were first tested with CRF stimuli consisting of

116 drifting Gabor gratings with 1.5 degrees of visual angle (dva) in diameter, presented at the joint

117 CRF location of simultaneously recorded neurons (Methods). In each of the four recordings,

118 neurons across cortical depth exhibited very similar preferences to grating orientations (Figure

119 1D). The modal pairwise difference in preferred orientation between neurons across layers was

120 near zero (*pen1* = 5°; *pen2* = 5°; *pen3* = 1°; *pen4* = 1°). This result confirmed that our recordings

121 were nearly perpendicular and included neurons predominantly from single orientation columns.

122 Gratings were sized to largely restrict the stimuli within the CRFs of recorded neurons to maximize

123 the extent to which V1 responses were driven in large proportion by feedforward circuitry. In V1,

124 geniculocortical inputs initially drive neurons within layers 4Cα and 4Cβ (Hubel and Wiesel,

125 1972; Hendrickson et al., 1978; Blasdel and Lund, 1983), which then propagate to supragranular

126 and infragranular layers (Callaway, 1998; Callaway, 2004).

128 **Nonclassical RF stimulation and border ownership**

129 Within each of the recordings, we also examined the responses of V1 neurons to nCRF stimuli. It

130 is known that neurons in primate visual cortex integrate visual information from far beyond their

131 CRFs (Albright and Stoner, 2002; Allman et al., 1985; Fitzpatrick, 2000; Roelfsema, 2006;

132 Angelucci et al., 2017). One example of contextual modulation is the selectivity of neurons to

133 border ownership (B_{own}). Specifically, neurons in early visual areas respond differently to identical

134 edges (borders) of objects when those objects lie at different locations outside the CRFs (Zhou et

135 al., 2000; Franken and Reynolds, 2021; Hesse and Tsao, 2016; Hesse and Tsao, 2023) (Figure 2).

136 As with other forms of nCRF modulation, B_{own} is thought to emerge either by feedback from higher

137 areas (Craft et al., 2007; Jehee et al., 2007; Layton et al., 2012; Eguchi and Stringer, 2016;

Wagatsuma et al., 2021; Mehrani and Tsotsos, 2021) or from horizontal connections within the same area (Zhaoping, 2005; Kogo et al., 2010). In V1, feedback and horizontal inputs arrive principally within supragranular and infragranular layers, i.e., they avoid layer 4 (Rockland and Pandya, 1979; Rockland and Lund, 1983; Rockland and Virga, 1989; Anderson and Martin, 2009; Markov et al., 2014; Federer et al., 2021; Gilbert and Wiesel, 1983; Shmuel et al., 2005; Siu et al., 2021). Thus, V1 activity generated by nCRF stimulation is known to involve different circuits compared to the activity generated by CRF stimulation.

In this study, border ownership test stimuli consisted of uniformly white or black objects (squares, 8 x 8 dva) on a black or white background, respectively, as in previous studies (Zhou et al., 2000) (Methods). One border of the object fell within the CRFs of the recorded neurons. Within the CRF, the oriented border could either be a dark-light edge or light-dark edge, resulting in two different local contrast polarity (LC) conditions (Figure 2A). Outside the CRF, the stimulus configuration differed for the same LC condition such that the oriented border could belong to opposite sides of an object relative to the CRF. Crucially, across these two object side conditions, the stimulus falling within the CRF was identical (Figure 2B) and was identical within an area that extended well beyond the CRF (8 x 16, dva). In this stimulus configuration, differences in evoked responses between the two object side conditions signify selectivity to border ownership (Zhou et al., 2000). The statistical significance was determined by a two-factor ANOVA on the mean spike counts for each condition, with object side (Side 1 and Side 2) and local contrast (LC1 and LC2) as the factors, along with their interaction (Methods).

We observed neurons with B_{own} in our recordings from macaque V1. Figures 2C and 2D shows examples of four neurons selective to B_{own}. Each of these neurons exhibited a preference for LC, with two neurons preferring one contrast polarity and two preferring the other. In addition,

each neuron responded differently depending on which side of the object appeared relative to the

CRF, thus exhibiting B_{own}. Three of the neurons responded more vigorously to borders belonging

to the lower-left side of object (Side 1) ($p < 10^{-14}$, 10^{-10}, 10^{-3} for neurons 1, 2, and 4, respectively),

while one neuron responded more vigorously to the border belonging to the upper-right side of the

object (Side 2) ($p = 0.014$, neuron 3). Furthermore, one of the neurons showed a consistent

preference for Side 1 across both LCs (neuron 4).

To quantify the B_{own}, as in previous studies (Zhou et al., 2000), we calculated a B_{own}

modulation index, defined as the difference in neuronal responses to borders belonging to opposite

sides of objects, normalized by the maximum response for each neuron (Methods). Responses to

the same side of object, but under opposite LC conditions, were averaged together to control for

global luminance differences in the overall display. Across the 621 neurons recorded in four

sessions, we found that 133 neurons (21.4% overall; *pen1* = 28.2%; *pen2* = 27.9%; *pen3* = 18.4%;

pen4 = 8.4%) responded differently to identical borders yet belonging to opposite sides of the

object, a proportion consistent with previous studies in awake monkeys (Zhou et al., 2000).

Neurons with B_{own} were found within all cortical laminar compartments, including input layer 4C

(5/6: 28/144 = 19.4%; 4C: 48/212 = 22.6%; 4A/B: 19/114 = 16.7%; 2/3: 28/99 = 28.3%) (Table

S1), and the proportion was independent of laminar compartment ($\chi^2(3) = 4.78$, $p = 0.19$, chi-

square test of homogeneity). Of all V1 neurons, 96 (15.5%) exhibited B_{own} depending on the LC

(5/6: 11.1%; 4C: 17.9%; 4A/B: 14.0%; 2/3: 20.2%), while 37 (6.0%) showed significant effects

independent of local contrast (5/6: 8.3%; 4C: 4.7%; 4A/B: 2.6%; 2/3: 8.1%). Thus, the majority of

V1 neurons with B_{own} exhibited selectivity for only one local contrast, and neurons with B_{own}

invariant to local contrast were rare, also consistent with previous observations (Zhou et al., 2000).

Columnar organization of border ownership selectivity

We next considered whether neurons with B_{own} comprised a distinct group, or instead reflected a general tendency of neurons to prefer one side of the object, with statistically significant neurons falling at the tails of the distribution. We leveraged the large number of simultaneously recorded single neurons to measure the distribution of B_{own} modulation across laminar compartments in each session. We found that in each of the four recordings, the distributions of B_{own} modulation generally appeared unimodal, with neurons exhibiting B_{own} falling at the distribution tails (Figure 2E). Surprisingly, we also found that within individual recordings, neurons across the cortical depth tended to share similar preferences for the side of the object. That is, for each recording, the B_{own} modulation index differed significantly from 0 (Median modulation index: *pen1* = 0.14, p < 10^{-12}; *pen2* = 0.08, p = 0.04; *pen3* = -0.18, p < 10^{-15}; *pen4* = 0.10, p < 10^{-5}, sign test against zero-median). For example, the first recording (*pen1*) was performed in a column in which most neurons preferred an orientation of 135°. In addition, within that column, neurons tended to exhibit higher responses to a border when it belonged to the lower-left side of an object (side 1), as opposed to the upper-right side (side 2). Thus, V1 exhibited a continuum of B_{own}, and neurons within single columns tended to prefer the same side of objects located outside of the CRF.

Interlaminar cross-correlations among neurons within single V1 columns

The large number of simultaneously recorded neurons from each session also allowed us to measure interlaminar information flow among single neurons under different visual stimulation conditions. Temporally precise cross-correlations in spike trains offer a unique means of assessing functional interactions among neurons in neural circuits (Perkel et al., 1967). Cross-correlations can be interpreted as suggesting one of myriad putative circuit arrangements among neuronal

ensembles (Moore et al., 1970; Aertsen and Gerstein, 1985; Melssen and Epping, 1987; Ostojic et

al., 2009). The identification of such interactions has played an important role in elucidating neural

circuits in the mammalian visual system (Ts'o et al., 1986; Schwarz and Bolz, 1991; Reid and

Alonso, 1995; Alonso et al., 1996; Alonso and Martinez, 1998; Usrey et al., 1998; Nelson et al.,

1992; Briggs et al., 2013; Roe and Ts'o, 1999; Senzai et al., 2019; Denman and Contreras, 2014;

Siegle et al., 2021; Jia et al., 2022; Trepka et al., 2022; Smith and Kohn, 2008; Sibille et al., 2022).

In particular, it has been useful in specifying the flow of signals through local neural circuits

(Trepka et al., 2022) and distributed networks (Siegle et al., 2021; Jia et al., 2022; Sibille et al.,

2022). Notably, past studies have demonstrated that neuronal cross-correlations are dynamic and

can depend on stimulus context and behavioral variables (Hembrook-Short et al., 2019; Martin

and von der Heydt, 2015; Hung et al., 2007; Hirabayashi and Miyashita, 2005; Hirabayashi et al.,

2010; Steinmetz et al., 2000).

219   Given that V1 activity generated by nCRF stimulation is thought to involve different

circuitry than activity generated by CRF stimulation (Angelucci et al., 2017), we next asked if the

pattern of cross-correlations measured under classical and nonclassical stimulus conditions might

differ in a manner consistent with models of their underlying circuitry. Specifically, modulation

of visual responses during nCRF stimulation, such as border ownership stimuli, is thought to

emerge from either feedback from higher areas or horizontal connections within the same areas

(Bair et al., 2003; Bullier et al., 2001; Hupe et al., 1998; Chen et al., 2017; Gieselmann and Thiele,

2022; Keller et al., 2020; Klink et al., 2017; Nassi et al., 2013; Nurminen et al., 2018; Angelucci

et al., 2017; Angelucci and Bressloff, 2006; Ichida et al., 2007; Schwabe et al., 2006; Pak et al.,

2020; Vangeneugden et al., 2019; Zhang et al., 2014; Adesnik et al., 2012; Chisum et al., 2003;

Stettler et al., 2002; Liang et al., 2017; Craft et al., 2007; Jehee et al., 2007; Chen et al., 2014; Self

et al., 2013; van Kerkoerle et al., 2014), both arriving principally in superficial (layers 1-3) and

deep (layers 5-6) layers (Rockland and Lund, 1983; Rockland and Pandya, 1979; Rockland and

Virga, 1989; Markov et al., 2014; Federer et al., 2021; Anderson and Martin, 2009; Gilbert and

Wiesel, 1983; Shmuel et al., 2005; Siu et al., 2021). In contrast, CRF responses are driven

predominantly by feedforward circuitry and geniculocortical inputs to layers 4C (Hubel and

Wiesel, 1972; Hendrickson et al., 1978; Blasdel and Lund, 1983; Callaway, 1998; Callaway,

2004). This suggests that the pattern of cross-correlations among neurons across V1 layers may

differ under CRF and nCRF stimulus conditions.

Across recording sessions, we calculated cross-correlograms (CCGs) from all possible

pairwise combinations of neuronal spike trains during both CRF (Gabor gratings) and nCRF

(border ownership) stimulation conditions. Figure 3A shows an example recording session (M1,

pen1) in which 159 visually responsive neurons were recorded during CRF stimulation, and CCGs

were computed for 7,956 pairwise combinations of neurons (Methods). CCGs were determined to

be significant if the jitter-corrected CCG peak occurred within 10ms of zero-time lag, and

exceeded 7 standard deviations (SD) above the mean of the noise distribution (Siegle et al., 2021;

Trepka et al., 2022) (Methods). In this recording, 13.8% of the total CCGs computed were found

to be significant. Eight significant pairs are highlighted in Figure 3A, with their corresponding

CCGs shown in Figure 3B. The peak lag (PL), defined as the relative time delay in occurrence of

peak correlation between two spike trains, was restricted to within 10ms. This metric approximates

the synchrony and/or the direction of information flow between neuronal pairs (Aertsen and

Gerstein, 1985; Moore et al., 1970; Melssen and Epping, 1987; Ostojic et al., 2009). These

examples illustrate the tendency of CCGs to corroborate the circuitry within V1 columns under

CRF conditions in that 1) PLs generally increase with larger distances between pairs, and 2) layer

4 neurons tend to lead superficial neurons (Trepka et al., 2022). We exploited this latter point to

test the extent to which the sequence of information flow across layers differs between CRF and

nCRF conditions.

Comparison of cross-correlations during CRF and nCRF stimulation.

From the 677 and 621 neurons recorded across the 4 recording sessions during CRF and nCRF

stimulation, there were 41,759 and 28,430 pairwise combinations included, respectively

(Methods). Of those, we found that 6,937 (16.6%) and 4,826 (17.0%) of the CCGs were significant

(CRF: *pen1*: 1099/7956 = 13.8%, *pen2*: 2402/11431 = 21.0%, *pen3*: 2174/17780 = 12.2%, *pen4*:

1262/4592 = 27.5%; nCRF: *pen1*: 447/2473 = 18.1%, *pen2*: 2750/11823 = 23.3%, *pen3*: 839/6998

= 12.0%, *pen4*: 790/7136 = 11.1%). Notably, the proportion of significant connections were nearly

identical under the two stimulus conditions, and also were within the range observed in previous

studies of macaque V1 (Chu et al., 2014; Hembrook-Short et al., 2019; Smith and Kohn, 2008;

Kohn and Smith, 2005).

As noted above, modulation of visual responses during nCRF stimulation, such as border

ownership stimuli, is thought to be generated principally by feedback/horizontal connections

arriving within superficial and deep layers, which contrasts with feedforward input driving CRF

stimulation (Figure 3C). Thus, we asked whether cross-correlations between pairs of neurons

situated across laminar compartments depended on the type of visual stimulation. We focused our

analysis on CCGs computed from pairs of neurons in which one neuron was located within laminar

compartments receiving feedback/horizontal inputs (FH$_i$), namely layers 2/3 and 5/6, and the other

was located within compartments relatively devoid of those inputs, namely 4C and 4A/B. This

analysis compared the relative spike timing between neuronal pairs during CRF and nCRF

stimulation. If indeed FH_i contribute more to activity during the latter than the former, then we

should expect this to be reflected in lead-lag relationships of the CCGs.

A comparison of V1 CCGs from key interlaminar neuronal pairs during CRF and nCRF

stimulation is shown for an example recording session (M1, *pen1*) in Figure 3D. During CRF

stimulation, the timing of spikes within compartments 4C and 4A/B generally preceded, or was

simultaneous with, spikes within FH_i compartments (CRF: $4C\text{-}2/3_{PL} = 2ms$, $p < 10^{-7}$; $4A/B\text{-}2/3_{PL}$

$= 2ms$, $p < 10^{-28}$; $4C\text{-}5/6_{PL} = 0ms$, $p = 0.52$; $4A/B\text{-}5/6_{PL} = 0ms$, $p = 0.64$; Sign test against zero-

median). In contrast, during nCRF stimulation, relative spike timing was shifted in favor of

neurons in FH_i compartments. Specifically, during nCRF stimulation, the timing of spikes within

compartments 4C and 4A/B lagged or was simultaneous with, spikes within FH_i compartments

(nCRF: $4C\text{-}2/3_{PL} = 0ms$, $p = 0.80$; $4A/B\text{-}2/3_{PL} = 0ms$, $p = 0.70$; $4C\text{-}5/6_{PL} = -2.5ms$, $p = 0.11$; 4A/B-

$5/6_{PL} = 0ms$, $p = 0.56$; Sign test against zero median). To further quantify the apparent shift in lead-

lag relationships, we calculated an asymmetry index for each CCG. The asymmetry index is

computed as the difference in the CCG integral on either side of zero-time lag, and summarizes

the lead-lag relationship independent of the CCG peak (Jia et al., 2022) (Methods). Consistent with

the analyses of PL, we found that CCG asymmetries differed significantly between CRF and nCRF

stimulation for each of the interlaminar comparisons. In each case, the distribution of asymmetries

during nCRF stimulation shifted to favor neurons in FH_i laminar compartments, when compared

to CRF stimulation ($4C\text{-}2/3$: $CRF_{Asym} = 0.036$, $nCRF_{Asym} = -0.004$, $\Delta = -0.04$, $p < 10^{-6}$; $4A/B\text{-}2/3$:

$CRF_{Asym} = 0.046$, $nCRF_{Asym} = -0.005$, $\Delta = -0.051$, $p < 10^{-11}$; $4C\text{-}5/6$: $CRF_{Asym} = -0.010$, $nCRF_{Asym}$

$= -0.044$, $\Delta = -0.034$, $p = 0.0015$; $4A/B\text{-}5/6$: $CRF_{Asym} = 0.012$, $nCRF_{Asym} = -0.028$, $\Delta = -0.04$, $p <$

10^{-5}; Wilcoxon rank-sum test). Thus, CCG data from this single recording session indicated that

relative spike timing was earlier within FH_i laminar compartments during nCRF stimulation than

during CRF stimulation.

Across all recordings, we identified a total of 2,573 and 2,118 significant neuronal pairs

from the four interlaminar combinations during CRF (*pen1* = 274; *pen2* = 1187; *pen3* = 560; *pen4*

= 302) or nCRF stimulation (*pen1* = 210; *pen2* = 1226; *pen3* = 382; *pen4* = 300), respectively, and

compared their CCG asymmetries. Similar to the pattern observed in the example recording, we

observed a consistent shift in lead-lag relationships for each of the interlaminar comparisons

(Figure 4). Specifically, during nCRF stimulation, the distribution of asymmetries during nCRF

stimulation shifted to favor neurons in FH_i laminar compartments, when compared to CRF

stimulation (4C-2/3: CRF_{Asym} = 0.028, $nCRF_{Asym}$ = 0.012, Δ = -0.016, $p < 10^{-9}$; 4A/B-2/3: CRF_{Asym}

= 0.038, $nCRF_{Asym}$ = 0.022, Δ = -0.016, $p < 10^{-9}$; 4C-5/6: CRF_{Asym} = -0.033, $nCRF_{Asym}$ = -0.047, Δ

= -0.014, $p < 10^{-5}$; 4A/B-5/6: CRF_{Asym} = -0.018, $nCRF_{Asym}$ = -0.05, Δ = -0.032, $p < 10^{-20}$; Wilcoxon

rank-sum test). Thus, relative spike timing was earlier within FH_i laminar compartments during

nCRF stimulation than during CRF stimulation in the combined dataset, suggesting that the

proportion of information flow from FH_i layers to input layers increased during nCRF stimulation.

This observation was consistent across individual recording sessions (Table S2). Notably, we

observed a similar pattern when instead of asymmetry, we compared CCG peak lags

(Supplementary Figure S2A), and when we compared the total number of both significant and

nonsignificant pairs (CRF = 15,920; nCRF = 11,263) (Supplementary Figure S2B), and when we

compared CCGs with a more stringent significance criterion (10 SDs) in the combined dataset

(Supplementary Figure S2C-D).

Lastly, we asked whether interlaminar information flow was related specifically to border

ownership modulation, independent of the comparisons between CRF and nCRF stimulation.

Since CRF and nCRF stimuli differed in ways potentially unrelated to nonclassical modulation

(e.g., variations within the CRF), we sought to assess the extent to which information flow from

FH_i to input layers predicted the magnitude of nCRF modulation. To this end, we computed two

modulation indices from the same B_{own} test stimuli (Figure 2A), specifically the B_{own} modulation

index and the local contrast (LC) index. In contrast to the B_{own} modulation index, which reflects

nCRF modulation, the LC index measures selectivity to luminance contrast polarity (dark-light vs.

light-dark) within the CRF. We hypothesized that if the relative timing of spiking activity within

FH_i laminar compartments is indicative of nCRF modulation, we should expect earlier relative

spike timing in FH_i compartments, i.e., lower CCG asymmetries with respect to other

compartments, to be associated with greater B_{own}. In contrast, we should not expect that

relationship with the LC index.

Consistent with our hypothesis, we observed a negative relationship between the B_{own}

modulation index and CCG asymmetry across the population and within individual sessions (linear

regression: population slope (β) = -0.32, p < 10^{-27}; individual sessions: *pen1* = -0.52, p < 10^{-5}; *pen2*

= -0.06, p = 0.14; *pen3* = -0.25, p < 10^{-4}; *pen4* = -0.20, p = 0.008) (Figure 5A). This result suggests

that the magnitude of B_{own} modulation for neuronal pairs depended on the proportion of

information flow from FH_i compartment to input layers. In contrast, the opposite was true for the

LC index. That is, LC index was positively correlated with CCG asymmetry across the population

(linear regression: population slope (β) = 0.46, p<10^{-28}) (Figure 5B). However, this relationship

was not consistent across sessions (individual sessions: *pen1* = -0.30, p = 0.09; *pen2* = 0.33, p <

10^{-5}; *pen3* = 0.23, p = 0.006; *pen4* = 0.14, p = 0.04). To explore both results further, we built

generalized linear models (GLMs) to assess the dependence of B_{own} or LC modulation on CCG

asymmetry to incorporate variations across interlaminar combinations and recording sessions. For

LC modulation, this analysis indicated that the magnitude of LC modulation was not reliably

predicted by interlaminar information flow (GLM: coefficient = -0.04, p = 0.30; R^2 = 0.27) (Table

S3). In contrast, for B_{own} modulation, this analysis confirmed that the magnitude of modulation

was reliably predicted by interlaminar information flow, independent of variations across

interlaminar combinations and recording sessions (GLM: coefficient = -0.16 p = $1.1*10^{-6}$; R^2 =

0.11). These results demonstrate that interlaminar information flow was related specifically to the

magnitude of border ownership modulation.

Discussion

We measured the sensitivity of neurons to nCRF stimulation, specifically border ownership, within large, local populations in macaque V1. Consistent with previous studies on awake animals, we observed V1 neurons with border ownership selectivity even under anesthesia. We observed that neurons within single columns tended to prefer the same side of objects located outside of the CRF. In addition, we found that cross-correlations among pairs of neurons situated across FH_i and other laminar compartments differed between CRF and nCRF stimulation. Specifically, neurons within FH_i laminar compartments were more likely to lead pairwise interactions with neurons in other compartments during nCRF stimulation than during CRF stimulation. Moreover, the magnitude of border ownership modulation was predicted by greater information flow from FH_i laminar compartments, independent of the comparisons between CRF and nCRF stimulation. These results demonstrate that the flow of signals between neurons in FH_i and other laminar compartments depends on the degree to which those neurons integrate visual information from beyond the CRF. Below we discuss some potential limitations of the results as well as their implications for the mechanisms underlying contextual modulation.

The observation that neurons within FH_i laminar compartments can lead those in layer 4 during nCRF stimulation may appear surprising. However, several anatomical pathways could mediate the propagation of B_{own} signals from FH_i compartments to layer 4. In macaque V1, layers 5/6 send dense projections to 4A/B (Lund, 1988; Callaway, 1998). In particular, layer 6 pyramidal neurons, especially the subset classified as Type 1 cells, project substantially to layer 4C (Wiser and Callaway, 1996; Fitzpatrick et al., 1985). Projections from layers 2/3 to 4A/B have also been reported (Blasdel et al., 1985; Callaway and Wiser, 1996), and neurons in 4A/B often extend apical dendrites into layers 2/3 (Lund, 1988; Yoshioka et al., 1994). Although direct projections from

layers 2/3 to 4C are generally sparse (Callaway, 1998), a subset of neurons in the lower part of

layer 3 can give off collateral axons to 4C (Lund and Yoshioka, 1991). Additionally, some 4C

neurons extend dendrites into 4B, enabling potential dendritic integration of inputs from more

superficial layers (Somogyi and Cowey, 1981; Mates and Lund, 1983; Yabuta and Callaway,

1998). Moreover, layers 2/3 may influence 4C neurons disynaptically, without requiring dense

monosynaptic connections. Importantly, while CCGs can suggest possible circuit arrangements,

functional connectivity may arise through mechanisms not fully captured by traditional anatomical

tracing. Our observation of functional inputs from layers 2/3 to layer 4 aligns with prior findings

in rodent V1, where CCG analysis (Senzai et al., 2019) or photostimulation (Xu et al., 2016)

revealed similar pathways. Nonetheless, future studies will be needed to determine the extent of

input to layer 4 neurons from more precise FHi laminae.

Each of the laminar compartments designated in this study comprised more than a single

distinct anatomical layer. This was done to mitigate the uncertainty in determining exact laminar

boundaries, to achieve comparable numbers of neurons across compartments, and to maximize

statistical power. In addition, combining layers allowed us to combine feedback with horizontal

input layers. It is known that feedback and horizontal connections exhibit layer specificity within

superficial and deep layers, e.g., horizontal connections are most prominent in layer 2/3 and 5

(Rockland and Lund, 1983; Gilbert and Wiesel, 1983), where feedback connections target

primarily in layers 1, upper layer 2, and 5/6 (Rockland and Pandya, 1979; Federer et al., 2021;

Shmuel et al., 2005). Yet, they both largely avoid layer 4 (Angelucci et al., 2017). Nonetheless,

given that the number of anatomically distinct layers within macaque V1 well exceeds the four

designated laminar compartments used here, it is important to consider the extent to which

collapsing layers limits the conclusions one can draw from our observations. For example,

although layer 6 receives feedback input, it also receives feedforward input directly from the LGN

(Hubel and Wiesel, 1972; Hendrickson et al., 1978; Blasdel and Lund, 1983; Callaway, 1998). In

addition, layer 4A is generally considered an input layer whereas 4B is not (Callaway, 1998; Hubel

and Wiesel, 1972; Blasdel and Lund, 1983; Hendrickson et al., 1978). Thus, the designation of any

compartment as nominally input, or feedback was relative and not absolute. That is, superficial

(layers 2/3) and deep (layers 5/6) layers receive higher proportions of feedback/horizontal inputs

than the 4A/B and 4C compartments. Indeed, previous studies of primate visual cortex have used

a similar approach in comparing input and feedback/horizontal laminar compartments (e.g. Hansen

et al., 2012; Nandy et al., 2017; van Kerkoerle et al., 2017; Pettine et al., 2019). Thus, the observed

changes in pattern of cross-correlations between neurons situated across compartments with

different relative feedforward and feedback/horizontal input can be interpreted in light of those

input differences.

Another important consideration is that a key challenge of studying classical and

nonclassical RF effects simultaneously in large neuronal populations is the difficulty in achieving

alignment of all CRFs with the test stimuli. Although our probe penetrations were largely normal

to the surface and neurons showed minimal deviations in orientation selectivity across the cortical

depth, V1 neurons nonetheless exhibit noteworthy RF scatter (location/size) even within single

cortical columns (Tootell et al., 1988; Gur et al., 2005; Li et al., 2022). This means that the

positioning of stimuli relative to the CRF inevitably varied across simultaneously recorded neurons

(e.g., Figure 1C). Moreover, this means that although neurons were generally well driven by CRF

grating stimuli and exhibited clear orientation tuning, CRF stimulation was not guaranteed to fall

solely within the CRF of all neurons. Nonetheless, involvement of the surround, especially far

surround was considerably less extensive for CRF stimulation than for nCRF stimulation.

Furthermore, in the latter case, regardless of variability in the CRF location, stimuli falling within

the CRF were always matched across nCRF stimulus conditions, and the surrounds were also

identical within 8° of the CRF (Figure 2A, B).

The experiments described here were performed in anesthetized animals. Several other

types of contextual modulation have been demonstrated in anesthetized animals (Ramsden et al.,

2001; Hupe et al., 1998; Bair et al., 2003; Allman et al., 1985; Gilbert and Wiesel, 1990; Nothdurft

et al., 1999; Webb et al., 2003; Zarella and Ts'o, 2016; Muller et al., 2003; Henry et al., 2013;

Bijanzadeh et al., 2018), and can involve inter-cortical feedback (Nurminen et al., 2018).

Nonetheless, anesthesia is known to affect certain types of contextual responses in area V1. For

example, the "figure-ground segregation" effect, where neurons respond more strongly to a texture

in a "figure" region than to the same texture in a "ground" region, has been shown to be suppressed

under anesthesia (Lamme et al., 1998) and when figure stimuli are not perceived (Super et al.,

2001). However, there has been debate over whether this effect depends on attention and awareness

(Marcus and Van Essen, 2002; Jones et al., 2015; Poltoratski and Tong, 2020; Poort et al., 2012;

Huang et al., 2020). It is worth noting that in that case, RFs are centered on the figure regions, and

the modulation likely involves detecting feature discontinuities around boundaries, followed by

region-filling and background suppression. Each step could be influenced differently by behavioral

variables. Nevertheless, the center figure enhancement emerges late in the V1 response, ~55ms

after response onset (Poort et al., 2012). In contrast, border ownership involves RFs centered on

the boundaries, with modulation emerging much earlier at ~10-35ms after response onset

(Sugihara et al., 2011). Therefore, although both forms of modulation may involve

feedback/horizontal input, the underlying circuitry may differ. Our finding of border ownership

modulation in anesthetized animals aligns with previous studies showing that border ownership

modulation can act separately even without attention (Qiu et al., 2007; O'Herron and von der

Heydt, 2009) or before object shape recognition (Williford and von der Heydt, 2016; Ko and von

der Heydt, 2018), perhaps signifying a representation of "proto-objects" in the early visual cortex

through automatic, pre-attentive grouping mechanism (von der Heydt, 2015; von der Heydt, 2023;

Self et al., 2019).

Border ownership enables neurons with small CRFs in the early visual cortex to assign the

occluding border between image regions to a foreground object, which is crucial for natural scene

segmentation and object recognition (Nakayama et al., 1995). Human imaging studies have

demonstrated the existence of B_{own} in both lower (Fang et al., 2009) and higher visual areas

(Kourtzi and Kanwisher, 2001; Andrews et al., 2002), yet the underlying circuitry is still poorly

understood. Several computational models have been proposed to explain B_{own}, including

feedforward models (Walker et al., 1999; Sakai and Nishimura, 2006; Super et al., 2010),

horizontal models (Zhaoping, 2005; Kogo et al., 2010), and feedback models (Craft et al., 2007;

Jehee et al., 2007; Jeurissen et al., 2016). Yet cues for determining B_{own} often lie far from the CRF

and beyond the extents of geniculocortical and horizontal V1 connections (Angelucci et al., 2017).

Furthermore, conduction along horizontal fibers appears too slow (0.1-0.4 mm/ms) (Grinvald et

al., 1994; Bringuier et al., 1999; Girard et al., 2001) to account for the rapid (~10-35ms) emergence

of B_{own} and its independence from object size (Zhang and von der Heydt, 2010; Sugihara et al.,

2011). In contrast, feedback inputs can be conducted through fibers 10 times faster (20-60 mm/ms)

(Girard et al., 2001), even via monosynaptic connections (Siu et al., 2021), and contribute to

modulating the early responses of V1 neurons (Hupe et al., 2001; Bair et al., 2003). Thus, although

we combined feedback and horizontal recipient layers, the differences in interlaminar signal flow

we observed were likely driven predominantly by feedback inputs.

In primates, V1 receives inter-cortical feedback connections primarily from areas V2, V3,

V4, MT (Rockland et al., 1994; Markov et al., 2014). Thus, V1 B_{own} signals could be generated

via inputs from these areas (Zhou et al., 2000; Hesse and Tsao, 2016; Hesse and Tsao, 2023;

Franken and Reynolds, 2021; Zhu et al., 2020). Our results resonate with a recent study reporting

that in area V4, B_{own} emerges earliest in deep layers compared to input layers (Franken and

Reynolds, 2021). This suggests that the earliest component of B_{own} in V4 is not inherited from

feedforward inputs, i.e., from V1/V2, but is generated within deep layers de nova, or through

feedback connections that arrive at deep layers (Franken and Reynolds, 2021). This further

supports the view that B_{own} in V1 emerges via feedback from higher visual areas. In addition,

similar to what we observed in V1, columnar organization of B_{own} was observed in V4 (Franken

and Reynolds, 2021). Indeed, neurons with B_{own} are clustered into patches in a wide range of

primate visual areas, including area V2, V3, V3A, V4, V4A (Hesse and Tsao, 2023). Thus, the

evidence of B_{own} modularity we observed in V1 could arise from organized feedback from the B_{own}

modules in those areas. This is supported by recent anatomical evidence suggesting that feedback

terminals in V1 are clustered and functionally specific (Angelucci et al., 2002; Federer et al., 2021;

Shmuel et al., 2005; Siu et al., 2021).

Figure legends

Figure 1. Classical receptive field (CRF) stimulation in single V1 columns.

(A) Neuropixels probe (384 channels/3.84mm) penetrations made into the lateral surface or

underlying calcarine sulcus of macaque V1. Bottom: Diagram of designated laminar

compartments. (B) Current-Source density (CSD) profile from an example recording session (M1,

pen1), showing the early current sink (red) indicative of layer 4C. Laminar compartment

boundaries (dashed lines) were determined using histological data and the CSD profile. (C) Top:

71 well-defined classical receptive fields (CRF) vertically stacked along the cortical depth, mapped

with a 10 x 10 array (4 x 4 dva) of probe stimuli (gratings, 0.4 dva in diameter). Bottom: Drifting

Gabor grating (1.5 dva in diameter) used for CRF stimulation, positioned within the joint receptive

fields of recorded neurons. Zero (x, y) denotes the center of the CRF population. (D) Heatmap of

visual responses from 159 neurons to 36 drift directions of Gabor gratings, vertically stacked along

cortical depth (M1, *pen1*). Top: Distribution of differences in preferred orientation for all pairwise

combinations of neurons.

Figure 2. Nonclassical receptive field (nCRF) stimulation and border ownership.

(A) Border ownership test stimuli for nCRF stimulation. Stimuli consisted of uniformly white or

black objects (8º squares) on black or white backgrounds, respectively. One border of the object

fell within the CRFs (red circle) of the recorded neurons, with orientation varying in 45º steps.

Within the CRF, depending on local contrast polarity (LC), the border could either be a dark-light

edge (LC1, Left column) or a light-dark edge (LC2, Right column). For the same LC condition,

the border within the CRF could be the bottom-left edge (Side 1, top row) or top-right edge (Side

2, bottom row) of the object. (B) Enlarged view of stimulus configuration within the CRF, showing

identical stimulus regions between Side 1 and Side 2 for each LC condition (8 x 16 dva). (C) Four

example B_{own} neurons recorded from V1. For each neuron, dot raster plots (top) and instantaneous

firing rates (bottom) show responses to the four stimulus conditions depicted in (A), respectively.

Magenta represents Side 1; green represents Side 2. (D) Bar plots showing the mean spike rates of

the four neurons across the four stimulus conditions. (E) Top: Cartoon depicting the preferred

border orientation for each of the four recording sessions (*pen*1-4). Arrows indicate the preferred

side of objects relative to the border in the CRF. Bottom: B_{own} modulation index for each neuron

plotted at its recorded cortical depth across the recordings. Color bars on the ordinate represent

laminar compartments. Black dots denote statistically significant B_{own} neurons, determined by

ANOVA with object side and local contrast as factors ($p < 0.05$). Middle: Marginal distribution of

the B_{own} modulation index, red arrows indicating the population median. $*p < 0.05$; $***p < 10^{-5}$.

Figure 3. Interlaminar cross-correlations during CRF and nCRF stimulation.

(A) Example recording (M1, *pen1*) showing 159 visually responsive neurons (circles) recorded

simultaneously during CRF stimulation, plotted at their relative cortical depth. The abscissa is

magnified for visualization. Also shown are eight example neuronal pairs with significant CCGs

during CRF stimulation. Neuronal pairs with zero-time lag CCGs are connected by dashed lines,

while pairs with nonzero-time lag CCGs are connected by arrowed lines, with the arrow pointing

to the lagging neuron. (B) Eight corresponding CCGs from (A). Neuronal pairs are separated by

their laminar compartment combinations. The reference neuron (1st neuron in the CCG function)

is designated as j, and the target neuron as k. PL (peak lag, ms) is defined as the relative time delay

in peak correlation between the two spike trains. (C) Diagram illustrating feedforward and

feedback (and horizontal) projections within the designated laminar compartments. Feedforward

inputs arrive within laminar compartments 4C, while feedback (and horizontal) projections

principally target superficial and deep layers. (D) Heatmap of all significant CCGs during either

CRF stimulation (Gabor gratings, top row) or nCRF stimulation (border ownership, middle row),

for the four key interlaminar combinations (columns), calculated from one example recording (M1,

pen1). Individual CCGs are normalized by their absolute maximum for visualization, and different

CCGs are vertically stacked. Bottom row: Distribution of CCG asymmetry for each of the four

key interlaminar compartments as in (A), across CRF stimulation (gray) and nCRF stimulation

(magenta). Arrows indicate the population median. $**p < 0.002$; $***p < 10^{-5}$.

Figure 4. Comparison of interlaminar CCG asymmetries during CRF and nCRF

stimulation.

Combined dataset for each of the four key interlaminar combinations. Histograms and cumulative

distributions of asymmetry for all significant CCGs during CRF (gray) and nCRF (magenta)

stimulation. Arrows indicate the population median. $***p < 10^{-5}$.

Figure 5. Dependence of nCRF modulation on interlaminar signal flow.

(A) Scatter plot of the B_{own} modulation index versus CCG asymmetry for all significant CCGs,

aggregated across the four key interlaminar combinations and all recording sessions. CCG

asymmetries were calculated during nCRF stimulation. The geometric mean of the B_{own} index for

each neuronal pair was used. Red lines denote the fitted regression line and the 95% confidence

interval in linear regression. (B) Similar to (A), but using the local contrast (LC) index instead of

the B_{own} index. The LC index was calculated as the difference in neuronal responses to borders

with opposite contrast polarities, normalized by the maximum response.

Supplementary figure legends and tables

Supplementary Figure S1. Histological data for Neuropixels recordings in macaque V1.

(A) Neuropixels probe (384 channels/3.84mm) penetrations made into the lateral surface or underlying calcarine sulcus of macaque V1. Bottom: Diagram of designated laminar compartments. (B) Histological reconstruction of probe tracks based on DiI derivative staining for two example recordings (*pen3-4*). Here, Neuropixels probes passed through the lateral surface and reached the calcarine sulcus of V1. Only neurons in the calcarine V1 from these two recordings were tested and included in the analysis. Designated boundaries are shown. Scale bar, 500 µm.

Supplementary Figure S2. Comparison of interlaminar cross-correlations during CRF and nCRF stimulation under different CCG criteria.

(A) Histogram distributions of CCG peak lags for each of the four key interlaminar combinations in the combined dataset during CRF (gray) and nCRF (magenta) stimulation. The shift (Δ) in median CCG peak lags during nCRF stimulation relative to CRF stimulation is shown. Significant CCGs are defined as those with peaks exceeding 7 SD above the noise level (flank). (B) Similar to (A), but comparing CCG asymmetry across all CCGs, including both significant and non-significant ones. (C) Comparison of CCG peak lags using a stricter criterion, where significant CCGs are defined as those with peaks exceeding 10 SD above the noise level (flank). (D) Comparison of CCG asymmetry using the stricter criterion of 10 SD.

Table S1. Number of neurons selective to B_{own} (LC) across recordings.

Layer	Total	pSide <0.05		pSide ≥0.05		Layer	Total	pSide <0.05		pSide ≥0.05	
		p_{LC} ≥0.05	p_{LC} <0.05	p_{LC} <0.05	p_{LC} ≥0.05			p_{LC} ≥0.05	p_{LC} <0.05	p_{LC} <0.05	p_{LC} ≥0.05
		pen1						*pen2*			
2/3	29	6	5	6	12	2/3	55	1	14	18	22
4A/B	19	0	5	8	6	4A/B	34	0	7	10	17
4C	38	3	7	16	12	4C	54	4	15	22	13
5/6	40	2	7	18	13	5/6	27	5	2	4	16
		pen3						*pen4*			
2/3	15	1	1	4	9	2/3	0	0	0	0	0
4A/B	27	0	4	9	14	4A/B	34	3	0	2	29
4C	60	1	12	24	23	4C	60	2	4	5	49
5/6	40	2	7	2	29	5/6	37	3	0	2	32
		pen all									
2/3	99	8	20	28	43						
4A/B	114	3	16	29	66						
4C	212	10	38	67	97						
5/6	144	12	16	26	90						

Table S2. Comparison of CCG asymmetries during CRF and nCRF for each recording.

4C-2/3	CRF	nCRF	Δ	p	4A/B-2/3	CRF	nCRF	Δ	p
pen1	0.036	-0.004	-0.040	$2.8*10^{-7}$	*pen1*	0.046	-0.005	-0.050	$7.5*10^{-12}$
pen2	0.037	0.013	-0.024	$3.7*10^{-9}$	*pen2*	0.030	0.022	-0.008	0.066
pen3	0.017	0.033	0.016	0.29	*pen3*	0.049	0.052	0.003	0.99
4C-5/6	CRF	nCRF	Δ	p	4A/B-5/6	CRF	nCRF	Δ	p
pen1	-0.010	-0.044	-0.033	0.0015	*pen1*	0.012	-0.028	-0.040	$2.1*10^{-6}$
pen2	-0.009	-0.027	-0.018	$1.5*10^{-5}$	*pen2*	-0.011	-0.018	-0.007	0.059
pen3	-0.036	-0.062	-0.026	$5.8*10^{-7}$	*pen3*	-0.025	-0.087	-0.063	$9.1*10^{-15}$
pen4	-0.041	-0.093	-0.051	$3.1*10^{-8}$	*pen4*	-0.030	-0.089	-0.060	$9.1*10^{-12}$

Table S3. GLMs for dependence of border ownership and local contrast on CCG asymmetry.

$$B_{own}(pair) \sim 1 + CCG\ asymmetry + laminar\ combination + pen$$

$$LC(pair) \sim 1 + CCG\ asymmetry + laminar\ combination + pen$$

GLM	Border ownership (B_{own})				Local contrast (LC)			
	Estimate	SE	tStat	p	Estimate	SE	tStat	p
Intercept	0.166	0.008	21.379	0	0.362	0.010	35.352	0

CCG asymmetry	**-0.158**	0.032	-4.880	**$1.1*10^{-6}$**	**-0.044**	0.043	-1.037	**0.300**
4C-5/6 *rel.4C-2/3*	0.014	0.007	1.948	0.05	-0.117	0.009	-12.606	$3.5*10^{-35}$
4A/B-2/3 *rel.4C-2/3*	$2.5*10^{-5}$	0.006	0.004	0.99	-0.016	0.008	-1.877	0.061
4A/B-5/6 *rel.4C-2/3*	0.009	0.007	1.351	0.18	-0.136	0.009	-15.093	0
pen2$_{rel.pen1}$	-0.009	0.007	-1.219	0.22	0.010	0.010	0.980	0.327
pen3 $_{rel.pen1}$	0.038	0.009	4.435	$9.7*10^{-6}$	0.035	0.011	3.082	0.002
pen4 $_{rel.pen1}$	0.049	0.010	5.147	$2.9*10^{-7}$	-0.088	0.013	-6.910	$6.4*10^{-12}$

577   Number of observations: 2118; Error degrees of freedom: 2110

578   GLM(B_{own}): R^2=0.11; F-statistic vs. constant model: 37.2, p-value=0.

579   GLM(LC): R^2=0.27; F-statistic vs. constant model: 109, p-value=0.

Methods

Experimental Model and Subject Details

Anesthetized recordings were conducted in two adult male rhesus macaques (*Macaca mulatta*, M1, 13 kg; M2, 8 kg). All experimental procedures were in accordance with the National Institutes of Health Guide for the Care and Use of Laboratory Animals, the Society for Neuroscience Guidelines and Policies, and with approved Institutional Animal Care and Use Committee (IACUC) protocol (#APLAC-9900) of Stanford University.

Electrophysiological Recordings

Prior to each recording session, treatment with dexamethasone phosphate (2 mg per 24 h) was instituted 24 h to reduce cerebral edema. After administration of ketamine HCl (10 mg per kilogram body weight, intramuscularly), monkeys were ventilated with 1-2% isoflurane in a 1:1 mixture of N_2O and O_2 to maintain general anesthesia. Electrocardiogram, respiratory rate, body temperature, blood oxygenation, end-tidal CO_2, urine output and inspired/expired concentrations of anesthetic gasses were monitored continuously. Normal saline was given intravenously at a variable rate to maintain adequate urine output. After a cycloplegic agent (atropine sulfate, 1%) was administered, the eyes were focused with contact lenses on an LCD monitor. Vecuronium bromide (60 μg/kg/h) was infused to prevent eye movements.

With the anesthetized monkey in the stereotaxic frame, an occipital craniotomy was performed over the opercular surface of V1. The dura was reflected to expose a small (~3 mm^2) patch of cortex. Next, a region relatively devoid of large surface vessels was selected for implantation, and the Neuropixels probe was inserted with the aid of a surgical microscope. Given the width of the probe (70 µm x 20 µm), insertion of it into the cortex sometimes required multiple

attempts if it flexed upon contacting the pia. The junction of the probe tip and the pia could be

visualized via the (Zeiss) surgical scope, and the relaxation of pia dimpling was used to indicate

penetration, after which the probe was lowered at least 3-4 mm. Prior to probe insertion, it was

dipped in a solution of the DiI derivative FM1-43FX (Molecular Probes, Inc) for subsequent

histological visualization of the electrode track.

Given the length of the probe (1 cm), and the complete distribution of electrode contacts

throughout its length, recordings could be made either in the opercular surface cortex (M1, *pen* 1-

2) or within the underlying calcarine sulcus (M2, *pen* 3-4), by selecting a subset of contiguous

active contacts (n = 384) from the total number (n=986). Recordings were made at 1 to 2 sites in

one hemisphere of each monkey. At the end of the experiment, monkeys were euthanized with

pentobarbital (150 mg/kg) and perfused with normal saline followed by 1 liter of 1% (wt/vol)

paraformaldehyde in 0.1 M phosphate buffer, pH 7.4.

Visual Stimulation

Visual stimuli were presented at manually mapped receptive field (RF) locations for each

recording session on an LCD monitor (Model NEC-4010, dimensions: 88.5 cm (H) x 49.7 cm (V),

resolution: 1360 x 768 pixels, frame rate: 60 Hz) positioned 114 cm from the monkeys. RF

eccentricities were ~4-6° (M1) and ~6-10° (M2). Visual stimuli were generated using customized

MATLAB scripts with the Psychophysics Toolbox extensions (version PTB-3) (Kleiner et al.,

2007). A photodiode was used to measure stimulus timing.

Classical receptive field (CRF) stimuli consisted of drifting Gabor gratings (2 deg./sec.,

100% Michelson contrast) with a diameter of 1.5 degrees of visual angle (dva), positioned within

the joint receptive fields (RFs) of recorded neurons. This size was selected to largely constrain the

stimuli within the CRFs of recorded neurons, typically ~0.5-1 dva, as determined manually online.

Gratings drifted in 36 different directions, from 0 to 360° in 10° steps, in a pseudorandom order.

Each stimulus condition was presented for 1 s and repeated 5 or 10 times. A blank screen with

equal luminance to the Gabor patch was presented for 0.25 s during the stimulus interval. Stimuli

were presented either monocularly (*pen* 1, 3) or both monocularly and binocularly (*pen* 2, 4). Four

spatial frequencies (0.5, 1, 2, 4 cycles/deg.) were tested. The optimal eye and spatial frequency

conditions were determined offline for further analysis.

Nonclassical receptive field (nCRF) stimuli used in this study, i.e., Border ownership test

stimuli consisted of uniformly white or black objects (square, 8 x 8 dva) presented on a black or

white background, respectively, like previous studies (Zhou et al., 2000). One border of the object

fell within the CRFs of the recorded neurons at an orientation that varied in 45º steps. Objects were

either 4 x 4 dva or 8 x 8 dva in size, and only conditions with 8 x 8 dva were selected for quantifying

border ownership selectivity to tolerate the variation in the CRF locations. Stimuli were presented

monocularly to the preferred eye, to avoid the surround stimulating the CRF of the non-preferred

eye. For each of the 4 orientation conditions, there were 4 basic conditions (Figure 2A). Within

the CRF, the oriented border could either be a dark-light edge or light-dark edge, resulting in two

different local contrast (LC) polarity conditions, namely, LC1 and LC2. For the same LC

condition, the stimulus configuration outside of the CRF differed such that the oriented border

could belong opposite sides of objects, resulting in two different sides of object conditions, namely,

side 1 and side 2. Each stimulus condition was presented for 1 s and repeated 10 or 20 times. A

blank screen with equal luminance to the mean of object and background luminance was presented

for 0.5s during the stimulus interval.

Receptive field mapping

In some sessions, the receptive fields of V1 neurons were tested systemically for offline use. CRFs

were mapped from a 10 x 10 array (4 x 4 dva) using probe stimuli. The array was centered at the

manually determined RFs location. The stimuli consisted of circular sine wave gratings of 0.4 dva

in diameter, and 100% Michelson contrast. Gratings were presented at the optimal orientation

determined manually online for each recording session and drifted at speed of 2 deg./sec. Each

stimulus condition was presented for 0.2s, interleaved with 0.3s of blank screen with equal

luminance, and repeated 15 times.

Mean spike counts from 0.05 to 0.25s after each stimulus onset were calculated, resulting

in a 10 x 10 matrix containing mean spike rates to each grid location. This 2D matrix was then

interpolated to a 40 x 40 matrix (MATLAB function 'interp2') and then smoothed with a 2D

Gaussian filter (MATLAB function 'imgaussfilt', σ=2). The outline of the CRF was defined as the

contour isoline at 80% of the maximum response using MATLAB function 'contour3'. Neurons

with only 1 contour region at the threshold were selected.

Data acquisition and spike sorting

Raw spike-band data were sampled and recorded at 30kHz. They were then median-

subtracted and high-pass filtered at 300 Hz during the pre-processing stage. Spike sorting was

carried out with Kilosort 3 (Pachitariu et al., 2024) to find spike times and assign each spike to

different templates (neurons). Default parameters in Kilosort 3 were used for spike sorting,

specifically, Ops.th = [9,9]; Ops.lam = 20; Ops.AUCsplit = 0.8; Ops.ThPre = 8; Ops.sig = 20;

Ops.nblocks = 5; Ops.spkTh = -6. The raw sorted data were then manually curated in Phy

(*https://github.com/cortex-lab/phy*) to remove templates with very few spikes or with atypical

waveforms and to perform minimal templates merging and splitting. Double-counted spikes from

the algorithm fitting the residuals to a new template were identified using established criteria

(Siegle et al., 2021), by counting spikes from two templates separated by less than 50μm (~5

channels) and with spike activity occurring within 5 samples (0.167ms). Double-counted spikes

from the template smaller in amplitude were removed from further analysis. Visual responsiveness

of each unit was assessed under CRF, nCRF, or RF mapping stimulation. For each type of

stimulation, a paired-sample t test was performed to determine whether the mean spike counts after

each stimulus presentation exceeded the preceding blank period at a significance level of 0.01.

Only neurons showing significant visual responsiveness were included for further analysis.

Layer Assignment

The laminar locations of our recorded neurons were estimated based on a combination of

functional analysis and histology results (Zhu et al., 2024; Carr et al., 2025). For each recording,

we first performed the current source density (CSD) analysis on the stimulus-triggered average of

local field potentials (LFP). LFP were low-pass filtered at 200 Hz and recorded at 2500 Hz. LFP

signals recorded from every 4 nearby channels were averaged and realigned to the onset of visual

stimulus. CSD was estimated as the second-order derivatives of signals along the probe axis using

the common five-point formula (Nicholson and Freeman, 1975). The result was then smoothed

across space ($\sigma = 120$ μm) to reduce the artifacts caused by varied electrode impedance. We located

the lower boundary of the major sink (the reversal point of sink and source) as the border between

layer 4C and layer 5/6. We also considered anatomical data in order to localize recorded neurons

within gray matter, allowing for minor adjustments (±1 group channel) in layer boundary

placement. Subsequent layer boundaries were determined by offsetting the cortical thickness

derived from histological slices (Supplementary Fig S1).

Single neuron properties during CRF stimulation

To assess the orientation tuning for each neuron, the spike rates during 0.1 to 1s after each stimulus

onset were calculated and averaged for each orientation condition (N= 18). The orientation tuning

responses were first smoothed with a Hanning window (half-width at half-height of $20°$), and then

fitted with a von Mises function (Swindale, 1998)

701
$$y = a_0 + a_1 * e^{a_2*(\cos(2*x - 2*a_3) - 1)}$$

The location of the peak of the fitted curve was determined as the preferred orientation. Only

neurons well fit by the function ($R^2 > 0.7$) were included for determining pairwise differences in

preferences for all combinations in each recording session.

Modulation index for border ownership

As in previous studies (Zhou et al., 2000), we calculated a B_{own} modulation index to quantify the

selectivity of each neuron for B_{own}. The B_{own} modulation index is defined as

709
$$B_{own}\ index = \frac{0.5 * (R_{LC1} + R_{LC2})_{Side1} - 0.5 * (R_{LC1} + R_{LC2})_{Side2}}{R_{max}}$$

Here, $R_{LC,\ Side}$ represents the average spike rate from 0.05 to 0.5 s after stimulus onset for one of

four basic conditions, which are derived from the combination of two different *Local Contrast*

conditions (LC1 and LC2) and two *Side* conditions (Side 1 and Side 2). Baseline activity was not

subtracted. Since the border within the CRF can vary in 45º steps from 0º to 180º, we only selected

the optimal orientation for each recording population to calculate the B_{own} index. R_{max} was defined

as the maximum response for each neuron across the four basic conditions and four orientations.

We used R_{max} instead of summing responses across the four basic conditions to account for the

slight variations in orientation preferences among neurons within each recording session. The

statistical significance of B_{own} was evaluated using a two-factor ANOVA (MATLAB function

'anovan') on the average spike rates for each condition. The two factors were object *Side* (with

two levels: Side 1 and Side 2) and *Local Contrast* (with two levels: LC 1 and LC2), including their

interactions. A significance level of 0.05 was used. Neurons were considered to exhibit significant

B_{own} modulation if they showed a significant main effect for the object *Side* factor. Whether they

were invariant to local contrast was determined by the main effect of the *Local Contrast* factor.

Cross-correlograms (CCGs)

To measure correlated firing, we computed the cross-correlations between spike trains of

all pairs of simultaneously recorded neurons. For each recording session, we calculated cross-

correlograms (CCGs) from pairwise combinations of neuronal spike trains during either CRF

(Gabor gratings) or nCRF (border ownership) stimulation conditions, respectively. We only

included neuronal pairs with geometric mean firing rates more than 0.5 spikes/sec for further

analysis. For both conditions, we focused on the spiking activity within the 0.25 -1s window during

each stimulus presentation, to mitigate the influence of the transient visual response after stimulus

onset. The CCG function for a given pair of neurons was defined as follows:

734
$$CCG(\tau)_{j-k} = \frac{\frac{1}{M}\sum_{i=1}^{M}\sum_{t=1}^{N-\tau} x_j^i(t) \times x_k^i(t+\tau)}{\theta(\tau)\sqrt{\lambda_j \lambda_k}}$$

where j denotes the first/reference neuron and k denotes the second/target neuron in the CCG

function. τ is the time lag between the two spike trains x_j^i and x_k^i from neuron j and k, during trial

i. The value of $x_j^i(t)$ and $x_k^i(t)$ is 1 if there is a spike at time bin t and is zero otherwise. M is the

total number of trials; N is the number of time bins within a trial. $\theta(\tau) = N - |\tau|$, is a triangular

function that corrects for the degree of overlap in the two spike trains at each time lag. λ_j and λ_k

are the mean firing rates of neuron j and k computed over the same bins used to compute the CCG

at each time lag. The normalization by the geometric mean of spike rates is commonly used by

previous studies to account for the influence of firing rates on the CCG peaks (Bair et al., 2001;

Kohn and Smith, 2005).

To correct for correlations due to stimulus-locking or slow fluctuations in the population

responses (e.g., gamma-band activity), we computed a jitter-corrected CCG by subtracting a

jittered CCG from the original CCG:

$$CCG_{corrected} = CCG_{original} - CCG_{jittered}$$

The jittered CCG ($CCG_{jittered}$) reflects the expected value of CCG computed from all possible jitters

of each spike train within a given jitter window (Harrison and Geman, 2009; Smith and Kohn,

2008). The jittered spike train preserves both the PSTH of the original spike train across trials and

the spike counts in the jitter window within each trial. As a result, jitter correction removes the

correlation between PSTHs (stimulus-locking) and correlation on timescales longer than the jitter

window (slow population correlations). Here, a 25-ms jitter window was chosen based on previous

studies (Jia et al., 2013; Siegle et al., 2021). Jitter-corrected CCG ($CCG_{corrected}$) was then smoothed

with a 5ms kernel [0.05, 0.25, 0.4, 0.25, 0.05] (Kohn and Smith, 2005; Smith and Kohn, 2008)

before further analysis.

We classified a CCG as significant if the peak of the jitter-corrected CCG occurred within

10 ms of zero and exceeded 7 standard deviations (SD) above the mean of the noise distribution,

similar to previous studies (Siegle et al., 2021). The noise distribution for a CCG was defined as

the flanks of the jitter-corrected CCG ($\{CCG(\tau)|100 \geq |\tau| \geq 50\ ms\}$). All analyses presented in

the main text involve only significant, jitter-corrected CCGs. In the supplementary material, we

also applied a more stringent criterion, requiring that significant CCG peaks exceed 10 SD above

the noise, and a less stringent criterion that includes both significant and non-significant CCGs.

These ensure that our results are robust across the choices of threshold (Supplementary Figure S2).

To assess the directionality of functional interactions or signal flow between pairs of

neurons, we focused on two metrics derived from the CCG functions. One is the peak lag (PL),

which describes the time lag of peak correlation between the spike trains of neuron k (target)

relative to neuron j (reference). PL is positive if the reference neuron j leads target neuron k in

spike timing, negative if j lags k, and zero if j and k fire synchronously. The second metric we used

to assess the signal flow is CCG asymmetry, computed by subtracting the sum of CCG values

during the [-13, 0] ms time window from the sum of CCG values during the [0, 13] ms. Similar to

PL, a positive CCG asymmetry indicates that reference neuron j leads k, while a negative

asymmetry indicates that j lags k. The 13ms window was chosen as half of the 25ms jitter window,

consistent with previous studies (Jia et al., 2022). Compared to peak lag, CCG asymmetry

additionally captures the strength of the functional interactions between pairs of neurons and is

less dependent on the exact shape of the CCG peak, which is often more complex in corticocortical

functional interactions (Alonso and Martinez, 1998).

Relationship between B$_{own}$ modulation and interlaminar signal flow

To determine whether the magnitude of border ownership or local contrast modulation is

predicted by interlaminar signal flow during nCRF stimulation, we first performed linear

regression (MATLAB, 'fitlm', ordinary least squares) and then built generalized linear models

(GLMs) using predictors including CCG asymmetry, interlaminar combinations, and recording

sessions. We focused on the four key interlaminar combinations (4C-2/3, 4A/B-2/3, 4C-5/6, 4A/B-

5/6), where the latter laminar compartments receive feedback/horizontal inputs (FH_i). Only

neuronal pairs with significant CCGs were included as individual samples in the GLM. The

resulting GLM equations were:

$$B_{own}(pair) \sim 1 + asymmetry + \ laminar\ combination + pen$$

$$LC(pair) \sim 1 + asymmetry + \ laminar\ combination + pen$$

B_{own} (pair) and LC (pair) were calculated as the geometric mean of the B_{own} and LC indices for the

constituent neurons in each pair. Specifically, $B_{own}(pair) = \sqrt{abs(B_{own}(j)) * abs(B_{own}(k))}$,

where $abs(B_{own}(j))$ and $abs(B_{own}(k))$ represent the absolute B_{own} modulation indices for constituent

neurons j and k. A similar calculation was performed for LC (pair). CCG asymmetries were

calculated from the nCRF stimulation condition where neurons in the input layers were selected

as reference neurons and the ones in FH_i compartments were target neurons in the CCG. Additional

predictors, including interlaminar combinations and recording sessions, were included in the

models to incorporate variations when pooling the dataset.

Statistical tests

B_{own} for each neuron was assessed with ANOVA. The proportion of B_{own} neurons within

different layers was assessed using chi-squared test of homogeneity. The sign of populational

border ownership preferences from individual recording session was tested with sign test. The sign

of populational CCG peak lag for a given laminar compartment combination was assessed with

sign test. The comparisons of peak lag and CCG asymmetry between CRF and nCRF stimulation

condition for each laminar compartment combination were evaluated using Wilcoxon rank-sum

tests. The dependence between CCG asymmetry and B_{own} for pair of neurons was assessed using

linear regression and GLM.

Acknowledgment

We thank Jonathan C. Horton for extensive help with the recordings and histology, Danielle A. Lopes, Stephen Cital, Shellie Hyde and Sam Baker for technical assistance, Tim Harris and Karel Svoboda for providing the Neuropixels probes. This work was supported by NIH Grant EY029759, Brain and Behavior Research Foundation Grant to X.M.; NIH Grants EY014924, NS116623, and a Ben Barres Professorship to T.M.

Author Contributions

Conceptualization, S.Z., X.C. and T.M.; Investigation, S.Z., X.C. and T.M.; Software, S.Z. and E.T.; Formal Analysis, S.Z. and Y.J.O.; Visualization, S.Z., Y.J.O.; Writing – Original Draft, S.Z. and T.M.; Writing – Review and Editing, S.Z., Y.J.O., E.T., X.C., T.M.; Funding Acquisition, X.C. and T.M.

Declaration of Interests

The authors declare no competing interests.

Data and Code Availability

All raw data generated in this study will be deposited in a public repository upon acceptance and made publicly available prior to publication.

The code developed for data analysis has been deposited to GitHub (https://github.com/szhu-007/Zhu_V1_interlaminar-signal-flow) and is freely available for access.

References

ADESNIK, H., BRUNS, W., TANIGUCHI, H., HUANG, Z. J. & SCANZIANI, M. 2012. A neural circuit for spatial summation in visual cortex. *Nature,* 490**,** 226-31.

AERTSEN, A. M. & GERSTEIN, G. L. 1985. Evaluation of neuronal connectivity: sensitivity of cross-correlation. *Brain Res,* 340**,** 341-54.

ALBRIGHT, T. D. & STONER, G. R. 2002. Contextual influences on visual processing. *Annu Rev Neurosci,* 25**,** 339-79.

ALLMAN, J., MIEZIN, F. & MCGUINNESS, E. 1985. Stimulus specific responses from beyond the classical receptive field: neurophysiological mechanisms for local-global comparisons in visual neurons. *Annu Rev Neurosci,* 8**,** 407-30.

ALONSO, J. & CHEN, Y. 2009. Receptive field. *Scholarpedia,* 4**,** 5393.

ALONSO, J. M. & MARTINEZ, L. M. 1998. Functional connectivity between simple cells and complex cells in cat striate cortex. *Nature Neuroscience,* 1**,** 395-403.

ALONSO, J. M., USREY, W. M. & REID, R. C. 1996. Precisely correlated firing in cells of the lateral geniculate nucleus. *Nature,* 383**,** 815-9.

ANDERSON, J. C. & MARTIN, K. A. 2009. The synaptic connections between cortical areas V1 and V2 in macaque monkey. *J Neurosci,* 29**,** 11283-93.

ANDREWS, T. J., SCHLUPPECK, D., HOMFRAY, D., MATTHEWS, P. & BLAKEMORE, C. 2002. Activity in the fusiform gyrus predicts conscious perception of Rubin's vase-face illusion. *Neuroimage,* 17**,** 890-901.

ANGELUCCI, A., BIJANZADEH, M., NURMINEN, L., FEDERER, F., MERLIN, S. & BRESSLOFF, P. C. 2017. Circuits and Mechanisms for Surround Modulation in Visual Cortex. *Annu Rev Neurosci,* 40**,** 425-451.

ANGELUCCI, A. & BRESSLOFF, P. C. 2006. Contribution of feedforward, lateral and feedback connections to the classical receptive field center and extra-classical receptive field surround of primate V1 neurons. *Prog Brain Res,* 154**,** 93-120.

ANGELUCCI, A., LEVITT, J. B., WALTON, E. J., HUPE, J. M., BULLIER, J. & LUND, J. S. 2002. Circuits for local and global signal integration in primary visual cortex. *J Neurosci,* 22**,** 8633-46.

BAIR, W., CAVANAUGH, J. R. & MOVSHON, J. A. 2003. Time course and time-distance relationships for surround suppression in macaque V1 neurons. *J Neurosci,* 23**,** 7690-701.

BAIR, W., ZOHARY, E. & NEWSOME, W. T. 2001. Correlated firing in macaque visual area MT: time scales and relationship to behavior. *J Neurosci,* 21**,** 1676-97.

BIJANZADEH, M., NURMINEN, L., MERLIN, S., CLARK, A. M. & ANGELUCCI, A. 2018. Distinct Laminar Processing of Local and Global Context in Primate Primary Visual Cortex. *Neuron,* 100**,** 259-274 e4.

BLASDEL, G. G. & LUND, J. S. 1983. Termination of afferent axons in macaque striate cortex. *J Neurosci,* 3**,** 1389-413.

BLASDEL, G. G., LUND, J. S. & FITZPATRICK, D. 1985. Intrinsic connections of macaque striate cortex: axonal projections of cells outside lamina 4C. *J Neurosci,* 5**,** 3350-69.

BRIGGS, F., MANGUN, G. R. & USREY, W. M. 2013. Attention enhances synaptic efficacy and the signal-to-noise ratio in neural circuits. *Nature,* 499**,** 476-80.

BRINGUIER, V., CHAVANE, F., GLAESER, L. & FREGNAC, Y. 1999. Horizontal propagation of visual activity in the synaptic integration field of area 17 neurons. *Science,* 283**,** 695-9.

BULLIER, J., HUPE, J. M., JAMES, A. C. & GIRARD, P. 2001. The role of feedback connections in shaping the responses of visual cortical neurons. *Prog Brain Res,* 134**,** 193-204.

CALLAWAY, E. M. 1998. Local circuits in primary visual cortex of the macaque monkey. *Annu Rev Neurosci,* 21**,** 47-74.

CALLAWAY, E. M. 2004. Feedforward, feedback and inhibitory connections in primate visual cortex. *Neural Netw,* 17**,** 625-32.

CALLAWAY, E. M. & WISER, A. K. 1996. Contributions of individual layer 2-5 spiny neurons to
local circuits in macaque primary visual cortex. *Vis Neurosci,* 13, 907-22.
CARR, N., ZHU, S., CHEN, X., LEE, K., PERLISS, A., MOORE, T. & CHANDRASEKARAN, C.
2025. Neuropixels reveal structure-function relationships in monkey V1 in vivo. *bioRxiv*.
CHEN, M., YAN, Y., GONG, X., GILBERT, C. D., LIANG, H. & LI, W. 2014. Incremental integration
of global contours through interplay between visual cortical areas. *Neuron,* 82, 682-94.
CHEN, R., WANG, F., LIANG, H. & LI, W. 2017. Synergistic Processing of Visual Contours across
Cortical Layers in V1 and V2. *Neuron,* 96, 1388-1402 e4.
CHISUM, H. J., MOOSER, F. & FITZPATRICK, D. 2003. Emergent properties of layer 2/3 neurons
reflect the collinear arrangement of horizontal connections in tree shrew visual cortex. *J*
Neurosci, 23, 2947-60.
CHU, C. C., CHIEN, P. F. & HUNG, C. P. 2014. Tuning dissimilarity explains short distance
decline of spontaneous spike correlation in macaque V1. *Vision Res,* 96, 113-32.
CRAFT, E., SCHÜTZE, H., NIEBUR, E. & VON DER HEYDT, R. 2007. A neural model of figure-
ground organization. *J. Neurophysiol,* 97, 4310-4326.
DENMAN, D. J. & CONTRERAS, D. 2014. The structure of pairwise correlation in mouse primary
visual cortex reveals functional organization in the absence of an orientation map. *Cereb*
Cortex, 24, 2707-20.
EGUCHI, A. & STRINGER, S. M. 2016. Neural network model develops border ownership
representation through visually guided learning. *Neurobiol Learn Mem,* 136, 147-165.
FANG, F., BOYACI, H. & KERSTEN, D. 2009. Border ownership selectivity in human early visual
cortex and its modulation by attention. *J Neurosci,* 29, 460-5.
FEDERER, F., TA'AFUA, S., MERLIN, S., HASSANPOUR, M. S. & ANGELUCCI, A. 2021.
Stream-specific feedback inputs to the primate primary visual cortex. *Nat Commun,* 12,
228.
FITZPATRICK, D. 2000. Seeing beyond the receptive field in primary visual cortex. *Curr Opin*
Neurobiol, 10, 438-43.
FITZPATRICK, D., LUND, J. S. & BLASDEL, G. G. 1985. Intrinsic connections of macaque striate
cortex: afferent and efferent connections of lamina 4C. *J Neurosci,* 5, 3329-49.
FRANKEN, T. P. & REYNOLDS, J. H. 2021. Columnar processing of border ownership in primate
visual cortex. *Elife,* 10.
GIESELMANN, M. A. & THIELE, A. 2022. Stimulus dependence of directed information exchange
between cortical layers in macaque V1. *Elife,* 11.
GILBERT, C. D. & WIESEL, T. N. 1983. Clustered intrinsic connections in cat visual cortex. *J*
Neurosci, 3, 1116-33.
GILBERT, C. D. & WIESEL, T. N. 1990. The influence of contextual stimuli on the orientation
selectivity of cells in primary visual cortex of the cat. *Vision Res,* 30, 1689-701.
GIRARD, P., HUPE, J. M. & BULLIER, J. 2001. Feedforward and feedback connections between
areas V1 and V2 of the monkey have similar rapid conduction velocities. *J Neurophysiol,*
85, 1328-31.
GRINVALD, A., LIEKE, E. E., FROSTIG, R. D. & HILDESHEIM, R. 1994. Cortical point-spread
function and long-range lateral interactions revealed by real-time optical imaging of
macaque monkey primary visual cortex. *J Neurosci,* 14, 2545-68.
GUR, M., KAGAN, I. & SNODDERLY, D. M. 2005. Orientation and direction selectivity of neurons
in V1 of alert monkeys: functional relationships and laminar distributions. *Cereb Cortex,*
15, 1207-21.
HANSEN, B. J., CHELARU, M. I. & DRAGOI, V. 2012. Correlated variability in laminar cortical
circuits. *Neuron,* 76, 590-602.
HARRISON, M. T. & GEMAN, S. 2009. A rate and history-preserving resampling algorithm for
neural spike trains. *Neural Comput,* 21, 1244-58.

HARTLINE, H. K. 1938. THE RESPONSE OF SINGLE OPTIC NERVE FIBERS OF THE
VERTEBRATE EYE TO ILLUMINATION OF THE RETINA. *American Journal of*
Physiology-Legacy Content, 121**,** 400-415.
HEMBROOK-SHORT, J. R., MOCK, V. L., USREY, W. M. & BRIGGS, F. 2019. Attention
Enhances the Efficacy of Communication in V1 Local Circuits. *J Neurosci,* 39**,** 1066-1076.
HENDRICKSON, A. E., WILSON, J. R. & OGREN, M. P. 1978. The neuroanatomical organization
of pathways between the dorsal lateral geniculate nucleus and visual cortex in Old World
and New World primates. *J Comp Neurol,* 182**,** 123-36.
HENRY, C. A., JOSHI, S., XING, D., SHAPLEY, R. M. & HAWKEN, M. J. 2013. Functional
characterization of the extraclassical receptive field in macaque V1: contrast, orientation,
and temporal dynamics. *J Neurosci,* 33**,** 6230-42.
HESSE, J. K. & TSAO, D. Y. 2016. Consistency of Border-Ownership Cells across Artificial
Stimuli, Natural Stimuli, and Stimuli with Ambiguous Contours. *J Neurosci,* 36**,** 11338-
11349.
HESSE, J. K. & TSAO, D. Y. 2023. Functional modules for visual scene segmentation in macaque
visual cortex. *Proc Natl Acad Sci U S A,* 120**,** e2221122120.
HIRABAYASHI, T. & MIYASHITA, Y. 2005. Dynamically modulated spike correlation in monkey
inferior temporal cortex depending on the feature configuration within a whole object. *J*
Neurosci, 25**,** 10299-307.
HIRABAYASHI, T., TAKEUCHI, D., TAMURA, K. & MIYASHITA, Y. 2010. Triphasic dynamics of
stimulus-dependent information flow between single neurons in macaque inferior temporal
cortex. *J Neurosci,* 30**,** 10407-21.
HUANG, L., WANG, L., SHEN, W., LI, M., WANG, S., WANG, X., UNGERLEIDER, L. G. &
ZHANG, X. 2020. A source for awareness-dependent figure-ground segregation in human
prefrontal cortex. *Proc Natl Acad Sci U S A,* 117**,** 30836-30847.
HUBEL, D. H. & WIESEL, T. N. 1972. Laminar and columnar distribution of geniculo-cortical fibers
in the macaque monkey. *J Comp Neurol,* 146**,** 421-50.
HUNG, C. P., RAMSDEN, B. M. & ROE, A. W. 2007. A functional circuitry for edge-induced
brightness perception. *Nat Neurosci,* 10**,** 1185-90.
HUPE, J. M., JAMES, A. C., GIRARD, P., LOMBER, S. G., PAYNE, B. R. & BULLIER, J. 2001.
Feedback connections act on the early part of the responses in monkey visual cortex. *J*
Neurophysiol, 85**,** 134-45.
HUPE, J. M., JAMES, A. C., PAYNE, B. R., LOMBER, S. G., GIRARD, P. & BULLIER, J. 1998.
Cortical feedback improves discrimination between figure and background by V1, V2 and
V3 neurons. *Nature,* 394**,** 784-7.
ICHIDA, J. M., SCHWABE, L., BRESSLOFF, P. C. & ANGELUCCI, A. 2007. Response facilitation
from the "suppressive" receptive field surround of macaque V1 neurons. *J Neurophysiol,*
98**,** 2168-81.
JEHEE, J. F., LAMME, V. A. & ROELFSEMA, P. R. 2007. Boundary assignment in a recurrent
network architecture. *Vision Res,* 47**,** 1153-65.
JEURISSEN, D., SELF, M. W. & ROELFSEMA, P. R. 2016. Serial grouping of 2D-image regions
with object-based attention in humans. *Elife,* 5.
JIA, X., TANABE, S. & KOHN, A. 2013. gamma and the coordination of spiking activity in early
visual cortex. *Neuron,* 77**,** 762-74.
JIA, X. X., SIEGLE, J. H., DURAND, S., HELLER, G., RAMIREZ, T. K., KOCH, C. & OLSEN, S.
R. 2022. Multi-regional module-based signal transmission in mouse visual cortex. *Neuron,*
110**,** 1585-+.
JONES, H. E., ANDOLINA, I. M., SHIPP, S. D., ADAMS, D. L., CUDEIRO, J., SALT, T. E. &
SILLITO, A. M. 2015. Figure-ground modulation in awake primate thalamus. *Proc Natl*
Acad Sci U S A, 112**,** 7085-90.

JUN, J. J., STEINMETZ, N. A., SIEGLE, J. H., DENMAN, D. J., BAUZA, M., BARBARITS, B.,
LEE, A. K., ANASTASSIOU, C. A., ANDREI, A., AYDIN, C., BARBIC, M., BLANCHE, T.
J., BONIN, V., COUTO, J., DUTTA, B., GRATIY, S. L., GUTNISKY, D. A., HAUSSER, M.,
KARSH, B., LEDOCHOWITSCH, P., LOPEZ, C. M., MITELUT, C., MUSA, S., OKUN, M.,
PACHITARIU, M., PUTZEYS, J., RICH, P. D., ROSSANT, C., SUN, W. L., SVOBODA, K.,
CARANDINI, M., HARRIS, K. D., KOCH, C., O'KEEFE, J. & HARRIS, T. D. 2017. Fully
integrated silicon probes for high-density recording of neural activity. *Nature,* 551**,** 232-
236.
KAPADIA, M. K., ITO, M., GILBERT, C. D. & WESTHEIMER, G. 1995. Improvement in visual
sensitivity by changes in local context: parallel studies in human observers and in V1 of
alert monkeys. *Neuron,* 15**,** 843-56.
KELLER, A. J., ROTH, M. M. & SCANZIANI, M. 2020. Feedback generates a second receptive
field in neurons of the visual cortex. *Nature,* 582**,** 545-549.
KLEINER, M., BRAINARD, D. & PELLI, D. 2007. What's new in Psychtoolbox-3? *Perception,* 36**,**
14-14.
KLINK, P. C., DAGNINO, B., GARIEL-MATHIS, M. A. & ROELFSEMA, P. R. 2017. Distinct
Feedforward and Feedback Effects of Microstimulation in Visual Cortex Reveal Neural
Mechanisms of Texture Segregation. *Neuron,* 95**,** 209-220 e3.
KNIERIM, J. J. & VAN ESSEN, D. C. 1992. Neuronal responses to static texture patterns in area
V1 of the alert macaque monkey. *J Neurophysiol,* 67**,** 961-80.
KO, H. K. & VON DER HEYDT, R. 2018. Figure-ground organization in the visual cortex: does
meaning matter? *J Neurophysiol,* 119**,** 160-176.
KOGO, N., STRECHA, C., VAN GOOL, L. & WAGEMANS, J. 2010. Surface construction by a 2-
D differentiation-integration process: a neurocomputational model for perceived border
ownership, depth, and lightness in Kanizsa figures. *Psychol Rev,* 117**,** 406-39.
KOHN, A. & SMITH, M. A. 2005. Stimulus dependence of neuronal correlation in primary visual
cortex of the macaque. *J Neurosci,* 25**,** 3661-73.
KOURTZI, Z. & KANWISHER, N. 2001. Representation of perceived object shape by the human
lateral occipital complex. *Science,* 293**,** 1506-9.
LAMME, V. A. 1995. The neurophysiology of figure-ground segregation in primary visual cortex.
J Neurosci, 15**,** 1605-15.
LAMME, V. A., ZIPSER, K. & SPEKREIJSE, H. 1998. Figure-ground activity in primary visual
cortex is suppressed by anesthesia. *Proc Natl Acad Sci U S A,* 95**,** 3263-8.
LAYTON, O. W., MINGOLLA, E. & YAZDANBAKHSH, A. 2012. Dynamic coding of border-
ownership in visual cortex. *J Vis,* 12**,** 8.
LEE, T. S., YANG, C. F., ROMERO, R. D. & MUMFORD, D. 2002. Neural activity in early visual
cortex reflects behavioral experience and higher-order perceptual saliency. *Nat Neurosci,*
5**,** 589-97.
LI, W., PIECH, V. & GILBERT, C. D. 2006. Contour saliency in primary visual cortex. *Neuron,* 50**,**
951-62.
LI, Y., WANG, T., YANG, Y., DAI, W., WU, Y., LI, L., HAN, C., ZHONG, L., LI, L., WANG, G.,
DOU, F. & XING, D. 2022. Cascaded normalizations for spatial integration in the primary
visual cortex of primates. *Cell Rep,* 40**,** 111221.
LIANG, H., GONG, X., CHEN, M., YAN, Y., LI, W. & GILBERT, C. D. 2017. Interactions between
feedback and lateral connections in the primary visual cortex. *Proc Natl Acad Sci U S A,*
114**,** 8637-8642.
LUND, J. S. 1973. Organization of neurons in the visual cortex, area 17, of the monkey (Macaca
mulatta). *J Comp Neurol,* 147**,** 455-96.
LUND, J. S. 1988. Anatomical organization of macaque monkey striate visual cortex. *Annu Rev
Neurosci,* 11**,** 253-88.

LUND, J. S. & YOSHIOKA, T. 1991. Local circuit neurons of macaque monkey striate cortex: III.
Neurons of laminae 4B, 4A, and 3B. *J Comp Neurol,* 311**,** 234-58.
MARCUS, D. S. & VAN ESSEN, D. C. 2002. Scene segmentation and attention in primate cortical
areas V1 and V2. *J Neurophysiol,* 88**,** 2648-58.
MARKOV, N. T., VEZOLI, J., CHAMEAU, P., FALCHIER, A., QUILODRAN, R., HUISSOUD, C.,
LAMY, C., MISERY, P., GIROUD, P., ULLMAN, S., BARONE, P., DEHAY, C.,
KNOBLAUCH, K. & KENNEDY, H. 2014. Anatomy of hierarchy: feedforward and feedback
pathways in macaque visual cortex. *J Comp Neurol,* 522**,** 225-59.
MARTIN, A. B. & VON DER HEYDT, R. 2015. Spike synchrony reveals emergence of proto-
objects in visual cortex. *Journal of Neuroscience,* 35**,** 6860-6870.
MATES, S. L. & LUND, J. S. 1983. Neuronal composition and development in lamina 4C of
monkey striate cortex. *J Comp Neurol,* 221**,** 60-90.
MEHRANI, P. & TSOTSOS, J. K. 2021. Early recurrence enables figure border ownership. *Vision*
Res, 186**,** 23-33.
MELSSEN, W. J. & EPPING, W. J. 1987. Detection and estimation of neural connectivity based
on crosscorrelation analysis. *Biol Cybern,* 57**,** 403-14.
MOORE, G. P., SEGUNDO, J. P., PERKEL, D. H. & LEVITAN, H. 1970. Statistical signs of
synaptic interaction in neurons. *Biophys J,* 10**,** 876-900.
MULLER, J. R., METHA, A. B., KRAUSKOPF, J. & LENNIE, P. 2003. Local signals from beyond
the receptive fields of striate cortical neurons. *J Neurophysiol,* 90**,** 822-31.
NAKAYAMA, K., HE, Z. J. & SHIMOJO, S. 1995. Visual Surface Representation: A Critical Link
between Lower-Level and Higher-Level Vision. *In:* KOSSLYN, S. M. & OSHERSON, D.
N. (eds.) *An Invitation to Cognitive Science, Volume 2: Visual Cognition.* The MIT Press.
NANDY, A. S., NASSI, J. J. & REYNOLDS, J. H. 2017. Laminar Organization of Attentional
Modulation in Macaque Visual Area V4. *Neuron,* 93**,** 235-246.
NASSI, J. J., LOMBER, S. G. & BORN, R. T. 2013. Corticocortical feedback contributes to
surround suppression in V1 of the alert primate. *J Neurosci,* 33**,** 8504-17.
NELSON, J. I. & FROST, B. J. 1985. Intracortical facilitation among co-oriented, co-axially aligned
simple cells in cat striate cortex. *Exp Brain Res,* 61**,** 54-61.
NELSON, J. I., SALIN, P. A., MUNK, M. H., ARZI, M. & BULLIER, J. 1992. Spatial and temporal
coherence in cortico-cortical connections: a cross-correlation study in areas 17 and 18 in
the cat. *Vis Neurosci,* 9**,** 21-37.
NICHOLSON, C. & FREEMAN, J. A. 1975. Theory of current source-density analysis and
determination of conductivity tensor for anuran cerebellum. *J Neurophysiol,* 38**,** 356-68.
NOTHDURFT, H. C., GALLANT, J. L. & VAN ESSEN, D. C. 1999. Response modulation by
texture surround in primate area V1: correlates of "popout" under anesthesia. *Vis*
Neurosci, 16**,** 15-34.
NURMINEN, L., MERLIN, S., BIJANZADEH, M., FEDERER, F. & ANGELUCCI, A. 2018. Top-
down feedback controls spatial summation and response amplitude in primate visual
cortex. *Nat Commun,* 9**,** 2281.
O'HERRON, P. J. & VON DER HEYDT, R. 2009. Short-term memory for figure-ground
organization in the visual cortex. *Neuron,* 61**,** 801-809.
O'KUSKY, J. & COLONNIER, M. 1982. A laminar analysis of the number of neurons, glia, and
synapses in the adult cortex (area 17) of adult macaque monkeys. *J Comp Neurol,* 210**,**
278-90.
OSTOJIC, S., BRUNEL, N. & HAKIM, V. 2009. How connectivity, background activity, and
synaptic properties shape the cross-correlation between spike trains. *J Neurosci,* 29**,**
10234-53.
PACHITARIU, M., SRIDHAR, S., PENNINGTON, J. & STRINGER, C. 2024. Spike sorting with
Kilosort4. *Nat Methods,* 21**,** 914-921.

PAK, A., RYU, E., LI, C. & CHUBYKIN, A. A. 2020. Top-Down Feedback Controls the Cortical
Representation of Illusory Contours in Mouse Primary Visual Cortex. *J Neurosci,* 40**,** 648-
660.
PERKEL, D. H., GERSTEIN, G. L. & MOORE, G. P. 1967. Neuronal spike trains and stochastic
point processes. II. Simultaneous spike trains. *Biophys J,* 7**,** 419-40.
PETTINE, W. W., STEINMETZ, N. A. & MOORE, T. 2019. Laminar segregation of sensory coding
and behavioral readout in macaque V4. *Proc Natl Acad Sci U S A,* 116**,** 14749-14754.
POLTORATSKI, S. & TONG, F. 2020. Resolving the Spatial Profile of Figure Enhancement in
Human V1 through Population Receptive Field Modeling. *J Neurosci,* 40**,** 3292-3303.
POORT, J., RAUDIES, F., WANNIG, A., LAMME, V. A., NEUMANN, H. & ROELFSEMA, P. R.
2012. The role of attention in figure-ground segregation in areas V1 and V4 of the visual
cortex. *Neuron,* 75**,** 143-56.
QIU, F. T., SUGIHARA, T. & VON DER HEYDT, R. 2007. Figure-ground mechanisms provide
structure for selective attention. *Nature Neuroscience,* 10**,** 1492-1499.
RAMSDEN, B. M., HUNG, C. P. & ROE, A. W. 2001. Real and illusory contour processing in area
V1 of the primate: a cortical balancing act. *Cereb Cortex,* 11**,** 648-65.
REID, R. C. & ALONSO, J. M. 1995. Specificity of monosynaptic connections from thalamus to
visual cortex. *Nature,* 378**,** 281-4.
ROCKLAND, K. S. & LUND, J. S. 1983. Intrinsic laminar lattice connections in primate visual
cortex. *J Comp Neurol,* 216**,** 303-18.
ROCKLAND, K. S. & PANDYA, D. N. 1979. Laminar origins and terminations of cortical
connections of the occipital lobe in the rhesus monkey. *Brain Res,* 179**,** 3-20.
ROCKLAND, K. S., SALEEM, K. S. & TANAKA, K. 1994. Divergent feedback connections from
areas V4 and TEO in the macaque. *Vis Neurosci,* 11**,** 579-600.
ROCKLAND, K. S. & VIRGA, A. 1989. Terminal arbors of individual "feedback" axons projecting
from area V2 to V1 in the macaque monkey: a study using immunohistochemistry of
anterogradely transported Phaseolus vulgaris-leucoagglutinin. *J Comp Neurol,* 285**,** 54-
72.
ROE, A. W. & TS'O, D. Y. 1999. Specificity of color connectivity between primate V1 and V2. *J
Neurophysiol,* 82**,** 2719-30.
ROELFSEMA, P. R. 2006. Cortical algorithms for perceptual grouping. *Annu Rev Neurosci,* 29**,**
203-27.
SAKAI, K. & NISHIMURA, H. 2006. Surrounding suppression and facilitation in the determination
of border ownership. *J Cogn Neurosci,* 18**,** 562-79.
SCHWABE, L., OBERMAYER, K., ANGELUCCI, A. & BRESSLOFF, P. C. 2006. The role of
feedback in shaping the extra-classical receptive field of cortical neurons: a recurrent
network model. *J Neurosci,* 26**,** 9117-29.
SCHWARZ, C. & BOLZ, J. 1991. Functional specificity of a long-range horizontal connection in
cat visual cortex: a cross-correlation study. *J Neurosci,* 11**,** 2995-3007.
SELF, M. W., JEURISSEN, D., VAN HAM, A. F., VAN VUGT, B., POORT, J. & ROELFSEMA, P.
R. 2019. The Segmentation of Proto-Objects in the Monkey Primary Visual Cortex. *Curr
Biol,* 29**,** 1019-1029 e4.
SELF, M. W., VAN KERKOERLE, T., SUPER, H. & ROELFSEMA, P. R. 2013. Distinct roles of
the cortical layers of area V1 in figure-ground segregation. *Curr Biol,* 23**,** 2121-9.
SENZAI, Y., FERNANDEZ-RUIZ, A. & BUZSAKI, G. 2019. Layer-Specific Physiological Features
and Interlaminar Interactions in the Primary Visual Cortex of the Mouse. *Neuron,* 101**,** 500-
513 e5.
SHMUEL, A., KORMAN, M., STERKIN, A., HAREL, M., ULLMAN, S., MALACH, R. & GRINVALD,
A. 2005. Retinotopic axis specificity and selective clustering of feedback projections from
V2 to V1 in the owl monkey. *J Neurosci,* 25**,** 2117-31.

SIBILLE, J., GEHR, C., BENICHOV, J. I., BALASUBRAMANIAN, H., TEH, K. L., LUPASHINA, T., VALLENTIN, D. & KREMKOW, J. 2022. High-density electrode recordings reveal strong and specific connections between retinal ganglion cells and midbrain neurons. *Nature Communications,* 13.

SIEGLE, J. H., JIA, X., DURAND, S., GALE, S., BENNETT, C., GRADDIS, N., HELLER, G., RAMIREZ, T. K., CHOI, H., LUVIANO, J. A., GROBLEWSKI, P. A., AHMED, R., ARKHIPOV, A., BERNARD, A., BILLEH, Y. N., BROWN, D., BUICE, M. A., CAIN, N., CALDEJON, S., CASAL, L., CHO, A., CHVILICEK, M., COX, T. C., DAI, K., DENMAN, D. J., DE VRIES, S. E. J., DIETZMAN, R., ESPOSITO, L., FARRELL, C., FENG, D., GALBRAITH, J., GARRETT, M., GELFAND, E. C., HANCOCK, N., HARRIS, J. A., HOWARD, R., HU, B., HYTNEN, R., IYER, R., JESSETT, E., JOHNSON, K., KATO, I., KIGGINS, J., LAMBERT, S., LECOQ, J., LEDOCHOWITSCH, P., LEE, J. H., LEON, A., LI, Y., LIANG, E., LONG, F., MACE, K., MELCHIOR, J., MILLMAN, D., MOLLENKOPF, T., NAYAN, C., NG, L., NGO, K., NGUYEN, T., NICOVICH, P. R., NORTH, K., OCKER, G. K., OLLERENSHAW, D., OLIVER, M., PACHITARIU, M., PERKINS, J., REDING, M., REID, D., ROBERTSON, M., RONELLENFITCH, K., SEID, S., SLAUGHTERBECK, C., STOECKLIN, M., SULLIVAN, D., SUTTON, B., SWAPP, J., THOMPSON, C., TURNER, K., WAKEMAN, W., WHITESELL, J. D., WILLIAMS, D., WILLIFORD, A., YOUNG, R., ZENG, H., NAYLOR, S., PHILLIPS, J. W., REID, R. C., MIHALAS, S., OLSEN, S. R. & KOCH, C. 2021. Survey of spiking in the mouse visual system reveals functional hierarchy. *Nature,* 592**,** 86-92.

SILLITO, A. M., GRIEVE, K. L., JONES, H. E., CUDEIRO, J. & DAVIS, J. 1995. Visual cortical mechanisms detecting focal orientation discontinuities. *Nature,* 378**,** 492-6.

SIU, C., BALSOR, J., MERLIN, S., FEDERER, F. & ANGELUCCI, A. 2021. A direct interareal feedback-to-feedforward circuit in primate visual cortex. *Nat Commun,* 12**,** 4911.

SMITH, M. A. & KOHN, A. 2008. Spatial and temporal scales of neuronal correlation in primary visual cortex. *J Neurosci,* 28**,** 12591-603.

SOMOGYI, P. & COWEY, A. 1981. Combined Golgi and electron microscopic study on the synapses formed by double bouquet cells in the visual cortex of the cat and monkey. *J Comp Neurol,* 195**,** 547-66.

STEINMETZ, P. N., ROY, A., FITZGERALD, P. J., HSIAO, S. S., JOHNSON, K. O. & NIEBUR, E. 2000. Attention modulates synchronized neuronal firing in primate somatosensory cortex. *Nature,* 404**,** 187-90.

STETTLER, D. D., DAS, A., BENNETT, J. & GILBERT, C. D. 2002. Lateral connectivity and contextual interactions in macaque primary visual cortex. *Neuron,* 36**,** 739-50.

SUGIHARA, T., QIU, F. T. & VON DER HEYDT, R. 2011. The speed of context integration in the visual cortex. *J Neurophysiol,* 106**,** 374-85.

SUPER, H., ROMEO, A. & KEIL, M. 2010. Feed-forward segmentation of figure-ground and assignment of border-ownership. *PLoS One,* 5**,** e10705.

SUPER, H., SPEKREIJSE, H. & LAMME, V. A. 2001. Two distinct modes of sensory processing observed in monkey primary visual cortex (V1). *Nat Neurosci,* 4**,** 304-10.

SWINDALE, N. V. 1998. Orientation tuning curves: empirical description and estimation of parameters. *Biol Cybern,* 78**,** 45-56.

TOOTELL, R. B., SWITKES, E., SILVERMAN, M. S. & HAMILTON, S. L. 1988. Functional anatomy of macaque striate cortex. II. Retinotopic organization. *J Neurosci,* 8**,** 1531-68.

TRAUTMANN, E. M., HESSE, J. K., STINE, G. M., XIA, R., ZHU, S., O'SHEA, D. J., KARSH, B., COLONELL, J., LANFRANCHI, F. F., VYAS, S., ZIMNIK, A., AMEMATSRO, E., STEINEMANN, N. A., WAGENAAR, D. A., PACHITARIU, M., ANDREI, A., LOPEZ, C. M., O'CALLAGHAN, J., PUTZEYS, J., RADUCANU, B. C., WELKENHUYSEN, M., CHURCHLAND, M., MOORE, T., SHADLEN, M., SHENOY, K., TSAO, D., DUTTA, B. &

HARRIS, T. 2025. Large-scale high-density brain-wide neural recording in nonhuman primates. *Nat Neurosci,* 28, 1562-1575.

TREPKA, E. B., ZHU, S. D., XIA, R. B., CHEN, X. M. & MOORE, T. 2022. Functional interactions among neurons within single columns of macaque V1. *Elife,* 11.

TS'O, D. Y., GILBERT, C. D. & WIESEL, T. N. 1986. Relationships between horizontal interactions and functional architecture in cat striate cortex as revealed by cross-correlation analysis. *J Neurosci,* 6, 1160-70.

USREY, W. M., REPPAS, J. B. & REID, R. C. 1998. Paired-spike interactions and synaptic efficacy of retinal inputs to the thalamus. *Nature,* 395, 384-7.

VAN KERKOERLE, T., SELF, M. W., DAGNINO, B., GARIEL-MATHIS, M. A., POORT, J., VAN DER TOGT, C. & ROELFSEMA, P. R. 2014. Alpha and gamma oscillations characterize feedback and feedforward processing in monkey visual cortex. *Proc Natl Acad Sci U S A,* 111, 14332-41.

VAN KERKOERLE, T., SELF, M. W. & ROELFSEMA, P. R. 2017. Layer-specificity in the effects of attention and working memory on activity in primary visual cortex. *Nat Commun,* 8, 13804.

VANGENEUGDEN, J., VAN BEEST, E. H., COHEN, M. X., LORTEIJE, J. A. M., MUKHERJEE, S., KIRCHBERGER, L., MONTIJN, J. S., THAMIZHARASU, P., CAMILLO, D., LEVELT, C. N., ROELFSEMA, P. R., SELF, M. W. & HEIMEL, J. A. 2019. Activity in Lateral Visual Areas Contributes to Surround Suppression in Awake Mouse V1. *Curr Biol,* 29, 4268-4275 e7.

VON DER HEYDT, R. 2015. Figure-ground organization and the emergence of proto-objects in the visual cortex. *Frontiers in Psychology,* 6, 1695.

VON DER HEYDT, R. 2023. Visual cortical processing—From image to object representation. *Frontiers in Computer Science,* 5.

VON DER HEYDT, R., PETERHANS, E. & BAUMGARTNER, G. 1984. Illusory contours and cortical neuron responses. *Science,* 224, 1260-2.

WAGATSUMA, N., HU, B., VON DER HEYDT, R. & NIEBUR, E. 2021. Analysis of spiking synchrony in visual cortex reveals distinct types of top-down modulation signals for spatial and object-based attention. *PLoS Comput Biol,* 17, e1008829.

WALKER, G. A., OHZAWA, I. & FREEMAN, R. D. 1999. Asymmetric suppression outside the classical receptive field of the visual cortex. *J Neurosci,* 19, 10536-53.

WEBB, B. S., TINSLEY, C. J., BARRACLOUGH, N. E., PARKER, A. & DERRINGTON, A. M. 2003. Gain control from beyond the classical receptive field in primate primary visual cortex. *Vis Neurosci,* 20, 221-30.

WILLIFORD, J. R. & VON DER HEYDT, R. 2016. Figure-Ground Organization in Visual Cortex for Natural Scenes. *eNeuro,* 3.

WISER, A. K. & CALLAWAY, E. M. 1996. Contributions of individual layer 6 pyramidal neurons to local circuitry in macaque primary visual cortex. *J Neurosci,* 16, 2724-39.

XU, X., OLIVAS, N. D., IKRAR, T., PENG, T., HOLMES, T. C., NIE, Q. & SHI, Y. 2016. Primary visual cortex shows laminar-specific and balanced circuit organization of excitatory and inhibitory synaptic connectivity. *J Physiol,* 594, 1891-910.

YABUTA, N. H. & CALLAWAY, E. M. 1998. Functional streams and local connections of layer 4C neurons in primary visual cortex of the macaque monkey. *J Neurosci,* 18, 9489-99.

YAN, Y., ZHAOPING, L. & LI, W. 2018. Bottom-up saliency and top-down learning in the primary visual cortex of monkeys. *Proc Natl Acad Sci U S A,* 115, 10499-10504.

YOSHIOKA, T., LEVITT, J. B. & LUND, J. S. 1994. Independence and merger of thalamocortical channels within macaque monkey primary visual cortex: anatomy of interlaminar projections. *Vis Neurosci,* 11, 467-89.

ZARELLA, M. D. & TS'O, D. Y. 2016. Cue combination encoding via contextual modulation of V1 and V2 neurons. *Eye Brain,* 8, 177-193.

ZHANG, N. R. & VON DER HEYDT, R. 2010. Analysis of the context integration mechanisms
underlying figure-ground organization in the visual cortex. *Journal of Neuroscience,* 30**,**
6482-6496.
ZHANG, S., XU, M., KAMIGAKI, T., HOANG DO, J. P., CHANG, W. C., JENVAY, S., MIYAMICHI,
K., LUO, L. & DAN, Y. 2014. Selective attention. Long-range and local circuits for top-
down modulation of visual cortex processing. *Science,* 345**,** 660-5.
ZHAOPING, L. 2005. Border ownership from intracortical interactions in visual area v2. *Neuron,*
47**,** 143-53.
ZHOU, H., FRIEDMAN, H. S. & VON DER HEYDT, R. 2000. Coding of border ownership in
monkey visual cortex. *J Neurosci,* 20, 6594-611.
ZHU, S., XIA, R., CHEN, X. & MOORE, T. 2024. Comparison of orientation encoding across
layers within single columns of primate V1 revealed by high-density recordings. *Front*
Neural Circuits, 18**,** 1399571.
ZHU, S. D., ZHANG, L. A. & VON DER HEYDT, R. 2020. Searching for object pointers in the
visual cortex. *Journal of Neurophysiology,* 123**,** 1979-1994.
ZIPSER, K., LAMME, V. A. & SCHILLER, P. H. 1996. Contextual modulation in primary visual
cortex. *J Neurosci,* 16**,** 7376-89.
1247

Figures and legend

Figure 1. Classical receptive field (CRF) stimulation in single V1 columns.
(A) Neuropixels probe (384 channels/3.84mm) penetrations made into the lateral surface or underlying calcarine sulcus of macaque V1. Bottom: Diagram of designated laminar compartments. (B) Current-Source density (CSD) profile from an example recording session (M1, *pen1*), showing the early current sink (red) indicative of layer 4C. Laminar compartment boundaries (dashed lines) were determined using histological data and the CSD profile. (C) Top: 71 well-defined classical receptive fields (CRF) vertically stacked along the cortical depth, mapped with a 10 x 10 array (4 x 4 dva) of probe stimuli (gratings, 0.4 dva in diameter). Bottom: Drifting Gabor grating (1.5 dva in diameter) used for CRF stimulation, positioned within the joint receptive fields of recorded neurons. Zero (x, y) denotes the center of the CRF population. (D) Heatmap of visual responses from 159 neurons to 36 drift directions of Gabor gratings, vertically stacked along cortical depth (M1, *pen1*). Top: Distribution of differences in preferred orientation for all pairwise combinations of neurons.

Figure 2. Nonclassical receptive field (nCRF) stimulation and border ownership.
(A) Border ownership test stimuli for nCRF stimulation. Stimuli consisted of uniformly white or black objects (8º squares) on black or white backgrounds, respectively. One border of the object fell within the CRFs (red circle) of the recorded neurons, with orientation varying in 45º steps. Within the CRF, depending on local contrast polarity (LC), the border could either be a dark-light edge (LC1, Left column) or a light-dark edge (LC2, Right column). For the same LC condition, the border within the CRF could be the bottom-left edge (Side 1, top row) or top-right edge (Side 2, bottom row) of the object. (B) Enlarged view of stimulus configuration within the CRF, showing identical stimulus regions between Side 1 and Side 2 for each LC condition (8 x 16 dva). (C) Four example B_{own} neurons recorded from V1. For each neuron, dot raster plots (top) and instantaneous firing rates (bottom) show responses to the four stimulus conditions depicted in (A), respectively. Magenta represents Side 1; green represents Side 2. (D) Bar plots showing the mean spike rates of the four neurons across the four stimulus conditions. (E) Top: Cartoon depicting the preferred border orientation for each of the four recording sessions (*pen*1-4). Arrows indicate the preferred side of objects relative to the border in the CRF. Bottom: B_{own} modulation index for each neuron plotted at its recorded cortical depth across the recordings. Color bars on the ordinate represent laminar compartments. Black dots denote statistically significant B_{own} neurons, determined by ANOVA with object side and local contrast as factors ($p < 0.05$). Middle: Marginal distribution of the B_{own} modulation index, red arrows indicating the population median. *$p < 0.05$; ***$p < 10^{-5}$.

Figure 3. Interlaminar cross-correlations during CRF and nCRF stimulation.

(A) Example recording (M1, *pen1*) showing 159 visually responsive neurons (circles) recorded simultaneously during CRF stimulation, plotted at their relative cortical depth. The abscissa is magnified for visualization. Also shown are eight example neuronal pairs with significant CCGs during CRF stimulation. Neuronal pairs with zero-time lag CCGs are connected by dashed lines, while pairs with nonzero-time lag CCGs are connected by arrowed lines, with the arrow pointing to the lagging neuron. (B) Eight corresponding CCGs from (A). Neuronal pairs are separated by their laminar compartment combinations. The reference neuron (1st neuron in the CCG function) is designated as *j*, and the target neuron as *k*. PL (peak lag, ms) is defined as the relative time delay in peak correlation between the two spike trains. (C) Diagram illustrating feedforward and feedback (and horizontal) projections within the designated laminar compartments. Feedforward inputs arrive within laminar compartments 4C and 6, while feedback (and horizontal) projections principally target superficial and deep layers. (D) Heatmap of all significant CCGs during either CRF stimulation (Gabor gratings, top row) or nCRF stimulation (border ownership, middle row), for the four key interlaminar combinations (columns), calculated from one example recording (M1, *pen1*). Individual CCGs are normalized by their absolute maximum for visualization, and different CCGs are vertically stacked. Bottom row: Distribution of CCG asymmetry for each of the four key interlaminar compartments as in (A), across CRF stimulation (gray) and nCRF stimulation (magenta). Arrows indicate the population median. **p < 0.002; ***p < 10^{-5}.

Figure 4. Comparison of interlaminar CCG asymmetries during CRF and nCRF stimulation.
Combined dataset for each of the four key interlaminar combinations. Histograms and cumulative distributions of asymmetry for all significant CCGs during CRF (gray) and nCRF (magenta) stimulation. Arrows indicate the population median. ***$p < 10^{-5}$.

Figure 5. Dependence of nCRF modulation on interlaminar signal flow.
(A) Scatter plot of the B_{own} modulation index versus CCG asymmetry for all significant CCGs, aggregated across the four key interlaminar combinations and all recording sessions. CCG asymmetries were calculated during nCRF stimulation. The geometric mean of the B_{own} index for each neuronal pair was used. Red lines denote the fitted regression line and the 95% confidence interval in linear regression. (B) Similar to (A), but using the local contrast (LC) index instead of the B_{own} index. The LC index was calculated as the difference in neuronal responses to borders with opposite contrast polarities, normalized by the maximum response.

Supplementary figures and legends

Supplementary Figure S1. Histological data for Neuropixels recordings in macaque V1.
(A) Neuropixels probe (384 channels/3.84mm) penetrations made into the lateral surface or underlying calcarine sulcus of macaque V1. Bottom: Diagram of designated laminar compartments. (B) Histological reconstruction of probe tracks based on DiI derivative staining for two example recordings (*pen3-4*). Here, Neuropixels probes passed through the lateral surface and reached the calcarine sulcus of V1. Only neurons in the calcarine V1 from these two recordings were tested and included in the analysis. Designated boundaries are shown. Scale bar, 500 μm.

Supplementary Figure S2. Comparison of interlaminar cross-correlations during CRF and nCRF stimulation under different CCG criteria.

(A) Histogram distributions of CCG peak lags for each of the four key interlaminar combinations in the combined dataset during CRF (gray) and nCRF (magenta) stimulation. The shift (Δ) in median CCG peak lags during nCRF stimulation relative to CRF stimulation is shown. Significant CCGs are defined as those with peaks exceeding 7 SD above the noise level (flank). (B) Similar to (A), but comparing CCG asymmetry across all CCGs, including both significant and non-significant ones. (C) Comparison of CCG peak lags using a stricter criterion, where significant CCGs are defined as those with peaks exceeding 10 SD above the noise level (flank). (D) Comparison of CCG asymmetry using the stricter criterion of 10 SD.