Limnospira (Cyanobacteria) chemical fingerprint reveals local molecular adaptation

Théotime Roussel[1,3], Cédric Hubas[2], Sébastien Halary[1], Mathias Chynel[2], Charlotte Duval[1], Jean-Paul Cadoret[3], Tarik Meziane[2], Léa Vernès[3], Claude Yéprémian[1], Cécile Bernard[1*], Benjamin Marie[1*]

[1] UMR7245 MCAM MNHN-CNRS, Muséum National d'Histoire Naturelle, CP 39, 12 rue Buffon, F-75231 Paris Cedex 05, France

[2] UMR 7208 BOREA MNHN-CNRS-SU-IRD, Muséum d'Histoire Naturelle, CP 53, 61 rue Buffon, F-75231 Paris Cedex 05, France

[3] Algama, 81 rue Réaumur, 75002 PARIS France

* Correspondance: benjamin.marie@mnhn.fr; cecile.bernard@mnhn.fr (+33140793212, +33140793183, BM; CB)

Marie B., https://orcid.org/0000-0001-9880-5541

Bernard C., http://orcid.org/0000-0001-7032-3989

Abstract

Limnospira can colonize a wide variety of environments (*e.g.* freshwater, brackish, alkaline or alkaline-saline water) and develop permanent blooms that limit overshadowed adjacent photo-trophs diversity, especially in alkaline and saline environments. Previous phylogenomic analysis of *Limnospira* allowed us to distinguish two major phylogenetic clades (I and II) but failed to clearly segregate strains according to their respective habitats in terms of salinity or of biogeography. In the present work, we attempt to determine whether *Limnospira* display metabolic signatures specific to its different habitats, particularly brackish or alkaline-saline ecosystems, and question the impact of accessory genes repertoire on respective chemical adaptation.

The study of the metabolomic diversity of 93 strains of *Limnospira* of the Paris Museum Collection, grown under standardized lab culture conditions, showed clearly distinct chemical fingerprints that were correlated with the respective biogeographic origin of the strains. The molecules that most discriminate the different *Limnospira* geographic groups are sugars, lipids, peptides, photosynthetic pigments, and antioxidant molecules. Interestingly, these molecule enrichments might represent adaptation traits to the local conditions encountered in their respective sampling environments concerning salinity, light and oxidative stress. We hypothesize that within extreme environments, such as those colonized by *Limnospira*, flexible genes that are necessary for the adaptation to specific local environmental conditions (*e.g.* salinity, light, oxidative pressure) are selected, leading to the specific production of certain metabolites involved in stress defence mechanisms.

Importance

Cyanobacteria are a group of photoautotrophic microorganisms that have colonized a wide range of environments around the world. Phylogenomic analysis of *Limnospira platensis* al-

lowed to distinguish two major phylogenetic clades (I and II) but failed to clearly segregate

strains following their habitats in terms of salinity or biogeography. One can presume that the

genes found within this variable portion of the genome of these clades could be involved in *L.*

platensis adaptation to local environmental conditions. In the present paper, we attempt to

determine whether *Limnospira platensis* display metabolic signatures specific for its different

habitats, particularly brackish or alkaline-saline ecosystems, and question the impact of acces-

sory gene repertoire on respective chemical adaptation.

Keywords: Cyanobacteria, metabolomics, genomics, adaptation, micro-diversity.

Introduction

The capacity of phytoplanktonic organism to durably colonize their environment appears to be balanced by their specific ability to perform efficient primary production avoiding being overshadowed by competitors and dealing with physico-chemical environmental constraints. This adaptive power is strongly dependent on the energetic cost they must pay for it, in terms of genomic or physiological expenses[1].

Cyanobacteria are a group of photoautotrophic microorganisms that have colonized a wide range of environments around the world[2-3]. The large diversity of cyanobacteria species is accompanied by a great diversity of primary and/or secondary/accessory metabolites, many of which presenting noticeable biological activities of central interest in key cellular or ecological functions[4-5]. The genus *Limnospira* (anc. *Arthrospira*), better known under its commercial/vernacular name 'spirulina', is one of the most extensively cyanobacteria taxa studied so far (*e.g.* more than 300 publications per year since 2020 on PubMed), especially due to its industrial potential (*e.g.* agri-food, pharmaceuticals, cosmetics)[6-7]. Indeed, cultivated *Limnospira* present a remarkably high growth rate under appropriate culture conditions (more than one division per day)[8-9], and are very rich in proteins (up to 70% of the dry weight)[10], unsaturated fatty acids[11], vitamins (at least 9)[12-13], and various molecules with antioxidant properties (*e.g.* polyphenols, carotenoids, ergothioneine, biopterins)[12,14]. To this aim, physiological aptitudes of certain *Limnospira* strains have been examined (*i.e.,* effect by modifying a culture parameter on biomass or molecule production)[8,15-18], whereas only few studies have focused on the global ecological trait of this taxon[19].

In a recent study based on a polyphasic approach combined with comparative genomics, we showed that the *Limnospira* genus is monospecific, represented by one single species, *Limnospira platensis*, which is cosmopolitan[20]. Yet, *Limnospira platensis* seems to develop a

singular ecological strategy, as this species is able to colonize a wide variety of environments

(*e.g.* freshwater, brackish, alkaline or alkaline-saline water)[19-20] in which it dominate the

phytoplanktonic community by developing permanent blooms that limit overshadowed

adjacent phototrophs diversity[19,21].

Phylogenomic analysis of *L. platensis* allowed to distinguish two major phylogenetic clades (I

and II), but failed to clearly segregate strains following their habitats in terms of salinity or

biogeography[20]. Such discrepancies have already been observed in other taxa, for instance for

the widespread extremophilic bacteria *Salinibacter ruber*[22]. However, in this case, non-

targeted metabolites analysis succeeded to segregate the strains, highlighting biogeographical

metabolic signatures related to local adaptations. Recently, the same approach has suggested

the existence of distinct chemical signatures between different ecotypes of the oceanic

picocyanobacterium *Prochlorococcus marinus*, each being adapted to a specific niche

characterized in terms of light intensity and temperature[23].

Concerning the species *L. platensis*, the flexible part of the genome remarkably represents up-

to 55% of the genome of each strain, and differs from one strain to another, carries large

variety of genes (carrying some gene with regulatory and defensive functions)[20], many of

which present so-far unknown functions and could potentially be related to specific

capabilities of the strains in term of physiology and ecology. One can presume that the genes

found within this variable portion of the genomes could be involved in *L. platensis* adaptation

to local environment conditions. These genes could indeed be responsible of carrying traits

that allow the species to adjust to specific local condition variation and support the

maintaining of constant blooms. Ultimately, this local adaptation might give *L. platensis* an

advantage to overpass other phytoplankton competitors within these environments.

101 In the present paper, we attempt to determine whether *Limnospira platensis* display metabolic

102 signatures specific for its different habitats, particularly brackish or alkaline-saline

103 ecosystems, and question the impact of accessory genes repertoire on respective chemical

104 adaptation. In this aim, we have investigated a wide panel of strains from different

105 environments and maintained in collections under identical culture conditions. We used two

106 types of extraction and chemical analysis approaches pursuing a large diversity of metabolites

107 potentially involved in various adaptative traits. First, we applied a targeted approach,

108 focusing on photosynthetic pigments and fatty acids known to be involved in adaptation to

109 different environmental conditions (*e.g.* light, salinity, pH). Then, we performed in parallel a

110 non-targeted approach to obtain an overall chemical fingerprint of each strain and look for

111 other potential differences.

Material and Methods

Strains and culture conditions

115 93 strains of *L. platensis* were studied. 89 strains were obtained from the Paris Museum

116 Collection (PMC)[24]: 72 from two Camargue area, a humid region located in the Rhône delta

117 (South of France) mostly consisting of marshland and brine lagoons, 13 from the lake Dziani

118 Dzaha, an alkaline and thalassohaline lake located on the island Petite-Terre of Mayotte, three

119 from lake Natron, an alkaline-saline lake located between Tanzania and Kenya, and one strain

120 from historical collection with no environmental data available. Three additional strains were

121 provided by other collections: *L. platensis* PCC 8005 and *L. platensis* PCC 7345 (Pasteur

122 Culture Collection of Cyanobacteria, France) from unknown locations, and *L. platensis* SAG

123 85.79 (Culture Collection of Algae at Göttingen University, Germany) from the lake Natron.

124 one strain provided by Spirulina Solutions (https://spirulinasolutions.fr/): Paracas R14 from a

125 freshwater lake in Peru. All information of strains, sampling site properties, year of isolation,

and genomic information are given in Supplementary Table S1. These 93 strains were

maintained as monospecific but non-axenic cultures, growing in 30 mL of ergonomically

improved Spirulina medium[24] at 24°C with a photon flux density of 30 μmol photon.$m^{-2}.s^{-1}$

under a 16:8-h light:dark photoperiod. Each month, 1/10 of each culture were used to

inoculated new cultures. The remaining 9/10 were centrifugated (10 min, 3,220 g, 3 times).

The supernatants were discarded, and the pellet were stored at -80°C during at least 3h, and

then lyophilization was performed (-47°C, 0.03 mBar for 16 h) (Labconco, Kansas City, MO,

USA). Six cycles of culture-biomass sampling have been made. The six freeze-dried biomass

samples of each strain have been mixed together and homogenized prior to chemical analyses.

Pigments

Hydrophilic pigments of the 93 strains were analyzed by spectrophotometry according to

Yéprémian et al.[25-26]. Ten mg of freeze-dried biomass were incubated with 10 mL of a saline

buffer solution (NaCl 0.15 $mol.L^{-1}$, KCl 27 $mmol.L^{-1}$, Na_2HPO_4 80 $mmol.L^{-1}$, KH_2PO_4 20

$mmol.L^{-1}$, and Na_2-EDTA 10 $mmol.L^{-1}$; pH = 7.5) during 16h, at 4°C in the dark. Samples

were then centrifuged (10 min, 3,220 g), and 1 mL of the supernatant was analyzed with a

spectrophotometer (CARY-60, Agilent) at 565, 620, and 650 nm, corrected by subtracting the

value recorded at 750 nm (< 0.01 absorbance unit). The values reported as concentration

$(mg.g^{-1})$ of C-phycocyanin and allophycocyanin (absence of phycoerythrin) were obtained

with calculus according to de Marasc and Houmard[27].

Lipophilic pigments of the 93 strains were analyzed by high performance liquid

chromatography (HPLC). Complete procedure is detailed in supplementary material and

methods. The relative abundance of each pigment (%) was calculated from its respective

concentration in the sample (mg.g^{-1} dw for chlorophyll *a* derivative, and μg.g^{-1} dw for other

pigments).

Fatty acids

Ten mg of freeze-dried biomass were used for the extraction of fatty acids on 91 strains (all

93 except *L. platensis* PMC 1041.18 and PMC 1044.18). Prior to extraction, an internal

standard (tricosanoic acid: 23:0) was added in each sample. Lipids were extracted according

to the protocol of Bligh and Dyer[29] as modified by Meziane et al.[30]. Complete procedure is

detailed in supplementary material and methods. Fatty acids were identified with a mass

spectrometer (Agilent 5977B GC/MSD) according to the comparison of retention times of

commercial fatty acid standards (Supelco 37). We then reported the values as % of total FA

and concentration (mg.g^{-1} dw).

Mass spectrometry

The intracellular metabolite contents of 88 strains (all 93 except *L. platensis* PMC 1287.21,

PMC 1288.21, PMC 1298.21, PMC 1300.21, and PMC 1302.21) were analyzed as previously

described[31]. Complete procedure is detailed in supplementary material and methods.

Intracellular metabolites contents were analyzed using an electrospray ionization hybrid

quadrupole time-of-flight (ESI-QqTOF) high-resolution mass spectrometer (Compact,

Brucker, Bremen, Germany) in the range 50–1500 *m/z*. Metabolite annotations were made

from MS2 data by generating a molecular network for the comparison of fragmentation

profiles using the MetGem software (version 1.3.6) with GNPS algorithm (Supplementary

figure S1).

Statistical analyses

The MetaboAnalyst 5.0 platform was used to perform data matrix normalization: Pareto for

ions and annotated ions matrices; mean-centered and divided by the standard deviation of

each variable for fatty acids and pigments matrices to better visualize the differences on less

concentrated molecules.

All three matrices (annotated ions, fatty acids, and pigments) were combined in a single

matrix further called "total metabolite matrix". The weight of each group of variables was

adjusted to avoid the analysis being influenced by the largest group. The weights are identical

for the variables of the same group.

Clade based on presence/absence of coding gene sequences was obtained with dendrogram

(Jaccard's dissimilarity index, Ward reconstruction, Silouhette analysis to find the optimal

number of clusters) using software R (version 1.4.1103).

A hierarchical clustering on principal components (HCPC) using Euclidian distance and a

principal component analysis (PCA) were performed to find different groups of strain using

software R (version 1.4.1103). Analysis of similarities (ANOSIM) was performed on the total

metabolite matrix to test differences between factors. Multiple correspondence analysis

(MCA) was performed to identify correlation between tested factors using software R

(version). For each tested factor, multiple factor analyses (MFA) and between class analyses

(BCA) were performed simultaneously (BC-MFA, code available at

https://github.com/Hubas-prog/BC-MFA) on the combined matrix, as describe in Michelet et

al.[32]. Percentage of total inertia explained (TIE) by the instrumental variable was

systematically calculated. A threshold based on the square cosine of the coordinates of the

variables was applied for highlighting the most discriminating variables with a square cosine

superior to 0.4 ($cos^2 > 0.4$). A Monte-Carlo test was systematically performed to test the

significance of the BCA ordination, using software R (version). For each factor, principal

component analyses (PCA) and heatmap analyses were performed on each of the three

matrices using the MetaboAnalyst 5.0 platform.

Differences between each class of factor have been research for each of the 215 molecules

with Kruskal-Wallis testes following by pairwise Wilcoxon testes. For total fatty acid means

comparison, normality was present (Shapiro-Wilk test, $p > 0.05$) and homogeneity of the

variances was not present (Bartlett test, $p < 0.05$), thus a Welch test was performed. All testes

were run using software R (version).

Results

Pigment composition

Fourty photosynthetic pigments were detected and quantified by HPLC-DAD in the 93

analyzed *L. platensis* strains: two phycobiliproteins (C-phycocyanin and allophycocyanin), 12

chlorophylls or derivatives (chlorophylls *a*, *d*, and pheophytins), and 26 carotenoids (five

carotenes, four carotenoid glycosides, six xanthophylls, one unknown carotene, and ten

unknown carotenoids) (Figure 1A; Supplementary Table S2). A mean of 27.9 ± 1.8 pigments

was detected per strain, ranging from 20 to 32 for the strains *L. platensis* PMC 1223.20, PMC

1278.20 and PMC 1279.20, and *L. platensis* PMC 1041.18, respectively (Supplementary

Table 2). 12 pigments were detected in all strains: C-phycocyanin, allophycocyanin,

chlorophyll *a,* chlorophyll *a* allomer, chlorophyll *a* epimer, two pheophytins *a*, β,φ-carotene,

β,ε-carotene-like, zeaxanthin, echinenone, echinenone/cantaxanthin-like, and two unknown

carotenoids (UC10, and UC9) (Supplementary Table S3). The most abundant pigments were

C-phycocyanin ($64.5 \pm 3.1\%$ of total pigment composition) and allophycocyanin ($25 \pm 1.6\%$

of total pigment composition). The chlorophyll *a* allomer were the most abundant form of

chlorophyll *a* in 90 out of 93 strains, representing $6.6 \pm 2.3\%$ of total pigment composition.

The three other strains (*L. platensis* PMC 1223.20, PMC 1278.20, and PMC 1279.20) had the

usual form of chlorophyll *a* as main chlorophyll *a*, representing 13.6 ± 2.3% of total pigment

composition (Supplementary Table 2). Carotenoids represented 1.19 ± 0.6% of total pigment

composition. The most abundant carotenoids were zeaxanthin (34.2 ± 5.5% of total carotenoid

composition), myxoxanthophyll (14.9 ± 5.6% of total carotenoid composition), β,β-carotene

(9.9 ± 1.3% of total carotenoid composition), and UC9 (425, 448, 470nm) (8.7 ± 3.6% of total

carotenoid composition) (Supplementary Table S2).

Fatty acid composition

Fourteen fatty acids were detected and quantified by GC-FID and GC-MS in the 91 analyzed

L. platensis strains. A mean of 52.32 ± 23.77 mg fatty acids.g^{-1} dry weight (dw) was

quantified, ranging from 13.78 to 110.59 mg.g^{-1} for the strains *L. platensis* PMC 1259.20 and

L. platensis PMC 894.15 respectively (Figure 1B; Supplementary Table S3). Among the 14

fatty acids detected, three were saturated fatty acid (14:0, 16:0, and 18:0), two were branched

fatty acids (15:0iso, and 15:0anteiso), four were mono-unsaturated fatty acids (16:1ω7,

16:1ω9, 18:1ω7, and 18:1ω9), and five were poly-unsaturated fatty acids (16:2ω6, 18:2ω6,

18:3ω6, 20:2ω6, and 20:3ω6) (Supplementary Table 1). Five fatty acids represented > 90% of

the total fatty acids: 16:0 (38.6 ± 1.8%), 18:3ω6 (24.6 ± 2.1%), 18:2ω6 (17.1 ±1.5%), 16:1ω9

(6.2 ± 1.1%), and 16:1ω7 (5.6 ± 0.8%) (Supplementary Table S3).

Metabolite composition by LC-MS

A total of 2,851 ions were detected by HPLC-MS in the 88 analyzed *L. platensis* strains. A

mean of 1,311.7 ± 175.1 ions was detected, ranging from 1,023 to 1,686 for the strains *L.*

platensis PMC 1044.18 and *L. platensis* SAG 85.79, respectively (Supplementary Table S4).

1013 ions (35.5%) had an area intensity superior a threshold of 5,000 count in single MS and

were analyzed by HPLC-MS/MS. Among them, 221 ions (15.9% of fragmented ion, and

5.6% of total ions) have been clustered and annotated with MetGem and GNPS (*e.g.*

phospholipids, saccharides, peptides, …) and 25 have been annotated at the molecular levels

(*e.g.* ergothioneine, glutathione, biopterin, nicotinamide adenine dinucleotide) (Figure 1C;

Supplementary Table S4), whereas other were annotated according to MS/MS fragmentation

pattern similarity revealed by molecular networking (Supplementary Figure S1). Analyses of

the 2,851 total ions and the 221 annotated ions showed similar results (Data not shown). We

further focused only of the 221 annotated ions (Figure 1C).

Groups of strains

Based on fatty acids, pigments and annotated ions, a complete matrix further called "total

metabolite matrix" was constructed for analyses (see Materials and methods). Considering the

sampling site, Multiple Factor Analysis (MFA) and the Hierarchical Clustering on Principal

Components (HCPC) based on the total metabolite matrix highlighted three groups of strains

(Figure 2A and 2B, respectively). Those three groups were statistically different (Figure 2A,

ANOSIM $p < 0.001$) and were also similarly discriminate regarding both fatty acids, pigments

and annotated ions separately (Supplementary figure S2A-C). The strains isolated from

different sampling sites were clustered in different groups: Camargue region in group 1, Lake

Dziani Dzaha in group 3, and other sampling site in group 2 (Figure 2C). These analyses

showed that strains coming from the lake Natron, Peru, and from the three unknown locations

were grouped together in group 3 (Figure 2A-B). For further analyses, all those strains were

put in a new sampling site group named "Others". The different environmental factors were

correlated to each of the three groups. Group 1 was correlated to the range of pH of 7-8,

salinity of 10-35‰, brackish habitat, and years of isolation 2019 and 2020. Group 3 was

correlated to the range of pH of 9-11, salinity of 35-70‰, alkaline-saline habitat, and years of

isolation 2010 and 2014. Group 2 was correlated to other pH, salinity, habitat, and year of

isolation prior to 2000 (Figure 2 C-D). The three groups highlighted by the HCPC that

corresponded to the sampling sites were used for further analyses.

Sampling sites

MFA and BCA analyses were done considering the sampling site and the three groups that

were previously clearly separated (Figure 3A-B). The primary axis of the BCA analysis

accounted for 63.1% of the variability. It primarily distinguished the *L. platensis* samples

taken at Dziani (group 3) from the others, based on their chemical composition (Figure 3B).

The second axis explained 37% of total inertia and mainly opposed HCPC group 2 from all

other strains (Figure 3B). Among fatty acids, pigments, and annotated ions, 31 most

discriminants were defined with a $cos^2 > 0.4$ (Figure 3C): two fatty acids, nine pigments, and

20 annotated ions (26 related to axis 1, and five to axis 2). In total, 126 molecules showed

significative difference between at least two groups (Kruskal-Wallis test, $p < 0.001$). The two

fatty acids were the PUFAs with the longest carbon chains, 20:2ω6 and 20:3ω6 that were

significantly more present in the strains from lake Dziani Dzaha in concentration and

proportion of total fatty acids (Figure 4, Kruskal-Wallis test, $p < 0.001$). Based on fatty acids

only, two groups of 42 and 49 strains have been discriminated by hierarchal clustering, not

related to any of the environmental factor (Supplementary Figure 2B). They significantly

differed by the total fatty acid biomass (30 ± 8 mg.g$^-$and 71 ± 15 mg.g^{-1}, respectively).

Six of the pigments were only present in the strains from lake Dziani Dzaha (Figures 1A and

4, Kruskal-Wallis test, $p < 0.001$). Among them, three were recognized as resembling

caloxanthin and nostoxanthin and an analog of β,β-carotene , and three were unknown

carotenoids. The strains from lake Dziani Dzaha did not possessed the usual form of β,β-

carotene, that was only present in the strains from Camargue and other locations. The

allophycocyanin were significantly more present in the strains from other locations (Figure 4,

Kruskal-Wallis test, p < 0.001). Among the 20 most discriminant annotated metabolites

(Figure 5), 13 were significantly more abundant in the strains from lake from lake Dziani

Dzaha, including four saccharides (MW = 254.1022, 596.2184, 684.2349 Da, and sucrose),

three little peptides of 2 or 3 amino acids (MW = 373.2214, 359.2057, and 260.1376 Da), two

phospholipids (MW = 467.304, and 479.3041 Da), one lipid (MW = 455.3042 Da), and two

unsaturated fatty acid glycerol annotated as monoolein and alpha-linolenoyl-glycerol

(Kruskal-Wallis test, p < 0.001). Three molecules were significantly less abundant in the

strains from lake Dziani Dzaha: two saccharides (MW = 285.1065, and 268.0803 Da), and S-

methylglutathione (Kruskal-Wallis test, p < 0.001). Four molecules were significantly more

present in the strains from other locations: two peptides (MW = 359.1869, and 373.1986 Da),

one fatty acid ester (MW = 588.3765 Da), and oxidized glutathione (Kruskal-Wallis test, p <

0.001).

Gene content

The analysis of the presence/absence of coding gene sequences of the strains with a

sequenced genome showed two groups of strains (I and II). The first group was divided in two

subgroups (I.1 and I.2) (Supplementary table S1; Supplementary figure S3). Clade I.1

contained 23 strains from the Camargue region. Clade I.2 contained four strains from Lake

Dziani Dzaha, two strains from Lake Natron (*L. platensis* PMC 1042.18 and SAG 85.79), and

the strain *L. platensis* PCC 8005. The Clade II contained the last three strains (*L. platensis*

PCC 7345, PMC 289.06, and Paracas R14).

BC-MFA analyses were done considering the coding gene clade and the three clades (I.1, I.2

and II) were clearly separated (Supplementary Figure 4A-B). The first axis of the BCA

analysis explained 70.5% and the second axis explained 29.5% of the variance. Chemical

composition of all three clades were separated by the first axis of the BC-MFA, while all

clade II was separated from clades I.1 and I.2 by the second axis (Supplementary Figure 4B.

Among fatty acids, pigments, and annotated ions, 18 most discriminant molecules were

defined with a $\cos^2 > 0.4$ (Supplementary Figure 4C): seven fatty acids, seven pigments, and

four annotated ions (17 related to axis 1, and one to axis 2). In total, 63 molecules showed

significative difference between at least two groups (Kruskal-Wallis test, $p < 0.01$).

The seven fatty acids, as well as the seven pigments were less abundant in the clade I.1

(Supplementary Figure 5, Kruskal-Wallis test, $p < 0.01$). Among the four most discriminant

annotated ions, two saccharides (MW = 536.1602 and 430.133 Da) were significantly more

abundant in the strains of clade I.1. The two other annotated ions were significantly more

abundant in the strains of clade II: oxidized glutathione and a fatty acid ester (MW =

588.3765 Da) (Supplementary Figure 5, Kruskal-Wallis test, $p < 0.01$).

Discussion

Analyses of pigment, lipid and small metabolite contents of various *Limnospira platensis*

strains cultivated under identical laboratory conditions provide unprecedent data that offers a

plain answer to the foremost question: is the diversity of these molecules driven by selective

pressures on *Limnospira platensis* (clade I or II)[20] and/or by the environment from which the

strains are isolated? Our results clearly show that the sampling sites characterized by diverse

environmental conditions (*e.g.* salinity, pH, light), are the variables explaining the distribution

of *Limnospira platensis* in three distinct chemotype groups. The molecules that most

discriminate these three groups of the *Limnospira platensis* strains studied, under the standard

conditions of lab cultures, are sugars, lipids, peptides, photosynthetic pigments, and

antioxidant molecules. Interestingly, these molecule enrichments might constitute

physiological adaptation traits to the local conditions encounter in their respective sampling

environments in relation to salinity, light and oxidative stress.

Osmoregulation and salinity resistance

The ponds of the Camargue in France and Lake Dziani Dzaha in Mayotte are environments

with different salinities. In the brackish ponds of Camargue, close to the Mediterranean Sea,

salinity varies between 12-35‰. Lake Dziani Dzaha is a thalassohaline lake. Here, salinity is

much higher, reaching up to 70‰[33]. Strains from Lake Dziani Dzaha have greater sugar

diversity, have more long-chain polyunsaturated fatty acids compared with strains isolated

from other habitats under our experimental conditions. They are also the only ones to produce

nostoxanthin and caloxanthin. This specific chemical signature of strains from Lake Dziani

Dzaha testifies to the adaptation of the *Limnospira platensis* species to resist to higher salinity

in this specific environment.

In cyanobacteria in general, several families of molecules are involved in osmoregulation and

detoxification of stress due to high salinity[34-35]. Cyanobacteria produce various sugars (*e.g.*

glucose, sucrose, trehalose or glycine betaine) to prevent the osmotic stress caused by high

salinity[36]. In *Synechococcus*, long-chain polyunsaturated fatty acids help combat the oxidative

stress caused by high salinity as they appear to be produced to a greater extent with increasing

salinity[37]. Interestingly, it has also been shown that certain carotenoids, nostoxanthin and

caloxanthin, are produced by a bacterium of the genus *Sphingomonas* that appears resistant to

high salinity[38]. In addition, the nostoxanthin and caloxanthin-producing bacterium

Sphingomonas nostxanthinifaciens may have a role in reducing the amount of reactive oxygen

species (ROS) in *Arabidopsis thaliana* when the two organisms are co-cultured in high salt

concentrations[39]. Taken together, these finding support the idea that sugar and carotenoid

molecules might support high salinity adaptation process in *Limnospira platensis*.

Light intensity

Facing high light, cyanobacteria modify chlorophyll, phycobiliprotein and carotenoid

composition to improve electron transfer and manage excess energy[40-42]. In *Synechococcus*,

various carotenoids are crucial for high light survival. Strains missing carotenoid biosynthesis

genes (*cru* genes) have lower growth rates and higher ROS levels compared to carotenoid-

producing strains[40].

The different *Limnospira platensis* strains relied on a wide diversity of chlorophylls (n = 5-

12) and carotenoids (n = 13-18) that could accordingly be involved in photo-pigment

adaptation. One of the most striking results is the presence of six exclusive carotenoids that

appear exclusively in strains from Lake Dziani Dzaha. This lake has a mean surface irradiance

of 2,000 µmol photon.s^{-1}.m^{-2}, which decreases by 99% in the first 20 cm of surface water to

around 20 µmol photon.s^{-1}.m^{-2}.[33] *Limnospira* living in the photic zone of the lake are therefore

subjected to both high and strong variations in light intensity over shallow depths. We found

18 carotenoid pigments that have an unknown chemical structure (putative analogues of other

pigments, n = 7, or totally unknown pigment, n = 11). This observation illustrates the ability

of *Limnospira* to diversify its pigment content in response to a given environment. The great

diversity of pigments produced by *Limnospira platensis* was shown by the characterization of

up to 48 hydrophobic pigments (22 carotenoids and 26 chlorophylls) by LC-APCI-FT-ICR-

MS[43], some of which could very likely correspond to those unknown pigments observed in

our study.

Oxidative stress

In a general manner, both high salinity, light and UV exposures are interdependently involved

in ROS production and subsequent cellular oxidative stress[44]. In addition to primary

mechanisms listed above, aiming at reducing the adverse effect of oxidative stress,

cyanobacteria also produce a large variety of other molecules (other than carotenoids and

long-chain polyunsaturated fatty acids). These molecules are specifically involved in

buffering ROS production and combating oxidative stress deleterious effects, such as

ergothioneine, glutathione, biopterins and their derivatives[14,45-46]. The cellular extracts of all

Limnospira platensis strains presently investigated contained significant amounts of each of

these molecules, that might constitute a set of molecular resource for this organism to deal

with remarkable local cellular stressors.

***Limnospira platensis* presents a peculiar ecological strategy**

Cyanobacteria have colonized many types of habitats (*e.g.* oceanic waters, fresh waters,

alkaline waters, benthic or pelagic, terrestrial, symbiotic)[2]. Aquatic species have for example

developed different ecological strategies to adapt to their environment. Some cyanobacteria

live in communities. These include benthic cyanobacteria, which form biofilms (*e.g.*

Phormidium, *Lyngbya*, *Oscillatoria*)[47], and pelagic cyanobacteria, which can form blooms

(*e.g. Microcystis*, *Aphanizomenon*, *Raphidiopsis, ...*)[48-50]. The genomes of these cyanobacteria

are remarkably large (> 5Mb) and contain many BGCs[51-53], producing various bio-active

secondary metabolites (including cyclic peptides, lipo-peptides, alkaloids, macrolides,

polyketides, …) that enable cyanobacteria to interact with their microbial community (*e.g.*

anti-grazing, anti-algal, communication blocking, competition for resources)[5,54]. However, the

production of these specific molecules has a high energy cost for the producing cells, as there

are involving large gene sets and the synthesis of many enzymes[55]. In contrary, some marine

picocyanobacteria (*e.g. Prochlorococcus*, *Synechococcus*) are exclusively dominating one

type of habitat (*i.e.* the worldwide oligothrophic seas) and exhibit a unique diversity of

photosynthetic pigments, allowing them to exploit a wide range of light micro-niches[56-57]. The

genomes of these cyanobacteria are remarkably small (1.5-2.5Mb) and contain no BGCs[51,58].

In these cyanobacteria, adaptation to an ecological niche is encoded in both core and flexible

parts of the genomes[59-60].

For *Limnospira platensis*, we denote a potential third strategy. *Limnospira platensis* genomes

are relatively large, ranging between 5.5 and 6.5Mb[20,61], being. comparable to the genome

size of community-dwelling cyanobacteria proliferating in eutrophic waters. Yet, they

remarkably contain no or only one BGC of secondary metabolites (potentially involved in the

biosynthesis of cyanobactins called Arthrospiramides), as the specialized marine

picocyanobacteria *Synechococcus* and *Prochlorococcus*[51,62-63]. Unlike marine *Synechococcus*

and *Prochlorococcus*, *Limnospira platensis* proliferates in many different habitat types:

freshwater, brackish water, alkaline water, and saline alkaline water[20,64], and *Limnospira*

platensis strains belonging to the same phylogenetic clade do not share similar habitats[20].

Moreover, in lakes where phytoplanktonic communities are dominated by *Limnospira*

platensis, this species frequently generates persistent proliferations. These blooms are often

accompanied by minimal surrounding diversity (except de picoeukaryote *Picocystis*

salinarum)[21,63], serving as a distinctive indicator of extreme environments (*e.g.* hypersaline

conditions, high light coupled with high turbidity, elevated temperatures, high pH levels, and

high sulfur contents). In such environments, only a limited number of micro-organism clades

can develop. This absence of surrounding microbial diversity, comprising potential

competitors, could explain the absence of BGCs in the *Limnospira platensis* genomes, as it

has less phototroph organisms with which to interact through allelopathic compounds

production, as various cyanobacteria from high diversity freshwater temperate lakes do[51].

Interestingly, such an ecological strategy could support the high biomass production observed

both *in situ* and *in vitro*, as lab culture conditions present many similarities (in terms of both

biotic and abiotic factors) with those of the extreme environment in which *Limnospira*

proliferates. Indeed, it has been shown by transcriptomics that the essential transcribed

functions expressed *in vitro* are related to primary metabolism (metabolism of sugar, fatty

acids, nitrogen, CO_2, photosynthesis, and antioxidant processes)[64].

Interestingly, the genomic investigation of *Limnospira platensis* strains shows that both well-

conserved (core) and flexible (pan) genome parts discriminate strains according to their

respective origin location (Supplementary figure S3). This latter variable part of the genome

is remarkably rich in transposable and regulatory elements and contains various genes, most

of which have as yet unknown molecular function, would be involved in regulatory processes

(transcription, translation, post-transcriptional, or post-translational), traducing the evolutive

dynamic character of this part of the genome[20]. The regulatory genes would result in slight

modifications of primary metabolism to resist different environmental stresses (*e.g.* the

production of osmoregulatory, antioxidant and photoprotective suites). Thus, one can

supposed that either punctual mutations on key metabolic genes or variation in gene content

of this flexible part of of *Limnospira platensis* strain genomes may enable genotype local

adaptation to the environmental conditions of a habitat. So far, the relationship between small

metabolite content (*e.g.* anti-oxidants, saccharides and carotenoids) - chemotype - and niche

partitioning - ecotype - still remained largely undetermined due to the lack of distinctive

genetic marker associated with the various pigment production. In addition, such an

assumption also required to be now further experimentally validated.

Conclusion

Taken together, these ecological and evolutionary concerns offer new comprehension on the

mechanisms that support the remarkable success encountered by *Limnospira* in natural as well

as in lab or industrial environments. One step further would be now to try to experimentally

identify the genetic and phenotypic traits that are defining the different *Limnospira* ecotypes

and to determine whether *Limnospira* clade I.1, I.2 and II genotypes can colonize similar

ecological niches, as this taxon exhibits ubiquitous geographic distribution, or represent

already well distinct ecologically and functionally distinguishable taxa.

Also, key questions remain on the understanding of the mechanism that allow *Limnospira in*

situ and *in vitro* blooms to avoid density-dependent regulations by cyanophages, as those

encountered by other blooming cyanobacteria[65]. Such adaptive mechanisms would help

explaining how *Limnospira platensis* can form permanent blooms in many environments. To

this end, two distinct hypotheses can be further tested: Either *Limnospira platensis* genome

present a mechanism to protect itself from phage lyses, or the fraction of the population

potentially lysed by phages would be replaced by rapidly dividing *Limnospira platensis*.

Author Contributions: Conceptualization, T.R., J.-P.C., L.V., C.B. and B.M.; methodology,

T.R., C.H., M.C., C.Y., B.M.; investigation, T.R., C.H., T.M.; resources, C.D.; writing-

original draft preparation, T.R., C.B. and B.M.; writing—review and editing, all authors. All

authors have read and agreed to the published version of the manuscript.

Funding: This research was funded by ANRT, through a convention Cifre n° 2020/058

awarded to the Company Algama employing T. Roussel for a PhD thesis.

Acknowledgments: We would like to thank the UMR 7245 MCAM, Muséum National

d'Histoire Naturelle, Paris, France for laboratories facilities and Algama for funds. The MS

spectra were acquired at the Plateau technique de spectrométrie de masse bio-organique,

Muséum National d'Histoire Naturelle, Paris, France. The authors thank the Pasteur

Collection of Cyanobacteria, the Culture Collection of Algae at Göttigen University, and

Gilles Planchon for sending strains.

Conflicts of Interest: The authors declare no conflict of interest.

References

1-   Reynolds, C. S. (2006). *The ecology of phytoplankton*. Cambridge University Press.

2-   Whitton, B.A. (ed) (2012) Ecology of Cyanobacteria II: Their Diversity in Space and

507   Time. Dordrecht: Springer Netherlands. Available online: https://doi.org/10.1007/978-

508   94-007-3855-3.

509   3-   Whitton, B.A. & Potts, M. (eds) (2007) The Ecology of Cyanobacteria: Their Diversity

510   in Time and Space. Dordrecht: The Netherlands, Springer Science & Business Media.

511   4-   Leflaive, J. & Ten-Hage, L. (2007) Algal and cyanobacterial secondary metabolites in

512   freshwaters: a comparison of allelopathic compounds and toxins. Freshwater Biology,

513   52(2), 199–214. Available online: https://doi.org/10.1111/j.1365-2427.2006.01689.x.

514   5-   Demay, J., Bernard, C., Reinhardt, A. & Marie, B. (2019) Natural Products from Cya-

515   nobacteria: Focus on Beneficial Activities. Marine Drugs, 17(6), 320. Available online:

516   https://doi.org/10.3390/md17060320.

517   6-   Nuhu, A. A. (2013). Spirulina (Arthrospira): An important source of nutritional and

518   medicinal compounds. *Journal of Marine biology*, *2013*, 1-8.

519   7-   Shao, W., Ebaid, R., El-Sheekh, M., Abomohra, A., & Eladel, H. (2019). Pharmaceuti-

520   cal applications and consequent environmental impacts of Spirulina (Arthrospira): An

521   overview. *Grasas y Aceites*, *70*(1), e292-e292.

8-   Vonshak, A. & Tomsaelli, L. (2000) Arthrospira (Spirulina): Systematics and Ecolo-

physiology. In Whittons, B.A. & Potts, M. The ecology of Cyanobacteria: Their Diver-

sity in Time and Space.

9-   Vonshak, A. (ed) (1997) Spirulina platensis (Arthrospira): Physiology, Cell-biology,

and Biotechnology. London, Bristol, PA: Taylor & Francis.

10-   Avila-Leon, I., Chuei Matsudo, M., Sato, S. & De Carvalho, J.C.M. (2012) Arthrospira

platensis biomass with high protein content cultivated in continuous process using urea

as nitrogen source: High protein biomass by continuous process with urea. Journal of

Applied Microbiology, 112(6), 1086–1094. Available online:

https://doi.org/10.1111/j.1365-2672.2012.05303.x.

11-   Ötleş, S., & Pire, R. (2001). Fatty acid composition of Chlorella and Spirulina microal-

gae species. *Journal of AOAC international*, *84*(6), 1708-1714.

12-   Martinez Andrade, K., Lauritano, C., Romano, G. & Ianora, A. (2018) Marine Microal-

gae with Anti-Cancer Properties. Marine Drugs, 16(5), 165. Available online:

https://doi.org/10.3390/md16050165.

13-   Edelmann, M., Aalto, S., Chamlagain, B., Kariluoto, S. & Piironen, V. (2019) Ribofla-

vin, niacin, folate and vitamin B12 in commercial microalgae powders. Journal of Food

Composition and Analysis, 82, 103226. Available online:

https://doi.org/10.1016/j.jfca.2019.05.009.

14-   Pfeiffer, C., Bauer, T., Surek, B., Schmig, E. & Gründemann, D. (2011) Cyanobacteria

produce high levels of ergothioneine. Food Chemistry, 129(4), 1766–1769. Available

online: https://doi.org/10.1016/j.foodchem.2011.06.047.

15-   Markou, G. (2014) Effect of Various Colors of Light-Emitting Diodes (LEDs) on the

Biomass Composition of Arthrospira platensis Cultivated in Semi-continuous Mode.

Applied Biochemistry and Biotechnology, 172(5), 2758–2768. Available online:
https://doi.org/10.1007/s12010-014-0727-3.

16- Bezerra, R.P., Matsudo, M.C., Sato, S., Perego, P., Converti, A. & Carvalho, J.C.M.D.
(2012) Effects of photobioreactor configuration, nitrogen source and light intensity on
the fed-batch cultivation of Arthrospira (Spirulina) platensis. Bioenergetic aspects. Bi-
omass and Bioenergy, 37, 309–317. Available online:
https://doi.org/10.1016/j.biombioe.2011.11.007.

17- Manirafasha, E., Murwanashyaka, T., Ndikubwimana, T., Rashid Ahmed, N., Liu, J.,
Lu, Y., Zeng, X., Ling, X. & Jing, K. (2018) Enhancement of cell growth and phycocy-
anin production in Arthrospira (Spirulina) platensis by metabolic stress and nitrate fed-
batch. Bioresource Technology, 255, 293–301. Available online:
https://doi.org/10.1016/j.biortech.2017.12.068.

18- Tanaka, K., Kishi, M., Assaye, H. & Toda, T. (2020) Low temperatures in dark period
affect biomass productivity of a cyanobacterium Arthrospira platensis. Algal Research,
52, 102132. Available online: https://doi.org/10.1016/j.algal.2020.102132.

19- Sili, C., Torzillo, G. & Vonshak, A. (2012) Arthrospira (Spirulina). In Whitton, B.A.
(ed) Ecology of Cyanobacteria II: Their Diversity in Space and Time. Dordrecht:
Springer Netherlands. Available online: https://doi.org/10.1007/978-94-007-3855-3.

20- Roussel, T., Halary, S., Duval, C., Piquet, B., Cadoret, J. P., Vernès, L., ... & Marie, B.
(2023). Monospecific renaming within the cyanobacterial genus Limnospira (Spirulina)
and consequences for food authorization. *Journal of Applied Microbiology*, *134*(8),
lxad159.

21- Cellamare, M., Duval, C., Drelin, Y., Djediat, C., Touibi, N., Agogué, H., ... & Bernard,
C. (2018). Characterization of phototrophic microorganisms and description of new cy-

anobacteria isolated from the saline-alkaline crater-lake Dziani Dzaha (Mayotte, Indian

Ocean). *FEMS microbiology ecology*, *94*(8), fiy108.

22- Rossello-Mora, R., Lucio, M., Pena, A., Brito-Echeverría, J., Lopez-Lopez, A., Valens-

Vadell, M., ... & Schmitt-Kopplin, P. (2008). Metabolic evidence for biogeographic iso-

lation of the extremophilic bacterium Salinibacter ruber. *The ISME journal*, *2*(3), 242-

253.

23- Kujawinski EB, Braakman R, Longnecker K, Becker JW, Chisholm SW, Dooley K,

Kido Soule MC, Swarr GJ, Halloran K. Metabolite diversity among representatives of

divergent *Prochlorococcus* ecotypes. mSystems. 2023 Oct 26;8(5):e0126122. doi:

10.1128/msystems.01261-22. Epub 2023 Oct 10. Erratum in: mSystems. 2024 Mar

19;9(3):e0012024.

24- Hamlaoui, S., Yéprémian, C., Duval, C., Marie, B., Djédiat, C., Piquet, B., Bernard, C.

& Duperron, S. (2022) The Culture Collection of Cyanobacteria and Microalgae at the

French National Museum of Natural History: A Century Old But Still Alive and Kick-

ing! Including in Memoriam: Professor Alain Cout.. Cryptogamie, Algologie, 43(3).

Available online: https://doi.org/10.5252/cryptogamie-algologie2022v43a3.

25- Yéprémian, C., Catherine, A., Bernard, C., Congestri, R., Elersek, T. & Pilkaityte, R.

(2017a) Phycocyanin Extraction and Determination. In Meriluoto, J., Spoof, L., &

Codd, G.A. (eds) Handbook of Cyanobacterial Monitoring and Cyanotoxin Analysis.

1st edition. Wiley, 335–338. Available online:

https://doi.org/10.1002/9781119068761.ch35.

26- Yéprémian, C., Catherine, A., Bernard, C., Congestri, R., Elersek, T. & Pilkaityte, R.

(2017b) Chlorophyll a Extraction and Determination. In Meriluoto, J., Spoof, L., &

Codd, G.A. (eds) Handbook of Cyanobacterial Monitoring and Cyanotoxin Analysis.

1st   edition.   Wiley,   331–334.   Available   online:

https://doi.org/10.1002/9781119068761.ch34.

27- De Marsac, N. T., & Houmard, J. (1988). [34] Complementary chromatic adaptation:

physiological conditions and action spectra. In *Methods in enzymology* (Vol. 167, pp.

318-328). Academic Press.

28- Brotas, V. & Plante-Cuny, M.R. (2003) The use of HPLC pigment analysis to study

microphytobenthos communities. Acta Oecologica, 24(1), S109-S115. Available online:

https://doi.org/10.1016/S1146-609X(03)00013-4.

29- Bligh EG, Dyer WJ. A rapid method of total lipid extraction and purification. Can J

Biochem Physiol. 1959 Aug;37(8):911-7. doi: 10.1139/o59-099. PMID: 13671378.

30- Meziane, T., Lee, S.Y., Mfilinge, P.L., Shin, P.K.S., Lam, M.H.W. & Tsuchiya, M.

(2007) Inter-specific and geographical variations in the fatty acid composition of man-

grove leaves: implications for using fatty acids as a taxonomic tool and tracers of organ-

ic   matter.   Marine   Biology,   150(6),   1103–1113.   Available   online:

https://doi.org/10.1007/s00227-006-0424-z.

31- Le Moigne, D., Demay, J., Reinhardt, A., Bernard, C., Kim Tiam, S., & Marie, B.

(2021). Dynamics of the Metabolome of Aliinostoc sp. PMC 882.14 in Response to

Light and Temperature Variations. *Metabolites*, *11*(11), 745.

32- Michelet, C., Zeppilli, D., Hubas, C., Baldrighi, E., Cuny, P., Dirberg, G., Militon, C.,

Walcker, R., Lamy, D., Jézéquel, R., Receveur, J., Gilbert, F., Houssainy, A.E., Dufour,

A., Heimbürger-Boavida, L.-E., Bihannic, I., Sylvi, L., Vivier, B. & Michaud, E. (2021)

First Assessment of the Benthic Meiofauna Sensitivity to Low Human-Impacted Man-

groves   in   French   Guiana.   Forests,   12(3),   338.   Available   online:

https://doi.org/10.3390/f12030338.

33- Leboulanger, C., Agogué, H., Bernard, C., Bouvy, M., Carré, C., Cellamare, M., Duval,

C., Fouilland, E., Got, P., Intertaglia, L., Lavergne, C., Le Floc'h, E., Roques, C. &

Sarazin, G. (2017) Microbial Diversity and Cyanobacterial Production in Dziani Dzaha

Crater Lake, a Unique Tropical Thalassohaline Environment. PLOS ONE. Edited by J.-

F.   Humbert,   12(1),   e0168879.   Available   online:

https://doi.org/10.1371/journal.pone.0168879.

34- Hagemann M. Molecular biology of cyanobacterial salt acclimation. FEMS Microbiol

Rev. 2011 Jan;35(1):87-123. doi: 10.1111/j.1574-6976.2010.00234.x.

35- Duval, C., Thomazeau, S., Drelin, Y., Yepremian, C., Bouvy, M., Couloux, A., ... &

Bernard, C. (2018). Phylogeny and salt-tolerance of freshwater Nostocales strains: con-

tribution to their systematics and evolution. *Harmful algae*, *73*, 58-71.

36- Mackay, M.A., Norton, R.S. & Borowitzka, L.J. (1984) Organic Osmoregulatory So-

lutes   in   Cyanobacteria.   Microbiology,   130(9),   2177–2191.   Available   online:

https://doi.org/10.1099/00221287-130-9-2177.

37- Verma, E., Singh, S., Niveshika & Mishra, A.K. (2019) Salinity-induced oxidative

stress-mediated change in fatty acids composition of cyanobacterium Synechococcus sp.

PCC7942. International Journal of Environmental Science and Technology, 16(2), 875–

886. Available online: https://doi.org/10.1007/s13762-018-1720-0.

38- Kikukawa, H., Okaya, T., Maoka, T., Miyazaki, M., Murofushi, K., Kato, T., Hirono-

Hara, Y., Katsumata, M., Miyahara, S. & Hara, K.Y. (2021) Carotenoid Nostoxanthin

Production by Sphingomonas sp. SG73 Isolated from Deep Sea Sediment. Marine

Drugs, 19(5), 274. Available online: https://doi.org/10.3390/md19050274.

39- Jiang, L., Seo, J., Peng, Y., Jeon, D., Lee, J.H., Kim, C.Y. & Lee, J. (2023) A nostoxan-

thin-producing bacterium, Sphingomonas nostoxanthinifaciens sp. nov., alleviates the

salt stress of Arabidopsis seedlings by scavenging of reactive oxygen species. Frontiers

in Microbiology, 14, 1101150. Available online:

https://doi.org/10.3389/fmicb.2023.1101150.

40- Zhu, Y., Graham, J.E., Ludwig, M., Xiong, W., Alvey, R.M., Shen, G. & Bryant, D.A.

(2010) Roles of xanthophyll carotenoids in protection against photoinhibition and oxi-

dative stress in the cyanobacterium Synechococcus sp. strain PCC 7002. Archives of

Biochemistry and Biophysics, 504(1), 86–99. Available online:

https://doi.org/10.1016/j.abb.2010.07.007.

41- Kumar, M., Kulshreshtha, J. & Singh, G.P. (2011) Growth and biopigment accumula-

tion of cyanobacterium Spirulina platensis at different light intensities and temperature.

Brazilian Journal of Microbiology, 42(3), 1128–1135. Available online:

https://doi.org/10.1590/S1517-83822011000300034.

42- Masamoto, K. & Furukawa, K. (1997) Accumulation of zeaxanthin in cells of the cya-

nobacterium, Synechococcus sp. Strain PCC 7942 grown under high irradiance. Journal

of Plant Physiology, 151(3), 257–261. Available online: https://doi.org/10.1016/S0176-

1617(97)80250-7.

43- Sommella, E., Conte, G. M., Salviati, E., Pepe, G., Bertamino, A., Ostacolo, C., ... &

Campiglia, P. (2018). Fast profiling of natural pigments in different spirulina (arthrospi-

ra platensis) dietary supplements by DI-FT-ICR and evaluation of their antioxidant po-

tential by pre-column DPPH-UHPLC assay. *Molecules*, *23*(5), 1152.

44- Khorobrykh, S., Havurinne, V., Mattila, H. & Tyystjérvi, E. (2020) Oxygen and ROS in

Photosynthesis. Plants, 9(1), 91. Available online:

https://doi.org/10.3390/plants9010091.

45- Noguchi, Y., Ishii, A., Matsushima, A., Haishi, D., Yasumuro, K., Moriguchi, T., Wada,

T., Kodera, Y., Hiroto, M., Nishimura, H., Sekine, M. & Inada, Y. (1999) Isolation of

Biopterin-α-glucoside from Spirulina (Arthrospira) platensis and Its Physiologic Func-

tion. Marine Biotechnology, 1, 207–210. Available online:

https://doi.org/10.1007/PL00011769.

46- Latifi, A., Ruiz, M. & Zhang, C.-C. (2009) Oxidative stress in cyanobacteria. FEMS

Microbiology Reviews, 33(2), 258–278. Available online:

https://doi.org/10.1111/j.1574-6976.2008.00134.x.

47- Quiblier, C., Wood, S., Echenique-Subiabre, I., Heath, M., Villeneuve, A. & Humbert,

J.-F. (2013) A review of current knowledge on toxic benthic freshwater cyanobacteria –

Ecology, toxin production and risk management. Water Research, 47(15), 5464–5479.

Available online: https://doi.org/10.1016/j.watres.2013.06.042.

48- Cirés, S. & Ballot, A. (2016) A review of the phylogeny, ecology and toxin production

of bloom-forming Aphanizomenon spp. and related species within the Nostocales (cya-

nobacteria). Harmful Algae, 54, 21–43. Available online:

https://doi.org/10.1016/j.hal.2015.09.007.

49- Jones, A.C., Monroe, E.A., Podell, S., Hess, W.R., Klages, S., Esquenazi, E., Niessen,

S., Hoover, H., Rothmann, M., Lasken, R.S., Yates, J.R., Reinhardt, R., Kube, M., Bur-

kart, M.D., Allen, E.E., Dorrestein, P.C., Gerwick, W.H. & Gerwick, L. (2011) Ge-

nomic insights into the physiology and ecology of the marine filamentous cyanobacte-

rium Lyngbya majuscula. Proceedings of the National Academy of Sciences, 108(21),

8815–8820. Available online: https://doi.org/10.1073/pnas.1101137108.

50- Harke, M.J., Steffen, M.M., Gobler, C.J., Otten, T.G., Wilhelm, S.W., Wood, S.A. &

Paerl, H.W. (2016) A review of the global ecology, genomics, and biogeography of the

toxic cyanobacterium, Microcystis spp. Harmful Algae, 54, 4–20. Available online:

https://doi.org/10.1016/j.hal.2015.12.007.

51- Shih, P.M., Wu, D., Latifi, A., Axen, S.D., Fewer, D.P., Talla, E., Calteau, A., Cai, F.,

Tandeau De Marsac, N., Rippka, R., Herdman, M., Sivonen, K., Coursin, T., Laurent,

T., Goodwin, L., Nolan, M., Davenport, K.W., Han, C.S., Rubin, E.M., Eisen, J.A.,

Woyke, T., Gugger, M. & Kerfeld, C.A. (2013) Improving the coverage of the cyano-

bacterial phylum using diversity-driven genome sequencing. Proceedings of the Nation-

al Academy of Sciences, 110(3), 1053–1058. Available online:

https://doi.org/10.1073/pnas.1217107110.

52- Leikoski, N., Liu, L., Jokela, J., Wahlsten, M., Gugger, M., Calteau, A., Permi, P., Ker-

feld, C.A., Sivonen, K. & Fewer, D.P. (2013) Genome Mining Expands the Chemical

Diversity of the Cyanobactin Family to Include Highly Modified Linear Peptides.

Chemistry & Biology, 20(8), 1033–1043. Available online:

https://doi.org/10.1016/j.chembiol.2013.06.015.

53- Halary, S., Duperron, S., Kim Tiam, S., Duval, C., Bernard, C. & Marie, B. (2023) In-

tra-population genomic diversity of the bloom-forming cyanobacterium, Aphani-

zomenon gracile, at low spatial scale. ISME Communications, 3(1), 57. Available

online: https://doi.org/10.1038/s43705-023-00263-3.

54- Leao, P.N., Engene, N., Antunes, A., Gerwick, W.H. & Vasconcelos, V. (2012) The

chemical ecology of cyanobacteria. Natural Product Reports, 29(3), 372. Available

online: https://doi.org/10.1039/c2np00075j.

55- Fischbach, M.A. & Walsh, C.T. (2006) Assembly-Line Enzymology for Polyketide and

Nonribosomal Peptide Antibiotics: Logic, Machinery, and Mechanisms. Chemical Re-

views, 106(8), 3468–3496. Available online: https://doi.org/10.1021/cr0503097.

56- Grébert, T., Dor., H., Partensky, F., Farrant, G.K., Boss, E.S., Picheral, M., Guidi, L.,

Pesant, S., Scanlan, D.J., Wincker, P., Acinas, S.G., Kehoe, D.M. & Garczarek, L.

(2018) Light color acclimation is a key process in the global ocean distribution of Syn-

echococcus cyanobacteria. Proceedings of the National Academy of Sciences, 115(9).

Available online: https://doi.org/10.1073/pnas.1717069115.

57-   Sanfilippo, J.E., Garczarek, L., Partensky, F. & Kehoe, D.M. (2019) Chromatic Accli-

mation in Cyanobacteria: A Diverse and Widespread Process for Optimizing Photosyn-

thesis. Annual Review of Microbiology, 73(1), 407–433. Available online:

https://doi.org/10.1146/annurev-micro-020518-115738.

58-   Biller, S.J., Berube, P.M., Berta-Thompson, J.W., Kelly, L., Roggensack, S.E., Awad,

L., Roache-Johnson, K.H., Ding, H., Giovannoni, S.J., Rocap, G., Moore, L.R. &

Chisholm, S.W. (2014) Genomes of diverse isolates of the marine cyanobacterium Pro-

chlorococcus.   Scientific   Data,   1(1),   140034.   Available   online:

https://doi.org/10.1038/sdata.2014.34.

59-   Biller, S.J., Berube, P.M., Lindell, D. & Chisholm, S.W. (2015) Prochlorococcus: the

structure and function of collective diversity. Nature Reviews Microbiology, 13(1), 13–

27. Available online: https://doi.org/10.1038/nrmicro3378.

60-   Doré, H., Guyet, U., Leconte, J., Farrant, G.K., Alric, B., Ratin, M., Ostrowski, M., Fer-

rieux, M., Brillet-Guéguen, L., Hoebeke, M., Siltanen, J., Le Corguillé, G., Corre, E.,

Wincker, P., Scanlan, D.J., Eveillard, D., Partensky, F. & Garczarek, L. (2023) Differ-

ential global distribution of marine picocyanobacteria gene clusters reveals distinct

niche-related adaptive strategies. The ISME Journal, 17(5), 720–732. Available online:

https://doi.org/10.1038/s41396-023-01386-0.

61-   Hicks, M., Tran-Dao, T.-K., Mulroney, L. & Bernick, D.L. (2021) De-novo Assembly

of Limnospira fusiformis Using Ultra-Long Reads. Frontiers in Microbiology, 12,

657995. Available online: https://doi.org/10.3389/fmicb.2021.657995.

62-   Donia, M.S. & Schmidt, E.W. (2011) Linking Chemistry and Genetics in the Growing

Cyanobactin Natural Products Family. Chemistry & Biology, 18(4), 508–519. Available

online: https://doi.org/10.1016/j.chembiol.2011.01.019.

63- Krienitz, L., Bock, C., Kotut, K. & Préschold, T. (2012) Genotypic diversity of Dicty-
osphaerium-morphospecies (Chlorellaceae, Trebouxiophyceae) in African inland wa-
ters, including the description of four new genera. Fottea, 12(2), 231–253. Available
online: https://doi.org/10.5507/fot.2012.017.

64- Kumaresan, V., Sannasimuthu, A., Arasu, M.V., Al-Dhabi, N.A. & Arockiaraj, J.
(2018) Molecular insight into the metabolic activities of a protein-rich micro alga, Ar-
throspira platensis by de novo transcriptome analysis. Molecular Biology Reports,
45(5), 829–838. Available online: https://doi.org/10.1007/s11033-018-4229-1.

65- Grasso, C.R., Pokrzywinski, K.L., Waechter, C., Rycroft, T., Zhang, Y., Aligata, A.,
Kramer, M. & Lamsal, A. (2022) A Review of Cyanophage–Host Relationships: High-
lighting Cyanophages as a Potential Cyanobacteria Control Strategy. Toxins, 14(6),
385. Available online: https://doi.org/10.3390/toxins14060385.

Figures

Figure 1. Heatmap analyses of pigments, lipids and metabolite matrices. Heatmap analysis of

the 40 pigments concentrations of the 93 analyzed *L. platensis* strains (normalization

according each feature) (A). Heatmap analysis of the normalized 14 fatty acid concentrations

of the 91 analyzed *L. platensis* strains (B). Total fatty acid concentration (mg.g^{-1} dw)

discriminates 2 groups of lower (1) and higher (2) lipid contents (Welch test, $p < 0.001$). (C)

Heatmap analysis of the 161 HPLC-MS/MS annotated ions of the 88 analyzed *L. platensis*

strains, expressed in normalized relative abundances.

Figure 2. Principal component analysis (PCA) (A) representing the total matrix metabolite composition of the strains regarding the groups given by the Hierarchical clustering on principal components (HCPC) (B). Characterization of each group regarding the sic tested factors (C) and on the Multiple correspondence analysis (D).

Figure 3. Multiple factor analysis (MFA: A), and Between class analysis with a $\cos^2$ threshold of 0.4 (BCA: B, $\cos^2$ plot: C) based on chemical composition of the strains. The tested factor is the sampling site.

Figure 4. Most discriminant fatty acids and pigments (cos^2 > 0.4) comparison between sampling sites (Kruskal-Wallis test, pairwise Wilcoxon testes). Pigment concentrations are given in mg.g^{-1} dw for phycobiliproteins and chlorophylls a, and µg.g^{-1} dw for other pigments(A). Fatty acid concentrations are given in mg.g^{-1} dw, and proportions in % of total fatty acids (B).

Figure 5. Most discriminant annotated ions relative abundance ($cos^2 > 0.4$) comparison

785 between sampling sites (Kruskal-Wallis test, pairwise Wilcoxon testes).