1 **PAbFold: Linear Antibody Epitope Prediction using AlphaFold2**
2

3 Jacob DeRoo[A], James S. Terry[B], Ning Zhao[E], Timothy J. Stasevich[D], Christopher D. Snow[A,C*] and Brian J. Geiss[A,B*]
4
5
6 [A]School of Biomedical Engineering, Colorado State University, Fort Collins CO USA
7 [B]Department of Microbiology, Immunology, & Pathology, Colorado State University, Fort Collins CO USA
8 [C]Department of Chemical & Biological Engineering, Colorado State University, Fort Collins CO USA
9 [D]Department of Biochemistry and Molecular Biology, Colorado State University, Fort Collins CO USA
10 [E]Department of Biochemistry and Molecular Genetics, University of Colorado-Anschutz Medical Campus, Aurora, CO USA
11
12
13 *Corresponding Authors: Brian.Geiss@colostate.edu and Christopher.Snow@colostate.edu
14
15 Keywords: AlphaFold2, antibody, linear epitope, epitope-prediction, scFv, competition ELISA

16
17
Abstract

Defining the binding epitopes of antibodies is essential for understanding how they bind to their antigens and
perform their molecular functions. However, while determining linear epitopes of monoclonal antibodies can
be accomplished utilizing well-established empirical procedures, these approaches are generally labor- and
time-intensive and costly. To take advantage of the recent advances in protein structure prediction algorithms
available to the scientific community, we developed a calculation pipeline based on the localColabFold
implementation of AlphaFold2 that can predict linear antibody epitopes by predicting the structure of the
complex between antibody heavy and light chains and target peptide sequences derived from antigens. We
found that this AlphaFold2 pipeline, which we call PAbFold, was able to accurately flag known epitope
sequences for several well-known antibody targets (HA / Myc) when the target sequence was broken into
small overlapping linear peptides and antibody complementarity determining regions (CDRs) were grafted
onto several different antibody framework regions in the single-chain antibody fragment (scFv) format. To
determine if this pipeline was able to identify the epitope of a novel antibody with no structural information
publicly available, we determined the epitope of a novel anti-SARS-CoV-2 nucleocapsid targeted antibody
using our method and then experimentally validated our computational results using peptide competition
ELISA assays. These results indicate that the AlphaFold2-based PAbFold pipeline we developed is capable of
accurately identifying linear antibody epitopes in a short time using just antibody and target protein
sequences. This emergent capability of the method is sensitive to methodological details such as peptide
length, AlphaFold2 neural network versions, and multiple-sequence alignment database. PAbFold is available
at https://github.com/jbderoo/PAbFold.

38
Introduction
Understanding where and how an antibody binds to its target protein is important for understanding how the
antibody performs its function, whether that function is neutralizing a pathogen during an immune response,
binding an epitope in immunoassays, or labeling a target molecule in a live-cell imaging experiment. However,
determining the binding epitope of an antibody can be a time and labor-intensive endeavor with significant
expense. Traditionally, antibody epitopes on target proteins have been identified by performing deletion

analysis on the target protein to determine if the antibody loses reactivity for the deletion mutants in various
immunoassays, which provides the general region of the target protein the antibody binds to. With the advent
of widely available chemical peptide synthesis, sequence-specific synthetic peptides can be used for
competitive immunoassays (such as enzyme-linked immunosorbent assays (ELISA)) to establish sequences that
can effectively compete with the antigen for antibody binding. Peptide mapping experiments are a powerful
method for determining the fine sequence of linear antibody epitopes, but these experiments can be relatively
expensive and the time between experimental design and data acquisition can be weeks to months due to the
need to design and chemically synthesize peptides. Once a peptide has been identified that binds with high
affinity and specificity to an antibody antigen binding fragment (Fab), crystal structures can be determined
that demonstrate intermolecular interactions between the peptide and antibody. These can then provide a
molecular-level explanation for an antibody's binding mode. Finally, with the advent of rapid single B-cell
sequencing technologies to analyze humoral immune responses towards vaccination or infection, determining
where specific antibody clones bind on an antigen becomes even more challenging due to the need to isolate
or synthesize specific antibody genes, produce antibodies, and then perform deletion or epitope mapping
experiments described above to fully understand how and where antibodies bind. These challenges make
determining antibody epitopes expensive and time-consuming, and limit the number of antibodies that are
characterized in detail.
62
Antibodies that bind to linear epitopes represent an important subset to molecular biology, as they can be
added to recombinant proteins for use in various types of immunoassays. A number of linear antibodies have
been developed for use in various immunoassays (ELISA, western blot, immunofluorescence, etc.). The
development of computational methods for linear epitope determination could increase the number and
quality of new linear epitopes available to the field. Most epitope prediction tools (such as BepiPred (1),
ElliPro (2), and ABCpred (3)) are generally designed to predict regions of an antigen that could be recognized
by any antibody rather than a specific antibody. These programs also provide no insight into the structural
match of the epitope and antibody, potentially making decisions without key structural information that
otherwise may be relevant. The challenge in predicting epitopes for a *specific* antibody lies in the complexity of
protein-protein interaction dynamics, which includes conformational changes, binding affinities, and
thermodynamic stability. Structure based approaches including HADDOCK (4, 5) and ZDOCK (4, 6) can be used
to dock peptides into antibody structures, but these require known peptides for binding. Significant progress
has been made to address this problem via deep learning: some of the new and exciting tools are GearBind (7),
PALM and A2binder (8), and DSMBind (9). We point the reader to this review for an excellent overview of some
of the tools that have existed for some time, along with a comparison of these tools (10).
78
Determining antibody-epitope interactions is, at its most basic level, a structural biology problem. Determining
what molecular interactions are present between an antibody and its antigen can define the epitope,
determine what portions of the epitope and CDR sequences are responsible for molecular interactions, and
provide clues to antibody specificity and affinity. With the advent of highly accurate structural predictions,
including the AlphaFold2 (AF2) neural networks (11, 12), the ability to accurately predict protein structures,
and potential protein-protein interactions, has dramatically increased. AlphaFold2 was trained on existing
protein structures and can effectively model new protein structures. Numerous antibodies, antibody Fab
regions, and other related constructs with bound target peptides or proteins have been crystalized and
deposited into the Protein Data Bank (PDB) (for example (13–16)). These PDB entries represent a valuable
training set that may increase the likelihood that AlphaFold2 can successfully predict the structure for

antibody-epitope complexes (12, 17–19). The authors of AlphaFold2 multimer (12) comment on the difficulty
of predicting antibody-epitope complexes, and results for this are indeed mixed at best (17–19). One way in
which this current report is distinct is our focus on linear epitopes. We hypothesize that the lack of strong
competing structure within the short peptide may boost AF2 prediction of scFv-epitope binding predictions
relative to conformational epitopes. This problem has precedent, as AlphaFold2 has previously been used to
study the interactions between proteins and peptides (17, 18). AlphaFold2's ability to correctly dock
independent protein chains can be repurposed to predict how strongly two proteins interact together and
extends to predicting the interaction between an antibody and short flexible peptides (linear epitopes) drawn
from a larger protein antigen.
98
To maximize compute efficiency, it is helpful to minimize the size of the system subject to structure prediction.
The computational expense of AlphaFold2 scales with the square of the length of the concatenated sequences
involved. Fortunately, not all portions of the antibody are critical. Antigen binding by antibodies is primarily
dictated by the antigen binding fragment (Fab) containing the variable light (V_L) and variable heavy (V_H)
fragments. Conversion of full antibody sequences into single chain variable fragments (scFv) can significantly
reduce structure prediction complexity and compute time. A wildtype scFv sequence can easily be generated
directly from translated antibody heavy and light chain DNA sequences. Briefly, the sequences are first divided
into framework and complementarity determining regions (CDRs) using Kabat (20) or IMGT (21) nomenclature.
A flexible linker sequence (GGGGSGGGGSGGGGS, 15 a.a.) is then added between the new C-terminus of the
truncated light chain and the original N-terminus of the shortened heavy chain to generate a single protein
sequence that incorporates both antigen-binding chains. The resulting fusion protein often functions in a
similar fashion to the original antibody. Another well-known protein engineering strategy for antibodies is
"loop grafting", where the CDR loops from one antibody are grafted onto a different framework region. We
have recently used this approach to develop scFvs with improved *in vivo* performance (22). The structures of
the novel scFv chimeras can be rapidly and confidently be predicted by AlphaFold2 due to their small size and
the extensive immunoglobin representation within sequence databases and the PDB. Excluding the time
needed to obtain a multiple sequence alignment (MSA), predicting the structure for a single scFv in complex
with a 10-a.a. peptide requires only 1.5 minutes on an NVIDIA A5000 graphics processing unit (GPU). This
modest compute time allows a GPU-laden server or workstation to handle large-scale structure prediction of
hundreds of related systems. As for the MSA input, a high quality MSA can quickly be obtained via ColabFold
(23), which relies on the MMseqs2 MSA server. In our workflow, we repeatedly predict the structure for a fixed
single scFv sequence in complex with varying peptide partners. In this case, we do not expect the peptide
portion of the MSA to be useful. Therefore, to avoid sending hundreds of nearly identical MSA requests to
MMseqs2 MSA server, and to avoid varying information in the MSA, we slightly modified the LocalColabFold
code to include the option to cache the MSA (install available on the GitHub). We generate one cached MSA
per epitope scan, where each residue in the query peptide is a glycine.
125
Several recent papers have attempted to use AlphaFold2 to identify antibody epitopes (24–26), but have
primarily focused on computational identification and have not verified their results using new antibodies that
are not within the PDB training set. While there are many other structure prediction models other than
AlphaFold2 (27, 28), including some specifically dedicated to predicting antibodies or antibody-like structures
(29–32), we chose AlphaFold2 to directly test its ability to correctly identify and place epitopes into an
antibody binding cleft. We selected AlphaFold2 due to its widespread use throughout the literature, as well as

its ease of installation and modification via the LocalColabFold implementation (23). In this project we test a
method we call PAbFold, a LocalColabFold-based pipeline to identify epitopes for several well-known linear-
epitope antibodies from sequence information only. There was a strong correlation between AlphaFold2's
confidence in the peptide structure (pLDDT) (33) and the experimentally verified epitope binding sequence.
Additionally, we found that AlphaFold2 very accurately predicted the linear epitope of a novel SARS-CoV-2
nucleocapsid-specific antibody (mBG17) with minimal prior epitope information. The molecular interactions
predicted by AlphaFold2 were experimentally validated using peptide mapping ELISA experiments. Overall,
this work demonstrates that AlphaFold2 has compelling promise for linear antibody epitope discovery from
sequence information alone. We also have observed that this emergent linear epitope prediction ability is
sensitive to the peptide length and that the performance was optimal when using AlphaFold2-multimer
version 2 and older MSAs generated by MMSEQS version 2202 server, rather than the more recent
AlphaFold2-multimer version 3 models and MMSEQS version 2302 server.
144
Materials and Methods:
Software:
All structure predictions were completed on a single AMD EPYC 7443 server with two NVIDIA RTX A5000 GPU
cards. PAbFold code was written in Python 3.7 and Bash. The only extra Python dependencies are NumPy and
Matplotlib. AlphaFold2 calculations were run using an installation of LocalColabFold (23). Briefly, PAbFold
contains 3 stages. In the first stage, a python script 'A_PeptideMapping_prep_submission_files.py' writes
FASTA input files for ColabFold. Each FASTA file contains the entire sequence of the subject scFv, a colon ":",
and then the candidate linear epitope which represents a small section of the target antigen protein that
changes dependent upon both the epitope length (default 10 a.a.) and a sliding window (default 1 a.a.).
154
After completion of the ColabFold jobs, two different analysis methods are presented in this paper, and both
are accessible via the 'B_PeptideMapping_plddt_perres_analysis.py' python script. The first is the 'Simple
Max' method, which assesses each peptide window with only the output model that is top ranked by
ColabFold (on the basis of ipTM). The AlphaFold2 confidence pLDDT (33) is recorded for each residue within
the peptide. Other than the N- and C-terminal residues, each residue is observed within multiple windows. We
proceed to calculate (and plot) the maximum pLDDT observed for each residue across the set of sliding
window peptides that contain that residue. Thus, in the 'Simple Max' method each residue is considered
independently. To obtain aggregate scores for each peptide window, we sum the maximum pLDDT associated
with each member residue. This method is sensitive in that any isolated high-confidence residue placements in
the top ranked AlphaFold2 peptide prediction can increase the score, but a high aggregate peptide score could
arise from multiple, mutually inconsistent peptide binding poses. Our second, complementary analysis method
instead focuses on recognizing full peptide poses of elevated AlphaFold2 confidence. We refer to the second
method as the 'Consensus method' because it begins by averaging the per-residue pLDDT across the five
AlphaFold2 models. We then compute the average pLDDT for each peptide. For visual inspection, scripts
output a heat map for the average per-residue pLDDT and a bar-chart that for the subsequent per-peptide
average pLDDT. In this case, we simply rank top peptides based on the per-peptide average pLDDT. Scripts are
available at https://github.com/jbderoo/PAbFold.
172
173 ***Antibody sequences:*** Sequences and references for antibodies, scFvs, and antigens can be found in
174 **Supplemental Table 1A**. To create an scFv, the complementarity determining regions or loops of an antibody

175 are identified via the Kabat numbering scheme. The loops are then spliced onto the scFv backbones of the
176 15F11 and 2E2 as previously described by our group (22). The scFv sequences are aligned with their CDR loops
177 and flexible linkers highlighted in **Supplemental Table 1B**.
178
Monoclonal Antibody Production:
Anti-SARS-CoV-2 nucleocapsid protein (NP) monoclonal mouse antibody mBG17 was previously developed and
characterized (34). Briefly, two BALB/c mice immunized with recombinant NP were sacrificed and primary
splenocytes isolated. Splenocytes were fused with Sp2/0 Ag14 myeloma cells and individual hybridoma clones
were isolated after eleven days. Hybridoma clones were tested for antibody production against NP via enzyme-
linked immunosorbent assay (ELISA) and western blot. Clones were further tested for isotype and cross-
reactivity, and V_H and V_L sequences were determined. The hybridoma clone mBG17 was identified as a SARS-
CoV-2 nucleocapsid-specific antibody targeting linear epitope via ELISA and western blot (34). Generation of
recombinant mBG17 and production of recombinant antibody in 293F cells was performed as described in
(35). The approximate epitope region for mBG17 was determined via western blot with modified recombinant
NP proteins containing 40 to 50 amino acid deletions. The epitope location was determined to reside between
SARS-CoV-2 nucleocapsid residues a.a. 381-419 based on loss of western blot signal with the a.a. 381-419
deletion (34).
192
Peptide Competition ELISA:
The anti-SARS-CoV-2 nucleocapsid protein mBG17 antibody epitope was experimentally identified using
competition enzyme-linked immunosorbent assay (ELISA). Using the previously determined 39 nucleocapsid
protein amino acid range for the mBG17 epitope as a starting point, seven overlapping peptides were
synthesized (Thermo Scientific) spanning the 39 amino acid region with overlaps of 5 amino acids. These
peptides were termed Fragment 1 through 7 (**Table 1**). A 96-well ELISA plate was coated with 0.1ug/ml of
recombinant SARS-CoV-2 NP (34) overnight at 4^{o}C. The plate was blocked with 4% (w/v) dry non-fat milk in 1X
PBS with 0.1% (v/v) Tween-20 for 1 h shaking at room temperature. While blocking, inhibited recombinant
mBG17 antibody samples were produced by incubating 40 µL of antibody with 40 µg (approximately 30 nMol)
of a single peptide fragment for one hour at room temperature. Following this, peptide-incubated mBG17 was
applied to the blocked nucleocapsid protein coated plate in triplicate and allowed to incubate for 1 h at room
temperature while shaking. The plate was rinsed with 0.1% (v/v) Tween-20 in 1X PBS and washed three more
times for 5 minutes shaking at room temperature. The plate was then incubated with HRP-conjugated goat
anti-mouse polyclonal antibody solution diluted at 1:20,000 in 1X PBS for 1 h shaking at room temperature.
After another rinse and three more washes the plate was developed with 1-Step™ Ultra TMB-ELISA Solution
(ThermoFisher) before stopping the reaction with an equal volume of 2M H_2SO_4. Solution absorbance at 450
nm was measured using a PerkinElmer Victor X5 multilabel plate reader. Absorbances were averaged within
fragment-inhibited sample groups and corrected with the average value of the negative control. These
absorbances were then normalized against the absorbance from the group with the highest value before
multiplying by 100 to obtain percentage of potential signal.
213
The effect of single alanine substitutions on fragment 5 (DDFSKQLQQS) peptide binding was determined by
competition ELISA using a series of ten alanine-substituted peptides (**Table 1**) at a range of concentrations to
determine relative competition activity. A modified version of the previously described inhibition ELISA was

performed using the unmodified Fragment 5 peptide and the ten alanine-substituted peptides. During the
mBG17 inhibition step, the mBG17 antibody solution was incubated with a 4-fold serial dilution of peptides
beginning at 40 µg and continuing to ~2.5 ng before being applied to the NP coated plates in triplicate. The
remainder of the competition ELISA was carried out as described above.
221

Table 1:

Peptide Name	Peptide Sequence
Nucleocapsid a. a. 381-390 (Frag 1)	ALPQRQKKQQ
Nucleocapsid a. a. 386-395 (Frag 2)	QKKQQTVTLL
Nucleocapsid a. a. 391-400 (Frag 3)	TVTLLPAADL
Nucleocapsid a. a. 396-405 (Frag 4)	PAADLDDFSK
Nucleocapsid a. a. 401-410 (Frag 5)	DDFSKQLQQS
Nucleocapsid a. a. 406-415 (Frag 6)	QLQQSMSSAD
Nucleocapsid a. a. 411-419 (Frag 7)	MSSADSTQA
Nucleocapsid D401A	ADFSKQLQQS
Nucleocapsid D402A	DAFSKQLQQS
Nucleocapsid F403A	DDASKQLQQS
Nucleocapsid S404A	DDFAKQLQQS
Nucleocapsid K405A	DDFSAQLQQS
Nucleocapsid Q406A	DDFSKALQQS
Nucleocapsid L407A	DDFSKQAQQS
Nucleocapsid Q408A	DDFSKQLAQS
Nucleocapsid Q409A	DDFSKQLQAS
Nucleocapsid S410A	DDFSKQLQQA

222
223
224 **Results:**
225
Assessment of AlphaFold2 generated scFv structures. We first verified that AlphaFold2 could generate scFv
structures that have similar structures to their parent monoclonal antibodies. We chose the 9E10 clone of the
anti-Myc antibody as an initial test system, as the scFv sequence is available (36) and has a well-known linear
epitope (EQKLSEEDL)(37). We predicted the wild-type Myc scFv structure and aligned this model to the
corresponding Fab crystal structure (PDB entry 2orb) via the align command in PyMOL (**Supplemental Figure**
1A). The AlphaFold2 predicted scFv was very similar (RMSD value of 0.42Å) to the anti-Myc Fab structure,
suggesting that the predicted scFv structure was a suitable starting point for epitope prediction. We also
examined the structures of the Myc CDRs loop grafted onto the 15F11 (38) and 2E2 (22) frameworks, as we
have previously observed that loop grafting onto these frameworks can enhance protein folding and solubility
(22). The loop-grafted Myc-2E2 and Myc-15F11 and structures were also similar to the Myc Fab structure (PDB
2ORB) (37) with similar RMSD values of 0.45Å (**Supplemental Figure 1B**), indicating that they are also
reasonable starting points for epitope prediction.
238

Development of Python-based scripts for automated scFv:peptide structure prediction. We developed a series
of Python scripts that automate the process of epitope prediction and analysis with AF2.
A_Peptide_Mapping_prep_submission_files.py accepts a linear scFv sequence and a linear full-length antigen
sequence, and processes the antigen sequence into a series of short peptides with custom peptide length and
sliding window sizes (default parameters are 10 amino acid peptides with a 1 amino acid sliding window). It
then adds lines for each scFv:peptide pair to a FASTA file. Structures are then predicted via LocalColabFold for
each scFv:peptide pair with AlphaFold2 in parallel on two NVIDIA RTX A5000 GPUs. The python script
B_PeptideMapping_plddt_perres_analysis.py parses the AlphaFold2 output structures to extract per-residue
pLDDT for the peptide residues in each scFv:peptide pair. Conf_plot_and_top10.py will plot the maximum
pLDDT (across all host peptides) scores as a function of amino acid position within the antigen sequence and
ranks predicted peptides based on ΣpLDDT scores for the 'Simple max' method. To use the 'Consensus'
method, include the –all-models flag when running B_PeptideMapping_plddt_perres_analysis.py. We also
supply a python script that replicates how we present the data called all_model_analysis.py for use. An
overview of the method is shown in **Figure 1**. AF2's failure to predict whole antigen structure coupled with the
scFv is highlighted in **Supplemental Figure 2**.
254

A Anti-Myc Fab
(2or9)

V_H
Myc
V_L

C_H1
C_L

AlphaFold2 prediction
for designed anti-Myc
scFv:Myc complex

Flexible
Linker
Myc

20   100

pLDDT

B Epitope scanning

Target protein:

MDFFRVVENQPPATMPL...

MDFFRVVENQ--------
-DFFRVVENQP-------
--FFRVVENQPP------
---FRVVENQPPA-----
----RVVENQPPAT----
-----VVENQPPATM---
------VENQPPATMP--

Sliding window:
overlapping linear
epitope candidates

C

Consensus (Heat Map)

\<pLDDT\> Sliding Window Sub-Index

Per-Residue pLDDT

Consensus (Sliding Window)

\<\<pLDDT\>\>

0   100   200   300   400

Window in Target Protein

Simple Max

Per-Residue Maximum pLDDT

Experimental Epitope

0   100   200   300   400

Residue in Target Protein

83.6: EQKLISEEDL
78.9: QKLISEEDLL
69.6: AAKLVSEKLA
67.6: TMPLNVSFTN
65.8: NFYQQQQQSE

Figure 1. PAbFold pipeline for linear epitope prediction. A) Antibody V_H and V_L protein sequences are used to generate scFv sequences, either based on the native antibody sequences or loop grafting complementarity determining regions (CDRs) onto either the 2E2 or 15F11 antibody framework regions (2E2 shown). **B)** The target antigen sequence is parsed into a list of small overlapping peptide sequences, with peptide step and window size parameters adjusted as needed. Rank ordered peptides are output, and partial epitope sequences are underlined manually to highlight the identification of the correct sequence. **C)** The scFv sequences from Panel A are co-folded with each of the peptide sequences derived from the target antigen in parallel batch mode on a GPU server. pLDDT scores from each structure prediction experiment are collected and scores are presented in their sliding window, both as a heat map organized along the length of the target antigen sequence and a bar chart that shows the per-peptide average pLDDT (Consensus Method). Additionally, the Simple Max data is presented in the third and final panel.

Testing of scFv:peptide structure prediction method using the Myc Epitope. We first tested the PAbFold method with the anti-Myc-scFv described in (39), using the full-length human Myc proto-oncogene protein sequence as the antigen. We initially used an antigen peptide length of 10 and a 1 amino acid sliding window. Given these parameters, the 9 a.a. Myc epitope motif (EQKLSEEDL) appeared intact within one of the 10-mer peptides, with subsets of the 8, 9, 11, and 12 a.a. appearing in neighboring sliding peptide windows. PAbFold generated predicted structures, each of which took an average of ~200 seconds to process. The entire process took approximately 12 hours on our GPU server. AlphaFold2 placed all peptides into or near the traditional antigen binding site between the CDR loops (**Supplemental Figure 3**). The average confidence (mean pLDDT across residues) for these peptides ranged from 20 to 90. When we inspect the consensus confidence for each residue in each sliding window (**Figure 2A**), the expected Myc peptide epitope (EQKLSEEDL) was one of several peptides with high average pLDDT. The second highest ranked peptide in this analysis (QKLSEEDLL) was a near perfect match for the expected epitope. We consider this window to be a successful prediction. Perhaps surprisingly, the peptide window with the exact match (EQKLSEEDL) did not score particularly well due to its average pLDDT of 51.0. In this instance, the expected epitope sequence did not stand out when plotting the maximum observed per-residue pLDDT for each residue (**Figure 2A, bottom**).

We proceeded to test predictions with two engineered scFv chimeras where loop grafting was used to place the Myc recognition CDRs onto two antibody framework regions with high *in vivo* performance, generating Myc-15F11 and Myc-2E2 scFv sequences. Epitope prediction performance was markedly improved with the chimeric scFvs (**Figure 2B and 2C**). Specifically, the QKLSEEDLL peptide window became the top ranked peptide on the basis of average consensus pLDDT. In the case of Myc-2E2 (**Figure 2C**), the average confidence for the correctly predicted epitope was particularly high compared to alternate peptide windows, and another close match to the expected epitope (EEQKLSEED) was ranked within the top 5 peptides (**Figure 2D**). Ranking epitopes using the Simple Max analysis was similar; the region containing the correct epitope was nearly top ranked for Myc-15F11 and was top ranked for Myc-2E2 (**Figure 2E**). Thus, AlphaFold2 was able to more clearly detect authentic Myc antibody epitope using CDRs loop grafted onto the 2E2 or 15F11 frameworks, relative to the native Myc scFv framework.

To investigate the superior epitope recognition performance of the chimeric Myc scFvs, we aligned the Cα coordinates for the predicted scFv structures (predicted with and without the target epitope) to the reference crystal structure and calculated the RMSD for all backbone positions (N, Cα, C, O) and the loops (**Supplemental Figure 4**). Notably, regardless of the Myc scFv variant, the CDR loop RMSD improved by more than 1Å when the epitope was present. Secondly, consistent with the improved epitope prediction performance for the

299 chimeric scFvs (15F11 and 2E2), the epitope peptide QKLISEEDL was placed more accurately for those
300 predicted structures than in the WT scFv (**Supplemental Figure 4**). We could not discern an obvious structural
301 difference between the WT and chimeric scFvs that explains the structure prediction performance gap.
302

303
Figure 2. The Alphafold2-based PAbFold method predicted the Myc linear epitope in different scFv backbones. The anti-Myc V_H and
V_L antibody sequences were used to generate either **A)** wild-type Myc scFv or loop grafted chimeric **B)** Myc-15F11 or **C)** Myc-2E2 scFv
variants. The Myc proto-oncogene protein sequence (Genbank NP_001341799.1) was used as the target antigen and processed into
10 amino acid overlapping peptides with a 1 amino acid sliding window. The structure for each scFv:peptide pair was predicted with
AlphaFold2 in batch mode on two NVIDIA A5000 GPUs. Average consensus pLDDT values for each scFv:peptide window are
illustrated, as well as the maximum pLDDT observed for each residue in any window (bottom). **D)** Top ranking binding peptides
based on average consensus pLDDT. **E)** Top ranked binding peptides based on summing per-residue maximum pLDDT. For D and E,
underlining represents overlap with the reported Myc epitope (EQKLSEEDL).
312

Assessment of peptide length, sliding window size, and position on AlphaFold2 scFv:peptide structure
prediction. Our initial selection of the 10 a.a. window was intended to match or slightly exceed the size of
known epitopes such as Myc and HA. We next assessed how different peptide sizes and sliding window lengths
would affect epitope prediction accuracy and run time. We re-ran the Myc-2E2-scFv:peptide complex
prediction calculations varying peptide size between 8, 9, 10, and 11 (with a fixed sliding window size of 2) or
varying the sliding window size to 1 or 5 (with a fixed peptide size of 10). We observed that using a sliding
window of 2 a.a. provided nearly the same level of accuracy and resolution as the 1 a.a. Ultimately, we
determined that our original peptide size of 10 amino acids and sliding window of 1 a.a. provided highest
resolution data possible (**Supplemental Figure 5**), and therefore maintained a peptide size of 10 and a sliding
window length of 1 for our remaining experiments.
323

We then predicted the complex structure for Myc-2E2 with various negative control peptides: A_{10}, $(GS)_5$,
$(GGGGS)_2$, and G_{10} to determine how non-binding peptides are docked and scored (**Supplemental Figure 5I**
and 5J). We again observed that AlphaFold2 placed all peptides into the traditional antigen binding between
the CDR loops, but the reported peptide scores for the negative controls were particularly low (29 – 41). These
results indicate that AlphaFold2 "knows" where antigens bind in antibody or scFv structures and attempts to
model any peptide partner into this region, but the low pLDDT scores indicate confidence in the interactions
are quite low.
331

We also tested if AlphaFold2 could detect the Myc epitope if it was inserted as an epitope tag within different
positions of a heterologous protein. We created a synthetic antigen by adding the Myc epitope within the 99-
a.a. unrelated HIV-1 Gag protease protein sequence at either the N- or C-terminus or in the middle of the
protein sequence, and used PAbFold to detect the Myc peptide (**Supplemental Figure 6**). In each case, the
average consensus pLDDT was highest for the inserted epitope, such that the authentic epitope would be top
ranked and prioritized for testing. Thus, as expected for a sliding window analysis, the epitope position within
the antigen was no barrier to detection.
339
Testing of the PAbFold method using the HA Epitope. Based on our success detecting the Myc epitope, we
sought to determine if our method could detect a different well-known linear peptide, HA, derived from
positions 114-126 within the Influenza A virus hemagglutinin protein (YDVPDYASLR). Using an anti-HA scFv
sequence that had been previously generated (22, 39), we generated new HA-15F11 and HA-2E2 scFvs loop
grafted sequences. We used the same procedure described above to predict structures for influenza A virus HA
derived peptides on HA-scFv (**Supplemental Figure 7A**), HA-15F11-scFv (**Supplemental Figure 7B**) and HA-2E2-
scFv (**Supplemental Figure 7C**). In the HA case, the expected epitope was ranked highly for all three scFv
variants, but when assessing entire peptides by average consensus pLDDT was only ranked in the top 5 for the
HA-15F11-scFv. These results, in combination with the Myc results described above, indicate that AlphaFold2
can accurately detect linear antibody epitopes in antigen sequences, and that grafting CDR loops onto
alternative scFv backbones may increase the noise-to-signal ratio, making the identification of correct epitopes
more accurate.
352
353   Like the Myc system, trends are observed with the HA system regarding loop placement. Although not as
354   extreme, the loops for all HA scFvs undergo movement that make it more closely match the crystal structure
355   (PDB entry 1frg). Again, the epitope placement of predicted structures of the chimeric scFvs more closely
356   mimicked the deposited crystal structure than the WT scFv (**Supplemental Figure 4B**).
357
Determination and experimental validation of a novel linear antibody epitope. The Myc and HA monoclonal
antibodies are well known and several crystal structures (Myc PDB: 2or9, peptide bound (2009) | HA PDB:1frg,
peptide bound (1994)) have been solved (22, 37, 39, 40), raising the possibility that AlphaFold2 has
incorporated these antibody or epitope structures into its training set. The AlphaFold2 training set was
reported to exclude chains of less than 10, which would eliminate the myc and HA epitope peptides.
Nonetheless, to guard against the possibility that the AlphaFold2 models have incorporated specific knowledge
into the training set thereby directly probing if PAbFold epitope scanning can predict a linear antibody epitope
without *a priori* knowledge of the antibody or antigen sequence, we tested if PAbFold can predict the epitope
sequence of a recently developed antibody lacking structural information available in the Protein Data Bank.
The mBG17 mouse monoclonal antibody was generated in response to the COVID-19 pandemic, the antibody
V_H and V_L sequences were determined, and the epitope was localized to a. a. 381-419 via Western blot analysis
of deletion mutants of the nucleocapsid protein (34). mBG17 was not included in AlphaFold2's training or test
set, making it an ideal test case for *de novo* epitope prediction.
371
The mBG17 monoclonal antibody was converted to wild-type scFv, 15F11-scFv, and 2E2-scFv using the same
procedures used for Myc and HA scFv. As an additional control calculation (labeled "3-body"), we used
AlphaFold2 to predict the structure for a 3-protein complex (the peptide, and the disconnected nontruncated

mBG17 V_H and V_L variable domain sequences). All 4 Fab variants (WT scFv mBG17, 15F11-mBG17 scFv, 2E2-
mBG17 scFv, and 3-body mBG17) were screened against all 10 a.a. peptides with a 1 a.a. sliding window, as
with Myc and HA. In all 4 cases, AlphaFold2 predicted that the top ranked peptides were located in the a.a.
381-419 region of the SARS-CoV-2 nucleocapsid protein, and more specifically residues a.a. 400-415 (**Figure**
3A, 3B, 3C, and 3D). The top scoring peptide for all three scFv variants was the 402-411 window
(DFSKQLQQSM) (**Figure 3E and 3F**). The strong AF2 preference for peptides from this C-terminal segment was
particularly evident in the average consensus pLDDT analysis.
382
We next sought to experimentally verify the minimal linear epitope for mBG17 to determine how closely the
AlphaFold2 prediction corresponded to our experimental data. Seven 10 a.a. peptides that overlapped by 5
a.a. each were synthesized and used in competition ELISAs with mBG17 monoclonal antibody and recombinant
SARS-CoV-2 nucleocapsid protein (**Figure 3G and 3H**). The peptide corresponding to a.a. 401-410 showed
almost complete competition of mBG17 binding to the SARS-CoV-2 nucleocapsid protein in the ELISA, whereas
none of the other peptides were able to compete for mBG17 binding to nucleocapsid. Peptides a.a. 296-405
and a.a. 406-415 overlap a.a. 401-410 at the N- and C-terminus, respectively, but neither was able to compete,
indicating that mBG17 binds a.a. 401-410 on both sides of a.a. 405 and a.a. 406. An alignment of all the
peptides used in the overlapping peptide competition ELISA experiments showed that peptide sequence
DDFSKQLQQS represents the experimentally determined epitope for mBG17, nearly identical to the epitope
predicted by AlphaFold2 (**Figure 3H:** DDFSKQLQQS). These results demonstrate that the PAbFold pipeline was
able to very accurately predict the region that an antibody binds to a novel linear epitope that is not present in
AlphaFold2's training set.
396

Figure 3: The AlphaFold2-driven PAbFold epitope scan method can accurately identify a linear epitope for a novel SARS-CoV-2 antibody. Antibody VH and VL sequences for SARS-CoV-2 nucleocapsid protein targeted antibody were used to generate scFv sequences **A)** WT, **B)** 15F11, **C)** 2E2 or native VH and VL sequences **D)** 3 body). Variant scFv sequence in complex with peptide windows from the SARS-CoV-2 nucleocapsid protein (Genbank Accession: YP_009724397) were subjected to AlphaFold2 structure prediction. The top 5 peptides ranked by either the **E)** Consensus method or the **F)** Simple Max method, with the underlined sequence highlighting the experimentally verified sequences and a cartoon schematic for each system shown. **G)** Competition ELISA schematic for assessing the ability of synthetic peptides derived from the SARS-CoV-2 nucleocapsid protein. **H)** Amino acid windows showing binding interference, with mBG17 binding to SARS-CoV-2 nucleocapsid protein (n = 3). Percentage of binding values were calculated from the no-peptide control. Alignment of synthetic peptides corresponding to SARS-CoV-2 nucleocapsid a. a. 381-419. Peptide a. a. 401-410, which demonstrated mBG17 competition.

Fine-characterization of the mBG17 epitope and comparison to the predicted AlphaFold2 model. To further
experimentally characterize the binding of the mBG17 to the a.a. 401-410 (DDFSKQLQQS) peptide and
compare experimental data with the predicted AlphaFold2 model, we designed and synthesized ten additional
peptides, each containing an alanine point mutation at one position in the a.a. 401-410 peptide. The peptides
are labeled D1A, D2A, F3A, S4A, K5A, Q6A, L7A, Q8A, Q9A, and S10A. Competition ELISAs were performed
using increasing concentrations of each peptide to better assess differential binding (**Figure 4A**). As expected,
WT (a.a. 401-410) peptide showed strong competition, although Q9A showed slightly better competition. This
could be attributed to alanine's propensity to be in an alpha-helical coil ($\text{Prop}_{A, \text{AHC}} = 0$) vs glutamine's
propensity to escape it ($\text{Prop}_{Q, \text{AHC}} = 0.39$) (41), thus further stabilizing the Q9A alpha helix. D1A showed no
change in competition, indicating that D1 was not involved in binding. Peptides with substitutions K5A, Q6A,
and S10A showed minor reductions in competition, S4A showed a moderate reduction on competition,
whereas resides D2A, F3A, L7A, and Q8A all showed strong reductions in competition. These data indicate that
the key interactions between mBG17 and the a.a. 401-410 peptide are residues D2, F3, L7, and Q8, with S4
playing a moderate role and D1, K5, Q6, Q9 and S10 playing negligible roles in binding.
422
Finally, we compared the experimental data shown above with the best scoring mBG17:DDFSKQLQQ model
generated by AlphaFold2 (**Figure 4B and 4C**). The AlphaFold2 model suggests that residue D2 forms a hydrogen
bond with mBG17 a.a. Y34, residue F3 forms a hydrophobic interaction with mBG17 a.a. L185, residue S4 lacks
a hydrogen bond partner, residue L7 forms a hydrophobic interaction at the base of the binding cleft with
mBG17 a.a. A104, and residue Q8 hydrogen bonds with the backbone carbonyl of Y34 and the backbone amide
of W35. Residues that experimentally showed no or minimal effects on competition (D1, K5, Q6, Q9) are all
predicted to interact primarily with the solvent and lacked visible interactions between the peptide and scFv
sequence. In summary, the AlphaFold2-driven PAbFold prediction was remarkably consistent with the
experimental alanine scanning data, suggesting that the prediction of the mBG17 linear epitope location was
accurate due to the correct prediction of the structural details for how that linear epitope binds to the
antibody.
434

435
436
Figure 4. The Alphafold2-Driven PAbFold method accurately predicts molecular interactions between a linear epitope and a scFv
A) Competition ELISA assessing the ability of synthetic alanine mutant peptides derived from the SARS-CoV-2 nucleocapsid protein
(a. a. 401-410: DDFSKQLQQS) to interfere with mBG17 binding to SARS-CoV-2 nucleocapsid protein (n = 3). Percentage of binding
values were calculated from the no-peptide control. **B)** AlphaFold2 model for mBG17-15F11 scFv bound to a. a. 401-410 peptide (the
average peptide pLDDT was 83.5). Residues that display sharply reduced binding to mBG17 upon mutation to alanine in competition
ELISAs (D2, F3, S4, L7, Q8) are shown as warm-colored thick sticks. Predicted hydrogen bonds between the peptide and the scFv are
depicted by yellow bars. Sites where mutation to alanine was less disruptive to binding (Q6, K5, S10A, D1A, and Q9A) are depicted
as thin sticks with cool colors. The carbon atoms of residues in panel B are colored according to the corresponding data in panel A. **C)**
The same AlphaFold2 model for the mBG17-15F11 scFv bound to a.a. 401-410 colored with confidence (pLDDT) as predicted by AF2.
446

Discussion

In this project we assessed the ability of an AlphaFold2-based linear epitope scan pipeline we call PAbFold (Peptide:Antibody Fold) to predict linear antibody epitopes using just antibody and antigen sequences. We first assessed the quality of scFv models produced by AlphaFold2. We then developed a series of Python scripts that accept scFv and whole antigen protein sequences as inputs, parse the antigen protein sequences into short overlapping peptides, run batch predictions for each scFv:peptide pair, and output two peptide scoring schemes based on the peptide per-residue pLDDT scores as a metric for AlphaFold2 model confidence. Binding of the expected epitope to the WT-Myc scFv could only be detected via the consensus method, but either analysis method could readily detect the expected epitope bound to the chimeric Myc scFvs. Conversely, the alternate analysis method (Simple Max) performed better with respect to ranking the expected HA epitope binding to the WT and chimeric anti-HA scFv variants. In the HA case, performance was comparable for both the WT and chimeric scFv variants.

It is important to note that binding of scFv variants to sequences other than the expected epitopes may be statistically unlikely but not impossible. For example, consider the peptide ATMPLNVSFT near the N-terminus of the Myc proto-oncogene protein sequence. In the context of the WT anti-Myc scFv this peptide had slightly higher average consensus pLDDT (52.4 rather than 51.0) than a peptide (QKLISEEDLL) that closely matched the expected epitope. In the absence of direct experimental evidence, predicted affinity for this unexpected sequence is not necessarily incorrect, though the lack of comparable predicted binding to the 15F11 and 2E2 chimeric scFv variants further decreases the likelihood. In the future, it might be useful to assess peptide binding via consensus across scFv variants.

Lastly, we tested this process on a novel antibody generated by our group targeting the SARS-CoV-2 nucleocapsid protein (mBG17) and found the method performed significantly better than with Myc and HA. Either analysis method could very easily flag peptide windows containing the authentic experimentally validated epitope. This worked for the WT scFv, the chimeric scFv variants, and even a structure with disconnected heavy and light chain domains. Experimentally, we cleanly validated the AlphaFold2 prediction using a peptide competition ELISA assay to experimentally determine the mBG17 epitope. Confidence in the AlphaFold2 prediction was further buoyed via alanine scanning peptide competition ELISAs that verified the importance of the key binding interactions predicted by AlphaFold2.

Identification of antibody V_H and V_L sequences from monoclonal B-cells has become a routine task, with sequence information obtainable via various sequencing technologies such as next generation sequencing and nanopore sequencing for a relatively low cost. As a result, the determination of the epitope in service of a deeper understanding of how antibodies bind their antigen is an increasingly notable bottleneck. An experimental epitope determination campaign can take weeks or months of work, but with the advent of AlphaFold2 and the epitope prediction method we describe here, an antibody and its antigen could be sequenced in a few days (often through contract research organizations for low cost) and accurate linear epitope predictions generated within less than a day, dramatically epitope validation throughput as well as providing detailed predictions for the molecular features of antibody-epitope interaction.

Conformational epitopes are structured antigens that are found during many immune responses, and prediction of these epitopes from antibody and antigen sequences would be a significant boon to the field of biology. For example, conformational epitope prediction coupled with single-cell B-cell sequencing would allow for detailed analysis of antibody maturation during immune responses to vaccines or pathogen infection, helping better define how the immune response to infection evolves over time and how evolution of antigen sequences affects the antibody response. In this work we did not focus on using AlphaFold2 to predict conformational epitopes primarily because of the complex structures that conformational epitopes possess. Literature reports suggest that prediction of the complexes between antibodies and both whole antigens and conformational epitope proteins has proven to be very difficult for AlphaFold2, and indeed the authors themselves make this observation (12, 42, 43). Notably, the structures that proved most difficult to predict for AF2 and other tools in the CASP15-CAPRI154 challenges were antibody-antigen complexes (44). Reports suggest that a mix of both statistics-based approaches (neural networks like AF2) and physics-based approaches (such as Rosetta) predict optimal antibody-antigen complexes (45). Indeed, if we attempt to predict binding of our scFvs to intact antigen proteins (**Supplemental Figure 2**), we find no predictive capability. When predicting scFv:peptide complexes, it may be the case that AlphaFold2 is able to thoroughly evaluate an induced fit for the peptide due to both its length (small sample space) and its propensity to not adopt a strong competing structure. In contrast, a larger and complicated structure may be more challenging to move during the AlphaFold2 structure prediction or recycle steps. Additional complexities may arise in extreme induced conformational changes during docking. Recent reports indicate that progress is being made in predicting the binding locations of conformational epitopes (46, 47).

We observed that the ability of AlphaFold2 to successfully predict the epitope peptide binding is quite delicate. First, epitope prediction was highly sensitive to the peptide length (**Supplemental Figure 5**), with minimal predictive power for peptide length other than 10 a.a. Further investigation of this sensitivity would be a useful avenue for future research. Perhaps with enhanced sampling, epitopes can be detected within longer peptides (e.g. 11 a.a., 12 a.a., etc.). Methodological tuning of this type could ultimately help illuminate the path to increasingly difficult protein-protein binding prediction problems. Similarly, we have likewise determined that epitope scanning performance was sensitive to changes in the underlying AlphaFold2 neural networks and the MSA. Specifically, unless otherwise noted, all data in this report was obtained using ColabFold version 1.5.2 and the 5 neural networks that comprise AlphaFold2 multimer version 2 (mm2). Likewise, the MSAs we use were obtained from the MMSEQS server (and cached) when the default sequence databases were UniRef30 2202 and PDB70 220313. They have since been updated to PDB30 2302 and PDB100 230517. For a complete description, see the change logs on the github for ColabFold (https://github.com/sokrypton/ColabFold#colabfold---v152).

Insofar as protein-peptide prediction is an emergent "off-label" capability for AlphaFold2 that is not part of the training sets, further training of the models or other changes can degrade performance. Benchmarking performance can be difficult when there are multiple moving targets. The most recent calculations we have analyzed were using ColabFold version 1.5.2 which was current as of February 19, 2023. The changes from ColabFold 1.5.2 to 1.5.5 (current as of this writing) are limited to version control and ensuring ColabFold still works on Google Colab, and therefore will not change the calculation performance. Relative to ColabFold 1.3 (the current method at the outset of this project), ColabFold 1.5.2 embodied two substantial changes. First,

ColabFold 1.5.2 used the updated AlphaFold multimer (mm) version 3 by default. Second, the backend server MMSEQS ((48) and (https://github.com/soedinglab/MMseqs2)) that supplies MSAs also underwent updates, namely the database updates. Upon evaluation, we found that the recent default methods (ColabFold 1.5.2) still predicted the epitope successfully for the mBG17 system (**Supplemental Figure 8**). However, the ColabFold 1.5.2 default methods had a pronounced decline in PAbFold performance for the HA and Myc systems. Specifically, the combination of mm3 and the revamped ColabFold MSA server tended to be less discriminating compared to the default settings for ColabFold 1.3 (ColabFold 1.3 was the most up to date version when this project was initialized). The updated configuration flagged diverse peptide sequences with elevated pLDDT values (**Supplemental Figures 9 and 10**) resulting in the loss of successful epitope predictive power. While testing ColabFold 1.5.2 with the most recent MSA server, but reverting the AlphaFold2 models to mm2, the outcome improved, with experimentally validated sequences rising to the top more frequently than when using mm3, but still falling short in ranking the experimentally validated epitope sequence embedded within the antigen. However, when previously cached MSAs were paired with mm2 (using ColabFold 1.5.2), performance was maximized. Furthermore, we attempted to recreate the MSA databases locally with similar but not identical results to queueing the server with databases UniRef30 2202 and PDB70 220313 (**Supplemental Figure 11**). Additionally, the MMSEQS team ((48) and (https://github.com/soedinglab/MMseqs2)) graciously rebuilt a server we could query using LocalColabFold that mimicked the original UniRef30 2202 and PDB70 220313 database set up as closely as possible on their end. The MSA that was generated from these databases was used, and still did not perform as well as the original MSAs that were generated upon first retrieval and generation (**Supplemental Figure 12**). As a negative control, we repeated all calculations without using any MSAs, and only relying upon the sequence to make a structural prediction. As expected, all epitopes were scored very poorly (**Supplemental Figure 13**). Despite our significant efforts, it is unclear why our initial results cannot be perfectly recapitulated, but the difference has been traced to detailed MSA contents (**Supplemental Figure 14**), resulting in differences in correct epitope identification. These results are summarized in (**Supplemental Figure 15**).

One key lesson of this research effort is that caching the MSAs proved to be very useful as a method to guard against changes in the performance of 3[rd] party tools. We recommend that future methods development work using LocalColabFold adopt the strategy of caching MSAs when feasible. It is also our hope that by describing the latent ability of AlphaFold2 to predict scFv-binding epitopes that this ability will be preserved and enhanced in future iterations.

Acknowledgements

The authors would like to thank members of the Snow, Stasevich, and Geiss groups for helpful discussions at TagTeam meetings. This project was funded in part by NIH R01AI132668 (Geiss) and NIH R56AI155897 and R01AI168459 (Snow, Stasevich, Geiss). The authors would also like to thank Dr. Milot Mirdita for being our point of contact for the ColabFold and MMSEQS teams, and assisting us with setting up databases and generating MSAs for our use.

References Cited

1. Larsen JEP, Lund O, Nielsen M. 2006. Improved method for predicting linear B-cell epitopes. Immunome Res 2:2.
2. Ponomarenko J, Bui HH, Li W, Fusseder N, Bourne PE, Sette A, Peters B. 2008. ElliPro: A new structure-based tool for the prediction of antibody epitopes. BMC Bioinformatics 9:1–8.
3. Saha S, Raghava GPS. 2006. Prediction of continuous B-cell epitopes in an antigen using recurrent neural network. Proteins 65:40–8.
4. Ambrosetti F, Jiménez-García B, Roel-Touris J, Bonvin AMJJ. 2020. Modeling Antibody-Antigen Complexes by Information-Driven Docking. Structure 28:119-129.e2.
5. Dominguez C, Boelens R, Bonvin AMJJ. 2003. HADDOCK: A protein-protein docking approach based on biochemical or biophysical information. J Am Chem Soc 125:1731–1737.
6. Chen R, Li L, Weng Z. 2003. ZDOCK: an initial-stage protein-docking algorithm. Proteins 52:80–7.
7. Cai H, Zhang Z, Wang M, Wu Y, Ying T, Tang J. 2021. Pretrainable Geometric Graph Neural Network for Antibody Affinity Maturation.
8. He H, He B, Guan L, Zhao Y, Chen G, Zhu Q, Yu-chian C. 2023. De novo generation of antibody CDRH3 with a pre-trained generative large language model.
9. Jin W, Chen X, Vetticaden A, Sarzikova S, Raychowdhury R, Uhler C, Hacohen N. 2023. DSMBind: SE(3) denoising score matching for unsupervised binding energy prediction and nanobody design. bioRxiv 1–24.
10. Jaszczyszyn I, Bielska W, Gawlowski T, Dudzic P, Satława T, Kończak J, Wilman W, Janusz B, Wróbel S, Chomicz D, Galson JD, Leem J, Kelm S, Krawczyk K. 2023. Structural modeling of antibody variable regions using deep learning—progress and perspectives on drug discovery. Front Mol Biosci 10:1–8.
11. Jumper J, Evans R, Pritzel A, Green T, Figurnov M, Ronneberger O, Tunyasuvunakool K, Bates R, Žídek A, Potapenko A, Bridgland A, Meyer C, Kohl SAA, Ballard AJ, Cowie A, Romera-Paredes B, Nikolov S, Jain R, Adler J, Back T, Petersen S, Reiman D, Clancy E, Zielinski M, Steinegger M, Pacholska M, Berghammer T, Bodenstein S, Silver D, Vinyals O, Senior AW, Kavukcuoglu K, Kohli P, Hassabis D. 2021. Highly accurate protein structure prediction with AlphaFold. Nature 596:583–589.
12. Evans R, O'Neill M, Pritzel A, Antropova N, Senior A, Green T, Žídek A, Bates R, Blackwell S, Yim J, Ronneberger O, Bodenstein S, Zielinski M, Bridgland A, Potapenko A, Cowie A, Tunyasuvunakool K, Jain R, Clancy E, Kohli P, Jumper J, Hassabis D. 2022. Protein complex prediction with AlphaFold-Multimer. bioRxiv 2021.10.04.463034.
13. Ofek G, Tang M, Sambor A, Katinger H, Mascola JR, Wyatt R, Kwong PD. 2004. Structure and Mechanistic Analysis of the Anti-Human Immunodeficiency Virus Type 1 Antibody 2F5 in Complex with Its gp41 Epitope. J Virol 78:10724–10737.
14. Ekiert DC, Bhabha G, Elsliger M, Friesen RHE, Jongeneelen M, Throsby M, Goudsmit J, Wilson IA. 2009. Antibody recognition of a highly conserved influenza virus epitope : implications for universal prevention and therapy. Science (80-) 324:246–251.

15. Stanfield RL, Gorny MK, Williams C, Zolla-Pazner S, Wilson IA. 2004. Structural Rationale for the Broad
Neutralization of HIV-1 by Human Monoclonal Antibody 447-52D. Structure 12:193–204.

16. Zhou T, Xu L, Dey B, Hessell AJ, Van Ryk D, Xiang SH, Yang X, Zhang MY, Zwick MB, Arthos J, Burton DR,
Dimitrov DS, Sodroski J, Wyatt R, Nabel GJ, Kwong PD. 2007. Structural definition of a conserved
neutralization epitope on HIV-1 gp120. Nature 445:732–737.

17. Ko J, Lee J. 2021. Can AlphaFold2 predict protein-peptide complex structures accurately? bioRxiv
2021.07.27.453972.

18. Tsaban T, Varga JK, Avraham O, Ben-Aharon Z, Khramushin A, Schueler-Furman O. 2022. Harnessing
protein folding neural networks for peptide–protein docking. Nat Commun 13:1–12.

19. Ghani U, Desta I, Jindal A, Khan O, Jones G, Hashemi N, Kotelnikov S, Padhorny D, Vajda S, Kozakov D.
2022. Improved Docking of Protein Models by a Combination of Alphafold2 and ClusPro. bioRxiv
2021.09.07.459290.

20. Johnson G, Wu T Te. 2000. Kabat Database and its applications: 30 years after the first variability plot.
Nucleic Acids Res 28:214–218.

21. Warr GW, Clem LW, Söderhäll K. 2003. The international imMunoGeneTics database IMGT. Dev Comp
Immunol 27:1.

22. Zhao N, Kamijo K, Fox PD, Oda H, Morisaki T, Sato Y, Kimura H, Stasevich TJ. 2019. A genetically encoded
probe for imaging nascent and mature HA-tagged proteins in vivo. Nat Commun 10.

23. Mirdita M, Schütze K, Moriwaki Y, Heo L, Ovchinnikov S, Steinegger M. 2022. ColabFold: making protein
folding accessible to all. Nat Methods 19:679–682.

24. Desta IT, Kotelnikov S, Jones G, Ghani U, Abyzov M, Kholodov Y, Standley DM, Beglov D, Vajda S,
Kozakov D. 2023. The ClusPro AbEMap web server for the prediction of antibody epitopes. Nat Protoc
18.

25. Zeng Y, Wei Z, Yuan Q, Chen S, Yu W, Lu Y, Gao J, Yang Y. 2023. Identifying B-cell epitopes using
AlphaFold2 predicted structures and pretrained language model. Bioinformatics 39.

26. Desta IT, Kotelnikov S, Jones G, Ghani U, Abyzov M, Kholodov Y, Standley DM, Sabitova M, Beglov D,
Vajda S, Kozakov D. 2023. Mapping of antibody epitopes based on docking and homology modeling.
Proteins Struct Funct Bioinforma 91:171–182.

27. Lin Z, Akin H, Rao R, Hie B, Zhu Z, Lu W, Smetanin N, Verkuil R, Kabeli O, Shmueli Y, dos Santos Costa A,
Fazel-Zarandi M, Sercu T, Candido S, Rives A. 2023. Evolutionary-scale prediction of atomic-level protein
structure with a language model. Science (80-) 379:1123–1130.

28. Ahdritz G, Bouatta N, Kadyan S, Xia Q, Gerecke W, O TJ, Berenberg D, Fisk I, Zanichelli N, Zhang B,
Nowaczynski A, Wang B, Stepniewska-Dziubinska MM, Zhang S, Ojewole A, Efe Guney M, Biderman S,
Watkins AM, Ra S, Ribalta Lorenzo P, Nivon L, Weitzner B, Andrew Ban Y-E, Sorger PK, Mostaque E,
Zhang Z, Bonneau R, AlQuraishi M, Allen Hamilton B, Bio C. 2022. OpenFold: Retraining AlphaFold2
yields new insights into its learning mechanisms and capacity for generalization. bioRxiv
2022.11.20.517210.

29. Lee JH, Yadollahpour P, Watkins A, Frey NC, Leaver-Fay A, Ra S, Cho K, Gligorijevi´cgligorijevi´c V, Regev
A, Bonneau R. 2023. EquiFold: Protein Structure Prediction with a Novel Coarse-Grained Structure
Representation. bioRxiv 2022.10.07.511322.

30. Ruffolo JA, Gray JJ. 2022. Fast, accurate antibody structure prediction from deep learning on massive set
of natural antibodies. Biophys J 121:155a-156a.

31. Abanades B, Wong WK, Boyles F, Georges G, Bujotzek A, Deane CM. 2023. ImmuneBuilder: Deep-
Learning models for predicting the structures of immune proteins. Commun Biol 6:575.

32.   Ruffolo JA, Sulam J, Gray JJ. 2022. Antibody structure prediction using interpretable deep learning.
651   Patterns 3:100406.
652   33.   Mariani V, Biasini M, Barbato A, Schwede T. 2013. IDDT: A local superposition-free score for comparing
653   protein structures and models using distance difference tests. Bioinformatics 29:2722–2728.
654   34.   Terry JS, Anderson LB, Scherman MS, McAlister CE, Perera R, Schountz T, Geiss BJ. 2021. Development
655   of a SARS-CoV-2 nucleocapsid specific monoclonal antibody. Virology 558:28–37.
656   35.   DeLuca KF, Mick JE, DeLuca JG. 2022. Production and purification of recombinant monoclonal
657   antibodies from human cells based on a primary sequence. STAR Protoc 3:101915.
658   36.   Interactions P. 2002. A Single-Chain Antibody / Epitope System for Functional Analysis of. Society
659   12729–12738.
660   37.   Krauß N, Wessner H, Welfle K, Welfle H, Scholz C, Seifert M, Zubow K, Aÿ J, Hahn M, Scheerer P, Skerra
661   A, Höhne W. 2008. The structure of the anti-c-myc antibody 9E10 Fab fragment/epitope peptide
662   complex reveals a novel binding mode dominated by the heavy chain hypervariable loops. Proteins
663   Struct Funct Genet 73:552–565.
664   38.   Sato Y, Kujirai T, Arai R, Asakawa H, Ohtsuki C, Horikoshi N, Yamagata K, Ueda J, Nagase T, Haraguchi T,
665   Hiraoka Y, Kimura A, Kurumizaka H, Kimura H. 2016. A Genetically Encoded Probe for Live-Cell Imaging
666   of H4K20 Monomethylation. J Mol Biol 428:3885–3902.
667   39.   Fujiwara K, Poikonen K, Aleman L, Valtavaara M, Saksela K, Mayer BJ. 2002. A single-chain
668   antibody/epitope system for functional analysis of protein-protein interactions. Biochemistry 41:12729–
669   38.
670   40.   Churchill MEA, Stura EA, Pinilla C, Appel JR, Houghten RA, Kono DH, Balderas RS, Fieser GG, Schulze-
671   Gahmen U, Wilson IA. 1994. Crystal structure of a peptide complex of anti-influenza peptide antibody
672   Fab 26/9: Comparison of two different antibodies bound to the same peptide antigen. J Mol Biol.
673   41.   Pace CN, Scholtz JM. 1998. A helix propensity scale based on experimental studies of peptides and
674   proteins. Biophys J 75:422–427.
675   42.   Polonsky K, Pupko T, Freund NT. 2023. Evaluation of the Ability of AlphaFold to Predict the Three-
676   Dimensional Structures of Antibodies and Epitopes. J Immunol 211:1578–1588.
677   43.   Guarra F, Colombo G. 2023. Computational Methods in Immunology and Vaccinology: Design and
678   Development of Antibodies and Immunogens. J Chem Theory Comput 19:5315–5333.
679   44.   Giulini M, Schneider C, Cutting D, Desai N, Deane CM, Bonvin AMJJ. 2023. Towards the accurate
680   modelling of antibody-antigen complexes from sequence using machine learning and information-
681   driven docking. bioRxiv 2023.11.17.567543.
682   45.   Hummer AM, Abanades B, Deane CM. 2022. Advances in computational structure-based antibody
683   design. Curr Opin Struct Biol 74:102379.
684   46.   Shashkova TI, Umerenkov D, Salnikov M, Strashnov P V., Konstantinova A V., Lebed I, Shcherbinin DN,
685   Asatryan MN, Kardymon OL, Ivanisenko N V. 2022. SEMA: Antigen B-cell conformational epitope
686   prediction using deep transfer learning. Front Immunol 13:1–11.
687   47.   Lo YT, Shih TC, Pai TW, Ho LP, Wu JL, Chou HY. 2021. Conformational epitope matching and prediction
688   based on protein surface spiral features. BMC Genomics 22:1–16.
689   48.   Mirdita M, Steinegger M, Breitwieser F, Söding J, Levy Karin E. 2021. Fast and sensitive taxonomic
690   assignment to metagenomic contigs. Bioinformatics 37:3029–3031.
691   49.   Kabsch W. 1978. A solution for the best rotation to relate two sets of vectors. Acta Crystallogr Sect A
692   34:827–828.
693   50.   Lawrence J, Bernal J, Witzgall C. 2019. A purely algebraic justification of the Kabsch-Umeyama

694   algorithm. J Res Natl Inst Stand Technol 124:1–6.

Supporting Information

Contents:

Supplemental Table 1A

>mBG17 scFv

MAEVKLEESGGGLVQPGGSMKFSCVASGFTFSDYWMNWVRQSPDKGLEWVAEIRLKSNNYATHYAASVKGRFTISRDDSKSS

VYLQMNNLRAEDSGIYYCTRSAMDYWGQGTSVTVSSGGGGSGGGGSGGGGSDIVMSQSPSSLAVSVGEKITMSCKSSQSLLY

TSDQKNYLAWFQQKPGQSPKLLIFWASTRDSGVPDRFTGSGSGTDFTLTISSVKAEDLAVYYCQQFYNYPRTFGGGTKLEI

>mBG17-15F11

MAEVKLVESGGGLVKPGGSLKLSCAASGFTFSDYWMNWVRQTPEKRLEWVAEIRLKSNNYATHYAASVKGRFTISRDNAKNTL

YLQMSSLRSEDTAIYYCARSAMDYWGQGTTLTVSSGGGGSGGGGSGGGGSDIVLTQSPASLTVSLGQRATISCKSSQSLLYTSDQ

KNYLAWYQQKPGQPPKLLIYWASTRDSGIPARFSGSGSGTDFTLNIHPVEEEDAATYYCQQFYNYPRTFGAGTKLEI

>mBG17-2E2

MAEVQLVESGGDLVKPGGSLKLSCAASGFTFSDYWMNWVRQTPDKRLEWVAEIRLKSNNYATHYAASVKGRFTISRDNAKNTL

YLQMSSLKSEDTAMYYCARSAMDYWGQGTSVTVSSGGGGSGGGGSGGGGSDIVLTQSPASLAVSLGQRATISCKSSQSLLYTS

DQKNYLAWYQQKPGQPPKLLIYWASTRDSGIPARFSGSGSGTDFTLNIHPVEEEDAATYYCQQFYNYPRTFGGGTKLEI

>mBG17 Fab VH:VL

MYLGLNCVFIVFLLKGVQSEVKLEESGGGLVQPGGSMKFSCVASGFTFSDYWMNWVRQSPDKGLEWVAEIRLKSNNYATHYA

ASVKGRFTISRDDSKSSVYLQMNNLRAEDSGIYYCTRSAMDYWGQGTSVTVSS:MDSQAQVLMLLLLWVSGTCGDIVMSQSP

SSLAVSVGEKITMSCKSSQSLLYTSDQKNYLAWFQQKPGQSPKLLIFWASTRDSGVPDRFTGSGS

>mBG17 epitope

DDFSKQLQQS

>mBG17 target protein sequence – SARS CoV-2 Nucleocapsid protein

MSDNGPQNQRNAPRITFGGPSDSTGSNQNGERSGARSKQRRPQGLPNNTASWFTALTQHGKEDLKFPRGQGVPINTNSSPD

DQIGYYRRATRRIRGGDGKMKDLSPRWYFYYLGTGPEAGLPYGANKDGIIWVATEGALNTPKDHIGTRNPANNAAIVLQLPQG

TTLPKGFYAEGSRGGSQASSRSSSRSRNSSRNSTPGSSRGTSPARMAGNGGDAALALLLLDRLNQLESKMSGKGQQQQGQTV

TKKSAAEASKKPRQKRTATKAYNVTQAFGRRGPEQTQGNFGDQELIRQGTDYKHWPQIAQFAPSASAFFGMSRIGMEVTPSG

TWLTYTGAIKLDDKDPNFKDQVILLNKHIDAYKTFPPTEPKKDKKKKADETQALPQRQKKQQTVTLLPAADLDDFSKQLQQSMS

SADSTQA

>HA scFv

MAEVKLVESGGDLVKPGGSLKLSCAASGFTFSSYGMSWVRQTPDKRLEWVATISRGGSYTYYPDSVKGRFTISRDNAKNTLYLQ

MSSLKSEDTAMYYCARRETYDEKGFAYWGQGTTVTVSSGGGGSGGGGSGGGGSDIELTQSPSSLTVTAGEKVTMSCKSSQSLL

NSGNQKNYLTWYQQKPGQPPKLLIYWASTRESGVPDRFTGSGSGRDFTLTISSVQAEDLAVYYCQNDNSHPLTFGAGTKLEL

>HA-15F11

MEVKLVESGGGLVKPGGSLKLSCAASGFTFSSYGMSWVRQTPEKRLEWVATISRGGSYTYYPDSVKGRFTISRDNAKNTLYLQM

SSLRSEDTAIYYCARRETYDEKGFAYWGQGTTLTVSSGGGGSGGGGSGGGGSDIVLTQSPASLTVSLGQRATISCKSSQSLLNSGN

QKNYLTWYQQKPGQPPKLLIYWASTRESGIPARFSGSGSGTDFTLNIHPVEEEDAATYYCQNDNSHPLTFGAGTKLEI

>HA-2E2

MAEVQLVESGGDLVKPGGSLKLSCAASGFTFSSYGMSWVRQTPDKRLEWVATISRGGSYTYYPDSVKGRFTISRDNAKNTLYLQ
MSSLKSEDTAMYYCARRETYDEKGFAYWGQGTSVTVSSGGGGSGGGGSGGGGSDIVLTQSPASLAVSLGQRATISCKSSQSLLN
SGNQKNYLTWYQQKPGQPPKLLIYWASTRESGIPARFSGSGSGTDFTLNIHPVEEEDAATYYCQNDNSHPLTFGGGTKLEI
770
>HA target protein sequence – influenza hemmaglutanin A
MKTIIALSYILCLVSAQKLPGSENRTATLCLGHHAVQNGTLVKTITNDQIEVTNATELVQSSSTGRICDNPHRVLDGRDCTLIDALLG
DPHCDSFQNKEWDLFIERSKAYSNCYPYDVPDYASLRSLVASSGTLEFTTEGFDWTGVTQNGTSYSCKRGSANSFFSRLNWLHK
LNYKYPAQNVTMPNDDKFDKLYIWGVHHPSTDNDQTSLYVQTSGRVTVSTKRSQQTVVPDIGSRPWVRGISSRISIHWTIVKP
GDILLINSTGNLIAPRGYFKIRNGKSSIMKSDALIGNCNSECITPNGSIPNDKPFQNVNRITYGDCPRYVKQSTLKLATGMRNVPEK
QTRGIFGAIAGFIENGWEGMVDGWYGFRHRNSEGTGQAADLKSTQAAIDQINGKLNRLIKKTNEKFHQIEKEFSEVEGRIQDLE
KYVEDTKVDLWSYNAELLVALENQHTIDLTDSEMNKLFERTRKQLRENAEDMGNGCFKIYHRCDNACIGSIRNGTYNHNVYRD
EALNNRFKIKGVELKSGYKDWILWISFAISCFLLCVGLMGLIMWTCQKGNIRCIRCNICH
779
780 >HA epitope
781 YPYDVPDYA
782
783 >Myc scFv
784 MEVKLVESGGDLVKPGGSLKLSCAASGFTFSHYGMSWVRQTPDKRLEWVATIGSRGTYTHYPDSVKGRFTISRDNDKNALYLQ
785 MNSLKSEDTAMYYCARRSEFYYYGNTYYYSAMDYWGQGASVTVSSGGGGSGGGGSGGGGSDIVLTQSPASLAVSLGQRATIS
786 CRASESVDNYGFSFMNWFQQKPGQPPKLLIYAISNRGSGVPARFSGSGSGTDFSLNIHPVEEDDPAMYFCQQTKEVPWTFGG
787 GTKLEI
788
789 >Myc-15F11
790 MEVKLVESGGGLVKPGGSLKLSCAASGFTFSHYGMSWVRQTPEKRLEWVATIGSRGTYTHYPDSVKGRFTISRDNAKNTLYLQ
791 MSSLRSEDTAIYYCARRSEFYYYGNTYYYSAMDYWGQGTTLTVSSGGGGSGGGGSGGGGSDIVLTQSPASLTVSLGQRATISCR
792 ASESVDNYGFSFMNWYQQKPGQPPKLLIYAISNRGSGIPARFSGSGSGTDFTLNIHPVEEEDAATYYCQQTKEVPWTFGAGTKL
793 EI
794
795 >Myc-2E2
796 MAEVQLVESGGDLVKPGGSLKLSCAASGFTFSHYGMSWVRQTPDKRLEWVATIGSRGTYTHYPDSVKGRFTISRDNAKNTLYL
797 QMSSLKSEDTAMYYCARRSEFYYYGNTYYYSAMDYWGQGTSVTVSSGGGGSGGGGSGGGGSDIVLTQSPASLAVSLGQRATIS
798 CRASESVDNYGFSFMNWYQQKPGQPPKLLIYAISNRGSGIPARFSGSGSGTDFTLNIHPVEEEDAATYYCQQTKEVPWTFGGGT
799 KLEI
800
>Myc target protein sequence
MDFFRVVENQPPATMPLNVSFTNRNYDLDYDSVQPYFYCDEEENFYQQQQQSELQPPAPSEDIWKKFELLPTPPLSPSRRSGL
CSPSYVAVTPFSLRGDNDGGGGSFSTADQLEMVTELLGGDMVNQSFICDPDDETFIKNIIIQDCMWSGFSAAAKLVSEKLASYQ
AARKDSGSPNPARGHSVCSTSSLYLQDLSAAASECIDPSVVFPYPLNDSSSPKSCASQDSSAFSPSSDSLLSSTESSPQGSPEPLVL
HEETPPTTSSDSEEEQEDEEEIDVVSVEKRQAPGKRSESGSPSAGGHSKPPHSPLVLKRCHVSTHQHNYAAPPSTRKDYPAAKRV
KLDSVRVLRQISNNRKCTSPRSSDTEENVKRRTHNVLERQRRNELKRSFFALRDQIPELENNEKAPKVVILKKATAYILSVQAEEQK
LISEEDLLRKRREQLKHKLEQLRNSCA
808
809 >Myc epitope
810 EQKLISEEDL

Supplemental Table 1B

```
Kabat numbering       1--------10--------20--------30--------40--------50-------
MYC           -MEVKLVESGGDLVKPGGSLKLSCAASGFTFSHYGMSWVRQTPDKRLEWVATIG--SRGT
MYC-2E2       MAEVQLVESGGDLVKPGGSLKLSCAASGFTFSHYGMSWVRQTPDKRLEWVATIG--SRGT
MYC-15F11     -MEVKLVESGGGLVKPGGSLKLSCAASGFTFSHYGMSWVRQTPEKRLEWVATIG--SRGT
mBG17         MAEVKLEESGGGLVQPGGSMKFSCVASGFTFSDYWMNWVRQSPDKGLEWVAEIRLKSNNY
mBG17-2E2     MAEVQLVESGGDLVKPGGSLKLSCAASGFTFSDYWMNWVRQTPDKRLEWVAEIRLKSNNY
mBG17-15F11   MAEVKLVESGGGLVKPGGSLKLSCAASGFTFSDYWMNWVRQTPEKRLEWVAEIRLKSNNY
HA-scFv       MAEVKLVESGGDLVKPGGSLKLSCAASGFTFSSYGMSWVRQTPDKRLEWVATISRG--GS
HA-2E2        MAEVQLVESGGDLVKPGGSLKLSCAASGFTFSSYGMSWVRQTPDKRLEWVATISRG--GS
HA-15F11      -MEVKLVESGGGLVKPGGSLKLSCAASGFTFSSYGMSWVRQTPEKRLEWVATISRG--GS

Kabat numbering       ---------65---70--------80-----------90---95---------------102
MYC           YTHYPDSVKGRFTISRDNDKNALYLQMNSLKSEDTAMYYCARRSEFYYYGNTYYYSAMDY
MYC-2E2       YTHYPDSVKGRFTISRDNAKNTLYLQMSSLKSEDTAMYYCARRSEFYYYGNTYYYSAMDY
MYC-15F11     YTHYPDSVKGRFTISRDNAKNTLYLQMSSLRSEDTAIYYCARRSEFYYYGNTYYYSAMDY
mBG17         ATHYAASVKGRFTISRDDSKSSVYLQMNNLRAEDSGIYYCTRS-------------AMDY
mBG17-2E2     ATHYAASVKGRFTISRDNAKNTLYLQMNSLKSEDTAMYYCARS-------------AMDY
mBG17-15F11   ATHYAASVKGRFTISRDNAKNTLYLQMSSLRSEDTAIYYCARS-------------AMDY
HA-scFv       YTYYPDSVKGRFTISRDNAKNTLYLQMSSLKSEDTAMYYCARRET-------YDEKGFAY
HA-2E2        YTYYPDSVKGRFTISRDNAKNTLYLQMSSLKSEDTAMYYCARRET-------YDEKGFAY
HA-15F11      YTYYPDSVKGRFTISRDNAKNTLYLQMSSLRSEDTAIYYCARRET-------YDEKGFAY

              -------110-                1--------10------------24---------
MYC           WGQGASVTVSSGGGGSGGGGSGGGGSDIVLTQSPASLAVSLGQRATISCRASESVDNYG
MYC-2E2       WGQGTSVTVSSGGGGSGGGGSGGGGSDIVLTQSPASLAVSLGQRATISCRASESVDNYG
MYC-15F11     WGQGTTLTVSSGGGGSGGGGSGGGGSDIVLTQSPASLTVSLGQRATISCRASESVDNYG
mBG17         WGQGTSVTVSSGGGGSGGGGSGGGGSDIVMSQSPSSLAVSVGEKITMSCKSSQSLLYTSD
mBG17-2E2     WGQGTSVTVSSGGGGSGGGGSGGGGSDIVLTQSPASLAVSLGQRATISCKSSQSLLYTSD
mBG17-15F11   WGQGTTLTVSSGGGGSGGGGSGGGGSDIVLTQSPASLTVSLGQRATISCKSSQSLLYTSD
HA-scFv       WGQGTTVTVSSGGGGSGGGGSGGGGSDIELTQSPSSLTVTAGEKVTMSCKSSQSLLNSGN
HA-2E2        WGQGTSVTVSSGGGGSGGGGSGGGGSDIVLTQSPASLAVSLGQRATISCKSSQSLLNSGN
HA-15F11      WGQGTTLTVSSGGGGSGGGGSGGGGSDIVLTQSPASLTVSLGQRATISCKSSQSLLNSGN

              -----34----40--------50--------60--------70--------80-------
MYC           -FSFMNWFQQKPGQPPKLLIYAISNRGSGVPARFSGSGSGTDFSLNIHPVEEDDPAMYFC
MYC-2E2       -FSFMNWYQQKPGQPPKLLIYAISNRGSGIPARFSGSGSGTDFTLNIHPVEEEDAATYYC
MYC-15F11     -FSFMNWYQQKPGQPPKLLIYAISNRGSGIPARFSGSGSGTDFTLNIHPVEEEDAATYYC
mBG17         QKNYLAWFQQKPGQSPKLLIFWASTRDSGVPDRFTGSGSGTDFTLTISSVKAEDLAVYYC
mBG17-2E2     QKNYLAWYQQKPGQPPKLLIYWASTRDSGIPARFSGSGSGTDFTLNIHPVEEEDAATYYC
mBG17-15F11   QKNYLAWYQQKPGQPPKLLIYWASTRDSGIPARFSGSGSGTDFTLNIHPVEEEDAATYYC
HA-scFv       QKNYLTWYQQKPGQPPKLLIYWASTRESGVPDRFTGSGSGRDFTLTISSVQAEDLAVYYC
HA-2E2        QKNYLTWYQQKPGQPPKLLIYWASTRESGIPARFSGSGSGTDFTLNIHPVEEEDAATYYC
HA-15F11      QKNYLTWYQQKPGQPPKLLIYWASTRESGIPARFSGSGSGTDFTLNIHPVEEEDAATYYC

              -90--------100----
MYC           QQTKEVPWTFGGGTKLEI
MYC-2E2       QQTKEVPWTFGGGTKLEI
MYC-15F11     QQTKEVPWTFGAGTKLEI
mBG17         QQFYNYPRTFGGGTKLEI
mBG17-2E2     QQFYNYPRTFGGGTKLEI
mBG17-15F11   QQFYNYPRTFGAGTKLEI
HA-scFv       QNDNSHPLTFGAGTKLEL
HA-2E2        QNDNSHPLTFGGGTKLEI
HA-15F11      QNDNSHPLTFGAGTKLEI
```

Legend: Heavy chain loops   linker   Light chain loops

A

Myc scFv WT Myc scFv 15F11 Myc scFv 2E2

Myc Fab
PDB entry 2orb

Myc Fab
PDB entry 2orb

Myc Fab
PDB entry 2orb

B

Structure	RMSD$_{C\alpha}$ [Å] vs 2orb
Myc scFv WT	0.42
Myc scFv 15F11	0.45
Myc scFv 2E2	0.45

815
816 **Supplemental Figure 1. Alignment of AlphaFold2 predicted scFv structures to an anti-c-Myc Fab crystal structure. A)** Alignments of
817 AlphaFold2-derived wild-type Myc scFv, Myc-2E2 scFv, and Myc-15F11 scFv structures with a Myc Fab crystal structure (PDB: 2orb).
818 Predicted scFv structures are shown in dark blue, 2orb Myc Fab structures are shown in light blue. **B)** RMSD values comparing
819 structural similarities between the wild-type Myc scFv, Myc-2E2 scFv, and Myc-15F11 scFv structures with a Myc Fab crystal structure
820 (PDB: 2orb) were computed by the PyMOL align command.
821

822
823 **Supplemental Figure 2**: Alphafold2's best attempt to dock whole sequences with the respective sequence's scFv. **A)** The whole HA
824 protein structure and scFv complex as predicted by AF2, with the correct epitope sequence highlighted in magenta. **B)** Shows the
825 same structure by highlighted by confidence (pLDDT) of the structure with AF2. Similarly, the entire Myc protein-scFv complex are
826 shown with **C)** the correct epitope highlighted in magenta and **D)** the confidence of the structure shown, and again for the mBG17 N-
827 protein-scFv complex in **E)** and **F)**.
828

Supplemental Figure 3: AlphaFold2 places all peptides near the CDR loops. The predicted Cα coordinates for all scFv (excluding the
flexible linker) were extracted, and all were aligned together using the Kabsch algorithm (49, 50). With the scFvs structurally aligned,
an all-against-all RMSD was calculated for the epitope peptides. To visually represent each peptide as a single point, the coordinates
for all epitope atoms were averaged. The "central" exemplar epitope (cyan) is the peptide with the smallest sum of RMSD to all other
peptides. **A)** The average and quartile for peptide placement relative to the central peptide via Box-and-Whisker plot reveals that
AlphaFold2 largely places all epitopes in the same area. The Myc CDRH3 runs through the middle of a traditional paratope pocket, it
isn't a "cradle" for the epitope to sit on. AlphaFold2 places peptides on both sides of the CDRH3, causing significant spread in the
peptide placement. **B)** An example of an exemplar, most-central predicted peptide structure (cyan) for the peptide PKSCASQDSS
(cyan) bound to the Myc-2E2 scFv (green) that is distant from an example outlier peptide (magenta, peptide PHSPLVLKRC, center-to-
center distance 14.8 Å). All peptide placements are still in contact with CDRH3, consistent with a strong AlphaFold2 bias to place
peptides in a typical antibody binding site. **C)** The Myc-2E2 scFv (pale-green) and the average epitope placement (cyan) peptide
alongside the crystal structure solution of the Myc epitope (grey). Remaining peptide placements are represented as a cloud of
spheres at the mean peptide position. Each peptide sphere is colored and sized by epitope pLDDT (ranging from 20 to 90). Although

AlphaFold2 frequently placed peptides on the opposite side of the CDRH3 from the Myc epitope (grey), it was not confident in these
peptide placements (low, small, blue pLDDT spheres). In contrast, some of the peptides placed around the CDRH3, and in positions
similar to the native epitope (grey) were placed with higher pLDDT confidence (increasingly large spheres trending from green to
yellow to orange and red). **D)** The top ranked peptide as predicted by PAbFold with sequence QKLISEEDLL (red) and the crystal
structure solution of the Myc epitope (grey).
848

scFv	Apo			Docked		
	BB Ca RMSD	Loop all backbone RMSD	Epitope all atom RMSD	BB Ca RMSD	Loop all backbone RMSD	Epitope all atom RMSD
Myc	0.65	2.87	NA	0.47	1.75	6.69
Myc-15F11	0.62	3.06	NA	0.51	1.51	2.45
Myc-2E2	0.61	2.96	NA	0.51	1.61	2.68

scFv	Apo			Docked		
	BB Ca RMSD	Loop all backbone RMSD	Epitope all atom RMSD	BB Ca RMSD	Loop all backbone RMSD	Epitope all atom RMSD
HA	0.56	1.39	NA	0.58	1.25	3.2
HA-15F11	0.56	1.32	NA	0.6	1.26	3.1
HA-2E2	0.58	1.21	NA	0.6	1.27	3.1

850
851

Supplemental Figure 4: RMSD comparison (all numbers have units of Å) for AlphaFold2 predicted scFv structures compared to reference crystal structures, **A)** 2or9 (Myc) and **B)** 1frg (HA), respectively. The loops of the scFv more closely mimic the crystal structure when the epitope peptide is present. The backbone also undergoes subtle changes during docking that make it slightly more similar to the crystal structure. These structures were aligned by identifying the framework residues in all structures, then aligning the framework region Cα with the Kabsch algorithm (49, 50). Specifically excluded from this process were the heavy and light CDR loops of the structures, as well as the flexible linker structure that connects the heavy and light chains due to the inherent floppy, unstructured nature of this region. After aligning the framework regions of the AlphaFold2 predicted structures and the crystal structures (2or9 and 1frg respectively), an RMSD of these Cα was calculated and is reported as the first column 'BB Cα RMSD'. Without further alignment, loop placement was analyzed with an all backbone RMSD by calculating the RMSD between the C, Cα, N, and O along the backbone of all residues in the scFv that were not used for the framework superimposition. This RMSD is reported in the second column as 'Loop all backbone RMSD'. Finally, to investigate peptide predicted placement and potential scFv:epitope interactions, an all-atom RMSD was calculated between the crystal structure and the AF2 predicted peptide structure (no additional alignment). Because the apo structure lacks a peptide position, this is only reported in the 'Docked' category and is in the 3[rd] column labeled 'Epitope all atom RMSD'. One script was written for each scFv (Myc and HA), and can be found in the Zenodo deposition of our data (https://zenodo.org/records/10884181) because this analysis is not a key part of PAbFold. Briefly this analysis reveals that all three HA scFv variants have predicted framework regions and loop regions in the apo structures that closely match the reference structure (0.56-0.58 Å and 1.21-1.39 Å). Accordingly, when the cognate epitope peptide is present, it can be placed with relatively high accuracy for all three scFvs (3.1-3.2 Å), with only small changes in the loops (1.39 Å to 1.25 Å, 1.32 Å to 1.26 Å, and 1.21 Å to 1.27 Å). In contrast, the apo structures for the three Myc scFvs have a much higher deviation in the loop regions (2.87 to 3.06 Å). When the epitope peptide is added, there is significant motion in the loops consistent with an "induced fit" description. In the two chimeric Myc scFvs (Myc-15F11 and Myc-2E2) the final loop RMSD is reduced to 1.51-1.61 Å, and the epitope peptide is successfully predicted (2.45-2.68 Å). However, despite a lower apo-state loop RMSD (2.87 Å), the loop RMSD for the wild-type Myc scFv only drops to 1.75 Å, and the epitope peptide placement does not match the experimental structure (6.69 Å). This is consistent with the failure of the wild-type Myc scFv AlphaFold2 predictions in main text Figure 2.

878
879
Supplemental Figure 5. Assessment of peptide size and sliding window sizes on epitope prediction efficacy. Myc-2E2 scFv:peptide
structures were predicted with peptides of 8 (**A**), 9 (**B**), 10 (**C**), 11 (**D**), and 12 (**E**) amino acid lengths derived from the Myc protein
with a sliding window of 2 amino acids, and pLDDT scores from each predicted structure were plotted against the Myc amino acid
position and sliding window length target. **F)** Negative control peptides bind to antibody binding sites, but with poor pLDDT scores.
Similarly, with a fixed peptide length of 10 and a sliding window step size of 1 (**F**), 2 (**G**), and 5 (**H**), we can see the practical epitope
detection outcome was similar for a sliding window of 1 and 2, but resolution and accuracy were reduced for a sliding window step
size of 5. To more fully illustrate the strong learned bias that AlphaFold2 has for placing any peptides among the CDR loops, we
predicted the structure of Myc-2E2 in complex with several control peptides. These negative control peptides bind to the generally
expected antibody binding site, but with poor pLDDT. **I)** GSx5 in magenta (GSGSGSGSGS) had a score (mean peptide from Simple Max
method pLDDT) of 29.5. (GGGGS)$_2$ in orange (GGGGSGGGGS) had a score of 31.9. G$_{10}$ in red (GGGGGGGGGG) had a score of 33.
Lastly, **J)** A$_{10}$ in cyan (AAAAAAAAAA) had a score of 41 and is the only negative control peptide to have an alpha-helical secondary
structure (presumably due to the increased alpha helical propensity of alanine).

892
893
894 **Supplemental Figure 6: PAbFold epitope detection is independent of position within target sequence.** The Myc epitope
895 (EQKLISEEDL) was added into the beginning, middle, or end of the 99-a.a. HIV protease sequence (Genbank Accession: NP_705926.1)
896 prior to epitope scanning structure prediction. Positions of the Myc epitope sequence added to in the **A)** N-terminus **B)** middle and
897 **C)** C-terminus of the HIV protease sequence. **D)** Highlights the ranked sequences recovered from each experiment in A, B, and C.
898
899
900

902
903
Supplemental Figure 7: Alphafold2 can accurately predict the HA linear epitope in different scFv backbones. The anti-HA VH and VL
antibody sequences were used to generate either **A)** wild-type scFv or CDR loop grafted onto the **B)** 15F11 or **C)** 2E2 antibody
backbones. The Influenza A virus hemagglutinin protein sequence (Genbank AUT17530.1) was used as the target antigen and
processed into 10 amino acid overlapping peptides with a 1 amino acid sliding window. The structures for each scFv:peptide pair
were predicted with Alphafold2, and pLDDT values for each scFv:peptide pair are shown. **D)** The top-ranking epitope sequences via
pLDDT scores are reported via the consensus method. Sequence underlining represents overlap with the known HA epitope (HA a.a.
114-125: YDVPDYASL). **E)** The top-ranking epitope sequences via pLDDT scores are reported via the simple max method.
911
912

913
914
915 **Supplemental Figure 8:** A comparison of Alphafold2 multimer version 3 and multimer version 2 applied to the mBG17 system. The
916 experimental epitope, DDFSKQLQQS, is still easily identified with all three scFv backbones (wildtype, 15F11, and 2E2).

917
Supplemental Figure 9: Myc comparison of epitope identification accuracy, comparing model types. Performance variation with
AlphaFold2 model (multiple versions 2 and 3) and MSA versions (most up to date version of the ColabFold MSA server uses UniRef30
(2302) and PDB100 (220517)) vs the old MSA server (when this data was initially generated, ColabFold MSA server used UniRef30
(2202) and PDB70 (220313)). The left column is the WT scFv, the middle column is the CDR loops spliced onto the 15F11 backbone,
and the right column is the CDR loops spliced onto the 2E2 backbone. Performance was ablated when using MM3 and the new MSA,
and significantly degraded when using MM2 with the new MSA. For AF2-MM2 Old MSA, see Figure 2.
924

925
Supplemental Figure 10: HA comparison of epitope identification accuracy, comparing model types. A comparison of the differing
AlphaFold2 models with the Myc system (multimer version 3 and 2) along with a comparison of the new MSA (most up to date
version of the ColabFold MSA server uses UniRef30 (2302) amd PDB100 (220517)) vs the old MSA server (when this data was initially
generated, ColabFold MSA server used UniRef30 (2202) and PDB70 (220313)). The left column is the WT scFv, the middle column is
the CDR loops spliced onto the 15F11 backbone, and the right column is the CDR loops spliced onto the 2E2 backbone. For AF2-MM2
Old MSA, see Supplemental Figure 7.
932
933

Local Fall 2022 remake

Supplemental Figure 11: Local remake of the databases used by the MMSEQS server. Databases were downloaded (UniRef30 (2202) and PDB70 (220313)) and were queried locally to produced MSA's for testing. These runs all were done with the multimer version 2 model of Alphafold 2. The left column is the WT scFv, the middle column is the CDR loops spliced onto the 15F11 backbone, and the right column is the CDR loops spliced onto the 2E2 backbone. The first row is the HA system, the second row is the Myc system, and the final row is the mBG17 system.

MMSEQS 2022 Rebuild

Supplemental Figure 12: Server remake of the MMSEQS databases. The databases were rebuilt by the MMSEQS team UniRef30 (2202) and PDB70 (220313)) on the Colabfold MSA server and were queried produced MSA's for testing. These runs all were done with the multimer version 2 model of Alphafold 2. The left column is the WT scFv, the middle column is the CDR loops spliced onto the 15F11 backbone, and the right column is the CDR loops spliced onto the 2E2 backbone. The first row is the HA system, the second row is the Myc system, and the final row is the mBG17 system.

Single Sequence

948
949
950 **Supplemental Figure 13: Single Sequence mode (no MSA's) of epitope prediction with AF2.** These runs all were done with the
951 multimer version 2 model of Alphafold 2 in single sequence mode (i.e. no MSA was used) as a negative control, to highlight the
952 importance of a quality MSA. The left column is the WT scFv, the middle column is the CDR loops spliced onto the 15F11 backbone,
953 and the right column is the CDR loops spliced onto the 2E2 backbone. The first row is the HA system, the second row is the Myc
954 system, and the final row is the mBG17 system.

Myc-2E2 MSA
Venn Diagram

Supplemental Figure 14: MSA overlap between the 4 generation methods. Here we highlight the number of unique entries that are shared amongst all of the MSA methods, those being: **1)** using the databases right now via colabfold (PDB30 2302 and PDB100 230517) (green) **2)** the databases after they had been accessed via colabfold and cached for repeated use (UniRef30 (2202) and PDB70 (220313)) (yellow), **3)** downloading the databases locally (UniRef30 (2202) and PDB70 (220313)) and attempting to create the MSAs ourselves (red), and **4)** querying the databases after the MMSEQS team rebuilt them for our use via colabfold (UniRef30 (2202) and PDB70 (220313)) (blue).

		MMSEQS 2022 Fall	Local 2022 Rebuild	MMSEQS 2022 Rebuild	MMSEQS 2023 Winter	Single Sequence
HA	WT	M	✔	-	✔	-
	15F11	✔	✔	-	M	-
	2E2	M	✔	-	-	-
Myc	WT	M	✔	-	-	-
	15F11	✔	-	-	-	-
	2E2	✔	-	✔	✔	-
mBG17	WT	✔	✔	✔	✔	-
	15F11	✔	✔	✔	✔	-
	2E2	✔	✔	✔	✔	-

968
969
970 **Supplemental Figure 15: Comparison of how well each MSA generation scheme accurately identified the experimentally derived**
971 **epitope within the top 5 epitope sequences.** A green checkmark shows that it was found by both the consensus model and the top
972 single model, a yellow "M" means the simple max method correctly identified the experimental epitope in the top 5 epitopes, and
973 the red dash means both methods failed. The consensus model did not identify the epitope correctly when the simple max method
974 failed to. The colored background behind the titles is the same color as Supplemental Figure 14 to help guide the eye.