Varying phylogenetic signal in susceptibility to four bacterial pathogens across species of

Drosophilidae

Hongbo Sun[1] *, Mark A. Hanson[1], Sarah K. Walsh[1,2,3], Ryan M. Imrie[1,3], Ben Raymond[1+], Ben

Longdon[1+]

[1] Centre for Ecology and Conservation; University of Exeter; Cornwall; United Kingdom,

[2] Environment and Sustainability Institute; University of Exeter; Cornwall; United Kingdom,

[3] Current address: MRC-University of Glasgow Centre for Virus Research, Glasgow; United

Kingdom.

* Corresponding author.

+ These authors contributed equally to this work.

Author information:

Hongbo Sun* hs766@exeter.ac.uk https://orcid.org/0009-0008-7272-7205

Mark A. Hanson m.hanson@exeter.ac.uk https://orcid.org/0000-0002-6125-3672

Sarah K. Walsh Sarah.Walsh.2@glasgow.ac.uk https://orcid.org/0000-0001-6598-095X

Ryan M. Imrie Ryan.Imrie@glasgow.ac.uk https://orcid.org/0000-0002-5055-2888

Ben Raymond b.raymond@exeter.ac.uk https://orcid.org/0000-0002-3730-0985

Ben Longdon b.longdon2@exeter.ac.uk https://orcid.org/0000-0001-6936-1697

Abstract

Bacterial infections are one of the major sources threatening public health. Pathogen host shifts –

where a pathogen jumps from one host species to another – are important sources of emerging

infectious diseases. However, compared to viruses, we know relatively little about the factors that

determine whether a bacteria can infect a novel host, such as how host phylogenetics constrains

variation in pathogen-host range and the link between host phylogeny and the infectivity and

virulence of a pathogen. Here, we experimentally examined susceptibility to bacterial infections using

a panel of 36 Drosophilidae species and four pathogens (*Providencia rettgeri*, *Pseudomonas*

entomophila, *Enterococcus faecalis*, *Staphylococcus aureus*). The outcomes of infection differed

greatly among pathogens and across host species. The host phylogeny explains a considerable amount

of variation in susceptibility, with the greatest phylogenetic signal for *P. rettgeri* infection, explaining

94% of the variation in mortality. Positive correlations were observed between mortality and bacterial

load for three out of the four pathogens. Correlations in susceptibility between the four pathogens

were positive but largely non-significant, suggesting susceptibility is mostly pathogen-specific. These

results suggest that susceptibility to bacterial pathogens may be predicted by the host phylogeny, but

the effect may vary in magnitude between different bacteria.

Keywords: Bacterial infection, Susceptibility, Phylogeny, Host shift, Host-microbe interaction

Introduction

Emerging and re-emerging infectious diseases are a major threat to public health [1-3]. Multi drug-resistant bacterial infections are a major cause of mortality, responsible for millions of deaths globally [4]. Recent studies have highlighted that host shifts – where a pathogen jumps from one host species to another – are important sources of emerging diseases, not only by viruses, but also by bacterial pathogens [5, 6]. There is an increasing awareness that bacterial pathogens also commonly undergo host shifts [6-8], which has been linked to their recent emergence as causative agents of disease in several host systems, including humans, livestock, wildlife, and plants [7-9]. In *Staphylococcus aureus*, a human pathogen associated with several severe hospital and community-acquired infections [10], humans are the primary transmission hub for host shifts into a range of livestock species, and cows have been considered the main animal reservoirs for human epidemics [11-13]. Another extensively studied example is the jump of *Mycoplasma gallisepticum* from poultry to wild house finches [14]. Here, the *Mycoplasma* host shift has driven the evolution of resistance in the house finch hosts [15], led to changes in bacterial gene content [7] and the evolution of virulence [16]. In plants the bacterial pathogen *Xylella fastidiosa* is found in over 600 host species [17]. A single gene loss event is associated with the host shift of *X. fastidiosa* from coffee plants to olive trees [18]. Most studies on bacterial host shifts are based on genomic data following natural host shifts, and there are limited experimental studies of bacterial host shifts that move beyond two-host systems.

Perhaps the best experimental studies relating to bacterial host shifts come from insects and their bacterial endosymbionts. In a study of coccinellid beetles and a male-killing isolate of *Spiroplasma*, the bacterium's ability to distort offspring sex ratios in novel hosts was negatively correlated with the genetic distance from the bacteria's natural host [19]. Similar patterns have been found for *Wolbachia*, a bacterial endosymbiont that infects ~40-60% of all arthropods [20-22]. Artificial transfers of *Wolbachia* are most successful between closely related hosts [23, 24]. This tallies with field data on *Wolbachia* distributions that found strong effects of the host phylogeny, with high rates of *Wolbachia* sharing between closely related species, and a decline in co-occurrence in different hosts with increasing phylogenetic distance between these hosts [9]. Viruses show similar trends, with

phylogenetic distance and clade effects explaining large amounts of variation in susceptibility to

several viruses [25, 26]. When comparing different viruses from the same family, positive inter-

specific correlations in susceptibility have been observed, suggesting similar mechanisms underlie

differences in susceptibility observed among species [26, 27]. Whether such patterns hold true for

free-living bacteria, which are less tightly linked to their hosts, remains to be tested.

Here, we have used a large cross-infection experiment to understand how susceptibility to bacterial

pathogens varies across host species. We infected a large panel of 36 species of Drosophilidae with

four bacterial pathogens: two Gram-positives, *Staphylococcus aureus*, *Enterococcus faecalis*, and two

Gram-negatives: *Providencia rettgeri* and *Pseudomonas entomophila*. In *Drosophila melanogaster*,

the major immune response to bacterial pathogens involves the Toll and IMD pathways. The Toll

pathway primarily controls resistance to fungal and Gram-positive bacterial infections, whereas the

Imd pathway mediates response to Gram-negative bacterial infections [28, 29]. Both of these *NF-κB*

pathways result in the upregulation of antimicrobial peptides (AMPs) and other defence response

genes [30, 31]. Genes in these pathways evolve rapidly. However, it remains unclear how differences

in immune genes among species contributes to differences in infection outcome [32, 33]. Recent

studies have demonstrated that AMPs show specific importance for certain bacterial infections and

that different alleles of AMPs are linked to differences in susceptibility both within and across species

[34, 35].

The bacterial pathogens used here have been well characterised in *D. melanogaster* [36, 37]. *E.*

faecalis was initially derived from a human clinical isolate [38, 39]. The virulence of *E. faecalis*

varies with different strains [40]. The *E. faecalis* FA2-2(pAM714) used in this study produces toxins

(e.g. cytolysin), that act as virulence factors to improve bacterial colonisation and decrease host

survival. in *Drosophila* [39], the Toll pathway, particularly the host defence peptides Bomanin and

Baramicin, have been shown to play a role in response to *E. faecalis* [41-43]. *S. aureus* was initially

isolated from humans and has been used in mice and *Drosophila* infection models [36, 44]. *S. aureus*

is a multispecies pathogen that can cause disease in humans and livestock. The dynamic gain and loss

of host-specific adaptive genes, usually located on mobile genetic elements (MGEs), enable *S. aureus*

to jump between different species [13, 45]. In *D. melanogaster*, resistance to *S. aureus* specifically

relies on the melanisation response [46, 47]. *P. entomophila* was isolated from a wild female *D.*

melanogaster and is a versatile soil bacterium that can infect *D. melanogaster* as well as insects from

different orders. Upon infection, *D. melanogaster* show differential expression of genes including

AMPs, oxidative stress response genes and detoxification genes [48-51]. Genes involved in host

damage prevention, signalling, renewal, and regulation also affect the susceptibility of *D.*

melanogaster when *D. melanogaster* is orally challenged with *P. entomophila* [52]. *P. entomophila*

causes digestive obstruction and eventually leads to the host's death by damage and perforation of the

gut through the production of insecticidal protease-based toxins [49, 51]. *P. rettgeri* was initially

isolated from the haemolymph of wild *D. melanogaster* [53]. The IMD pathway, and particularly the

AMP Diptericin has been shown to be an important part of the immune response to *P. rettgeri* [35, 54,

55]. A single nucleotide polymorphism at residue 69 in *Diptericin A*, "S69R" of *DptA*, is associated

with increased susceptibility to *P. rettgeri* bacterial infection [34, 35].

Here, we examine the role of the host phylogeny in determining the outcomes of bacterial infections.

We investigated how variation in bacterial virulence varies across the host phylogeny as well as the

relationship between virulence and bacterial load for each of the four bacterial pathogens.

Methodology

Bacterial strains and culture

We used four bacterial pathogens in this study. Two Gram-positive bacteria, *Staphylococcus aureus* PIG1 (*S. aureus*) [44] and *Enterococcus faecalis* FA2-2/pAM714 (*E. faeclis*) [38, 39], and two Gram-negative bacteria: *Providencia rettgeri* Dmel (*P. rettgeri*) [53] and *Pseudomonas entomophila* (*P. entomophila*) [49]. Bacterial cultures were initiated from frozen glycerol stocks onto LB (Lysogeny Broth) agar, with the exception of *P. entomophila,* which were plated on egg yolk agar (10g of agar, 10g of LB, 1 L of water and 20 mL of egg yolk (Sigma)). Colonies of *P. entomophila* with halos on egg yolk agar (indicating phospholipase activity) were used to prepare inocula. Insect inocula were cultured in LB broth, with the exception of *E. faecalis* which used BHI (Brain heart infusion) broth. All bacteria were grown at a 30℃ incubator, shaking 180 rpm/min for liquid culture. Inocula were prepared in a three-step process in order to standardise physiology in the stationary phase. Initially, colonies were used to prepare cultures of 5 mL in 30 mL sterile universals. These were diluted in liquid media to OD_{600} 0.2, and then 50 µL of this was inoculated into new vials. Following overnight growth, cultures were diluted with culture media to the desired OD value (OD 2.0 for *E. faecalis* and OD 1.0 for the other bacteria based on pilot mortality data).

Fly stocks and rearing.

All flies were maintained in multi-generation stock bottles (Fisher brand) at 22°C, ~50% relative humidity in a 12-hour light-dark cycle. Each stock bottle contained 50 ml of one of four varieties of *Drosophila* media which were chosen to optimise rearing conditions (full details in Supplementary Table 1). Male flies were used to avoid any effect of sex or mating status, which has been shown to influence the susceptibility of female flies to various pathogens [56-59]. Before inoculation, 0-1 day old male flies were transferred to vials containing cornmeal media. These flies were then transferred to fresh media every three or four days for a week (age 7–8 days, flip once before infection), at which point they were inoculated. Flies were kept at 22°C, 70% relative humidity in a 12-hour light-dark cycle throughout the experiments. Flies were inoculated under CO_2 anaesthesia via septic pin prick with 10µm diameter stainless steel needles (26002-10, Fine Science Tools, CA, USA). These needles

were bent approximately 250µm from the end to provide a depth stop and dipped in the inoculum

before being pricked into the anepisternal cleft in the thorax of anaesthetised flies. Inoculation by this

method bypasses the gut immune barrier but avoids differences in inoculation dose due to variations

in feeding rate in oral inoculation. This method has been demonstrated to be highly repeatable (see

results) and gives a consistent initial dose (our estimate of initial dose (CFUs) introduced into

individual fly for each pathogen: OD = 1 *P. rettgeri* :$10^{\wedge}(1.94\pm0.08)$; OD = 1 *P. entomophila* :

$10^{\wedge}(1.50\pm0.14)$; OD= 2 *E. faecalis* :$10^{\wedge}(2.67\pm0.12)$; OD = 1 *S. aureus* : $10^{\wedge}(2.75\pm0.12)$. All

inoculations were done by a single experimenter (HS).

For the survival assay, we inoculated male flies of 36 fly species. We inoculated three replicate vials

for 165 species-bacteria combinations, two replicate vials for 13 combinations and one replicate vial

for two combinations. For each combination, one of 36 species was infected with either a control

Ringer's solution [60] or one of the four bacterial pathogens listed above. Vials contained a mean of

14 flies (range = 5-18 flies). Following pin prick infection, the number of dead flies in each vial was

recorded daily for 14 days. Flies were tipped into new vials of food every two days.

We measured bacterial load in 18 species of fly; these species were selected based on their survival

results and covered a broad range of survival values for each pathogen. We measured colony forming

units (CFUs) for individual male flies of each species (mean = 15 replicate flies, range 1-16) infected

with each species-bacteria combination (there is only one replicate for *D. teisseri* in *P. entomophila*

infection (Figure 3) as most of flies died due to infection before collection, there are at least 7 flies per

species with a mean of 15 flies for all other treatments.). CFUs were measured at 16 hours post-

infection. Flies were prepared and infected as above. Before homogenisation, anaesthetised flies were

moved to 96-well tubes containing a 4mm steel ball and 200 µL 70% ethanol, flies were then gently

cleaned using 70% ethanol and homogenised in 200 µL sterile LB or sterile BHI (for flies infected

with *E. faecalis*) in a Tissue Lyser II at 30 Hz for three minutes. Homogenates were diluted 10^{-1} to 10^{-5}

and plated on LB agar or BHI agar in two technical replicates before being cultured overnight at 30°C.

Bacterial loads were measured as colony-forming units (CFUs) per fly, following the formula:

CFU per fly = (Colony counts × Dilution factor × Homogenate volume) / Plating volume.

Host Phylogeny

The method used to infer the host phylogeny has been described elsewhere [61] and details are

provided in the supplementary methods.

Sequencing of *Diptericin A*

To determine the *Diptericin A* allele of each species (which has been linked to susceptibility to

bacterial pathogens [35]), *Diptericin* genes were sequenced across species to determine their S69R

allele (see supplementary methods).

Statistical analysis

We used phylogenetic mixed models to look at the effects of host species relatedness on bacterial

susceptibility across the 36 host species. We fitted all models using a Bayesian approach in the R

package MCMCglmm v2.29 [62] in RStudio (R v4.2.2) [63]. We used a multivariate model with

mortality following inoculation with each of the four bacteria or control treatments as the response

variables. Mortality was calculated as each vial's mean proportion of dead flies over 14 days.

The models took the following form:

(1) $y_{hit} = \beta_{1:t} + u_{p:ht} + u_{s:ht} + e_{hit}$

(2) $y_{hit} = \beta_{1:t} + u_{p:ht} + e_{hit}$

Where y is the mortality of the i^{th} biological replicate of host species h, for treatment t (each of the

four bacteria and the non-infected control). β are the fixed effects, with β_1 being the intercepts for

each trait.

We also ran bivariate models for each of the four pathogens across the 18 host species tested to

investigate how mortality was related to bacterial load. These models had the following form:

$$(3) \quad y_{hit} = \beta_{1:t} + u_{p:ht} + e_{hit}$$

Where y is the response of the i^{th} biological replicate of host species h, for treatment t (either

mortality or the bacterial load (CFUs) of an individual fly infected with that pathogen). U are random

effects for the phylogenetic species effects (p), non-phylogenetic species effects (s) and residuals (e).

$u_{s:ht}$ was removed from models (2) and (3) as the models struggled to separate the phylogenetic and

non-phylogenetic effects. The random effects and the residuals are assumed to be multivariate normal

with a zero mean and a covariance structure $V_p \otimes \mathbf{A}$ for the phylogenetic effects and $V_e \otimes \mathbf{I}$ for the

residuals ($\otimes$ here is the Kronecker product). $\mathbf{A}$ is the phylogenetic relatedness matrix, $\mathbf{I}$ is an identity

matrix and the $\mathbf{V}$ are 5×5 (co)variance matrices describing the (co)variances between mortality when

infected with each of the four bacteria or the control. In model (3) the $\mathbf{V}$ are 2×2 (co)variance matrices

describing the (co)variances between mortality and bacterial load when infected with each of the four

bacteria. Specifically, the matrices V_p and V_s describe the phylogenetic and nonphylogenetic between-

species variances in mortality for each treatment and the covariances between them, whereas the

residual covariance matrix V_e describes within-species variance that includes both actual within-

species effects and measurement error. The off-diagonal elements in V_e (the covariances) are unable to

be estimated because no vial has multiple measurements – so they were set to zero. Model (1) was

also run with two additional fixed effects of body size and *Diptericin A* S69R site status (see

supplementary methods).

The MCMC chain was run for 13 million iterations with a burn-in of 3 million iterations and a

thinning interval of 5000 iterations. We confirmed our experimental design and sample sizes had

sufficient power to detect effects by down sampling a similar dataset [27]. The results presented were

obtained using parameter expanded priors for the $\mathbf{V}_p$ and $\mathbf{V}_s$ matrices [66]. Results were tested for

sensitivity to the use of different priors by being run with different prior structures. The alternative

priors gave qualitatively similar results, except for the correlations where we found mean estimates

were higher with the majority being significant. To determine what may be driving this we simulated

response data based on mean estimates from the parameter expanded prior model using the

'simulate.MCMCglmm' function. We then refitted models on this data with either parameter

expanded or alternative priors (inverse wishart or flat). The alternative priors had higher estimates of

the correlations in the refitted models suggesting they may be inadvertently influencing the estimates.

The proportion of the between-species variance that the phylogeny can explain was calculated from

model (1) using the equation $V_p / (V_p + V_s)$, where V_p and V_s represent the phylogenetic and species-

specific components of between-species variance [67], respectively, and are equivalent to

phylogenetic heritability or Pagel's lambda [68, 69]. The repeatability of bacterial load measurements

was calculated from model (2) as $V_p/(V_p + V_e)$, where V_e is the residual variance of the model [70].

Interspecific correlations were calculated from model (2) and (3) V_p matrix as $cov_{x,y}/ \sqrt{(var_x \times var_y)}$.

Parameter estimates reported are means of the posterior density, and 95% credible intervals (CIs) were

taken to be the 95% highest posterior density intervals.

We ran additional models using Abbots correction on data, and a binomial model on day 14 mortality

(see supplementary methods). We confirmed our results were robust to these different analyses

(supplementary Table S5-S6, Figure S3) and so results presented are the main analysis described

above.

Results

Patterns of susceptibility to bacterial pathogens across host species

To investigate how susceptibility to bacterial pathogens varies across host species, we measured survival in 7402 flies from 36 species of Drosophilidae inoculated with one of four bacterial pathogens or a control. We found large differences in the mean virulence of the bacteria; the mean mortality was 70% for *P. entomophila* infection, 46% for *P. rettgeri*, 39% for *S. aureus*, and 36% for *E. faecalis* (compared to only 4% in control inoculated flies). There were dramatic differences in the survival of the different species with the majority of individuals of certain species dying, whilst others showed little mortality when infected with the same pathogen (Figure 1). For example, just over half of the species showed rapid mortality following *P. entomophila* infection, with over 50% of flies dying within one day post-infection in 19 out of 36 fly species (Figure 1). Conversely, survival remained high for most species infected with *E. faecalis,* with the exceptions of *D. yakuba* and *D. sucinea* where >85% of flies had died by the third day post-infection (Figure 1).

Figure 1. Survival of 36 species of Drosophilidae after infection with four bacterial pathogens.

Each point represents the mean survival of three replicates, colours represent the inoculation

treatment.

The host phylogeny explains varying amounts of variation in mortality

Different clades of hosts show different mortality levels with each bacteria, with hosts with high and low mortality clustering together on the host phylogeny (Figure 2). For example, a clade within the subgenus Drosophila (delineated by *D. micromelanica* down to *D. flavomontana* in Figure 2) shows low mortality when infected with *P. rettgeri* compared to the other major host clades. We used a phylogenetic mixed model to determine the proportion of variation in mortality explained by the host phylogeny for each pathogen (Table 1). The estimates vary among the four pathogens, with the host phylogeny explaining large amounts of variation for three out of four bacterial infections. We find that almost all of the variation in mortality is explained by the phylogeny following *P. rettgeri* infections (94%).

The phylogeny explains a substantial proportion of the variation in mortality for *S. aureus* (64%) but less for *P. entomophila* (36%) and *E. faecalis* (14%). However, these have broad 95% CI's around estimates of the posterior mean, which are pushed up against zero for *P. entomophila* and *E. faecalis*. If we compare the phylogenetic signal in mortality for each of the pathogens we find that the phylogeny explains a greater amount of variation in mortality to *P. rettgeri* infection than for the other pathogens (although the effect is marginal for *P. rettgeri* - *S. aureus*), and *S. aureus* has greater phylogenetic signal than *E. faecalis* (Table S2). Measurements were highly repeatable (Table 1) with a large proportion of the variance being explained by the inter-specific phylogenetic component, with little within species variation or measurement error (Repeatability = $V_p/(V_P + V_e)$).

Pathogen	Variance explained by phylogeny	Repeatability
P. rettgeri	0.94 (95% CI: 0.81, 1.00)	0.84 (95% CI: 0.75,0.93)
P. entomophila	0.36 (95% CI: 0.00, 0.74)	0.70 (95% CI: 0.48,0.89)
E. faecalis	0.14 (95% CI: 0.00, 0.50)	0.90 (95% CI: 0.80,0.97)
S. aureus	0.64 (95% CI: 0.24, 0.94)	0.80 (95% CI: 0.63,0.92)

Table 1. Estimates of the mean proportion of variation explained by the host phylogeny (taken from model (1)), and the repeatability of estimates (taken from model (2)).

Figure 2: Susceptibility to four bacterial pathogens varies across 36 Drosophilidae species. Bar

length and colour represent mean mortality change across pathogen and fly species with error bars

representing standard errors. The phylogeny of Drosophilidae host species is presented on the left,

with the scale bar representing the number of nucleotide substitutions per site. Names of host species

are given on the right; the column label shows different treatments (bacterial or control inoculation).

Patterns of susceptibility are largely distinct for each bacteria

In *D. melanogaster*, the host immune response differs markedly between infections by Gram-positive

and Gram-negative bacteria [71, 72]. If mechanisms of host defence depend on the induced immune

response, we might expect positive correlations in host susceptibility among species for bacteria of the

same Gram type. We found the correlations between bacteria all had positive estimates (Table 2),

although the 95% CIs overlapped zero for all except for the correlations between *S. aureus–E. faecalis*

and *E. faecalis–P. rettgeri.* The correlation coefficient was less than 0.4 for all correlations bar the

estimate between the two Gram-positive bacteria *S. aureus* and *E. faecalis* where $r = 0.66$. Overall,

this suggests that although hosts may show similar trends in susceptibility to all the bacteria tested

(correlations are largely positive); host susceptibility generally differs by pathogen, with shifts in the

rank order of species susceptibilities to the different pathogens (i.e. as correlations are all less than

one, so species do not show overall high or low susceptibity to bacteria *per se*, but rather pathogen

specific susceptibilities).

Bacteria	Inter-specific correlation *r*
S. aureus - E. faecalis	**0.66 (95% CI: 0.37, 0.90)**
S. aureus - P. rettgeri	0.35 (95% CI: -0.04, 0.70)
S. aureus - P. entomophila	0.34 (95% CI: -0.13, 0.73)
E. faecalis - P. rettgeri	**0.39 (95% CI: 0.02, 0.72)**
E. faecalis - P. entomophila	0.31 (95% CI: -0.19, 0.74)
P. rettgeri - P. entomophila	0.37 (95% CI: -0.05, 0.75)
P. rettgeri - Control	-0.06 (95% CI: -0.65, 0.61)
P. entomophila – Control	-0.04 (95% CI: -0.65, 0.66)
E. faecalis - Control	-0.15 (95% CI: -0.85, 0.66)
S. aureus - Control	-0.12 (95% CI: -0.80, 0.64)

Table 2. Correlations in mortality between different pathogen and control treatments. Numbers

represent the mean estimates of correlation coefficients from Model (2) with 95% credible intervals

shown in brackets. Significant correlation coefficients (with 95% CI that do not span zero) are

highlighted in bold.

Mortality correlates with bacterial load in most cases

To test whether increased mortality is due to higher bacterial loads, we selected 18 fly species that varied in their mortality and measured their bacterial load to each pathogen at 16 hours post-infection. We found that 24% of the flies infected by *P. entomophila* had died by 16 hours post-infection (<2% of flies infected with each of the other bacteria died by this time point). We excluded dead flies from the analysis below to avoid confounding bacterial growth rates, as these flies may have died prematurely from injury, and bacterial growth rates may have developed differently in carcasses or flies failing to mount a full immune response. however, we note that the results were robust if we included the bacterial load measured in corpses. We quantified bacterial load in 1328 individual flies (Figure 3) and ran models combining this with the mortality data above for these 18 fly species.

Mortality estimates from this model using a subset of the data from the survival experiment (Table 3) are consistent with observations in survival experiments (Figures 1 and 2). Mean bacterial loads are relatively consistent across the four pathogens (Table 3). We found strong positive correlations between mortality and bacterial load for *P. rettgeri* and *P. entomophila* and a slightly lower positive correlation for *S. aureus* with 95% CI's that just overlap zero (Table 4, Figure S1). However, we found no evidence of a significant positive correlation between mortality and bacterial load for *E. faecalis* (Table 4, Figure S1). However, the 95% CI's around estimates were broad so there was no clear indication that bacteria were behaving differently.

We found some evidence of microbiome proliferation in pathogen infected insects when measuring CFU in four species (*D. americana*, *S. lebanonensis*, *D. putrida* and *S. pattersoni*). We note there was an elevated count of likely microbiome-specific colonies in these species based on colony morphology and growth rate compared to colonies of pathogenic bacteria on LB/BHI agar. We included these colonies in our final bacterial load counts to avoid subjective calls on microbiome or pathogen colony identification, but will note that excluding these species from our analysis did not alter the results qualitatively.

Pathogen	Bacterial load	Mortality
P. rettgeri	4.63 (95% CI: 3.67, 5.50)	0.46 (95% CI: 0.17, 0.76)
P. entomophila	3.78 (95% CI: 2.97, 4.46)	0.62 (95% CI: 0.33, 0.90)
E. faecalis	3.36 (95% CI: 2.80, 3.92)	0.37 (95% CI: -0.02, 0.74)
S. aureus	3.12 (95% CI: 2.32, 3.85)	0.36 (95% CI: 0.06, 0.68)

Table 3. Estimations of mortality and bacterial load for each pathogen. Numbers represent the mean estimates of mortality (mean proportion dead) and bacterial load (CFU per fly on a Log10 scale) for each pathogen from Model (3) with 95% credible intervals shown in brackets.

Figure 3: Mortality and bacterial load to four bacterial pathogens across 18 fly species. Bar

length and colour represent mean mortality (left) or bacterial load (right) across 18 fly species bearing

either *P. rettgeri* (A), *P. entomophila* (B), *E. faecalis* (C) or *S. aureus* (D) infection. Bacterial loads

are data point at 16 hours post infection, with the error bars representing standard errors of a mean of

15 replicate flies for each species. The phylogeny of Drosophilidae host species is presented on the

left, with the scale bar representing the number of nucleotide substitutions per site. Names of host

species are presented on the right.

Pathogen	Inter-specific correlation r
P. rettgeri	0.59 (95% CI's: 0.16, 0.94)
P. entomophila	0.54 (95% CI's: 0.10, 0.90)
E. faecalis	0.22 (95% CI's: -0.43, 0.74)
S. aureus	0.48 (95% CI's: -0.03, 0.93)

Table 4. Correlations between mortality and bacterial load for each pathogen. Numbers represent the mean estimates of correlation coefficients from Model (3) with 95% credible intervals shown in brackets. Significant correlation coefficients are where the 95% CI do not span zero.

Testing for effect of *Diptericin A* allelic variation and body size

We observed an exceptionally high proportion of variance explained by host phylogeny for *P. rettgeri* (Table 1). This suggests some phylogenetically inherited factor, or factors, strongly determines susceptibility to this pathogen. Previous studies linked the gene *Diptericin A* (*DptA*) to defence against *P. rettgeri*, including resistant or susceptible alleles at a polymorphic residue both within and across species [35, 54].

We successfully amplified and sequenced *DptA* from 29 species and annotated the allele status at a previously identified polymorphic residue (residue 69 of *D. melanogaster DptA*). Four residues were found in our panel at this site (Figure S2). These were dominated by the resistant allele encoding serine (S) in 75% of species, and three alternate alleles each found in only 2-3 species, including a known susceptible allele encoding arginine (R), and alleles encoding either glutamine (Q) or limited replication of each individual allele, we fitted a fixed effect of *DptA* residue 69 by pathogen with species either being resistant (S and Q) or susceptible (R and N). We found no significant effect of the susceptible *DptA* alleles for any of the four pathogens (Table S4) but note there was limited power to detect an effect with only five species having susceptible alleles. There was little change in the variance explained by the phylogeny when the *DptA* allele status was included as a fixed effect (Table S4) indicating this allele does not readily explain the phylogenetic patterns observed.

The wing size of *Drosophila* can provide a proxy for body size [73] which is known to be phylogenetically dependent [74]. We included wing size as a fixed effect in models to see if wing size explained some of the variation in host susceptibility. We found no effect of wing size on susceptibility for any pathogen (mean and 95% CIs for each pathogen: *P. rettgeri*: -0.34 (95% CI:-0.75, 0.02), *P. entomophila*: -0.02 (95% CI:-0.54, 0.45), *E. faecalis*: 0.01 (95% CI:-0.37, 0.37), *S. aureus*: 0.03 (-0.37, 0.42)).

Discussion

We examined how susceptibility to bacterial pathogens varies across the host phylogeny, using a panel of 36 Drosophilidae species and four bacterial pathogens. Using phylogenetic mixed models we found the host phylogeny can explain a large proportion of variation in susceptibility, but differs in the extent of variation explained between pathogens. Distinct infection patterns were seen between the different pathogens, with all correlations in susceptibility between the pathogens being less than one, suggesting susceptibility generally varies between bacterial pathogens. However, despite being largely insignificant, all correlations had positive estimates, and the strongest correlation was seen between the two Gram-positive bacteria (Table 2) suggesting that differences in susceptibility may have shared underlying mechanisms. Positive correlations between mortality and bacterial load for *P. rettgeri*, *P. entomophila* and *S. aureus* suggest virulence is a direct result of bacterial accumulation in the host. The lack of correlation between virulence and bacterial load for *E. faecalis* may have a biological underpinning, such as the production of toxins, or inhibited replication capabilities of the human *E. faecalis* strain we used, which may not be well adapted to invertebrate hosts [40]; alternately this may be due to limited statistical power. These results (Figure S1) also demonstrate that different species can have differences in mortality with similar bacterial loads. As such, examination of whether potential tolerance mechanisms underly differences in susceptibility [75] could be an interesting follow up to this work.

Our survival results find a substantial amount of variation in susceptibility can be explained by the host phylogeny, with the largest effect for *P. rettgeri,* where almost all of the variation in susceptibility is explained by the host phylogeny. One potential explanation is that susceptibility is linked to variation in immune genes with certain host clades having lost or gained different aspects of the immune system. Previous studies have shown that a specific AMP, *Diptericin*, plays an essential role in defence against *P. rettgeri* infection. Flies encoding arginine (R) at the *Diptericin* S69R polymorphic site are more susceptible than flies encoding serine (S) [35, 54]. Due to the limited statistical replication of the different *Diptericin* alleles (75% of the 29 species with allele data had the "S" resistance allele, with each of the other 3 alleles only replicated in 2 or 3 species) we had little

power to detect effects here. However, there is also substantial variation in susceptibility just among

species with the "S" allele, suggesting this *Diptericin* allele is not the sole driver of the patterns

observed here. While we did not find an effect of the *Diptericin* S69R allele, we cannot rule out a role

for *Diptericin* in explaining variation in host susceptibility, although other mechanisms may well also

play a role in explaining among species differences in susceptibility. Further study could consider a

broader range of species encoding different variants at the S69R site, variation elsewhere in the

Diptericin gene, or differences in induction of *Diptericin* and other AMPs across species given rapid

induction is important for host survival [35, 36, 54].

We found that among species correlations in susceptibility between bacteria all had positive estimates.

The strongest significant positive correlation was observed between the two Gram-positive bacteria:

S. aureus and *E. faecalis*. One explanation for this could be differences in immune defence across the

host phylogeny. In *D. melanogaster* Gram-positive bacteria are largely targeted by the Toll pathway

[30, 31, 76, 77]. Toll pathway members are highly conserved across arthropods and many Toll

pathway genes have large copy number variations across species, which may be linked to their

functional diversity [78, 79]. We also found a significant positive correlation across Gram types

between *E. faecalis* and *P. rettgeri*. Several studies have demonstrated there is cross-talk between the

Toll and IMD pathways, which may play a part in explaining the shared susceptibility [80-82].

However, it is also possible the positive correlations are driven by other factors, for example, the

suitability of the host physiological environment for bacterial proliferation.

We found the lowest proportion of variance explained by the phylogeny for *E. faecalis*, for which we

also found no correlation between bacterial loads and host mortality. Susceptibility to *E. faecalis* has

been shown to involve the action of Bomanin and Baramicin host defence peptides, whose mechanism

remains unclear, but could contribute to promoting tolerance to infection rather than pathogen

suppression [41, 42, 83]. A previous GWAS study in *D. melanogaster* found variation in susceptibility

to *E. faecalis* was due to genes not classically associated with immunity [43], suggesting factors

unrelated to bacterial suppression may be important in surviving infection by this pathogen. Further, a

study of immune priming showed that flies primed with a low dose of *E. faecalis* were better able to

survive subsequent infection by *E. faecalis*, but this was not related to bacterial clearance [84]. *E.*

faecalis virulence is known to involve the action of toxins, including Cytolysin and Enterocin V, and

induced immune responses including AMPs and iron sequestration have minimal impact on *E.*

faecalis virulence [54, 85, 86]. The disconnect between mortality and replication could certainly be

explained by toxin-based virulence if low doses of toxins are sufficient to cause mortality as in other

Gram-positive insect pathogens [87] and cytolysin, for example, is hemolytic in animal cells in the

nanomolar range. Differences in the host response to bacterial toxins such as presence of Toll pathway

effector Baramicin A may explain the phylogenetic patterns of susceptibility to *E. faecalis* [42].

We found evidence of microbiome growth in four species (*D. americana*, *S. lebanonensis*, *D. putrida*

and *S. pattersoni*) when measuring bacterial loads in homogenized pathogen infected insects.

Anecdotally, these species exhibited a relatively high survival to most of the bacteria (Figure 3). The

microbiome can play an important role in priming the immune response and defending against

bacterial infections [88, 89]. Gut probiotics such as *Lactobacillus* have been demonstrated to reduce

fly mortality caused by pathogenic bacteria [90]. It has been reported that there is limited variation in

the microbiome across species of Drosophilidae in laboratory culture [91, 92]. As such, a more

comprehensive study of how the microbiome varies across these host species, to what extent the

microbiome is related to host phylogeny, and whether this relates to differences in susceptibility is

warranted.

Although we have studied a relatively large number of host species, only four bacterial taxa were

used, limiting our ability to draw broad conclusions on differences between bacterial taxa (for

example differences between Gram type). Practical constraints resulted in fewer species being used to

estimate the bacterial load data, limiting our ability to assess phylogenetic signal for this trait, or

explore the role of tolerance in determining the patterns we observe among host speices. In addition,

further work is needed to understand the mechanisms underlying the differences in susceptibility

among species, and whether the same or different factors affect susceptibility in different host clades.

In conclusion, we have found that the host phylogeny can explain varying amounts of variation in

susceptibility depending on the pathogen. These varying phylogenetic effects (or lack thereof) have

important consequences for understanding susceptibility across host species and its impact on

pathogen host shifts [26, 61], which is helpful for our understanding of overcoming increasing

bacterial infection issues in public health. For some pathogens, certain clades of related hosts will be

particularly susceptible, meaning pathogens will be more likely to jump into those species, and they

could also act as disease reservoirs [93-96]. However, patterns may be less clear for other pathogens,

with relatedness not being a strong predictor of susceptibility across species. As such, caution should

be used when estimating the host range of novel pathogens based on susceptibility in closely related

host species, without suitable information on patterns of susceptibility across host species for the

bacterial taxa in question.

Acknowledgements

Bacterial strains were kindly provided by Brian Lazzaro and Ashley Frank. Thanks to Megan Wallace and Michael Jardine for their assistance in the code writing and experimental process for this study. Thanks to editor and two anonymous reviewers for the thoughtful and constructive comments.

Supplementary materials

Supplementary methods

Table S1 Drosophilidae host species, wing length and rearing diet information.

Table S2 Estimations of the differences between phylogenetic signal of each pathogen pair.

Table S3 *Diptericin A* sanger sequencing details (primer, information of *DptA* S69R site encoded amino acid, GenBank accession number).

Table S4 Statistical coefficients for models fitting or not fitting *Diptericin* alleles as fixed effects terms.

Table S5 Estimates of the mean proportion of variation [Abbot's corrected mortality, and the day 14 binomial mortality data] explained by the host phylogeny.

Table S6 Correlations in mortality [Abbot's corrected mortality, and the day 14 binomial mortality data] between different pathogens.

Figure S1 Correlations between bacterial load and mortality.

Figure S2 *Diptericin* allele variation across Drosophilidae species.

Figure S3 Correlation between mean mortality and Abbot's corrected mortality.

All data files and R scripts for analysis used in this study are available in an online repository:

https://doi.org/10.6084/m9.figshare.25648563

Author Contributions

Conceptualization: HS, BR, BL

Formal Experiments: HS

Data curation: HS, MAH

Formal analysis: HS, RMI, BR, BL

Investigation: HS , MAH, RMI, SKW

Methodology: HS, MAH, BR, BL

Supervision: BR, BL

Visualization: HS, RMI, SKW, BL

Writing- Manuscript draft: HS, MAH, BR, BL

Writing- Reviewing and editing: HS, MAH, BR, BL

Competing interests: We do not have any competing interests.

Funding:

This work has been supported by the following funding resources:

Hongbo Sun is supported by China Scholarship Council and University of Exeter PhD Scholarships

https://www.exeter.ac.uk/study/pg-research/csc-scholarships/.

Ben Longdon and Ryan Imrie are supported by a Sir Henry Dale Fellowship jointly funded by the

Wellcome Trust and the Royal Society (grant no. 109356/Z/15/Z) https://wellcome.ac.uk/funding/sir-

henry-dale-fellowships.

Ben Raymond is supported by NERC NE/V012053/1 and BBSRC BB/S002928/1.

Sarah K. Walsh is supported by a studentship funded by the Biotechnology and Biological Sciences

Research Council (BBSRC) South West Biosciences Doctoral Training Partnership (BB/M009122/1).

Mark A. Hanson is supported by Swiss National Science Foundation (SNSF) grant P500PB_211082

and a Wellcome Trust ECA Grant (grant no. 227559/Z/23/Z).

Reference

Bonneaud, C., Longdon, B. 2020 Emerging pathogen evolution: Using evolutionary theory to understand the fate of novel infectious pathogens. *EMBO Rep*. **21**, e51374. (10.15252/embr.202051374)

Petrosillo, N. 2019 Emerging infections and future threats. *Erciyes Medical Journal*. (10.14744/etd.2019.57805)

Watkins, K. 2018 Emerging Infectious Diseases: a Review. *Curr Emerg Hosp Med Rep*. **6**, 86-93. (10.1007/s40138-018-0162-9)

Murray, C. J. L., Ikuta, K. S., Sharara, F., Swetschinski, L., Robles Aguilar, G., Gray, A., Han, C., Bisignano, C., Rao, P., Wool, E.*, et al.* 2022 Global burden of bacterial antimicrobial resistance in 2019: a systematic analysis. *The Lancet*. **399**, 629-655. (https://doi.org/10.1016/S0140-6736(21)02724-0)

Richardson, E. J., Bacigalupe, R., Harrison, E. M., Weinert, L. A., Lycett, S., Vrieling, M., Robb, K., Hoskisson, P. A., Holden, M. T. G., Feil, E. J.*, et al.* 2018 Gene exchange drives the ecological success of a multi-host bacterial pathogen. *Nat Ecol Evol*. **2**, 1468-1478. (10.1038/s41559-018-0617-0)

Mohapatra, S., Menon, N. G. 2022 Factors responsible for the emergence of novel viruses: An emphasis on SARS-CoV-2. *Curr Opin Environ Sci Health*. **27**, 100358. (10.1016/j.coesh.2022.100358)

Delaney, N. F., Balenger, S., Bonneaud, C., Marx, C. J., Hill, G. E., Ferguson-Noel, N., Tsai, P., Rodrigo, A., Edwards, S. V. 2012 Ultrafast evolution and loss of CRISPRs following a host shift in a novel wildlife pathogen, Mycoplasma gallisepticum. *PLoS Genet*. **8**, e1002511. (10.1371/journal.pgen.1002511)

Murray, S., Pascoe, B., Méric, G., Mageiros, L., Yahara, K., Hitchings, M. D., Friedmann, Y., Wilkinson, T. S., Gormley, F. J., Mack, D.*, et al.* 2017 Recombination-Mediated Host Adaptation by Avian Staphylococcus aureus. *Genome Biol Evol*. **9**, 830-842. (10.1093/gbe/evx037)

9 Sanaei, E., Albery, G. F., Yeoh, Y. K., Lin, Y. P., Cook, L. G., Engelstadter, J. 2023 Host phylogeny

and ecological associations best explain Wolbachia host shifts in scale insects. *Mol Ecol*. **32**, 2351-

2363. (10.1111/mec.16883)

10 Lowy, F. D. 1998 Staphylococcus aureus infections. *N Engl J Med*. **339**, 520-532.

(10.1056/nejm199808203390806)

11 Cuny, C., Friedrich, A., Kozytska, S., Layer, F., Nübel, U., Ohlsen, K., Strommenger, B., Walther,

B., Wieler, L., Witte, W. 2010 Emergence of methicillin-resistant Staphylococcus aureus (MRSA) in

different animal species. *Int J Med Microbiol*. **300**, 109-117. (10.1016/j.ijmm.2009.11.002)

12 Price, L. B., Stegger, M., Hasman, H., Aziz, M., Larsen, J., Andersen, P. S., Pearson, T., Waters, A.

E., Foster, J. T., Schupp, J.*, et al.* 2012 Staphylococcus aureus CC398: host adaptation and emergence

of methicillin resistance in livestock. *mBio*. **3**, (10.1128/mBio.00305-11)

13 Howden, B. P., Giulieri, S. G., Wong Fok Lung, T., Baines, S. L., Sharkey, L. K., Lee, J. Y. H.,

Hachani, A., Monk, I. R., Stinear, T. P. 2023 Staphylococcus aureus host interactions and adaptation.

Nat Rev Microbiol. **21**, 380-395. (10.1038/s41579-023-00852-y)

14 Fischer, J. R., Stallknecht, D. E., Luttrell, P., Dhondt, A. A., Converse, K. A. 1997 Mycoplasmal

conjunctivitis in wild songbirds: the spread of a new contagious disease in a mobile host population.

Emerg Infect Dis. **3**, 69-72. (10.3201/eid0301.970110)

15 Bonneaud, C., Balenger, S. L., Zhang, J., Edwards, S. V., Hill, G. E. 2012 Innate immunity and the

evolution of resistance to an emerging infectious disease in a wild bird. *Mol Ecol*. **21**, 2628-2639.

(10.1111/j.1365-294X.2012.05551.x)

16 Fleming-Davies, A. E., Williams, P. D., Dhondt, A. A., Dobson, A. P., Hochachka, W. M., Leon, A.

E., Ley, D. H., Osnas, E. E., Hawley, D. M. 2018 Incomplete host immunity favors the evolution of

virulence in an emergent pathogen. *Science*. **359**, 1030-1033. (doi:10.1126/science.aao2140)

17 European Food Safety, A., Delbianco, A., Gibin, D., Pasinato, L., Boscia, D., Morelli, M. 2022

Update of the Xylella spp. host plant database - systematic literature search up to 31 December 2021.

EFSA J. **20**, e07356. (10.2903/j.efsa.2022.7356)

18 Donegan, M. A., Coletta-Filho, H. D., Almeida, R. P. P. 2023 Parallel host shifts in a bacterial plant

pathogen suggest independent genetic solutions. *Mol Plant Pathol*. **24**, 527-535.

(10.1111/mpp.13316)

19 Tinsley, M. C., Majerus, M. E. N. 2007 Small steps or giant leaps for male-killers? Phylogenetic

constraints to male-killer host shifts. *BMC Evolutionary Biology*. **7**, 238. (10.1186/1471-2148-7-238)

20 Weinert, L. A., Araujo-Jnr, E. V., Ahmed, M. Z., Welch, J. J. 2015 The incidence of bacterial

endosymbionts in terrestrial arthropods. *Proc Biol Sci*. **282**, 20150249. (10.1098/rspb.2015.0249)

21 Hilgenboecker, K., Hammerstein, P., Schlattmann, P., Telschow, A., Werren, J. H. 2008 How many

species are infected with Wolbachia?--A statistical analysis of current data. *FEMS Microbiol Lett*. **281**,

215-220. (10.1111/j.1574-6968.2008.01110.x)

22 Zug, R., Hammerstein, P. 2012 Still a host of hosts for Wolbachia: analysis of recent data suggests

that 40% of terrestrial arthropod species are infected. *PLoS One*. **7**, e38544.

(10.1371/journal.pone.0038544)

23 Russell, J. A., Goldman-Huertas, B., Moreau, C. S., Baldo, L., Stahlhut, J. K., Werren, J. H.,

Pierce, N. E. 2009 Specialization and geographic isolation among Wolbachia symbionts from ants and

lycaenid butterflies. *Evolution*. **63**, 624-640. (10.1111/j.1558-5646.2008.00579.x)

24 Sanaei, E., Charlat, S., Engelstädter, J. 2021 Wolbachia host shifts: routes, mechanisms, constraints

and evolutionary consequences. *Biol Rev Camb Philos Soc*. **96**, 433-453. (10.1111/brv.12663)

25 Longdon, B., Brockhurst, M. A., Russell, C. A., Welch, J. J., Jiggins, F. M. 2014 The evolution and

genetics of virus host shifts. *PLoS Pathog*. **10**, e1004395. (10.1371/journal.ppat.1004395)

26 Longdon, B., Hadfield, J. D., Webster, C. L., Obbard, D. J., Jiggins, F. M. 2011 Host phylogeny

determines viral persistence and replication in novel hosts. *PLoS Pathog*. **7**, e1002260.

(10.1371/journal.ppat.1002260)

27 Imrie, R. M., Roberts, K. E., Longdon, B. 2021 Between virus correlations in the outcome of

infection across host species: Evidence of virus by host species interactions. *Evol Lett*. **5**, 472-483.

(10.1002/evl3.247)

28 Huan, Y., Kong, Q., Mou, H., Yi, H. 2020 Antimicrobial Peptides: Classification, Design,

Application and Research Progress in Multiple Fields. *Front Microbiol*. **11**, 582779.

(10.3389/fmicb.2020.582779)

29 Hoffmann, J. A., Reichhart, J. M. 2002 Drosophila innate immunity: an evolutionary perspective.

Nat Immunol. **3**, 121-126. (10.1038/ni0202-121)

30 Lemaitre, B., Hoffmann, J. 2007 The host defense of Drosophila melanogaster. *Annu Rev Immunol*.

25, 697-743. (10.1146/annurev.immunol.25.022106.141615)

31 Buchon, N., Silverman, N., Cherry, S. 2014 Immunity in Drosophila melanogaster--from microbial

recognition to whole-organism physiology. *Nat Rev Immunol*. **14**, 796-810. (10.1038/nri3763)

32 Sackton, T. B., Lazzaro, B. P., Schlenke, T. A., Evans, J. D., Hultmark, D., Clark, A. G. 2007

Dynamic evolution of the innate immune system in Drosophila. *Nature Genetics*. **39**, 1461-1468.

(10.1038/ng.2007.60)

33 Obbard, D. J., Welch, J. J., Kim, K.-W., Jiggins, F. M. 2009 Quantifying Adaptive Evolution in the

Drosophila Immune System. *PLOS Genetics*. **5**, e1000698. (10.1371/journal.pgen.1000698)

34 Unckless, R. L., Howick, V. M., Lazzaro, B. P. 2016 Convergent Balancing Selection on an

Antimicrobial Peptide in Drosophila. *Curr Biol*. **26**, 257-262. (10.1016/j.cub.2015.11.063)

35 Hanson, M. A., Grollmus, L., Lemaitre, B. 2023 Ecology-relevant bacteria drive the evolution of

host antimicrobial peptides in Drosophila. *SCIENCE*. **381**, 284-+. (10.1126/science.adg5725)

36 Duneau, D., Ferdy, J. B., Revah, J., Kondolf, H., Ortiz, G. A., Lazzaro, B. P., Buchon, N. 2017

Stochastic variation in the initial phase of bacterial infection predicts the probability of survival in D.

melanogaster. *Elife*. **6**, (10.7554/eLife.28298)

37 Needham, A. J., Kibart, M., Crossley, H., Ingham, P. W., Foster, S. J. 2004 Drosophila

melanogaster as a model host for Staphylococcus aureus infection. *Microbiology*. **150**, 2347-2355.

(https://doi.org/10.1099/mic.0.27116-0)

38 Clewell, D. B., Tomich, P. K., Gawron-Burke, M. C., Franke, A. E., Yagi, Y., An, F. Y. 1982

Mapping of Streptococcus faecalis plasmids pAD1 and pAD2 and studies relating to transposition of

Tn917. *J Bacteriol*. **152**, 1220-1230. (10.1128/jb.152.3.1220-1230.1982)

39 Cox, C. R., Gilmore, M. S. 2007 Native microbial colonization of Drosophila melanogaster and its

use as a model of Enterococcus faecalis pathogenesis. *Infect Immun*. **75**, 1565-1576.

(10.1128/IAI.01496-06)

40 Gentry-Weeks, C. R., Karkhoff-Schweizer, R., Pikis, A., Estay, M., Keith, J. M. 1999 Survival of

Enterococcus faecalis in mouse peritoneal macrophages. *Infect Immun*. **67**, 2160-2165.

(10.1128/iai.67.5.2160-2165.1999)

41 Hanson, M. A., Cohen, L. B., Marra, A., Iatsenko, I., Wasserman, S. A., Lemaitre, B. 2021 The

Drosophila Baramicin polypeptide gene protects against fungal infection. *PLOS Pathogens*. **17**,

e1009846. (10.1371/journal.ppat.1009846)

42 Huang, J., Lou, Y., Liu, J., Bulet, P., Cai, C., Ma, K., Jiao, R., Hoffmann, J. A., Liégeois, S., Li, Z.,

Ferrandon, D. 2023 A Toll pathway effector protects Drosophila specifically from distinct toxins

secreted by a fungus or a bacterium. *Proceedings of the National Academy of Sciences*. **120**,

e2205140120. (doi:10.1073/pnas.2205140120)

43 Chapman, J. R., Dowell, M. A., Chan, R., Unckless, R. L. 2020 The Genetic Basis of Natural

Variation in Drosophila melanogaster Immune Defense against Enterococcus faecalis. *Genes*. **11**, 234.

44 Liu, G. Y., Essex, A., Buchanan, J. T., Datta, V., Hoffman, H. M., Bastian, J. F., Fierer, J., Nizet, V.

2005 Staphylococcus aureus golden pigment impairs neutrophil killing and promotes virulence

through its antioxidant activity. *J Exp Med*. **202**, 209-215. (10.1084/jem.20050846)

45 Matuszewska, M., Murray, G. G. R., Harrison, E. M., Holmes, M. A., Weinert, L. A. 2020 The

Evolutionary Genomics of Host Specificity in Staphylococcus aureus. *Trends in Microbiology*. **28**,

465-477. (https://doi.org/10.1016/j.tim.2019.12.007)

46 Binggeli, O., Neyen, C., Poidevin, M., Lemaitre, B. 2014 Prophenoloxidase Activation Is Required

for Survival to Microbial Infections in Drosophila. *PLOS Pathogens*. **10**, e1004067.

(10.1371/journal.ppat.1004067)

47 Dudzic, J. P., Hanson, M. A., Iatsenko, I., Kondo, S., Lemaitre, B. 2019 More than black or white:

melanization and toll share regulatory serine proteases in Drosophila. *Cell reports*. **27**, 1050-1061.

e1053.

48 Mulet, M., Gomila, M., Lemaitre, B., Lalucat, J., García-Valdés, E. 2012 Taxonomic

characterisation of Pseudomonas strain L48 and formal proposal of Pseudomonas entomophila sp.

nov. *Syst Appl Microbiol*. **35**, 145-149. (10.1016/j.syapm.2011.12.003)

49 Vodovar, N., Vinals, M., Liehl, P., Basset, A., Degrouard, J., Spellman, P., Boccard, F., Lemaitre, B.

2005 Drosophila host defense after oral infection by an entomopathogenic Pseudomonas species. *Proc*

Natl Acad Sci U S A. **102**, 11414-11419. (10.1073/pnas.0502240102)

50 Vodovar, N., Vallenet, D., Cruveiller, S., Rouy, Z., Barbe, V., Acosta, C., Cattolico, L., Jubin, C.,

Lajus, A., Segurens, B.*, et al.* 2006 Complete genome sequence of the entomopathogenic and

metabolically versatile soil bacterium Pseudomonas entomophila. *Nature Biotechnology*. **24**, 673-679.

(10.1038/nbt1212)

51 Dieppois, G., Opota, O., Lalucat, J., Lemaitre, B. 2015 Pseudomonas entomophila: A Versatile

Bacterium with Entomopathogenic Properties. In *Pseudomonas: Volume 7: New Aspects of*

Pseudomonas Biology. (ed.^eds. J.-L. Ramos, J. B. Goldberg, A. Filloux), pp. 25-49. Dordrecht:

Springer Netherlands.

52 Prakash, A., Monteith, K. M., Vale, P. F. 2022 Mechanisms of damage prevention, signalling and

repair impact disease tolerance. *Proc Biol Sci*. **289**, 20220837. (10.1098/rspb.2022.0837)

53 Juneja, P., Lazzaro, B. P. 2009 Providencia sneebia sp. nov. and Providencia burhodogranariea sp.

nov., isolated from wild Drosophila melanogaster. *Int J Syst Evol Microbiol*. **59**, 1108-1111.

(59/5/1108 [pii]10.1099/ijs.0.000117-0)

54 Hanson, M. A., Dostalova, A., Ceroni, C., Poidevin, M., Kondo, S., Lemaitre, B. 2019 Synergy and

remarkable specificity of antimicrobial peptides in vivo using a systematic knockout approach. *Elife*.

8, (10.7554/eLife.44341)

55 Unckless, R. L., Howick, V. M., Lazzaro, B. P. 2016 Convergent balancing selection on an

antimicrobial peptide in Drosophila. *Current Biology*. **26**, 257-262.

56 Short, S. M., Lazzaro, B. P. 2010 Female and male genetic contributions to post-mating immune

defence in female Drosophila melanogaster. *Proc Biol Sci*. **277**, 3649-3657.

(10.1098/rspb.2010.0937)

57 Schwenke, R. A., Lazzaro, B. P. 2017 Juvenile Hormone Suppresses Resistance to Infection in

Mated Female Drosophila melanogaster. *Curr Biol*. **27**, 596-601. (10.1016/j.cub.2017.01.004)

58 Duneau, D. F., Kondolf, H. C., Im, J. H., Ortiz, G. A., Chow, C., Fox, M. A., Eugenio, A. T.,

Revah, J., Buchon, N., Lazzaro, B. P. 2017 The Toll pathway underlies host sexual dimorphism in

resistance to both Gram-negative and Gram-positive bacteria in mated Drosophila. *BMC Biol*. **15**,

124. (10.1186/s12915-017-0466-3)

59 Gupta, V., Frank, A. M., Matolka, N., Lazzaro, B. P. 2022 Inherent constraints on a polyfunctional

tissue lead to a reproduction-immunity tradeoff. *BMC Biology*. **20**, 127. (10.1186/s12915-022-01328-

w)

60 Larson, E. L., Strom, M. S., Evans, C. A. 1980 Analysis of three variables in sampling solutions

used to assay bacteria of hands: type of solution, use of antiseptic neutralizers, and solution

temperature. *Journal of Clinical Microbiology*. **12**, 355-360. (doi:10.1128/jcm.12.3.355-360.1980)

61 Longdon, B., Hadfield, J. D., Day, J. P., Smith, S. C., McGonigle, J. E., Cogni, R., Cao, C., Jiggins,

F. M. 2015 The causes and consequences of changes in virulence following pathogen host shifts.

PLoS Pathog. **11**, e1004728. (10.1371/journal.ppat.1004728)

62 Jarrod, H. 2010 MCMC Methods for Multi-Response Generalized Linear Mixed Models: The

MCMCglmm R Package. *Journal of Statistical Software*. **33**,

63 Team, R. C. R: A Language and Environment for Statistical Computing. Vienna, Austria: R

Foundation for Statistical Computing 2023.

64 Rosenheim, J. A., Hoy, M. A. 1989 Confidence Intervals for the Abbott's Formula Correction of

Bioassay Data for Control Response. *Journal of Economic Entomology*. **82**, 331-335.

(10.1093/jee/82.2.331)

65 Abbott, W. S. 1925 A Method of Computing the Effectiveness of an Insecticide. *Journal of*

Economic Entomology. **18**, 265-267. (10.1093/jee/18.2.265a)

66 Tokuda, T., Goodrich, B., Mechelen, I. V., Gelman, A., Tuerlinckx, F. Year Visualizing

Distributions of Covariance Matrices *. 2011; 2011.

67 Freckleton, R. P., Harvey, P. H., Pagel, M. 2002 Phylogenetic analysis and comparative data: a test

and review of evidence. *Am Nat*. **160**, 712-726. (10.1086/343873)

68 Elizabeth A. Housworth, Emília P. Martins, Michael Lynch. 2004 The Phylogenetic Mixed Model.

The American Naturalist. **163**, 84-96. (10.1086/380570)

69 Pagel, M. 1999 Inferring the historical patterns of biological evolution. *Nature*. **401**, 877-884.

(10.1038/44766)

70 Falconer, D., Mackay, T. 1996 Introduction to quantitative genetics. Essex. *UK: Longman Group*.

12,

71 De Gregorio, E., Spellman, P. T., Tzou, P., Rubin, G. M., Lemaitre, B. 2002 The Toll and Imd

pathways are the major regulators of the immune response in Drosophila. *Embo j*. **21**, 2568-2579.

(10.1093/emboj/21.11.2568)

72 Troha, K., Im, J. H., Revah, J., Lazzaro, B. P., Buchon, N. 2018 Comparative transcriptomics

reveals CrebA as a novel regulator of infection tolerance in D. melanogaster. *PLOS Pathogens*. **14**,

e1006847. (10.1371/journal.ppat.1006847)

73 Gilchrist, G., Huey, R., Serra, L. 2001 Rapid evolution of wing size clines in Drosophila

subobscura. *Genetica*. **112-113**, 273-286. (10.1023/A:1013358931816)

74 Rainford, J. L., Hofreiter, M., Mayhew, P. J. 2016 Phylogenetic analyses suggest that

diversification and body size evolution are independent in insects. *BMC Evolutionary Biology*. **16**, 8.

(10.1186/s12862-015-0570-3)

75 McCarville, J. L., Ayres, J. S. 2018 Disease tolerance: concept and mechanisms. *Current Opinion*

in Immunology. **50**, 88-93. (https://doi.org/10.1016/j.coi.2017.12.003)

76 Rutschmann, S., Kilinc, A., Ferrandon, D. 2002 Cutting edge: the toll pathway is required for

resistance to gram-positive bacterial infections in Drosophila. *J Immunol*. **168**, 1542-1546.

(10.4049/jimmunol.168.4.1542)

77 Ambrosi, P., Chahda, J. S., Koslen, H. R., Chiel, H. J., Mizutani, C. M. 2014 Modeling of the

dorsal gradient across species reveals interaction between embryo morphology and Toll signaling

pathway during evolution. *PLoS Comput Biol*. **10**, e1003807. (10.1371/journal.pcbi.1003807)

78 Palmer, W. J., Jiggins, F. M. 2015 Comparative Genomics Reveals the Origins and Diversity of

Arthropod Immune Systems. *Mol Biol Evol*. **32**, 2111-2129. (10.1093/molbev/msv093)

79 Sackton, T. B., Lazzaro, B. P., Schlenke, T. A., Evans, J. D., Hultmark, D., Clark, A. G. 2007

Dynamic evolution of the innate immune system in Drosophila. *Nat Genet*. **39**, 1461-1468.

(10.1038/ng.2007.60)

80 Nishide, Y., Kageyama, D., Yokoi, K., Jouraku, A., Tanaka, H., Futahashi, R., Fukatsu, T. 2019

Functional crosstalk across IMD and Toll pathways: insight into the evolution of incomplete immune

cascades. *Proc Biol Sci*. **286**, 20182207. (10.1098/rspb.2018.2207)

81 Kounatidis, I., Ligoxygakis, P. 2012 Drosophila as a model system to unravel the layers of innate

immunity to infection. *Open Biol*. **2**, 120075. (10.1098/rsob.120075)

82 Valanne, S., Myllymäki, H., Kallio, J., Schmid, M. R., Kleino, A., Murumägi, A., Airaksinen, L.,

Kotipelto, T., Kaustio, M., Ulvila, J.*, et al.* 2010 Genome-wide RNA interference in Drosophila cells

identifies G protein-coupled receptor kinase 2 as a conserved regulator of NF-kappaB signaling. *J*

Immunol. **184**, 6188-6198. (10.4049/jimmunol.1000261)

83 Xu, R., Lou, Y., Tidu, A., Bulet, P., Heinekamp, T., Martin, F., Brakhage, A., Li, Z., Liégeois, S.,

Ferrandon, D. 2023 The Toll pathway mediates Drosophila resilience to Aspergillus mycotoxins

through specific Bomanins. *EMBO reports*. **24**, e56036. (https://doi.org/10.15252/embr.202256036)

84 Cabrera, K., Hoard, D. S., Gibson, O., Martinez, D. I., Wunderlich, Z. 2023 Drosophila immune

priming to Enterococcus faecalis relies on immune tolerance rather than resistance. *PLoS Pathog*. **19**,

e1011567. (10.1371/journal.ppat.1011567)

85 Iatsenko, I., Marra, A., Boquete, J.-P., Peña, J., Lemaitre, B. 2020 Iron sequestration by transferrin

1 mediates nutritional immunity in Drosophila melanogaster. *Proceedings of the National Academy of*

Sciences. **117**, 7317-7325. (doi:10.1073/pnas.1914830117)

86 Wadhawan, A., Silva, C. J. S. d., Nunes, C. D., Edwards, A. M., Dionne, M. S. 2022 E. faecalis

acquires resistance to antimicrobials and insect immunity via common mechanisms. *bioRxiv*.

2022.2008.2017.504265. (10.1101/2022.08.17.504265)

87 Cornforth, D. M., Matthews, A., Brown, S. P., Raymond, B. 2015 Bacterial cooperation causes

systematic errors in pathogen risk assessment due to the failure of the independent action hypothesis.

PLoS Pathogens. **11**, e1004775. (10.1371/journal.ppat.1004775.s002)

88 Panwar, R. B., Sequeira, R. P., Clarke, T. B. 2021 Microbiota-mediated protection against

antibiotic-resistant pathogens. *Genes & Immunity*. **22**, 255-267. (10.1038/s41435-021-00129-5)

89 Barron, A. J., Lesperance, D. N. A., Doucette, J., Calle, S., Broderick, N. A. 2023 Microbiome

derived acidity protects against microbial invasion in Drosophila. *bioRxiv*.

(10.1101/2023.01.12.523836)

90 Blum, J. E., Fischer, C. N., Miles, J., Handelsman, J. 2013 Frequent Replenishment Sustains the

Beneficial Microbiome of Drosophila melanogaster. *MBIO*. **4**, (10.1128/mBio.00860-13)

91 Chandler, J. A., Morgan Lang, J., Bhatnagar, S., Eisen, J. A., Kopp, A. 2011 Bacterial Communities

of Diverse Drosophila Species: Ecological Context of a Host–Microbe Model System. *PLOS*

Genetics. **7**, e1002272. (10.1371/journal.pgen.1002272)

92 Brooks, A. W., Kohl, K. D., Brucker, R. M., van Opstal, E. J., Bordenstein, S. R. 2016

Phylosymbiosis: Relationships and Functional Effects of Microbial Communities across Host

Evolutionary History. *PLOS Biology*. **14**, e2000225. (10.1371/journal.pbio.2000225)

93 Thines, M. 2019 An evolutionary framework for host shifts – jumping ships for survival. *New*

Phytologist. **224**, 605-617. (https://doi.org/10.1111/nph.16092)

94 van Baarlen, P., van Belkum, A., Summerbell, R. C., Crous, P. W., Thomma, B. P. H. J. 2007

Molecular mechanisms of pathogenicity: how do pathogenic microorganisms develop cross-kingdom

host jumps? *FEMS Microbiology Reviews*. **31**, 239-277. (10.1111/j.1574-6976.2007.00065.x)

95 Bonneaud, C., Weinert, L. A., Kuijper, B. 2019 Understanding the emergence of bacterial

pathogens in novel hosts. *Philos Trans R Soc Lond B Biol Sci*. **374**, 20180328.

(10.1098/rstb.2018.0328)

96 Bacigalupe, R., Tormo-Mas, M. Á., Penadés, J. R., Fitzgerald, J. R. 2019 A multihost bacterial

pathogen overcomes continuous population bottlenecks to adapt to new host species. *Science*

Advances. **5**, eaax0063. (doi:10.1126/sciadv.aax0063)

A *P. rettgeri*

B *P. entomophila*

C *E. faecalis*

D *S. aureus*

Mortality(%)

Mean load(CFU)