Varying phylogenetic signal to four bacterial pathogens across species of Drosophilidae

Hongbo Sun[1] *, Mark A. Hanson[1], Sarah K. Walsh[1,2,3], Ryan M. Imrie[1,3], Ben Raymond[1+], Ben

Longdon[1+]

[1] Centre for Ecology and Conservation; Faculty of Environment, Science, and Economy; Biosciences;

University of Exeter; Cornwall; United Kingdom,

[2] Environment and Sustainability Institute; University of Exeter; Cornwall; United Kingdom,

[3] MRC-University of Glasgow Centre for Virus Research, Glasgow; United Kingdom.

*　Corresponding author.

+ These authors contributed equally to this work.

Author information:

Hongbo Sun* hs766@exeter.ac.uk https://orcid.org/0009-0008-7272-7205

Mark A. Hanson m.hanson@exeter.ac.uk https://orcid.org/0000-0002-6125-3672

Sarah K. Walsh Sarah.Walsh.2@glasgow.ac.uk https://orcid.org/0000-0001-6598-095X

Ryan M. Imrie Ryan.Imrie@glasgow.ac.uk https://orcid.org/0000-0002-5055-2888

Ben Raymond b.raymond@exeter.ac.uk https://orcid.org/0000-0002-3730-0985

Ben Longdon b.longdon2@exeter.ac.uk https://orcid.org/0000-0001-6936-1697

Abstract

Pathogen host shifts – where a pathogen jumps from one host species to another – are important sources of emerging infectious diseases. Although viral pathogens are a major source of emerging infectious diseases, there is a growing focus on the emergence of novel bacterial infections through host shifts. However, compared to viruses we know relatively little about the factors that determine whether a bacteria can infect a novel host, such as how host phylogenetics constrains variation in pathogen host range. Here, we experimentally examined susceptibility to bacterial infections using a panel of 36 Drosophilidae species and four bacterial pathogens (*Providencia rettgeri*, *Pseudomonas entomophila*, *Enterococcus faecalis*, *Staphylococcus aureus*). The outcomes of infection differed greatly among pathogens and across host species. The host phylogeny explains a considerable amount of variation in susceptibility, with the greatest phylogenetic signal for *P. rettgeri* infection, with 94% of variation in mortality being explained by the host phylogeny. Positive correlations were observed between mortality and bacterial load for three out of the four pathogens. Correlations in susceptibility between the four pathogens were positive but largely non-significant, suggesting susceptibility is mostly pathogen-specific. These results suggest that susceptibility to bacterial pathogens may be predicted by the host phylogeny, but the effect may vary in magnitude between different bacteria.

Keywords: Bacterial infection, Susceptibility, Phylogeny, Host shift, Host-microbe interaction

Author Summary

Bacterial host shifts, where bacterial pathogens jump into and establish infection in a new host

species, are an important source of emerging infectious diseases. Here, we use four bacterial

pathogens to infect a panel of 36 species of fruit fly that shared a common ancestor ~50 million years

ago. We found large amounts of variation in susceptibility across host species to these bacterial

pathogens. This variation is explained by the host phylogeny to varying degrees, with hosts clustering

into clades with distinct levels of susceptibility. However, the magnitude of the effect varies between

the four bacterial pathogens. For three out of the four pathogens tested mortality was correlated with

bacterial load, suggesting it is a result of bacterial proliferation in the host. This has important

implications for our understanding of the emergence of bacterial pathogens, and to date there are few

large-scale experimental studies of bacterial host shifts.

Introduction

Emerging and re-emerging infectious diseases are a major threat to public health [1-3]. Recent studies

have highlighted that host shifts – where a pathogen jumps from one host species to another – are

important sources of emerging diseases, not only by viruses, but also by bacterial pathogens [4, 5].

Despite a major focus on viral pathogens, there is an increasing awareness that bacterial pathogens

also commonly undergo host shifts [6-8], which has been linked to their recent emergence as

causative agents of disease in several host systems, including humans, livestock, wildlife, and plants

[6-8].

Most research on bacterial host shifts has considered pathogens of humans or wildlife. In

Staphylococcus aureus, a human pathogen associated with several severe hospital and community-

acquired infections [9], humans are the primary transmission hub for host shifts into a range of

livestock species, and cows have been considered the main animal reservoirs for human epidemics

[10-12]. Another extensively studied example is the jump of *Mycoplasma gallisepticum* from poultry

to wild house finches [13]. Here, the *Mycoplasma* host shift has driven the evolution of resistance in

the house finch hosts [14], led to changes in bacterial gene content [6] and the evolution of virulence

[15]. In plants the bacterial pathogen *Xylella fastidiosa* is found in over 600 host species [16]. A single

gene loss event is associated with the host shift of *X. fastidiosa* from coffee plants to olive trees [17].

Most studies on bacterial host shifts are based on genomic data following natural host shifts, and there

are limited experimental studies of bacterial host shifts that move beyond two-host systems.

Perhaps the best experimental studies relating to bacterial host shifts come from insects and their

bacterial endosymbionts. In a study of coccinellid beetles and a male-killing isolate of *Spiroplasma*,

the bacterium's ability to distort offspring sex ratios in novel hosts was negatively correlated with the

genetic distance from the bacteria's natural host [18]. Similar patterns have been found for *Wolbachia*,

a bacterial endosymbiont that infects ~40-60% of all arthropods [19-21]. Artificial transfers of

Wolbachia are most successful between closely related hosts [22, 23]. This tallies with field data on

Wolbachia distributions that found strong effects of the host phylogeny, with high rates of *Wolbachia*

sharing between closely related species, and a decline in co-occurrence in different hosts with

increasing phylogenetic distance between these hosts [8]. Viruses show similar trends, with

phylogenetic distance and clade effects explaining large amounts of variation in susceptibility to

several viruses [24, 25]. When comparing different viruses from the same family, positive inter-

specific correlations in susceptibility have been observed, suggesting similar mechanisms underlying

among-species differences in susceptibility [25, 26]. However, it is unclear if similar phylogenetic

patterns could exist for free-living pathogenic bacteria (i.e. non-endosymbionts), which are less tightly

linked to their hosts.

Here, we have used a large cross-infection experiment to understand how susceptibility to bacterial

pathogens varies across host species. We infected a large panel of 36 species of Drosophilidae with

four bacterial pathogens: two Gram-positives, *Staphylococcus aureus*, *Enterococcus faecalis*, and two

Gram-negatives: *Providencia rettgeri* and *Pseudomonas entomophila*. In *Drosophila melanogaster*,

the major immune response to bacterial pathogens involves the Toll and IMD pathways. The Toll

pathway primarily controls resistance to fungal and Gram-positive bacterial infections, whereas the

Imd pathway mediates response to Gram-negative bacterial infections [27, 28]. Both pathways result

in the initiation of the *NF-κB* signalling pathway and the upregulation of antimicrobial peptides

(AMPs) and other defence response genes [29, 30]. Genes in these pathways evolve rapidly; however

how differences among species contribute to functional outcomes remains unclear [31, 32]. Recent

studies have demonstrated that AMPs show specificity to certain bacterial infections and that different

alleles of AMPs are linked to differences in susceptibility both within and across species [33, 34].

The bacterial pathogens used here have been well characterised in *D. melanogaster* [35, 36]. *E.*

faecalis was initially derived from a human clinical isolate [37, 38]. *E. faecalis* produces toxins (e.g.

cytolysin), that act as virulence factors to improve bacterial colonisation and decrease host survival, in

Drosophila [38], the Toll pathway, particularly the AMPs Bomanin and Baramicin have been shown

to play a role is resistance to *E. faecalis* [39-41] . *S. aureus* was initially isolated from humans and

has been used in mice and *Drosophila* infection models [35, 42]. In *D. melanogaster*, resistance to *S.*

aureus specifically relies on the melanisation response [43, 44]. *P. entomophila* was isolated from a

wild female *D. melanogaster* – upon infection flies show differential expression of genes including

AMPs, oxidative stress response genes and detoxification genes [45, 46]. *P. entomophila* causes

digestive obstruction and eventually leads to the host's death by damage and perforation of the gut

through the production of insecticidal protease-based toxins [46, 47]. *P. rettgeri* was initially isolated

from the haemolymph of wild *D. melanogaster* [48]. The Toll pathway response, and particularly the

AMP Diptericin have been shown to be an important part of the immune response to *P. rettgeri* [33,

34, 49]. A single nucleotide polymorphism at residue 69 in *Diptericin A*, "S69R" of *Dpt* A, is

associated with increased susceptibility to *P. rettgeri* bacterial infection [34, 50].

Here, we examine the role of the host phylogeny in determining the outcomes of bacterial infections.

We investigated how variation in bacterial virulence varies across the host phylogeny and the

relationship between virulence and bacterial load for each of the four bacterial pathogens.

Methodology

Bacterial strains and culture

We used four bacterial pathogens in this study. Two Gram-positive bacteria, *Staphylococcus aureus* PIG1 (*S. aureus*) [42] and *Enterococcus faecalis* FA2-2/pAM714 (*E. faeclis*) [37, 38], and two Gram-negative bacteria: *Providencia rettgeri* Dmel (*P. rettgeri*) [48] and *Pseudomonas entomophila* (*P. entomophila*) [46] . Bacterial cultures were initiated from frozen glycerol stocks onto LB (Lysogeny Broth) agar, with the exception of *P. entomophila,* which were plated on egg yolk agar (10g of agar, 10g of LB, 1 L of water and 20 mL of egg yolk (Sigma)). Colonies of *P. entomophila* with halos on egg yolk agar (indicating phospholipase activity) were used to prepare inocula. Insect inocula were cultured in LB broth, with the exception of *E. faecalis* which used BHI (Brain heart infusion) broth. All bacteria were grown at a 30℃ incubator, shaking 180 rpm/min for liquid culture. Inocula were prepared in a three-step process in order to standardise physiology in the stationary phase. Initially, one or two colonies were used to prepare overnight cultures of 5 mL in 30 mL sterile universals, here inoculation success with *E. faecalis* was improved by using two colonies. These were diluted in liquid media to OD_{600} 0.2, and then 50 μL of this was inoculated into new vials. Following overnight growth, stationary phase cultures were diluted with culture media to the desired OD value (OD 2.0 for *E. faecalis* and OD 1.0 for the other bacteria).

Fly stocks and rearing.

All flies were maintained in multi-generation stock bottles (Fisher brand) at 22°C, 70% relative humidity in a 12-hour light-dark cycle. Each stock bottle contained 50 ml of one of four varieties of *Drosophila* media which were chosen to optimise rearing conditions (full details in Supplementary Table 1). Male flies were used to avoid any effect of sex or mating status, which has been shown to influence the susceptibility of female flies to various pathogens [51-53]. Before inoculation, 0-1 day old male flies were transferred to vials containing cornmeal media. These flies were then transferred to fresh media every three or four days for a week (age 7–8 days, flip once before infection), at which point they were inoculated. Flies were kept at 22°C, 70% relative humidity in a 12-hour light-dark cycle throughout the experiments. Flies were inoculated under CO_2 anaesthesia via septic pin prick

with 12.5µm diameter stainless steel needles (Fine Science Tools, CA, USA). These needles were bent

approximately 250µm from the end to provide a depth stop and dipped in the inoculum before being

pricked into the anepisternal cleft in the thorax of anaesthetised flies. Inoculation by this method

bypasses the gut immune barrier but avoids differences in inoculation dose due to variations in

feeding rate in oral inoculation.

For the survival assay, we inoculated male flies of 36 fly species. We inoculated three replicate vials

for 165 species-bacteria combinations, two replicate vials for 13 combinations and one replicate vial

for two combinations, with each combination one of 36 species infected with either a control Ringer's

solution [54] or one of the four bacterial pathogens listed above. Vials contained a mean of 14 flies

(range = 5-18 flies). Following injection, the number of dead flies in each vial was recorded daily for

14 days. Flies were tipped into new vials of food every two days.

We measured bacterial load in 18 species of fly; these species were selected based on their survival

results and covered a broad range of survival values for each pathogen. We measured colony forming

units (CFUs) for individual male flies of each species (mean = 15 flies, range 1-16) infected with each

species-bacteria combination. CFUs were measured at 16 hours post-infection; data points for dead

flies were excluded to avoid a bifurcation in bacterial load. Flies were prepared and infected as above.

Before homogenisation, anaesthetised flies were moved to 96-well tubes containing a 4mm steel ball

and 200 µL 70% ethanol, flies were then gently cleaned using 70% ethanol and homogenised in 200

µL sterile LB or sterile BHI (for flies infected with *E. faecalis*) in a Tissue Lyser II at 30 Hz for three

minutes. Homogenates were diluted 10^{-1} to 10^{-5} and plated on LB agar or BHI agar in two technical

replicates before being cultured overnight at 30°C. Bacterial loads were measured as colony-forming

units (CFUs) per fly, following the formula:

CFU per fly = (Colony counts × Dilution factor × Homogenate volume)/ Plating volume.

Host Phylogeny

The method used to infer the host phylogeny has been described elsewhere [55]. Briefly, publicly

available sequences of the *28S*, *Adh*, *Amyrel*, *COI*, *COII*, *RpL32*, and *SOD* genes were collected from

GenBank (see www.doi.org/10.6084/m9.figshare.6653192 for a full breakdown of genes and

accessions by species). Gene sequences were aligned in Geneious 9.1.8 (https://www.geneious.com)

using a progressive pairwise global alignment algorithm with free end gaps and a 70% similarity IUB

cost matrix. Gap open penalties, gap extension penalties, and refinement iterations were kept as

default.

Phylogenetic reconstruction was performed using BEAST version 1.10.4 [56] as the subsequent

phylogenetic mixed model (described below) requires a tree with the exact root-tip distances for all

taxa. Genes were partitioned into separate ribosomal (*28S*), mitochondrial (*COI*, *COII*), and nuclear

(*Adh*, *Amyrel*, *RpL32*, *SOD*) groups. The mitochondrial and nuclear groups were further partitioned

into groups for codon position 1+2 and codon position 3, with unlinked substitution rates and base

frequencies across codon positions. Each group was fitted to separate relaxed uncorrelated lognormal

molecular clock models using random starting trees and four-category gamma-distributed HKY

substitution models. The BEAST analysis was run twice, with 1 billion Markov Chain Monte Carlo

(MCMC) generations sampled every 100,000 iterations, using a birth-death process tree-shape prior.

Model trace files were evaluated for chain convergence, sampling, and autocorrelation using Tracer

version 1.7.1 [57]. A maximum clade credibility tree was inferred from the posterior sample with a

10% burn-in. The reconstructed tree was visualised using ggtree version 2.0.4 [58].

Sequencing of *Diptericin A*

To determine the *Diptericin A* allele of each species (which has been linked to susceptibility to

bacterial pathogens [34]), Diptericin genes were detected across species using a recursive tBLASTn

approach with a high E-value threshold to compensate for the very short query length (E < 1),

followed by manual curation. Using this method, we could detect Diptericin sequence from all species

included in this study. Then we designed degenerate primers of *Diptericin A* for each of species except

for seven species for which no *Diptericin A* sequence data was available (full primer details in

Supplementary Table 2). Fly were homogenised in 48 μL extraction buffer (0.5 μL Tris-Cl pH 8.2

(1M), 0.1 µL EDTA (0.5M), 0.25 µL NaCl (5M) and 47.15 µL ddH₂O) and 2µL proteinase

K(5mg/µL) then incubated at 37 °C for 30 minutes, followed by 95 °C for 5 minutes. PCRs were then

carried out under the following conditions: 95°C for 30secs, followed by 35 cycles of: 95°C for 20

secs, 54°C for 20 secs, 68°C for 1minute, followed by a final 5 minutes at 68°C. PCR products were

purified using Monarch® PCR&DNA Cleanup Kit, and Sanger sequenced through Eurofins Genomic

tube sequencing service, *Diptericin A* S69R site allele analysis were carried out by Geneious prime

v10.2.6. All sequences have been uploaded to GenBank (full accession number detailed in

Supplementary Table 2).

Statistical analysis

We used phylogenetic mixed models to look at the effects of host species relatedness on bacterial

susceptibility across the 36 host species. We fitted all models using a Bayesian approach in the R

package MCMCglmm v2.29 [59] in RStudio (R v4.2.2) [60]. We used a multivariate model with

mortality following inoculation with each of the four bacteria or control treatments as the response

variables. Mortality was calculated as each vial's mean proportion of dead flies over 14 days. The

models took the following form:

(1) $y_{hit} = \beta_{1:t} + u_{p:ht} + u_{s:ht} + e_{hit}$

(2) $y_{hit} = \beta_{1:t} + u_{p:ht} + e_{hit}$

Where y is the mortality of the i^{th} biological replicate of host species h, for treatment t (each of the

four bacteria and the non-infected control). β are the fixed effects, with β_1 being the intercepts for

each trait.

We also ran bivariate models for each of the four pathogens across the 18 host species tested to

investigate how mortality was related to bacterial load. These models had the following form:

(3) $y_{hit} = \beta_{1:t} + u_{p:ht} + e_{hit}$

Where y is the response of the i^{th} biological replicate of host species h, for treatment t (either

mortality or the bacterial load (CFU) of an individual fly infected with that pathogen). U are random

effects for the phylogenetic species effects (p), non-phylogenetic species effects (s) and residuals (e).

$u_{s:ht}$ was removed from models (2) and (3) as the models struggled to separate the phylogenetic and

non-phylogenetic effects. The random effects and the residuals are assumed to be multivariate normal

with a zero mean and a covariance structure $V_p \otimes \mathbf{A}$ for the phylogenetic effects and $V_e \otimes \mathbf{I}$ for the

residuals ($\otimes$ here is the Kronecker product). $\mathbf{A}$ is the phylogenetic relatedness matrix, $\mathbf{I}$ is an identity

matrix and the $\mathbf{V}$ are 5×5 (co)variance matrices describing the (co)variances between mortality when

infected with each of the four bacteria or the control. In model (3) the $\mathbf{V}$ are 2×2 (co)variance matrices

describing the (co)variances between mortality and bacterial load when infected with each of the four

bacteria. Specifically, the matrices V_p and V_s describe the phylogenetic and nonphylogenetic between-

species variances in mortality for each treatment and the covariances between them, whereas the

residual covariance matrix V_e describes within-species variance that includes both actual within-

species effects and measurement error. The off-diagonal elements in V_e (the covariances) are unable to

be estimated because no vial has multiple measurements – so they were set to zero.

Model (1) was also run with two additional fixed effects. Firstly, we included wing size, measured as

the length of the IV longitudinal vein from the tip of the proximal segment to the join of the distal

segment with vein [61], which provided a proxy for body size [62] and was included in a further

model as *wingsizeβ$_{2:t}$*. This was done to ensure that any phylogenetic signal in body size did not

explain the differences in susceptibility between species [63]. Secondly, we ran an additional model

which included the effect of polymorphisms at the *Diptericin A* S69R site ($DptA\beta_{3:t}$) which has

previously been implicated in resistance to bacterial pathogens [34, 49]. As variance explained by the

phylogeny is after accounting for any fixed effects [64], we examined if variance explained by the

phylogeny changed with the inclusion of the *Diptericin* allele (we would expect the phylogenetic

variance to decrease if the fixed effects have an effect on mortality and are clustered on the

phylogeny).

The MCMC chain was run for 13 million iterations with a burn-in of 3 million iterations and a

thinning interval of 5000 iterations. We confirmed our experimental design and sample sizes had

sufficient power to detect effects by down sampling a similar dataset [26]. The results presented were

obtained using parameter expanded priors for the $\mathbf{V}_p$ and $\mathbf{V}_s$ matrices [65]. Results were tested for

sensitivity to the use of different priors by being run with different prior structures. The alternative

priors gave qualitatively results, except for the correlations where we found mean estimates were

higher with the majority being significant. To determine what may be driving this we simulated

response data based on mean estimates from the parameter expanded prior model using the

'simulate.MCMCglmm' function. We then refitted models on this data with either parameter

expanded or alternative priors (inverse wishart or flat). The alternative priors had higher estimates of

the correlations in the refitted models suggesting they may be inadvertently influencing the estimates.

The proportion of the between-species variance that the phylogeny can explain was calculated from

model (1) using the equation $V_p/(V_p + V_s)$, where V_p and V_s represent the phylogenetic and species-

specific components of between-species variance [66], respectively, and are equivalent to

phylogenetic heritability or Pagel's lambda [67, 68]. The repeatability of bacterial load measurements

was calculated from model (2) as $V_p/(V_p + V_e)$, where V_e is the residual variance of the model [69].

Interspecific correlations were calculated from model (2) and (3) V_p matrix as $cov_{x,y}/\sqrt{(var_x \times var_y)}$.

Parameter estimates reported are means of the posterior density, and 95% credible intervals (CIs) were

taken to be the 95% highest posterior density intervals.

Results

Patterns of susceptibility to bacterial pathogens across host species

To investigate how susceptibility to bacterial pathogens varies across host species, we measured survival in 7402 flies from 36 species of Drosophilidae inoculated with one of four bacterial pathogens or a control. We found large differences in the mean virulence of the bacteria; the mean mortality was 70% for *P. entomophila* infection, 46% for *P. rettgeri*, 39% for *S. aureus*, and 36% for *E. faecalis* (compared to only 4% in control inoculated flies). There were dramatic differences in the survival of the different species with the majority of individuals of certain species dying, whilst others showed little mortality when infected with the same pathogen (Figure 1). For example, just over half of species showed rapid mortality following *P. entomophila* infection, with over 50% of flies dying within one day post-infection in 19 out of 36 fly species (Figure 1). Conversely, survival remained high for most species infected with *E. faecalis,* with the exceptions of *D. yakuba* and *D. sucinea* where >85% of flies had died by the third day post-infection (Figure 1).

Figure 1. Survival of 36 species of Drosophilidae after infection with four bacterial pathogens.

Each point represents the mean survival of three replicates, colours represent the inoculation

treatment.

296 **The host phylogeny explains varying amounts of variation in mortality**

297 Different clades of hosts show different mortality levels with each bacteria, with hosts with high and

298 low mortality clustering together on the host phylogeny (Figure 2). For example, a clade within the

299 subgenus Drosophila (delineated by *D. micromelanica* down to *D. flavomontana* in Figure 2) shows

300 low mortality when infected with *P. rettgeri* compared to the other major host clades. We used a

301 phylogenetic mixed model to determine the proportion of variation in mortality explained by the host

302 phylogeny for each pathogen (Table 1). The estimates vary among the four pathogens, with the host

303 phylogeny explaining large amounts of variation for three out of four bacterial infections. We find that

304 almost all of the variation in mortality is explained by the phylogeny following *P. rettgeri* infections

305 (94%). The phylogeny explains a substantial proportion of the variation in mortality for *S. aureus*

306 (64%) and *P. entomophila* (36%) but less for *E. faecalis* (14%).

Pathogen	Variance explained by phylogeny	Repeatability
P. rettgeri	0.94 (95% CI: 0.81, 1.00)	0.84 (95% CI: 0.75,0.93)
P. entomophila	0.36 (95% CI: 0.00, 0.74)	0.70 (95% CI: 0.48,0.89)
E. faecalis	0.14 (95% CI: 0.00, 0.50)	0.90 (95% CI: 0.80,0.97)
S. aureus	0.64 (95% CI: 0.24, 0.94)	0.80 (95% CI: 0.63,0.92)

307 **Table 1. Estimates of the mean proportion of variation explained by the host phylogeny (taken**

308 **from model (1)), and the repeatability of estimates (taken from model (2)).**

309

Figure 2: Susceptibility to four bacterial pathogens varies across 36 Drosophilidae species. Bar length and colour represent mean mortality change across pathogen and fly species with error bars representing standard errors. The phylogeny of Drosophilidae host species is presented on the left, with the scale bar representing the number of nucleotide substitutions per site. Names of host species are given on the right; the column label shows different treatments (bacterial or control inoculation).

Patterns of susceptibility are largely distinct for each bacteria

In *D. melanogaster*, the host immune response differs markedly between infections by Gram-positive

and Gram-negative bacteria [70, 71]. If mechanisms of host defence depend on the induced immune

response, we might expect positive correlations in host susceptibility among species for bacteria of the

same Gram type. We found the correlations between bacteria all had positive estimates (Table 2),

although the 95% CIs overlapped zero for all except for the correlations between *S. aureus–E. faecalis*

and *E. faecalis–P. rettgeri.* The correlation coefficient was less than 0.4 for all correlations bar the

estimate between the two Gram-positive bacteria *S. aureus* and *E. faecalis* where $r = 0.66$. Overall,

this suggests that host susceptibility generally differs by pathogen, with shifts in the rank order of

species susceptibilities to the different pathogens (i.e. species do not show overall high or low

susceptibity to bacteria *per se*, but rather pathogen specific susceptibilities).

Bacteria	Inter-specific correlation r
S. aureus - E. faecalis	**0.66 (95% CI: 0.37, 0.90)**
S. aureus - P. rettgeri	0.35 (95% CI: -0.04, 0.70)
S. aureus - P. entomophila	0.34 (95% CI: -0.13, 0.73)
E. faecalis - P. rettgeri	**0.39 (95% CI: 0.02, 0.72)**
E. faecalis - P. entomophila	0.31 (95% CI: -0.19, 0.74)
P. rettgeri - P. entomophila	0.37 (95% CI: -0.05, 0.75)

Table 2. Correlations in mortality between different pathogens. Numbers represent the mean

estimates of correlation coefficients from Model (2) with 95% credible intervals shown in brackets.

Significant correlation coefficients (with 95% CI that do not span zero) are highlighted in bold.

Mortality correlates with bacterial load in most cases

To test whether increased mortality is due to higher bacterial loads, we selected 18 fly species that varied in their mortality and measured their bacterial load to each pathogen at 16 hours post-infection. We found that 24% of the flies infected by *P. entomophila* had died by 16 hours post-infection (<2% of flies infected with each of the other bacteria died by this time point). We excluded dead flies from the analysis presented below but note that the results were robust if we included the bacterial load measured in corpses. We quantified bacterial load in 1328 individual flies (Figure 3) and ran models combining this with the mortality data above for these 18 fly species.

Mortality estimates from this model using a subset of the data from the survival experiment (Table 3) are consistent with observations in survival experiments (Figures 1 and 2). Mean bacterial loads are relatively consistent across the four pathogens (Table 3). We found strong positive correlations between mortality and bacterial load for *P. rettgeri* and *P. entomophila* and a slightly lower positive correlation for *S. aureus* with 95% CI's that just overlap zero (Table 4, Figure S1). However, we found no evidence of a significant positive correlation between mortality and bacterial load for *E. faecalis* (Table 4, Figure S1).

We found some evidence of microbiome proliferation in pathogen infected insects when measuring CFU in four species (*D. americana*, *S. lebanonensis*, *D. putridia* and *S. pattersoni*). We confirmed that excluding these species from our analysis did not alter the results qualitatively.

Pathogen	Bacterial load	Mortality
P. rettgeri	4.63 (95% CI: 3.67, 5.50)	0.46 (95% CI: 0.17, 0.76)
P. entomophila	3.78 (95% CI: 2.97, 4.46)	0.62 (95% CI: 0.33, 0.90)
E. faecalis	3.36 (95% CI: 2.80, 3.92)	0.37 (95% CI: -0.02, 0.74)
S. aureus	3.12 (95% CI: 2.32, 3.85)	0.36 (95% CI: 0.06, 0.68)

Table 3. Estimations of mortality and bacterial load for each pathogen. Numbers represent the mean estimates of mortality (mean proportion dead) and bacterial load (CFU per fly on a Log10 scale) for each pathogen from Model (3) with 95% credible intervals shown in brackets.

Figure 3: Mortality and bacterial load to four bacterial pathogens across 18 fly species. Bar

length and colour represent mean mortality (left) or bacterial load (right) across 18 fly species bearing

either *P. rettgeri* (A), *P. entomophila* (B), *E. faecalis* (C) or *S. aureus* (D) infection. Bacterial loads

are data point at 18 hours post infection, with the error bars representing standard errors of a mean of

15 replicate flies for each species. The phylogeny of Drosophilidae host species is presented on the

left, with the scale bar representing the number of nucleotide substitutions per site. Names of host

species are presented on the right.

Pathogen	Inter-specific correlation r
P. rettgeri	0.59 (95% CI's: 0.16, 0.94)
P. entomophila	0.54 (95% CI's: 0.10, 0.90)
E. faecalis	0.22 (95% CI's: -0.43, 0.74)
S. aureus	0.48 (95% CI's: -0.03, 0.93)

361 **Table 4. Correlations between mortality and bacterial load for each pathogen.** Numbers represent

362 the mean estimates of correlation coefficients from Model (3) with 95% credible intervals shown in

363 brackets. Significant correlation coefficients are where the 95% CI do not span zero.

Testing for effect of *Diptericin A* allelic variation and body size

We observed an exceptionally high proportion of variance explained by host phylogeny for *P. rettgeri* (Table 1). This suggests some phylogenetically inherited factor, or factors, strongly determines susceptibility to this pathogen. Previous studies linked the gene *Diptericin A* (*Dpt A*) to defence against *P. rettgeri*, including resistant or susceptible alleles at a polymorphic residue both within and across species [34, 49].

To test if this was the case here, we attempted Sanger sequencing of the *Dpt A* gene across our species panel. We successfully amplified and sequenced *Dpt A* from 29 species and mapped the allele at a previously identified polymorphic residue (residue 69 of *D. melanogaster Dpt A*). Four residues were found in our panel at this site (Figure S2). These were dominated by the resistant allele encoding serine (S) in 75% of species, and three alternate alleles each found in only 2-3 species, including a known susceptible allele encoding arginine (R), and alleles encoding either glutamine (Q) or asparagine (N), preliminarily associated with resistance and susceptibility respectively [34]. Given the limited replication of each individual allele, we fitted a fixed effect of *Dpt A* residue 69 by pathogen with species either being resistant (S and Q) or susceptible (R and N). We found no significant effect of the susceptible *Dpt A* alleles for any of the four pathogens (Table S3) but note there was limited power to detect an effect with only five species having susceptible alleles. There was little change in the variance explained by the phylogeny when the Diptericin allele status was included as a fixed effect (Table S3) indicating the alleles are largely not explaining the phylogenetic patterns observed.

Likewise, we found no effect of wing size on susceptibility for any pathogen (mean and 95% CIs for each pathogen: *P. rettgeri*: -0.34 (95% CI:-0.75, 0.02), *P. entomophila*: -0.02 (95% CI:-0.54, 0.45), *E. faecalis*: 0.01 (95% CI:-0.37, 0.37), *S. aureus*: 0.03 (-0.37, 0.42)).

Discussion

We examined how susceptibility to bacterial pathogens varies across the host phylogeny, using a panel of 36 Drosophilidae species and four bacterial pathogens. Using phylogenetic mixed models we found the host phylogeny can explain a large proportion of variation in susceptibility, but differs in the extent of variation explained between pathogens. Distinct infection patterns were seen between the different pathogens, with all correlations in susceptibility between the pathogens being less than one, suggesting susceptibility generally varies between bacterial pathogens. However, despite being largely insignificant, all correlations had positive estimates, and the strongest correlation was seen between the two Gram-positive bacteria (Table 2) suggesting that differences in susceptibility may have shared underlying mechanisms. Positive correlations between mortality and bacterial load for *P. rettgeri*, *P. entomophila* and *S. aureus* suggest virulence is a direct results of bacterial accumulation in the host. The lack of correlation between virulence and bacterial load for *E. faecalis* may have a biological underpinning, such as the production of toxins, or may be due to limited statistical power.

Our survival results find a substantial amount of variation in susceptibility can be explained by the host phylogeny, with the largest effect for *P. rettgeri,* where almost all of the variation in susceptibility is explained by the host phylogeny. One potential explanation is that susceptibility is linked to variation in immune genes with certain host clades having lost or gained different aspects of the immune system. Previous studies have shown that a specific AMP, Diptericin, plays an essential role in defence against *P. rettgeri* infection. Flies encoding arginine (R) at the *Diptericin* S69R polymorphism are more susceptible than flies encoding serine (S) at this site [34, 49]. Due to the limited replication of the different Diptericin alleles (75% of the 29 species with allele data had the "S" resistance allele, with each of the other 3 alleles only replicated in 2 or 3 species) we had little power to detect effects here. However, there is also substantial variation in susceptibility just among species with the "S" allele, suggesting this Diptericin allele is not the sole driver of the patterns observed here. Further study across a broader range of species that vary in their Diptericin alleles are required, perhaps combined with measurements of the speed of induction of Diptericin and other AMPs, given rapid induction has been demonstrated to be important for host survival [34, 35, 49].

We found the among species correlations in susceptibility between bacteria all had positive estimates.

The strongest significant positive correlation was observed between the two Gram-positive bacteria:

S. aureus and *E. faecalis*. One explanation for this could be differences in immune defence across the

host phylogeny. In *D. melanogaster* Gram-positive bacteria are largely targeted by the Toll pathway

[29, 30, 72, 73]. Toll pathway members are highly conserved across arthropods and many Toll

pathway genes have large copy number variations across species, which may be linked to their

functional diversity [74, 75]. We also found a significant positive correlation across Gram types

between *E. faecalis* and *P. rettgeri*. Several studies have demonstrated there is cross talk between the

Toll and IMD pathways which may play a part in explaining the shared susceptibility [76-78].

However, it is also possible the positive correlations are driven by other factors, for example,

suitability of the host physiological environment for bacterial proliferation.

We found the lowest proportion of variance explained by the phylogeny for *E. faecalis*, for which we

also found no correlation between bacterial loads and host mortality. Susceptibility to *E. faecalis* has

been shown involve the action of Bomanin and Baramicin antimicrobial peptides, whose mechanism

remains unclear, but could contribute to promoting tolerance to infection rather than pathogen

suppression [39-41]. A previous GWAS study in *D. melanogaster* found variation in susceptibility to

E. faecalis was due to genes not classically associated with immunity [79], suggesting factors

unrelated to bacterial suppression may be important in surviving infection by this pathogen. Further, a

study of immune priming showed that flies primed with a low dose of *E. faecalis* were better able to

survive subsequent infection by *E. faecalis*, but this was not related to bacterial clearance [80]. *E.*

faecalis virulence is known to involve the action of toxins, including Cytolysin and Enterocin V, and

induced immune responses including AMPs and iron sequestration have minimal impact on *E.*

faecalis virulence [49, 81, 82]. The disconnect between mortality and replication could certainly be

explained by toxin-based virulence if low doses of toxins are sufficient to cause mortality as in other

Gram-positive insect pathogens [83] and cytolysin, for example, is hemolytic in animal cells in the

nanomolar range. Differences in the host response to bacterial toxins such as presence of Toll pathway

effector Baramicin A may explain the strong phylogenetic patterns of susceptibility to *E. faecalis*[41].

We found evidence of microbiome growth in four species (*D. americana*, *S. lebanonensis*, *D. putridia*

and *S. pattersoni*) when measuring bacterial loads in homogenized pathogen infected insects.

Anecdotally, these species exhibited a relatively high survival to most of the bacteria (Figure 3). The

microbiome can play an important role in priming the immune response and defending against

bacterial infections [84, 85]. Gut probiotics such as *Lactobacillus* have demonstrated to reduce fly

mortality caused by pathogenic bacteria [86]. It has been reported that there is limited variation in the

microbiome across species of Drosophilidae in laboratory culture [87, 88]. As such, a more

comprehensive study of how the microbiome varies across these host species, and whether this relates

to differences in susceptibility, is warranted.

In conclusion, we have found that the host phylogeny can explain varying amounts of variation in

susceptibility depending on the pathogen. These varying phylogenetic effects (or lack thereof) have

important consequences for understanding susceptibility across host species and its impact on

pathogen host shifts. For some pathogens, certain clades of related hosts will be particularly

susceptible, meaning pathogens will be more likely to jump into those species, and they could also act

as disease reservoirs. However, patterns may be less clear for other pathogens, with relatedness not

being a strong predictor of susceptibility across species. Further work is needed to understand the

mechanisms underlying the differences in susceptibility among species, and whether the same or

different factors affect susceptibility in different host clades.

Acknowledgements

Bacterial strains were kindly provided by Brian Lazzaro and Ashley Frank. Thanks to Megan Wallace

and Michael Jardine for their assistance in the code writing and experimental process for this study.

Supplementary materials

Table S1 Drosophilidae host species, wing length and rearing diet information.

Table S2 *Diptericin A* sanger sequencing details (primer, information of *Dpt A* S69R site encoded amino acid, GenBank accession number)

Table S3 Statistical coefficients for models fitting or not fitting Diptericin alleles as fixed effects terms.

Figure S1 Correlations between Bacterial load and mortality

Figure S2 Diptericin allele variation across Drosophilidae species

All data files and R scripts used in this study are available in an online repository:

https://figshare.com/s/4836105c5cef7aaad217

Author Contributions

Conceptualization: HS, BR, BL

Formal Experiments: HS

Data curation: HS, MA. H

Formal analysis: HS, RM. I, BR, BL

Investigation: HS, MA. H, RM. I, SK. W

Methodology: HS, MA. H, BR, BL

Supervision: BR, BL

Visualization: HS, RM. I, SK. W, BL

Writing- Manuscript draft: HS, MA. H, BR, BL

Writing- Reviewing and editing: HS, MA. H, RM. I, SK. W, BR, BL

Competing interests: We do not have any competing interests.

Funding:

This work has been supported by the following funding resources:

Hongbo Sun is supported by China Scholarship Council and University of Exeter PhD Scholarships

https://www.exeter.ac.uk/study/pg-research/csc-scholarships/

Ben Longdon and Ryan Imrie are supported by a Sir Henry Dale Fellowship jointly funded by the

Wellcome Trust and the Royal Society (grant no. 109356/Z/15/Z) https://wellcome.ac.uk/funding/sir-

henry-dale-fellowships.

Ben Raymond is supported by NERC NE/V012053/1 and BBSRC BB/S002928/.

Sarah K. Walsh is supported by a studentship funded by the Biotechnology and Biological Sciences

Research Council (BBSRC) South West Biosciences Doctoral Training Partnership (BB/M009122/1).

Mark A. Hanson is supported by Swiss National Science Foundation (SNSF) grant P500PB_211082

and a Wellcome Trust ECA Grant (grant no. 227559/Z/23/Z).

Reference

1.   Bonneaud C, Longdon B. Emerging pathogen evolution: Using evolutionary theory to
understand the fate of novel infectious pathogens. EMBO Rep. 2020;21(9):e51374. Epub 20200830.
doi: 10.15252/embr.202051374. PubMed PMID: 32864788; PubMed Central PMCID:
PMCPMC7507376.
2.   Petrosillo N. Emerging infections and future threats. Erciyes Medical Journal. 2019. doi:
10.14744/etd.2019.57805.
3.   Watkins K. Emerging Infectious Diseases: a Review. Curr Emerg Hosp Med Rep. 2018;6(3):86-
93. Epub 20180622. doi: 10.1007/s40138-018-0162-9. PubMed PMID: 32226656; PubMed Central
PMCID: PMCPMC7100414.
4.   Richardson EJ, Bacigalupe R, Harrison EM, Weinert LA, Lycett S, Vrieling M, et al. Gene
exchange drives the ecological success of a multi-host bacterial pathogen. Nat Ecol Evol.
2018;2(9):1468-78. Epub 20180723. doi: 10.1038/s41559-018-0617-0. PubMed PMID: 30038246;
PubMed Central PMCID: PMCPMC7610605.
5.   Mohapatra S, Menon NG. Factors responsible for the emergence of novel viruses: An
emphasis on SARS-CoV-2. Curr Opin Environ Sci Health. 2022;27:100358. Epub 20220328. doi:
10.1016/j.coesh.2022.100358. PubMed PMID: 35369608; PubMed Central PMCID:
PMCPMC8958772.
6.   Delaney NF, Balenger S, Bonneaud C, Marx CJ, Hill GE, Ferguson-Noel N, et al. Ultrafast
evolution and loss of CRISPRs following a host shift in a novel wildlife pathogen, Mycoplasma
gallisepticum. PLoS Genet. 2012;8(2):e1002511. Epub 20120209. doi:
10.1371/journal.pgen.1002511. PubMed PMID: 22346765; PubMed Central PMCID:
PMCPMC3276549.
7.   Murray S, Pascoe B, Méric G, Mageiros L, Yahara K, Hitchings MD, et al. Recombination-
Mediated Host Adaptation by Avian Staphylococcus aureus. Genome Biology and Evolution.
2017;9(4):830-42. doi: 10.1093/gbe/evx037.
8.   Sanaei E, Albery GF, Yeoh YK, Lin YP, Cook LG, Engelstadter J. Host phylogeny and ecological
associations best explain Wolbachia host shifts in scale insects. Mol Ecol. 2023;32(9):2351-63. Epub
20230303. doi: 10.1111/mec.16883. PubMed PMID: 36785954.
9.   Lowy FD. Staphylococcus aureus infections. N Engl J Med. 1998;339(8):520-32. doi:
10.1056/nejm199808203390806. PubMed PMID: 9709046.
10.   Cuny C, Friedrich A, Kozytska S, Layer F, Nübel U, Ohlsen K, et al. Emergence of methicillin-
resistant Staphylococcus aureus (MRSA) in different animal species. Int J Med Microbiol. 2010;300(2-
3):109-17. Epub 20091216. doi: 10.1016/j.ijmm.2009.11.002. PubMed PMID: 20005777.
11.   Price LB, Stegger M, Hasman H, Aziz M, Larsen J, Andersen PS, et al. Staphylococcus aureus
CC398: host adaptation and emergence of methicillin resistance in livestock. mBio. 2012;3(1). Epub
20120221. doi: 10.1128/mBio.00305-11. PubMed PMID: 22354957; PubMed Central PMCID:
PMCPMC3280451.
12.   Howden BP, Giulieri SG, Wong Fok Lung T, Baines SL, Sharkey LK, Lee JYH, et al.
Staphylococcus aureus host interactions and adaptation. Nat Rev Microbiol. 2023;21(6):380-95. Epub
20230127. doi: 10.1038/s41579-023-00852-y. PubMed PMID: 36707725; PubMed Central PMCID:
PMCPMC9882747.
13.   Fischer JR, Stallknecht DE, Luttrell P, Dhondt AA, Converse KA. Mycoplasmal conjunctivitis in
wild songbirds: the spread of a new contagious disease in a mobile host population. Emerg Infect Dis.
1997;3(1):69-72. doi: 10.3201/eid0301.970110. PubMed PMID: 9126448; PubMed Central PMCID:
PMCPMC2627586.
14.   Bonneaud C, Balenger SL, Zhang J, Edwards SV, Hill GE. Innate immunity and the evolution of
resistance to an emerging infectious disease in a wild bird. Mol Ecol. 2012;21(11):2628-39. Epub
20120419. doi: 10.1111/j.1365-294X.2012.05551.x. PubMed PMID: 22512302.

15.   Fleming-Davies AE, Williams PD, Dhondt AA, Dobson AP, Hochachka WM, Leon AE, et al.
Incomplete host immunity favors the evolution of virulence in an emergent pathogen. Science.
2018;359(6379):1030-3. doi: doi:10.1126/science.aao2140.
16.   European Food Safety A, Delbianco A, Gibin D, Pasinato L, Boscia D, Morelli M. Update of the
Xylella spp. host plant database - systematic literature search up to 31 December 2021. EFSA J.
2022;20(6):e07356. Epub 20220615. doi: 10.2903/j.efsa.2022.7356. PubMed PMID: 35734284;
PubMed Central PMCID: PMCPMC9198695.
17.   Donegan MA, Coletta-Filho HD, Almeida RPP. Parallel host shifts in a bacterial plant pathogen
suggest independent genetic solutions. Mol Plant Pathol. 2023;24(6):527-35. Epub 20230329. doi:
10.1111/mpp.13316. PubMed PMID: 36992605; PubMed Central PMCID: PMCPMC10189759.
18.   Tinsley MC, Majerus MEN. Small steps or giant leaps for male-killers? Phylogenetic
constraints to male-killer host shifts. BMC Evolutionary Biology. 2007;7(1):238. doi: 10.1186/1471-
2148-7-238.
19.   Weinert LA, Araujo-Jnr EV, Ahmed MZ, Welch JJ. The incidence of bacterial endosymbionts in
terrestrial arthropods. Proc Biol Sci. 2015;282(1807):20150249. doi: 10.1098/rspb.2015.0249.
PubMed PMID: 25904667; PubMed Central PMCID: PMCPMC4424649.
20.   Hilgenboecker K, Hammerstein P, Schlattmann P, Telschow A, Werren JH. How many species
are infected with Wolbachia?--A statistical analysis of current data. FEMS Microbiol Lett.
2008;281(2):215-20. Epub 20080228. doi: 10.1111/j.1574-6968.2008.01110.x. PubMed PMID:
18312577; PubMed Central PMCID: PMCPMC2327208.
21.   Zug R, Hammerstein P. Still a host of hosts for Wolbachia: analysis of recent data suggests
that 40% of terrestrial arthropod species are infected. PLoS One. 2012;7(6):e38544. Epub 20120607.
doi: 10.1371/journal.pone.0038544. PubMed PMID: 22685581; PubMed Central PMCID:
PMCPMC3369835.
22.   Russell JA, Goldman-Huertas B, Moreau CS, Baldo L, Stahlhut JK, Werren JH, Pierce NE.
Specialization and geographic isolation among Wolbachia symbionts from ants and lycaenid
butterflies. Evolution. 2009;63(3):624-40. Epub 20081119. doi: 10.1111/j.1558-5646.2008.00579.x.
PubMed PMID: 19054050.
23.   Sanaei E, Charlat S, Engelstädter J. Wolbachia host shifts: routes, mechanisms, constraints
and evolutionary consequences. Biol Rev Camb Philos Soc. 2021;96(2):433-53. Epub 20201030. doi:
10.1111/brv.12663. PubMed PMID: 33128345.
24.   Longdon B, Brockhurst MA, Russell CA, Welch JJ, Jiggins FM. The evolution and genetics of
virus host shifts. PLoS Pathog. 2014;10(11):e1004395. Epub 20141106. doi:
10.1371/journal.ppat.1004395. PubMed PMID: 25375777; PubMed Central PMCID:
PMCPMC4223060.
25.   Longdon B, Hadfield JD, Webster CL, Obbard DJ, Jiggins FM. Host phylogeny determines viral
persistence and replication in novel hosts. PLoS Pathog. 2011;7(9):e1002260. Epub 20110922. doi:
10.1371/journal.ppat.1002260. PubMed PMID: 21966271; PubMed Central PMCID:
PMCPMC3178573.
26.   Imrie RM, Roberts KE, Longdon B. Between virus correlations in the outcome of infection
across host species: Evidence of virus by host species interactions. Evol Lett. 2021;5(5):472-83. Epub
20210715. doi: 10.1002/evl3.247. PubMed PMID: 34621534; PubMed Central PMCID:
PMCPMC8484721.
27.   Huan Y, Kong Q, Mou H, Yi H. Antimicrobial Peptides: Classification, Design, Application and
Research Progress in Multiple Fields. Front Microbiol. 2020;11:582779. Epub 20201016. doi:
10.3389/fmicb.2020.582779. PubMed PMID: 33178164; PubMed Central PMCID: PMCPMC7596191.
28.   Hoffmann JA, Reichhart JM. Drosophila innate immunity: an evolutionary perspective. Nat
Immunol. 2002;3(2):121-6. doi: 10.1038/ni0202-121. PubMed PMID: 11812988.
29.   Lemaitre B, Hoffmann J. The host defense of Drosophila melanogaster. Annu Rev Immunol.
2007;25:697-743. doi: 10.1146/annurev.immunol.25.022106.141615. PubMed PMID: 17201680.

30.   Buchon N, Silverman N, Cherry S. Immunity in Drosophila melanogaster--from microbial
recognition to whole-organism physiology. Nat Rev Immunol. 2014;14(12):796-810. Epub
2014/11/26. doi: 10.1038/nri3763. PubMed PMID: 25421701; PubMed Central PMCID:
PMCPMC6190593.
31.   Sackton TB, Lazzaro BP, Schlenke TA, Evans JD, Hultmark D, Clark AG. Dynamic evolution of
the innate immune system in Drosophila. Nature Genetics. 2007;39(12):1461-8. doi:
10.1038/ng.2007.60.
32.   Obbard DJ, Welch JJ, Kim K-W, Jiggins FM. Quantifying Adaptive Evolution in the Drosophila
Immune System. PLOS Genetics. 2009;5(10):e1000698. doi: 10.1371/journal.pgen.1000698.
33.   Unckless RL, Howick VM, Lazzaro BP. Convergent balancing selection on an antimicrobial
peptide in Drosophila. Current Biology. 2016;26(2):257-62.
34.   Hanson MA, Grollmus L, Lemaitre B. Ecology-relevant bacteria drive the evolution of host
antimicrobial peptides in Drosophila. SCIENCE. 2023;381(6655):284-+. doi: 10.1126/science.adg5725.
PubMed PMID: WOS:001046763100001.
35.   Duneau D, Ferdy JB, Revah J, Kondolf H, Ortiz GA, Lazzaro BP, Buchon N. Stochastic variation
in the initial phase of bacterial infection predicts the probability of survival in D. melanogaster. Elife.
2017;6. Epub 20171012. doi: 10.7554/eLife.28298. PubMed PMID: 29022878; PubMed Central
PMCID: PMCPMC5703640.
36.   Needham AJ, Kibart M, Crossley H, Ingham PW, Foster SJ. Drosophila melanogaster as a
model host for Staphylococcus aureus infection. Microbiology. 2004;150(7):2347-55. doi:
https://doi.org/10.1099/mic.0.27116-0.
37.   Clewell DB, Tomich PK, Gawron-Burke MC, Franke AE, Yagi Y, An FY. Mapping of Streptococcus
faecalis plasmids pAD1 and pAD2 and studies relating to transposition of Tn917. J Bacteriol.
1982;152(3):1220-30. doi: 10.1128/jb.152.3.1220-1230.1982. PubMed PMID: 6292164; PubMed
Central PMCID: PMCPMC221629.
38.   Cox CR, Gilmore MS. Native microbial colonization of Drosophila melanogaster and its use as
a model of Enterococcus faecalis pathogenesis. Infect Immun. 2007;75(4):1565-76. Epub 20070112.
doi: 10.1128/IAI.01496-06. PubMed PMID: 17220307; PubMed Central PMCID: PMCPMC1865669.
39.   Xu R, Lou Y, Tidu A, Bulet P, Heinekamp T, Martin F, et al. The Toll pathway mediates
Drosophila resilience to Aspergillus mycotoxins through specific Bomanins. EMBO reports.
2023;24(1):e56036. doi: https://doi.org/10.15252/embr.202256036.
40.   Hanson MA, Cohen LB, Marra A, Iatsenko I, Wasserman SA, Lemaitre B. The Drosophila
Baramicin polypeptide gene protects against fungal infection. PLOS Pathogens. 2021;17(8):e1009846.
doi: 10.1371/journal.ppat.1009846.
41.   Huang J, Lou Y, Liu J, Bulet P, Cai C, Ma K, et al. A Toll pathway effector protects Drosophila
specifically from distinct toxins secreted by a fungus or a bacterium. Proceedings of the National
Academy of Sciences. 2023;120(12):e2205140120. doi: doi:10.1073/pnas.2205140120.
42.   Liu GY, Essex A, Buchanan JT, Datta V, Hoffman HM, Bastian JF, et al. Staphylococcus aureus
golden pigment impairs neutrophil killing and promotes virulence through its antioxidant activity. J
Exp Med. 2005;202(2):209-15. Epub 20050711. doi: 10.1084/jem.20050846. PubMed PMID:
16009720; PubMed Central PMCID: PMCPMC2213009.
43.   Binggeli O, Neyen C, Poidevin M, Lemaitre B. Prophenoloxidase Activation Is Required for
Survival to Microbial Infections in Drosophila. PLOS Pathogens. 2014;10(5):e1004067. doi:
10.1371/journal.ppat.1004067.
44.   Dudzic JP, Hanson MA, Iatsenko I, Kondo S, Lemaitre B. More than black or white:
melanization and toll share regulatory serine proteases in Drosophila. Cell reports. 2019;27(4):1050-
61. e3.
45.   Mulet M, Gomila M, Lemaitre B, Lalucat J, García-Valdés E. Taxonomic characterisation of
Pseudomonas strain L48 and formal proposal of Pseudomonas entomophila sp. nov. Syst Appl
Microbiol. 2012;35(3):145-9. Epub 20120210. doi: 10.1016/j.syapm.2011.12.003. PubMed PMID:
22326814.

46.   Vodovar N, Vinals M, Liehl P, Basset A, Degrouard J, Spellman P, et al. Drosophila host
defense after oral infection by an entomopathogenic Pseudomonas species. Proc Natl Acad Sci U S A.
2005;102(32):11414-9. Epub 20050801. doi: 10.1073/pnas.0502240102. PubMed PMID: 16061818;
PubMed Central PMCID: PMCPMC1183552.
47.   Dieppois G, Opota O, Lalucat J, Lemaitre B. Pseudomonas entomophila: A Versatile Bacterium
with Entomopathogenic Properties. In: Ramos J-L, Goldberg JB, Filloux A, editors. Pseudomonas:
Volume 7: New Aspects of Pseudomonas Biology. Dordrecht: Springer Netherlands; 2015. p. 25-49.
48.   Juneja P, Lazzaro BP. Providencia sneebia sp. nov. and Providencia burhodogranariea sp. nov.,
isolated from wild Drosophila melanogaster. Int J Syst Evol Microbiol. 2009;59(Pt 5):1108-11. Epub
2009/05/02. doi: 59/5/1108 [pii]
10.1099/ijs.0.000117-0. PubMed PMID: 19406801.
49.   Hanson MA, Dostalova A, Ceroni C, Poidevin M, Kondo S, Lemaitre B. Synergy and
remarkable specificity of antimicrobial peptides in vivo using a systematic knockout approach. Elife.
2019;8. Epub 20190226. doi: 10.7554/eLife.44341. PubMed PMID: 30803481; PubMed Central
PMCID: PMCPMC6398976.
50.   Unckless RL, Howick VM, Lazzaro BP. Convergent Balancing Selection on an Antimicrobial
Peptide in Drosophila. Curr Biol. 2016;26(2):257-62. Epub 20160114. doi: 10.1016/j.cub.2015.11.063.
PubMed PMID: 26776733; PubMed Central PMCID: PMCPMC4729654.
51.   Short SM, Lazzaro BP. Female and male genetic contributions to post-mating immune
defence in female Drosophila melanogaster. Proc Biol Sci. 2010;277(1700):3649-57. Epub 20100623.
doi: 10.1098/rspb.2010.0937. PubMed PMID: 20573620; PubMed Central PMCID: PMCPMC2982248.
52.   Schwenke RA, Lazzaro BP. Juvenile Hormone Suppresses Resistance to Infection in Mated
Female Drosophila melanogaster. Curr Biol. 2017;27(4):596-601. Epub 20170209. doi:
10.1016/j.cub.2017.01.004. PubMed PMID: 28190728; PubMed Central PMCID: PMCPMC5319889.
53.   Duneau DF, Kondolf HC, Im JH, Ortiz GA, Chow C, Fox MA, et al. The Toll pathway underlies
host sexual dimorphism in resistance to both Gram-negative and Gram-positive bacteria in mated
Drosophila. BMC Biol. 2017;15(1):124. Epub 20171221. doi: 10.1186/s12915-017-0466-3. PubMed
PMID: 29268741; PubMed Central PMCID: PMCPMC5740927.
54.   Larson EL, Strom MS, Evans CA. Analysis of three variables in sampling solutions used to
assay bacteria of hands: type of solution, use of antiseptic neutralizers, and solution temperature.
Journal of Clinical Microbiology. 1980;12(3):355-60. doi: doi:10.1128/jcm.12.3.355-360.1980.
55.   Longdon B, Hadfield JD, Day JP, Smith SC, McGonigle JE, Cogni R, et al. The causes and
consequences of changes in virulence following pathogen host shifts. PLoS Pathog.
2015;11(3):e1004728. Epub 20150316. doi: 10.1371/journal.ppat.1004728. PubMed PMID:
25774803; PubMed Central PMCID: PMCPMC4361674.
56.   Drummond AJ, Suchard MA, Xie D, Rambaut A. Bayesian phylogenetics with BEAUti and the
BEAST 1.7. Mol Biol Evol. 2012;29(8):1969-73. Epub 20120225. doi: 10.1093/molbev/mss075.
PubMed PMID: 22367748; PubMed Central PMCID: PMCPMC3408070.
57.   Rambaut A, Drummond AJ, Xie D, Baele G, Suchard MA. Posterior Summarization in Bayesian
Phylogenetics Using Tracer 1.7. Syst Biol. 2018;67(5):901-4. doi: 10.1093/sysbio/syy032. PubMed
PMID: 29718447; PubMed Central PMCID: PMCPMC6101584.
58.   Yu G. Using ggtree to Visualize Data on Tree-Like Structures. Current Protocols in
Bioinformatics. 2020;69(1):e96. doi: https://doi.org/10.1002/cpbi.96.
59.   Jarrod H. MCMC Methods for Multi-Response Generalized Linear Mixed Models: The
MCMCglmm R Package. Journal of Statistical Software. 2010;33.
60.   R: A Language and Environment for Statistical Computing [Internet]. R Foundation for
Statistical Computing. 2023. Available from: https://www.R-project.org/.
61.   Gilchrist G, Huey R, Serra L. Rapid evolution of wing size clines in Drosophila subobscura.
Genetica. 2001;112-113:273-86. doi: 10.1023/A:1013358931816.

62.   Huey RB, Moreteau B, Moreteau J-C, Gibert P, Gilchrist GW, Ives AR, et al. Sexual size
dimorphism in a Drosophila clade, the D. obscura group. Zoology. 2006;109(4):318-30. doi:
https://doi.org/10.1016/j.zool.2006.04.003.
63.   R. P. Freckleton, P. H. Harvey, M. Pagel. Phylogenetic Analysis and Comparative Data: A Test
and Review of Evidence. The American Naturalist. 2002;160(6):712-26. doi: 10.1086/343873.
PubMed PMID: 18707460.
64.   WILSON AJ. Why h2 does not always equal VA/VP? Journal of Evolutionary Biology.
2008;21(3):647-50. doi: https://doi.org/10.1111/j.1420-9101.2008.01500.x.
65.   Tokuda T, Goodrich B, Mechelen IV, Gelman A, Tuerlinckx F, editors. Visualizing Distributions
of Covariance Matrices *2011.
66.   Freckleton RP, Harvey PH, Pagel M. Phylogenetic analysis and comparative data: a test and
review of evidence. Am Nat. 2002;160(6):712-26. doi: 10.1086/343873. PubMed PMID: 18707460.
67.   Elizabeth A. Housworth, Emília P. Martins, Michael Lynch. The Phylogenetic Mixed Model.
The American Naturalist. 2004;163(1):84-96. doi: 10.1086/380570. PubMed PMID: 14767838.
68.   Pagel M. Inferring the historical patterns of biological evolution. Nature.
1999;401(6756):877-84. doi: 10.1038/44766.
69.   Falconer D, Mackay T. Introduction to quantitative genetics. Essex. UK: Longman Group.
1996;12.
70.   De Gregorio E, Spellman PT, Tzou P, Rubin GM, Lemaitre B. The Toll and Imd pathways are the
major regulators of the immune response in Drosophila. Embo j. 2002;21(11):2568-79. doi:
10.1093/emboj/21.11.2568. PubMed PMID: 12032070; PubMed Central PMCID: PMCPMC126042.
71.   Troha K, Im JH, Revah J, Lazzaro BP, Buchon N. Comparative transcriptomics reveals CrebA as
a novel regulator of infection tolerance in D. melanogaster. PLOS Pathogens. 2018;14(2):e1006847.
doi: 10.1371/journal.ppat.1006847.
72.   Rutschmann S, Kilinc A, Ferrandon D. Cutting edge: the toll pathway is required for resistance
to gram-positive bacterial infections in Drosophila. J Immunol. 2002;168(4):1542-6. doi:
10.4049/jimmunol.168.4.1542. PubMed PMID: 11823479.
73.   Ambrosi P, Chahda JS, Koslen HR, Chiel HJ, Mizutani CM. Modeling of the dorsal gradient
across species reveals interaction between embryo morphology and Toll signaling pathway during
evolution. PLoS Comput Biol. 2014;10(8):e1003807. Epub 20140828. doi:
10.1371/journal.pcbi.1003807. PubMed PMID: 25165818; PubMed Central PMCID:
PMCPMC4148200.
74.   Palmer WJ, Jiggins FM. Comparative Genomics Reveals the Origins and Diversity of Arthropod
Immune Systems. Mol Biol Evol. 2015;32(8):2111-29. Epub 20150422. doi: 10.1093/molbev/msv093.
PubMed PMID: 25908671; PubMed Central PMCID: PMCPMC4833078.
75.   Sackton TB, Lazzaro BP, Schlenke TA, Evans JD, Hultmark D, Clark AG. Dynamic evolution of
the innate immune system in Drosophila. Nat Genet. 2007;39(12):1461-8. Epub 20071107. doi:
10.1038/ng.2007.60. PubMed PMID: 17987029.
76.   Nishide Y, Kageyama D, Yokoi K, Jouraku A, Tanaka H, Futahashi R, Fukatsu T. Functional
crosstalk across IMD and Toll pathways: insight into the evolution of incomplete immune cascades.
Proc Biol Sci. 2019;286(1897):20182207. doi: 10.1098/rspb.2018.2207. PubMed PMID: 30963836;
PubMed Central PMCID: PMCPMC6408883.
77.   Kounatidis I, Ligoxygakis P. Drosophila as a model system to unravel the layers of innate
immunity to infection. Open Biol. 2012;2(5):120075. doi: 10.1098/rsob.120075. PubMed PMID:
22724070; PubMed Central PMCID: PMCPMC3376734.
78.   Valanne S, Myllymäki H, Kallio J, Schmid MR, Kleino A, Murumägi A, et al. Genome-wide RNA
interference in Drosophila cells identifies G protein-coupled receptor kinase 2 as a conserved
regulator of NF-kappaB signaling. J Immunol. 2010;184(11):6188-98. Epub 20100426. doi:
10.4049/jimmunol.1000261. PubMed PMID: 20421637.

79.   Chapman JR, Dowell MA, Chan R, Unckless RL. The Genetic Basis of Natural Variation in
Drosophila melanogaster Immune Defense against Enterococcus faecalis. Genes. 2020;11(2):234.
PubMed PMID: doi:10.3390/genes11020234.
80.   Cabrera K, Hoard DS, Gibson O, Martinez DI, Wunderlich Z. Drosophila immune priming to
Enterococcus faecalis relies on immune tolerance rather than resistance. PLoS Pathog.
2023;19(8):e1011567. Epub 20230811. doi: 10.1371/journal.ppat.1011567. PubMed PMID:
37566589; PubMed Central PMCID: PMCPMC10446173.
81.   Iatsenko I, Marra A, Boquete J-P, Peña J, Lemaitre B. Iron sequestration by transferrin 1
mediates nutritional immunity in Drosophila melanogaster. Proceedings of the National Academy of
Sciences. 2020;117(13):7317-25. doi: doi:10.1073/pnas.1914830117.
82.   Wadhawan A, Silva CJSd, Nunes CD, Edwards AM, Dionne MS. E. faecalis acquires resistance
to antimicrobials and insect immunity via common mechanisms. bioRxiv. 2022:2022.08.17.504265.
doi: 10.1101/2022.08.17.504265.
83.   Cornforth DM, Matthews A, Brown SP, Raymond B. Bacterial cooperation causes systematic
errors in pathogen risk assessment due to the failure of the independent action hypothesis. PLoS
Pathogens. 2015;11(4):e1004775. doi: 10.1371/journal.ppat.1004775.s002.
84.   Panwar RB, Sequeira RP, Clarke TB. Microbiota-mediated protection against antibiotic-
resistant pathogens. Genes & Immunity. 2021;22(5):255-67. doi: 10.1038/s41435-021-00129-5.
85.   Barron AJ, Lesperance DNA, Doucette J, Calle S, Broderick NA. Microbiome derived acidity
protects against microbial invasion in Drosophila. bioRxiv. 2023. Epub 20230112. doi:
10.1101/2023.01.12.523836. PubMed PMID: 36711873; PubMed Central PMCID: PMCPMC9882140.
86.   Blum JE, Fischer CN, Miles J, Handelsman J. Frequent Replenishment Sustains the Beneficial
Microbiome of Drosophila melanogaster. MBIO. 2013;4(6). doi: 10.1128/mBio.00860-13. PubMed
PMID: WOS:000329174500030.
87.   Chandler JA, Morgan Lang J, Bhatnagar S, Eisen JA, Kopp A. Bacterial Communities of Diverse
Drosophila Species: Ecological Context of a Host–Microbe Model System. PLOS Genetics.
2011;7(9):e1002272. doi: 10.1371/journal.pgen.1002272.
88.   Brooks AW, Kohl KD, Brucker RM, van Opstal EJ, Bordenstein SR. Phylosymbiosis:
Relationships and Functional Effects of Microbial Communities across Host Evolutionary History. PLOS
Biology. 2016;14(11):e2000225. doi: 10.1371/journal.pbio.2000225.
787