Divergence in a Eukaryotic Transcription Factor's co-TF Dependence Involves Multiple Intrinsically Disordered Regions

Lindsey F. Snyder[1], Emily M. O'Brien[2], Jia Zhao[2#], Jinye Liang[2], Baylee J. Bruce[1],

Yuning Zhang[3^], Wei Zhu[4], Thomas H. Cassier[2], Nicholas J. Schnicker[5, 6],

Xu Zhou[7], Raluca Gordân[3, 4, 8, 9, 10], Bin Z. He[1,2,*]

[1] Interdisciplinary Graduate Program in Genetics, University of Iowa, Iowa City, IA, USA; [2] Department of Biology, University of Iowa, Iowa City, IA, USA; [3] Department of Biostatistics & Bioinformatics, Duke University, Durham, NC, USA; [4] Department of Molecular Genetics & Microbiology, Duke University, Durham, NC, USA; [5] Protein and Crystallography Facility, University of Iowa, Iowa City, IA, USA; [6] Department of Molecular Physiology and Biophysics, University of Iowa, Iowa City, IA, USA; [7] Department of Pediatrics, Division of Gastroenterology, Hepatology and Nutrition, Boston Children's Hospital and Harvard Medical School, Boston, MA, USA; [8] Department of Computer Science, Duke University, Durham, NC, USA; [9] Department of Cell Biology, Duke University, Durham, NC, USA; [10] Department of Genomics and Computational Biology, University of Massachusetts Chan Medical School, Worcester, MA, USA

[#] Current address: Laboratory of Immunophysiology, The Ragon Institute of Mass General, MIT, and Harvard; Department of Biology, Massachusetts Institute of Technology. Cambridge, MA, USA.

[^] Current address: Department of Genetics, Washington University School of Medicine in St. Louis, St. Louis, MO, USA.

[*] correspondence should be addressed to bin-he@uiowa.edu

Running title: Divergence in a eukaryotic TF's dependence on co-TF

ABSTRACT

Combinatorial control by transcription factors (TFs) is central to eukaryotic gene regulation, yet

its mechanism, evolution, and regulatory impact are not well understood. Here we use natural

variation in the yeast phosphate starvation (PHO) response to examine the genetic basis and

species variation in TF interdependence. In *Saccharomyces cerevisiae*, main TF Pho4 relies on

the co-TF Pho2 to regulate ~28 genes, whereas in the related pathogen *Candida glabrata*, Pho4

has reduced Pho2 dependence and regulates ~70 genes. We found *C. glabrata* Pho4 (CgPho4)

binds the same motif with 3-4 fold higher affinity. Machine learning and yeast one-hybrid assay

identify two intrinsically disordered regions (IDRs) in CgPho4 that boost its activation domain's

activity. In ScPho4, an IDR next to the DNA binding domain both allows for enhanced activity

with Pho2 and inhibits activity without Pho2. This study reveals how IDR divergence drives TF

interdependence evolution by influencing activation potential and autoinhibition.

INTRODUCTION

Transcription factors (TFs) are the cornerstone of gene regulatory networks. In eukaryotes, TFs often work collaboratively to regulate gene expression. This combinatorial control is crucial for enhancing specificity, because most eukaryotic TFs recognize short and degenerate motifs – typically less than 10 bps – that appear hundreds to tens of thousands of times in the genome [1]. TFs with the same family of DNA binding domains (DBDs) also recognize highly similar motifs [2]. Despite these, TFs often bind only a fraction of their motifs *in vivo* and regulate an even smaller subset of the genes they bind to [3]; paralogous TFs regulate distinct sets of genes [4,5]. The key to achieving this specific regulation is the requirement of two or more TFs to jointly regulate the target genes [6–8]. Another important function of combinatorial control and TF interdependence is to allow cells to integrate multiple upstream signals. In fly development, for example, a combination of tissue-specific "selector" and morphogen gradient – both are TFs – precisely determine the expression pattern and define the cell fate [9].

The importance of combinatorial control and the molecular mechanisms behind TF interdependence are traditionally studied using mutations that disrupt TF interactions. For instance, a missense mutation that disrupts the interaction between two cardiac TFs, GATA4 and TBX5, was shown to cause dysregulation of cardiac genes and lead to malformation of heart tissues [10]. Similarly, studies of combinatorial control evolution often focus on losses (and gains) of TF interactions, like the MADS-box TF, Mcm1, and its co-TFs, MATa2 and MATα2 in the yeast mating type pathway [11,12]. A far less explored type of change in the combinatorial control is a change in the TF interdependence itself, that is, when TFs involved in cooperative regulation evolved to be more or less dependent on the co-TF(s). Such changes are expected to dramatically rewire the network by altering the specificity and signals required for gene regulation, which could have profound implications in disease and evolution. In relation to this possibility, not all eukaryotic TFs require other TFs to function. The yeast TF, Gal4, regulates its target genes on its own [13]. This raises several intriguing and unanswered questions: if TF

interdependence and combinatorial control evolve, what genetic changes underlie the

divergence, what aspects of the TF activities are impacted, and what are the consequences on

the regulatory output?

Intriguingly, natural variation in co-TF dependence exists in the yeast phosphate

starvation (PHO) response network. In the model yeast *Saccharomyces cerevisiae*, the main TF

of the PHO response, Pho4 (hereinafter as ScPho4), strongly depends on the co-TF, Pho2, to

induce 27/28 of its target genes [3]. In a related human yeast pathogen, *Candida glabrata*, its

Pho4 (hereinafter as CgPho4) is far less dependent on Pho2 and induces twice as many genes

(Fig. 1A) [14,15]. The level of Pho2-dependence varies quantitatively among Pho4 orthologs in

other yeasts and is correlated with the number of genes induced in a common genome

background [15]. This latter observation is consistent with the role of combinatorial control in

enhancing specificity. What remains unknown is what genetic differences between Pho4

orthologs underlie the divergence in co-TF dependence, and what TF activities are impacted by

those variations.

Here, we propose two non-mutually exclusive models for the difference in Pho2-

dependence between ScPho4 and CgPho4 (Fig. 1B). The first "enhanced activity" model is

based on studies showing that ScPho4 requires Pho2 to **(1)** bind cooperatively to the target

gene promoters [16], **(2)** recruit the histone acetyltransferase (HAT) complex [17] and **(3)** help recruit

general TFs and the PolII complex [18]. Under this model, we hypothesize that CgPho4 binds

more tightly to DNA than ScPho4, and is more capable of recruiting general transcription factors

(TFs) and the PolII complex, thereby becoming less Pho2-dependent. The second

"autoinhibition" model is based on a study suggesting that ScPho4 is auto-inhibited, and that

interaction with Pho2 unmasks its activation domain and allows it to function [19]. This model

predicts that CgPho4 either lacks or has far weaker effects of the auto-inhibition and hence

doesn't depend on Pho2.

In this study, we tested these two models by comparing the DNA binding and activation

abilities of the two Pho4 orthologs, and systematically swapping regions between the two Pho4s

in a series of 50 chimeric TFs, then quantifying their activities with and without Pho2. Our results

support both models as contributing to the Pho2-dependence variation. To our surprise, while

CgPho4 DBD binds to the same consensus motif with a 3-4-fold higher affinity than ScPho4

DBD, swapping DBD alone failed to yield the expected results. Instead, the differences in Pho2-

dependence originated primarily from differences in the Intrinsically Disordered Regions (IDRs)

in the two TFs that modulate both their Activation Domain (AD) and DNA binding domain (DBD)

activities. Therefore, our results reveal that evolution in a eukaryotic TF protein, particularly

through changes in the IDR, can lead to divergence in co-TF dependence, which in turn results

in a more than two-fold change in the size of the target network.

97   **RESULTS**

Domain organization and sequence divergence between ScPho4 and CgPho4

Based on genetic and biochemical studies, ScPho4 encodes the following functional domains

from its N- to its C-terminus (Fig. 1C): a regulatory region (R1, aa 1-42) interacting with the

negative regulator Pho80 [20], the activation domain (AD, aa 43-99) [21,22], a region encoding the

nuclear export and import signals (NLS, aa 100-176) [23,24], a protein-interaction domain

interacting with both Pho80 and the co-TF, Pho2 (referred together as P2ID, aa 177-242) [20,25],

and the bHLH DNA binding domain (DBD, aa 243-312) [21,26]. Regarding ScPho4-ScPho2

interaction, a previous study mapped the region in ScPho4 required for the interaction to aa

200-218 [25]. The same 18 aa stretch also contains the phosphorylation site that directly controls

the TF-TF interaction [23].

108   Both ScPho4 and CgPho4 were predicted to be mostly intrinsically disordered outside

the DBD, with some low confidence helices in R1 and AD (Supplementary Fig. 1). CgPho4 is

significantly longer than ScPho4 (533 vs 312 amino acids, Fig. 1C). The NLS and P2ID regions

contain four functionally important phosphorylation sites targeted by the cyclin-dependent

kinase complex Pho80/85 [24,27]. Those located in the NLS control Pho4's nuclear translocation,

while phosphorylation of the one in P2ID disables ScPho4 from interacting with Pho2 [23]. All five

Pho80/85 targeted phosphorylation sites are clearly identifiable in the alignment (Supplementary

Fig. 2); in fact, they were previously found to be conserved in orthologs outside the

Saccharomycotina subphylum [28]. Thus, while the protein length and sequence diverged

significantly between the two Pho4 orthologs, the domain architecture and post-translational

modification motifs appear to be highly conserved in evolution.

CgPho4 binds the same consensus motif with a higher affinity compared to ScPho4

To test the enhanced activity model, we first compared the DNA binding ability of the two Pho4

orthologs. First, we asked if the two Pho4 orthologs recognize the same DNA sequence. The N-

terminal stretch of the first α-helix in the bHLH domain, known as the basic region, contains the

residues determining sequence specificity. All four residues known to recognize the nucleotide

bases and five additional residues that recognize the phosphate backbone in ScPho4 [26] are all

conserved in CgPho4 except for an R252K change (number based on ScPho4, Fig. 2A). Next,

we used our published Chromatin-Immunoprecipitation (ChIP-seq) data for both Pho4 orthologs

to identify their respective motifs [15] (Materials and Methods). Consistent with the amino acid

sequence conservation, the results showed that they bind the same E-box motif "CACGTG",

with no obvious differences in the flanking nucleotides (Fig. 2B). These motifs were based on a

relatively small number of ChIP peaks (74 for ScPho4 and 118 for CgPho4). To

comprehensively characterize and compare the binding preferences of the two proteins, we

applied Protein Binding Microarray (PBM) to map the binding landscape in the entire 7-mer

space (universal PBM, Fig. 2C). The result revealed a lack of divergence in their sequence

preference (Pearson's r = 0.89). To complement the short oligo length in the universal PBM and

further examine differences in the flanking base preference, we designed a second genome

context library, which includes 36-bp sequences centered on ChIP-identified binding sites for

both Pho4's in their native genome along with the flanking nucleotides [29]. Similar to the uPBM,

the genome context PBM revealed no evidence for binding specificity differences between

ScPho4 and CgPho4 (Supplementary Fig. 3, Spearman's ρ = 0.84 and 0.86 for sequences

containing the consensus E-box or non-consensus variants, respectively, Materials and

Methods). We conclude that ScPho4 and CgPho4 have the same sequence specificity.

Next, we tested the hypothesis that CgPho4 binds more tightly than ScPho4 does. To do

so, we purified the DBD region of both Pho4s (Materials and Methods) and measured their

binding affinity to a 17-bp oligo based on the *S. cerevisiae PHO5* promoter with the consensus

E-box motif "CACGTG". Biolayer Interferometry (BLI) measurements showed that CgPho4 DBD

binds the consensus motif >3 times more tightly than ScPho4 DBD does (K_D = 5.2 nM vs 1.2 nM

for ScPho4 and CgPho4, respectively; Student's t-test for log K_D difference *P* < 0.01, Fig. 2D,

E). We confirmed this K_D difference using Electrophoretic Mobility Shift Assay (K_D = 5.5 nM vs

1.9 nM for ScPho4 and CgPho4, respectively; Supplementary Fig. 4). We conclude that CgPho4

recognizes the same E-box motif and binds with a higher affinity than ScPho4. It is worth noting

that this observed difference in affinity, while statistically significant, is modest ($\Delta\Delta G$ = -0.87

kcal/mol, based on the mean BLI measurements, assuming a temperature of 25°C). Its potential

impact on gene regulation depends critically on the effective concentration of Pho4 in the

nucleus.

CgPho4 encodes two additional Activation Enhancer Domains (AEDs)

We next tested the hypothesis that CgPho4 has increased transactivation potential compared to

ScPho4 under the enhanced activity model. We first predicted regions with activation potential

in both Pho4 orthologs using PADDLE, a Convolutional Neural Network trained on 150

activation domains (ADs) from 164 TFs in *S. cerevisiae* [30]. PADDLE recovered the

experimentally identified AD in ScPho4 between aa 60-102, which contained the 9aaTAD motif

previously described as the minimum residues required for activation [31] (Fig. 3A, orange

triangle). No other regions with significant activation potential were predicted in ScPho4. By

contrast, three regions in CgPho4 were predicted to have activation capabilities, including one

corresponding to the AD in ScPho4, which was predicted to have strong activity (Z-score > 6,

Fig. 3B). Two additional regions, one overlapping R1 and the other spanning the NLS and P2ID,

were predicted to have medium activation strength (Z-score > 4). We refer to these two regions

as E1 and E2, respectively from here on. Interestingly, the best match to the 9aaTAD motif

pattern in CgPho4 was found in E2 rather than in the AD (Fig. 3B).

To determine the activity of the predicted regions with activation potential in both Pho4s,

we set up a yeast one-hybrid (Y1H) assay, in which each candidate region was fused to the

Gal4 DBD and its activation potential was measured by a genome-integrated *GAL1*pr-mCherry

reporter in *S. cerevisiae* (Fig. 3C, Materials and Methods). To see if fusing the candidate region

with Gal4 DBD created a new activation domain, we applied PADDLE to all constructs and

observed no peaks either in the Gal4 DBD or in the connecting regions (Supplementary Fig. 5).

The yeast one-hybrid result confirmed that both ScPho4 and CgPho4 ADs were able to activate

the reporter above the background level (7- and 5.6-fold, Holm-Bonferroni corrected $P < 0.01$).

We were able to further localize the required residues for activation in ScPho4's AD to a region

of 32 aa centered on the 9aaTAD motif (Fig. 3A, C). Neither E1 nor E2 from either Pho4

activated the reporter on its own (Fig. 3C). This was surprising for CgPho4's E1 and E2, since

both had a medium strength Z-score prediction, and the latter also contained a match to the

9aaTAD (Fig. 3B). We hypothesized that E1 and E2 in CgPho4 could enhance the activity of the

main AD. To test this, we first fused ScE1 and CgE1 to their respective AD, and found that only

the latter resulted in a significant enhancement in the activation potential (Fig. 3D). Next, we

fused CgE1 or CgE2 in both orientations to the minimal AD of ScPho4 (ScAD_9aa). We found

that indeed, both regions were able to significantly enhance the activity of the ScAD_9aa in an

orientation-independent manner (Fig. 3D). It is worth noting that this enhancement activity does

not involve the endogenous ScPho4 or ScPho2. ScPho4 is inactive in the phosphate replete

condition in which the Y1H experiment was conducted. To test the involvement of ScPho2, we

repeated the assay in a host lacking *pho2*. No significant difference in the boosting effects of

either CgE1 or CgE2 was observed, although CgE1's effect is slightly lower without Pho2

(Supplementary Fig. 6).

In summary, we found that CgPho4 encodes two Activation Enhancer Domains (AEDs),

both of which are in the Intrinsically Disordered Region (IDR). The AEDs have little activation

capability on their own but can significantly enhance the activation potential of the AD of either

of the Pho4 orthologs. Since these two AEDs are present in CgPho4 but not in ScPho4, we

hypothesize that they contribute to CgPho4 being less dependent on Pho2.

A dual fluorescence reporter assay accurately measures the activity of Pho4 chimeras

with and without Pho2 for dissecting the divergence in co-TF dependence

So far, we compared DNA binding and activation potential between CgPho4 and ScPho4 by

isolating individual regions and testing them either *in vitro* or in a synthetic *in vivo* system. While

they support the enhanced activity model, it remains unclear whether and how they contribute to

divergence in Pho2-dependence in the native Pho4 protein context. They also do not test or rule

out the autoinhibition model, which necessitates intramolecular interactions. To answer these

questions, we divided ScPho4 and CgPho4 into five corresponding parts (Fig. 1C, 4A), with

breakpoints chosen to be on the edge of well-aligned regions to avoid breaking known or

predicted functional domains and secondary structures. Boundaries for all regions used in this

and other experiments can be found in Table 1. We created all 2^5=32 combinations as well as

28 additional ones with alternative breakpoints (Fig. 4A). All constructs were C-terminally tagged

with mNeon to quantify the protein levels, and were expressed from the native *ScPHO4*

promoter and UTRs (Fig. 4B). We also created two *S. cerevisiae* host strains, in which the

ScPHO5 CDS was replaced by an mCherry reporter, and *pho80* was knocked out so that all

Pho4 chimeras were constitutively nuclear localized [24,27]. One of the two host strains had *pho2*

knocked out. For each chimera, we measured its *PHO5pr*-mCherry and Pho4-mNeon levels

using flow cytometry in both hosts. While mCherry and mNeon levels for the same Pho4

construct varied between experiments, the ratios between the two were consistent and

characteristic of the specific construct (Fig. 4C, D). We therefore defined the activity of a Pho4

chimera as the ratio between the median fluorescence intensity (MFI) of mCherry and mNeon,

which we will refer to as A_{PHO2} or $A_{pho2\Delta}$ from hereon.

The two Activation Enhancer Domains (AEDs) in CgPho4 increased the activity of the

chimeric TFs but were insufficient to make them Pho2-independent

The chimeric Pho4 constructs exhibited varied A_{PHO2} and $A_{pho2\Delta}$ values (Fig. 5A). Interestingly,

some showed higher A_{PHO2} than either ScPho4 or CgPho4. To identify the potential genetic basis

for the divergence in Pho2-dependence, we first calculated the activity difference when one or

two regions of ScPho4 were replaced with their counterpart(s) from CgPho4. The results were

plotted as two heat maps separated based on the presence of Pho2 (Fig. 5B). For example, a

chimera with its NLS from CgPho4 (NLS:Cg) and the rest from ScPho4 led to estimates of the

activity difference between NLS:Cg and NLS:Sc on ScPho4 background with or without Pho2.

Based on our previous results, we expected R1, NLS, and DBD in CgPho4 to enhance

the activity of the chimera with and without Pho2. In particular, the first two encode CgE1 and a

majority of CgE2_9aa, which we showed enhanced the activity of the main AD (Fig. 3); DBD:Cg

is expected to increase the binding ability of the chimera (Fig. 2). We indeed found R1:Cg and

NLS:Cg to enhance A_{PHO2} on ScPho4's background (Fig. 5C, "R1, AD, NLS" group); contrary to

our expectation, however, DBD:Cg reduced the chimera's activity with Pho2 (Fig. 5C, bHLH).

Also unexpectedly, R1:Cg and NLS:Cg had a much smaller effect on $A_{pho2\Delta}$ (Fig. 5B bottom, 5C

right), suggesting that despite their ability to increase A_{PHO2}, the chimeras were still dependent

on Pho2. In fact, none of the 1- or 2-region swaps led to large increases in $A_{pho2\Delta}$. It is worth

noting that this result also argues against a classic lock-and-key model for autoinhibition, which

we expect to be broken by swapping one of the regions involved.

To quantitatively examine the contribution of R1, AD, and NLS regions from CgPho4 and

whether they interact non-additively (epistasis), we fit a linear model to the data to estimate the

main and interaction terms for the regions, i.e., $Y = X_0 + R1 + AD + NLS + R1{:}AD + R1{:}NLS +$

AD:NLS + R1:AD:NLS. In this model, the first term represents the activity of ScPho4, the next

three terms the main effect of each CgPho4 region (":Cg" omitted for brevity), and the rest are

interaction terms. We found that R1:Cg and NLS:Cg each had significant, positive effects on

their own on both A_{PHO2} and $A_{pho2\Delta}$; although the magnitude was much smaller for $A_{pho2\Delta}$ (Fig. 5D,

Holm-Bonferroni corrected $P < 0.05$). Both also had a significant and positive interaction term

with AD:Cg on A_{PHO2} (corrected $P < 0.05$ for R1:AD and AD:NLS); the estimates trended in the

same direction but were not significant for $A_{pho2\Delta}$ (Fig. 5D). One explanation for the observed

epistasis may be that the two CgPho4 AEDs work more efficiently with AD:Cg than with AD:Sc.

However, it could also be explained by the disruption of the native conformation and function of

the interacting regions resulting in a lower activity in the species-mixed constructs. Note that to

minimize such effects, we designed the breakpoints to avoid any predicted secondary structures

(Supplementary Fig. 1, Supplementary Fig. 2).

Since the NLS:Cg region encodes both the NLS/NES and a key part of the second AED

(CgE2_9aa), we asked which of these two mechanisms was responsible for its positive effect on

A_{PHO2}. To test the possibility that the enhanced A_{PHO2} was due to a stronger nuclear localization

activity of NLS:Cg, we performed fluorescence microscopy to quantify the concentration of the

nuclear-localized Pho4 proteins and the ratio of nuclear vs total Pho4 proteins in six constructs

that bear either NLS:Sc or NLS:Cg. No significant difference between the two groups was found

(Supplementary Fig. 7, F-test $P > 0.1$). We thus conclude that the positive effect of NLS from

CgPho4 on A_{PHO2} was mainly due to its ability to boost activation.

We also examined the effect of swapping CgPho4's DBD. On its own, swapping DBD:Cg

decreased both A_{PHO2} and $A_{pho2\Delta}$ compared to ScPho4 (Fig. 5B), suggesting that DBD:Cg may

be incompatible with one or multiple regions in ScPho4.

In summary, we found that R1:Cg and NLS:Cg increased the activity of the chimera with

Pho2 but were insufficient to remove the Pho2 dependence alone or in combination. In the

presence of Pho2, R1:Cg and NLS:Cg both showed positive epistasis with AD:Cg on ScPho4's

background. Lastly, no 1- or 2-region swaps from CgPho4 into ScPho4 increased $A_{pho2\Delta}$ to the

level of CgPho4. Together, we conclude that CgPho4's two AEDs can increase the activity of

the chimera in the presence of Pho2, although they fail to restore the activity without Pho2 to

CgPho4's level, suggesting that additional mechanisms are at play. CgPho4's DBD, alone or in

combination with R1 and the NLS region, did not contribute to increased activity.

A double-edged sword: Pho2 Interaction Domain (P2ID) in ScPho4 allows for enhanced

activity with Pho2 but restricts its activity without Pho2

In the 1- and 2- region swap heatmap (Fig. 5B), the P2ID showed a puzzling pattern: swapping

CgPho4's P2ID (P2ID:Cg) into the ScPho4 background had a dominant negative effect on A_{PHO2}

while offering little increase in $A_{pho2\Delta}$. This suggests that P2ID:Sc is essential for ScPho4's

function. Conversely, swapping ScPho4's P2ID (P2ID:Sc) into CgPho4 increased A_{PHO2} beyond

that of CgPho4 (18 to 22) but reduced $A_{pho2\Delta}$ (17 to 4.8). This suggests that P2ID:Sc is a key

factor to Pho2-dependence. To investigate the unique property of the P2ID further, we plotted

all chimeras based on their A_{PHO2} and $A_{pho2\Delta}$ values (Fig. 6A). Strikingly, the chimeras fell into

three distinct groups based on the identity of their P2IDs. The first group had P2ID from

CgPho4, and their activities were not dependent on Pho2, falling along the diagonal line. The

second group had P2ID from ScPho4. These had strong activity with Pho2 and low to no activity

without Pho2, like ScPho4 does. The third group included additional chimeras with mixed P2ID

regions, and they filled the intermediate space between the first two groups. Notably, many

chimeras in the second group showed higher A_{PHO2} than ScPho4. Most of them contained the

R1 and NLS regions from CgPho4, consistent with our results above showing that these two

regions enhanced the activity of the Pho4 chimeras in the presence of Pho2 (Fig. 5A, B).

The evidence above suggests that P2ID:Sc has two effects: allowing for collaborative

regulation with Pho2 and restricting Pho4's activity without Pho2. Can these two functions be

separated? To answer this question, we compared the mixed P2ID chimeras to those with

whole P2IDs from either Pho4. We found that swapping P2ID:Sc into CgPho4 resulted in a 1.2-

fold increase in A_{PHO2} and a 71% reduction in $A_{pho2\Delta}$ (Fig. 6B, row 2 vs 3). Interestingly,

swapping just the second half of P2ID:Sc (aa 205-242) into CgPho4 resulted in a ~40%

reduction in A_{PHO2} and 75% reduction in $A_{pho2\Delta}$. By contrast, just swapping the first half of

P2ID:Sc (aa 177-204) resulted in 1.76-fold increase in A_{PHO2} and maintained the same level of

$A_{pho2\Delta}$ (1.1-fold). From this, we deduced that the first half of P2ID:Sc mainly functions to interact

with Pho2 while the second half appears to limit Pho4's activity without Pho2. The same trend

was observed in another set of chimeras with lower activities (Fig. 6B, rows 6-9).

Several chimeras had very low activities even with Pho2 (A_{PHo2} < 3.6, or 20% of A_{PHO2} for

ScPho4). Most of these nonfunctional chimeras have P2ID:Cg and DBD:Sc (Fig. 6C, rows 3-7).

We hypothesize that DBD:Sc requires P2ID:Sc and Pho2 to function in the context of the full

length Pho4. This is contrary to the conventional view that DBDs can function on their own,

which is supported by our own *in vitro* results (Fig. 2). Nonetheless, we reasoned that if the

above hypothesis is correct, putting P2ID:Sc back by inserting it in between P2ID:Cg and

DBD:Sc should rescue the non-functional chimeras in the presence of Pho2. That is what we

observed (Fig. 6C, rows 8-12), with some of the chimeras even exceeding the A_{PHO2} of ScPho4

and CgPho4. However, all of them still required Pho2. These results support the above

hypothesis, showing that the dual-functional P2ID is essential for ScPho4 to function.

Given that chimeras with P2ID:Cg have equal activities with and without Pho2 (Fig. 6A), we

asked if CgPho4 still physically interacts with Pho2 in *S. cerevisiae*. Using the yeast two-hybrid

assay, we were unable to detect an interaction between CgPho4ΔDBD (aa 2-463) and a region

of ScPho2 known to interact with multiple TFs, including ScPho4 [32] (Fig. 6D). By contrast,

ScPho4ΔDBD (aa 2-250) was able to interact with the same region of ScPho2 as previously

found (Fig. 6D). This is consistent with our observation above, where chimeras with P2ID:Cg

and lacking the region from ScPho4 mediating Pho2-interaction cannot be enhanced by Pho2

(Fig. 6A).

In summary, our chimera dissection revealed three key regions behind the difference in

co-TF dependence. Notably, all three are IDRs. Among them, a region adjacent to the DBD in

ScPho4 (P2ID:Sc) functions as a double-edged sword: it both allows Pho4 to gain activity with

Pho2's help and restricts it when Pho2 is absent. We showed that these two functions are

encoded by physically separate parts, offering a path for determining their respective

mechanisms of actions. CgPho4 lacks the ability to interact with and use Pho2's help. Instead,

the two AEDs we identified earlier – both in IDR – conferred higher activity independent of

Pho2. This, combined with the lack of the autoinhibition by P2ID:Sc, makes CgPho4 as active

as ScPho4 and not dependent on the co-TF.

328   **DISCUSSION**

Combinatorial control plays crucial roles in eukaryotic gene regulation. Mutations disrupting TF

interactions can cause dysregulation and disease [10]. However, how mutations can alter TF

interdependence itself, whether in disease or evolution, is less understood. In this study, we

investigated the molecular basis for natural variation in co-TF dependence in the yeast

phosphate starvation (PHO) response. We found three key differences between two

orthologous Pho4s with varying dependence on the co-TF Pho2 (Fig. 7): **(1)** <u>DNA Binding</u>

<u>Affinity</u>: CgPho4 binds the same consensus DNA motif with 3-4-fold higher affinity than ScPho4;

(2) <u>Activation potential</u>**:** CgPho4 has two unique activation enhancing domains (AEDs) that

increase the activation potential of both Pho4s. **(3)** <u>Autoinhibition</u>: ScPho4 contains an IDR next

to its DBD that both allows it to interact with Pho2 to gain enhanced activity, and inhibits its

activity in the absence of Pho2. Therefore, our results support both the enhanced activity model

and the autoinhibition model for the difference in Pho2-dependence between Pho4 orthologs.

Among the three differences, the contribution of CgPho4's two AEDs to the TF's activity

and dependence on Pho2 is well supported by the yeast one-hybrid data and chimeric Pho4

results (Fig. 3, 5). We are not aware of existing examples or proposed mechanisms for such a

phenomenon. One hypothesis for how AEDs work is that they are weaker ADs with the same

biochemical activities, i.e., recruiting cofactors through protein-protein interactions. As such,

they are not sufficient for activation by themselves but can increase the activity of the nearby AD

(the effect can be non-additive). Alternatively, AEDs may affect the conformation of the AD and

have no activity on their own. Further tests by synthetic constructs and biochemical assays,

such as co-IP followed by mass-spec, will provide mechanistic insight into how AEDs function.

By contrast, the significance of the binding affinity differences between the two Pho4s

remains unclear (Fig. 2E). Binding kinetics predicts that if the nuclear concentrations of both

Pho4 are much higher than their K_D, a 3-4-fold difference in affinity would have little impact.

Conversely, if the nuclear concentration of Pho4 is close to its K_D, the same difference could

significantly affect gene induction. Existence of the second scenario is supported by a study

showing that nuclear ScPho4 levels are much lower at intermediate phosphate concentrations

than in extreme starvation conditions, leading to differential binding and induction of ScPho4's

targets [33]. In our chimeric Pho4 experiments, replacing ScPho4's DBD with that from CgPho4

reduced Pho4's activity rather than increasing it both with and without Pho2 (Fig. 5). It seems to

suggest that the binding affinity difference has no functional impact *in vivo*. However, it is worth

noting that we assayed the activity of chimeric Pho4s in the *pho80Δ* background where all Pho4

proteins are constitutively inside the nucleus at near-maximal levels, hence not allowing us to

test the effect of the affinity difference at a lower level of nuclear Pho4. Future studies will need

to measure Pho4 activities at varying nuclear concentrations to test the above hypothesis.

Surprisingly, we found that ScPho4's DBD requires both ScPho4's P2ID and Pho2 for

full activity (Fig. 6C), challenging the view that TFs are modular, where DBD are expected to

function independently. Although ours and others' studies confirmed that ScPho4 DBD can bind

DNA *in vitro* (Fig. 2) and even *in vivo* when overexpressed on its own [34], they may not fully

reflect its physiological activity, which depends on the nuclear concentration, nucleosomal

context, and interactions with other TF regions and cofactors. Further experiments are needed

to resolve the paradox and re-examine the assumption of TF modularity.

Only one of the three functional differences identified between the two Pho4 orthologs is

in a structured region. The two AEDs and the dual-functional P2ID:Sc are both within IDRs.

IDRs are abundant in eukaryotic TFs, with over 80% having at least one such region [35].

Approximately 75% of ScPho4 and CgPho4 were predicted to be IDR (Supplementary Fig. 1).

While most studies of TF function and evolution focus on the structured regions like the DBD,

our work joins a small number of studies highlighting the significance of IDR divergence in TF

evolution [36,37]. On one hand, IDRs play crucial roles in nearly all aspects of a TF's function,

including recruiting cofactors, interacting with co-TFs, contributing to binding specificity and

forming molecular condensates [38–43]. On the other hand, IDRs evolve much faster than

structured regions, potentially offering more raw variation for natural selection to act on. Our

current understanding of how IDR evolves and affects TF function is limited by challenges in

aligning their sequence and studying their functions. Progress in experimental techniques like

phase separation [43] and large language models trained on protein sequences offer new

opportunities to investigate TF IDR's function and evolution [44].

Our study focused on dissecting the genetic basis for Pho2-dependence variation in two

Pho4 orthologs. We previously showed that this trait varies across Pho4 orthologs, with reduced

Pho2-dependence potentially evolving independently in more than one lineage [15]. We wonder

whether the IDR-associated divergence identified above also correlated with the level of Pho2-

dependence more broadly. Preliminary analyses of the activation potential and P2ID length

across eight Pho4 orthologs with different Pho2-dependence levels supported this speculation

(Supplementary Fig. 8).

Lastly, our study illustrates a mechanism for the evolution of combinatorial control in

eukaryotes. Unlike previous research that focused on the gain and loss of TF interactions in

gene regulatory networks [45,46], we show that TF's dependence on co-TF itself can evolve. This

divergence can lead to significant rewiring of the regulatory network, as seen with Pho4, where

reduced co-TF dependence correlates with an expanded target gene network [15]. This contrasts

with the dominant pattern seen in the literature on combinatorial control evolution: the Johnson

lab, for example, showed that changes in TF combinations in the yeast mating type pathway

altered the mode of regulation but maintained the overall network output [11,12,47,48]. While cases

of TF interdependence evolution are still rare, it is interesting to compare our example with

Gain-of-Function mutations in kinases: in the proto-oncogene ABL, a fusion with another gene

called BCL makes the merged protein independent of the upstream and downstream activators,

which drives excessive cell proliferation and leads to Chronic Myeloid Leukemia [49]. Whether

similar changes in TF dependence on co-TFs lead to misregulation and in turn causes diseases

or novel phenotypes is an interesting question for future studies.

In summary, our study provides a detailed molecular picture of how co-TF dependence

is mediated and how it evolves, particularly through IDR changes. Further exploration of both

questions is essential for understanding gene regulation and regulatory evolution in eukaryotes.

METHODS

Breakpoints used for the chimeric Pho4 constructs and individual regions are listed in Table 1.

Plasmids and strains are listed in Supplementary Data 1 and Table 2. Computational and

statistical analysis scripts performed in this study are available at https://github.com/binhe-

lab/E013-Pho4-evolution, which will be archived using Zenodo and minted with a DOI at the

time of publication.

Bioinformatic analyses of Pho4 orthologs

Pho4 ortholog sequences were from the Yeast Gene Order Browser (http://ygob.ucd.ie/), and

were aligned using ProbCons [50] via JalView's Web Service [51,52]. The alignment was manually

edited to align the five Pho80/85 motifs [24]. Secondary structures for ScPho4 and CgPho4 were

predicted using the PSIPRED 4.0 [53]. 9aaTAD motifs were predicted using the webapp

https://www.med.muni.cz/9aaTAD/ [31]. For ScPho4, one match was found using the moderately

stringent pattern, located at aa 75-83; for CgPho4, one match was identified using the most

stringent pattern, at aa 270-278, with three more using the moderately stringent patterns (aa 20-

28, 24-32, 282-290). Among the latter three, aa 24-32 had a lower % match at 67% vs 83% for

the others.

Identifying Pho4 binding motifs from ChIP-seq data

Chromatin Immunoprecipitation (ChIP-seq) for ScPho4 and CgPho4 were previously performed

in *S. cerevisiae*, with both Pho4 expressed from the same endogenous ScPho4 locus with

native regulatory sequences [15]. ChIP identified peaks for both Pho4s were downloaded from the

supplementary files of the above publication. The sequences under each peak were extracted

from *S. cerevisiae* genome (sacCer3, NCBI refseq assembly GCF_000146045.2), which were

submitted to the peak-motifs tool without control sequences on the RSAT Fungi server

(https://rsat.france-bioinformatique.fr/fungi/) [54]. The top motif was reported for each Pho4

ortholog.

Protein expression and purification for DNA binding assays

All recombinant proteins were expressed using pET-11a (Sigma #69436-3) based vectors in

BL21(DE3) *E. coli* cells. Vector maps are available upon request. The DNA binding domain

(DBD) constructs included ScPho4 DBD-6xHIS (aa 236-312) and CgPho4 DBD-6xHIS (aa 452-

533) cloned downstream of the T7 promoter in pET-11a. The transformed bacterial strains were

grown overnight in LB + 100 ug/mL ampicillin. The overnight culture was diluted to OD 0.1 and

the cells were grown to OD 0.6 and induced with 1mM IPTG for 2 hours. Induced cells were

collected by centrifugation at 6000 rpm at 4ºC for 35 minutes, and were snap frozen in liquid

nitrogen, then stored at -80ºC until purification. Our preliminary experiments showed that the

protein of interest was largely in the insoluble fraction. Therefore, we performed a refold protocol

on the Ni-NTA column. Briefly, frozen pellets were resuspended in 1xPBS, pH7.4 with cOmplete

EDTA-free protease inhibitor (Sigma #11836170001) and sonicated for 45 cycles of 1 sec on / 2

sec off repeated 3 times at 50% power to lyse the cells. Sonicated samples were centrifuged at

35k rpm for 30 minutes. The supernatant was discarded, and the pellet was resuspended in the

solubilization buffer (20 mM TrisBase, 0.5 M NaCl, 5 mM imidazole, 5.5 M Guanidine

Hydrochloride, 1 mM 2-mercaptoethanol pH 8.0) and stirred at ~25°C for an hour. The

solubilized pellet was centrifuged at 35k rpm for 30 minutes and the supernatant was filtered

and loaded onto a 5mL Ni-NTA column. The column was washed with a urea buffer (20 mM

TrisBase, 0.5 M NaCl, 20 mM imidazole, 5.5 M Urea, 1 mM 2-mercaptoethanol pH 8.0). Then,

the protein was refolded with a reverse urea gradient from buffer A (20 mM TrisBase, 0.5 M

NaCl, 20 mM imidazole, 5.5 M Urea, 1 mM 2-mercaptoethanol pH 8.0) to buffer B (20 mM

TrisBase, 0.5 M NaCl, 20 mM imidazole, 1 mM 2-mercaptoethanol pH 8.0) on a BioRad FPLC.

The refolded protein was eluted off of the Ni-NTA column using a gradient from buffer C (20 mM

TrisBase, 0.5 M NaCl, 20 mM imidazole, 1 mM 2-mercaptoethanol pH 8.0) to buffer D (20 mM

TrisBase, 0.5 M NaCl, 500 mM imidazole, 1 mM 2-mercaptoethanol pH 8.0). Fractions

containing the protein of interest were identified by gel electrophoresis, pooled, and diluted with

a no salt buffer (25 mM Na_2HPO_4 pH 7.0, 0.5 mM THP) and run on a Heparin column

equilibrated with low salt buffer (25 mM Na_2HPO_4 pH 7.0, 0.15 M NaCl, 0.5 mM THP). Protein

was eluted with a gradient from low to high salt buffer (25 mM Na_2HPO_4 pH 7.0, 1.5 M NaCl, 0.5

mM THP). Protein-containing fractions were concentrated using a 3 kDa cutoff Amicon

centrifugal filter (Sigma UFC8003) and loaded onto a Superdex 75 size exclusion column

equilibrated with a storage buffer (25 mM Na_2HPO_4 pH 7.0, 0.5 M NaCl, 0.5 mM THP).

Fractions containing the expected size products were collected, analyzed by gel electrophoresis,

and stored in the storage buffer at 4°C.

N-GST-CgPho4 full length was constructed for the Protein Binding Microarray assay.

The corresponding N-GST-ScPho4 purification has been described before [55]. Both proteins

were grown and lysed as described in [55]. Briefly, BL21 *E. coli* cells containing the constructs

were induced at OD600 0.8 with 1mM IPTG and collected by centrifugation after 3 hours of

induction at 30ºC. Pellets were snap frozen in liquid nitrogen and stored at -80ºC until

purification. Cells were lysed with rLysozyme (Millipore 71110) for 20 minutes at room temp in

the lysis buffer (1xPBS, pH7.4) with the cOmplete protease inhibitor tablet and 1 mM PMSF.

The protein was run on a GST column equilibrated with the lysis buffer and eluted with a

gradient to buffer B (50mM Tris, 10mM glutathione, pH 8). GST-CgPho4 fractions containing the

protein were pooled and loaded onto a heparin column equilibrated with low salt buffer (25mM

Na_2HPO_4 pH 7.4, 150mM NaCl, 0.5mM THP) and eluted with a gradient to high salt buffer

(25mM sodium phosphate dibasic pH 7.4, 1.5M NaCl, 0.5mM THP). Fractions containing the

protein were pooled and concentrated with a 30 kDa cutoff Amicon centrifugal filter (Sigma

UFC9030) and loaded onto a Superdex 200 size exclusion column equilibrated with the storage

buffer (25mM HEPES pH 7.4, 500mM NaCl, 0.5mM THP). Protein was run on an SDS-PAGE

and pure fractions were pooled and concentrated. 10% glycerol was added before snap freezing

and storage at -80ºC.

Universal Protein Binding Microarray (uPBM)

The uPBM was performed following the PBM protocol as described in [56]. Briefly, after the primer

extension step is used to double-strand the DNA molecules on the array, the chambers are

blocked with 2% milk. After washing, proteins are incubated with the array for 1 hour. Alexa

Fluor 488-conjugated anti-GST antibody (Invitrogen A-11131) was used to detect binding. The

array was scanned using a GenePix 4400A scanner (Molecular Devices). GST-ScPho4 and

GST-CgPho4 (full length) were prepared as described above. Both proteins were assayed at a

final concentration of 1 uM as determined by the optical absorbance. An 8x15k array was used

to assay all possible 9-mers, from which a robust 7-mer enrichment score is derived. The non-

parametric enrichment score, or E-score, is invariant to differences in the concentration of the

proteins used in the assay, and thus are suitable for comparisons of relative affinities between

arrays. E-score ranges from -0.5 (lowest enrichment) to +0.5 (highest enrichment). Scores

greater than 0.35 correspond to specific TF-DNA binding [29,57]. Data analysis was performed

using custom Perl scripts as described in [56] to extract and normalize fluorescence based

intensity.

Genomic Context Protein Binding Microarray (gcPBM)

All probes in the DNA library are 60 bp in length with 24 bp complementary to the primer used

for double stranding and 36 bp of genomic region centered on the E-box motif or its variant. The

library contains 1) 5711 *S. cerevisiae* genomic regions with putative Pho4 binding sites; 2) 150

negative controls from the *S. cerevisiae* genome not specifically bound by Pho4 [55]; 3) 4000 DNA

sequences used in previous MITOMI experiments to calibrate the binding affinities [58]; 4) 150

genomic regions from *C. glabrata* that contain Pho4 consensus and nonconsensus binding sites

[15]; 5) 100 probes from the library of sequences we tested with Biolayer Interferometry; and 6)

150 negative controls from the *C. glabrata* genome not bound by CgPho4. We used the

NNNNGTG, CACNNN, and GTGNNN libraries from Maerkl and Quake (2007) in our gcPBM

design. The MITOMI and BLI probes required the addition of random flanks to maintain the

36bp length;10 different flanking sequences were generated for each sequence. Each probe is

represented by six replicates, including three replicates in each orientation and were distributed

randomly across the array. The custom 8x60k (8 chambers, 60,000 DNA spots per chamber)

was synthesized by Agilent Technologies. The gcPBM was performed and analyzed following

the PBM protocol as described above and in [29,55]. The log transformed median intensity of the 6

replicate probes were used for comparisons. Because the log signal intensities are not directly

comparable between the two Pho4 proteins, which were assayed on separate arrays, we used

Spearman's rank correlation coefficient to quantify the level of concordance in their sequence

preference. Any difference in preference should result in a change in the ranks within each

Pho4's data.

Biolayer interferometry (BLI)

A library of 17-bp dsDNA was used for the assay. The consensus probe had the sequence

"CTAGTCC**CACGTG**TGAG", with the E-box motif bolded, and was identical to the DNA used in

the crystal structure of ScPho4's DBD [26]. Nine half-site variants, including "AACGTG,

TACGTG, … CAAGTG, ACGGTG, CATGTG" were constructed on this background. For each

probe, the complementary ssDNA oligos were synthesized by Integrated DNA Technologies

(IDT). One of the two probes was biotinylated on the 5' end. To anneal them into dsDNA, 1 pmol

(5 uL of 200 uM) of the biotinylated oligo was mixed with 2 pmol (10 uL of 200 uM) of the

complementary, unmodified strand in the Nuclease-Free Duplex Buffer (30 mM HEPES, pH 7.5;

100 mM potassium acetate). The mixture was heated to 95ºC and then left at room temperature

for it to cool down. Annealed probes were stored at -20ºC until use.

ScPho4 DBD and CgPho4 DBD were purified and stored as described above. Protein

quality was checked weekly for signs of degradation using Dynamic Light Scattering (DLS) on a

DynaPro NanoStar instrument. Before each experiment, the concentration of the protein prep

was measured in triplicates using a NanoDrop instrument. The mean concentration was used to

prepare the protein dilutions and calculate K_D.

BLI experiments were performed on an Octet RED 96 instrument at 30ºC with 1000 rpm

shaking. For each protein-probe pair, eight streptavidin (SA) biosensors were hydrated for 15-20

min at room temperature in the 1x kinetics buffer (1x PBS pH 7.4, 0.01% BSA, 0.002% Tween-

20, 0.005% Sodium azide). A black 96 well flat-bottom plate was loaded with experimental

components also diluted in the 1x kinetics buffer. To begin the experiment, biosensors were

equilibrated in a 1x kinetics buffer for 60 seconds to reach the baseline. Seven biosensors were

then submerged in 35 nM biotinylated annealed DNA for 30 seconds, while one biosensor was

submerged in the 1x kinetics buffer as a no-DNA control. Biosensors were then submerged in 1

ug/ml biocytin for 60 seconds to block any empty streptavidin pocket on their surface, before

being dipped back into a 1x kinetics buffer for 60 seconds for a baseline measurement. Loaded

and blocked biosensors were then submerged into a gradient of protein concentrations

calculated for each probe based on the K_D. To obtain reliable measurements, we use a

concentration range that spans 10x to 1/10 of the K_D values. Biosensors stayed in the protein

solution for 900-1000 seconds or until equilibrium was reached.

Data analysis was performed in the ForteBio Data analysis v11 software. After

subtracting the background and aligning the y-axis, the processed data were subjected to either

a steady state analysis or a kinetic curve fitting. For steady state analysis, the equilibrium-level

signal from each biosensor was plotted against the protein concentration, from which K_D was

calculated. Kinetic curve fitting was done using a one-site specific binding with Hill slope model

as implemented in the Data analysis v11 software. The latter more effectively fit the BLI data

and thus were utilized for most of the analysis. For the consensus sequence, we found the

kinetic curve-based estimates for both ScPho4 and CgPho4 to have higher variance, likely due

to the fast kinetics not adequately captured by the model [59]. Steady-state analysis gave

consistent mean K_D as the kinetic analysis did, but resulted in lower variance, and was used

instead. Two to four replicates were performed for each 17bp DNA library sequence, using at

least two independent protein preps.

Electrophoretic Mobility Shift Assay (EMSA)

IR700 labeled 17bp consensus DNA (same as the consensus DNA for BLI above, Integrated

DNA technologies) was diluted to 0.1 nM and mixed 1:1 with a 2x dilution series of the DBD of

interest starting at 32nM in binding buffer (20mM Tris-HCL, pH 8, 150mM KCl, 10% glycerol,

5mM MgCl2, 1mM EDTA, and 1mM DTT). The mixture was incubated for 1 hour at 4ºC. A 10%

native PAGE gel in 1x Tris Glycine buffer (2.5 mM Tris pH 8.3, 19.2 mM glycine) was prerun at

200V for 20 min. The DNA/protein mixture was then loaded onto the gel and ran for 25-35

minutes. The gel was imaged using the Odyssey FC imager in the 700 nm channel. To estimate

K_D, the unbound band in each protein-containing lane was quantified using the ImageStudio

software (with background subtraction) and divided by the no protein control. This value was

then subtracted from 1 and plotted against the protein concentrations. A nonlinear curve fitting

was performed in Prism v10.2.1 using the one site specific binding model. K_D estimates and the

standard deviations were reported.

Yeast media and growth

Yeast cells were grown in Yeast extract-Peptone Dextrose (YPD) medium or Synthetic

Complete (SC) medium, using Yeast Nitrogen Base without amino acids (Sigma Y0626)

supplemented with 2% glucose and amino acid mix. Phosphate starvation medium was made

using Yeast Nitrogen Base with ammonium sulfate, without phosphates, without sodium chloride

(MP Biomedicals, 114027812) and supplemented to a final concentration of 2% glucose, 1.5

mg/ml potassium chloride, 0.1 mg/ml sodium chloride and amino acids, as described previously

[15]. Phosphate concentration in the medium was measured using a Malachite Green Phosphate

Assay kit (Sigma, MAK307).

Yeast strain and plasmid construction

The hosts containing the endogenous *PHO5*pr-mCherry reporter were constructed by replacing

the endogenous coding sequence with mCherry using CRISPR/Cas9. Briefly, guide RNAs

targeting *PHO5* were designed in Benchling (https://www.benchling.com) and cloned into

bRA89 (AddGene 100950), a plasmid that contains the Cas9 protein and gRNA scaffold [60].

Homology arms to the *PHO5* 5'UTR and 3' UTR were added onto an mCherry donor DNA using

PCR. The donor DNA and plasmid were co-transformed into the host using the standard LiAc

transformation protocol. Transformants were selected for the CRISPR plasmid using

hygromycin resistance and screened with PCR. Positive clones were validated using Sanger

sequencing and fluorescence microscopy. Pho2 was then knocked out using a *HIS3* cassette

with homology arms to the *PHO2* 5'UTR and 3'UTR.

The chimeric Pho4 plasmid library was constructed using Golden Gate. The fragments

of CgPho4 and ScPho4 were PCR amplified using Phusion Flash polymerase (Thermo

Scientific F548S) with unique overhangs for Golden Gate. They were then assembled and

inserted into a pRS315-based backbone that contains the Sc*PHO4* promoter, a C-terminal in-

frame mNeon tag followed by the Sc*PHO4* 3' UTR and terminator. Primers were designed using

the NEBridge Golden Gate assembly tool. The inserts were verified with PCR and Sanger

sequencing and then transformed into the yeast hosts using either the standard LiAc protocol [61]

or the Zymo yeast transformation kit (Zymo research T2001) and plated onto SD-leu media.

Flow cytometry

Cells were inoculated into a 96 deep-well plate (Fisher Scientific 07-200-700) and grown

overnight in SD-Leu or SC medium supplemented to final concentrations of 0.13 mg/ml adenine

and 0.1 mg/ml tryptophan to reduce autofluorescence [62]. Cultures were diluted in the morning to

an OD600 of 0.15 and grown to an OD600 of 0.6. These cells were directly subjected to flow

cytometry at a flow rate of 25 uL/min on an Attune NxT flow cytometer (ThermoFisher) fitted

with an autosampler. Data were collected using the Attune NxT software v3.1 for FSC, SSC and

the appropriate fluorescence channels. Calibration beads (Spherotech RCP305A) were run

routinely to ensure that experiments from different days were comparable. Pho4-mNeon was

measured in the BL1 channel with 488 nm excitation and 510±10 nm emission; mCherry was

measured in the YL2 channel using 561 nm excitation and 615±25 nm emission. Voltages for

each channel were set by the brightest sample and negative control so the sample signals were

between 10^2-10^5. Events were gated based on FSC-H / SSC-H to remove non-cells, then on the

FSC-W / FSC-H to isolate singlets (Supplementary Fig. 9). At least 10,000 gated events were

collected per sample. Each strain was measured at least three times, and on two different dates.

A nonfluorescent strain was included in every experiment for subtracting the autofluorescence.

Further gating and analyses were performed in R using the FlowClust and FlowCore packages.

Detailed analysis scripts are available in the project github repository.

Fluorescent microscopy for Pho4 nuclear localization

Six Pho4 constructs were chosen, with three bearing NLS:Sc (SSSSS, CCSCC, SSSCC) and

three bearing NLS:Cg (SSCSS, CCCCC, SSCCS). They were transformed into either a *PHO2*

wild type or *pho2Δ* background. Yeast cells were grown to mid-log phase in SD-Leu and fixed

using 4% paraformaldehyde for 10 minutes, washed twice with 1xPBS before DAPI staining.

4',6-diamidino-2-phenylindole (DAPI, Sigma, D9542) was diluted in the respective medium and

added to the fixed cells at a final concentration of 10 ug/mL. Cells were incubated with DAPI in

the dark for 30 minutes before DAPI was removed and cells were washed twice with 1xPBS.

Fixed cells were mounted on a 1.2% agarose pad on a depression slide for image. Fluorescent

imaging was performed on a Leica epifluorescence microscope. A 405 nm laser was used for

DAPI excitation and 430-550 nm for emission. For Pho4-mNeON, 488nm was used for

excitation and 530 nm for emission. A bright field image was also taken to show the cell

boundaries. A total of three images were recorded for each field of view.

Microscopy analysis was performed in ImageJ (v 1.53). To quantify the nuclear fraction

of Pho4 proteins, ten cells were randomly selected for each construct in each host background,

and the experiment was repeated two times, resulting in a total of 15 or 20 cells quantified in

PHO2 and *pho2Δ* backgrounds, respectively. Cell boundaries were manually traced using the

bright field channel, and nucleus using the DAPI channel. Both were added to the ROI manager,

and the raw integrated density (sum of pixel values) was quantified for both areas. The ratio of

nuclear vs whole cell integrated density was used for plotting and statistical analysis.

Yeast one-hybrid and yeast two-hybrid

Indicated regions from CgPho4, ScPho4, and ScPho2 were PCR amplified and cloned into

pGBD-C3 [63] containing the Gal4 DBD using Gibson Assembly. Plasmids were verified with PCR

and Sanger sequencing. The *GAL1*pr-mCherry reporter was created using a pRS306 based

integrative plasmid digested with StuI and inserted into the *ura3* locus of a *gal4Δ gal80Δ S.*

cerevisiae strain. The reporter was tested using flow cytometry according to the protocol above

using only the red (YL2) channel. Plasmids were then transformed into this yeast strain using

the standard LiAc method [61] or the Zymo yeast transformation kit (Zymo research T2001) and

selected for on SD-trp. For the yeast two-hybrid, the Gal4 activation domain (AD) and Gal4 DNA

binding domain (DBD) fusion plasmids were constructed by PCR amplifying the indicated

regions and cloning into pGBD-C3 and pGAD-C3 [63] using Gibson Assembly. The plasmids were

co-transformed and selected for on SD -leu -trp. Positive colonies were patched onto fresh

plates and grown at 30⁰C for 24 hours before replica plating onto SD -leu -trp -ade -his to test

for interactions. Plates were imaged after 45-50 hours.

Statistical analyses

All replicates are biological. For yeast strains, the same strain was grown and measured in

separate vials. Independent transformants were tested for the flow cytometry host strains and

randomly selected constructs and were found to generate consistent results. For recombinant

proteins, the same constructs were independently transformed into the bacteria and separate

batches of purified proteins were used for the measurements.

Binding affinity comparison by BLI (Fig. 2): K_D estimates from either steady state or kinetic curve

fitting analyses were log10 transformed and a Student's t-test was used to compare ScPho4 vs

CgPho4 DBDs against the same 17-bp DNA. Raw two-sided _P_-values and Holm-Bonferroni

corrected _P_ values for the 10 tests were reported; _Activation_ (Fig. 3): median fluorescence

intensity (MFI) for the _GAL1pr_-mCherry reporter was recorded from flow cytometry for each

sample. For Fig. 3C, each Gal4 DBD fusion construct was compared to the background level.

Because the host (with _GAL1pr_-mCherry but no Gal4 DBD plasmid) and Gal4 DBD alone

showed a similar level of low MFI, we combined them as the reference group. A linear model

was fit in R with the following command `lm(MFI ~ Genotype)`, where "Genotype" is a factor

representing the constructs. This model estimates a common standard deviation for all

constructs and tests the significance of each construct against the reference group with a two-

sided t-test. The raw _P_-values were corrected for multiple testing using the Holm-Bonferroni

procedure. Constructs with a corrected _P_-value <= 0.05 were considered significant. For Fig. 3D,

the first two groups were tested using a two-sample t-test, while the third involved multiple levels,

and were tested as described above for Fig. 3C. A Holm-Bonferroni correction was applied to

the raw *P*-values for all six tests together. *Epistasis between regions R1, AD, and NLS of*

CgPho4 (Fig. 5D): to determine the main and interaction terms in these swaps from CgPho4 on

ScPho4's background, we fit a linear model, $Y = X_0 + R1 + AD + NLS + R1{:}AD + R1{:}NLS +$

$AD{:}NLS + R1{:}AD{:}NLS$ in R, using the command `lm(A ~ R1 * AD * NLS)`, where "A" is either

A_{PHO2} or $A_{pho2\Delta}$, and the independent variables are coded as ScPho4 = 0, CgPho4 = 1. The raw

P-values were adjusted for multiple testing using the Holm-Bonferroni procedure. *Split P2ID* (Fig.

6B): The same procedure as applied to the activation region test above (Fig. 3) was applied to

the split P2ID swaps. Two sets of four chimeras were chosen, each with a reference construct

having P2ID:Cg, one with the entire P2ID swapped for the ScPho4 version, and two with the

first or second half of P2ID swapped for the ScPho4 version. The latter three were compared to

the reference using a linear model `lm(MFI ~ Genotype)`, for A_{PHO2} and $A_{pho2\Delta}$ separately. Holm-

Bonferroni correction was applied to the two sets combined (6 tests in total), again separately

for A_{PHO2} and $A_{pho2\Delta}$. A corrected $P < 0.05$ was considered significant. *Nuclear fraction of Pho4*

chimeras (Supplementary Fig. 7): The ratio of the sum of pixel values in the GFP channel inside

the nucleus vs the whole cell was treated as the response variable, and the identity of the NLS

region ("Cg" vs "Sc") as the predictor. First, a two-way ANOVA was performed with the formula

`lm(Nuc_frac ~ NLS + Host)`, where Host is either *PHO2* or *pho2Δ*. Second, given that ratios

are often non-normally distributed, we performed the non-parametric Kruskal-Wallis test with the

command `kruskal.test(Nuc_frac ~ NLS)` separately in the *PHO2* and *pho2Δ* hosts. The

ANOVA F-test *P*-value (0.13) was reported in the results. The two *P*-values for the second test

were 0.42 (*PHO2*) and 0.36 (*pho2Δ*).

698   **DATA AVAILABILITY**

Source data for figures in this paper are provided as a Source Data file and are available at

https://doi.org/10.5281/zenodo.14501732. Raw microscopy images for quantifying Pho4 nuclear

concentration are available at https://doi.org/10.6084/m9.figshare.28437011 . Protein Binding

Microarray data are available through the Gene Expression Omnibus (GEO) under GSE293214

and GSE293355. No restrictions apply to any of the data generated in this study.

CODE AVAILABILITY

Scripts for figures and statistical tests are available at https://doi.org/10.5281/zenodo.14501732.

REFERENCES

707
1. Stewart, A. J., Hannenhalli, S. & Plotkin, J. B. Why Transcription Factor Binding Sites Are

Ten Nucleotides Long. *Genetics* **192**, 973–985 (2012).

2. Berger, M. F. *et al.* Variation in homeodomain DNA binding revealed by high-resolution

analysis of sequence preferences. *Cell* **133**, 1266–1276 (2008).

3. Zhou, X. & O'Shea, E. K. Integrated approaches reveal determinants of genome-wide

binding and function of the transcription factor Pho4. *Mol. Cell* **42**, 826–836 (2011).

4. Merhej, J. *et al.* A Network of Paralogous Stress Response Transcription Factors in the

Human Pathogen Candida glabrata. *Front. Microbiol.* **7**, (2016).

5. Sánchez-Higueras, C. *et al.* In vivo Hox binding specificity revealed by systematic changes

to a single cis regulatory module. *Nat. Commun.* **10**, 3597 (2019).

6. Slattery, M. *et al.* Cofactor binding evokes latent differences in DNA binding specificity

between Hox proteins. *Cell* **147**, 1270–1282 (2011).

7. Todeschini, A.-L., Georges, A. & Veitia, R. A. Transcription factors: specific DNA binding

and specific gene regulation. *Trends Genet.* **30**, 211–219 (2014).

8. Avsec, Ž. *et al.* Base-resolution models of transcription-factor binding reveal soft motif

syntax. *Nat. Genet.* (2021) doi:10.1038/s41588-021-00782-6.

9. Guss, K. A., Nelson, C. E., Hudson, A., Kraus, M. E. & Carroll, S. B. Control of a Genetic

Regulatory Network by a Selector Gene. *Science* **292**, 1164–1167 (2001).

10. Ang, Y.-S. *et al.* Disease Model of GATA4 Mutation Reveals Transcription Factor

Cooperativity in Human Cardiogenesis. *Cell* **167**, 1734-1749.e22 (2016).

11. Baker, C. R., Booth, L. N., Sorrells, T. R. & Johnson, A. D. Protein modularity, cooperative

binding, and hybrid regulatory states underlie transcriptional network diversification. *Cell* **151**,

80–95 (2012).

12. Britton, C. S., Sorrells, T. R. & Johnson, A. D. Protein-coding changes preceded cis-

regulatory gains in a newly evolved transcription circuit. *Science* **367**, 96–100 (2020).

13. Traven, A., Jelicic, B. & Sopta, M. Yeast Gal4: a transcriptional paradigm revisited. *EMBO*

Rep. **7**, 496–499 (2006).

14. Kerwin, C. L. & Wykoff, D. D. Candida glabrata PHO4 Is Necessary and Sufficient for Pho2-

Independent Transcription of Phosphate Starvation Genes. *Genetics* **182**, 471–479 (2009).

15. He, B. Z., Zhou, X. & O'Shea, E. K. Evolution of reduced co-activator dependence led to

target expansion of a starvation response pathway. *eLife* **6**, (2017).

16. Barbaric, S., Münsterkötter, M., Goding, C. & Hörz, W. Cooperative Pho2-Pho4 Interactions

at the PHO5 Promoter Are Critical for Binding of Pho4 to UASp1 and for Efficient

Transactivation by Pho4 at UASp2. *Mol. Cell. Biol.* **18**, 2629–2639 (1998).

17. Nourani, A., Utley, R. T., Allard, S. & Côté, J. Recruitment of the NuA4 complex poises the

PHO5 promoter for chromatin remodeling and activation. *EMBO J.* **23**, 2597–2607 (2004).

18. Magbanua, J. P., Ogawa, N., Harashima, S. & Oshima, Y. The transcriptional activators of

the PHO regulon, Pho4p and Pho2p, interact directly with each other and with components

of the basal transcription machinery in Saccharomyces cerevisiae. *J. Biochem. (Tokyo)* **121**,

1182–1189 (1997).

19. Shao, D., Creasy, C. L. & Bergman, L. W. Interaction of Saccharomyces cerevisiae Pho2

with Pho4 increases the accessibility of the activation domain of Pho4. *Mol. Gen. Genet.*

MGG **251**, 358–364 (1996).

20. Jayaraman, P. S., Hirst, K. & Goding, C. R. The activation domain of a basic helix-loop-helix

protein is masked by repressor interaction with domains distinct from that required for

transcription regulation. *EMBO J.* **13**, 2192–2199 (1994).

21. Ogawa, N. & Oshima, Y. Functional domains of a positive regulatory protein, PHO4, for

transcriptional control of the phosphatase regulon in Saccharomyces cerevisiae. *Mol. Cell.*

Biol. **10**, 2224–2236 (1990).

22. McAndrew, P. C., Svaren, J., Martin, S. R., Hörz, W. & Goding, C. R. Requirements for

Chromatin Modulation and Transcription Activation by the Pho4 Acidic Activation Domain.

Mol. Cell. Biol. **18**, 5818–5827 (1998).

23. Komeili, A. & O'Shea, E. K. Roles of Phosphorylation Sites in Regulating Activity of the

Transcription Factor Pho4. *Science* **284**, 977–980 (1999).

24. O'Neill, E. M., Kaffman, A., Jolly, E. R. & O'Shea, E. K. Regulation of PHO4 nuclear

localization by the PHO80-PHO85 cyclin-CDK complex. *Science* **271**, 209–212 (1996).

25. Hirst, K., Fisher, F., McAndrew, P. C. & Goding, C. R. The transcription factor, the Cdk, its

cyclin and their regulator: directing the transcriptional response to a nutritional signal. *EMBO*

J. **13**, 5410–5420 (1994).

26. Shimizu, T. *et al.* Crystal structure of PHO4 bHLH domain–DNA complex: flanking base

recognition. *EMBO J.* **16**, 4689–4697 (1997).

27. Kaffman, A., Herskowitz, I., Tjian, R. & O'Shea, E. K. Phosphorylation of the transcription

factor PHO4 by a cyclin-CDK complex, PHO80-PHO85. *Science* **263**, 1153–1156 (1994).

28. Gomes-Vieira, A. L. *et al.* Evolutionary conservation of a core fungal phosphate

homeostasis pathway coupled to development in Blastocladiella emersonii. *Fungal Genet.*

Biol. **115**, 20–32 (2018).

29. Gordân, R. *et al.* Genomic regions flanking E-box binding sites influence DNA binding

specificity of bHLH transcription factors through DNA shape. *Cell Rep.* **3**, 1093–1104 (2013).

30. Sanborn, A. L. *et al.* Simple biochemical features underlie transcriptional activation domain

diversity and dynamic, fuzzy binding to Mediator. *eLife* **10**, e68068 (2021).

31. Piskacek, S. *et al.* Nine-amino-acid transactivation domain: establishment and prediction

utilities. *Genomics* **89**, 756–768 (2007).

32. Bhoite, L. T. *et al.* Mutations in the Pho2 (Bas2) Transcription Factor That Differentially

Affect Activation with Its Partner Proteins Bas1, Pho4, and Swi5. *J. Biol. Chem.* **277**, 37612–

37618 (2002).

33. Ming Yip, H., Cheng, S., Olson, E. J., Crone, M. & Maerkl, S. J. Perfect adaptation achieved

by transport limitations governs the inorganic phosphate response in S. cerevisiae. *Proc.*

Natl. Acad. Sci. U. S. A. **120**, e2212151120 (2023).

34. Donovan, B. T. *et al.* Basic helix-loop-helix pioneer factors interact with the histone octamer

to invade nucleosomes and generate nucleosome-depleted regions. *Mol. Cell* **83**, 1251-

1263.e6 (2023).

35. Liu, J. *et al.* Intrinsic Disorder in Transcription Factors. *Biochemistry* **45**, 6873–6888 (2006).

36. Lynch, V. J., May, G. & Wagner, G. P. Regulatory evolution through divergence of a

phosphoswitch in the transcription factor CEBPB. *Nature* **480**, 383–386 (2011).

37. Hsu, I. S. *et al.* A functionally divergent intrinsically disordered region underlying the

conservation of stochastic signaling. *PLOS Genet.* **17**, e1009629 (2021).

38. Krois, A. S., Dyson, H. J. & Wright, P. E. Long-range regulation of p53 DNA binding by its

intrinsically disordered N-terminal transactivation domain. *Proc. Natl. Acad. Sci.* **115**,

E11302–E11310 (2018).

39. Brodsky, S. *et al.* Intrinsically Disordered Regions Direct Transcription Factor In Vivo

Binding Specificity. *Mol. Cell* (2020) doi:10.1016/j.molcel.2020.05.032.

40. Ferrie, J. J., Karr, J. P., Tjian, R. & Darzacq, X. "Structure"-function relationships in

eukaryotic transcription factors: The role of intrinsically disordered regions in gene

regulation. *Mol. Cell* **0**, (2022).

41. Ferrie, J. J. *et al.* p300 is an obligate integrator of combinatorial transcription factor inputs.

Mol. Cell S1097-2765(23)01023–7 (2023) doi:10.1016/j.molcel.2023.12.004.

42. Mindel, V. *et al.* Intrinsically disordered regions of the Msn2 transcription factor encode

multiple functions using interwoven sequence grammars. *Nucleic Acids Res.* gkad1191

(2023) doi:10.1093/nar/gkad1191.

43. Ji, D. *et al.* FOXA1 forms biomolecular condensates that unpack condensed chromatin to

function as a pioneer factor. *Mol. Cell* **84**, 244-260.e7 (2024).

44. Rives, A. *et al.* Biological structure and function emerge from scaling unsupervised learning

to 250 million protein sequences. *Proc. Natl. Acad. Sci.* **118**, e2016239118 (2021).

45. Tuch, B. B., Galgoczy, D. J., Hernday, A. D., Li, H. & Johnson, A. D. The evolution of

combinatorial gene regulation in fungi. *PLoS Biol.* **6**, e38 (2008).

46. Wapinski, I. *et al.* Gene duplication and the evolution of ribosomal protein gene regulation in

yeast. *Proc. Natl. Acad. Sci. U. S. A.* **107**, 5505–5510 (2010).

47. Tsong, A. E., Tuch, B. B., Li, H. & Johnson, A. D. Evolution of alternative transcriptional

circuits with identical logic. *Nature* **443**, 415–420 (2006).

48. Nocedal, I. & Johnson, A. D. How Transcription Networks Evolve and Produce Biological

Novelty. *Cold Spring Harb. Symp. Quant. Biol.* **80**, 265–274 (2015).

49. Sherbenou, D. W. & Druker, B. J. Applying the discovery of the Philadelphia chromosome. *J.*

Clin. Invest. **117**, 2067–2074 (2007).

50. Do, C. B., Mahabhashyam, M. S. P., Brudno, M. & Batzoglou, S. ProbCons: Probabilistic

consistency-based multiple sequence alignment. *Genome Res.* **15**, 330–340 (2005).

51. Waterhouse, A. M., Procter, J. B., Martin, D. M. A., Clamp, M. & Barton, G. J. Jalview

Version 2--a multiple sequence alignment editor and analysis workbench. *Bioinforma. Oxf.*

Engl. **25**, 1189–1191 (2009).

52. Troshin, P. V. *et al.* JABAWS 2.2 distributed web services for Bioinformatics: protein

disorder, conservation and RNA secondary structure. *Bioinformatics* **34**, 1939–1940 (2018).

53. Buchan, D. W. A. & Jones, D. T. The PSIPRED Protein Analysis Workbench: 20 years on.

Nucleic Acids Res. **47**, W402–W407 (2019).

54. Thomas-Chollier, M. *et al.* RSAT peak-motifs: motif analysis in full-size ChIP-seq datasets.

Nucleic Acids Res. gkr1104 (2011) doi:10.1093/nar/gkr1104.

55. Zhang, Y., Ho, T. D., Buchler, N. E. & Gordân, R. Competition for DNA binding between

paralogous transcription factors determines their genomic occupancy and regulatory

functions. *Genome Res.* **31**, 1216–1229 (2021).

56. Berger, M. F. & Bulyk, M. L. Universal protein-binding microarrays for the comprehensive

characterization of the DNA-binding specificities of transcription factors. *Nat. Protoc.* **4**, 393–

411 (2009).

57. Berger, M. F. & Bulyk, M. L. Protein Binding Microarrays (PBMs) for Rapid, High-

Throughput Characterization of the Sequence Specificities of DNA Binding Proteins. in

Gene Mapping, Discovery, and Expression vol. 338 245–260 (Humana Press, New Jersey,

2006).

58. Maerkl, S. J. & Quake, S. R. A Systems Approach to Measuring the Binding Energy

Landscapes of Transcription Factors. *Science* **315**, 233–237 (2007).

59. Weeramange, C. J., Fairlamb, M. S., Singh, D., Fenton, A. W. & Swint-Kruse, L. The

strengths and limitations of using biolayer interferometry to monitor equilibrium titrations of

biomolecules. *Protein Sci.* pro.3827 (2020) doi:10.1002/pro.3827.

60. Anand, R., Memisoglu, G. & Haber, J. Cas9-mediated gene editing in Saccharomyces

cerevisiae. (2017).

61. Gietz, R. D. & Schiestl, R. H. High-efficiency yeast transformation using the LiAc/SS carrier

DNA/PEG method. *Nat. Protoc.* **2**, 31–34 (2007).

62. Lam, F. H., Steger, D. J. & O'Shea, E. K. Chromatin decouples promoter threshold from

dynamic range. *Nature* **453**, 246–250 (2008).

63. James, P., Halladay, J. & Craig, E. A. Genomic libraries and a host strain designed for

highly efficient two-hybrid selection in yeast. *Genetics* **144**, 1425–1436 (1996).

856
ACKNOWLEDGEMENTS

We would like to thank Dr. Miles Pufall's lab for teaching us EMSA. We thank Dr. Jan Fassler

for sharing many yeast plasmids for the yeast one- and two-hybrids. Drs. Jan Fassler, Miles

Pufall, Craig Ellermeier and Todd Washington all critically read the thesis chapter by LFS, on

which this manuscript is based. We thank Kyle Malcolm for helping with the early development

of the BLI assay. Christian Weinrich discovered the P2ID's effect on Pho2-dependence during

his rotation. We thank Dr. Yann Vanrobaeys for helping with an initial analysis using PADDLE.

We would like to acknowledge the use of resources at the Protein and Crystallography Facility

within the Carver College of Medicine at the University of Iowa and thank Lokesh Gakhar, Zhen

Xu, and Devin Reusch for assistance with protein purification and BLI assays. This work was

primarily supported by NIH R35-GM137831 (to BZH). LFS was supported on NIH Predoctoral

Training Grant T32GM008629; BJB was supported on NIH Predoctoral Training Grant

T32GM1454441; RG was supported by NIH R01-GM135658 and BZH was also supported by a

startup fund from the University of Iowa.

AUTHOR CONTRIBUTIONS

L.F.S. and B.Z.H. designed the experiments. L.F.S. and E.M.O. constructed the chimeras and

performed the flow cytometry. L.F.S. performed the EMSA and the yeast two hybrid

experiments. J.L. performed the microscopy for nuclear localization analysis. E.M.O. and B.J.B.

performed experiments and analyses for the revision. Y.Z. and W.Z. performed the PBM

experiments with input from R.G.. J.Z., T.J.C. established critical methods. X. Z. generated

preliminary results for the work. N.J.S. gave input on and performed BLI experiments. L.F.S,

Y.Z., and B.Z.H. analyzed the data. L.F.S. and B.Z.H. wrote the manuscript with edits from all

co-authors.

COMPETING INTERESTS

The authors declare no conflicting interests.

                                              TABLES

    Table 1. Breakpoints for chimeric Pho4 and individual regions tested in this study

Origin	Region Name	Start	End	Length	Used in figure(s)
CgPho4	R1	1	44	44	Fig. 1,4,5,6,S2,S4
CgPho4	AD	45	112	68	Fig. 1,4,5,6,S2,S4
CgPho4	NLS	113	282	170	Fig. 1,4,5,6,S2,S4
CgPho4	P2ID	283	458	176	Fig. 1,4,5,6,S2,S4
CgPho4	DBD	459	533	75	Fig. 1,4,5,6,S2,S4
ScPho4	R1	1	42	42	Fig. 1,4,5,6,S2,S4
ScPho4	AD	43	99	57	Fig. 1,4,5,6,S2,S4
ScPho4	NLS	100	176	77	Fig. 1,4,5,6,S2,S4
ScPho4	P2ID	177	242	66	Fig. 1,4,5,6,S2,S4
ScPho4	DBD	243	312	70	Fig. 1,4,5,6,S2,S4
CgPho4	P2ID first half	283	327	45	Fig. 6, S2
CgPho4	P2ID second half	328	458	131	Fig. 6, S2
ScPho4	P2ID first half	177	204	28	Fig. 6, S2
ScPho4	P2ID second half	205	242	38	Fig. 6, S2
CgPho4	E1	2	44	43	Fig. 3,S5
CgPho4	AD	45	126	82	Fig. 3,S5
CgPho4	E2	252	463	212	Fig. 3,S5
CgPho4	AD.1	45	75	31	Fig. 3,S5
CgPho4	AD.2	76	126	51	Fig. 3,S5
CgPho4	E2_9aa	252	288	37	Fig. 3,S5
ScPho4	E1	2	42	41	Fig. 3,S5
ScPho4	AD	43	101	59	Fig. 3,S5
ScPho4	AD_9aa	70	101	32	Fig. 3,S5
ScPho4	E2	156	250	95	Fig. 3,S5
ScPho2	AD	404	559	156	Fig. 3
ScPho4	Alternative NLS	100	156	57	Fig. 5
CgPho4	Alternative P2ID	253	463	211	Fig. 5
ScPho4	ΔbHLH	2	250	249	Fig. 6 Y2H
CgPho4	ΔbHLH	2	463	462	Fig.6 Y2H
ScPho2	Pho4int	112	407	296	Fig. 6 Y2H
CgPho4	purified DBD	452	533	82	Fig. 2,S4
ScPho4	purified DBD	236	312	77	Fig. 2,S4

    Table 2. Yeast strains

ID	Genotype	Source
yH373	pho4::NAT; pho2::HIS3;	this study

	pho80::TRP1; pho5::mCherry	
yH555	pho4::NAT; his3::pRS303-HIS3; pho80::TRP1; pho5::mCherry	this study
yH549	gal4delta gal80delta LYS2::GAL1-HIS3 GAL2-ADE2 MET2::GAL7-lacZ	PMID: 8978031, gift from Fassler lab
yH696	gal4delta gal80delta LYS2::GAL1-HIS3; GAL2-ADE2; MET2::GAL7-lacZ; ura3::pRS306-*GAL1pr*-mCherry-URA3	this study
yH156	pho80::TRP1	PMID: 28485712
various	chimeric Pho4 plasmids transformed into yH373 and yH555	this study
various	yeast two-hybrid plasmids transformed into yH696	this study
various	yeast one-hybrid plasmids transformed into yH696	this study

FIGURE LEGENDS

Fig. 1 Different co-TF (Pho2) dependence between orthologous TF (Pho4) in yeast

species. (A) Cartoon summary of previous work: *S. cerevisiae* Pho4 (ScPho4) depends on the

co-TF, Pho2, for DNA binding and activation of Phosphate Starvation (PHO) Response genes

(top row), while Pho4 from *C. glabrata* (CgPho4) can bind and activate without the co-TF. See

text for references. (B) Two non-mutually exclusive models to explain the divergence in Pho2-

dependence. (C) The tiles in the middle show the conservation scores at aligned positions of the

two Pho4 orthologs; the box diagrams represent ScPho4 and CgPho4, with gray lines

connecting matching residues to the alignment tiles. Five regions were delineated based on

knowledge about ScPho4: R1 = regulatory, interact with Pho80; AD = activation domain; NLS =

nuclear import and export signals; P2ID = Pho2-interaction domain (also includes a region

contacting Pho80); DBD = basic Helix-Loop-Helix DNA binding domain. The lines and cylinders

above and below the region diagrams show predicted α helices in each protein, with lighter

color corresponding to lower prediction confidence. Prediction was done using PSIPRED 4.0.

Fig. 2 CgPho4 DBD recognizes the same motif as ScPho4 DBD and has a higher affinity towards the consensus DNA. (A) Alignment of the DBD region between the two Pho4 orthologs. Top: sequence logo for the bHLH domain (PROSITE PS50888). Middle: coordinates above are based on ScPho4. Filled and open triangles indicate residues in ScPho4 DBD making either base or phosphate backbone contacts (Shimizu *et al.*, 1997). Bottom: predicted α-helices in both Pho4 DBDs. (B) WebLogo motifs for ScPho4 and CgPho4 are derived from ChIP-seq peaks. (C) Universal Protein Binding Microarray (uPBM) comparing the binding of full length ScPho4 and CgPho4 against all possible 9-mers. A rank-order based E-score is plotted for all 7-mers, where a score greater than 0.35 indicates specific TF-DNA binding (Gordân *et al.*, 2013). Red dots in the top right corner represent oligos with the "CACGTG" motif. Pearson's correlation coefficient and a *p*-value based on the t-distribution are shown at the top. (D) Binding to the 17bp oligo containing a consensus E-box motif is measured by Biolayer interferometry (BLI). Fractional binding is plotted vs the concentration of purified ScPho4 or CgPho4 DBD. One of the representative binding curves is shown here. (E) Dissociation constants (K_D) are determined from the binding curves and shown as dots (n = 5 biological replicates). The bars and the red dots are the means and the red lines the 95% confidence intervals from bootstrapping. Two-sided Student's t-test was performed on the log transformed K_D. A Wilcoxon rank-sum test on raw K_D similarly yielded $P < 0.01$.

Fig. 3 CgPho4 encodes two Activation Enhancer Domains (AEDs) that can enhance the activation potential of the main AD. (A, B) PADDLE prediction of activation potential in ScPho4 and CgPho4. Predictions were made in 53 aa blocks and plotted using the middle of each block as the x coordinate. As a result, the first and last 26 aa positions have no scores. Z-scores > 4 and 6 indicate regions with medium and strong activation potential, respectively. The five regions in each protein are labeled as in Fig. 1. Orange triangles mark the 9aaTAD motif

matches. Below the plots are diagrams of the Gal4DBD fusion constructs used to test the

activation potential, with their boundaries labeled on the right. (C) Activation potential of each

region was tested in the context of a Gal4 DBD fusion driving a *GAL1pr*-mCherry reporter.

Shown are the mean (bar) and individual data points (n=3, except for the host, for which n = 6,

biological replicates). Host has the mCherry reporter but no plasmid; Gal4_DBD has the

reporter and the Gal4 DBD alone. A linear model with construct as the independent variable

was used to compare the Gal4 DBD fusions to the two negative controls combined; an asterisk

indicates significant difference at a 0.05 level after Holm-Bonferroni correction. (D) CgE1 and

CgE2_9aa can enhance the activation capability of ScAD_9aa in an orientation-independent

manner. Plot design and replicate numbers are the same as in C. An asterisk indicates

significance difference at a 0.05 level by an unpaired, two-sided Student's t-test between the

indicated groups after Holm-Bonferroni correction.

Fig. 4 A dual fluorescence reporter system for accurately quantifying the activity of Pho4

chimeras with and without Pho2. (A) ScPho4 and CgPho4 were divided into five

corresponding parts. Names of each region were based on the annotations for ScPho4.

Predicted alpha helices were drawn as cartoons outside of the box diagrams. All $2^5 = 32$

combinations (2 were the original proteins) were made in the main chimera set, and an

additional set of chimeras were constructed to test specific hypotheses. (B) A diagram showing

the design of the dual fluorescence reporter system. The chimeric Pho4 fragments were cloned

downstream of the endogenous *ScPHO4* promoter and were tagged at the C-terminus with an

in-frame mNeon fluorescent protein, followed by the endogenous *ScPHO4* 3'UTR and

terminator. The plasmid contains a CEN/ARS element, resulting in stable inheritance and low

biological variation. (C) The *PHO5pr-mCherry* levels correlate with the ScPho4-mNeon levels

between biological replicates. A 95% confidence band centered on the linear regression line in

blue is shown as a gray ribbon. (D) A_{PHO2} and $A_{pho2\Delta}$ values for ScPho4 and CgPho4 across 6

different days of experiments showing consistent results (n = 3 for all groups except ScPho4

with Pho2, where n = 12, biological replicates, separate inoculates at the overnight growth

stage.)

Fig. 5. R1 and NLS of CgPho4 confer stronger activity than their counterparts in ScPho4.

(A) Pho4 chimeras (black), endogenous ScPho4 (green) and CgPho4 (blue) were plotted based

on their activity with and without Pho2. The solid line indicates equal activity (independent of

Pho2). (B) Heatmap showing the difference in activity between the CgPho4 region(s) and the

counterpart of ScPho4 measured in the ScPho4 background. The upper half shows the

difference with Pho2 and the lower half without Pho2. The triangles (e.g., a) show the difference

between single regions while the squares on the top show the differences when swapping two

regions (see key to the right). Red color indicates a higher activity of the region from CgPho4

than the counterpart in ScPho4 and blue color indicates decreased activity. (C) A subset of the

chimeras were shown with their A_{PHO2} and $A_{pho2\Delta}$ values. Bars represent the means; lines are 95%

confidence intervals based on bootstrap; plus signs are individual data points (n=6, biological

replicates), which were not shown for the endogenous ScPho4 and CgPho4 (n=36, biological

replicates). Vertical dotted lines equal the endogenous ScPho4 level. (D) Epistasis analysis of

CgPho4 R1, AD and NLS. A linear model was fit to the data with all two and three-way

interactions. Shown are the individual region and interaction coefficient estimates (bar and point)

for the CgPho4 region(s). Line ranges centered on the estimates (bar and point) indicate the

standard errors of the coefficients as calculated by the linear model fit. Dark gray indicates the

term is significant at 0.05 level by student's t-test after Holm-Bonferroni correction.

Fig. 6 P2ID in ScPho4 allows for enhanced activation with Pho2 via physical interaction

with the co-TF. (A) Scatter plot of all chimeric Pho4 constructs based on their A_{PHO2} and $A_{pho2\Delta}$

values, colored by the identity of their P2ID region. Mixed P2ID uses an additional breakpoint

that separates the region into two halves (Supplementary Fig. 2). (B) Comparing the effects of

swapping the whole P2ID from ScPho4 into CgPho4 vs individual halves. Bars represent the

means; lines are 95% confidence intervals based on bootstrap; plus signs are individual data

points (n=6, biological replicates), which were not shown for the endogenous ScPho4 and

CgPho4 (n=36, biological replicates). Two sets were tested (rows 2-5 and 6-9). Two-sided

Student's t-tests were performed as indicated by the bars and lines to the right. Holm-Bonferroni

correction was applied to the raw P-values for both sets, separately for A_{PHO2} and $A_{pho2\Delta}$. An asterisk

indicates corrected P < 0.05. (C) Testing the hypothesis that ScPho4 DBD requires the P2ID:Sc

and Pho2 to be fully functional. Plot designs, including bars, line ranges and replicate numbers

are the same as in (B). Compared with the chimeras with P2ID:Cg + P2ID:Sc, which are all non-

functional even with Pho2, adding back P2ID:Sc (4+5) rescued the activity with Pho2, but not

without. (D) Pho4-Pho2 physical interaction was tested with a yeast two-hybrid assay. The bait

and target regions were fused to the activation domain (AD) and DNA binding domain (DBD) of

the yeast Gal4 protein. Interaction between them reconstitutes a functional Gal4 and allows

cells to grow on the dropout media (right column), while all strains are able to grow on the rich

media (left column). Vector indicates a Gal4 DBD or Gal4 AD only plasmid without a bait or

target fused to it and is included as a control. Two independent transformants are shown.

Fig. 7 Summary of Pho4 protein divergence and its contribution to the Pho2-dependence

variation in ScPho4 and CgPho4. Left (1): CgPho4 DBD recognizes the same DNA

sequences as ScPho4 DBD, but binds the consensus motif with a 3-4 fold higher affinity. While

consistent with the enhanced activity model, further studies found no evidence for the enhanced

activity to be a major factor for the reduced Pho2-dependence. (2) Two regions in the IDR of

CgPho4 (red asterisks) can enhance the activity of the main activation domain (AD) but are not

sufficient on their own to activate gene expression. These two Activation Enhancing Domains

(AEDs) directly contribute to CgPho4's reduced Pho2-dependence. Right (3): An IDR (green

asterisk) adjacent to the DBD in ScPho4, named P2ID, can both interact with ScPho2 and boost

the activity of Pho4 and also repress its activity when Pho2 is absent, providing support for the

autoinhibition model.

A

▲ Base pair recognition   △ Phosphate recognition

B

ScPho4   CgPho4

C

r = 0.89, *p* < 0.01

D

E

p < 0.01
n = 5

A

ScPho4 | R1 | AD | NLS | P2ID | DBD | 312 aa
CgPho4 | R1 | AD | NLS | P2ID | DBD | 533 aa

Helix ▮ Disordered ▬ Activation ▮ Medium ▮ Strong

30 main set   28 variations

B

ScPHO4 promoter

Pho4 chimera — mNeon
Chimera plasmid library
CEN/ARS

transform

Pho4 — 2 — mCherry
ScPHO5 promoter

Pho4 — mCherry
ScPHO5 promoter

flow cytometry

$A_{PHO2} = \dfrac{\text{(mCherry)}}{\text{(mNeon)}}$

$A_{pho2\Delta} = \dfrac{\text{(mCherry)}}{\text{(mNeon)}}$

C

ScPho4

PHO5pr-mCherry vs Pho4-mNeon

D

host   ▮ *PHO2*   ▮ *pho2Δ*

mCherry/mNeon — Day of Experiment

ScPho4

CgPho4

 CgPho4 AED boost activity

 P2ID:Sc dual-function