Alternative oxidase alleviates mitochondrial oxidative stress during limited nitrate

reduction in *Arabidopsis thaliana*

Daisuke Otomaru[a], Natsumi Ooi[a,b], Kota Monden[a], Takamasa Suzuki[c], Ko Noguchi[d], Tsuyoshi

Nakagawa[a] and Takushi Hachiya[a†]

[a]Department of Molecular and Functional Genomics, Interdisciplinary Center for Science

Research, Shimane University, 1060 Nishikawatsu-cho, Matsue, Shimane 690-8504, Japan

[b]Division of Biological Science, Nara Institute of Science and Technology, 8916-5 Takayama,

Ikoma, Nara 630-0192, Japan

[c]College of Bioscience and Biotechnology, Chubu University, 1200 Matsumoto-cho, Kasugai,

Aichi 487-8501, Japan

[d]Department of Life Sciences, Tokyo University of Pharmacy and Life Sciences, 1432-1

Horinouchi Hachioji, Tokyo 192-0392, Japan

[†]**To whom correspondence should be addressed.**

Takushi Hachiya

E-mail: takushi.hachiya@life.shimane-u.ac.jp

Tel.: +81-852-32-6288

Fax: +81-852-32-6109

Funding

This study was supported by KAKENHI Grants from the Japan Society for the Promotion of

Science [Grant-in-Aid for Scientific Research (C) No. 20K05771/23K04978 to TH].

Abstract

The conversion of nitrate to ammonium, i.e., nitrate reduction, is a major consumer of reductants in plants. Previous studies have reported that the mitochondrial alternative oxidase (AOX) is upregulated under limited nitrate reduction conditions, including no/low nitrate or when ammonium is the sole nitrogen (N) source. Electron transfer from ubiquinone to AOX bypasses the proton-pumping complexes III and IV, thereby consuming reductants efficiently. Thus, upregulated AOX under limited nitrate reduction may dissipate excessive reductants and thereby attenuate oxidative stress. Nevertheless, so far there is no firm evidence for this hypothesis due to the lack of experimental systems to analyze the direct relationship between nitrate reduction and AOX. We therefore developed a novel culturing system for *A. thaliana* that manipulates activities of nitrate reduction and AOX separately without causing N starvation, ammonium toxicity, and lack of nitrate signal. Using this system we examined genome-wide gene expression and plant growth to better understand the relationship between AOX and nitrate reduction. Here we show that AOX alleviates mitochondrial oxidative stress and sustains plant growth under limited nitrate reduction.

1 | INTRODUCTION

Plants use nitrate and ammonium as main sources of nitrogen (N). The conversion of nitrate to ammonium, i.e., nitrate reduction, requires eight moles of electrons per mole of nitrate, which accounts for about half of the energy required for protein synthesis from nitrate (Hachiya et al., 2007). Thus, ammonium is an energetically superior N source, but most crops prefer nitrate (Miller & Cramer, 2004). In herbaceous plants, nitrate reduction generally occurs in the shoots (Andrews et al., 1992; Hachiya et al., 2020; Scheurwater et al., 2002), and acts as a major sink of reductants generated by the chloroplasts (Rasmusson et al., 2020). Hence, when the nitrate supply is limited, excessive reductants may accumulate within illuminated photosynthetic cells. Recent studies have shown that the $NAD(P)H/NAD(P)^+$ ratio and the levels of hydrogen peroxide are increased in *A. thaliana* leaves of ammonium-grown plants compared with nitrate-grown plants (Podgórska et al., 2013; Podgórska et al., 2020; Rasmusson et al, 2020). Moreover, N starvation treatments have been found to enlarge the ascorbate pool in several plant species (Iwagami et al., 2022). Taken together, these findings suggest that a decrease in nitrate reduction could lead to oxidative stress in plants.

It is widely accepted that nitrate reduction significantly alters the action of the plant mitochondrial electron transport chain (mETC) (Rasmusson et al., 2020). In fact, the amount of electrons consumed in nitrate reduction appears to be comparable to that in the mETC (Noctor & Foyer, 1998; Rasmusson et al., 2020). In photosynthetic tissues, the expression and activities of several enzymes that bypass energy conservation steps in mETC are induced under limited nitrate reduction conditions; i.e., if the plant is experiencing N starvation or if ammonium is the sole N source (Escobar et al., 2006; Hachiya et al., 2010; Noguchi & Terashima, 2006; Qiao et al., 2022; Watanabe et al., 2010). One enzyme is the alternative oxidase (AOX), which allows direct electron transfer from ubiquinone to molecular oxygen, bypassing the proton-pumping complexes III and IV (Noguchi & Yoshida, 2008). Therefore, upregulation of AOX under limited nitrate reduction may dissipate excessive reductants without being limited to steep proton gradients

across the mitochondrial inner membrane, thereby attenuating reactive oxygen species (ROS)

production and oxidative stress. Actually, in *Arabidopsis* plants under low nitrate conditions, the

shoot expression of antioxidant enzyme genes was induced by disrupting the major isoform

AOX1a (Watanabe et al., 2010). Antisense suppression of *Arabidopsis AOX1a* was found to

greatly decrease the reducing state of ascorbate under ammonium but not under nitrate (Podgórska

et al., 2020). In barley grown under low nitrate, chemical inhibition of AOX activity elevated the

levels of hydrogen peroxide (Wang et al., 2016). Moreover, in tobacco cell cultures subjected to

N starvation, antisense suppression of *AOX1* caused carbohydrate accumulation (Sieger et al.,

2005). This implies that, under limited nitrate reduction, oxidation of NADH via AOX could

replenish the NAD^+ to drive the glycolysis.

The above studies suggest that AOX is linked to nitrate reduction. However, these

studies manipulated nitrate reduction activity by transferring plants to the growth conditions that

included no/low nitrate or the use of ammonium as the sole N source. Reduced nitrate reduction

may therefore be accompanied by N starvation or ammonium toxicity (Hachiya et al., 2021a),

causing plant growth suppression and/or the initiation of stress responses. Moreover, since nitrate

acts as a signal to alter genome-wide gene expression (Liu et al., 2017; Wang et al., 2004), a

decrease in the nitrate supply also depletes the nitrate signal. For these reasons, it is impossible

to distinguish whether the above-mentioned effects of AOX deficiency are caused by reduced

nitrate reduction, N starvation, ammonium toxicity, or the lack of nitrate signal. To solve this

problem, we developed a novel culturing system to manipulate the degree of nitrate reduction and

AOX activity without causing N starvation, ammonium toxicity, and the lack of nitrate signal.

Using this system we examined genome-wide gene expression and plant growth to better

understand the relationship between AOX and nitrate reduction.

2 | MATERIALS AND METHODS

2.1 Plant materials and growth conditions

In this study we used *A. thaliana*, accession Columbia (Col-0) as a control line as well as the

aox1a-1 (SALK_084897, Giraud et al., 2008), *aox1a-2* (SAIL_030_D08, Giraud et al., 2008), *nr*

(*nia1-1/chl3-5*, Wang et al., 2004), *aox1a-1 nr*, and *aox1a-2 nr* mutants. The homozygous

multiple mutants of *aox1a-1 nr* and *aox1a-2 nr* were produced via crossing. The surface-sterilized

seeds were placed in the dark at 4°C for three days to break seed dormancy. For *in vitro* culturing,

we first grew plants on a sterile cellophane sheet (Hachiya et al., 2021b) placed on a solid medium

containing 30 mL half-strength Murashige and Skoog (1/2-MS) salts without nitrogen

supplemented with 2.5 mM $(NH_4)_2SO_4$, 0.1% (w/v) MES-KOH (pH 6.7), 0.5% (w/v) sucrose,

and 0.25% (w/v) gellan gum (Fujifilm Wako, Osaka, Japan). Plants were grown horizontally at

22°C for 18 days under a fixed photosynthetic photon flux density (PPFD) of 30 µmol m^{-2} s^{-1}

(i.e., constant light). Plants-on-cellophane were then transferred to a medium containing 30 mL

of 1/2-MS salts without nitrogen supplemented with 0.05% (w/v) MES-KOH (pH 5.7), 0.5%

(w/v) sucrose, and 0.25% (w/v) gellan gum (Fujifilm). The resulting mixture was incubated at

22°C for 24 h, also under a PPFD of 30 µmol m^{-2} s^{-1} (i.e., constant light). Finally, the plants-on-

cellophane were transferred to a medium containing 30 mL of 1/2-MS salts supplemented with

0.05% (w/v) MES-KOH (pH 5.7), 0.5% (w/v) sucrose, and 0.25% (w/v) gellan gum (Fujifilm)

and then incubated at 22°C under a PPFD of 150–200 µmol m^{-2} s^{-1} (constant light). For soil

cultivation, plants were grown for 14 or 24 days under a PPFD of 100–130 µmol m^{-2} s^{-1} (i.e., on

a 16h/8h light/dark cycle) at 22–23°C using an Arasystem 360 kit (Betatech BVBA, Gent,

Belgium) in a 1 : 1 mixture of nutrient-rich soil (Supermix A; Sakata Seed, Kyoto, Japan) and

vermiculite.

2.2 Determination of in vitro nitrate reductase activities

First, we assessed *in vitro* nitrate reductase (NR) activity using a previously published method

(Hachiya et al., 2016) with slight modification. Frozen samples were first ground in a Multi-Bead

Shocker (Yasui Kikai, Osaka, Japan) using zirconia beads. The resulting powder was then mixed

with 10 vol. of extraction buffer (50 mM HEPES-KOH, pH 7.6, 1 mM EDTA, 7 mM cysteine)

and the extracts were centrifuged at 20,400×g at 4°C for 10 min. Next, a 30 μl aliquot of the

supernatant was added to 90 μl of assay buffer (50 mM HEPES-KOH, pH 7.6, 133 μM NADH,

2 mM EDTA, 6.67 mM KNO_3). After incubation at 30°C for 2 and 17 min, 24 μl of 1% (w/v)

sulfanilamide solution in 1 N HCl and 24 μl of 0.02% (w/v) *N*-(1-naphthyl)ethylenediamine

dihydrochloride solution were added to 48 μl aliquots of reaction mixture. Finally, nitrite content

was determined based on absorbance readings at 540 nm. *In vitro* NR activity was calculated by

determining the nitrite amount produced over a 15 min period.

2.3 Determination of in vivo nitrate reductase activities

First, we submerged fresh shoots in 50 vols. of reaction buffer (i.e., 100 mM sodium phosphate

buffer, pH 7.4, 10 mM KNO_3, 4% (v/v) n-propanol), followed by vacuum-infiltration. The

reaction mixture was then incubated at 30°C for 1 h in the dark. The amount of nitrate produced

was then visualized by mixing the supernatant with 1% (w/v) sulfanilamide solution in 1 N HCl,

and 0.02% (w/v) *N*-(1-naphthyl)ethylenediamine dihydrochloride solution in a 2 : 1 : 1 ratio.

2.4 Determination of nitrate concentration

Nitrate content was quantified as per a previously published protocol (Hachiya et al., 2016) with

slight modifications. Nitrate was first extracted with 10 vol. of deionized water at 100°C for 20

min. Next, 10 μL of the supernatant was mixed with 40 μL of a reaction reagent (i.e., 5% (w/v)

salicylic acid in concentrated sulfuric acid), and the resulting mixture was then incubated at room

temperature for 20 min. A mock treatment of 40 μL of concentrated sulfuric acid was also

produced. Finally, 1 mL of 8% (w/v) NaOH solution was added to the mixture, and nitrate was

determined based on absorbance at 410 nm.

2.5 Determination of total protein

Total protein was determined according to a previously described method (Hachiya et al., 2016).

Frozen samples were first homogenized with a Multi-Beads Shocker (Yasui Kikai) using zirconia

beads. Total proteins were then extracted with 10 vol. of sample buffer [2% (w/v) SDS, 62.5 mM

Tris-HCl (pH 6.8), 10% (v/v) glycerol, and 0.0125% (w/v) bromophenol blue] and Halt^{TM}

protease inhibitor cocktail (ThermoFisher Scientific, Tokyo, Japan), followed by incubation at

95°C for 5 min. Extracts were then centrifuged at 20,400×g at 8°C for 10 min and 10 μL aliquots

were suspended in 500 μL deionized water. Next, 100 μL 0.15% (w/v) sodium deoxycholate was

added, and the mixture was incubated at room temperature for 10 min. 100 μL of 72% (v/v)

trichloroacetic acid was then added, followed by incubation at room temperature for 15 min and

centrifugation at 20,400×g for 10 min. Precipitates were then air-dried and suspended in 25 μL

water. This suspension was used to determine total protein concentrations using Takara BCA

Protein Assay Kits (TaKaRa, Kusatsu, Japan).

2.6 Immunodetection of AOX and ACTIN proteins

12.35 μL of total protein extract was mixed with 0.65 μl of 1 M DTT, followed by incubation at

95°C for 3 min. 5 μL of mixture (~0.43 mg fresh weight) was then subjected to SDS-PAGE using

a 12% Mini-PROTEAN TGX Gel (Bio-Rad, Hercules, CA, USA) before transferring to a Trans-

Blot Turbo Mini PVDF membrane using the Trans-Blot Turbo Transfer System (Bio-Rad).

Membranes were incubated for 1h in blocking buffer (50 mM Tris-HCl, and 150 mM NaCl, pH

7.6, 0.1% (v/v) Tween-20) containing 5% (w/v) ECL Prime Blocking Agent (Cytiva, Tokyo,

Japan). Membranes were then incubated for 1h with a 1/200 dilution of monoclonal antibodies

against AOX (AS10 699, Agrisera, Vännäs, Sweden) or a 1/2,500 dilution of polyclonal

antibodies against ACTIN (AS13 2640, Agrisera). Finally, the antigen–antibody complex was

detected using a 1/10,000 dilution of horseradish peroxidase conjugated with anti-mouse IgG

sheep antibodies (Cytiva) for AOX or anti-rabbit IgG donkey antibodies (Cytiva) for ACTIN by

ECL Prime chemiluminescent detection (Cytiva). Proteins blotted on membranes were visualized

using TaKaRa CBB Protein Safe Stain (TaKaRa) and chemiluminescence and CBB staining were

visualized using ImageQuant LAS 500 mini (Cytiva).

2.7 RNA extraction

Samples were first homogenized as described above. Next, total RNA was extracted using an

RNeasy Plant Mini Kit (Qiagen, Tokyo, Japan). For RNA-seq library preparation, RNA was

purified using on-column DNase digestion (Qiagen).

2.8 RNA-seq

RNA quality was first evaluated using a Qubit RNA IQ Assay Kit (ThermoFisher Scientific).

RNA samples with RNA IQs 8.7–10.0 were used for library preparation. cDNA libraries were

constructed using a NEBNext Ultra II RNA Library Prep Kit with Sample Purification Beads

(New England Biolabs, Tokyo, Japan), a NEBNext Poly(A) mRNA Magnetic Isolation Module

(New England Biolabs), and NEBNext Multiplex Oligos for Illumina (New England Biolabs).

cDNA libraries were then sequenced using a NextSeq 500 (Illumina, Tokyo, Japan), and resulting

bcl files were converted to fastq files using bcl2fastq (Illumina). The RNA-seq raw data are

available in ArrayExpress database under accession number E-MTAB-14027. Finally, obtained

reads were analyzed using iDEP version 096 (Ge et al., 2018), Metascape (Zhou et al., 2019), and

GeneCloud (Krouk et al., 2015) using the default settings.

2.9 RT-qPCR

Reverse transcription (RT) was performed using a ReverTraAce qPCR RT Master Mix with

gDNA Remover (Toyobo, Osaka Japan). Synthesized cDNA was then diluted tenfold with water

and used for quantitative PCR (qPCR). RT-qPCR was performed on a QuantStudio 1

(ThermoFisher Scientific) with KOD SYBR qPCR Mix (Toyobo). Relative transcript levels were

calculated using the comparative cycle threshold method with *ACTIN3* as an internal standard.

Primer sequences are shown in Table S1.

2.10. Statistical analysis

The Tukey–Kramer multiple comparison test were conducted using R software v.2.15.3.

3 | RESULTS and DISCUSSION

3.1 Manipulation of nitrate reduction and AOX activities

We manipulated the activities of nitrate reduction and AOX via the transfer experiment (Figure

1A) using Col-0 and the mutants deficient in either or both of NR and AOX activities. First, plants

were grown for 18 days with ammonium as the sole N source to ensure uniform growth

independently of NR activity. Higher medium pH and lower light intensity were used to reduce

ammonium toxicity and AOX expression (Escobar et al., 2006; Hachiya et al., 2010; Hachiya et

al., 2021a; Yoshida et al., 2011), allowing uniform growth regardless of AOX activity. Second,

plants were subjected to N starvation at pH 5.7 for 24 h to induce AOX expression (Escobar et

al., 2006; Hachiya et al., 2010). Finally, plants were transferred to a medium containing adequate

nitrate and incubated at an elevated light intensity, which induces NR expression and fills

photosynthetically-derived reductants to the cell (Noguchi & Yoshida, 2008; Wang et al., 2004;

Yoshida et al., 2010). The NR activity of Col-0 increased rapidly from 0 to 7 h after nitrate supply

and slightly more from 7 to 24 h (Figure S1A). Both *in vitro* and *in vivo* NR activities were

increased 7 h after nitrate supply in the shoots of Col-0 and *aox1a-1*, but not in *nr* and *aox1a-1*

nr (Figures 1B and 1C). Thus, we focused on the 7 h after nitrate supply to identify the early

effects of nitrate reduction for subsequent experiments. We found that nitrate accumulated in the

shoots of all lines after nitrate supply, with concentrations ranging from 14.7 to 21.6 μmol g^{-1}

(Figure 1D), indicating an adequate supply of nitrate for signaling. Actually, expression of *NIA2*

encoding the major NR isoform in leaves, which is inducible by nitrate signal (Konishi &

Yanagisawa, 2013; Wilkinson & Crawford, 1991), was strongly induced 7 h after nitrate supply

(Figure S1B). Higher nitrate concentrations in shoots of *nr* and *aox1a-1 nr* than Col-0 and *aox1a-*

1 would reflect a deficiency in nitrate reduction (Figure 1D). Shoot protein concentrations were

comparable among all lines before and after nitrate supply, ranging from 27.0 to 29.7 mg g^{-1}

(Figure 1E), suggesting that no N starvation occurred. RT-qPCR and Western blot analyses

confirmed that the signals corresponding to *AOX1a* and AOX were negligible in *aox1a-1* and

aox1a-1 nr (Figures S1C and S1D), suggesting that knockout of *AOX1a* is sufficient to diminish

AOX activity, as reported earlier (Giraud et al., 2008; Hachiya et al., 2010). Taken together, these

culturing conditions permit comparisons of plants differing nitrate reduction and AOX activity

without causing ammonium toxicity, N starvation, and lack of nitrate signaling.

3.2 AOX alleviates mitochondrial oxidative stress under limited nitrate reduction

Next, to dissect the role of AOX in limited nitrate reduction, we performed three independent

RNA-seq analyses using shoots from Col-0, *aox1a-1*, *aox1a-2*, *nr*, *aox1a-1 nr*, and *aox1a-2 nr* 7

h after nitrate supply (Table S2). A k-means clustering analysis classified transcripts into four

groups according to their expression patterns (Figure 2A and Table S3). In cluster D, *AOX1a*

deficiency consistently induced gene expression in the *nr* background, but not in the Col-0

background (Figures 2A and 2B). Moreover, 42 of 141 genes in cluster D were statistically

significantly upregulated in *aox1a-1 nr* and *aox1a-2 nr* relative to *nr* (Figure 2C and Table S4).

Further enrichment analyses of the 42 genes identified significant overrepresentation of the terms

"toxin catabolic process," "glutathione metabolism" (Figure 2D), "interpro-ipr004046

(glutathione S-transferase, C-terminal)," and "gst" (Figure 2E). These terms were derived from

the glutathione S-transferase (GST) genes *AT2G29460* (*GSTU4*), *AT1G17170* (*GSTU24*),

AT1G17180 (*GSTU25*), and *AT1G02920* (*GSTF7*) (Table S4). *GSTU4*, a plant-specific tau class

GST, may contribute to hydrogen peroxide degradation by using glutathione as an electron donor

(Santamaría et al., 2018). The terms "response to hypoxia" and "response to oxidative stress"

were also enriched in these 42 genes (Figure 2D). Furthermore, genes induced by hypoxia

(Tamura & Bono, 2022) and H_2O_2 treatments (Hieno et al., 2019) were significantly upregulated

in *nr*, and their induction was enhanced by *AOX1a* deficiency (Figures 2F and 2G and Tables S5

and S6). The NAC transcription factor ANAC017 mediates ROS-related mitochondrial retrograde

signaling, thereby activating mitochondrial dysfunction stimulation genes including *AOX1a*,

UPOX, and *ANAC013* (De Clercq et al., 2013; Ng et al., 2013). The expression of ANAC017-

inducible genes (Ng et al., 2013) and mitochondrial dysfunction stimulation genes (De Clercq et

al., 2013) reached a maximum level in *aox1a nr and aox1a-2 nr* (Figures 2H and 2I and Tables

S7 and S8). Further RT-qPCR analyses revealed that the expression levels of hypoxia-inducible

genes (Figures 2J–M) and oxidative stress marker genes (Figures 2N–Q) were induced by nitrate

supply in *nr*, and this induction was enhanced by *AOX1a* deficiency. Meanwhile, before nitrate

supply, little difference was observed among all lines (Figures 2J–Q). Since hypoxia elevates

intracellular $NADH/NAD^+$ ratio via mETC inhibition (Igamberdiev et al., 2005), the upregulation

of hypoxia-inducible genes may reflect reductant accumulation. Consequently, our transcriptome

analysis suggests that AOX dissipates excessive reductants and mitigate oxidative stress under

limited nitrate reduction.

The plant mETC possesses type II NAD(P)H dehydrogenases located on the cytosol

side (ND_{ex}) or matrix side (ND_{in}) of the inner mitochondrial membrane (Rasmusson et al., 2020).

Since these dehydrogenases transfer electrons to ubiquinone and bypass the proton-pumping

complex I, they can dissipate excessive reductants like AOX. Of these, the expression of *NDB2*,

encoding the primary ND_{ex} contributor to NADH oxidation, was significantly induced in *nr* after

nitrate supply, and this induction was intensified by *AOX1a* deficiency (Figures 3A and 3B). Since

cytosolic NR has a much lower K_m for NADH than ND_{ex} (Sweetman et al., 2019), NDB2 may

operate only when nitrate reduction is limited. The *NDA2*, encoding the NADH-oxidizing ND_{in}

(Rasmusson et al., 2020), also showed a expression pattern similar to *NDB2* after nitrate supply

(Fig. 3A and 3C). The compensated induction of *NDB2* and *NDA2* by *AOX1a* deficiency (Figures

3) suggests a contribution of AOX1a-NDB2 and AOX1a-NDA2 modules to balance cellular

redox under limited nitrate reduction. Also, the STRING database ver. 12.0 integrating protein-

protein interactions (Szklarczyk et al., 2023) confirmed tight functional connections between

AOX1a, NDB2, and NDA2. Meanwhile, the expression of the uncoupling proteins (UCPs), which

act as an uncoupler and/or as an aspartate/glutamate exchanger across the inner mitochondrial

membrane (Monné et al., 2018), was little changed among the lines (Figure 3A).

3.3 AOX sustains plant growth under limited nitrate reduction

Finally, we analyzed shoot growth parameters following nitrate supply. Over 6 days, we found

that *AOX1a* deficiency significantly reduced (−15%) shoot fresh weight in *nr*, but not in Col-0

(Figure 4A). Meanwhile, rosette diameter was little affected by *AOX1a* deficiency (Figure 4B).

When plants were grown in nutrient-rich soil, shoot appearance and shoot fresh weight were

decreased in *aox1a-1 nr* and *aox1a-2 nr* relative to the others (Figures 4C–E). However, in NR-

deficient plants, leaves were pale green (Figure 4C). Thus, it is plausible that long-term limitation

of nitrate reduction could cause N starvation, which complicates the interpretation of the

relationship between nitrate reduction, AOX, and shoot growth.

4 | CONCLUSIONS

We have successfully developed a cultural system to manipulate activities of nitrate reduction

and AOX without causing ammonium toxicity, N starvation, and lack of nitrate signaling.

Analyses using this system suggest that AOX alleviates mitochondrial oxidative stress and sustain

plant growth under limited nitrate reduction. It awaits further analyses to address physiological

and biochemical details using the system based on knowledge obtained from the present study.

AUTHOR CONTRIBUTIONS

DO and NO performed most of the measurements and downstream analyses. KM supported most

of the measurements and downstream analyses. TS performed next-generation sequencing

analysis. KN and TN supervised all analyses and interpretations. TH designed the study,

performed a part of growth analysis, and supervised all analyses and interpretations. All authors

contributed to the writing and/or reviewing of the manuscript.

ACKNOWLEDGEMENTS

aox1a-2 seeds were provided by Dr. James Whelan (La Trobe University, Australia). The authors

thank Enago for English language review.

DATA AVAILABILITY STATEMENT

The RNA-seq raw data are available in ArrayExpress under accession number E-MTAB-14027

(https://www.ebi.ac.uk/biostudies/arrayexpress).

REFERENCES

Andrews, M., Morton, J. D., Lieffering, M., Bisset, L., 1992. The partitioning of nitrate

assimilation between root and shoot of a range of temperate cereals and pasture grasses. *Ann. Bot.*

70(3), 271–276. https://doi.org/10.1093/oxfordjournals.aob.a088469

De Clercq, I., Vermeirssen, V., Van Aken, O., Vandepoele, K., Murcha, M. W., Law, S. R., Inzé,

A., Ng, S., Ivanova, A., Rombaut, D., van de Cotte, B., Jaspers, P., Van de Peer, Y., Kangasjärvi,

J., Whelan, J., Van Breusegem, F., 2013. The membrane-bound NAC transcription factor

ANAC013 functions in mitochondrial retrograde regulation of the oxidative stress response in

Arabidopsis. *Plant Cell* 25(9), 3472–3490. https://doi.org/10.1105/tpc.113.117168

Escobar, M. A., Geisler, D. A., Rasmusson, A. G., 2006. Reorganization of the alternative

pathways of the Arabidopsis respiratory chain by nitrogen supply: opposing effects of ammonium

and nitrate. *Plant J.* 45(5), 775–788. https://doi.org/10.1111/j.1365-313X.2005.02640.x

Ge, S. X., Son, E. W., Yao, R., 2018. iDEP: an integrated web application for differential

expression and pathway analysis of RNA-Seq data. *BMC Bioinform.* 19, 534.

https://doi.org/10.1186/s12859-018-2486-6

Giraud, E., Ho, L. H. M., Clifton, R., Carroll, A., Estavillo, G., Tan, Y. F., Howell, K. A., Ivanova,

A., Pogson, B. J., Millar, A. H., Whelan, J., 2008. The absence of ALTERNATIVE OXIDASE1a

in Arabidopsis results in acute sensitivity to combined light and drought stress. *Plant Physiol.*

147(2), 595–610. https://doi.org/10.1104/pp.107.115121

Hachiya, T., Terashima, I., Noguchi, K., 2007. Increase in respiratory cost at high growth

temperature is attributed to high protein turnover cost in *Petunia × hybrida* petals. *Plant Cell*

Environ. 30(10), 1269–1283. https://doi.org/10.1111/j.1365-3040.2007.01701.x

Hachiya, T., Watanabe, C. K., Boom, C., Tholen, D., Takahara, K., Kawai-Yamada, M.,

Uchimiya, H., Uesono, Y., Terashima, I., Noguchi, K., 2010. Ammonium-dependent respiratory

increase is dependent on the cytochrome pathway in *Arabidopsis thaliana* shoots. *Plant Cell*

Environ. 33(11), 1888–1897. https://doi.org/10.1111/j.1365-3040.2010.02189.x

Hachiya, T., Ueda, N., Kitagawa, M., Hanke, G., Suzuki, A., Hase, T., Sakakibara, H., 2016.

Arabidopsis root-type ferredoxin:NADP(H) oxidoreductase 2 is involved in detoxification of

nitrite in roots. *Plant Cell Physiol.* 57(11), 2440–2450. https://doi.org/10.1093/pcp/pcw158

Hachiya, T., Okamoto, Y., Watanabe, M.. Takebayashi, Y., Kojima, M., Suzuki, T., Sakakibara,

H., 2020. Genome-wide responses to shoot nitrate satiety are attenuated by external ammonium

in *Arabidopsis thaliana*. *Soil Sci. Plant Nutr.* 66(2), 317–327.

https://doi.org/10.1080/00380768.2020.1717905

Hachiya, T., Inaba, J., Wakazaki, M., Sato, M., Toyooka, K., Miyagi, A., Kawai-Yamada, M.,

Sugiura, D., Nakagawa, T., Kiba, T., Gojon, A., Sakakibara, H., 2021a. Excessive ammonium

assimilation by plastidic glutamine synthetase causes ammonium toxicity in *Arabidopsis thaliana*.

Nat. Commun. 12(1), 4944. https://doi.org/10.1038/s41467-021-25238-7

Hachiya, T., Oya, T., Monden, K., Nagae, A., Nakagawa, T., 2021b. A cellophane-supported

Arabidopsis culture for seamless transfer between different media is useful for studying various

nitrogen responses. *Soil Sci. Plant Nutr.* 67(3), 277–282.

https://doi.org/10.1080/00380768.2021.1908094

Hieno, A., Naznin, H. A., Inaba-Hasegawa, K., Yokogawa, T., Hayami, N., Nomoto, M., Tada,

Y., Yokogawa, T., Higuchi-Takeuchi, M., Hanada, K., Matsui, M., Ikeda, Y., Hojo, Y., Hirayama,

T., Kusunoki, K., Koyama, H., Mitsuda, N., Yamamoto, Y. Y., 2019. Transcriptome analysis and

identification of a transcriptional regulatory network in the response to H_2O_2. *Plant Physiol.*

180(3), 1629–1646. https://doi.org/10.1104/pp.18.01426

Igamberdiev, A. U., Baron, K., Manac'h-little, N., Stoimenova, M., Hill, R. D., 2005. The

haemoglobin/nitric oxide cycle: involvement in flooding stress and effects on hormone signalling.

Ann. Bot. 96(4), 557–564. https://doi.org/10.1093/aob/mci210

Iwagami, T., Ogawa, T., Ishikawa, T., Maruta, T., 2022. Activation of ascorbate metabolism by

nitrogen starvation and its physiological impacts in *Arabidopsis thaliana*. *Biosci. Biotech.*

Biochem. 86(4), 476–489. https://doi.org/10.1093/bbb/zbac010

Konishi, M., Yanagisawa, S., 2013. Arabidopsis NIN-like transcription factors have a central role

in nitrate signalling. *Nat. Commun.* 4(1), 1617. https://doi.org/10.1038/ncomms2621

Krouk, G., Carré, C., Fizames, C., Gojon, A., Ruffel, S., Lacombe, B., 2015. GeneCloud reveals

semantic enrichment in lists of gene descriptions. *Mol. Plant* 8(6), 971–973.

https://doi.org/10.1016/j.molp.2015.02.005

Miller, A. J., Cramer, M. D., 2004. Root nitrogen acquisition and assimilation. *Plant Soil* 274, 1–

36. https://doi.org/10.1007/s11104-004-0965-1

Monné, M., Daddabbo, L., Gagneul, D., Obata, T., Hielscher, B., Palmieri, L., Miniero, D. V.,

Fernie, A. R., Weber, A. P. M., Palmieri, F., 2018. Uncoupling proteins 1 and 2 (UCP1 and UCP2)

from *Arabidopsis thaliana* are mitochondrial transporters of aspartate, glutamate, and

dicarboxylates. *J. Biol. Chem.* 293(11), 4213-4227. https://doi.org/10.1074/jbc.RA117.000771

Ng, S., Ivanova, A., Duncan, O., Law, S. R., Van Aken, O., De Clercq, I., Wang, Y., Carrie, C.,

Xu, L., Kmiec, B., Walker, H., Van Breusegem, F., Whelan, J., Giraud, E., 2013. A membrane-

bound NAC transcription factor, ANAC017, mediates mitochondrial retrograde signaling in

Arabidopsis. *Plant Cell* 25(9), 3450–3471. https://doi.org/10.1105/tpc.113.113985

Noctor, G., Foyer, C. H., 1998. A re-evaluation of the ATP : NADPH budget during C_3

photosynthesis: a contribution from nitrate assimilation and its associated respiratory activity?. *J.*

Exp. Bot. 49(329), 1895–1908. https://doi.org/10.1093/jxb/49.329.1895

Noguchi, K., Terashima, I., 2006. Responses of spinach leaf mitochondria to low N availability.

Plant Cell Environ. 29(4), 710–719. https://doi.org/10.1111/j.1365-3040.2005.01457.x

Noguchi, K., Yoshida, K., 2008. Interaction between photosynthesis and respiration in

illuminated leaves. *Mitochondrion* 8(1), 87–99. https://doi.org/10.1016/j.mito.2007.09.003

Podgórska, A., Gieczewska, K. Łukawska-Kuźma, K., Rasmusson, A. G., Gardeström, P., Szal,

B., 2013. Long-term ammonium nutrition of *Arabidopsis* increases the extrachloroplastic

NAD(P)H/NAD(P)$^+$ ratio and mitochondrial reactive oxygen species level in leaves but does not

impair photosynthetic capacity. *Plant Cell Environ.* 36(11), 2034–2045.

https://doi.org/10.1111/pce.12113

Podgórska, A., Mazur, R., Ostaszewska-Bugajska, M., Kryzheuskaya, K., Dziewit, K., Borysiuk,

K., Wdowiak, A., Burian, M., Rasmusson, A. G., Szal, B., 2020. Efficient photosynthetic

functioning of *Arabidopsis thaliana* through electron dissipation in chloroplasts and electron

export to mitochondria under ammonium nutrition. *Front. Plant Sci.* 11, 103.

https://doi.org/10.3389/fpls.2020.00103

Qiao, X., Ruan, M., Yu, T., Cui, C., Chen, C., Zhu, Y., Li, F., Wang, S., Na, X., Wang, X., Bi,

Y., 2022. UCP1 and AOX1a contribute to regulation of carbon and nitrogen metabolism and yield

in Arabidopsis under low nitrogen stress. *Cell. Mol. Life Sci.* 79, 69.

https://doi.org/10.1007/s00018-021-04036-w

Rasmusson, A. G., Escobar, M. A., Hao, M., Podgórska, A., Szal, B., 2020. Mitochondrial

NAD(P)H oxidation pathways and nitrate/ammonium redox balancing in plants. *Mitochondrion*

53, 158–165. https://doi.org/10.1016/j.mito.2020.05.010

Santamaría, M. E., Arnaiz, A., Velasco-Arroyo, B., Grbic, V., Diaz, I., Martinez, M., 2018.

Arabidopsis response to the spider mite *Tetranychus urticae* depends on the regulation of reactive

oxygen species homeostasis. *Sci. Rep.* 8(1), 9432. https://doi.org/10.1038/s41598-018-27904-1

Scheurwater, I., Koren, M., Lambers, H., Atkin, O. K., 2002. The contribution of roots and shoots

to whole plant nitrate reduction in fast- and slow-growing grass species. *J. Exp. Bot.* 53(374),

1635–1642. https://doi.org/10.1093/jxb/erf008

Sieger, S. M., Kristensen, B. K., Robson, C. A., Amirsadeghi, S., Eng, E. W. Y., Abdel-Mesih,

A., Møller, I. M., Vanlerberghe, G. C., 2005. The role of alternative oxidase in modulating carbon

use efficiency and growth during macronutrient stress in tobacco cells. *J. Exp. Bot.* 56(416),

1499–1515. https://doi.org/10.1093/jxb/eri146

Sweetman, C., Waterman, C. D., Rainbird, B. M., Smith, P. M. C., Jenkins, C. D., Day, D. A.,

Soole, K. L., 2019. AtNDB2 is the main external NADH dehydrogenase in mitochondria and is

important for tolerance to environmental stress. *Plant Physiol.* 181(2), 774-788.

https://doi.org/10.1104/pp.19.00877

Szklarczyk, D., Kirsch, R., Koutrouli, M., Nastou, K., Mehryary, F., Hachilif, R., Gable, A. L.,

Fang, T., Doncheva, N. T., Pyysalo, S., Bork, P., Jensen, L. J., von Mering, C., 2023. The

STRING database in 2023: protein–protein association networks and functional enrichment

analyses for any sequenced genome of interest. *Nucleic Acids Res.* 51(D1), D638-D646.

https://doi.org/10.1093/nar/gkac1000

Tamura, K., Bono, H., 2022. Meta-Analysis of RNA sequencing data of Arabidopsis and Rice

under hypoxia. *Life* 12(7), 1079. https://doi.org/10.3390/life12071079

Wang, F., Wang, X., Zhao, C., Wang, J., Li, P., Dou, Y., Bi, Y., 2016. Alternative pathway is

involved in the tolerance of highland barley to the low-nitrogen stress by maintaining the cellular

redox homeostasis. *Plant Cell Rep.* 35, 317–328. https://doi.org/10.1007/s00299-015-1886-6

Wang, R., Tischner, R., Gutiérrez, R. A., Hoffman, M., Xing, X., Chen, M., Coruzzi, G.,

Crawford, N. M., 2004. Genomic analysis of the nitrate response using a nitrate reductase-null

mutant of Arabidopsis. *Plant Physiol.* 136(1), 2512–2522. https://doi.org/10.1104/pp.104.044610

Watanabe, C. K., Hachiya, T., Takahara, K., Kawai-Yamada, M., Uchimiya, H., Uesono, Y.,

Terashima, I., Noguchi, K., 2010. Effects of AOX1a deficiency on plant growth, gene expression

of respiratory components and metabolic profile under low-nitrogen stress in *Arabidopsis*

thaliana. *Plant Cell Physiol.* 51(5), 810–822. https://doi.org/10.1093/pcp/pcq033

Wilkinson, J. Q., Crawford, N. M., 1991. Identification of the *Arabidopsis CHL3* gene as the

nitrate reductase structural gene *NIA2*. *Plant Cell* 3(5), 461-471.

https://doi.org/10.1105/tpc.3.5.461

Yoshida, K., Shibata, M., Terashima, I., Noguchi, K., 2010. Simultaneous determination of in

vivo plastoquinone and ubiquinone redox states by HPLC-based analysis. *Plant Cell Physiol.*

51(5), 836–841. https://doi.org/10.1093/pcp/pcq044

Yoshida, K., Watanabe, C. K., Hachiya, T., Tholen, D., Shibata, M., Terashima, I., Noguchi, Ko.,

2011. Distinct responses of the mitochondrial respiratory chain to long- and short-term high-light

environments in *Arabidopsis thaliana*. *Plant Cell Environ.* 34(4), 618–628.

https://doi.org/10.1111/j.1365-3040.2010.02267.x

Zhou, Y., Zhou, B., Pache, L., Chang, M., Khodabakhshi, A. H., Tanaseichuk, O., Benner, C.,

Chanda, S. K., 2019. Metascape provides a biologist-oriented resource for the analysis of systems-

level datasets. *Nat. Commun.* 10(1), 1523. https://doi.org/10.1038/s41467-019-09234-6

SUPPORTING INFORMATION

Additional supporting information can be found online in the Supporting Information section at

the end of this article.

FIGURE LEGENDS

FIGURE 1 Manipulation of activities of nitrate reduction and AOX without causing N starvation,

ammonium toxicity, and the lack of nitrate signal.

(A) Schematic diagram of experiment. (B-E) *in vitro* NR activity (B), *in vivo* NR activity (C),

nitrate concentration (D), and total protein concentration (E) in plant shoots before and 7 h after

nitrate supply. Two plants of each line (eight in total) per plate were grown, and two shoots were

pooled as one biological replicate. Data: mean ± SD (n = 3 (B), n = 5–9 (D), n = 4 (E)). For (C),

color intensity is proportional to NR activity. Different lowercase letters indicate significant

differences determined via Tukey–Kramer tests at $P < 0.05$. ND and NS denote "not detected"

and "not significant." FW denotes "fresh weight."

FIGURE 2 Transcriptomic alteration by *AOX1a* deficiency under limited nitrate reduction

conditions.

(A–I) One plant from each line (six in total) per plate was grown. Six shoots were pooled as one

biological replicate. Shoots from plants 7 h after nitrate supply were subjected to RNA-seq.

Numbers indicate the order of RNA-seq experiments. (A) Heat map from a k-means clustering of

normalized transcript levels. Magenta and green represent higher and lower expression levels,

respectively. (B) Plots of normalized transcript levels in cluster D. Normalized transcript levels

of splice variants were averaged for each gene. (C) Venn diagram showing gene number from

cluster D and genes significantly upregulated in *aox1a-1 nr* or *aox1a-2 nr* relative to *nr*. (D, E)

Enriched terms identified by Metascape (D) and GeneCloud (E) from the 42 genes in (C). (E)

shows the number of genes containing the term (left) and the fold enrichment (right). (F–I) Shoot

expression of genes upregulated under hypoxia (F), by H_2O_2 (G), by ANAC017 (H), and included

in mitochondrial dysfunction stimulon (I) 7h after nitrate supply. Normalized transcript levels of

splice variants were averaged for each gene using means from three independent experiments. (J–

Q) RT-qPCR analysis of marker genes upregulated under hypoxia (J–M) and by oxidative stress

(N–Q) in shoots before and 7 h after nitrate supply. Two plants of each line (eight in total) per

plate were grown, and four shoots were pooled as one biological replicate. Data: mean ± SD (n =

5). (B, F–Q) Different lowercase letters indicate significant differences determined via Tukey–

Kramer tests at $P < 0.05$.

FIGURE 3 Effects of *AOX1a* deficiency on gene expression of type II NADP(H) dehydrogenases

and uncoupling proteins under limited nitrate reduction conditions.

(A) Transcriptional changes in genes encoding type II NADP(H) dehydrogenases (*NDA1/2*,

NDB1/2/4, *NDC1*), uncoupling proteins (*UCP1/2/3*), and *CI76* encoding the complex I 76-kD

subunit in plant shoots 7h after nitrate supply. Transcriptional changes are represented as log2

fold change ratios against Col-0 based on the RPM (reads per million mapped reads) from the

RNA-seq. Red and blue represent up- (>2-fold) and downregulation (<0.5 fold), respectively.

NDB3 was removed due to low expression. (B,C) Relative transcript levels of *NDB2* (B) and

NDA2 (C) in plant shoots before and 7 h after nitrate supply. Data: mean ± SD (n = 5). Different

lowercase letters indicate significant differences determined via Tukey–Kramer tests at $P < 0.05$.

FIGURE 4 Effects of *AOX1a* deficiency on shoot growth under limited nitrate reduction
conditions.

(A, B) Shoot fresh weight (A) and rosette diameter (B) of plants 6 days after nitrate supply. One
plant of each line (four in total) per plate were grown and one shoot was pooled as one biological
replicate. (C–E) Shoot appearance of 24-day-old plants (C) and shoot fresh weights of 14-day-
old (D) and 24-day-old (E) plants grown in pots containing nutrient-rich soil. Data: mean ± SD
(n = 17 (A, B), n = 18 (D), n = 7 (E)). Different lowercase letters indicate significant differences
determined via Tukey–Kramer tests at $P < 0.05$.

Figure 1

A

Procedure for transfer experiment

	Condition 1 (preculture) →	Condition 2 (preculture) →	Condition 3 (NO₃⁻ supply)
Duration	18 days	24 hours	0, 2, 4, 7, 24 hours or 6 days
N source	5 mM NH₄⁺	No nitrogen	19.7 mM NO₃⁻ & 10.3 mM NH₄⁺
pH	6.7	5.7	5.7
PPFD	30 μmol m⁻² s⁻¹	30 μmol m⁻² s⁻¹	150-200 μmol m⁻² s⁻¹

Col-0
aox1a-1
nr
aox1a-1 nr

B — *In vitro* NR activity

C — *In vivo* NR activity

Col-0   *aox1a-1*   *nr*   *aox1a-1 nr*

7 h

D — NO₃⁻

E — Total protein

Figure 2

Figure 3

A

Transcriptional changes (log2 fold change)

Gene	aox1a-1			aox1a-2			nr			aox1a-1 nr			aox1a-2 nr		
	1st	2nd	3rd	1st	2nd	3rd	1st	2nd	3rd	1st	2nd	3rd	1st	2nd	3rd
NDA1	-0.2	-0.4	0.0	-0.2	0.1	-0.1	-0.1	-0.2	0.1	0.4	0.2	0.4	0.2	0.3	0.6
NDA2	-0.2	0.4	-1.1	-0.2	-0.2	-0.5	-0.0	0.8	-0.2	0.7	2.5	1.2	0.6	1.1	1.6
NDC1	0.0	0.2	-0.0	-0.1	0.3	0.0	0.0	0.1	0.1	-0.1	-0.0	0.0	0.1	0.1	-0.2
NDB1	-0.1	-0.0	-0.1	-0.0	-0.1	0.0	0.1	0.2	-0.1	-0.1	-0.1	0.1	-0.1	-0.0	-0.3
NDB2	0.4	-0.1	-0.5	0.4	0.1	-0.1	0.9	1.2	0.4	1.9	2.4	1.7	2.4	2.0	1.8
NDB4	-1.6	1.0	0.3	0.1	-	2.7	-1.0	1.9	1.6	0.3	3.3	2.7	-0.6	1.7	2.6
UCP1	-0.1	0.3	0.3	-0.1	0.1	0.2	-0.2	-0.1	0.1	-0.1	0.1	0.1	-0.1	0.1	0.3
UCP2	-0.1	-0.0	-0.3	-0.1	-0.1	0.1	-0.2	-0.2	0.1	-0.2	-0.3	-0.1	-0.2	-0.3	-0.2
UCP3	-0.4	0.2	-0.1	-0.4	0.3	0.0	-0.6	0.4	0.1	-0.7	-0.2	0.0	-0.4	0.2	-0.5
CI76	-0.1	0.1	-0.2	0.0	-0.0	-0.1	-0.1	0.1	-0.1	-0.0	-0.0	-0.2	-0.0	-0.0	-0.3

Figure 4

A Shoot FW

B Rosette diameter

C

Col-0   aox1a-1   aox1a-2

nr   aox1a-1 nr   aox1a-2 nr

2 cm

D Shoot FW

14-d

E Shoot FW

24-d

Col-0
aox1a-1
aox1a-2
nr
aox1a-1 nr
aox1a-2 nr